Non Parametric Hidden Markov Models with Finite State Space: Posterior Concentration Rates

Elodie Vernet

Laboratoire de Mathématiques d'Orsay, Univ. Paris-Sud, CNRS, Université Paris-Saclay, 91405 Orsay, France elodie.vernet@math.u-psud.fr

January 16, 2018

Abstract

The use of non parametric hidden Markov models with finite state space is flourishing in practice while few theoretical guarantees are known in this framework. Here, we study asymptotic guarantees for these models in the Bayesian framework. We obtain posterior concentration rates with respect to the L_1 -norm on joint marginal densities of consecutive observations in a general theorem. We apply this theorem to two cases and obtain minimax concentration rates. We consider discrete observations with emission distributions distributed from a Dirichlet process and continuous observations with emission distributions distributed from Dirichlet process mixtures of Gaussian distributions.

Contents

D Proof of Theorem [4.3](#page-13-2) (Dirichlet process mixtures of Gaussian distributions) [31](#page-30-0)

1 Introduction

Hidden Markov models (HMMs) are stochastic processes much used in practice in fields as diverse as genomics, speech recognition, econometrics or climate. A hidden Markov chain is a sequence $(X_t, Y_t)_{t \in \mathbb{N}}$ where the sequence $(X_t)_{t \in \mathbb{N}}$ is a non observed Markov chain and the sequence of observations $(Y_t)_{t \in \mathbb{N}}$ is a noisy version of the chain $(X_t)_{t \in \mathbb{N}}$. In this paper we consider the case where the state space of the underlying Markov chain is finite. In this situation, HMMs are often employed to classify dependent data with respect to the hidden states X_t , $t \in \mathbb{N}$. Their popularity is due to their tractability. Since their introduction in [\[Baum and Petrie, 1966\]](#page-33-0), many algorithms have been developed to infer these models. The books [Cappé et al., 2005], [\[MacDonald and Zucchini, 1997\]](#page-35-0) and [\[MacDonald and](#page-35-1) [Zucchini, 2009\]](#page-35-1) give an overview of this family of models.

Parametric HMMs suffer from a lack of robustness so that non parametric HMMs are used more and more in applications. Indeed two constraints weaken parametric HMMs: the necessary assumption of a bound on the number of states of the Markov chain and the limitations of the parametric modeling of emission distributions (the distributions of an observation Y_t given the hidden states X_t). To deal with these issues, HMMs with an infinite countable number of states for the Markov chain are applied in [\[Beal](#page-33-2) [and Krishnamurthy, 2012\]](#page-33-2) to gene expression time course clustering, [\[Jochmann, 2015\]](#page-35-2) to U.S. inflation dynamics and in [\[Fox et al., 2009\]](#page-34-0) to segmentation of visual motion capture data. To handle speaker diarization, [\[Fox et al., 2011\]](#page-34-1) proposes a model where the number of states of the Markov chain is not bounded and the emission distributions are not restricted to live in a parametric family. HMMs, where the number of states is known but the emission distributions set is not assumed to be parametric, are used in [\[Langrock et al., 2015\]](#page-35-3) for whales dive modeling, [\[Yau et al., 2011\]](#page-35-4) for genetic copy number variants, Whiting et al., 2003] for climate state identification, [Lefèvre, 2003] for speech recognition, [\[Gassiat et al., 2015\]](#page-34-2) for gene expression identification, see also the references herein. This last framework, namely HMMs where the number of states of the Markov chain is known and emission distributions may live in infinite-dimensional sets is the one we consider in this paper.

The use of non parametric HMMs is flourishing in practice while few theoretical properties are known. Many theoretical results exist for parametric HMMs particularly for the maximum likelihood estimator, see Cappé et al. [2005] and references herein for instance, see also [\[de Gunst and Shcherbakova, 2008\]](#page-34-3) for a Bernstein von Mises property of the posterior. In the non parametric framework, there exist few theoretical guarantees of the asymptotic behavior of estimators or posterior since identifiability for general HMMs with finite state space was still an issue until recently. General identifiability is proved in [\[Gassiat et al., 2015\]](#page-34-2) when the number of states of the Markov chain is known and in [\[Alexandrovich and Holzmann, 2014\]](#page-33-3) when this number is unknown. [\[Gassiat et al., 2015\]](#page-34-2) proves that under mild assumptions, the knowledge of the marginal joint density of at least three consecutive observations (Y_t, Y_{t+1}, Y_{t+2}) gives the parameters of the HMM up to label switching (i.e. the transition matrix of the Markov chain and the emission distributions). Here, we are interested in obtaining asymptotics in the Bayesian framework for the marginal joint density of consecutive observations.

In the Bayesian non parametric setting, asymptotic analysis typically takes the following two forms: posterior consistency and posterior concentration rates. The posterior is said to be consistent at a parameter θ^* if it concentrates its mass around θ^* , when the ob-

servations come from θ^* and the number of observations increases. Posterior consistency is related to the merging of posteriors distributions associated to two priors, see [\[Diaconis](#page-34-4) [and Freedman, 1986\]](#page-34-4). In a non parametric setup, where it is not feasible to construct a fully subjective prior (on an infinite dimensional space), it is a minimal requirement, see [\[Ghosh and Ramamoorthi, 2003\]](#page-34-5). To go further on, one can study the rate at which this concentration occurs. Obtaining a minimax posterior concentration rates is a criterion of optimality. In particular, minimax concentration rates lead to minimax Bayesian estimators [\[Ghosal et al., 2000\]](#page-34-6) and to minimax size of credible regions [\[Hoffmann et al., 2013\]](#page-34-7). The concentration rate analysis also allows a better understanding of the impact of the prior, see [\[Rousseau, 2015\]](#page-35-7) for a discussion.

In Bayesian HMMs where the number of states of the Markov chain is known, [\[Vernet](#page-35-8) [et al., 2015\]](#page-35-8) provides assumptions leading to posterior consistency for the L_1 -norm of the marginal density of consecutive observations. Here, we pursue the study of the asymptotic behavior of the posterior distribution in this framework and with the same topology. Namely, we study posterior concentration rates for non parametric HMMs with respect to the L_1 -norm of the marginal joint density of consecutive observations. We first give a general theorem relating the posterior concentration rate to the prior and the true model (Theorem [3.1\)](#page-6-0). Then we apply the theorem to different setups, where we obtain minimax rates (Section [4\)](#page-9-0). To the best of our knowledge, these are the first results on posterior concentration rates in non parametric HMMs.

Let us mention the few other asymptotic results we know in the framework of non parametric HMMs. In the non parametric frequentist framework with a finite and known number of states, [\[De Castro et al., 2015b\]](#page-34-8) offers an oracle inequality for a penalized least-squares estimator of the emission distributions. In the framework of HMMs with an unknown number of states and emission distributions living in a finite-dimensional set, posterior concentration rates are studied in [\[Gassiat and Rousseau, 2014\]](#page-34-9). [\[Gassiat and](#page-34-10) [Rousseau, 2013\]](#page-34-10) proposes asymptotics for the particular case of translated HMMs with finite state space. Finally, convergence with respect to smoothing distributions is studied in [\[De Castro et al., 2015a\]](#page-34-11).

The paper is organized as follows. In Section [2,](#page-2-0) we precise the studied model and the notations. In Section [3,](#page-5-0) we state general assumptions under which the posterior concentration rate is derived (Theorem [3.1\)](#page-6-0). We have chosen to define a set of assumptions as close as possible to those typically obtained in density estimation for i.i.d. models, see [\[Ghosal et al., 2000\]](#page-34-6). The proof of this theorem is given in Section [3.2.](#page-7-0) All the other proofs are postponed in the appendices. Theorem [3.1](#page-6-0) is applied in Section [4](#page-9-0) in two cases. In Section [4.1,](#page-10-0) the observations are assumed to be discrete and the prior on emission distributions is based on a Dirichlet process. We obtain a minimax rate which is $1/\sqrt{n}$ up to a power of $\log n$ in Corollary [4.2.](#page-11-1) In Section [4.2,](#page-12-0) the observations are assumed to be continuous and the emission distributions follow independently Dirichlet process mixtures of Gaussian distribution. Minimax rates of concentration are obtained for Hölder-type functional classes, see Corollary [4.4.](#page-13-3)

2 Bayesian hidden Markov models and notations

We consider observations coming from homogeneous hidden Markov models with finite state space. Hidden Markov chains are discrete time stochastic processes $(X_t, Y_t)_{t \in \mathbb{N}}$ satisfying the following properties. The sequence $(X_t)_{t\in\mathbb{N}}$ is a Markov chain. Conditionally on the hidden chain $(X_t)_{t\in\mathbb{N}}$, the observations Y_t are independent with Y_t only depending on X_t . The states $(X_t)_{t\in\mathbb{N}}$ are latent, they are called the hidden states. The statistician observes the sequence $(Y_t)_{t\leq n}$ where n is an integer. Throughout the paper, for any integer *n*, an *n*-uple (y_1, \ldots, y_n) is denoted $y_{1:n}$.

We first introduce the notations concerning the Markov chain $(X_t)_{t\in\mathbb{N}}$. For all $t \in \mathbb{N}, X_t$

belongs to $\{1, \ldots, k\}$, where k is assumed to be known in this paper. A transition matrix Q and an initial probability distribution μ describe the distribution of the underlying Markov chain

$$
X_1 \sim \sum_{1 \le i \le k} \mu_i \delta_i, \quad X_t | (X_{t-1} = i) \sim \sum_{1 \le j \le k} Q_{i,j} \delta_j
$$

where δ_i denotes the Dirac measure at i. The set of all initial probability distributions is the $k-1$ -simplex $\Delta_k = \{x \in [0,1]^k : \sum_{1 \leq i \leq k} x_i = 1\}$. We denote Δ_k^k the set of all transition matrices such that each row of the matrix is an element of Δ_k . In the following we need $\Delta_k(q) = \{ \mu \in \Delta_k : \mu_i \geq \underline{q} \,\,\forall i \}$ and $\Delta_k^k(q) = \{ Q \in \Delta_k^k : Q_{i,j} \geq \underline{q} \,\,\forall i,j \}$, for $q \in (0,1)$. When Q is in $\Delta_k^k(q)$, with $q \in (0,1)$, then the uniform mixing coefficients, defined in [\[Rio, 2000\]](#page-35-9), associated to the corresponding Markov chain are bounded by $\phi(m) \leq (1-q)^m$, moreover the corresponding Markov chain is irreducible and positive recurrent.

The observations Y_t are assumed to live in \mathbb{R}^d which is endowed with its Borel sigma field. The distribution of Y_t is assumed to be absolutely continuous with respect to some measure λ on \mathbb{R}^d . Conditionally on $(X_t)_{t\in\mathbb{N}}$, Y_t is distributed from a distribution $f_{X_t}\lambda$ depending on the state X_t :

$$
Y_t|(X_s)_{s\in\mathbb{N}} \sim Y_t|X_t \sim f_{X_t}\lambda.
$$

The distributions $f_i \lambda$, $1 \leq i \leq k$ are called the emission distributions. The set of probability density functions with respect to λ is denoted F. The vector $f = (f_1, \ldots, f_k) \in \mathcal{F}^k$ is formed with the k emission density functions.

Then the model is completely described by the parameters μ and $\theta = (Q, f)$ where $\mu \in \Delta^k$ and $\theta = (Q, f) \in \Delta^k_k \times \mathcal{F}^k =: \Theta$. The model can be visualized in Figure [1.](#page-3-0)

Figure 1: The model

Let $P^{\mu,\theta}$ be the probability distribution of the process $(X_t, Y_t)_{t \in \mathbb{N}}$ under (μ, θ) . Then for any $\theta \in \Theta$, initial probability μ , and measurable set A of $\{1,\ldots,k\}^{\ell} \times (\mathbb{R}^d)^{\ell}$, note that:

$$
P^{\mu,\theta}((X_{1:\ell},Y_{1:\ell})\in A)=\int\sum_{x_1,\dots,x_\ell=1}^k1_{(x_1,\dots,x_\ell,y_1,\dots,y_\ell)\in A}\mu_{x_1}Q_{x_1,x_2}\dots Q_{x_{\ell-1},x_\ell}
$$

$$
f_{x_1}(y_1)\dots f_{x_\ell}(y_\ell)\lambda(dy_1)\dots\lambda(dy_\ell).
$$

Note that when Q is in $\Delta_k^k(q)$, with q positive, there exists a unique stationary initial distribution μ^Q associated with Q. When μ is not specified, the stationary distribution associated with the transition matrix Q is considered in the place of μ . In other words, we define $P^{(Q,f)} := P^{\mu^Q,(Q,f)}$. The joint distribution of ℓ consecutive observations $((Y_1, \ldots Y_\ell))$ for instance) under the stationary process associated with θ is denoted P_ℓ^{θ} . Let p_ℓ^{θ} denote the density of P_ℓ^{θ} with respect to $\lambda^{\otimes \ell}$. Then,

$$
p_{\ell}^{\theta}(y_1,\ldots,y_{\ell})=\sum_{x_1,\ldots,x_{\ell}=1}^k\mu_{x_1}Q_{x_1,x_2}\ldots Q_{x_{\ell-1},x_{\ell}}f_{x_1}(y_1)\ldots f_{x_{\ell}}(y_{\ell}),\quad \lambda^{\otimes \ell} \text{ a.s.}.
$$

The log-likelihood for a sequence of observations $Y_{1:\ell}$ under a parameter θ is denoted

$$
L_{\ell}^{\theta} := \log (p_{\ell}^{\theta}(Y_1, \ldots, Y_{\ell})).
$$

The dependency of L^{θ}_{ℓ} with $Y_{1:\ell}$ is implicit and can be deduced from the context.

Working in the Bayesian framework, we put a prior Π on the set of parameters Θ. We choose a product probability measure $\Pi = \Pi_Q \otimes \Pi_f^{(k)}$ where Π_Q is a probability distribution on Δ_k^k and $\Pi_f^{(k)}$ is a probability distribution on \mathcal{F}^k . To a realization θ from Π , we implicitly associate a stationary initial distribution μ^{Q} . In other words, we generalize Π to a distribution on Δ_k × Θ such that under Π and conditionally on $θ = (Q, f), μ = μ^Q$. Then using the Bayes' theorem, the posterior is expressed by

$$
\Pi(\theta \in A | Y_{1:n}) = \frac{\int_A p_n^{\theta}(Y_{1:n}) \Pi(d\theta)}{\int_{\Theta} p_n^{\theta}(Y_{1:n}) \Pi(d\theta)}.
$$

We are interested in the asymptotic behaviour of the posterior that is to say when the number n of observations $Y_{1:n}$ tends to infinity. For this purpose, we take a frequentist point of view, assuming that the observations come from the true parameters μ^* and $\theta^* = (Q^*, f^*)$. We suppose that the true initial distribution μ^* is stationary. We also assume that there exists $q^* > 0$ such that

$$
Q^* \in \Delta_k^k(\underline{q}^*). \tag{1}
$$

[\[Vernet et al., 2015\]](#page-35-8) shows posterior consistency at θ^* under general assumptions. In this paper, we consider posterior concentration rates at θ^* . Recall that the posterior is said to concentrate at rate ϵ_n , a sequence decreasing to 0, for the loss $D(\cdot, \cdot)$ if there exists a constant $M > 0$ such that

$$
\Pi(\theta : D(\theta, \theta^*) \ge M\epsilon_n | Y_{1:n}) = o_{P^{\theta^*}}(1),
$$

where $Z = o_{P^{\theta^*}}(1)$ means that Z converges in probability to 0. We choose to study the concentration of the posterior from the density estimation point of view. We compare two parameters θ and $\tilde{\theta}$ by computing the L₁-distance between the joint densities $p_\ell^{\tilde{\theta}}$ and $p_\ell^{\tilde{\theta}}$. For two distributions P_1 and P_2 , let p_1 and p_2 be their respective densities with respect to a dominated measure ν . The L_1 -metric is defined by

$$
||p_1 - p_2||_{L_1(\nu)} = \int |p_1 - p_2|\nu
$$

and let

$$
KL(p_1, p_2) = \int p_1 \log \left(\frac{p_1}{p_2}\right) \nu
$$

be the Kullback-Leibler divergence between p_1 and p_2 . For an integer $\ell \geq 1$, we use the pseudo-distance D_ℓ on Θ defined by

$$
D_{\ell}(\theta, \tilde{\theta}) = \|p_{\ell}^{\theta} - p_{\ell}^{\tilde{\theta}}\|_{L_1(\lambda^{\otimes l})}.
$$

We study the posterior rate of concentration with respect to this pseudo-distance D_{ℓ} . On \mathcal{F}^k , we use the distance $d(\cdot, \cdot)$ such that for all $(f, \tilde{f}) \in (\mathcal{F}^k)^2$

$$
d(f, \tilde{f}) = \max_{1 \le i \le k} ||f_i - \tilde{f}_i||_{L_1(\lambda)},
$$

on \mathbb{R}^d , $d \geq 2$, we use the supremum norm $\|\cdot\|$. For a positive real ϵ , a pseudo distance D defined on a set A, let $N(\epsilon, A, D)$ be the covering number that is to say the minimum number of balls of radius ϵ (in the pseudo-distance D) needed to cover A. Throughout the paper the notation \leq means less or equal up to a multiplicative constant which is not important in the context.

3 General Theorem

3.1 Assumptions and main theorem

In this section, we state the general Theorem [3.1](#page-6-0) which gives posterior concentration rates with respect to the D_ℓ pseudo-metric. As in [\[Ghosal et al., 2000\]](#page-34-6) for instance, we propose a set of conditions which relates the rate $\epsilon_n/\underline{q}_n$ to the prior and the true model. We apply this theorem to the case of discrete observations in Section [4.1](#page-10-0) and to the case of continuous observations in Section [4.2](#page-12-0) where minimax rates are achieved. Now, we enumerate the assumptions of Theorem [3.1.](#page-6-0) Assumptions [\(A\)](#page-5-2) and [\(B\)](#page-5-3) concern the prior on the emission distributions $\Pi_f^{(k)}$ and the vector of the true emission distributions f^* . Assumptions [\(C\)](#page-6-1) and [\(D\)](#page-6-2) involve the prior on transition matrices Π_Q and the true transition matrix Q^* .

For a given sequence $\epsilon_n > 0$ tending to 0, we introduce a sequence $\tilde{\epsilon}_n$ such that $\tilde{\epsilon}_n \leq \epsilon_n$ and the sequence u_n of positive numbers such that

(i)
$$
u_n = 1
$$
, for all $n \in \mathbb{N}$; if $\tilde{\epsilon}_n \gtrsim n^{-s}$, for some $s < 1/2$,
\n(ii) $(1-\epsilon)^{3/2} \epsilon$, $u_n = N - 1$, (2)

(ii)
$$
u_n = (\log(n))^{3/2}
$$
, for all $n \in \mathbb{N}$; otherwise.

We consider the following assumptions

(A) there exist a positive constant C_f and a sequence B_n of subsets of \mathcal{F}^k such that

$$
\Pi_f^{(k)}(B_n) \gtrsim \exp(-C_f n \tilde{\epsilon}_n^2)
$$

and such that for all $f \in B_n$,

$$
\int f_i^*(y) \log^2 \left(\frac{f_j^*(y)}{f_j(y)} \right) \lambda(dy) \le \frac{\tilde{\epsilon}_n^2}{u_n}, \text{ for all } 1 \le i, j \le k,
$$
\n(A.1)

there exist a set $S \subset \mathcal{Y}$ and functions $\tilde{f}_1, \ldots, \tilde{f}_k$ satisfying

$$
\int_{S} \frac{|f_j^*(y) - f_j(y)|^2}{f_j^*(y)} \lambda(dy) \le \frac{\tilde{\epsilon}_n^2}{u_n}, \text{ for all } 1 \le j \le k,
$$
\n(A.2)

$$
\int_{S^c} \tilde{f}_j(y)\lambda(dy) \le \tilde{\epsilon}_n^2, \text{ for all } 1 \le j \le k,
$$
\n(A.3)

$$
\int_{S^c} f_j^*(y)\lambda(dy) \le \frac{\tilde{\epsilon}_n^2}{u_n}, \text{ for all } 1 \le j \le k,
$$
\n(A.4)

$$
\int_{S} f_{i}^{*}(y) \max_{1 \le j \le k} \log \left(\frac{\tilde{f}_{j}(y)}{f_{j}(y)} \right) \lambda(dy) \le \tilde{\epsilon}_{n}^{2}, \text{ for all } 1 \le i \le k,
$$
\n(A.5)

$$
\int_{S} \frac{|f_j^*(y) - \tilde{f}_j(y)|^2}{\tilde{f}_j(y)} \lambda(dy) \le \tilde{\epsilon}_n^2 \quad \text{for all } 1 \le j \le k,
$$
\n(A.6)

(B) there exist positive constants C and C' and a sequence $(\mathcal{F}_n)_{n\geq 1}$ of subsets of \mathcal{F}^k such that

$$
\Pi_f^{(k)}(\mathcal{F}_n^c) = o(\exp(-Cn\tilde{\epsilon}_n^2))
$$

and

$$
N\left(\frac{\epsilon_n}{12}, \mathcal{F}_n, d\right) \lesssim \exp\left(C'n\epsilon_n^2\right),
$$

(C) there exists a positive constant C_Q such that $C_Q + C_f + C_K < C$ with $C_K = 4 +$ $\log (2k/q^*) + 10^4 k^2/q^{*5},$

$$
\Pi_Q\left(\left\{Q: \|Q - Q^*\| \le \frac{\tilde{\epsilon}_n}{\sqrt{u_n}}\right\}\right) \gtrsim \exp(-C_Q n \tilde{\epsilon}_n^2),
$$

(D) there exists a sequence \underline{q}_n of $(0,1/k]$ such that

$$
\Pi_Q\Big(\big(\Delta_k^k(\underline{q}_n)\big)^c\Big) = o(\exp(-Cn\tilde{\epsilon}_n^2)).
$$

Under the above assumptions, we prove that the posterior distribution concentrates at the rate $\epsilon_n/\underline{q}_n$.

Theorem 3.1. Let $\epsilon_n \geq \tilde{\epsilon}_n > 0$ be two sequences tending to 0 such that $n\tilde{\epsilon}_n^2 \to +\infty$. Assume (A) , (B) , (C) and (D) .

Then there exists a positive constant M such that

$$
\Pi\left(\theta: D_{\ell}(\theta, \theta^*) \ge M \frac{\epsilon_n}{\underline{q}_n} \middle| Y_{1:n} \right) = o_{\mathbb{P}^{\theta^*}}(1). \tag{3}
$$

We now discuss Assumptions [\(A\)](#page-5-2) to [\(D\)](#page-6-2). We have purposely considered assumptions which are as similar as possible to those considered in the set up of density estimation with i.i.d. observations see [Ghosal et al. \[2000\]](#page-34-6). In particular Assumption [\(B\)](#page-5-3) is the typical entropy assumption on a sieve which captures most of the prior mass (see Assumptions (2.2) and (2.3) of [Ghosal et al. \[2000\]](#page-34-6)). Assumption [\(A\)](#page-5-2) is slightly more involved than the Kullback-Leibler condition (2.4) of [Ghosal et al. \[2000\]](#page-34-6) in the case of i.i.d. observations.

This paragraph explains the differences between Assumption [\(A\)](#page-5-2) and condition (2.4) of [Ghosal et al. \[2000\]](#page-34-6) and can be omitted at first reading. Following [Ghosal and van der](#page-34-12) [Vaart \[2007b\]](#page-34-12), posterior rates of convergence are obtained by controlling "Kullback-Leibler" neighborhoods and constructing some tests. Under [\(A\)](#page-5-2) and condition (2.4) of [Ghosal et al. \[2000\]](#page-34-6), the Kullback-Leibler neighborhoods are controlled. Recall that in [Ghosal et al. \[2000\]](#page-34-6), the control of $\mathbb{E}^{\theta^*}(L_n^{\theta^*}-L_n^{\theta})$ and $Var^{\theta^*}(L_n^{\theta^*}-L_n^{\theta})$ is obtained by controlling

$$
\int f_i^*(y) \log \left(\frac{f_i^*(y)}{f_i(y)} \right) \lambda(dy) \quad \text{and} \quad \int f_i^*(y) \log^2 \left(\frac{f_i^*(y)}{f_i(y)} \right) \lambda(dy).
$$

Here $\mathbb{E}^{\theta^*}(L_n^{\theta^*} - L_n^{\theta}) \lesssim n\tilde{\epsilon}_n^2$ and $Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta}) \lesssim n\tilde{\epsilon}_n^2 \log n$ if f and f^* satisfy Assumptions There $\mathbb{E}[(L_n - L_n) \gtrsim n\epsilon_n$ and V at $(L_n - L_n) \gtrsim n\epsilon_n$ log n if f and f satisfy Assumptions $(A.1)$ – $(A.6)$ and $||Q - Q^*|| \leq \tilde{\epsilon}_n/\sqrt{u_n}$. The non intuitive part of (A) comes from the introduction of \tilde{f}_j as an approximation of f_j^* , which may be different from f_j in [\(A.3\)](#page-5-6), [\(A.5\)](#page-5-7) and [\(A.6\)](#page-5-5). Indeed, whithout the introduction of $(f_{j,1\leq j\leq k}, S)$, the HMM structure of the likelihood would lead to a crude upper bound of $\mathbb{E}^{\theta^*}(\overline{L}_n^{\theta^*}-L_n^{\theta})$ of the form

$$
n \int f_i^*(y) \max_j \log \left(\frac{f_j^*(y)}{f_j(y)} \right) \lambda(dy),
$$

see Equations [\(20\)](#page-14-0) and [\(21\)](#page-14-1) in the proof of Lemma [3.2.](#page-7-1) Since i may be different from j we would loose the local quadratic approximation of the Kullback-Leibler divergence and we would only obtain $n\tilde{\epsilon}_n$ instead of $n\tilde{\epsilon}_n^2$ as an upper bound of $\mathbb{E}^{\theta^*}(L_n^{\theta^*}-L_n^{\theta})$. The details of the control of the Kullback-Leibler divergence are given in Appendix [A.](#page-13-1)

Not withstanding the technical aspects discussed above, Assumptions $(A.1)$ – $(A.6)$ are verified using techniques similar to those used in the case of density estimation with i.i.d. observations to control

$$
\int f_i^*(y) \log \left(\frac{f_i^*(y)}{f_i(y)} \right) \lambda(dy) \quad \text{and} \quad \int f_i^*(y) \log^2 \left(\frac{f_i^*(y)}{f_i(y)} \right) \lambda(dy).
$$

For $\Pi_f^{(k)} = (\Pi_f)^{\otimes k}$, and many families of individual prior models Π_f on the f_j 's, the rate obtained by bounding $\max_j KL(f_j^*, f_j)$ in the i.i.d. set up will be the same as in our setup. For instance, if $\mathcal{Y} = [0, 1]$ and f^* is bounded from below and above, a control of $||f_j - f_j^*||_\infty^2 \leq \tilde{\epsilon}_n^2$ or $||f_j - f_j^*||_2^2 \leq \tilde{\epsilon}_n^2$ and $f_j > c$ imply $(A.1)$ – $(A.6)$. This kind of controls have been derived under (hierarchical) Gaussian process priors or log linear priors as in [van der Vaart and van Zanten \[2009\]](#page-35-10), [Arbel et al. \[2013\]](#page-33-4) and [Rivoirard and Rousseau](#page-35-11) [\[2012\]](#page-35-11). Condition [\(A\)](#page-5-2) becomes more involved when $\mathcal Y$ is not compact. This case is treated under non parametric Gaussian mixtures in Section [4.2](#page-12-0) and in the case where $\mathcal{Y} = \mathbb{N}$ in Section [4.1.](#page-10-0)

Assumption [\(C\)](#page-6-1) is checked as soon as Π_Q admits a positive density with respect to the Lebesgue measure which is continuous at Q^* and $\tilde{\epsilon}_n \geq \sqrt{\log(n)/n}$. The rate ϵ_n is often equal to $\tilde{\epsilon}_n$ up to log n, the use of these two different rates is usual and allows more flexibility. Then the rate ϵ_n is only determined by the non parametric part of the model, i.e. $\Pi_f^{(k)}$ and f^* , as described above.

Following the previous explanation, when Π_Q , $\Pi_f^{(k)}$, f^* and Q^* are fixed, ϵ_n is specified by Assumption [\(A\)](#page-5-2) and [\(B\)](#page-5-3). This rate ϵ_n is deteriorated via \underline{q}_n which is set through Assumption [\(D\)](#page-6-2). The larger $\tilde{\epsilon}_n$ is, that is to say the more difficult the estimation of the non parametric part $(f^* \text{ with } \Pi_f^{(k)})$ is, the more stringent Assumption [\(D\)](#page-6-2) is. To avoid too small \underline{q}_n which leads to deteriorated posterior convergence rate $\epsilon_n/\underline{q}_n$, one may choose a prior Π_Q which is supported on $\Delta_k^k(q)$ for some $0 < q \leq q^*$. More examples of distribution Π _Ω are given in Section [4.](#page-9-0)

In the following section, we give the proof of Theorem [3.1.](#page-6-0)

3.2 Proof of Theorem [3.1](#page-6-0)

To obtain posterior concentration rates in the framework of HMMs with finite state space, we use the technique of proof of [Ghosal and van der Vaart \[2007b\]](#page-34-12). The key tools of this technique are a control of the prior mass on log-likelihood neighborhoods of θ^* and the existence of certain tests. We use the tests built in [Gassiat and Rousseau \[2014\]](#page-34-9). The main difficulty of the proof arises from the control of log-likelihood neighborhoods. These neighbourhoods are controlled thanks to Lemmas [3.2](#page-7-1) and [3.3.](#page-8-0) The proof of these lemmas are based on refinements of results of [Douc and Matias \[2001\]](#page-34-13) and [Douc et al. \[2004\]](#page-34-14).

In Lemma [3.2,](#page-7-1) we control $KL(p_n^{\theta^*})$ $\binom{\theta^*}{n}, p_n^{\theta}$. Its proof is given in Section [A.](#page-13-1)

Lemma 3.2. Let $0 < \tilde{\epsilon}_n$ be small enough. Assume that $\theta = (Q, f) \in \Theta$, is such that Assumptions [\(A.1\)](#page-5-4)–[\(A.6\)](#page-5-5) hold with $u_n = 1$ for all $n \in \mathbb{N}$ and

$$
||Q - Q^*|| \le \tilde{\epsilon}_n. \tag{4}
$$

Then there exists $N > 0$ such that for all $n \geq N$,

$$
\mathbb{E}^{\theta^*}(L_n^{\theta^*} - L_n^{\theta}) = KL(p_n^{\theta^*}, p_n^{\theta}) \le C_K n \tilde{\epsilon}_n^2,
$$

where C_K is defined in Assumption [\(C\)](#page-6-1).

As can be seen in the proof, Assumption [\(A.1\)](#page-5-4) can be replaced in Lemma [3.2](#page-7-1) by the following weaker assumption:

$$
\int_{S^c} f_i^*(y) \max_{1 \le j \le k} \log \left(\frac{f_j^*(y)}{f_j(y)} \right) \lambda(dy) \le C_K \tilde{\epsilon}_n^2, \text{ for all } 1 \le i \le k \tag{5}
$$

which is implied by Assumptions $(A.1)$ and $(A.4)$. Note however that Assumption $(A.1)$ is used in the control of the variance $Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta})$ in Lemma [3.3.](#page-8-0)

Lemma [3.3](#page-8-0) gives a control of $Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta})$. It is proved in Appendix [B.](#page-16-0)

Lemma 3.3. Let $0 < \epsilon_n$ be small enough and u_n be defined by Equation [\(2\)](#page-5-9) Assume that $\theta = (Q, f) \in \Theta$, is such that Assumptions [\(A.1\)](#page-5-4), [\(A.2\)](#page-5-10) and [\(A.4\)](#page-5-8) hold and

$$
||Q - Q^*|| \le \frac{\tilde{\epsilon}_n}{\sqrt{u_n}}.\tag{6}
$$

Then there exists a positive constant C_{KL^2} such that for all $\alpha \in (0,1)$ and $n \in \mathbb{N}$

$$
Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta}) = \mathbb{E}^{\theta^*} \left[\left(\log \frac{p^{\theta^*}(Y_{1:n})}{p^{\theta}(Y_{1:n})} - \mathbb{E}^{\theta^*} \left(\log \frac{p^{\theta^*}(Y_{1:n})}{p^{\theta}(Y_{1:n})} \right) \right)^2 \right]
$$

$$
\leq C_{KL^2} \frac{n}{\alpha} \left(\frac{\tilde{\epsilon}_n}{\sqrt{u_n}} \right)^{2-\alpha}.
$$

We now give the proof of Theorem [3.1.](#page-6-0)

Proof of Theorem [3.1.](#page-6-0) This proof follows the lines of the proof of Theorem 1 of [Ghosal](#page-34-12) [and van der Vaart \[2007b\]](#page-34-12) with two variants. These differences come from the tests (see Equations [\(7\)](#page-8-1) and [\(8\)](#page-8-2)) and the control of the Kullback-Leibler neighborhoods (Equation (11) .

Using the tests built in the proof of Theorem 4 in [Gassiat and Rousseau \[2014\]](#page-34-9), for all $M > 0$, there exists $\psi_n \in [0, 1]$ such that

$$
\mathbb{E}^{\theta^*}(\psi_n) \le N\left(\frac{\epsilon_n}{12}, \Delta_k^k(\underline{q}_n) \times \mathcal{F}_n, D_\ell\right) \exp\left(\frac{-n\epsilon_n^2 \underline{q}^{*2} k^4 M^2}{128l}\right) \tag{7}
$$

and

$$
\sup_{\substack{\theta \in \Delta_k^k(\underline{q}_n) \times \mathcal{F}_n \\ D_\ell(\theta, \theta^*) \ge M \epsilon_n / \underline{q}_n}} \mathbb{E}^{\theta} (1 - \psi_n) \le \exp\left(-\frac{n\epsilon_n^2 k^2 M^2}{128l}\right).
$$
\n
$$
(8)
$$

Since

$$
D_{\ell}(\theta, \tilde{\theta}) \le \sum_{1 \le i \le k} |\mu_i^Q - \mu_i^{\tilde{Q}}| + k(\ell - 1) \max_{1 \le i, j \le k} |Q_{i,j} - \tilde{Q}_{i,j}| + \ell d(f, \tilde{f}),
$$

we obtain

$$
N\left(\frac{\epsilon_n}{12}, \Delta_k^k(\underline{q}_n) \times \mathcal{F}_n, D_\ell\right) \le \left(\frac{24\ell k(k-1)}{\epsilon_n}\right)^{k(l-1)} N\left(\frac{\epsilon_n}{24\ell}, \mathcal{F}_n, d\right) \tag{9}
$$

which leads to

$$
\mathbb{E}^{\theta^*}(\psi_n) \lesssim \exp(-n\epsilon_n^2 (M^2 \tilde{C} - C')) \tag{10}
$$

for some constant \tilde{C} , using Assumption [\(B\)](#page-5-3). We replace Equation (8.4) of [Ghosal and](#page-34-12) [van der Vaart \[2007b\]](#page-34-12) by Equation [\(10\)](#page-8-4). Equation (8.5) of [Ghosal and van der Vaart](#page-34-12) [\[2007b\]](#page-34-12) is replaced by Equation [\(8\)](#page-8-2).

Let α_n be a sequence tending to 0, to be specified later. We define

 $\mathcal{B}_n(\theta^*)$

$$
:= \left\{ \theta : KL(p_n^{\theta^*}, p_n^{\theta}) \le C_K n \tilde{\epsilon}_n^2, \ Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta}) \le \frac{C_{KL^2} n}{\alpha_n} \left(\frac{\tilde{\epsilon}_n}{\sqrt{u_n}} \right)^{2-\alpha_n} \right\} \tag{11}
$$

in the place of $B_n(\theta^*, \bar{\epsilon}_n, 2)$ in the notation of [Ghosal and van der Vaart \[2007b\]](#page-34-12), setting $\bar{\epsilon}_n = \sqrt{C_K} \tilde{\epsilon}_n$. Using Assumptions [\(A\)](#page-5-2) and [\(C\)](#page-6-1), and Lemmas [3.2](#page-7-1) and [3.3](#page-8-0)

$$
\Pi\left(\mathcal{B}_n\left(\theta^*\right)\right) \gtrsim \exp(-(C_Q + C_f)n\tilde{\epsilon}_n^2) \gtrsim \exp(-Cn\tilde{\epsilon}_n^2).
$$

By choosing

- (i) $\alpha_n = 1/\log(n)$, for all $n \in \mathbb{N}$; if $\tilde{\epsilon}_n \gtrsim n^{-s}$ for some $s < 1/2$,
- (ii) $\alpha_n = \log(\log(n))/\log(n)$, for all $n \in \mathbb{N}$; otherwise,

and following the lines of the proof of Lemma 10 of [Ghosal and van der Vaart \[2007b\]](#page-34-12),

$$
P^{\theta^*}(\mathcal{D}_n < \Pi(B_n(\theta^*)) \exp(-2C_K n \tilde{\epsilon}_n^2)) = O\left(\frac{n(\tilde{\epsilon}_n/\sqrt{u_n})^{2-\alpha_n}}{\alpha_n (n \tilde{\epsilon}_n^2)^2}\right) = o(1),\tag{12}
$$

with $\mathcal{D}_n = \int_{\mathcal{B}_n(\theta^*)} p_n^{\theta}(Y_{1:n})/p_n^{\theta^*}$ $\int_{n}^{\theta^*}(Y_{1:n})\Pi(d\theta)$. Following the lines of the proof of Theorem 1 of [Ghosal and van der Vaart \[2007b\]](#page-34-12) with the above modifications, we obtain,

$$
\mathbb{E}^{\theta^*} \left(\Pi \left(\theta \in \Delta_k^k(\underline{q}_n) \times \mathcal{F}_n : D_\ell(\theta, \theta^*) \ge M \frac{\epsilon_n}{\underline{q}_n} | Y_{1:n} \right) \right) = o(1) \tag{13}
$$

for M large enough.

This concludes the proof since

$$
\mathbb{E}^{\theta^*} \left(\Pi((\Delta_k^k(\underline{q}_n) \times \mathcal{F}_n)^c | Y_{1:n}) \right) = o(1)
$$

using Equation [\(12\)](#page-9-1) and Lemma 1 of [Ghosal and van der Vaart \[2007b\]](#page-34-12) with

$$
\Pi((\Delta_k^k(\underline{q}_n) \times \mathcal{F}_n)^c) = o(\exp(-2n\bar{\epsilon}_n^2)\Pi(\mathcal{B}_n(\theta^*))
$$

as soon as $C > 2C_K + C_Q + C_f$ (using Assumptions [\(B\)](#page-5-3) and [\(D\)](#page-6-2)).

4 Applications

In this section, we apply Theorem [3.1](#page-6-0) to different priors and different classes of emission density functions. In all examples treated in Section [4,](#page-9-0) the prior on emission distributions is chosen to be a product of a distribution Π_f on \mathcal{F} :

$$
\Pi_f^{(k)} = (\Pi_f)^{\otimes k}.\tag{14}
$$

However, Theorem [3.1](#page-6-0) can also be applied to other priors such as priors restricted to translated emission density functions, the translation HMM is described in Equation [\(16\)](#page-12-1).

In Section [4.1,](#page-10-0) we consider discrete observations, i.e. $\mathcal{Y} = \mathbb{N}$. We assume that the prior Π_f on each emission distributions is a Dirichlet process. We compute the rate ϵ_n obtained with this prior when the true emission distributions have an exponential decay. In Section [4.2,](#page-12-0) the observations are assumed to live in $\mathbb R$ and the emission distributions are supposed to be absolutely continuous with respect to the Lebesgue measure. We consider a Dirichlet process mixture of Gaussian distributions as a prior Π_f on each emission density functions. We compute the rate ϵ_n obtained with this prior when the emission density functions belong to functional classes of β -Hölder types.

We always assume that

(Q0) Π_Q is absolutely continuous with respect to the Lebesgue measure on Δ_k^k with density π_Q , $\pi_Q(Q^*) > 0$ and $\pi_Q(Q) = \pi_q(Q_{1,\cdot}) \dots \pi_q(Q_{k,\cdot})$, for all $Q \in \Delta_k^k$ and where Q_i , denotes the *i*-th row of Q .

In Sections [4.1](#page-10-0) and [4.2,](#page-12-0) we consider three different priors Π_{Q} on transition matrices which corresponds to three different decays of π_Q near the boundary of Δ_k^k :

 \Box

(Q1) exponential tail:

$$
\pi_q(u_1,\ldots,u_k) \lesssim \exp(-\alpha_1/u_1)\ldots \exp(-\alpha_k/u_k),
$$

for all $u \in \Delta_k$, for some positive constants α_i , $1 \leq i \leq k$,

(Q2) exponential of exponential tail:

$$
\pi_q(u_1,\ldots,u_k) \lesssim \exp(-\beta_1 \exp(u_1^{-\alpha_1})) \ldots \exp(-\beta_k \exp(u_k^{-\alpha_k})),
$$

for all $u \in \Delta_k$, for some positive constants α_i and β_i , $1 \leq i \leq k$,

(Q3) truncated distribution:

$$
\Pi_Q(\Delta_k^k(\underline{q})) = 1,
$$

for some positive q .

Note that Assumption $(Q3)$ implies Assumption $(Q2)$ which implies $(Q1)$. In [\[Gassiat](#page-34-9)] [and Rousseau, 2014\]](#page-34-9), together with priors of type [\(Q1\),](#page-10-3) more general priors are also considered, since they assume

$$
\pi_q(u_1,\ldots,u_k) \lesssim u_1^{\alpha_1-1}\ldots u_k^{\alpha_k-1},
$$

for all $u \in \Delta_k$, for some positive constants α_i , $1 \leq i \leq k$. [Gassiat and Rousseau \[2014\]](#page-34-9) show that q_n , in Assumption [\(D\)](#page-6-2) and Theorem [3.1,](#page-6-0) is equal to a power of $1/n$ when the emission distributions belong to a parametric family. We do not consider this type of priors since they lead to deteriorated rates $\epsilon_n/\underline{q}_n$. Under [\(Q1\),](#page-10-3) [\[Gassiat and Rousseau,](#page-34-9) [2014\]](#page-34-9) obtain q_n equal to a power of $1/(\log n)$ when the emission distributions belong to a parametric family. We obtain the same rate q_n in the case of discrete observations and emission distributions with exponential decay (more generally, it would be the case as soon as $\epsilon_n = n^{-1/2} \log(n)^t$ for some positive t). However, in the case of emission distributions absolutely continuous with respect to Lebesgue measure (Section [4.2\)](#page-12-0), Assumption [\(D\)](#page-6-2) leads to a rate q_n at least polynomial in $1/n$ with priors satisfying [\(Q1\).](#page-10-3) While priors verifying [\(Q2\)](#page-10-2) or [\(Q3\)](#page-10-1) lead to a rate q_n equal to a power of $1/\log n$ as soon as ϵ_n is a power of n; and thus do not deteriorate the posterior concentration rate (up to $log(n)$). Note that Assumptions [\(Q0\)](#page-9-2) and [\(Q3\)](#page-10-1) are compatible if and only if $q \leq q^*$. Thus, the use of a prior verifying [\(Q3\)](#page-10-1) requires a knowledge of a lower bound q^* of $\min_{1 \leq i,j \leq k} Q^*_{i,j}$.

In [\[Vernet et al., 2015\]](#page-35-8), posterior consistency is derived under Assumption [\(Q3\)](#page-10-1) while weaker conditions on f^* and $\Pi_f^{(k)}$ (see Assumptions (A0), (A1) and (A2) in [\[Vernet et al.,](#page-35-8) 2015) compared to (A) , (B) and (C) . Here, we manage to obtain posterior concentration rates under Assumptions [\(Q1\)](#page-10-3) and [\(Q2\)](#page-10-2) which are weaker than Assumption [\(Q3\)](#page-10-1) because stronger conditions on $\Pi_f^{(k)}$ and f^* are assumed.

4.1 Discrete observations

In this section, we apply Theorem [3.1](#page-6-0) to the case of discrete observations so that λ is the count measure on N. HMMs with discrete observations are used in different applications, as in [\[Borchers et al., 2013\]](#page-33-5) for animal abundance estimation, in [\[Gassiat et al., 2015\]](#page-34-2)for gene expression identification, or in [\[Linderman et al., 2014\]](#page-35-12) for neural representation of spatial navigation, to cite a few.

In the framework of discrete distribution estimation with i.i.d. observations, [\[Han](#page-34-15) [et al., 2014\]](#page-34-15) have proved that no rates can be obtained with the L_1 loss without constraint on the considered distributions. Moreover, they obtain a minimax rate proportional to $1/\log n$ over the set $\{f \in \mathcal{F}: \sum_{i\in \mathbb{N}} -f(i)\log(f(i)) \leq C\}$. Rates of convergence are more widely studied in the case of the L_2 norm, for instance with monotony constraint in [\[Jankowski and Wellner, 2009\]](#page-35-13), with log-concave constraint in [\[Balabdaoui et al., 2013\]](#page-33-6), with convex constraint in [\[Durot et al., 2013\]](#page-34-16) and with envelope constraint in [\[Boucheron](#page-33-7) [et al., 2009\]](#page-33-7).

In the non parametric Bayesian framework, the Dirichlet process is a very popular prior. Here, we consider a Dirichlet process $DP(G)$ on the emission distributions, with G some finite positive measure on N:

$$
\Pi_f = DP(G).
$$

[\[Canale and Dunson, 2011\]](#page-33-8) propose other priors for discrete observations based on discretization of continuous mixtures of kernels and gives an overview of the priors used in the case of discrete observations.

We first verify Assumptions [\(A\)](#page-5-2), [\(B\)](#page-5-3) and [\(C\)](#page-6-1). As it can be seen from the proof (see Lemma [C.2](#page-26-0) in Appendix [C\)](#page-24-0), we need a heavy tail condition on G :

(P) there exists positive constants $a \leq A$ and $\alpha \geq 2$ such that for all $1 \leq j \leq k$ and for all $l \in \mathbb{N}$

$$
al^{-\alpha} \le G(l) \le Al^{-\alpha}
$$

.

Here, we consider the following class of discrete distributions which is based on an envelope constraint:

$$
\mathcal{D}(m, c, K) = \left\{ f \in \mathcal{F} : f(l) \le d \exp(-cl^m), \sum_{l \le N} \frac{-\log(f(l))}{l} \lesssim N^K, \right\}
$$
for all N large enough $\left\}, (15)$

where c, K and m are positive constants. We also consider the following assumption linking the tails of the true emission distributions:

(I) there exists $\delta > 0$ such that for all N large enough and all $1 \leq i, j \leq k$,

$$
\sum_{l\geq N} f_i^*(l) \log^2 \left(f_j^*(l) \right) \lesssim \exp(-N^m(c-\delta)).
$$

Under these assumptions, we obtain the following rates $\tilde{\epsilon}_n$ and ϵ_n .

Theorem 4.1. Assume there exist positive constants c, K and m such that for all $1 \leq$ $j \leq k$, $f_j^* \in \mathcal{D}(m, c, K)$ and that Assumptions [\(Q0\),](#page-9-2) [\(I\)](#page-11-2) and [\(P\)](#page-11-3) hold.

Then Assumptions (A) , (B) and (C) hold with

$$
\tilde{\epsilon}_n = \frac{1}{\sqrt{n}} (\log n)^{t_0}
$$
 and $\epsilon_n = \frac{1}{\sqrt{n}} (\log n)^t$,

where $t > 4t_0$ and $t_0 \geq 1/2 \max(1/m + 1, K/m)$.

Theorem [4.1](#page-11-0) leads to the following posterior concentration rates $(\epsilon_n/\underline{q}_n)$ which are minimax (up to $\log n$):

Corollary 4.2. Assume there exist positive constants c, K and m such that for all $1 \leq$ $j \leq k$, $f_j^* \in \mathcal{D}(m, c, K)$ and that Assumptions [\(Q0\),](#page-9-2) [\(I\)](#page-11-2) and [\(P\)](#page-11-3) hold. Moreover suppose that $\Pi_{\mathcal{Q}}$ satisfies

- [\(Q1\),](#page-10-3) then the posterior concentrates with rate $\frac{1}{\sqrt{2}}$ $\frac{1}{n}(\log n)^{t+2t_0}$;
- [\(Q2\),](#page-10-2) then the posterior concentrates with rate $\frac{1}{\sqrt{2}}$ $\frac{1}{\overline{n}}(\log n)^t;$
- [\(Q3\),](#page-10-1) then the posterior concentrates with rate $\frac{1}{\sqrt{2}}$ $\frac{1}{\overline{n}}(\log n)^t;$

with $t > 4t_0$ and $t_0 \geq 1/2 \max(1/m + 1, K/m)$.

4.2 Dirichlet process mixtures of Gaussian distributions–Adaptivity to Hölder function classes

Dirichlet process mixtures of Gaussian distributions are commonly used to model densities on \mathbb{R} or \mathbb{R}^d . In particular, there exist efficient algorithms to sample from the posterior distribution in the i.i.d. framework. In the translation HMM:

$$
Y_t = m_{X_t} + \epsilon_t,\tag{16}
$$

where $\epsilon_t \stackrel{\text{i.i.d.}}{\sim} g\lambda$, $m_j \in \mathbb{R}$ and X_t is a Markov chain with translation matrix Q ; [Yau et al.](#page-35-4) [\[2011\]](#page-35-4) use a Dirichlet process mixtures of Gaussian distributions on g. In the context of i.i.d. observations posterior concentration rates have been derived with such prior models in [Ghosal and van der Vaart \[2007a\]](#page-34-17), [Kruijer et al. \[2010\]](#page-35-14) and [Shen et al. \[2013\]](#page-35-15). In the framework of HMMs, we propose to apply Theorem [3.1](#page-6-0) when Π_f is a Dirichlet process mixture of Gaussian distributions.

We assume that the reference measure λ is the Lebesgue measure on R. We also assume that the prior on \mathcal{F}^k is a product of Dirichlet process mixture of Gaussian distributions:

$$
Y_t | X_t = j \sim f_j, \quad f_j = \int \phi_\sigma(\cdot - \mu) dP_j(\mu),
$$

$$
P_j \stackrel{\text{i.i.d.}}{\sim} DP(G), \text{ for all } 1 \le j \le k, \quad \sigma \sim \pi_\sigma \lambda,
$$

where ϕ_{σ} is the Gaussian density function with variance σ^2 and mean zero, $DP(G)$ is the Dirichlet process with finite positive base measure G and π_{σ} is a distribution on R.

We define the same functional classes as in [Kruijer et al. \[2010\]](#page-35-14):

$$
\mathcal{P}(\beta, L, \gamma) := \left\{ f \in \mathcal{F} : \log f \text{ is locally } \beta \text{-H\"older with derivatives } l_j = (\log f)^{(j)} \right\}
$$

and $|l_{\lfloor \beta \rfloor}(y) - l_{\lfloor \beta \rfloor}(x)| \le r!L(y)|y - x|^{\beta - \lfloor \beta \rfloor}, \text{ as soon as } |x - y| \le \gamma \right\}$ (17)

where $\beta > 0$, L is a polynomial function, $\gamma > 0$ and $|\beta|$ is the largest integer smaller than $β$. We also consider the following tail conditions:

(T1) there exist positive constants M_0 , τ_0 , γ_0 such that for all $1 \leq i \leq k$ and all $y \in \mathbb{R}$

 $f_i^*(y) \leq M_0 \exp(-\tau_0 |y|^{\gamma_0}),$

(T2) for all $1 \le i, j \le k$ there exist constants $T_{i,j}$, $M_{i,j}$, $\tau_{i,j}$, $\gamma_{i,j} < \gamma_0$ such that

$$
f_i^*(y) \le f_j^*(y) M_{i,j} \exp(\tau_{i,j}|y|^{\gamma_{i,j}}), \quad |y| \ge T_{i,j},
$$

(T3) for all $1 \leq i \leq k$, f_i^* is positive and there exist $c_i > 0$, $y_i^m < y_i^M$ such that f_i^* is non decreasing on $(-\infty, y_i^m)$, non increasing on $(y_i^M, +\infty)$ and $f_i^*(y) \geq c_i$ for $y \in (y_i^m, y_i^M)$.

Assumptions [\(T1\)](#page-12-2) and [\(T3\)](#page-12-3) are the same tail assumptions as those used in [Kruijer et al.](#page-35-14) [\[2010\]](#page-35-14). The new Assumption [\(T2\)](#page-12-4) links the tail of each emission distributions.

We now describe the assumptions concerning the prior on the emission distributions:

- (G1) $G([-y, y]^c) \lesssim \exp(-C_1 y^{a_1})$ for all sufficiently large $y > 0$, for some positive constant a_1 .
- (S1) Π_{σ} ($\sigma \leq x$) \lesssim exp($-C_2x^{-a_2}$) for all sufficiently small $x > 0$, for some positive constant a_2 ,

(S2) $\Pi_{\sigma}(\sigma > x) \lesssim x^{-a_3}$ for all sufficiently large $x > 0$, for some positive constant a_3 ,

(S3) there exists $a_6 \leq 1$ such that

$$
\Pi_{\sigma} (x \le \sigma \le x(1+s)) \gtrsim x^{-a_4} s^{a_5} \exp(-C_3 x^{-a_6})
$$

for all $s \in (0,1)$, sufficiently small $x > 0$, for some positive constants a_4 , a_5 and a_6 .

The gamma and Gaussian distributions satisfy Assumption [\(G1\).](#page-12-5) The inverse gamma distribution verifies [\(S1\),](#page-12-6) [\(S2\)](#page-13-4) and [\(S3\).](#page-13-5)

Theorem 4.3. Assume that there exist β , L and γ such that for all $1 \leq j \leq k$, $f_j^* \in$ $\mathcal{P}(\beta, L, \gamma)$ and Assumptions [\(Q0\),](#page-9-2) [\(T1\)](#page-12-2)[–\(T3\),](#page-12-3) [\(G1\)](#page-12-5) and [\(S1\)](#page-12-6)[–\(S3\)](#page-13-5) hold. Then Assumptions (A) , (B) and (C) hold with

 $\tilde{\epsilon}_n = n^{-\frac{\beta}{2\beta+1}}\log(n)^{t_0}$ and $\epsilon_n = n^{-\frac{\beta}{2\beta+1}}\log(n)^t$

 (18)

where $t > t_0 \geq (2 + 2/\gamma_0 + 1/\beta)/(1/\beta + 2)$.

The proof of Theorem [4.3](#page-13-2) is given in Appendix [D.](#page-30-0) Using Theorem [3.1](#page-6-0) and [4.3,](#page-13-2) we directly deduce posterior rate of convergence under the Assumptions of Theorem [4.3](#page-13-2) and the different types of priors Π_{Q} .

Corollary 4.4. Assume that there exist β , L and γ such that for all $1 \leq j \leq k$, $f_j^* \in$ $\mathcal{P}(\beta, L, \gamma)$ and that Assumptions [\(Q0\),](#page-9-2) [\(T1\)](#page-12-2)[–\(T3\),](#page-12-3) [\(G1\)](#page-12-5) and [\(S1\)](#page-12-6)[–\(S3\)](#page-13-5) hold. Moreover suppose that Π_{Q} satisfies

- [\(Q1\),](#page-10-3) then the posterior concentrates with rate $n^{\frac{-\beta+1}{2\beta+1}}(\log n)^{3t}$;
- [\(Q2\),](#page-10-2) then the posterior concentrates with rate $n^{-\frac{\beta}{2\beta+1}}(\log n)^{t+1/(\max_{1\leq i\leq k}\alpha_i)}$;
- [\(Q3\),](#page-10-1) then the posterior concentrates with rate $n^{-\frac{\beta}{2\beta+1}}(\log n)^t$;

with $t > (2 + 2/\gamma_0 + 1/\beta)/(1/\beta + 2)$.

The minimax rate, with respect to D_{ℓ} in the HMM framework for emission density functions belonging to functional classes of β -Hölder type, is larger than $n^{-\beta/(2\beta+1)}$. Indeed with a hidden Markov chain (X_t, Y_t) distributed from a parameter $\theta = (Q, f)$ such that $Q_{i,j} = 1/k$ and $f_i = f_1$ for all $1 \leq i, j \leq k$, the observations (Y_t) are i.i.d. from $f_1 \lambda$. Thus, priors satisfying $(Q0)$, $(T1)$ – $(T3)$, $(G1)$, $(S1)$ – $(S3)$ and $(Q2)$ lead to minimax rates (up to $log n$). As these priors do not depend on the regularity of the functional class considered, they ensure adaptive Bayesian density estimation in the framework of HMMs.

Acknowledgements

I want to thank Elisabeth Gassiat and Judith Rousseau for their valuable comments. This work was partly supported by the grants ANR Banhdits and Calibration.

Appendices

A Proof of Lemma [3.2](#page-7-1) : control of the Kullback Leibler divergence between θ^* and θ

First denote $\epsilon = \epsilon_n$. Using Assumptions [\(1\)](#page-4-0) and [\(4\)](#page-7-2), there exists $q > 0$ such that for all $1 \leq i, j \leq k, Q^*_{i,j} \geq \underline{q}$ and $Q_{i,j} \geq \underline{q}$, more precisely, \underline{q} can be chosen equal to $\underline{q}^*/2$ as soon

as *n* is large enough. Let $q_t^{\theta, Y_{1:t-1}}$ be the conditional density function of Y_t given $Y_{1:t-1}$ with respect to λ :

$$
q_t^{\theta, Y_{1:t-1}} = \sum_{i=1}^k f_i(\cdot) Q_{t,i}^{\theta, Y_{1:t-1}},
$$

where $Q_{t,i}^{\mu,\theta,Y_{1:t-1}} = \mathbb{P}^{\theta}(X_t = i|Y_{1:t-1}, X_1 \sim \mu)$, where $t \geq 1, 1 \leq i \leq k, \mu \in \Delta_k$. When μ is not specified $(Q_{t,i}^{\theta,Y_{1:t-1}})$, the stationary initial probability distribution is considered. Note that, when $\mu \in \Delta_k(\underline{q}), \, \tilde{\theta} \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$,

$$
Q_{t,i}^{\mu,\tilde{\theta},Y_{1:t-1}} = \frac{\sum_{j=1}^{k} Q_{t-1,j}^{\mu,\tilde{\theta},Y_{1:t-2}} Q_{j,i} f_j(Y_{t-1})}{\sum_{t=1}^{k} Q_{t-1,t}^{\mu,\tilde{\theta},Y_{1:t-2}} f_t(Y_{t-1})} \ge \underline{q},\tag{19}
$$

for all $1 \leq i \leq k, t \geq 1$.

$$
KL(p_n^{\theta^*}, p_n^{\theta})
$$

= $\mathbb{E}^{\theta^*} \left(\sum_{t=1}^n \int q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta, Y_{1:t-1}}(y)} \right) \lambda(dy) \right)$
= $\mathbb{E}^{\theta^*} \left(\sum_{t=1}^n \int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta, Y_{1:t-1}}(y)} \right) \lambda(dy) \right)$
+ $\mathbb{E}^{\theta^*} \left(\sum_{t=1}^n \int_{S^c} q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta^*, Y_{1:t-1}}(y)} \right) \lambda(dy) \right).$ (20)

Using Equation [\(19\)](#page-14-2), for all $1 \leq l \leq k$,

$$
\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta, Y_{1:t-1}}(y)} = \frac{\sum_{i=1}^k f_i^*(y) Q_{t,i}^{\theta^*, Y_{1:t-1}}}{\sum_{i=1}^k f_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}} \le \frac{k \max_{1 \le j \le k} f_j^*(y)}{\underline{q} f_l(y)}
$$
\n
$$
\le \max_{1 \le j \le k} \frac{k f_j^*(y)}{\underline{q} f_j(y)} \tag{21}
$$

then Assumptions [\(5\)](#page-7-3) and [\(A.4\)](#page-5-8) lead to

$$
\mathbb{E}^{\theta^*} \left(\int_{S^c} q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta, Y_{1:t-1}}(y)} \right) \lambda(dy) \right) \leq \left(\log \frac{k}{\underline{q}} + 1 \right) \epsilon^2. \tag{22}
$$

We now control the expectancy of the third line of Equation [\(20\)](#page-14-0)

$$
\mathbb{E}^{\theta^*}\left(\sum_{t=1}^n \int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{q_t^{\theta, Y_{1:t-1}}(y)}\right) \lambda(dy)\right) \n\leq \mathbb{E}^{\theta^*}\left(\sum_{t=1}^n \int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}\right) \lambda(dy)\right) \n+ \mathbb{E}^{\theta^*}\left(\sum_{t=1}^n \int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}{q_t^{\theta, Y_{1:t-1}}(y)}\right) \lambda(dy)\right).
$$
\n(23)

We control the expectancy of the third line of Equation [\(23\)](#page-14-3), using

$$
\frac{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}{q_t^{\theta, Y_{1:t-1}}(y)} = \frac{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}{\sum_{i=1}^k f_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}} \le \max_{1 \le i \le k} \frac{\tilde{f}_i(y)}{f_i(y)};
$$

by Lemma [A.1,](#page-15-0) and Assumption [\(A.5\)](#page-5-7), to obtain

$$
\mathbb{E}^{\theta^*} \left(\int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}{q_t^{\theta, Y_{1:t-1}}(y)} \right) \lambda(dy) \right) \le \epsilon^2.
$$
 (24)

We bound the expectancy of the second line of Equation [\(23\)](#page-14-3), using the inequality recalled at the top of page 1234 of [\[Kruijer et al., 2010\]](#page-35-14),

$$
\mathbb{E}^{\theta^*} \left(\int_S q_t^{\theta^*, Y_{1:t-1}}(y) \log \left(\frac{q_t^{\theta^*, Y_{1:t-1}}(y)}{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}} \right) \lambda(dy) \right) \n\leq \mathbb{E}^{\theta^*} \left(\int_S \frac{\left(q_t^{\theta^*, Y_{1:t-1}}(y) - \sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}} \right)^2}{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}} \lambda(dy) \right) \n+ \mathbb{E}^{\theta^*} \left(\int_{S^c} \sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}} - q_t^{\theta^*, Y_{1:t-1}}(y) \lambda(dy) \right).
$$
\n(25)

The expectancy of the second line of Equation [\(25\)](#page-15-1) is controlled as follows

$$
\mathbb{E}^{\theta^*}\left(\int_{S} \frac{\left(q_t^{\theta^*, Y_{1:t-1}}(y) - \sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}\right)^2}{\sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}}}\lambda(dy)\right) \n\leq 2 \mathbb{E}^{\theta^*}\left(\frac{k \max_{1 \leq i \leq k} \left(Q_{t,i}^{\theta^*, Y_{1:t-1}} - Q_{t,i}^{\theta, Y_{1:t-1}}\right)^2}{\frac{q}{2}}\right) + 2 \mathbb{E}^{\theta^*}\left(\int_{S} \max_{1 \leq i \leq k} \frac{\left(f_i^*(y) - \tilde{f}_i(y)\right)^2}{\frac{q \tilde{f}_i(y)}{\theta}}\lambda(dy)\right) \n\leq \left(\frac{16k(1+2k)}{\frac{q^4}{4}} + 1\right) \frac{2}{\frac{q}{2}} \epsilon^2 + \frac{16k\rho^{2(t-1)}}{\frac{q}{2}}\n\right)
$$
\n(26)

using Lemma [B.2](#page-18-0) and then Assumption [\(A.2\)](#page-5-10) and Lemma [B.2.](#page-18-0) The expectancy of the third line of Equation [\(25\)](#page-15-1) is controlled thanks to Assumption [\(A.3\)](#page-5-6):

$$
\mathbb{E}^{\theta^*} \left(\int_{S^c} \sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}} - q_t^{\theta^*, Y_{1:t-1}}(y) \lambda(dy) \right)
$$
\n
$$
\leq \mathbb{E}^{\theta^*} \left(\int_{S^c} \sum_{i=1}^k \tilde{f}_i(y) Q_{t,i}^{\theta, Y_{1:t-1}} \lambda(dy) \right)
$$
\n
$$
\leq \max_{1 \leq i \leq k} \int_{S^c} \tilde{f}_i(y) \lambda(dy) \leq \epsilon^2.
$$
\n(27)

We conclude the proof by combining Equations [\(20\)](#page-14-0), [\(22\)](#page-14-4), [\(23\)](#page-14-3), [\(24\)](#page-15-2), [\(25\)](#page-15-1), [\(26\)](#page-15-3) and [\(27\)](#page-15-4). \Box

Lemma A.1. For all $a_i, b_i, c_i, d_i \geq 0, 1 \leq i \leq k$,

$$
\frac{\sum_{1 \le i \le k} a_i b_i}{\sum_{1 \le j \le k} c_j d_j} = \frac{\sum_{1 \le i \le k} a_i / c_i * b_i / d_i * c_i d_i}{\sum_{1 \le j \le k} c_j d_j} \le \max_{1 \le i \le k} \frac{a_i}{c_i} \max_{1 \le j \le k} \frac{b_j}{d_j}.
$$

B Proof of Lemma [3.3](#page-8-0) with technical lemmas: control of $Var^{\theta^*}(L_n^{\theta^*} - L_n^{\theta^*})$ $\binom{\theta}{n}$

B.1 Proof of Lemma [3.3](#page-8-0)

First denote $\epsilon = \tilde{\epsilon}_n / \sqrt{u_n}$. Using Assumptions [\(1\)](#page-4-0) and [\(6\)](#page-8-5), there exists $q > 0$ such that for all $1 \leq i, j \leq k$, $Q_{i,j}^* \geq \underline{q}$ and $Q_{i,j} \geq \underline{q}$, more precisely, \underline{q} can be chosen equal to $\underline{q}^*/2$ as soon as *n* is large enough. Let *Var* be the variance of $L_n^{\overline{\theta}^*} - L_n^{\theta}$:

$$
Var := \mathbb{E}^{\theta^*} \left[\left(\log \frac{p_n^{\theta^*}(Y_{1:n})}{p_n^{\theta}(Y_{1:n})} - \mathbb{E}^{\theta^*} \left(\log \frac{p_n^{\theta^*}(Y_{1:n})}{p_n^{\theta}(Y_{1:n})} \right) \right)^2 \right].
$$

Denoting $Z_t = \log \left(\frac{P^{\theta^*}(Y_t|Y_{1:t-1})}{P^{\theta}(Y_t|Y_{1:t-1})} \right)$ $P^{\theta}(Y_t|Y_{1:t-1})$ $\big)$, then

$$
Var = \mathbb{E}^{\theta^*} \left(\left(\sum_{t=1}^n Z_t - \mathbb{E}^{\theta^*} \left(\sum_{t=1}^n Z_t \right) \right)^2 \right).
$$

We want to bound Var by $Cn\epsilon^{(2-\alpha)/2}$, for any $\alpha > 0$. In this purpose, we split the sum in two parts:

$$
Var \leq 2 \mathbb{E}^{\theta^*} \left[\left(\sum_{t=1}^n \left(Z_t - \mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) \right) \right)^2 \right] + 2 \mathbb{E}^{\theta^*} \left[\left(\sum_{t=1}^n \left(\mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) - \mathbb{E}^{\theta^*} (Z_t) \right) \right)^2 \right],
$$
\n
$$
= S_1 \tag{28}
$$

 S_1 is the expectancy of the square of a sum of martingale increments, for which the covariances are zero so that only n terms remain. S_2 is further controlled using the exponential forgetting of Markov chain. First, we control S_1 :

$$
S_1 = \sum_{t=1}^n \mathbb{E}^{\theta^*} \left(\left(Z_t - \mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) \right)^2 \right)
$$

+
$$
2 \sum_{1 \leq r < t \leq n} \mathbb{E}^{\theta^*} \left(\left(Z_r - \mathbb{E}^{\theta^*} \left(Z_r | Y_{1:r-1} \right) \right) \mathbb{E}^{\theta^*} \left(Z_t - \mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) | Y_{1:t-1} \right) \right) \tag{29}
$$

$$
\leq \sum_{t=1}^n \mathbb{E}^{\theta^*} \left(Z_t^2 \right)
$$

using the convexity of the square function and Jensen's inequality so that

$$
\mathbb{E}^{\theta^*}(\mathbb{E}^{\theta^*}(Z_t|Y_{1:t-1})^2) \leq \mathbb{E}^{\theta^*}(Z_t^2). \tag{30}
$$

As to S_2 :

$$
S_2 = \sum_{t=1}^n \mathbb{E}^{\theta^*} \left(\left(\mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) - \mathbb{E}^{\theta^*} \left(Z_t \right) \right)^2 \right) + 2 \sum_{1 \leq r < t \leq n} \mathbb{E}^{\theta^*} \left(\left(\mathbb{E}^{\theta^*} \left(Z_r | Y_{1:r-1} \right) - \mathbb{E}^{\theta^*} \left(Z_r \right) \right) \mathbb{E}^{\theta^*} \left(\mathbb{E}^{\theta^*} \left(Z_t | Y_{1:t-1} \right) - \mathbb{E}^{\theta^*} \left(Z_t \right) | Y_{1:r-1} \right) \right) \n\leq \sum_{t=1}^n \mathbb{E}^{\theta^*} \left(Z_t^2 \right) + 2 \sum_{1 \leq r < t \leq n} \sqrt{\mathbb{E}^{\theta^*} \left(Z_r^2 \right)} \sqrt{\mathbb{E}^{\theta^*} \left(\left| \mathbb{E}^{\theta^*} \left(Z_t | Y_{1:r-1} \right) - \mathbb{E}^{\theta^*} \left(Z_t \right) \right|^2 \right)},
$$
\n(31)

using Equation [\(30\)](#page-16-2) and Cauchy-Schwarz inequality to bound the second term.

Combining (28) , (29) et (31) , we obtain

$$
Var \le 4 \sum_{t=1}^{n} \mathbb{E}^{\theta^*} (Z_t^2) + 4 \sum_{1 \le r < t \le n} \sqrt{\mathbb{E}^{\theta^*} (Z_r^2)} \sqrt{\mathbb{E}^{\theta^*} (|\mathbb{E}^{\theta^*} (Z_t | Y_{1:r-1}) - \mathbb{E}^{\theta^*} (Z_t)|^2)}.
$$
 (32)

Then using Lemmas [B.1](#page-17-0) and [B.3,](#page-21-0)

$$
Var \le 4\left(\frac{16}{\underline{q}(1-\rho^2)} + Cn\epsilon^2\right) + 16\rho^{-\frac{5\alpha}{4}}\left(2\epsilon + \frac{10}{\underline{q}}\right)^{\alpha/2} \sum_{1 \le r < t \le n} \left(\frac{16\rho^{2(r-1)}}{\underline{q}^2} + \tilde{C}\epsilon^2\right)^{1-\alpha/4} \rho^{\frac{\alpha}{4}(t-r)}.\tag{33}
$$

Since,

$$
\sum_{1 \le r < t \le n} \left(\frac{16\rho^{2(r-1)}}{\underline{q}^2} + \tilde{C}\epsilon^2 \right)^{1-\alpha/4} \rho^{\frac{\alpha}{4}(t-r)} \le 2 \frac{\rho^{\alpha/4}}{1-\rho^{\alpha/4}} \left(\frac{16}{\underline{q}^2(1-\rho)} + n\tilde{C}\epsilon^{2-\alpha/2} \right) \tag{34}
$$

therefore there exists a constant $C_{KL^2} > 0$ only depending on k and $q^*(=2q)$ such that

$$
Var \le C_{KL^2} \frac{n}{\alpha} \left(\frac{\tilde{\epsilon}_n}{\sqrt{u_n}}\right)^{2-\alpha}.
$$
\n(35)

 $\rm\,B.2\quad Lemma~B.1:~control~of~\mathbb{E}^{\theta^*}\left(Z_t^2\right)$ $t^2\big)$

Lemma B.1. For all $\theta, \theta^* \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$

$$
\mathbb{E}^{\theta^*}(Z_t^2) \le \frac{16\rho^{2(t-1)}}{q^2} + 2 \max_{1 \le i \le k} \int f_i^*(y) \max_{1 \le j \le k} \log^2\left(\frac{f_j^*(y)}{f_j(y)}\right) \lambda(dy) + \frac{32 ||Q - Q^*||^2}{q^4(1-\rho)^2} + \frac{32}{q^4(1-\rho)^2} \int \min\left(\sum_{1 \le i \le k} \frac{|f_i^*(y) - f_i(y)|^2}{f_i^*(y)}, q^2 k^2 \sum_{1 \le j \le k} f_j^*(y)\right) \lambda(dy),
$$
\n(36)

with $\rho = \frac{1 - kq}{1 - (k-1)q} \leq 1 - q$. If moreover Assumptions [\(A.1\)](#page-5-4), [\(A.4\)](#page-5-8), [\(A.6\)](#page-5-5) and [\(6\)](#page-8-5) hold, then

$$
\mathbb{E}^{\theta^*}(Z_t^2) \le \frac{16\rho^{2(t-1)}}{\underline{q}^2} + \tilde{C}\frac{\tilde{\epsilon}_n^2}{u_n} \tag{37}
$$

where $\tilde{C} \leq 33(1+2k)/q^6$.

Proof of Lemma [B.1.](#page-17-0) Let $Q_{t,i}^{\mu,\theta,Y_{1:t-1}} = \mathbb{P}^{\theta}(X_t = i | Y_{1:t-1}, X_1 \sim \mu)$, where $t \geq 1, 1 \leq i \leq k$, $\mu \in \Delta_k$ and when μ is not specified, the stationary initial probability distribution is considered. Then the conditional density function of Y_t given $Y_{1:t-1}$ with respect to λ is:

$$
\sum_{i=1}^{k} f_i(\cdot) Q_{t,i}^{\theta, Y_{1:t-1}};
$$

so that

$$
\mathbb{E}^{\theta^*} \left(Z_t^2 \right) = \mathbb{E}^{\theta^*} \left(\left(\log \frac{\sum_{i=1}^k f_i^*(Y_t) Q_{t,i}^{\theta^*; Y_{1:t-1}}}{\sum_{j=1}^k f_j(Y_t) Q_{t,j}^{\theta^*; Y_{1:t-1}}} \right)^2 \right) \n\leq 2 \mathbb{E}^{\theta^*} \left(\max_{1 \leq j \leq k} \log^2 \left(\frac{f_j^*(Y_t)}{f_j(Y_t)} \right) \right) + \frac{2}{q^2} \mathbb{E}^{\theta^*} \left(\left(\sum_{j=1}^k |Q_{t,j}^{\theta^*; Y_{1:t-1}} - Q_{t,j}^{\theta; Y_{1:t-1}}| \right)^2 \right)
$$
\n(38)

using Equation [\(19\)](#page-14-2) and Lemma [A.1.](#page-15-0)

Combining Equation [\(38\)](#page-18-1) and Lemma [B.2](#page-18-0) (Equation [\(40\)](#page-18-2)), we obtain Equation [\(36\)](#page-17-2). Moreover, using Assumption [\(A.1\)](#page-5-4),

$$
\mathbb{E}^{\theta^*} \left(\max_{1 \le j \le k} \log^2 \left(\frac{f_j^*(Y_t)}{f_j(Y_t)} \right) \right) \le \epsilon^2. \tag{39}
$$

Finally, combining Equations [\(38\)](#page-18-1), [\(39\)](#page-18-3) and Lemma [B.2](#page-18-0) (Equation [\(41\)](#page-18-4)), we obtain Equation [\(37\)](#page-17-3). \Box

Lemma B.2. For all $\theta, \theta^* \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$ and $\mu, \mu^* \in \Delta_k(\underline{q})$,

$$
\mathbb{E}^{\theta^*} \left(\left(\sum_{j=1}^k |Q_{t,j}^{\theta^*, Y_{1:t-1}} - Q_{t,j}^{\theta, Y_{1:t-1}}| \right)^2 \right) \n\le 8\rho^{2(t-1)} + \frac{16 \|(Q - Q^* \|^2}{\underline{q}^2 (1 - \rho)^2} \n+ \frac{16}{\underline{q}^2 (1 - \rho)^2} \int \min \left(\sum_{1 \le i \le k} \frac{|f_i^*(y) - f_i(y)|^2}{f_i^*(y)}, \underline{q}^2 k^2 \sum_{1 \le j \le k} f_j^*(y) \right) \lambda(dy),
$$
\n(40)

with $\rho = \frac{1 - kq}{1 - (k-1)q} \leq 1 - q$.

If moreover \bar{A} ssumptions $(A.1)$, $(A.4)$, $(A.6)$ and (6) hold, then

$$
\mathbb{E}^{\theta^*} \left(\left(\sum_{j=1}^k |Q_{t,j}^{\theta^*, Y_{1:t-1}} - Q_{t,j}^{\theta, Y_{1:t-1}}| \right)^2 \right) \le 8\rho^{2(t-1)} + C' \frac{\tilde{\epsilon}_n^2}{u_n} \tag{41}
$$

where $C' \leq 16(1+2k)/q^4$.

Proof of Lemma [B.2.](#page-18-0) We first control $\sum_{i=1}^{k} |Q_{t,i}^{\theta^*, Y_{1:t-1}} - Q_{t,i}^{\mu^*, \theta, Y_{1:t-1}}|$. For this purpose, we are going to use a modified version of Proposition 1 of [Douc and Matias \[2001\]](#page-34-13). By Proposition 1 of [Douc and Matias \[2001\]](#page-34-13) and for all θ , θ^* in $\Delta_k^k(\underline{q}) \times \mathcal{F}^k$ we can control the L_1 -norm between two conditional probabilities of the state t when the initial probabilities are equal.

$$
\sum_{i=1}^{k} |Q_{t,i}^{\theta^*, Y_{1:t-1}} - Q_{t,i}^{\mu^*, \theta, Y_{1:t-1}}|
$$
\n
$$
\leq \sum_{j=1}^{k} \left| Q_{t-1,j}^{(Q_{2,\cdot}^{\theta^*, Y_1}, Q, f), Y_{2:t-1}} - Q_{t-1,j}^{(Q_{2,\cdot}^{\theta^*, Y_1}, Q^*, f^*), Y_{2:t-1}} \right| + \frac{1}{2} \left(\frac{1 - kq}{1 - (k-1)q} \right)^{t-1} \ast
$$
\n
$$
\sum_{u=1}^{k} \left| \frac{\sum_{j=1}^{k} (A^{Y_{t-1}, \theta} \dots A^{Y_2, \theta})_{j,u} (A^{Y_1, \theta} \mu^*)_u}{\sum_{v=1}^{k} (A^{Y_{t-1}, \theta} \dots A^{Y_2, \theta} A^{Y_1, \theta} \mu^*)_v} - \frac{\sum_{j=1}^{k} (A^{Y_{t-1}, \theta} \dots A^{Y_2, \theta})_{j,u} (A^{Y_1, \theta^*} \mu^*)_u}{\sum_{v=1}^{k} (A^{Y_{t-1}, \theta} \dots A^{Y_2, \theta} A^{Y_1, \theta^*} \mu^*)_v} \right|
$$
\n(42)

with
$$
(A_{i,j}^{Y,\theta})_{1 \le i,j \le k} = (Q_{j,i}f_j(Y))_{1 \le i,j \le k}
$$
. And
\n
$$
\sum_{u=1}^{k} \left| \frac{\sum_{j=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta} \mu^{*})_{u}}{\sum_{v=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta} \mu^{*})_{v}} - \frac{\sum_{j=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*})_{v}}{\sum_{v=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*} - A^{Y_{1},\theta} \mu^{*})_{u}}
$$
\n
$$
= \sum_{u=1}^{k} \left| \frac{\sum_{j=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*} - A^{Y_{1},\theta} \mu^{*})_{u}}{\sum_{v=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{i,u}(A^{Y_{1},\theta^{*}} \mu^{*} - A^{Y_{1},\theta} \mu^{*})_{w}}
$$
\n
$$
= \frac{\sum_{y=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*} - A^{Y_{1},\theta} \mu^{*})_{w}}{\sum_{v=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*})_{v}}
$$
\n
$$
\le 2 \sum_{u=1}^{k} \min \left(\left| \frac{\sum_{j=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*})_{v}}{\sum_{v=1}^{k} (A^{Y_{t-1},\theta} \dots A^{Y_{2},\theta})_{j,u}(A^{Y_{1},\theta^{*}} \mu^{*} - A^{Y_{1},\theta
$$

using Lemma [A.1.](#page-15-0) Combining Equations [\(42\)](#page-18-5) and [\(43\)](#page-19-0),

$$
\sum_{i=1}^{k} |Q_{t,i}^{\theta^*, Y_{1:t-1}} - Q_{t,i}^{\mu^*, \theta, Y_{1:t-1}}|
$$
\n
$$
\leq \sum_{j=1}^{k} \left| Q_{t-1,j}^{(Q_{2,:}^{\theta^*, Y_1}, Q, f), Y_{2:t-1}} - Q_{t-1,j}^{(Q_{2,:}^{\theta^*, Y_1}, Q^*, f^*), Y_{2:t-1}} \right| + \left(\frac{1 - kq}{1 - (k-1)q} \right)^{t-1} * \qquad (44)
$$
\n
$$
\left(\frac{\|Q - Q^*\|}{\frac{q}{2}} + \min\left(\frac{(1 - (k-1)q)}{\frac{q}{2}} \frac{\sum_{i=1}^{k} |f_i^*(Y_1) - f_i(Y_1)| \mu_i^*}{\sum_{j=1}^{k} f_j^*(Y_1) \mu_j^*}, k \right) \right).
$$

By repeating the arguments of Equation [\(43\)](#page-19-0), we show that

$$
\sum_{i=1}^{k} |Q_{2,i}^{(\mathcal{Q}_{t-1}^{\theta^*, Y_{1:t-2}}, \mathcal{Q}^*, f^*), Y_{1:t-1}} - Q_{2,i}^{(\mathcal{Q}_{t-1}^{\theta^*, Y_{1:t-2}}, \mathcal{Q}, f), Y_{1:t-1}}|
$$
\n
$$
= \sum_{i=1}^{k} \left| \frac{\sum_{i=1}^{k} A_{j,i}^{Y_{t-1}, \theta^*} Q_{t-1,i}^{\theta^*, Y_{1:t-2}}}{\sum_{i,u=1}^{k} A_{u,i}^{Y_{t-1}, \theta^*} Q_{t-1,i}^{\theta^*, Y_{1:t-2}}} - \frac{\sum_{i=1}^{k} A_{j,i}^{Y_{t-1}, \theta} Q_{t-1,i}^{\theta^*, Y_{1:t-2}}}{\sum_{i,u=1}^{k} A_{u,i}^{Y_{t-1}, \theta} Q_{t-1,i}^{\theta^*, Y_{1:t-2}}}\right|
$$
\n
$$
\leq 2 \left(\frac{||Q - Q^*||}{\underline{q}} + \min \left(\frac{(1 - (k-1)\underline{q})}{\underline{q}} \frac{\sum_{i=1}^{k} |f_i^*(Y_{t-1}) - f_i(Y_{t-1})| Q_{t-1,i}^{\theta^*, Y_{1:t-2}}}{\sum_{j=1}^{k} f_j^*(Y_{t-1}) Q_{t-1,j}^{\theta^*, Y_{1:t-2}}}, k \right) \right).
$$

By induction on [\(44\)](#page-19-1),

$$
\sum_{i=1}^{k} |Q_{t,i}^{\theta^*, Y_{1:t-1}} - Q_{t,i}^{\mu^*, \theta, Y_{1:t-1}}|
$$
\n
$$
\leq 2 \left(\frac{||Q - Q^*||}{\underline{q}} + \min \left(\frac{(1 - (k-1)\underline{q})}{\underline{q}} \frac{\sum_{i=1}^{k} |f_i^*(Y_{t-1}) - f_i(Y_{t-1})| Q_{t-1,i}^{\theta^*, Y_{1:t-2}}}{\sum_{j=1}^{k} f_j^*(Y_{t-1}) Q_{t-1,j}^{\theta^*, Y_{1:t-2}}}, k \right) \right)
$$
\n
$$
+ \sum_{u=3}^{t} \left(\frac{1 - k\underline{q}}{1 - (k-1)\underline{q}} \right)^{u-1}
$$
\n
$$
\left(\frac{||Q - Q^*||}{\underline{q}} + \min \left(\frac{(1 - (k-1)\underline{q})}{\underline{q}} \frac{\sum_{i} |f_i^*(Y_{t-u+1}) - f_i(Y_{t-u+1})| Q_{t-u+1,i}^{\theta^*, Y_{1:t-u}}}{\sum_{j} f_j^*(Y_{t-u+1}) Q_{t-u+1,j}^{\theta^*, Y_{1:t-u}}}, k \right) \right).
$$
\n(45)

Using Corollary 1 of [Douc et al. \[2004\]](#page-34-14), we can control the ℓ_1 -norm between two conditional probabilities of the state t for the same parameter θ but different initial probabilities: for all $\theta \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$, $\mu, \tilde{\mu} \in \Delta_k$ and for all $y_{1:l-1} \in \{y : \exists i, f_i^*(y) > 0\}^{l-1}$

$$
\sum_{i=1}^{k} \left| Q_{i,l}^{\mu,\theta,y_{1:l-1}} - Q_{i,l}^{\tilde{\mu},\theta,y_{1:l-1}} \right| \leq 2\rho^{l-1}.
$$
\n(46)

Combining Equations [\(45\)](#page-20-0) and [\(46\)](#page-20-1), we obtain

$$
\sum_{i=1}^{k} |Q_{t,i}^{\mu^*,\theta^*,Y_{1:t-1}} - Q_{t,i}^{\mu,\theta,Y_{1:t-1}}|
$$
\n
$$
\leq 2\left(\frac{||Q - Q^*||}{\underline{q}} + \min\left(\frac{1}{\underline{q}}\frac{\sum_{i=1}^{k}|f_i^*(Y_{t-1}) - f_i(Y_{t-1})|Q_{t-1,i}^{\theta^*,Y_{1:t-2}}}{\sum_{j=1}^{k}f_j^*(Y_{t-1})Q_{t-1,j}^{\theta^*,Y_{1:t-2}}},k\right)\right)
$$
\n
$$
+ 2\sum_{u=3}^{t} \left(\frac{1 - k\underline{q}}{1 - (k-1)\underline{q}}\right)^{u-1}
$$
\n
$$
\frac{\left(\frac{||Q - Q^*||}{\underline{q}} + \min\left(\frac{1}{\underline{q}}\frac{\sum_{i}|f_i^*(Y_{t-u+1}) - f_i(Y_{t-u+1})|Q_{t-u+1,i}^{\theta^*,Y_{1:t-u}}}{\sum_{j}f_j^*(Y_{t-u+1})Q_{t-u+1,i}^{\theta^*,Y_{1:t-u}}},k\right)\right)}{= \Delta_{t-u+1}} \qquad (47)
$$
\n
$$
\leq \frac{2}{\rho}\sum_{u=1}^{t-1} \rho^u\Delta_{t-u} + 4\rho^{t-1}.
$$

Then

$$
\left(\sum_{i=1}^{k} |Q_{t,i}^{\mu^*,\theta^*,Y_{1:t-1}} - Q_{t,i}^{\mu,\theta,Y_{1:t-1}}|\right)^2 \le \frac{8}{\rho^2} \left(\sum_{u=1}^{t-1} \rho^u\right) \left(\sum_{u=1}^{t-1} \rho^u \Delta_{t-u}^2\right) + 8\rho^{2(t-1)},\tag{48}
$$

using Cauchy-Schwarz inequality. Moreover, using Lemma [A.1,](#page-15-0)

$$
\mathbb{E}^{\theta^{*}}(\Delta_{t-u}^{2})
$$
\n
$$
\leq 2 \frac{\|Q - Q^{*}\|^{2}}{\underline{q}^{2}} + \frac{2}{\underline{q}^{2}} \mathbb{E}^{\theta^{*}} \left(\min \left(\left(\frac{\sum_{i} |f_{i}^{*}(Y_{t-u}) - f_{i}(Y_{t-u})| Q_{t-u,i}^{\theta^{*}, Y_{1:t-u-1}}}{\sum_{j} f_{j}^{*}(Y_{t-u}) Q_{t-u,j}^{\theta^{*}, Y_{1:t-u-1}}} \right)^{2}, (\underline{q}k)^{2} \right) \right)
$$
\n
$$
\leq 2 \frac{\|Q - Q^{*}\|^{2}}{\underline{q}^{2}}
$$
\n
$$
+ \frac{2}{\underline{q}^{2}} E \left(\int \sum_{1 \leq \iota \leq k} Q_{t-u,\iota}^{\theta^{*}, Y_{1:t-u-1}} f_{\iota}(y) \min \left(\left(\frac{\sum_{i} |f_{i}^{*}(y) - f_{i}(y)| Q_{t-u,i}^{\theta^{*}, Y_{1:t-u-1}}}{\sum_{j} f_{j}^{*}(y) Q_{t-u,j}^{\theta^{*}, Y_{1:t-u-1}}} \right)^{2}, (\underline{q}k)^{2} \right) \lambda(dy) \right)
$$
\n
$$
\leq 2 \frac{\|Q - Q^{*}\|^{2}}{\underline{q}^{2}} + \frac{2}{\underline{q}^{2}} \int \min \left(\sum_{1 \leq i \leq k} \frac{|f_{i}^{*}(y) - f_{i}(y)|^{2}}{f_{i}^{*}(y)} , \underline{q}^{2}k^{2} \sum_{1 \leq j \leq k} f_{j}^{*}(y) \right) \lambda(dy).
$$
\n(49)

Combining Equations [\(48\)](#page-20-2) and [\(49\)](#page-21-2), we obtain Equation [\(40\)](#page-18-2) which implies Equation [\(41\)](#page-18-4) under Assumptions [\(A.1\)](#page-5-4), [\(A.4\)](#page-5-8), [\(A.6\)](#page-5-5) and [\(6\)](#page-8-5). This concludes the proof of Lemma [B.2.](#page-18-0) \Box

 $\mathbf{B.3} \quad \textbf{Lemma B.3: control of } \mathbb{E}^{\theta^*}\big(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^2 \big)$ $\mathbf{B.3} \quad \textbf{Lemma B.3: control of } \mathbb{E}^{\theta^*}\big(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^2 \big)$ $\mathbf{B.3} \quad \textbf{Lemma B.3: control of } \mathbb{E}^{\theta^*}\big(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^2 \big)$

In the following lemma we show that $\mathbb{E}^{\theta^*}\left(\left(\left| \mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t) \right| \right)^2 \right)$ geometrically decreases to 0 when t tends to $+\infty$, using the exponential forgetting of the Markov chain. **Lemma B.3.** For all $\theta, \theta^* \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$ and $\alpha \in (0, 2)$,

$$
\mathbb{E}^{\theta^*} \left(|\mathbb{E}^{\theta^*} (Z_t | Y_{1:r-1}) - \mathbb{E}^{\theta^*} (Z_t) |^2 \right) \n\leq 8 \mathbb{E}^{\theta^*} (Z_t^2)^{\frac{2-\alpha}{2}} \rho^{-\frac{5\alpha}{2}} \rho^{\frac{\alpha}{2}(t-r)} \left(2 \max_{1 \leq j \leq k} \int f_j^*(y) \max_{1 \leq i \leq k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \frac{10}{q} \right)^{\alpha}.
$$
\n(50)

If moreover Assumption [\(A.1\)](#page-5-4) holds then

$$
\mathbb{E}^{\theta^*}(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^2) \leq 8\mathbb{E}^{\theta^*}(Z_t^2)^{\frac{2-\alpha}{2}}\rho^{-\frac{5\alpha}{2}}\rho^{\frac{\alpha}{2}(t-r)}\left(2\frac{\tilde{\epsilon}_n}{\sqrt{u_n}} + \frac{10}{\underline{q}}\right)^{\alpha}.
$$
 (51)

Proof of Lemma [B.3.](#page-21-0) Denote

$$
L_t = (X_t, Y_t, Q_{t, \cdot}^{\theta, Y_{1:t-1}}, Q_{t, \cdot}^{\theta^*, Y_{1:t-1}})
$$

for all $t \in \mathbb{N}$, then $(L_t)_{t \in \mathbb{N}}$ is the extended Markov chain with transition kernel Π_{θ} more precisely described in [\[Douc and Matias, 2001\]](#page-34-13) at page 384. Let

$$
h\colon \begin{cases} \{1,\ldots,k\} \times \{y: \exists 1 \leq j \leq k, f_j^*(y) > 0\} \times \{\mu \in \Delta_k : \mu_i > \underline{q} \,\,\forall i\}^2 \longrightarrow \mathbb{R} \\ l = (x,y,\mu,\mu^*) \longmapsto h(l) = \log\left(\frac{\sum_{i=1}^k \mu_i^* f_i^*(y)}{\sum_{i=1}^k \mu_i f_i(y)}\right) \end{cases}
$$

then

$$
h(L_t) = Z_t = \log \left(\frac{p^{\theta^*}(Y_t|Y_{1:t-1})}{p^{\theta}(Y_t|Y_{1:t-1})} \right)
$$

and for all $r \leq t$ and $0 < \alpha < 2$,

$$
\mathbb{E}^{\theta^*}(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^2) \n= \mathbb{E}^{\theta^*}(|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^{2-\alpha}|\mathbb{E}^{\theta^*}(Z_t|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(Z_t)|^{\alpha}) \n\leq 2^{2-\alpha}\mathbb{E}^{\theta^*}((\max(\mathbb{E}^{\theta^*}(|Z_t|), \mathbb{E}^{\theta^*}(|Z_t||Y_{1:r-1})))^{2-\alpha}|\mathbb{E}^{\theta^*}(h(L_t)|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(h(L_t))|^{\alpha}).
$$
\n(52)

The following term is geometrically decreasing, using Lemma [B.4](#page-22-0)

$$
|\mathbb{E}^{\theta^*}(h(L_t)|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(h(L_t))|
$$

\$\leq \int \int \int h(l_t) \Pi_{\theta}^{t-r}(l_r, dl_t) - \int h(l_t) \Pi_{\theta}^{t-r}(\tilde{l}_r, dl_t) \Big| P^{\theta}(dl_r|Y_{1:r-1}) P^{\theta}(d\tilde{l}_r).\$ (53)

More precisely, using Equation [\(53\)](#page-22-1) and Lemma [B.4](#page-22-0) with $m = \frac{t-r+1}{2}$ $\lfloor \frac{r+1}{2} \rfloor$ and $u = t - r$, we obtain

$$
|\mathbb{E}^{\theta^*}(h(L_t)|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(h(L_t))| \le \rho^{\frac{t-r}{2} - \frac{5}{2}} \left(2 \max_{1 \le j \le k} \int f_j^*(y) \max_{1 \le i \le k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \frac{10}{q} \right).
$$
\n
$$
\text{and } \rho^{\theta^*}(h(L_t)|Y_{1:r-1}) - \mathbb{E}^{\theta^*}(h(L_t))| \le \rho^{\frac{t-r}{2} - \frac{5}{2}} \left(2 \max_{1 \le j \le k} \int f_j^*(y) \max_{1 \le i \le k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \frac{10}{q} \right).
$$

Therefore using Equations [\(52\)](#page-21-3) and [\(54\)](#page-22-2),

$$
\mathbb{E}^{\theta^*} \left(|\mathbb{E}^{\theta^*} (Z_t | Y_{1:r-1}) - \mathbb{E}^{\theta^*} (Z_t) |^2 \right) \n\leq 2^{2-\alpha} \mathbb{E}^{\theta^*} \left(\max(\mathbb{E}^{\theta^*} ((|Z_t|), \mathbb{E}^{\theta^*} (|Z_t| | Y_{1:r-1})))^{2-\alpha} \right) \n\rho^{-\frac{5\alpha}{2}} \rho^{\frac{\alpha}{2}(t-r)} \left(2 \max_{1 \leq j \leq k} \int f_j^*(y) \max_{1 \leq i \leq k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \frac{10}{q} \right)^{\alpha}
$$
\n(55)

By convexity of the square function and concavity of $x \mapsto x^{\frac{2-\alpha}{2}}$, with $0 < \alpha < 2$,

$$
\mathbb{E}^{\theta^*} \left(\max(\mathbb{E}^{\theta^*}(|Z_t|), \mathbb{E}^{\theta^*}(|Z_t||Y_{1:r-1}))^{2-\alpha} \right)
$$
\n
$$
\leq \mathbb{E}^{\theta^*} \left(\max(\mathbb{E}^{\theta^*}(Z_t^2)^{1/2}, \mathbb{E}^{\theta^*}(Z_t^2|Y_{1:r-1})^{1/2})^{2-\alpha} \right)
$$
\n
$$
\leq \mathbb{E}^{\theta^*}(Z_t^2)^{\frac{2-\alpha}{2}} + \mathbb{E}^{\theta^*}(\mathbb{E}^{\theta^*}(Z_t^2|Y_{1:r-1})^{\frac{2-\alpha}{2}})
$$
\n
$$
\leq 2\mathbb{E}^{\theta^*}(Z_t^2)^{\frac{2-\alpha}{2}}.
$$
\n(56)

Combining Equations [\(55\)](#page-22-3) and [\(56\)](#page-22-4), we get Equation [\(50\)](#page-21-4). Besides, using Assumption [\(A.1\)](#page-5-4) and Cauchy–Schwarz inequality ,

$$
\max_{1 \le j \le k} \int f_j^*(y) \max_{1 \le i \le k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) \le \epsilon
$$
\n(57)

so that Equation [\(51\)](#page-21-5) holds.

Lemma [B.4](#page-22-0) is an improved version of Proposition 2 of [\[Douc and Matias, 2001\]](#page-34-13).

Lemma B.4. For all integers $u > 0$, $m < u$, for all $z, \tilde{z} \in \{1, ..., k\} \times \{y : \exists 1 \le j \le k\}$ $k, f_j^*(y) > 0$ $\} \times {\mu \in \Delta_k : \mu_i > \underline{q} \,\forall i$ $\} \times {\mu \in \Delta_k : \mu_i > \underline{q} \,\forall i}$ and for all $\theta, \theta^* \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$,

$$
\left| \int h(l) \Pi_{\theta}^{u}(z,dl) - \int h(l) \Pi_{\theta}^{u}(\tilde{z},dl) \right|
$$

\n
$$
\leq \frac{4}{\underline{q}} \rho^{u-1} + \frac{4}{\underline{q}} \rho^{m} + 2\rho^{m-2} \left(\max_{1 \leq j \leq k} \int f_{j}^{*}(y) \max_{1 \leq i \leq k} \left| \log \frac{f_{i}^{*}(y)}{f_{i}(y)} \right| \lambda(dy) + \log \left(\frac{1}{\underline{q}} \right) \right)
$$
\n(58)

Proof of Lemma [B.4.](#page-22-0) This lemma is an improved version of Proposition 2 of [Douc and](#page-34-13) [Matias \[2001\]](#page-34-13). We improve the result by defining h on

$$
\mathcal{Z} = \{1, \ldots, k\} \times \{y : \exists 1 \le j \le k, f_j^*(y) > 0\} \times \{\mu \in \Delta_k : \mu_i > \underline{q} \,\forall i\} \times \{\mu \in \Delta_k : \mu_i > \underline{q} \,\forall i\}
$$

 \Box

and using that if $z \in \mathcal{Z}$ then $\Pi_{\theta^*}(z, \mathcal{Z}) = 1$. Then we obtain

$$
\text{lip}(h, x, y) = \frac{1}{\underline{q}} \tag{59}
$$

since for all $(x, y, \mu, \mu^*), (x, y, \tilde{\mu}, \tilde{\mu}^*) \in \mathcal{Z}$

$$
|h(x, y, \mu, \mu^*) - h(x, y, \tilde{\mu}, \tilde{\mu}^*)| = \left| \log \left(\frac{\sum_{i=1}^k \mu_i^* f_i^*(y)}{\sum_{i=1}^k \mu_i f_i(y)} \right) - \log \left(\frac{\sum_{i=1}^k \tilde{\mu}_i^* f_i^*(y)}{\sum_{i=1}^k \tilde{\mu}_i f_i(y)} \right) \right|
$$

\n
$$
= \left| \log \left(\frac{\sum_{i=1}^k \mu_i^* f_i^*(y)}{\sum_{i=1}^k \tilde{\mu}_i^* f_i^*(y)} \right) - \log \left(\frac{\sum_{i=1}^k \mu_i f_i(y)}{\sum_{i=1}^k \tilde{\mu}_i f_i(y)} \right) \right|
$$

\n
$$
\leq \max_{1 \leq i \leq k} \left| \log \left(\frac{\mu_i^*}{\tilde{\mu}_i^*} \right) \right| + \max_{1 \leq i \leq k} \left| \log \left(\frac{\mu_i}{\tilde{\mu}_i} \right) \right|
$$

\n
$$
\leq \frac{1}{q} \left(\sum_{i=1}^k |\mu_i^* - \tilde{\mu}_i^*| + \sum_{i=1}^k |\mu_i - \tilde{\mu}_i| \right)
$$

using . Moreover

$$
k(h, x, y) = \log \frac{1}{q} + \max_{1 \le i \le k} \left| \log \left(\frac{f_i^*(y)}{f_i(y)} \right) \right| \tag{60}
$$

because, using Lemma [A.1,](#page-15-0)

$$
|h(x, y, \mu, \mu^*)| = \left| \log \left(\frac{\sum_{i=1}^k \mu_i^* f_i^*(y)}{\sum_{i=1}^k \mu_i f_i(y)} \right) \right|
$$

\n
$$
\leq \max_{1 \leq i \leq k} \left| \log \left(\frac{\mu_i^* f_i^*(y)}{\mu_i f_i(y)} \right) \right|
$$

\n
$$
\leq \max_{1 \leq i \leq k} \left| \log \left(\frac{\mu_i^*}{\mu_i} \right) \right| + \max_{1 \leq i \leq k} \left| \log \left(\frac{f_i^*(y)}{f_i(y)} \right) \right|
$$

\n
$$
\leq \log \frac{1}{\underline{q}} + \max_{1 \leq i \leq k} \left| \log \left(\frac{f_i^*(y)}{f_i(y)} \right) \right|.
$$

Moreover instead of using Proposition 1 of [\[Douc and Matias, 2001\]](#page-34-13) we use Corollary 1 of [Douc et al. \[2004\]](#page-34-14) so that for all $\theta \in \Delta_k^k(\underline{q}) \times \mathcal{F}^k$, $\mu, \tilde{\mu} \in \Delta_k(\underline{q})$ and for all $y_{1:l-1} \in$ ${y : \exists i, f_i^*(y) > 0}^{l-1}$

$$
\sum_{i=1}^{k} \left| Q_{i,l}^{\mu,\theta,y_{1:l-1}} - Q_{i,l}^{\tilde{\mu},\theta,y_{1:l-1}} \right| \leq 2\rho^{l-1}.
$$
 (61)

Then, let $z = (x, y, \mu, \mu^*) \in \mathcal{Z}$ and $\tilde{z} = (\tilde{x}, \tilde{y}, \tilde{\mu}, \tilde{\mu}^*) \in \mathcal{Z}$ using the proof of Proposition 1 of [\[Douc and Matias, 2001\]](#page-34-13):

$$
\begin{split}\n|\int h(l)\Pi^{u}(z,dl) - \int h(l)\Pi^{u}(\tilde{z},dl)| \\
&\leq \underbrace{\left|\int h(l)\Pi^{u}((x,y,\mu,\mu^{*}),dl) - \int h(l)\Pi^{u}((x,\tilde{y},\tilde{\mu},\tilde{\mu}^{*}),dl)\right|}_{=A} \\
&\quad + \underbrace{\left|\int h(l)\Pi^{u}(x,\tilde{y},\tilde{\mu},\tilde{\mu}^{*}),dl) - \int h(l)\Pi^{u}((\tilde{x},\tilde{y},\tilde{\mu},\tilde{\mu}^{*}),dl)\right|}_{=B}\n\end{split} \tag{62}
$$

where

$$
|A| \leq \left| \sum_{x_{2:u+1}=1}^{k} \int \text{lip}(h, x_{u+1}, y_{u+1}) \right|
$$

$$
\left(\sum_{i=1}^{k} |Q_{u,i}^{Q_1^{\mu,\theta,y},\theta,y_{2:u}} - Q_{u,i}^{Q_1^{\tilde{\mu},\theta,\tilde{y}},\theta,y_{2:u}}| + \sum_{i=1}^{k} |Q_{u,i}^{Q_1^{\mu^*,\theta^*,y},\theta^*,y_{2:u}} - Q_{u,i}^{Q_1^{\tilde{\mu}^*,\theta^*,\tilde{y}},\theta^*,y_{2:u}}| \right) \tag{63}
$$

$$
Q_{x,x_2}^* \dots Q_{x_u,x_{u+1}}^* f_{x_2}^*(y_2) \dots f_{x_{u+1}}^*(y_{u+1}) \lambda(dy_2) \dots \lambda(dy_{u+1}) \right|
$$

and for any $1 \leq m \leq u$,

$$
|B| \leq \sum_{x_{2:u+1}=1}^{k} \int \text{lip}(h, x_{u+1}, y_{u+1}) \left(\sum_{i=1}^{k} |Q_{m+1,m}^{Q_{u+1-m}^{i,\theta,y_{u-m+1}};\omega} - Q_{m+1,i}^{\nu,\theta,y_{u-m+1}} - Q_{m+1,i}^{\nu,\theta,y_{u-m+1}} - \sum_{i=1}^{k} |Q_{m+1,m}^{Q_{u+1-m}^{i,\theta,y_{u,m+1}};\omega} - Q_{m+1,i}^{\nu,\theta,y_{u-m+1}}| \right) + \sum_{i=1}^{k} |Q_{x,x_2}^{*} - Q_{\tilde{x},x_2}^{*} |Q_{x_2,x_3}^{*} \dots Q_{x_u,x_{u+1}}^{*} f_{x_2}^{*}(y_2) \dots f_{x_{u+1}}^{*}(y_{u+1}) \lambda(dy_{2:u+1}) + \sum_{x_{m:u+1}=1}^{k} \int k(h, x_{u+1}, y_{u+1}) |Q_{x,x_m}^{*} - Q_{\tilde{x},x_m}^{*} |Q_{x_m,x_{m+1}}^{*} \dots Q_{x_u,x_{u+1}}^{*} f_{x_m}^{*}(y_m) \dots f_{x_{u+1}}^{*}(y_{u+1}) \lambda(dy_{2:u+1}).
$$
\n(64)

Combining Equations [\(59\)](#page-23-0), [\(61\)](#page-23-1) and [\(63\)](#page-24-1),

$$
|A| \le \frac{4}{\underline{q}} \rho^{u-1} \tag{65}
$$

and using Equations [\(59\)](#page-23-0), [\(60\)](#page-23-2), [\(61\)](#page-23-1) and [\(64\)](#page-24-2)

$$
|B| \le \frac{4}{q} \rho^m + 2\rho^{m-2} \left(\max_{1 \le j \le k} \int f_j^*(y) \max_{1 \le i \le k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \log \left(\frac{1}{q} \right) \right) \tag{66}
$$

therefore using Equations (62) , (65) and (66) we obtain

$$
\left| \int h(l) \Pi^u(z, dl) - \int h(l) \Pi^u(\tilde{z}, dl) \right|
$$

\n
$$
\leq \frac{4}{\underline{q}} \rho^{u-1} + \frac{4}{\underline{q}} \rho^m + 2\rho^{m-2} \left(\max_{1 \leq j \leq k} \int f_j^*(y) \max_{1 \leq i \leq k} \left| \log \frac{f_i^*(y)}{f_i(y)} \right| \lambda(dy) + \log \left(\frac{1}{\underline{q}} \right) \right).
$$
\n
$$
\Box
$$

C Proof of Theorem [4.1](#page-11-0) (discrete observations)

Assumption [\(B\)](#page-5-3) will be checked using Proposition 2 of [\[Shen et al., 2013\]](#page-35-15) that we recall here.

Lemma C.1. [Proposition 2 of [\[Shen et al., 2013\]](#page-35-15)] Let H be a positive integer, A and ϵ be positive, denote

$$
\mathcal{H}_{H,A,\epsilon} = \left\{ f = \sum_{h=1}^{+\infty} \pi_h \delta_{z_h} \; : \; \sum_{h>H} \pi_h < \epsilon, z_h \in [0,A], h \leq H \right\}^k.
$$

Then

$$
DP(G) \left(\mathcal{H}_{H,A,\epsilon}^{c} \right) \leq k HG((A, +\infty)) + k \left(\frac{eG(\mathbb{N})}{H} \log \frac{1}{\epsilon} \right)^{H}
$$
(68)

$$
N(4l\epsilon, \mathcal{H}_{H,A,\epsilon}, d) \lesssim (A+1)^{kH} \epsilon^{-kH}.
$$
\n(69)

We now give the proof of Theorem [4.1.](#page-11-0)

Proof of Theorem [4.1.](#page-11-0) We first prove Assumption [\(A\)](#page-5-2) with $\tilde{f}_j = f_j$, for all $1 \leq j \leq k$ using Lemma [C.2](#page-26-0) with $\epsilon = \tilde{\epsilon}_n^2/u_n$, $L = L_n = \left(-\log(\tilde{\epsilon}_n^2/(u_n \log \log n))\right)/(c - \delta)\right)^{1/m}$ and $S_L = \{1, ..., L\}$. Using that $f_i^* \in \mathcal{D}(m, c, K)$ for all $1 \leq i \leq k$ and Assumption [\(P\),](#page-11-3) we get

$$
\sum_{l>L_n} \frac{f_i^*(l)}{(G(l))^2} \lesssim \sum_{l>L_n} \exp(-cl^m)l^{2\alpha} \lesssim \sum_{l>L_n} \exp(-(c-\delta)l^m)l^{m-1}
$$

$$
\lesssim \int_{L_n}^{\infty} \exp(-(c-\delta)x^m)x^{m-1}\lambda(dx) \lesssim \exp(-(c-\delta)L_n^m) \lesssim \tilde{\epsilon}_n^2
$$

which proves Equation [\(77\)](#page-26-1). Equation [\(78\)](#page-26-2) is proved similarly. Equation [\(76\)](#page-26-3) follows using Assumption [\(I\).](#page-11-2) Then, we can apply Lemma [C.2](#page-26-0) so that

$$
DP(G)^{\otimes k} \left(f_j \ : \forall 1 \le i, j \le k \sum_{l=1}^{+\infty} f_i^*(l) \log^2 \left(\frac{f_j^*(l)}{f_j(l)} \right) \le \frac{\epsilon_n^2}{u_n}, \sum_{l=1}^{L_n} \frac{\left(f_j^*(l) - f_j(l) \right)^2}{f_j^*(l)} \le \frac{\epsilon_n^2}{u_n}, \sum_{l>L_n} f_j(l) \le \frac{\epsilon_n^2}{u_n}, \sum_{l=1}^{L_n} \frac{\left(f_j^*(l) - f_j(l) \right)^2}{f_j(l)} \le \frac{\epsilon_n^2}{u_n} \right) \tag{70}
$$

$$
\gtrsim \prod_{j=1}^k \left(\left(\frac{\epsilon_n^2}{4u_n} \right)^{(L_n - 1 + G(\mathbb{N}))/2} f_j^*(l^*)^{L_n - 2} \left(\frac{1}{3} \right)^{L_n} \prod_{l=1}^{L_n} G(l) f_j^*(l)^{G(l)} \right).
$$

Moreover using that $f_i^* \in \mathcal{D}(m, c, K)$ for all $1 \leq i \leq k$ and Equation [\(P\),](#page-11-3) we obtain

$$
\log \prod_{l=1}^{L_n} G(l) \gtrsim -L_n \log(L_n),\tag{71}
$$

$$
\log \left(\prod_{l=1}^{L_n} f_j^*(l)^{G(l)} \right) \gtrsim \sum_{l=1}^{L_n} G(l) \log f_j^*(l) \gtrsim L_n^K. \tag{72}
$$

Combining Equations [\(70\)](#page-25-0) with $l^* = \arg \max_l (\min_{1 \leq j \leq k} f_j^*(l))$, [\(71\)](#page-25-1) and [\(72\)](#page-25-2), Assumption [\(A\)](#page-5-2) of Theorem [3.1](#page-6-0) is true if

$$
(-\log(\tilde{\epsilon}_n))^{\max(1/m+1,K/m)} \lesssim n\tilde{\epsilon}_n^2.
$$

Then we choose

$$
\tilde{\epsilon}_n = \frac{1}{\sqrt{n}} (\log n)^{t_0}
$$

with $2t_0 > \max(1/m + 1, K/m)$ and Assumption [\(A\)](#page-5-2) holds.

Using Assumption [\(Q0\),](#page-9-2) for $\tilde{\epsilon}_n$ small enough,

$$
\Pi_Q\left(\left\{Q: \|Q - Q^*\| \le \frac{\tilde{\epsilon}_n}{\sqrt{u_n}}\right\}\right) \ge \frac{\pi(Q^*)}{2}\lambda\left(\left\{Q: \|Q - Q^*\| \le \frac{\tilde{\epsilon}_n}{\sqrt{u_n}}\right\}\right) \gtrsim \left(\frac{\tilde{\epsilon}_n}{\sqrt{u_n}}\right)^{k(k-1)}\tag{73}
$$

so that Assumption [\(C\)](#page-6-1) holds.

Using Lemma [C.1](#page-24-5) with $A = \exp((\log n)^{2t_0+t/2}), H = (n\epsilon_n^2)/((\log n)^{2t_0+t})$ and $\epsilon_n =$ $(\log n)^t/\sqrt{n},$

$$
\Pi(\mathcal{F}_n^c)
$$

\n
$$
\lesssim (\log n)^{t-2t_0} \exp\left(-(\alpha+1)(\log n)^{2t_0+t/2}\right) + \exp\left(-(t-2t_0-1)(\log n)^{t-2t_0}(\log \log n)^2\right)
$$

\n
$$
= o(\exp(-C'(\log n)^{2t_0})) = o(\exp(-C'n\tilde{\epsilon}_n^2)),
$$
\n(74)

if $t > 4t_0$. Moreover

$$
\log\left(N\left(\frac{\epsilon_n}{12},\mathcal{F}_n,D_\ell\right)\right) \lesssim (\log n)^{3t/2} + (\log n)^{t-2t_0+1} \lesssim n\epsilon_n^2 \tag{75}
$$

so that Assumption [\(B\)](#page-5-3) holds. This concludes the proof of Theorem [4.1.](#page-11-0)

Lemma C.2. Let S_L be a subset of $\{1, \ldots, L\}$. If

$$
\max_{1 \le i,j \le k} \sum_{S_L^c} f_i^*(l) \log^2(f_j^*(l)) \le \frac{\epsilon}{8} \tag{76}
$$

 \Box

and

$$
\max_{1 \le i \le k} \sum_{S_L^c} \frac{f_i^*(l)}{(G(l))^2} \le \frac{\epsilon}{384 \log^2(2) k (G(\mathbb{N}))^2},\tag{77}
$$

and if there exists $\delta > 0$ such that

$$
\max_{1 \le i \le k} \sum_{l \in S_L^c} f_i^*(l) \le \epsilon^{1+\delta},\tag{78}
$$

then, for all $l^* \in S_L$ and all $\epsilon > 0$ small enough,

$$
P_G := DP(G)^{\otimes k} \left(f_j \ : \forall 1 \le i, j \le k \sum_{l=1}^{+\infty} f_i^*(l) \log^2 \left(\frac{f_j^*(l)}{f_j(l)} \right) \le \epsilon,
$$

$$
\sum_{l \in S_L} \frac{\left(f_j^*(l) - f_j(l) \right)^2}{f_j^*(l)} \le \epsilon, \sum_{l \in S_L^c} f_j(l) \le \epsilon, \sum_{l \in S_L} \frac{\left(f_j^*(l) - f_j(l) \right)^2}{f_j(l)} \le \epsilon \right)
$$

$$
\ge \prod_{j=1}^k \left(\left(\sqrt{\frac{\epsilon}{4}} \right)^{L-1+G(\mathbb{N})} \left(\frac{1}{3} \right)^L f_j^*(l^*)^{L-2} \left[\prod_{l \in S_L} G(l) f_j^*(l)^{G(l)} \right] \right).
$$

Proof of Lemma [C.2.](#page-26-0) Note that if for all $l \in S_L$ and for all $1 \leq j \leq k$,

$$
\left(1 - \sqrt{\frac{\epsilon}{4}}\right) f_j^*(l) \le f_j(l) \le \left(1 + \sqrt{\frac{\epsilon}{4}}\right) f_j^*(l)
$$

then for all $l \in S_L$,

$$
\log^2\left(\frac{f_j^*(l)}{f_j(l)}\right) \le \frac{\epsilon}{2}, \quad \frac{|f_j^*(l) - f_j(l)|^2}{f_j^*(l)^2} \le \frac{\epsilon}{2k} \quad \text{and} \quad \frac{|f_j^*(l) - f_j(l)|^2}{f_j(l)^2} \le \epsilon
$$

so that

$$
\sum_{l \in S_L} f_i^*(l) \log^2 \left(\frac{f_j^*(l)}{f_j(l)} \right) \le \frac{\epsilon}{2}, \sum_{l \in S_L} \frac{\left| f_j^*(l) - f_j(l) \right|^2}{f_j^*(l)} \le \epsilon \text{ and } \sum_{l \in S_L} \frac{\left| f_j^*(l) - f_j(l) \right|^2}{f_j(l)} \le \epsilon.
$$

Moreover using Assumptions [\(76\)](#page-26-3) and [\(78\)](#page-26-2) if for all $1 \le i, j \le k$,

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2(f_j(l)) < \frac{\epsilon}{8},
$$

then

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f_j^*(l)}{f(l)} \right) \le 2 \sum_{l \in S_L^c} f_i^*(l) \log^2(f_j^*(l)) + 2 \sum_{l \in S_L^c} f_i^*(l) \log^2(f_j(l)) \le \frac{\epsilon}{2}.
$$

Combining the two last remarks, we obtain

$$
P_G \geq \prod_{j=1}^k \left(DP(G) \left(f : \left(1 - \sqrt{\frac{\epsilon}{4}} \right) f_j^*(l) \leq f(l) \leq \left(1 + \sqrt{\frac{\epsilon}{4}} \right) f_j^*(l), \ \forall l \in S_L, \right) \right)
$$

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2(f(l)) \leq \frac{\epsilon}{8} \text{ and } \sum_{l \in S_L^c} f_j(l) \leq \epsilon \right)
$$

$$
\geq \prod_{j=1}^k \left(DP(G) \left(f : \exp \left(-\sqrt{\frac{\epsilon}{16 \max_{1 \leq i \leq k} \sum_{l \in S_L^c} f_i^*(l)} \right)} \right) \leq \sum_{m \in S_L^c} f(m) \leq \epsilon,
$$

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) \leq \frac{\epsilon}{32}, \ \forall 1 \leq i \leq k \text{ and}
$$

$$
\left(1 - \sqrt{\frac{\epsilon}{4}} \right) \frac{f_j^*(l)}{\sum_{m \in S_L} f(m)} \leq \frac{f(l)}{\sum_{m \in S_L} f(m)} \leq \left(1 + \sqrt{\frac{\epsilon}{4}} \right) \frac{f_j^*(l)}{\sum_{m \in S_L} f(m)}, \ \forall l \in S_L \right)
$$

indeed if

$$
\exp\left(-\sqrt{\frac{\epsilon}{16 \max_{1 \le i \le k} \sum_{l \in S_L^c} f_i^*(l)}}\right) \le \sum_{m \in S_L^c} f(m)
$$

and

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) \le \frac{\epsilon}{32}, \ \forall 1 \le i \le k
$$

then

$$
\sum_{l \in S_L^c} f_i^*(l) \log^2(f(l))
$$
\n
$$
\leq 2 \sum_{l \in S_L^c} f_i^*(l) \log^2\left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)}\right) + 2 \sum_{l \in S_L^c} f_i^*(l) \log^2\left(\sum_{m \in S_L^c} f(m)\right) \leq \frac{\epsilon}{8}.
$$

Using the tail free property of the Dirichlet process, $(f(l)/\sum_{m\in S_L^c}f(m))_{l\in S_L^c}, \sum_{m\in S_L^c}f(m)$ and $(f(1)/\sum_{m\in S_L}f(m),\ldots,f(L)/\sum_{m\in S_L}f(m))$ are independent. So that we obtain

$$
P_G \ge \int_{\exp\left(-\sqrt{\frac{\epsilon}{16 \max_{1 \le i \le k} \sum_{l \in S_L^c} f_i^*(l)}}\right)}^{\epsilon} \frac{\Gamma(G(\mathbb{N}))}{\Gamma(G(S_L))\Gamma(G(S_L^c))} a^{G(S_L^c) - 1} (1 - a)^{G(S_L) - 1}
$$

$$
Dir(G|_{S_L}) \left(x : \left(1 - \sqrt{\frac{\epsilon}{4}}\right) \frac{f_j^*(l)}{1 - a} \le x_l \le \left(1 + \sqrt{\frac{\epsilon}{4}}\right) \frac{f_j^*(l)}{1 - a}, \ \forall l \in S_L\right) \lambda(da)
$$

$$
DP(G|_{S_L^c}) \left(\sum_{l \in S_L^c} f_i^*(l) \log^2\left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)}\right) < \frac{\epsilon}{32}, \ \forall 1 \le i \le k\right)
$$

(79)

We first control the integral of Equation [\(79\)](#page-27-0). Note that if

$$
x \in V_{SL} := \left\{ x \in \Delta_{|S_L|} : x_l \in V_l, \ \forall l \in S_L \setminus \{l^*\} \right\}
$$

where for all $l \in S_L \setminus \{l^*\}$

$$
V_l := \left\{ x_l : \left(1 - \sqrt{\frac{\epsilon}{16}} f_j^*(l^*) \right) \frac{f_j^*(l)}{1 - a} \le x_l \le \left(1 + \sqrt{\frac{\epsilon}{16}} f_j^*(l^*) \right) \frac{f_j^*(l)}{1 - a} \right\}
$$

and if

$$
a \in V_A := \left\{ \sum_{l \in S_L} f_j^*(l) - \sqrt{\frac{\epsilon}{16}} f_j^*(l^*) \le 1 - a \le \sum_{l \in S_L} f_j^*(l) + \sqrt{\frac{\epsilon}{16}} f_j^*(l^*) \right\} \tag{80}
$$

then for all $l \in S_L$,

$$
\left(1 - \sqrt{\frac{\epsilon}{4}}\right) \frac{f_j^*(l)}{1 - a} \le x_l \le \left(1 + \sqrt{\frac{\epsilon}{4}}\right) \frac{f_j^*(l)}{1 - a},\tag{81}
$$

where $x_{l^*} = 1 - \sum_{l \in S_L, l \neq l^*} x_l$. So that

$$
D(a) \ge \frac{\Gamma(G(S_L))}{\prod_{m \in S_L} \Gamma(G(m))} 1_{V_A}(a) \int_{V_{SL}} \prod_{l \in S_L \setminus \{l^*\}} x_l^{G(l)-1} \left(1 - \sum_{m \in S_L \setminus \{l^*\}} x_m \right)^{G(l^*)-1} \lambda(dx)
$$
\n(82)

where

$$
\left(1 - \sum_{m \in S_L \setminus \{l^*\}} x_m\right)^{G(l^*)-1} \ge \left(\frac{f_j^*(l^*)}{1-a}\right)^{G(l^*)-1} \min\left((1/2)^{G(l^*)-1}, (3/2)^{G(l^*)-1}\right) \tag{83}
$$

and

$$
\int_{V_l} x_l^{G(l)-1} \lambda(dx_l) \ge \left(\frac{f_j^*(l)}{1-a}\right)^{G(l)} f_j^*(l^*) \sqrt{\frac{\epsilon}{4}} \min\left((1/2)^{G(l)-1}, (3/2)^{G(l)-1}\right),\tag{84}
$$

using Equation [\(81\)](#page-28-0). Then combining Equations [\(82\)](#page-28-1), [\(83\)](#page-28-2) and [\(84\)](#page-28-3); $D(a)$ is bounded by

$$
\mathbb{1}_{V_A}(a) \frac{\Gamma(G(S_L))}{\prod_{m \in S_L} \Gamma(G(m))} \left(\frac{2}{3}\right)^L \left(\frac{1}{2}\right)^{G(\mathbb{N})} \left(\frac{\epsilon}{4}\right)^{(L-1)/2} \left(\frac{1}{1-a}\right)^{G(S_L)-1} f_j^*(l^*)^{L-2} \prod_{l \in S_L} f_j^*(l)^{G(l)}.
$$

So that

$$
\int_{\exp}^{\epsilon} \left(-\sqrt{\frac{c}{16 \max_{1 \le i \le k} \sum_{l \in S_L^c} f_i^*(l)}} \right) \frac{\Gamma(G(\mathbb{N}))}{\Gamma(G(S_L)) \Gamma(G(S_L^c))} a^{G(S_L^c) - 1} (1 - a)^{G(S_L) - 1} D(a) \lambda(da)
$$
\n
$$
\ge \frac{\Gamma(G(\mathbb{N}))}{G(S_L^c) \Gamma(G(S_L^c)) \prod_{l \in S_L} \Gamma(G(l))} \left(\frac{\epsilon}{4} \right)^{(L-1)/2} \left(\frac{2}{3} \right)^L \left(\frac{1}{2} \right)^{G(\mathbb{N})} f_j^*(l^*)^{L-2} \prod_{l \in S_L} f_j^*(l)^{G(l)}
$$
\n
$$
\left(\left(\min \left(\sum_{m \in S_L^c} f_j^*(m) + \sqrt{\frac{\epsilon}{16}} f_j^*(l^*), \epsilon \right) \right)^{G(S_L^c)}
$$
\n
$$
- \left(\max \left(\exp \left(-\sqrt{\epsilon / \left(16 \max_{1 \le i \le k} \sum_{l \in S_L^c} f_i^*(l) \right)} \right), \sum_{m \in S_L^c} f_j^*(m) - \sqrt{\frac{\epsilon}{16}} f_j^*(l^*) \right) \right)^{G(S_L^c)} \right)
$$
\n
$$
\ge \left(\sqrt{\frac{\epsilon}{4}} \right)^{L-1+G(\mathbb{N})} \left(\frac{1}{3} \right)^L f_j^*(l^*)^{L-2} \prod_{l \in S_L} G(l) f_j^*(l)^{G(l)}.
$$
\n(85)

using that for all $0 < a < 1$,

$$
\frac{1}{a} \le \Gamma(a) \le \frac{2}{a}.\tag{86}
$$

and that under Assumption [\(78\)](#page-26-2), for ϵ small enough

$$
\exp\left(-\frac{\epsilon^{-\delta/2}}{4}\right) \ge \exp\left(-\sqrt{\frac{\epsilon}{16\max_{1\le i\le k}\sum_{l\in S_L^c}f_i^*(l)}}\right) \ge 0 \ge \sum_{m\in S_L^c}f_j^*(m) - \sqrt{\frac{\epsilon}{16}}f_j^*(l^*).
$$

We now control the last term of Equation [\(79\)](#page-27-0). Using Markov's inequality,

$$
DP(G|_{S_L^c}) \left(\sum_{l \in S_l^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) < \frac{\epsilon}{16}, \ \forall 1 \le i \le k \right)
$$
\n
$$
\ge 1 - DP(G|_{S_L^c}) \left(\sum_{i=1}^k \sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) > \frac{\epsilon}{16} \right)
$$
\n
$$
\ge 1 - \frac{\mathbb{E}^{\theta^*} \left(\sum_{i=1}^k \sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) \right)}{\frac{\epsilon}{16}}.
$$
\n
$$
(87)
$$

As $f(l)/\sum_{m\in S_L^c} f(m)$ is distributed from $\beta\big(G(l), G(S_L^c \setminus \{l\})\big),$

$$
\mathbb{E}^{\theta^*} \left[\log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) \right]
$$
\n
$$
= \frac{\Gamma(G(S_L^c))}{\Gamma(G(l))\Gamma(G(S_L^c \setminus \{l\}))} \left(\underbrace{\int_0^{1/2} \log^2(x) x^{G(l)-1} (1-x)^{G(S_L^c \setminus \{l\})-1} \lambda(dx)}_{I_1} + \underbrace{\int_1^1 \log^2(x) x^{G(l)-1} (1-x)^{G(S_L^c \setminus \{l\})-1} \lambda(dx)}_{I_2} \right),
$$
\n(88)

with

$$
I_1 \le 2 \int_0^{1/2} \log^2(x) x^{G(l)-1} \lambda(dx) = \frac{4 \log^2(2)(1/2)^{G(l)}}{G(l)^3} \left(\frac{G(l)^2}{2} + \frac{G(l)}{\log 2} + \frac{1}{\log^2 2} \right)
$$

$$
\le \frac{12 \log^2(2)(G(\mathbb{N}))^2}{G(l)^3},
$$
 (89)

and

$$
I_2 \le 2\log^2(2)\int_{1/2}^1 (1-x)^{G(S_L^c \setminus \{l\})-1} \lambda(dx) \le \frac{2\log^2(2)}{G(S_L^c \setminus \{l\})}.
$$
\n(90)

Combining Equations [\(86\)](#page-29-0) [\(88\)](#page-29-1), [\(89\)](#page-29-2) and [\(90\)](#page-29-3), we obtain

$$
\mathbb{E}^{\theta^*} \left(\sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) \right) \leq 24 G(\mathbb{N})^2 \log^2(2) \sum_{l \in S_L^c} \frac{f_i^*(l)}{(G(l))^2}.
$$
 (91)

Then using Assumption [\(77\)](#page-26-1) and Equations [\(87\)](#page-29-4) and [\(91\)](#page-29-5)

$$
DP(G|_{S_L^c}) \bigg(\sum_{l \in S_L^c} f_i^*(l) \log^2 \left(\frac{f(l)}{\sum_{m \in S_L^c} f(m)} \right) < \frac{\epsilon}{16k}, \ \forall 1 \le i \le k \bigg) \ge 1/2 \tag{92}
$$

Lemma [C.2](#page-26-0) follows combining Equations [\(79\)](#page-27-0), [\(85\)](#page-28-4) and [\(92\)](#page-29-6).

 \Box

D Proof of Theorem [4.3](#page-13-2) (Dirichlet process mixtures of Gaussian distributions)

Proof of Theorem [4.3](#page-13-2). Let $\sigma_n = \tilde{\epsilon}_n/(\log(1/\tilde{\epsilon}_n))$, $\tilde{\epsilon}_n = n^{-\beta/(2\beta+1)}(\log n)^{t_0}$. Following the computations of the proof of Theorem 4 of [\[Shen et al., 2013\]](#page-35-15) and using Assumption [\(S3\),](#page-13-5) Lemma [D.1](#page-30-1) ensures that Assumption [\(A\)](#page-5-2) holds with $t_0 \geq (2 + 2/\gamma_0 + 1)/(1/\beta + 2)$. Using Assumption [\(\(Q0\)\)](#page-9-2), Assumption [\(C\)](#page-6-1) holds. Using Theorem 5 of [\[Shen et al., 2013\]](#page-35-15), Assumptions [\(G1\),](#page-12-5) [\(S1\)](#page-12-6) and [\(S2\);](#page-13-4) Assumption [\(B\)](#page-5-3) holds with $\epsilon_n = n^{-\beta/(2\beta+1)}(\log(n))$ ^t, $t > t_0$. This concludes the proof of Theorem [4.3.](#page-13-2) \Box

The following lemma is a generalization of Lemma 4 of [\[Kruijer et al., 2010\]](#page-35-14) in the HMM context. In other words, we give a set of density functions $(f_i)_{1\leq i\leq k}$ satisfying Assumptions $(A.1)$ – $(A.6)$ in Lemma [D.1.](#page-30-1)

Lemma D.1. Assume that there exist β , L and γ such that for all $1 \leq j \leq k$, $f_j^* \in$ $\mathcal{P}(\beta, L, \gamma)$ and Assumptions [\(T1\)–](#page-12-2)[\(T3\)](#page-12-3) hold. Let σ be a positive real small enough.

Then for all $1 \leq j \leq k$, there exists a discrete measure $m_j = \sum_{i=1}^{N_j} \mu_j^i \delta_{z_j^i}$ supported on $\{x : f_j^*(x) \geq K_j \sigma^{2\beta + H_1}\}\$ with $H_1 > 2\beta$, K_j a constant small enough and $N_j =$ $O(\sigma^{-1}|\log \sigma|^{2/\gamma_0})$ such that Assumptions [\(A.1\)](#page-5-4)–[\(A.6\)](#page-5-5) hold with $f_j = \phi_{\sigma} * m_j$ for all $1 \leq$ $j \leq k$ and $\sigma^{2\beta} \leq \tilde{\epsilon}_n^2/u_n$.

Assumptions [\(A.1\)](#page-5-4)–[\(A.6\)](#page-5-5) also hold with $f_j = \phi_{\tilde{\sigma}} * \tilde{m}_j$, for all $\tilde{\sigma} \in [\sigma, \sigma + \sigma^{\delta'H_1+2}]$ and for all $\tilde{m}_j = \sum_{i=1}^{+\infty} \tilde{\mu}_j^i \delta_{\tilde{z}_j^i}$ such that $\tilde{\mu}_j^{1:N_j} \in \mathcal{B}(\mu_j, \sigma^{\delta'H_1+2})$ and $\tilde{z}_j^i \in \mathcal{B}(z_j^i, \sigma^{\delta'H_1+2})$, for all $1 \leq i \leq N_j$, where $\delta' \geq 1 + \beta/H_1$.

Proof of Lemma [D.1.](#page-30-1) The proof of Lemma [D.1](#page-30-1) is based on [\[Kruijer et al., 2010\]](#page-35-14). First notice that $f_j^* \in \mathcal{P}(\beta, L, \gamma)$ implies that for all integer $m \leq \beta$, $|l_m^j|$ is bounded by a polynomial. Then Assumption [\(T1\)](#page-12-2) implies that Assumption (C2) of [\[Kruijer et al.,](#page-35-14) [2010\]](#page-35-14) holds and stronglier implies that there exists $\delta > 0$ such that for all $1 \leq i, j, i \leq k$ and all integer $m \leq \beta$,

$$
\int |l_m^j(x)|^{(2\beta+\delta)/m} f_i^*(x)\lambda(dx) < \infty,
$$
\n
$$
\int |L^j(x)|^{2+\delta/\beta} f_i^*(x)\lambda(dx) < \infty,
$$
\n
$$
\int |l_m^j(x)|^{(2\beta+\delta)/m} f_i^*(x) \log f_i^*(x)\lambda(dx) < \infty,
$$
\n
$$
\int |L^j(x)|^{2+\delta/\beta} f_i^*(x) \log f_i^*(x)\lambda(dx) < \infty.
$$
\n
$$
(93)
$$

Let $\sigma > 0$, we consider

$$
S = \bigcap_{j=1}^{k} \left(A_{\sigma}^{j} \cap E_{\sigma}^{j} \right),
$$

where

$$
A^j_{\sigma} = \left\{ x : \left| l^j_m(x) \right| \leq B\sigma^{-m} |\log(\sigma)|^{-m/2}, \forall 1 \leq m \leq \lfloor \beta \rfloor, \left| L^j(x) \right| \leq B\sigma^{-\beta} |\log(\sigma)|^{-\beta/2} \right\}
$$

and

$$
E^j_{\sigma} = \left\{ x \ : \ f^*_j(x) \ge \sigma^{H_1} \right\}.
$$

Using Assumptions [17,](#page-12-7) [\(T1\),](#page-12-2) [\(T3\)](#page-12-3) and Lemma 2 of [\[Kruijer et al., 2010\]](#page-35-14), there exists k density functions h_{β}^{j} β_{β} such that for all $1 \leq j \leq k$ and all $x \in S$,

$$
h_{\beta}^{j} * \phi_{\sigma}(x) = f_{j}^{*}(x) \left(1 + O(R^{j}(x)\sigma^{\beta}) \right) + O((1 + R^{j}(x))\sigma^{H})
$$
\n(94)

where R^j is defined as in Equation (16) page 1232 of [\[Kruijer et al., 2010\]](#page-35-14) and H is as large as we want. Using Assumptions [17,](#page-12-7) [\(T1\)](#page-12-2) and [\(T3\),](#page-12-3) the proof of Lemma 2 of [\[Kruijer](#page-35-14) [et al., 2010\]](#page-35-14) is easily generalizable in this context so that

$$
\int_{S^c} h_\beta^j * \phi_\sigma(y) \lambda(dy) \lesssim \sigma^{2\beta}.
$$
\n(95)

The generalization can be proved using Equation [\(93\)](#page-30-2) and by replacing Equation (56) of [\[Kruijer et al., 2010\]](#page-35-14) by Equation [\(102\)](#page-32-0).

As at page 1251 in [\[Kruijer et al., 2010\]](#page-35-14), we denote

$$
\tilde{h}^{\beta}_j = \frac{1_{x:h^{\beta}_j(x) \geq \sigma^{H_2}} h^{\beta}_j}{\int_{x:h^{\beta}_j(x) \geq \sigma^{H_2}} h^{\beta}_j d\lambda}
$$

and using Lemma 12 of [\[Kruijer et al., 2010\]](#page-35-14), for all $1 \leq j \leq k$, there exist k discrete distributions $m_j = \sum_{i=1}^{N_j} \mu_j^i \delta_{z_j^i}$ supported in $\{x : f_j^*(x) \ge K_j \sigma^{2\beta + H_1}\}\$ for $H_1 > 2\beta$, a constant K_j small enough and with $N_j = O(\sigma^{-1} |\log \sigma|^{2/\gamma_0})$ such that

$$
\|\tilde{h}_j^{\beta} * \phi_{\sigma} - m_j * \phi_{\sigma}\|_{\infty} \leq \sigma^{-1} e^{-C |\log \sigma|^{2/\gamma_0}},
$$

$$
\|\tilde{h}_j^{\beta} * \phi_{\sigma} - m_j * \phi_{\sigma}\|_1 \leq \sigma^{-1} e^{-C' |\log \sigma|^{2/\gamma_0}}
$$
\n(96)

for any C', C large enough.

It is now sufficient to prove that Assumptions [\(A.1\)](#page-5-4) to [\(A.6\)](#page-5-5) hold with $\epsilon_n^{4/(2-\alpha)}$ = $O(\sigma^{2\beta})$, $f_j = \tilde{m}_j * \phi_{\tilde{\sigma}}$ and $\tilde{f}_j = h_j^{\beta}$ $\phi_j^{\beta} * \phi_{\tilde{\sigma}}$ for all $\tilde{\sigma} \in [\sigma, \sigma + \sigma^{\delta'H_1+2}]$ and all discrete distributions $\tilde{m}_j = \sum_{i=1}^{N_j} \tilde{\mu}_j^i \delta_{\tilde{z}_j^i}$ such that $\tilde{\mu}_j \in \mathcal{B}(\mu_j, \sigma^{\delta'H_1+2}) \cap \Delta_{N_j}$ and $\tilde{z}_j \in \mathcal{B}(z_j, \sigma^{\delta'H_1+2})$ where $\delta' \geq 1 + \beta/H_1$. By Lemma 2 of [\[Kruijer et al., 2010\]](#page-35-14),

$$
||f_j - m_j * \phi_\sigma||_1 \lesssim \sigma^{H_1 + 1 + \beta}, \quad ||f_j - m_j * \phi_\sigma||_\infty \lesssim \sigma^{H_1 + \beta}, \tag{97}
$$

• Proof of $(A,1)$. We cut the integral as in the following.

$$
\int f_i^*(y) \log^2 \frac{f_j^*(y)}{f_j(y)} \lambda(dy) \le \int_S f_i^*(y) \log^2 \frac{f_j^*(y)}{\tilde{f}_j(y)} \lambda(dy)
$$
\n
$$
+ \int_{S^c} f_i^*(y) \log^2 \frac{f_j^*(y)}{\tilde{f}_j(y)} \lambda(dy)
$$
\n
$$
+ \int f_i^*(y) \log^2 \frac{\tilde{f}_j(y)}{f_j(y)} \lambda(dy).
$$
\n(98)

The last integral can be controlled by $O(\sigma^{2\beta})$ as in the proof of Lemma 4 of [\[Kruijer](#page-35-14)] [et al., 2010\]](#page-35-14).

Using Equation [\(94\)](#page-30-3) we control the first integral of the bound of Equation [\(98\)](#page-31-0):

$$
\int_S f_i^*(y) \log^2 \frac{f_j^*(y)}{\tilde{f}_j(y)} \lambda(dy) \le \int_S f_i^*(y) \frac{|f_j^*(y) - \tilde{f}_j(y)|^2}{\min(\tilde{f}_j(y), f_j^*(y))} \lambda(dy) \lesssim \sigma^{2\beta}
$$

as soon as H is large enough, using Equation [\(94\)](#page-30-3).

Using that $f_j^* \lesssim \tilde{f}_j \lesssim \phi_\sigma * f_j^* \lesssim 1$ (see Remark 1 and the bottom of page 1252 of [\[Kruijer et al., 2010\]](#page-35-14)), we control the second integral in the bound of Equation [\(98\)](#page-31-0),

$$
\int_{S^c} f_i^*(y) \log^2 \frac{f_j^*(y)}{\tilde{f}_j(y)} \lambda(dy) \lesssim \int_{S^c} f_i^*(y) \lambda(dy) + \int_{S^c} f_i^*(y) \log^2 \left(f_j^*(y)\right) \lambda(dy)
$$

which is bounded by $O(\sigma^{2\beta})$ following the proof of [\(A.4\)](#page-5-8).

• Proof of $(A.2)$. We cut the integral into three parts:

$$
\int_{S} \frac{|f_i^*(y) - f_i(y)|^2}{f_i^*(y)} \lambda(dy)
$$
\n
$$
\lesssim \underbrace{\int_{S} \frac{|f_i^*(y) - h_{\beta}^i * \phi_{\sigma}(y)|^2}{f_i^*(y)} \lambda(dy)}_{I_1} + \underbrace{\int_{S} \frac{|h_{\beta}^i * \phi_{\sigma}(y) - \tilde{h}_{\beta}^i * \phi_{\sigma}(y)|^2}{f_i^*(y)} \lambda(dy)}_{I_3} + \underbrace{\int_{S} \frac{|\tilde{h}_{\beta}^i * \phi_{\sigma}(y) - f_i(y)|^2}{f_i^*(y)} \lambda(dy)}_{I_3}
$$

Using Equation [\(94\)](#page-30-3),

$$
I_1 \lesssim \sigma^{2\beta}.\tag{99}
$$

We now control I_2 using the bound

$$
\left| \frac{h_{\beta}^{i} * \phi_{\sigma}(y)}{\tilde{h}_{\beta}^{i} * \phi_{\sigma}(y)} - 1 \right| = O(\sigma^{2\beta}), \text{ for all } y \in S
$$

of page 1252 of [\[Kruijer et al., 2010\]](#page-35-14) and Equation [\(94\)](#page-30-3). Then

$$
I_2 = \int_S \left(\frac{|h_\beta^i * \phi_\sigma(y) - \tilde{h}_\beta^i * \phi_\sigma(y)|}{h_\beta^i * \phi_\sigma(y)} \right)^2 h_\beta^i * \phi_\sigma(y) \frac{h_\beta^i * \phi_\sigma(y)}{f_i^*(y)} \lambda(dy)
$$

$$
\leq \int_S (O(\sigma^{2\beta}))^2 h_\beta^i * \phi_\sigma(y) 2 \lambda(dy) \lesssim \sigma^{2\beta},
$$
 (100)

using Equation [\(94\)](#page-30-3). As to I_3 , using Equations [\(96\)](#page-31-1) and [\(97\)](#page-31-2), it is upper-bounded by

$$
2\frac{\|\tilde{h}_i^{\beta} * \phi_\sigma - m_i * \phi_\sigma\|_\infty \|\tilde{h}_i^{\beta} * \phi_\sigma - m_i * \phi_\sigma\|_1}{\sigma^{H_1}} + 2\frac{\|f_i - m_i * \phi_\sigma\|_\infty \|f_i - m_i * \phi_\sigma\|_1}{\sigma^{H_1}} \lesssim \sigma^{2\beta}
$$
\n(101)

when $2 > \gamma_0$, such a γ_0 can always be chosen (see the first line of page 1253 of [\[Kruijer et al., 2010\]](#page-35-14)).

- Proof of $(A.3)$. Assumption $(A.3)$ is proved in Equation [\(95\)](#page-31-3).
- Proof of $(A.4)$. It is sufficient to bound

$$
\int_{(E^i_{\sigma})^c} f^*_j(y) \lambda(dy)
$$

and

$$
\int_{(A^i_\sigma)^c} f^*_j(y) \lambda(dy).
$$

Using Assumption [\(T2\),](#page-12-4) for all $0 < \delta < 1$

$$
\int_{(E_{\sigma}^{i})^{c}} f_{j}^{*}(y) \lambda(dy)
$$
\n
$$
\leq \int_{\{y:f_{j}^{*}(y) < \sigma^{H_{1}} M_{j,i} \exp(\tau_{j,i}|y|^{\gamma_{j,i}})\}} f_{j}^{*}(y) \lambda(dy)
$$
\n
$$
\leq \int_{\{y:f_{j}^{*}(y) < \sigma^{H_{1}} M_{j,i} \exp(\tau_{j,i}|y|^{\gamma_{j,i}})\}} (f_{j}^{*}(y))^{1/\delta} (f_{j}^{*}(y))^{1-1/\delta} \lambda(dy)
$$
\n
$$
\leq \sigma^{H_{1}/\delta} \int (M_{j,i})^{1/\delta} \exp(\tau_{j,i}|y|^{\gamma_{j,i}}/\delta) (f_{j}^{*}(y))^{1-1/\delta} \lambda(dy) \lesssim \sigma^{2\beta}
$$
\n
$$
(102)
$$

as soon as $H_1 > 2\beta$, using Assumption [\(T1\).](#page-12-2) Moreover using [\(93\)](#page-30-2) and Markov inequality, as in the proof of Lemma 2 of [\[Kruijer et al., 2010\]](#page-35-14),

$$
\int_{(A^i_{\sigma})^c} f_j^*(y) \lambda(dy) \lesssim \sigma^{2\beta}.
$$
 (103)

• Proof of $(A.5)$. Using the same argument as in the bottom of the page 1252 of [\[Kruijer et al., 2010\]](#page-35-14), Equations [\(96\)](#page-31-1) and [\(97\)](#page-31-2),

$$
\int_{S} f_{i}^{*}(y) \max_{1 \leq j \leq k} \log \left(\frac{\tilde{f}_{j}(y)}{f_{j}(y)} \right) \lambda(dy)
$$
\n
$$
\leq \int_{S} f_{i}^{*}(y) \max_{1 \leq j \leq k} \frac{|\tilde{f}_{j}(y) - f_{j}(y)|}{f_{j}(y)} \lambda(dy)
$$
\n
$$
\leq \int_{S} f_{i}^{*}(y) \max_{1 \leq j \leq k} \frac{\|\tilde{f}_{j} - f_{j}\|_{\infty}}{\sigma^{H_{2}} - \|\tilde{f}_{j} - f_{j}\|_{\infty}} \lambda(dy) \lesssim \sigma^{2\beta}
$$
\n(104)

• Proof of [\(A.6\)](#page-5-5). Using that $f_j^* \lesssim \tilde{f}_j$ (see Assumption (C3) of [\[Kruijer et al., 2010\]](#page-35-14)) Equation [\(99\)](#page-32-1) implies [\(A.6\)](#page-5-5).

 \Box

References

- Grigory Alexandrovich and Hajo Holzmann. Nonparametric identification of hidden Markov models. *arXiv preprint arXiv:1404.4210*, 2014.
- Julyan Arbel, Ghislaine Gayraud, and Judith Rousseau. Bayesian optimal adaptive estimation using a sieve prior. Scand. J. Stat., 40(3):549–570, 2013.
- Fadoua Balabdaoui, Hanna Jankowski, Kaspar Rufibach, and Marios Pavlides. Asymptotics of the discrete log-concave maximum likelihood estimator and related applications. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 75 (4):769–790, 2013.
- Leonard E. Baum and Ted Petrie. Statistical inference for probabilistic functions of finite state Markov chains. The Annals of Mathematical Statistics, 37(6):1554–1563, 1966.
- Matthew Beal and Praveen Krishnamurthy. Gene expression time course clustering with countably infinite hidden Markov models. arXiv preprint arXiv:1206.6824, 2012.
- D. L. Borchers, W. Zucchini, M. P. Heide-Jørgensen, A. Cañadas, and R. Langrock. Using hidden Markov models to deal with availability bias on line transect surveys. Biometrics, 69(3):703–713, 2013.
- Stéphane Boucheron, Elisabeth Gassiat, et al. A Bernstein-von Mises theorem for discrete probability distributions. Electronic Journal of Statistics, 3:114–148, 2009.
- Antonio Canale and David B Dunson. Bayesian kernel mixtures for counts. Journal of the American Statistical Association, 106(496), 2011.
- O. Cappé, E. Moulines, and T. Rydén. *Inference in Hidden Markov Models*. Springer, 2005.
- Yohann De Castro, Elisabeth Gassiat, and Sylvain Le Corff. Consistent estimation of the filtering and marginal smoothing distributions in nonparametric hidden Markov models. arXiv preprint arXiv:1507.06510, 2015a.
- Yohann De Castro, Elisabeth Gassiat, and Claire Lacour. Minimax adaptive estimation of non-parametric hidden Markov models. arXiv preprint arXiv:1501.04787, 2015b.
- M. C. de Gunst and O. Shcherbakova. Asymptotic behavior of Bayes estimators for hidden Markov models with application to ion channels. Mathematical Methods of Statistics, 17(4):342–356, 2008.
- Persi Diaconis and David Freedman. On the consistency of Bayes estimates. The Annals of Statistics, pages 1–26, 1986.
- R. Douc and C. Matias. Asymptotics of the maximum likelihood estimator for general hidden Markov models. Bernoulli, 7:381–420, 2001.
- R. Douc, E. Moulines, and T. Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. The Annals of statistics, 32 (5):2254–2304, 2004.
- Cécile Durot, Sylvie Huet, François Koladjo, and Stéphane Robin. Least-squares estimation of a convex discrete distribution. Computational Statistics \mathcal{B} Data Analysis, 67: 282–298, 2013.
- Emily B Fox, Michael I Jordan, Erik B Sudderth, and Alan S Willsky. Sharing features among dynamical systems with beta processes. In Advances in Neural Information Processing Systems, pages 549–557, 2009.
- Emily B Fox, Erik B Sudderth, Michael I Jordan, and Alan S Willsky. A sticky HDP-HMM with application to speaker diarization. The Annals of Applied Statistics, pages 1020–1056, 2011.
- E. Gassiat and J. Rousseau. Non parametric finite translation hidden Markov models and extensions. Bernoulli, 2013. to appear.
- E. Gassiat, A. Cleynen, and S. Robin. Inference in finite state space non parametric hidden Markov models and applications. *Statistics and Computing*, pages 1–11, 2015.
- Elisabeth Gassiat and Judith Rousseau. About the posterior distribution in hidden Markov models with unknown number of states. Bernoulli, 20(4):2039–2075, 2014.
- Subhashis Ghosal and Aad van der Vaart. Posterior convergence rates of Dirichlet mixtures at smooth densities. Ann. Statist., 35(2):697–723, 2007a.
- Subhashis Ghosal and Aad van der Vaart. Convergence rates of posterior distributions for non-i.i.d. observations. Ann. Statist., 35(1):192–223, 2007b.
- Subhashis Ghosal, Jayanta K. Ghosh, and Aad W. van der Vaart. Convergence rates of posterior distributions. Ann. Statist., 28(2):500–531, 2000.
- J.K. Ghosh and R.V. Ramamoorthi. Bayesian Nonparametrics. Springer, 2003.
- Yanjun Han, Jiantao Jiao, and Tsachy Weissman. Minimax estimation of discrete distributions under ℓ_1 loss. CoRR, abs/1411.1467, 2014.
- Marc Hoffmann, Judith Rousseau, and Johannes Schmidt-Hieber. On adaptive posterior concentration rates. arXiv preprint arXiv:1305.5270, 2013.
- Hanna K Jankowski and Jon A Wellner. Estimation of a discrete monotone distribution. Electronic journal of statistics, 3:1567, 2009.
- Markus Jochmann. Modeling US inflation dynamics: A Bayesian nonparametric approach. Econometric Reviews, 34(5):537–558, 2015.
- Willem Kruijer, Judith Rousseau, and Aad van der Vaart. Adaptive Bayesian density estimation with location-scale mixtures. Electronic Journal of Statistics, 4:1225–1257, 2010.
- Roland Langrock, Thomas Kneib, Alexander Sohn, and Stacy L DeRuiter. Nonparametric inference in hidden Markov models using p-splines. Biometrics, 2015.
- Fabrice Lefèvre. Non-parametric probability estimation for HMM-based automatic speech recognition. Computer Speech & Language, $17(2)$:113-136, 2003.
- Scott W Linderman, Matthew J Johnson, Matthew A Wilson, and Zhe Chen. A nonparametric Bayesian approach to uncovering rat hippocampal population codes during spatial navigation. *arXiv preprint arXiv:1411.7706*, 2014.
- I. L. MacDonald and W. Zucchini. Hidden Markov and other models for discrete-valued time series. Chapman and Hall/CRC, London, UK, 1997.
- I. L. MacDonald and W. Zucchini. Hidden Markov models for time series: an introduction using R. Chapman and Hall/CRC, London, UK, 2009.
- Emmanuel Rio. Inégalités de Hoeffding pour les fonctions lipschitziennes de suites dépendantes. Comptes Rendus de l'Académie des Sciences-Series I-Mathematics, 330 (10):905–908, 2000.
- Vincent Rivoirard and Judith Rousseau. Posterior concentration rates for infinite dimensional exponential families. Bayesian Anal., 7(2):311–333, 2012.
- Judith Rousseau. On the frequentist properties of Bayesian nonparametric methods. Annual statistical reviews, 2015. to appear.
- Weining Shen, Surya T. Tokdar, and Subhashis Ghosal. Adaptive Bayesian multivariate density estimation with Dirichlet mixtures. Biometrika, 100(3):623–640, 2013.
- A. W. van der Vaart and J. H. van Zanten. Adaptive Bayesian estimation using a Gaussian random field with inverse gamma bandwidth. Ann. Statist., 37(5B):2655–2675, 2009.
- Elodie Vernet et al. Posterior consistency for nonparametric hidden Markov models with finite state space. Electronic Journal of Statistics, 9:717–752, 2015.
- Julian Peter Whiting, Martin Francis Lambert, and Andrew Viggo Metcalfe. Modelling persistence in annual australian point rainfall. Hydrology and Earth System Sciences, 7(2):197–211, 2003.
- C. Yau, O. Papaspiliopoulos, G.O. Roberts, and C. Holmes. Bayesian non-parametric hidden Markov models with applications in genomics. Journal of the Royal Statistical Society, 73:37–57, 2011.