

Exact two-body solutions and Quantum defect theory of two dimensional dipolar quantum gas

Jianwen Jie and Ran Qi*

Department of Physics, Renmin University of China, Beijing 100872, P. R. China

(Dated: November 12, 2018)

In this paper, we provide the two-body exact solutions of two dimensional (2D) Schrödinger equation with isotropic $\pm 1/r^3$ interactions. Analytic quantum defect theory are constructed base on these solutions and are applied to investigate the scattering properties as well as two-body bound states of ultracold polar molecules confined in a quasi-2D geometry. Interestingly, we find that for the attractive case, the scattering resonance happens simultaneously in all partial waves which has not been observed in other systems. The effect of this feature on the scattering phase shift across such resonances is also illustrated.

I. INTRODUCTION

In recent years, great progress has been made in the production of ultracold polar molecules thanks to the development of stimulated Raman adiabatic passage (STIRAP) technique [1, 2]. Extensive experimental and theoretical efforts have been devoted to exploring the production of dipolar gases and to revealing exotic quantum phases resulting from the anisotropic nature and long-range character of the dipolar interaction [3–8]. In three dimensions, the system is usually suffered from large chemical reaction rate leading to a very short life time [9–11]. The reaction rate, however, can be greatly suppressed by confining the system to lower dimensionality and makes it possible to realize a stable quantum many-body system [12]. As a result, a lot of efforts have been made in order to understand both two-body and many-body properties of dipolar quantum gases in two or one dimensional geometries [4, 13–25]. However, up to now, few analytical results are known even in the simple two-body problems due to the existence of long range $1/r^3$ tail in the interaction potential.

In this paper, following a similar method developed for the three dimensional scattering problem under repulsive $1/r^3$ and attractive $1/r^6$ interaction [26], we present exact solutions, in the form of a generalized Neumann expansion, for the 2D quantum scattering problem with either repulsive or attractive isotropic $1/r^3$ interactions. Both cases are relevant for current polar molecule experimental setups [14, 20]. With the help of these exact solutions, we will construct the analytic quantum defect theory (QDT) [27–30] for realistic dipolar scattering in the quasi-2D confinement and investigate the corresponding scattering properties as well as the two-body bound states.

We start in Sec. II by summarizing the explicit form of our exact solutions and also give the long range and short range asymptotic behavior analytically. In Sec. III, to be more self contained, we briefly review the general scat-

tering formula in 2D and provide the definition of phase shift and cross sections. In Sec. IV, we first introduce the experimental setups under our consideration for two different kinds of dipole-dipole scattering in quasi-2D geometry. Then we construct the analytic QDT for these two cases and present our results on the scattering properties. Finally, we conclude ourselves in Sec. V.

II. SOLUTIONS OF THE SCHRÖDINGER EQUATION

We consider the Schrödinger equation for $\pm 1/r^3$ type potentials in two spacial dimensions satisfied by the radial wave function $\psi_m(k, r)$:

$$\left[-\frac{d^2}{dr^2} + \frac{m^2 - 1/4}{r^2} \pm \frac{D}{r^3} - \bar{\varepsilon} \right] u_m(r) = 0, \quad (1)$$

where $u_m(r) = \psi_m(k, r)/\sqrt{r}$, $D = \mu d^2/\hbar^2$ is the dipolar length with μ and d being the reduced mass and dipole moment, $\bar{\varepsilon} = 2\mu\varepsilon/\hbar^2 = k^2$ with ε and k the scattering energy and wave number, and $m\hbar$ is angular momentum. The solutions for the three dimensional version of this type of equation are already provide in the repulsive case [26]. We find that the method used in [26] can be easily generalized to two spacial dimensions as well as to the attractive case and we will present the explicit form of our exact solutions to Eq. (1) in the following part of this section.

We found that there exists a pair of linearly independent solutions with energy-independent asymptotic behaviors near the origin ($r \ll D$). The explicit form can be written as

$$\bar{u}_{\varepsilon m}^{+1}(r) = \frac{G_{\varepsilon m}^{-1}(-\nu)\xi_{\varepsilon m}^+(r) - G_{\varepsilon m}^{-1}(\nu)\eta_{\varepsilon m}^+(r)}{\sin 2\pi\nu}, \quad (2)$$

$$\bar{u}_{\varepsilon m}^{+2}(r) = -G_{\varepsilon m}^{-1}(-\nu)\xi_{\varepsilon m}^+(r) - G_{\varepsilon m}^{-1}(\nu)\eta_{\varepsilon m}^+(r), \quad (3)$$

$$\bar{u}_{\varepsilon m}^{-1}(r) = \frac{-G_{\varepsilon m}^{-1}(-\nu)\xi_{\varepsilon m}^-(r) + G_{\varepsilon m}^{-1}(\nu)\eta_{\varepsilon m}^-(r)}{2 \sin \pi\nu}, \quad (4)$$

$$\bar{u}_{\varepsilon m}^{-2}(r) = \frac{G_{\varepsilon m}^{-1}(-\nu)\xi_{\varepsilon m}^-(r) + G_{\varepsilon m}^{-1}(\nu)\eta_{\varepsilon m}^-(r)}{2 \cos \pi\nu}. \quad (5)$$

Here and below the plus and minus signs on the superscripts correspond to the repulsive and attractive inter-

*Electronic address: qiran@ruc.edu.cn

actions, respectively. Functions $\xi_{\varepsilon m}^{\pm}(r)$ and $\eta_{\varepsilon m}^{\pm}(r)$ in (2)-(5) are another pair of linearly independent solutions that takes the form of a generalized Neumann expansion:

$$\xi_{\varepsilon m}^{\pm}(r) = \sum_{n=-\infty}^{\infty} r^{1/2} b_n^{\pm} J_{\nu+n}(kr), \quad (6)$$

$$\eta_{\varepsilon m}^{\pm}(r) = \sum_{n=-\infty}^{\infty} (-1)^n b_n^{\pm} r^{1/2} J_{-\nu-n}(kr), \quad (7)$$

where

$$b_j^{\pm} = (\pm\Delta)^j \frac{\Gamma(\nu)\Gamma(\nu-m+1)\Gamma(\nu+m+1)}{\Gamma(\nu+j)\Gamma(\nu-m+j+1)\Gamma(\nu+m+j+1)} c_j(\nu), \quad (8)$$

$$b_{-j}^{\pm} = (\pm\Delta)^j \frac{\Gamma(\nu-j+1)\Gamma(\nu-m-j)\Gamma(\nu+m-j)}{\Gamma(\nu+1)\Gamma(\nu-m)\Gamma(\nu+m)} c_j(-\nu), \quad (9)$$

with j being a positive integer, $\Delta = kD/2$ and

$$c_j(\nu) = Q(\nu+j-1)Q(\nu+j-2)\cdots Q(\nu)b_0. \quad (10)$$

The coefficient b_0 is a normalization constant which can be set to 1, and $Q(\nu)$ is given by a continued fraction

$$Q(\nu) = \frac{1}{1 - \varepsilon_s \frac{1}{\frac{Q(\nu+1)}{(\nu+1)[(\nu+1)^2 - m^2]} \frac{1}{(\nu+2)[(\nu+2)^2 - m^2]}}, \quad (11)$$

where ε_s is a scaled energy defined as

$$\varepsilon_s \equiv \Delta^2 = \frac{1}{4} \frac{\varepsilon}{(\hbar^2/2\mu)(1/D^2)}. \quad (12)$$

The $G_{\varepsilon m}(\nu)$ function in (2)-(5) is defined as

$$G_{\varepsilon m}(\nu) = \Delta^{-\nu} \frac{\Gamma(1+m+\nu)\Gamma(1-m+\nu)}{\Gamma(1-\nu)} C(\nu), \quad (13)$$

where $C(\nu) = \lim_{j \rightarrow \infty} c_j(\nu)$.

Finally, ν is a root of a characteristic function

$$\Lambda_m(\nu, \varepsilon_s) \equiv (\nu^2 - m^2) - \frac{\varepsilon_s}{\nu} [\bar{Q}(\nu) - \bar{Q}(-\nu)], \quad (14)$$

where $\bar{Q}(\nu)$ is defined as

$$\bar{Q}(\nu) = \frac{Q(\nu)}{(\nu+1)[(\nu+1)^2 - m^2]}. \quad (15)$$

The solution of ν for $\Lambda_m(\nu, \varepsilon_s) = 0$ could either be real or complex depending on the scattering energy and angular momentum.

The pair of solutions $\bar{u}_{\varepsilon m}^{\pm 1}$ and $\bar{u}_{\varepsilon m}^{\pm 2}$ have been defined in such a way that they have energy-independent behavior

near the origin ($r \ll D$), which are given as

$$\bar{u}_{\varepsilon m}^{+1}(r) \rightarrow r^{3/4} \sqrt{\frac{1}{\pi\sqrt{D}}} e^{-2\sqrt{\frac{E}{r}}}, \quad (16)$$

$$\bar{u}_{\varepsilon m}^{+2}(r) \rightarrow -r^{3/4} \sqrt{\frac{1}{\pi\sqrt{D}}} e^{2\sqrt{\frac{E}{r}}}, \quad (17)$$

$$\bar{u}_{\varepsilon m}^{-1}(r) \rightarrow r^{3/4} \sqrt{\frac{1}{\pi\sqrt{D}}} \sin\left(2\sqrt{\frac{D}{r}} - \frac{\pi}{4}\right), \quad (18)$$

$$\bar{u}_{\varepsilon m}^{-2}(r) \rightarrow r^{3/4} \sqrt{\frac{1}{\pi\sqrt{D}}} \cos\left(2\sqrt{\frac{D}{r}} - \frac{\pi}{4}\right), \quad (19)$$

for both positive and negative energies. Note that the solution $\bar{u}_{\varepsilon m}^{+1}$ approaches zero exponentially in the limit $r \rightarrow 0$ and thus is the physical solution for pure repulsive $1/r^3$ interaction.

For positive scattering energy $\varepsilon > 0$, the asymptotic behaviors of $\bar{u}_{\varepsilon m}^{\pm 1}$, $\bar{u}_{\varepsilon m}^{\pm 2}$ as $r \rightarrow \infty$ are given as

$$\bar{u}_{\varepsilon m}^{\pm 1}(r) \rightarrow \frac{1}{\sqrt{2\pi k}} \left[Z_{11}^{\pm} \sin\left(kr - \frac{l\pi}{2}\right) - Z_{12}^{\pm} \cos\left(kr - \frac{l\pi}{2}\right) \right], \quad (20)$$

$$\bar{u}_{\varepsilon m}^{\pm 2}(r) \rightarrow \frac{1}{\sqrt{2\pi k}} \left[Z_{21}^{\pm} \sin\left(kr - \frac{l\pi}{2}\right) - Z_{22}^{\pm} \cos\left(kr - \frac{l\pi}{2}\right) \right], \quad (21)$$

where $l = m - 1/2$. The matrix Z_{ij} are dimensionless functions of m and scaled energy ε_s which can be obtained analytically as

$$Z_{11}^{+} = \frac{2}{\sin 2\nu\pi} \left[\frac{\alpha_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} - \frac{\alpha_{\varepsilon m} \cos \pi\nu + \beta_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (22)$$

$$Z_{12}^{+} = \frac{2}{\sin 2\nu\pi} \left[\frac{\beta_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} - \frac{\beta_{\varepsilon m} \cos \pi\nu - \alpha_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (23)$$

$$Z_{21}^{+} = -2 \left[\frac{\alpha_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\alpha_{\varepsilon m} \cos \pi\nu + \beta_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right] \quad (24)$$

$$Z_{22}^{+} = -2 \left[\frac{\beta_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\beta_{\varepsilon m} \cos \pi\nu - \alpha_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (25)$$

$$Z_{11}^{-} = \frac{1}{\sin \nu\pi} \left[\frac{-\alpha_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\alpha_{\varepsilon m} \cos \pi\nu + \beta_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (26)$$

$$Z_{12}^{-} = \frac{1}{\sin \nu\pi} \left[\frac{-\beta_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\beta_{\varepsilon m} \cos \pi\nu - \alpha_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (27)$$

$$Z_{21}^{-} = \frac{1}{\cos \nu\pi} \left[\frac{\alpha_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\alpha_{\varepsilon m} \cos \pi\nu + \beta_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (28)$$

$$Z_{22}^{-} = \frac{1}{\cos \nu\pi} \left[\frac{\beta_{\varepsilon m}}{G_{\varepsilon m}(-\nu)} + \frac{\beta_{\varepsilon m} \cos \pi\nu - \alpha_{\varepsilon m} \sin \pi\nu}{G_{\varepsilon m}(\nu)} \right], \quad (29)$$

where

$$\alpha_{\varepsilon m} = \cos[\pi(\nu - m)/2]X_{\varepsilon m} - \sin[\pi(\nu - m)/2]Y_{\varepsilon m}, \quad (30)$$

$$\beta_{\varepsilon m} = \sin[\pi(\nu - m)/2]X_{\varepsilon m} + \cos[\pi(\nu - m)/2]Y_{\varepsilon m}, \quad (31)$$

and $X_{\varepsilon m}$, $Y_{\varepsilon m}$ are defined as

$$X_{\varepsilon m} = \sum_{n=-\infty}^{\infty} (-1)^n b_{2n}, \quad (32)$$

$$Y_{\varepsilon m} = \sum_{n=-\infty}^{\infty} (-1)^n b_{2n+1}, \quad (33)$$

These dimensionless Z_{ij}^{\pm} functions are the key quantities to calculate the scattering phase shifts which will be clear later.

For negative energy $\varepsilon < 0$, $\bar{u}_{\varepsilon m}^{\pm 1}$, $\bar{u}_{\varepsilon m}^{\pm 2}$ have the following asymptotic behaviors as $r \rightarrow \infty$

$$\bar{u}_{\varepsilon m}^{\pm 1}(r) \rightarrow \frac{1}{\sqrt{2\pi\kappa}} \left(W_{11}^{\pm} e^{\kappa r} + W_{12}^{\pm} e^{-\kappa r} \right), \quad (34)$$

$$\bar{u}_{\varepsilon m}^{\pm 2}(r) \rightarrow \frac{1}{\sqrt{2\pi\kappa}} \left(W_{21}^{\pm} e^{\kappa r} + W_{22}^{\pm} e^{-\kappa r} \right), \quad (35)$$

where the $\kappa = \sqrt{-\varepsilon}$ and W_{ij} are also dimensionless functions of m and ε_s :

$$W_{11}^+ = -\frac{D_m}{\sin 2\pi\nu} \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} - \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (36)$$

$$W_{12}^+ = -\frac{E_m}{2 \cos \pi\nu} \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} + \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (37)$$

$$W_{21}^+ = -D_m \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} + \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (38)$$

$$W_{22}^+ = -E_m \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} - \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right] \sin \pi\nu, \quad (39)$$

$$W_{11}^- = \frac{D_m}{2 \sin \pi\nu} \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} - \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (40)$$

$$W_{12}^- = \frac{1}{2} E_m \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} + \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (41)$$

$$W_{21}^- = \frac{1}{2 \cos \pi\nu} D_m \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} + \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right], \quad (42)$$

$$W_{22}^- = \frac{1}{2} E_m \left[\frac{1}{i^\nu G_{\varepsilon m}(\nu)} - \frac{1}{i^{-\nu} G_{\varepsilon m}(-\nu)} \right] \tan \pi\nu, \quad (43)$$

where we have defined

$$D_m = \sum_{n=-\infty}^{\infty} i^n b_n, \quad (44)$$

$$E_m = \sum_{n=-\infty}^{\infty} (-i)^n b_n. \quad (45)$$

Another important function which is usually called the χ function is defined as

$$\chi_m^{\pm}(\varepsilon_s) = W_{11}^{\pm}/W_{21}^{\pm}. \quad (46)$$




FIG. 1: Potential curve (upper panel) and corresponding quantum defect K^c (lower panel) for two model potentials $V_+(r)$ (left collum) and $V_-(r)$ (right collum). $E_D = \hbar^2/(mD^2)$ is the characteristic energy scale associate with the dipole length D .

This function is useful in determining binding energy of the two-body bound state which will be clear in the following sections.

Finally, from the asymptotic behavior as $r \rightarrow 0$ given in (16)-(19), it is easy to show that the solution pairs have the Wronskian given by

$$W(\bar{u}_{\varepsilon m}^{\pm 1}, \bar{u}_{\varepsilon m}^{\pm 2}) = -\frac{2}{\pi}, \quad W(\bar{u}_{\varepsilon m}^{-1}, \bar{u}_{\varepsilon m}^{-2}) = \frac{1}{\pi}, \quad (47)$$

Since the Wronskian is a constant that is independent of r , the asymptotic forms of solutions at large r should give the same result, which requires

$$\det(Z^+) = Z_{11}^+ Z_{22}^+ - Z_{12}^+ Z_{21}^+ = 4, \quad (48)$$

$$\det(Z^-) = Z_{11}^- Z_{22}^- - Z_{12}^- Z_{21}^- = 2, \quad (49)$$

$$\det(W^+) = W_{11}^+ W_{22}^+ - W_{12}^+ W_{21}^+ = -2, \quad (50)$$

$$\det(W^-) = W_{11}^- W_{22}^- - W_{12}^- W_{21}^- = -1. \quad (51)$$

These relationships have been verified in our calculations which provides a nontrivial check for our solution.

III. GENERAL SCATTERING THEORY OF TWO-BODY PROBLEM IN TWO DIMENSION

In this section, to make our following discussions more self contained, we will briefly review the general theory of purely 2D elastic scattering under arbitrary interaction potential that decays faster than $1/r^2$. At inter-particle distance $r \rightarrow \infty$ the wave function of two distinguishable colliding atoms is represented as a superposition of the

incident plane wave and scattered circular wave

$$\psi_{\mathbf{k}}(\mathbf{r}) \simeq e^{i\mathbf{k}\cdot\mathbf{r}} + f(k, \theta) \frac{e^{ikr}}{\sqrt{r}}, \quad (52)$$

where \mathbf{k} is the relative momentum of the two particles under scattering, $f(k, \theta)$ is the scattering amplitude and θ is the angle between \mathbf{r} and \mathbf{k} .

When expanding in the partial wave channels, we have

$$\psi_{\mathbf{k}}(\mathbf{r}) = \sum_{m=-\infty}^{\infty} \psi_m(k, r) e^{im\theta} \quad (53)$$

$$f(k, \theta) = \sum_{m=-\infty}^{\infty} f_m(k) e^{im\theta} \quad (54)$$

where $f_m(k)$ and $\psi_m(k, r)$ are m -wave scattering amplitude and radial wave function, and we have

$$\psi_m(k, r) = \psi_m^0(k, r) + f_m(k) \frac{e^{ikr}}{\sqrt{r}} \quad (55)$$

where $\psi_m^0(k, r) = i^m J_m(kr)$ is m partial wave component of incident plane wave $e^{i\mathbf{k}\cdot\mathbf{r}}$. Finally, using the asymptotic form of Bessel function $J_m(x)$ in the $x \rightarrow \infty$ limit:

$$J_m(x) \rightarrow \sqrt{\frac{2}{\pi x}} \sin\left(x - \frac{l\pi}{2}\right) \quad (56)$$

where $l = m - 1/2$, and thus we have

$$\begin{aligned} \psi_m(k, r) &\rightarrow i^m \sqrt{\frac{2}{\pi kr}} \left[\sin\left(kr - \frac{l\pi}{2}\right) - \frac{f_m(k)}{4} e^{i(kr - l\pi/2)} \right] \\ &\propto \sin\left[kr - \frac{l\pi}{2} + \delta_m(k)\right] \end{aligned} \quad (57)$$

where $\delta_m(k)$ is the m partial wave scattering phase shift which is related to scattering amplitude as

$$f_m(k) = \sqrt{\frac{2i}{\pi k}} \frac{1}{\cot \delta_m(k) - i} \quad (58)$$

The 2D total and partial cross section for two distinguishable particles are thus given as

$$\sigma(k) = \sigma_0(k) + 2 \sum_{m=1}^{\infty} \sigma_m(k) \quad (59)$$

$$\sigma_m(k) = 2\pi |f_m(k)|^2 = \frac{4}{k} \sin^2 \delta_m(k) \quad (60)$$

In the case of identical particles, the scattering wave function in Eq. (52) should be symmetrized(anti-symmetrized) for bosons(fermions) and the cross sections have an extra factor 2 in the r.h.s. of Eq. (60). In this case, only even(odd) partial wave has nonzero contributions for bosonic(fermionic) particles.




FIG. 2: Scattering phase shifts (left panel) and cross sections (right panel) of pure repulsive $1/r^3$ interaction for the first four partial waves.

IV. QUANTUM DEFECT THEORY FOR QUASI-2D DIPOLE-DIPOLE SCATTERING

We consider two different cases of dipole-dipole scattering when two polar molecules are strongly confined along z -axis with a confining length $a_{\perp} \ll D$, while moving freely in the $x - y$ plane. Case I : by applying a strong static electric field perpendicular to the $x - y$ plane, all dipole moments are aligned along z -axis. In this case the interaction between two polar molecules has an isotropic repulsive D/r^3 tail [13]. Case II : one implements a strong electric field fast rotating within the $x - y$ plane and thus generates a fast rotating dipolar moment in each polar molecule. In this case, the dipole-dipole interaction has an isotropic attractive $-D/r^3$ tail [20]. In both cases, the inter molecular interaction will only deviate from the simple $\pm D/r^3$ form at short distance when $r \lesssim a_{\perp} \ll D$ [13]. As a result, the effect of this deviation can be encoded in a simple short range boundary condition [27–30]. This justifies the implementation of quantum defect theory which will be constructed below.

For any two dimensional interaction with a long range $\pm D/r^3$ tail, the radial part of scattering wave function in m -partial wave channel with energy ε can be generally written as $\psi_m(k, r) = u_{\varepsilon m}(r)/\sqrt{r}$ with $u_{\varepsilon m}(r)$ given as

$$u_{\varepsilon m}(r) = A_{\varepsilon m} [\bar{u}_{\varepsilon m}^1(r) - K^c \bar{u}_{\varepsilon m}^2(r)] \quad (61)$$

where K^c is usually called the quantum defect. At positive scattering energy, combining the asymptotic behavior of $\bar{u}_{\varepsilon m}^{1,2}(r)$ given in (20), (21) with the definition of phase shift in (57), one immediately obtains the phase shift as

$$K_m \equiv \tan \delta_m = \frac{K^c Z_{22} - Z_{12}}{Z_{11} - K^c Z_{21}}. \quad (62)$$

As for two-body bound states, one should use the asymptotic behavior at negative energy as given in (34) and (35). Since a physical bound state must decay exponentially at large r , the binding energy E_b can be determined by requiring the coefficient of $e^{\kappa r}$ term in $u_{\varepsilon m}(r)$ to vanish. This leads to the following equation




FIG. 3: Phase shifts of first four partial waves for $-D/r^3$ interaction across the scattering resonance. Solid lines refer to exact results from Eq. (70) while open circles are from low energy expansion in Eq. (71)-(74).

for E_b

$$\chi_m(\varepsilon_s) = K^c \quad (63)$$

where the $\chi_m(\varepsilon_s)$ has been defined earlier in Eq. (46).

As analyzed above, since the interacting potential only deviate from $\pm D/r^3$ within some short distance $r_0 \ll D$, K_c will be determined by the boundary condition at r_0 which is insensitive to neither energy nor angular momentum. In Fig. 1 we first illustrate how K^c changes with the short range behavior of interacting potentials. We consider two different model potentials $V_{\pm}(r)$ with $\pm D/r^3$ tails but truncated at r_0 as shown in Fig. 1(a) and (b). For $V_+(r) = -V_0\theta(r_0 - r) + \theta(r - r_0)D/r^3$ we fix r_0 at $0.1D$ and change the short range potential depth V_0 , while for $V_-(r) = +\infty\theta(r_0 - r) - \theta(r - r_0)D/r^3$ we just change the truncation radius r_0 at which a hard wall boundary condition is implemented.

From Fig. 1, one can see that K^c has very different behaviors as one tunes the short range behaviors of the potential with repulsive and attractive $1/r^3$ tail. In the repulsive case, as shown in Fig. 1(c), K^c is nearly zero everywhere except in the vicinity of some extremely narrow

shape resonances. This behavior is mainly due to the existence of a large repulsive barrier which makes the wave function almost unaffected by the short range attractive part of the potential. In contrast, for potential with an attractive tail the value of K_c changes significantly and experiences a sequence of much wider resonances as one tunes the short range behavior, see 1(d). As a result, for the repulsive case, we will only consider the scattering in pure repulsive limit corresponding to $K^c = 0$, while for the attractive case, we investigate both scattering and bound state properties across a shape resonance where K^c can be tuned from $-\infty$ to ∞ .

A. Scattering for pure repulsive $1/r^3$ potential

In this case, one can set $K^c = 0$ and the phase shift is given as

$$\tan \delta_m^+ = -\frac{Z_{21}^+}{Z_{11}^+}. \quad (64)$$

In Fig. 2, we show the results of phase shift and partial cross section for the first four partial waves. One can see

clearly that, in the low energy regime $kD \ll 1$ the s-wave channel completely dominates over higher partial waves, while at higher energy when $kD \gtrsim 1$ all partial waves has non-negligible contributions. At very low scattering energy, the asymptotic behavior of phase shift can be obtained analytically from our exact solution. Below we provide the low energy expansion of $\tan \delta_m^+$ for the first a few partial waves:

$$\tan \delta_0^+ \simeq \frac{\pi}{2} \frac{1 + (4/\pi)k_s \ln(k_s \bar{a}_0^+)}{\ln(k_s \bar{a}_0^+) - \pi k_s} \quad (65)$$

$$\tan \delta_1^+ \simeq -\frac{2k_s}{3} \left[1 + \frac{3\pi}{16} k_s \ln(k_s \bar{a}_1^+) - \frac{4}{5} k_s^2 \right] \quad (66)$$

$$\tan \delta_2^+ \simeq -\frac{2k_s}{15} \left(1 - \frac{\pi}{32} k_s + \frac{52}{1575} k_s^2 \right) \quad (67)$$

$$\tan \delta_3^+ \simeq -\frac{2k_s}{35} \left(1 - \frac{\pi}{128} k_s + \frac{236}{99225} k_s^2 \right) \quad (68)$$

where $k_s = kD$, $\bar{a}_0^+ = \exp(3\gamma)/2$ is simply the dimensionless 2D s-wave scattering length and $\bar{a}_1^+ = \exp(3\gamma - 11/12)/2$ with γ being the Euler's constant. For all $m > 0$ partial waves, the leading order behavior agrees with that from the first order Born approximation:

$$\tan \delta_m^+ \simeq -\frac{2k_s}{4m^2 - 1}, \quad (69)$$

while the sub leading terms in Eq. (65)-(68) are new in this work. These analytic expressions provide very good estimation for the phase shifts at low energy $k_s \lesssim 1$ as shown in Fig. 2 (dashed lines) and will be useful in many-body calculations [20].

B. Scattering and bound state for interaction with an attractive $-1/r^3$ tail

In this case, a quantum defect K^c is required to fix the scattering property as well as the bound states. The m -partial wave scattering phase shift is given by Eq. (62) as

$$\tan \delta_m^- = \frac{K^c Z_{22}^- - Z_{12}^-}{Z_{11}^- - K^c Z_{21}^-}, \quad (70)$$

In Fig. 3, we show the scattering phase shift for the first four partial waves at different quantum defect K^c .

Below we provide the low energy expansion of $\tan \delta_m^-$




FIG. 4: Binding energy for the first four partial waves where (a)-(d) refers to $m = 0 - 4$. Solid line, open squares and open circles refer to numerical solution of (77), analytic formulas (78)-(81), and the leading s-wave behavior $E_b = -1/a_0^2$.

for the first a few partial waves:

$$\tan \delta_0^- \simeq \frac{\pi}{2} \frac{1 - \frac{4}{\pi} k_s \ln |k_s \bar{a}_0^-|}{\ln |k_s \bar{a}_0^-| + \pi k_s} \quad (71)$$

$$\tan \delta_1^- \simeq \frac{2k_s}{3} \left[1 - \frac{\frac{3\pi}{16} \ln |k_s \bar{a}_1^-| + \frac{4}{5} k_s}{1 - \left(\frac{\ln^2 |k_s \bar{a}_1^-|}{8} - B^c \right) k_s^2} k_s \right] \quad (72)$$

$$\tan \delta_2^- \simeq \frac{2k_s}{15} \left[1 + \frac{\frac{\pi}{32} + \frac{52}{1575} k_s}{1 + \frac{\ln |k_s \bar{a}_2^-|}{120} k_s^2} k_s \right] \quad (73)$$

$$\tan \delta_3^- \simeq \frac{2k_s}{35} \left[1 + \frac{\frac{\pi}{128} + \frac{236}{99225} k_s}{1 + \frac{\ln |k_s \bar{a}_3^-|}{1120} k_s^2} k_s \right] \quad (74)$$

where $\bar{a}_m^- = \exp(3\gamma - \pi K^c - \beta_m)/2$ with $\beta_0 = 0$, $\beta_1 = 11/12$, $\beta_2 = 23/6$ and $\beta_3 = 243/40$. Comparing with the threshold behavior of phase shift for an s-wave contact interaction, one can see that the s-wave scattering length is still well defined through Eq. (71) and is given as $a_0 = \bar{a}_0^- D$. The constant B^c is related to K^c as

$$B^c = \frac{199}{1152} + \frac{5 + 12(K^c)^2}{96} \pi^2. \quad (75)$$

Again, the leading order behavior agrees with that from the first order Born approximation for all $m > 0$ partial waves:

$$\tan \delta_m^- \simeq \frac{2k_s}{4m^2 - 1}. \quad (76)$$

The sub leading $k_s^2 \ln |k_s \bar{a}_1^-|$ term in Eq. (72) also agrees with an earlier result obtained from a perturbative approach in [20].

As for two-body bound states, the binding energy for m partial wave is determined by

$$\chi_m^-(\varepsilon_s) = K^c. \quad (77)$$

The low energy expansion for χ_m leads to the following approximate equation for the near threshold binding energy $E_b = -\kappa^2$ in the limit $-K^c \gg 1$:

For s-wave, the binding wave number κ satisfies

$$\frac{\Omega_\kappa}{\pi} - [2\Omega_\kappa(\Omega_\kappa^2 + \pi^2) + 15 - 18\zeta(3)] \frac{(1 + 2\kappa_s)\kappa_s^2}{4\pi} \simeq K^c \quad (78)$$

where $\kappa_s = \kappa D$, $\Omega_\kappa = \ln(\kappa_s e^{3\gamma}/2)$, and $\zeta(s)$ is the Riemann Zeta function. The leading order behavior in the limit $a_0 \rightarrow \infty$ is simply $E_b = -1/a_0^2$. This is consistent with the fact that the s-wave scattering length is well defined through low energy behavior of $m = 0$ phase shift.

For higher partial waves, we find:

$$E_b^{(1)} \simeq \frac{8}{\pi K^c t} \left(1 + \frac{\ln t}{t-1}\right)^{-1} \quad (79)$$

$$E_b^{(2)} \simeq \frac{120}{\pi K^c} \quad (80)$$

$$E_b^{(3)} \simeq \frac{1120}{\pi K^c} \quad (81)$$

where $t = \ln|\pi K^c/2| - 6\gamma + 11/6$ and $E_b^{(m)}$ refers to binding energy for m partial wave. In Fig. 4, we show the results for the first a few bound states from numerically solving (77) and compare with the analytic formulas in Eq. (78)-(81). From these analytic behaviors for near threshold binding energy, one can clearly see that the scattering resonances happen at $K^c \rightarrow \infty$ for all partial waves, at which a zero energy bound state appears for each partial wave channel. This resonant feature is also

reflected in the scattering phase shifts as a rapid phase change at low scattering energy near the resonance, as illustrated in Fig. 3. It takes place at large negative and large positive value of K^c for s-wave and higher partial waves, respectively. The location of this phase change corresponds to the pole of Eq. (71)-(74), which for s-wave simply gives to $k \sim 1/a_0$. Such features in two-body scattering phase shift suggests that near the scattering resonance, the non-zero partial waves may still have important contribution to the many-body interaction energy even in the dilute regime when $nD^2 \ll 1$ where n is the particle density. Such influences in many-body physics will be left for future studies.

V. CONCLUSION

In this work, we for the first time present analytical solutions for 2D Schrödinger equation with both repulsive and attractive inverse cubic interactions. We constructed quantum defect theory base on these solution, and investigated the scattering properties and two-body bound states of two polar molecules confined in quasi-2D geometry with two different experimental setups. We provide both exact numerical and simple analytic low energy formula for the phase shifts and two-body binding energies, which could be useful in future many-body studies. In the attractive case, we identified a resonant feature that took place simultaneously in all partial wave channels which could have important effects in corresponding many-body system.

Acknowledgements. We thank Peng Zhang, Paul S. Julienne and Hui Zhai for useful discussions. This work is supported by the Fundamental Research Funds for the Central Universities, and the Research Funds of Renmin University of China under Grant No. 15XNLF18.

-
- [1] K.-K. Ni, S. Ospelkaus, M. H. G. de Miranda, A. Peer, B. Neyenhuis, J. J. Zirbel, S. Kotochigova, P. S. Julienne, D. S. Jin, and J. Ye, *Science* **322**, 231 (2008).
- [2] S. Ospelkaus, K.-K. Ni, G. Quemener, B. Neyenhuis, D. Wang, M. H. G. de Miranda, J. L. Bohn, J. Ye, D. S. Jin, *Phys. Rev. Lett.* **104**, 030402 (2010).
- [3] K.-K. Ni, S. Ospelkaus, D. Wang, G. Quemener, B. Neyenhuis, M. H. G. de Miranda, J. L. Bohn, J. Ye, and D. S. Jin, *Nature (London)* **464**, 1324 (2010).
- [4] M. A. Baranov, *Phys. Rep.* **464**, 71 (2008).
- [5] L. D. Carr *et al.*, *New J. Phys.* **11**, 055049 (2009).
- [6] G. Queéééer, and P. S. Julienne, *Chem. Rev.* **112**, 4949 (2012).
- [7] R. Qi, Z.-Y. Shi, and H. Zhai, *Phys. Rev. Lett.* **110**, 045302 (2013); T. Shi, S.-H. Zou, H. Hu, C.-P. Sun, and S. Yi, *Phys. Rev. Lett.* **110**, 045301 (2013).
- [8] Y. Li and C. Wu, *J. Phys.: Condens. Matter* **26** 493203 (2014).
- [9] E. R. Hudson *et al.*, *Phys. Rev. A* **73**, 063404 (2006).
- [10] S. Ospelkaus *et al.*, *Science* **327**, 853 (2010).
- [11] R. V. Krems, *Phys. Chem. Chem. Phys.* **10**, 4079 (2008).
- [12] M. H. G. de Miranda, A. Chotia, B. Neyenhuis, D. Wang, G. Quéméner, S. Ospelkaus, J. L. Bohn, J. Ye, and D. S. Jin, *Nat. Phys.* **7**, 502 (2011).
- [13] B. Simon, *Ann. Phys. (NY)* **97**, 279 (1976).
- [14] R. G. Newton, *J. Math. Phys.* **27**, 2720 (1986).
- [15] C. Ticknor, *Phys. Rev. A* **81**, 042708 (2010).
- [16] M. Klawunn, A. Pikoński, and L. Santos, *Phys. Rev. A* **82**, 044701 (2010).
- [17] J. P. D’Incao and C. H. Greene, *Phys. Rev. A* **83**, 030702(R) (2011).
- [18] M. Rosenkranz and W. Bao, *Phys. Rev. A* **84**, 050701(R) (2011).
- [19] A. G. Volosniev, D. V. Fedorov, A. S. Jensen, and N. T. Zinner, *Phys. Rev. Lett.* **106**, 250401 (2011).
- [20] J. Levisen, N. R. Cooper, and G. V. Shlyapnikov, *Phys. Rev. A* **84**, 013603 (2011).
- [21] F. Deuretzbacher, J. C. Cremon, and S. M. Reimann,

- Phys. Rev. A **81**, 063616 (2010); *ibid*, Phys. Rev. A **87**, 039903(E) (2013).
- [22] N. T. Zinner, B. Wunsch, I. B. Mekhov, S.-J. Huang, D.-W. Wang, and E. Demler, Phys. Rev. A **84**, 063606 (2011); M. D. Girardeau, and G.E. Astrakharchik, Phys. Rev. Lett. **109**, 235305 (2012); A. G. Volosniev, J. R. Armstrong, D. V. Fedorov, A. S. Jensen, M. Valiente and N. T. Zinner, New J. Phys. **15** 043046(2013).
- [23] S. Sinha and L. Santos, Phys. Rev. Lett. **99**, 140406 (2007).
- [24] P. Giannakeas, V.S. Melezhik, and P. Schmelcher, arXiv:1302.5632.
- [25] N. Bartolo, D. J. Papoular, L. Barbiero, C. Menotti, A. Recati, Phys. Rev. A **88**, 023603 (2013).
- [26] B. Gao, Phys. Rev. A **59**, 2778 (1999); *ibid*, Phys. Rev. A **58**, 1728 (1998).
- [27] C. Greene, U. Fano, and G. Strinati, Phys. Rev. A **19**, 1485 (1979); C. Greene, A. R. P. Rau, and U. Fano, Phys. Rev. A **26**, 2441 (1982).
- [28] F. H. Mies, J. Chem. Phys. **80**, 2514 (1984).
- [29] B. Gao, Eite Tiesinga, C. J. Williams, and P. S. Julienne, Phys. Rev. A **72**, 042719 (2005).
- [30] B. Gao, Phys. Rev. A **78**, 012702 (2008); *ibid*, Phys. Rev. A **80**, 012702 (2009).