

ON p -ADIC MÖBIUS MAPS

YUEFEI WANG AND JINGHUA YANG

ABSTRACT. In this paper, we study three aspects of the p -adic Möbius maps. One is the group $\mathrm{PSL}(2, \mathcal{O}_p)$, another is the geometrical characterization of the p -adic Möbius maps and its application, and the other is different norms of the p -adic Möbius maps. Firstly, we give a series of equations of the p -adic Möbius maps in $\mathrm{PSL}(2, \mathcal{O}_p)$ between matrix, chordal, hyperbolic and unitary aspects. Furthermore, the properties of $\mathrm{PSL}(2, \mathcal{O}_p)$ can be applied to study the geometrical characterization, the norms, the decomposition theorem of p -adic Möbius maps, and the convergence and divergence of p -adic continued fractions. Secondly, we classify the p -adic Möbius maps into four types and study the geometrical characterization of the p -adic Möbius maps from the aspects of fixed points in \mathbb{P}_{Ber}^1 and the invariant axes which yields the decomposition theorem of p -adic Möbius maps. Furthermore, we prove that if a subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$ containing elliptic elements only, then all elements fix the same point in \mathbb{H}_{Ber} without using the famous theorem–Cartan fixed point theorem, and this means that this subgroup has potentially good reduction. In the last part, we extend the inequalities obtained by Gehring and Martin [24, 25], Beardon and Short [12] to the non-archimedean settings. These inequalities of p -adic Möbius maps are between the matrix, chordal, three-point and unitary norms. This part of work can be applied to study the convergence of the sequence of p -adic Möbius maps which can be viewed as a special cases of the work in [20] and the discrete criteria of the subgroups of $\mathrm{PSL}(2, \mathbb{C}_p)$.

1. INTRODUCTION

1.1. **Statement of results.** We call an element in the projective special linear group $\mathrm{PSL}(2, \mathbb{C}_p)$ the p -adic Möbius map, where \mathbb{Q}_p is the field of p -adic rational numbers and \mathbb{C}_p is the completion of the algebraic closure of \mathbb{Q}_p . The projective space $\mathbb{P}^1(\mathbb{C}_p)$ is totally disconnected and not locally compact, which implies that we can not adopt the method used in dealing with the complex settings easily. The main tool that we use is the projective Berkovich space \mathbb{P}_{Ber}^1 (see concrete definitions in section 3), since $\mathrm{PSL}(2, \mathbb{C}_p)$ acts on the hyperbolic Berkovich space \mathbb{H}_{Ber} isometrically and the projective Berkovich space is compact with respect to the weak topology.

Date: August 24, 2018.

2010 Mathematics Subject Classification. 51B10, 32P05, 37P05, 40A05.

Key words and phrases. p -adic Möbius maps, discrete criteria, the convergence theorem, three points norms, unitary norms, hyperbolic norms, chordal norms, inequalities.

We study the subgroup $\mathrm{PSL}(2, \mathcal{O}_p)$, where $\mathcal{O}_p = \{z \mid |z| \leq 1\}$ firstly, since the study of the unitary groups of the projective special linear group $\mathrm{PSL}(2, \mathbb{C})$ is a very important topic in the study of Möbius maps. There exist a lot of equations of Möbius maps in the unitary group. It is natural to generalize the equations to the non-archimedean settings. We give a series of equations of the p -adic Möbius maps in $\mathrm{PSL}(2, \mathcal{O}_p)$ between matrix, chordal, hyperbolic and unitary aspects. Furthermore, the properties of $\mathrm{PSL}(2, \mathcal{O}_p)$ can be applied to study the geometrical characterization, the norms and the decomposition theorem of p -adic Möbius maps and the convergence and divergence of p -adic continued fractions.

Let g be a p -adic Möbius map, $\rho_v(z, w)$ be the chordal metric on the projective line $\mathbb{P}^1(\mathbb{C}_p)$, $\rho(z, w)$ be the hyperbolic metric on the hyperbolic Berkovich space, $L(g) = p^{\rho(g(\zeta_{Gauss}), \zeta_{Gauss})}$, and $\|\cdot\|$ be the matrix norm (see concrete definitions in section 2).

Theorem 1.1. *For any p -adic Möbius map g , the following are equivalent:*

- (1) $g \in \mathrm{PSL}(2, \mathcal{O}_p)$;
- (2) $L(g) = 1$;
- (3) $\rho(\zeta_{Gauss}, g(\zeta_{Gauss})) = 0$;
- (4) g is a chordal isometry;
- (5) $\|g\| = 1$;
- (6) for any h in the Matrix ring $M(2, \mathbb{C}_p)$, $\|gh\| = \|hg\| = \|ghg^{-1}\| = \|h\|$.

In the second part, we study the characterization of p -adic Möbius maps. In [26], Kato introduced the idea of the Kleinian group to study the p -adic Möbius maps firstly, and in [6, 30], Vermitage and Parker, Qiu, Yang and Yin gave the discrete criteria of subgroups of $\mathrm{PSL}(2, \mathbb{C}_p)$. The method that they used are also derived from the Kleinian group. We not only follow the philosophy of the Kleinian group to study the p -adic Möbius maps, but also we lay emphasis on the study of the difference between them. First, we classify the p -adic Möbius maps into four types which is a bit different than those in the Kleinian group, and we study the geometrical characterization of the p -adic Möbius maps from the aspects of fixed points in \mathbb{P}_{Ber} and the invariant axes. Furthermore, we can decompose a p -adic Möbius map g into two involutions α, β (an elliptic element of order 2), namely $g = \alpha \circ \beta$, and the structure of the fixed points of α, β can reflect the type and other properties of g . This method can be used to prove that if a subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$ contains elliptic elements only, then all elements fix the same point in \mathbb{H}_{Ber} without using the famous theorem–Cartan fixed point theorem, and this means that this subgroup has potentially good reduction (see concrete definitions in section 2). In the proof of the results, we should face three difficulties which do not exist in the archimedean settings. One is that $\mathbb{P}^1(\mathbb{C}_p)$ and \mathbb{P}_{Ber} are not locally compact, another is that there exists a new kind of p -adic Möbius maps—the wild elliptic elements whose geometrical structure of the fixed points are complicated, and the other is that the prime

number p affects the structure of the fixed points of p -adic Möbius maps. We give a series of tables to compare the properties of p -adic Möbius maps and those of the Kleinian group.

	the Kleinian group	p -adic Möbius maps
type	loxodromic	loxodromic
	parabolic	parabolic
	elliptic	tame elliptic
		wild elliptic

Assuming $g(z) = \frac{az+b}{cz+d} = \alpha \circ \beta$ with $ad - bc = 1$, let F_g, F_α, F_β denote the fixed points of g, α, β in $\mathbb{P}^1 \cup \mathbb{H}^3$ respectively.

the Kleinian group	$F_g \subset \mathbb{P}^1 \cup \mathbb{H}^3$	$F_\alpha \cap F_\beta$
loxodromic	two points in \mathbb{P}^1	\emptyset
parabolic	one point in \mathbb{P}^1	unique point in \mathbb{P}^1
elliptic	a geodesic line in $\mathbb{P}^1 \cup \mathbb{H}^3$	unique point in \mathbb{H}^3

Let $\text{Int}(A)$ denote the interior of the set A . Assuming $g(z) = \frac{az+b}{cz+d} = \alpha \circ \beta$ with $ad - bc = 1$, let F_g, F_α, F_β denote the fixed points of g, α, β in \mathbb{P}_{Ber}^1 respectively.

the p -adic Möbius map	$F_g \subset \mathbb{P}^1 \cup \mathbb{H}_{Ber}$	$F_\alpha \cap F_\beta$
loxodromic	two points in \mathbb{P}^1	\emptyset
parabolic	$F_g \cap \mathbb{H}_{Ber} \neq \emptyset$	$\text{Int}(F_\alpha \cap F_\beta) \neq \emptyset$
tame elliptic	a geodesic line in \mathbb{P}_{Ber}^1	unique point in \mathbb{H}_{Ber}
wild elliptic	$\text{Int}(F_g) \neq \emptyset$	$\text{Int}(F_\alpha \cap F_\beta) \neq \emptyset$

Theorem 1.2. *If the subgroup $G \subset \text{PSL}(2, \mathbb{C}_p)$ contains elliptic elements only, then all the elements of G share at least one fixed point in \mathbb{H}_{Ber} . Furthermore, G has potentially good reduction, and G is equicontinuous on $\mathbb{P}^1(\mathbb{C}_p)$.*

A point $a \in \mathbb{P}^1(\mathbb{C}_p)$ is called the limit point of a subgroup G of $\text{PSL}(2, \mathbb{C}_p)$ if there exists a point $b \in \mathbb{P}^1(\mathbb{C}_p)$ and an infinite sequence $\{g_n | n \geq 1\} \subset G$, where $g_n \neq g_m$ if $n \neq m$ with $\lim g_n(b) = a$. The set consisting of all limit points is called the *limit set*. G is said to act discontinuously at $x \in \mathbb{P}^1(\mathbb{C}_p)$ if there is a neighborhood U of x such that $g(U) \cap U = \emptyset$ for all but finitely many $g \in G$. The set of points at which G acts discontinuously is called the *discontinuous set*.

Theorem 1.3. *If the limit set of G is empty, then G has potentially good reduction.*

Two points α, β are called *antipodal* points if there exists an element $u \in \mathrm{PSL}(2, \mathcal{O}_p)$ such that $u(0) = \alpha, u(\infty) = \beta$.

Theorem 1.4. *For any p -adic Möbius map g , there exists an element $u \in \mathrm{PSL}(2, \mathcal{O}_p)$ such that $g = uf$, where either f is a loxodromic element with antipodal fixed points, or $f = I$.*

In the last part, we extend the inequalities obtained by Gehring and Martin [24, 25], Beardon and Short [12] to the non-archimedean settings. These inequalities of p -adic Möbius maps are between the matrix, chordal, three-point and unitary norms. This part of work can be applied to study the convergence of the sequence of p -adic Möbius maps which can be viewed as a special cases of the work in [20] and the discrete criteria of the subgroups of $\mathrm{PSL}(2, \mathbb{C}_p)$.

For any two p -adic Möbius maps g, h , we define the **uniformly convergent metric** on the $\mathrm{PSL}(2, \mathbb{P}^1(\mathbb{C}_p))$ as follows

$$\rho_0(g, h) = \sup_{z \in \mathbb{P}^1(\mathbb{C}_p)} (g(z), h(z)).$$

Let $M(g) = \|g - g^{-1}\| / \|g\|$.

Theorem 1.5. *Let g be a p -adic Möbius map.*

- (1) *If $p = 3$, then $\rho_0(g, I) = M(g)$.*
- (2) *If $p = 2$, then $2^{-1}M(g) \leq \rho_0(g, I) \leq 2M(g)$.*

Let $\varepsilon(g) = \max\{\rho_v(g(z_0), z_0), \rho_v(g(z_1), z_1), \rho_v(g(z_2), z_2)\}$, where z_0, z_1, z_2 are three distinct roots of the equation $z^3 = 1$.

Theorem 1.6. *For any p -adic Möbius map g , we have $2^{-1}\varepsilon(g) \leq M(g) \leq 6\varepsilon(g)$.*

Let $\varepsilon_1(g) = \{\rho_v(g(0), 0), \rho_v(g(1), 1), \rho_v(g(\infty), \infty)\}$.

Theorem 1.7. *For any p -adic Möbius map, $2^{-1}\varepsilon_1(g) \leq M(g) \leq \varepsilon_1(g)$.*

If g is a parabolic element, we can improve the inequality. Let $\varepsilon_2(g) = \max\{\rho_v(g(0), 0), \rho_v(g(\infty), \infty)\}$.

Corollary 1.8. *If g is a parabolic element, then $2^{-1}\varepsilon_2(g) \leq M(g) \leq \varepsilon_2(g)$.*

Let $\mathcal{U} = \mathrm{PSL}(2, \mathcal{O}_p)$. We define $d(g, \mathcal{U}) = \inf\{\rho_0(g, u) | u \in \mathcal{U}\}$. $d(g, \mathcal{U})$ measures how far an element from the group \mathcal{U} . This result is quite different from that in the archimedean setting.

Theorem 1.9. *For any p -adic Möbius map, either $d(g, \mathcal{U}) = 0$, if $g \in \mathcal{U}$, or $d(g, \mathcal{U}) = 1$, if $g \notin \mathcal{U}$.*

As an application of this result, we derive a discrete criteria of subgroups of $\mathrm{PSL}(2, \mathbb{C}_p)$.

Theorem 1.10. *If G is a subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$ with $G \cap \mathcal{U} = \{I\}$, then G is a discrete subgroup.*

Corollary 1.11. *If a subgroup $G \subset \mathrm{PSL}(2, \mathbb{C}_p)$ contains unit element or loxodromic element only, then G is a discrete subgroup.*

The other application of these inequalities is to get the convergence theorem of p -adic Möbius maps.

Theorem 1.12. *Let $\{g_n\}$ be a sequence of p -adic Möbius maps, z_j , ($j = 1, 2, 3$) be three distinct points and $g_n(z_j) \rightarrow w_j$, where w_j are also three distinct points. Then the sequence $\{g_n\}$ converges to a p -adic Möbius map g uniformly, where $g(z_j) = w_j$, ($j = 1, 2, 3$).*

1.2. Motivation. Firstly the study of the Kleinian group in the archimedean case has been well developed for a rather long time. It is natural for one to consider a parallel theory in the non-archimedean settings. Here we wish to give a fairly clear picture of the p -adic Möbius maps from the point of view of the Kleinian group. We study three aspects of the p -adic Möbius maps. One is the group $\mathrm{PSL}(2; \mathcal{O}_p)$, another is the geometrical characterization of the p -adic Möbius maps and its application, and the other is different norms of the p -adic Möbius maps, since $\mathrm{PSL}(2; \mathcal{O}_p)$ is similar to the unitary group in the Kleinian group, the geometrical characterization of the $\mathrm{PSL}(2; \mathcal{O}_p)$ maps is useful in the study of the structure and dynamics of the subgroups of $\mathrm{PSL}(2; \mathbb{C}_p)$, and the study of norms of Möbius maps is a very important topic, and many mathematicians such as Gehring and Martin [24, 25], Beardon and Short [12] do a lot of works in this topic.

The other reason is that the three aspects can be viewed as tools to study other related topics. The properties of the norms of p -adic Möbius maps have a lot of applications in three topics. One is the discrete criteria of subgroups of $\mathrm{PSL}(2; \mathbb{C}_p)$ and another is the pointwise convergence of p -adic Möbius maps (see [26, 32, 34]) and the other is the p -adic continued fraction. The geometrical characterization shows that the subgroup G containing elliptic element only shares one unique point, which means that a non-elementary group should have an loxodromic element which is very useful to study the dynamics of the discrete subgroup of $\mathrm{PSL}(2; \mathbb{C}_p)$. The rapid development of the Berkovich space and the arithmetical dynamical system (see [3], [4], [11], [13], [14], [15], [16], [19], [21], [31]) also promote studying the p -adic Möbius maps. Especially, the study of p -adic Möbius maps can be applied in studying the quantum mechanics and quantum cosmology [22, 23]). In this paper we give the affirmative answers to all these three questions in the non-archimedean settings.

1.3. Outline of the paper. Outline of the paper. In section 1, we present our main results of the paper. In section 2, the basic theories of the p -adic analysis and the Berkovich space are briefly reviewed. In section 3, we obtain a few preliminary results of p -adic Möbius maps. Section 4 contains proofs of the equations of p -adic Möbius maps in $\mathrm{PSL}(2; \mathcal{O}_p)$. In section 5, we give the results of the geometrical characterization of the p -adic Möbius maps. In section 6, the inequalities of p -adic Möbius maps between matrix,

chordal, three points, hyperbolic, and unitary norms are derived. In section 7, we prove the decomposition theorem of the p -adic Möbius maps and discuss its application.

2. SOME PRELIMINARY RESULTS

2.1. The non-archimedean space $\mathbb{P}^1(\mathbb{C}_p)$. Let $p \geq 2$ be a prime number. Let \mathbb{Q}_p be the field of p -adic numbers and \mathbb{C}_p be the completion of the algebraic closure of \mathbb{Q}_p . Denote $|\mathbb{C}_p^*|$ the valuation group of \mathbb{C}_p . Then every element $r \in |\mathbb{K}^*|$ can be expressed as $r = p^s$ with $s \in \mathbb{Q}$. We have the **strong triangle inequality**

$$|x - y| \leq \max\{|x|, |y|\}$$

for $x, y \in \mathbb{C}_p$. If x, y and z are points of \mathbb{C}_p with $|x - y| < |x - z|$, then $|x - z| = |y - z|$.

For any $a \in \mathbb{C}_p$, and $r > 0$, we define

$$D(a, r)^- = \{z \in \mathbb{C}_p \mid |z - a| < r\}$$

to be the **open** disk of radius r and centered at a . Similarly,

$$D(a, r) = \{z \in \mathbb{C}_p \mid |z - a| \leq r\}$$

denote the **closed** disk of radius r and centered at a . Both $D(a, r)^-$ and $D(a, r)$ are closed and open topologically, and every point in disk $D(a, r)^-$ is the center. This denotes that if $x \in D(a, r)^-$, then $D(a, r)^- = D(x, r)^-$ (resp. $D(a, r) = D(x, r)$). By the strong triangle inequality, if two disks D_1 and D_2 in \mathbb{C}_p have non-empty intersection, then either $D_1 \subset D_2$, or $D_2 \subset D_1$.

For any $z, w \in \mathbb{P}^1(\mathbb{C}_p)$, we define the chordal distance

$$\rho_v(z, w) = \frac{|z - w|}{\max\{1, |z|\} \max\{1, |w|\}}$$

for $z, w \in \mathbb{C}_p$,

$$\rho_v(z, w) = \frac{1}{\max\{1, |w|\}}$$

for $w \in \mathbb{C}_p$ and $z = \infty$, and

$$\rho_v(z, w) = 0$$

for $z = w = \infty$.

It follows the definition of the chordal distance that if $|z| \leq 1, |w| \leq 1$, then $\rho_v(z, w) = |z - w|$, and if $|z| > 1, |w| \leq 1$, then by the strong triangle inequality, we have $|z - w| = |z|$, and hence $\rho_v(z, w) = \frac{|z - w|}{|z|} = 1$, and if $|z| > 1, |w| > 1$, then $\rho_v(z, w) = \frac{|z - w|}{|z||w|} = \left| \frac{1}{z} - \frac{1}{w} \right|$.

Lemma 2.1 ([2]). *Let $d_0 > 1$ be an integer which is not divisible by p , and $d = d_0 p^t$ be a natural number. Let ζ be the primitive d -th root of unity. Then $|\zeta - 1| = 1$.*

Lemma 2.2 ([2]). *Let ζ be the primitive p^d -th root of unity. Then $|\zeta - 1| = p^{-\frac{1}{p^d - 1(p-1)}}$.*

2.2. The Berkovich space. We shall use a few properties of about the structure of the Berkovich affine line and its topology. Here we give a brief introduction to the Berkovich space. More details can be found in [7].

The underlying point set for the **Berkovich affine line** \mathbb{A}_{Ber}^1 is the collection of all multiplicative seminorms $[\cdot]_x$ on the polynomial ring $\mathbb{C}_p[z]$ which extend the absolute value on \mathbb{C}_p . Recall that a **multiplicative seminorms** on the ring $\mathbb{C}_p[z]$ is a function $[\cdot]_x : \mathbb{C}_p[z] \rightarrow [0, +\infty)$.

- $[0]_x = 0, [1]_x = 1$;
- $[fg]_x = [f]_x [g]_x$ for all $f, g \in \mathbb{C}_p[z]$;
- $[f + g]_x \leq \max\{[f]_x, [g]_x\}$ for all $f, g \in \mathbb{C}_p[z]$.

It is a **norm** provided that $[f]_x = 0$ if and only if $f = 0$. The topology on \mathbb{A}_{Ber}^1 is the weakest one for which the mapping $x \mapsto [f]_x$ is continuous for all $f \in \mathbb{C}_p[z]$.

Recall the Berkovich's classification Theorem: Every point $x \in \mathbb{A}_{Ber}^1$ can be viewed as a nested sequence of disks $D(a_1, r_1) \supset D(a_2, r_2) \supset \dots$. Moreover, points in \mathbb{A}_{Ber}^1 can be divided into four types:

- A point in \mathbb{A}_{Ber}^1 corresponding to a nested sequence $\{D(a_i, r_i)\}$ of disks with $\lim r_i = 0$ is said to be of **type I**.
- A point in \mathbb{A}_{Ber}^1 corresponding to a nested sequence $\{D(a_i, r_i)\}$ of disks with nonempty intersection, for which $r = \lim r_i > 0$ belongs to the valuation group $|\mathbb{C}^*|$ of \mathbb{C} , is said to be of **type II**.
- A point in \mathbb{A}_{Ber}^1 corresponding to a nested sequence $\{D(a_i, r_i)\}$ of disks with nonempty intersection, for which $r = \lim r_i > 0$ does not belong to the valuation group $|\mathbb{C}_p^*|$ of \mathbb{C}_p , is said to be of **type III**.
- A point in \mathbb{A}_{Ber}^1 corresponding to a nested sequence $\{D(a_i, r_i)\}$ of disks with empty intersection is said to be of **type IV**.

We call points of type I, II, III the **nonsingular points**, and points of type IV the **singular points**. Every nonsingular point in \mathbb{A}_{Ber}^1 has a representation which is the intersection of the corresponding nested sequence of disks. So a nonsingular point in \mathbb{A}_{Ber}^1 can be identifying with a point a (type I) or a disk $D(a, r)$ (type II, III).

We define a partial order on \mathbb{A}_{Ber}^1 as follows. For $x, y \in \mathbb{A}_{Ber}^1$, define $x \preceq y$ if and only if $[f]_x \leq [f]_y$ for all $f \in \mathbb{C}_p[z]$. If x, y are two points in \mathbb{A}_{Ber}^1 identifying with disks $D(a, r)$ and $D(a', r')$ respectively, then $x \preceq y$ if and only if $D(a, r) \subset D(a', r')$.

For a point $x \in \mathbb{A}_{Ber}^1$, we denote the set of elements larger than x by

$$[x, \infty[= \{w \in \mathbb{A}_{Ber}^1 \mid x \preceq w\}.$$

Observe that $[x, \infty[$ is isomorphic, as an ordered set, to $[0, +\infty[\subset \mathbb{R}$.

Given two points x, y in \mathbb{A}_{Ber}^1 , we have that

$$[x, \infty[\cap [y, \infty[= [x \vee y, \infty[,$$

where $x \vee y$ is the smallest element larger than x and y . If x is different from y , then the element $x \vee y$ is a point of type II. We also denote

$$[x, y] = \{w \in \mathbb{A}_{Ber}^1 \mid x \preceq w \preceq x \vee y\} \cup \{w \in \mathbb{A}_{Ber}^1 \mid y \preceq w \preceq x \vee y\}.$$

The sets $]x, y]$, $[x, y[$ and $]x, y[$ are defined in the obvious way.

For a set $E \subset \mathbb{K}$, denote $\text{diam}(E) = \sup_{z, w \in E} |z - w|$ the diameter of E in the non-Archimedean metric. For $x \in \mathbb{A}_{Ber}^1$, which is corresponding to the nested sequence $\{D(a_i, r_i)\}$ of disks, the diameter of x is given by

$$\text{diam}(x) = \lim_{i \rightarrow \infty} \text{diam}(D(a_i, r_i)).$$

For a nonsingular element $x \in \mathbb{A}_{Ber}^1$ identifying with the disk $D(a, r)$, the diameter of x coincides with the diameter (radius r) of $D(a, r)$.

In order to endow the Berkovich affine line with a topology, we define an open disk of \mathbb{A}_{Ber}^1 by

$$\mathcal{D}(a, r)^- = \{x \in \mathbb{A}_{Ber}^1 \mid \text{diam}(a \vee x) < r\},$$

for $a \in \mathbb{K}$ and $r > 0$. Similarly, a closed disk of \mathbb{A}_{Ber}^1 is defined by

$$\mathcal{D}(a, r) = \{x \in \mathbb{A}_{Ber}^1 \mid \text{diam}(a \vee x) \leq r\}.$$

Let $\mathbb{P}^1(\mathbb{C}_p)$ be the projective line over \mathbb{C}_p , which can be viewed as $\mathbb{P}^1(\mathbb{C}_p) = \mathbb{C}_p \cup \{\infty\}$. We can also introduce the **Berkovich projective line** \mathbb{P}_{Ber}^1 over $\mathbb{P}^1(\mathbb{K})$ similarly. In [7], Baker and Rumely pointed out that \mathbb{P}_{Ber}^1 can be defined as $\mathbb{A}_{Ber}^1 \cup \{\infty\}$, where $\infty \in \mathbb{P}^1(\mathbb{K})$ is regarded as a point of type I. \mathbb{P}_{Ber}^1 can be also identifying with the disjoint union of a closed set \mathcal{X} which is homeomorphic to $\mathcal{D}(0, 1)$ and an open set \mathcal{Y} which is homeomorphic to $\mathcal{D}(0, 1)^-$. This provides a useful way to visualize \mathbb{P}_{Ber}^1 .

Lemma 2.3 ([7]). *The space \mathbb{P}_{Ber}^1 is uniquely path-connected. More precisely, given any two distinct points $x, y \in \mathbb{P}_{Ber}^1$, there is a unique arc $[x, y]$ in \mathbb{P}_{Ber}^1 from x to y , and if $x \in \mathcal{X}$ and $y \in \mathcal{Y}$, then the arc contains $\zeta_{\text{Gauss}} \in \mathbb{A}_{Ber}^1$, where ζ_{Gauss} is identifying with the disk $D(0, 1)$.*

We say that a metric space (X, d) is an **\mathbb{R} -tree**, if for any two points $x, y \in X$, there is a unique arc from x to y and this arc is the geodesic segment.

Lemma 2.4 ([7]). *Let $x, y \in \mathcal{D}(0, 1)$. Then the metric $d(x, y) = 2\text{diam}(x \vee y) - \text{diam}(x) - \text{diam}(y)$ makes $\mathcal{D}(0, 1)$ into an \mathbb{R} -tree.*

Therefore, we can define a metric on \mathbb{P}_{Ber}^1 as $d_p(x, y) = d(x, y)$ if $x, y \in \mathcal{X}$ which can be identified with $\mathcal{D}(0, 1)$, and $d_p(x, y) = d(x, y)$ if $x, y \in \mathcal{Y}$ which can be identified with $\mathcal{D}(0, 1)^-$, and $d_p(x, y) = d(x, \zeta_{\text{Gauss}}) + d(\zeta_{\text{Gauss}}, y)$ if $x \in \mathcal{X}$ and $y \in \mathcal{Y}$.

The **Berkovich hyperbolic space** \mathbb{H}_{Ber} is defined by

$$\mathbb{H}_{Ber} = \mathbb{P}_{Ber}^1 \setminus \mathbb{P}^1(\mathbb{C}).$$

Since $\infty \in \mathbb{P}_{Ber}^1$ is of type I, \mathbb{H}_{Ber} can be also viewed as $\mathbb{A}_{Ber}^1 \setminus \mathbb{C}$. Thus \mathbb{H}_{Ber} has a tree structure induced by \mathbb{A}_{Ber}^1 . Over \mathbb{H}_{Ber} we can define the **hyperbolic distance**,

$$\rho(x, y) = 2 \log \text{diam}(x \vee y) - \log \text{diam}(x) - \log \text{diam}(y), \quad x, y \in \mathbb{H}_{Ber}.$$

Lemma 2.5 ([7]). \mathbb{H}_{Ber} is a complete metric space under $\rho(x, y)$.

Lemma 2.6 ([7]). Suppose that $w, y, x \in \mathbb{H}_{Ber}$. Then $\rho(x, y) = \rho(x, w) + \rho(w, y)$ if and only if w belongs to $[x, y]$.

2.3. The action of a rational map ϕ over \mathbb{P}_{Ber}^1 . Let $\phi \in \mathbb{C}_p(T)$ be a nonconstant rational function of degree $d \geq 1$. Since type I points are dense in \mathbb{P}_{Ber}^1 , for any $x \in \mathbb{H}_{Ber}$, there exists a sequence x_n tending to x with respect to the Berkovich topology. We can define $\phi(z) = \lim_{n \rightarrow \infty} \phi(x_n)$.

If $d = 1$, ϕ has an algebraic inverse and thus induces an automorphism of \mathbb{P}_{Ber}^1 . Define $\text{Aut}(\mathbb{P}_{Ber}^1)$ to be the group of automorphisms of \mathbb{P}_{Ber}^1 . The following lemmas can be found in [7].

Lemma 2.7 ([7]). If $\phi(z) \in \mathbb{C}_p(z)$ is nonconstant, then $\phi : \mathbb{P}_{Ber}^1 \rightarrow \mathbb{P}_{Ber}^1$ takes points of each type (I, II, III, IV) to points of the same type. Thus $\phi(z)$ has a given type if and only if z does.

Lemma 2.8 ([7]). Let $f(z) \in \mathbb{C}_p(z)$ be a nonconstant rational function, and suppose that $x \in \mathbb{A}_{Ber}^1$ is a point of type II, corresponding to a disc $D(a, r)$ in \mathbb{C}_p under Berkovich's classification. Then $f(x)$ corresponds to the disc $D(b, R)$ if and only if there exist $a_1, a_2, \dots, a_m, b_1, b_2, \dots, b_n \in \mathbb{C}_p$ for which $D(b, R) \setminus \cup_{i=1}^n D(b_i, R)^-$ is the image under $f(z)$ of $D(a, r) \setminus \cup_{i=1}^m D(a_i, r)^-$.

2.4. Reduction on rational function over \mathbb{C}_p . Let $\mathcal{O}_p = \{z \in \mathbb{C}_p \mid |z| \leq 1\}$, $\mathcal{O}^* = \{z \in \mathbb{C}_p \mid |z| = 1\}$, $\mathcal{M} = \{z \mid |z| < 1\}$ and $k = \mathcal{O}_p / \mathcal{M}$. We also call k the residue field of \mathbb{C}_p . If $x \in \mathcal{O}_p$, we denote the reduction of x modulo \mathcal{M} by \bar{x} . For any $z \in \mathbb{C}_p$, there exists a homogeneous coordinate $[x, y]$ for z , where $x, y \in \mathcal{O}$ with at least one in \mathcal{O}^* . Reduction induces a well-defined map $\mathbb{P}^1(\mathbb{C}_p) \rightarrow \mathbb{P}^1(k)$ by $\overline{[x, y]} = [\bar{x}, \bar{y}]$. Any rational function $f(z) \in \mathbb{C}_p(z)$ can be written in homogeneous coordinates as $f([x, y]) = [g(x, y), h(x, y)]$ where $g, h \in \mathcal{O}_p[x, y]$ are relatively prime homogeneous polynomials of degree $d = \deg(f)$. We can ensure that at least one coefficient of either g or h has valuation zero (i.e., absolute value 1). The reduction map induces a map $\mathcal{O}_p[x, y] \rightarrow k[x, y]$.

Definition 2.9. Let $f(z) \in \mathbb{C}_p(z)$ be a map with homogenous presentation $f([x, y]) = [g(x, y), h(x, y)]$, where $g, h \in \mathcal{O}_p[x, y]$ are relatively prime homogeneous polynomials of degree $d = \deg(f)$, and at least one coefficient of g or h has absolute value 1. We say that f has good reduction if \bar{g} and \bar{h} have no common zeros in $k \times k$ besides $(x, y) = (0, 0)$.

If there is some linear fractional transformation $h \in \text{PSL}(2, \mathbb{C}_p)$ such that $h^{-1} \circ f \circ h$ has good reduction, we say that f has **potentially good reduction**.

Lemma 2.10 ([33]). *Let $f \in \mathrm{PSL}(2, \mathbb{C}_p)$ be a rational function of degree one. Then f has good reduction if and only if $f \in \mathrm{PSL}(2, \mathcal{O})$.*

Lemma 2.11 ([7]). *Let $f \in \mathbb{C}_p(z)$ be a non-constant rational function. Then f has good reduction if and only if $f^{-1}(\zeta_{\mathrm{Gauss}}) = \zeta_{\mathrm{Gauss}}$.*

3. THE P-ADIC MÖBIUS MAPS

We classify non-unit elements in $\mathrm{PSL}(2, \mathbb{C}_p) = \mathrm{SL}(2, \mathbb{C}_p)/\{\pm I\}$. Since the product of all eigenvalues of $g \in \mathrm{PSL}(2, \mathbb{C}_p)$ is one, either the absolute value of each eigenvalue of g is one or there exists at least one eigenvalue whose absolute value is larger than 1. Thus each non-unit element $g \in \mathrm{PSL}(2, \mathbb{C}_p)$ falls into the following four classes:

(a) g is said to be **parabolic** if the absolute value of any eigenvalue of g is 1, and g can not be conjugated to a diagonal matrix.

(b) g is said to be **loxodromic** if there exists at least eigenvalue of g whose absolute value is larger than 1.

(c) $g \neq I$ is said to be **elliptic** if the absolute value of any eigenvalue of g is 1, and g can be conjugated to a diagonal matrix.

In this paper, we classify the elliptic elements more precisely.

(d) g is said to be **tame elliptic** if the two eigenvalues λ_1, λ_2 of g satisfy $|\lambda_1 - 1| = |\lambda_2 - 1| = 1$.

(e) g is said to be **wild elliptic** if one of the eigenvalues of g lies in the disc $D(1, 1)^-$.

For $g = (a_{ij})$ in the matrix ring $M(m, \mathbb{C}_p)$, the norm of g is defined by $\|g\| = \max_{1 \leq i \leq m, 1 \leq j \leq m} \{|a_{ij}|\}$. Obviously, $\|g\| = 0$ implies that each $a_{ij} = 0$. It is easy to verify that $\|\alpha g\| = |\alpha| \|g\|$, $\|g + h\| \leq \max\{\|g\|, \|h\|\}$ and $\|gh\| \leq \|g\| \|h\|$.

For any element $g \in \mathrm{PSL}(2, \mathbb{C}_p)$, there exist two lifts g_1, g_2 in $\mathrm{SL}(2, \mathbb{C}_p)$ with $\|g_1\| = \|g_2\|$. We define $\|g\| = \|g_1\| = \|g_2\|$. If $g, h \in \mathrm{PSL}(2, \mathbb{C}_p)$ correspond the lifts $g_1, g_2 \in \mathrm{SL}(2, \mathbb{C}_p)$ and $h_1, h_2 \in \mathrm{SL}(2, \mathbb{C}_p)$ respectively, then we define $\|g - h\| = \inf_{1 \leq i \leq 2, 1 \leq j \leq 2} \|g_i - h_j\|$.

If $d = 1$, ϕ has an algebraic inverse and thus induces an automorphism of $\mathbb{P}_{\mathrm{Ber}}^1$. Define $\mathrm{Aut}(\mathbb{P}_{\mathrm{Ber}}^1)$ to be the group of automorphisms of $\mathbb{P}_{\mathrm{Ber}}^1$. The following lemmas can be found in [7].

Lemma 3.1 ([7]). *The path distance metric $\rho(x, y)$ on $\mathbb{H}_{\mathrm{Ber}}$ is independent of the choice of homogenous coordinates on $\mathbb{P}_{\mathrm{Ber}}^1$, in the sense that if $h(z) \in \mathbb{C}_p(z)$ is a p -adic Möbius map, then $\rho(h(x), h(y)) = \rho(x, y)$ for all $x, y \in \mathbb{H}_{\mathrm{Ber}}$.*

Lemma 3.2 ([30]). *Let w satisfy $|w| = \lambda|w - a|$, where $\lambda \in |\mathbb{C}_p^*|$, and $w, a \in \mathbb{C}_p$.*

- (1) *If $\lambda > 1$, then $w \in D(a, \frac{|a|}{\lambda}) \setminus D(a, \frac{|a|}{\lambda})^-$.*
- (2) *If $\lambda = 1$, then $w \in \mathbb{P}^1(\mathbb{C}_p) \setminus D(0, |a|)^- \cup D(a, |a|)^-$.*
- (3) *If $0 < \lambda < 1$, then $w \in D(0, \lambda|a|) \setminus D(0, \lambda|a|)^-$.*

Proposition 3.3. *A p -adic Möbius map g is*

- (1) *parabolic if it is conjugate to $z \rightarrow z + 1$;*
- (2) *elliptic if it is conjugate to $z \rightarrow kz$ for some k with $|k| = 1$, $k \neq 1$;*
- (3) *loxodromic if it is conjugate to $z \rightarrow kz$ for some k with $|k| > 1$.*

Proposition 3.4. *Let f and g be two p -adic Möbius maps neither of which is the identity. Then f and g are conjugate if and only if $\text{trace}(f) = \text{trace}(g)$.*

Proposition 3.5. *Let $g \neq I$ be a p -adic Möbius map. Then*

- (1) *g is parabolic if and only if $(a + d)^2 - 4 = 0$;*
- (2) *g is elliptic if and only if $0 < |(a + d)^2 - 4| \leq 1$;*
- (3) *g is loxodromic if and only if $|(a + d)^2 - 4| > 1$.*

Proof. Since the trace of the matrix is invariant under the conjugation, without loss of generality, we can assume that $g(z) = \frac{a}{d}z$, if g is elliptic or loxodromic which yields that $\text{tr}(g) = a + d = \lambda + 1/\lambda$, where λ is the eigenvalue of g . Thus $(a + d)^2 - 4 = (\lambda - 1/\lambda)^2$. By the non-archimedean property of the metric, we have $|\lambda - 1/\lambda| > 1$, if g is loxodromic, and $0 < |\lambda - 1/\lambda| < 1$, if g is elliptic. If g is parabolic, we can assume that $g(z) = z + 1$ which yields that $a + d = 2$.

Conversely, if $(a + d)^2 - 4 = 0$, then $\lambda - 1/\lambda = 0$ which denotes $\lambda^2 = 1$. Since $g \in \text{PSL}(2, \mathbb{C}_p)$, we have $\lambda = 1$ which yields that g is parabolic. If $|(a + d)^2 - 4| > 1$, then $|\lambda - 1/\lambda| > 1$ which denotes that $|\lambda| > 1$ or $|1/\lambda| > 1$. Thus g is loxodromic. If $0 < |(a + d)^2 - 4| < 1$, then $0 < |\lambda - 1/\lambda| < 1$ which yields $|\lambda| < 1$. Thus g is elliptic. □

4. THE PROPERTIES OF $\text{PSL}(2, \mathcal{O}_p)$

Lemma 4.1 ([33]). *If $f \in \text{PSL}(2, \mathcal{O}_p)$, we have $\rho_v(f(z), f(w)) = \rho_v(z, w)$.*

Lemma 4.2 ([30]). *Let g be any p -adic Möbius map. Then*

$$\rho(g(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}}) = 2 \log_p \|g\|.$$

Lemma 4.3. *Let g be any p -adic Möbius map. Then the best Lipschitz constant (relative the chordal metric) for g is given by $L(g) = p^{\rho(\zeta_{\text{Gauss}}, g(\zeta_{\text{Gauss}}))}$, namely $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. Furthermore, there exist at least two points $z, w \in \mathbb{P}^1(\mathbb{C}_p)$ such that $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$.*

Proof. We can assume that $g(z) = \frac{az+b}{cz+d}$. The element g has at least one fixed point a_g . If $|a_g| \leq 1$, let $h(z) = z - a_g$ and $\iota(z) = 1/z$, and then $\iota h g h^{-1} \iota^{-1}$ fixes ∞ . Since h and ι fix the point ζ_{Gauss} , we have

$$\rho(\iota h g h^{-1} \iota^{-1}(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}}) = \rho(\iota h g(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}}) = \rho(g(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}})$$

which yields that $L(g) = L(\iota h g h^{-1} \iota^{-1})$. If $|a_g| > 1$, let $h(z) = z - 1/a_g$ and $\iota(z) = 1/z$. Thus $\iota h g \iota h^{-1} \iota^{-1}$ fixes ∞ . Similarly, since h and ι fix the point ζ_{Gauss} , we have

$$\rho(\iota h g \iota h^{-1} \iota^{-1}(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}}) = \rho(\iota h g(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}}) = \rho(g(\zeta_{\text{Gauss}}), \zeta_{\text{Gauss}})$$

which yields that $L(g) = L(\iota h g \iota h^{-1} \iota^{-1})$. Therefore, without loss of generality, we can assume that $g(z) = \frac{az+b}{d}$ with $ad = 1$. If $|a| > 1$, we consider the inverse $g^{-1}(z) = \frac{d}{a}z - \frac{b}{a}$, since $L(g) = L(g^{-1})$. Thus we can assume that $|a| \leq 1$.

For any $z, w \in \mathbb{P}^1(\mathbb{C}_p)$, we have

$$\begin{aligned} \rho_v(g(z), g(w)) &= \frac{|g(z) - g(w)|}{\max\{1, |g(z)|\} \max\{1, |g(w)|\}} \\ &= \frac{|z - w|}{\max\{|az + b|, |d|\} \max\{|aw + b|, |d|\}} \\ &= \frac{|z - w|}{\max\{1, |z|\} \max\{1, |w|\}} \frac{\max\{1, |z|\} \max\{1, |w|\}}{\max\{|az + b|, |d|\} \max\{|aw + b|, |d|\}} \\ &= \rho_v(z, w) \frac{\max\{1, |z|\} \max\{1, |w|\}}{\max\{|az + b|, |d|\} \max\{|aw + b|, |d|\}} \\ &= \rho_v(z, w) \frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}}. \end{aligned}$$

When $|a| = 1$, let $D_1 = D(-\frac{b}{a}, 1)$ and $D_2 = D(0, 1)$. If $D_1 = D_2$, then $|\frac{b}{a}| \leq 1$ which yields that $|b| \leq |a| = 1$, namely $L(g) = 1$. If $z, w \in D_1 = D_2$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = 1,$$

namely $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $z \in D_1 = D_2$ and $w \notin D_1 = D_2$,

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{|w|}{|w + b/a|}.$$

Since $|\frac{b}{a}| \leq 1$, we have $|w| = |w + \frac{b}{a}|$ which yields that $\frac{|w|}{|w + b/a|} = 1$. Thus $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $z, w \notin D_1 = D_2$, we have

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{|w||z|}{|z + b/a||w + b/a|}.$$

Since $|\frac{b}{a}| \leq 1$, we have $|w| = |w + \frac{b}{a}|$ and $|z| = |z + \frac{b}{a}|$ which yields that

$$\frac{|z||w|}{|z + b/a||w + b/a|} = 1.$$

Thus $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $D_1 \cap D_2 = \emptyset$, then $|\frac{b}{a}| > 1$ which yields that $|b| > |a| = 1$, namely $L(g) = |b|^2$. If $z, w \in D_1$, then $|z| > 1, |w| > 1, |z + b/a| \leq 1, |w + b/a| \leq 1$ which yields that

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \left|\frac{b}{a}\right|^2 = |b|^2.$$

Thus $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $z, w \in D_2$, then $|z| \leq 1, |w| \leq 1, |z + b/a| < 1, |w + b/a| < 1$ which yield that

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{1}{|z + b/a||w + b/a|} = \left|\frac{a}{b}\right|^2 = |b|^{-2} \leq L(g).$$

Thus $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. If $z \in D_1$ and $w \in D_2$, then $|z| > 1, |w| \leq 1, |z + b/a| \leq 1, |w + b/a| > 1$ which yields that

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{|z|}{|w + b/a|} = 1 \leq L(g).$$

If $z \notin D_1 \cup D_2$, then $|z| > 1, |z + b/a| > 1$ which yields that

$$\frac{\max\{1, |z|\}}{\max\{|a^2z + ab|, 1\}} = \frac{|z|}{|z + b/a|} = 1.$$

Thus if $z, w \notin D_1 \cup D_2$, then $\rho_v(g(z), g(w)) = \rho_v(z, w) \leq L(g)\rho_v(z, w)$. If $z \notin D_1 \cup D_2, w \in D_1$, then $\rho_v(g(z), g(w)) = \rho_v(z, w)|w| = |b|\rho_v(z, w) \leq L(g)\rho_v(z, w)$. If $z \notin D_1 \cup D_2, w \in D_2$, then $\rho_v(g(z), g(w)) = \rho_v(z, w)/|w + b/a| = |a/b|\rho_v(z, w) \leq L(g)\rho_v(z, w)$. Therefore, $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$.

When $|a| < 1$, let $D_1 = D(-\frac{b}{a}, \frac{1}{|a|^2})$ and $D_2 = D(0, 1)$. If $D_2 \subset D_1$, then $|\frac{b}{a}| \leq \frac{1}{|a|^2}$ which yields that $L(g) = |d|^2 = |a|^{-2}$. If $z, w \in D_1$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = |a|^2 \max\{1, |z|\} \max\{1, |w|\} \leq \frac{1}{|a|^2} = L(g),$$

namely $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. If $z, w \notin D_1$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{|a|^2|z||w|}{|a|^4|z||w|} = |a|^{-2}$$

which yields that $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $z \in D_1, w \notin D_1$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{|a^2z + ab|, 1\} \max\{|a^2w + ab|, 1\}} = \frac{|a|^2|w| \max\{1, |z|\}}{|a|^2|w|} = \max\{1, |z|\} \leq \frac{1}{|a|^2} = L(g).$$

Thus $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. If $D_1 \cap D_2 = \emptyset$, then $|\frac{b}{a}| > \frac{1}{|a|^2}$ which yields that $|b| > \frac{1}{|a|}$. Thus $L(g) = |b|^2$. If $z \in D_2$, then

$$\frac{\max\{1, |z|\}}{\max\{1, |a^2z + ab|\}} \leq 1.$$

If $z \in D_1$, then

$$\frac{\max\{1, |z|\}}{\max\{1, |a^2z + ab|\}} = \frac{|b|}{|a|}.$$

If $z \notin D_1 \cup D_2$, then

$$\frac{\max\{1, |z|\}}{\max\{1, |a^2z + ab|\}} = \frac{|z|}{|a^2z + ab|}.$$

Thus if $z, w \in D_1$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{1, |a^2 z + ab|\} \max\{1, |a^2 w + ab|\}} = |b|^2$$

which yields that $\rho_v(g(z), g(w)) = L(g)\rho_v(z, w)$. If $z, w \in D_2$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{1, |a^2 z + ab|\} \max\{1, |a^2 w + ab|\}} \leq |a|^2 \leq 1 \leq L(g).$$

Thus $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. If $z_1 \in D_1, w \in D_2$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{1, |a^2 z + ab|\} \max\{1, |a^2 w + ab|\}} \leq |a||b| \leq L(g).$$

Thus $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. If $z, w \notin D_1 \cup D_2$, then

$$\frac{|a|^2 \max\{1, |z|\} \max\{1, |w|\}}{\max\{1, |a^2 z + ab|\} \max\{1, |a^2 w + ab|\}} = \frac{|a|^2 |z||w|}{|a^2 z + ab||a^2 w + ab|}.$$

If $|z| = |z + \frac{b}{a}|$, then $\frac{|z|}{|z + b/a|} = 1$. If $|z| < |z + b/a|$, then $\frac{|z|}{|z + b/a|} < 1$. If $|z| > |z + b/a|$, then $|z| = |b/a|$. Therefore, we have

$$\frac{|a|^2 |z||w|}{|a^2 z + ab||a^2 w + ab|} \leq |b|^2.$$

Thus $\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$. □

Remark 4.4. In [33], the Lipschitz constant of the rational map with respect to the chordal metric is derived by the resultant (see definition below). Let $\mathcal{O}_p^* = \{\alpha \in \mathbb{C}_p : |\alpha| = 1\}$. Let $g(z) = a_0 z^n + a_1 z^{n-1} + \dots + a_{n-1} z + a_n$ and $h(z) = b_0 z^m + b_1 z^{m-1} + \dots + b_{m-1} z + b_m$. The rational function $\phi(z) = \frac{g(z)}{h(z)} \in \mathbb{C}_p(z)$ is called a normalized form if all the coefficients of $g(z)$ and $h(z)$ are in \mathcal{O}_p and at least one coefficient of $g(z)$ and $h(z)$ are in \mathcal{O}_p^* . Let $\alpha_i, 1 \leq i \leq n$ be the roots of $g(z)$, and $\beta_j, 1 \leq j \leq m$ be the roots of $h(z)$. Let $\text{Res}(g(z), h(z)) = a_0^n b_0^m \prod_{i=1}^n \prod_{j=1}^m (\alpha_i - \beta_j)$ be the resultant of two polynomials, and $\text{Res}(\phi) = \text{Res}(g(z), h(z))$ be the resultant of the rational function ϕ . The absolute value of the resultant $\text{Res}(\phi)$ depends only on the map ϕ .

Lemma 4.5 ([33]). *Let $\phi : \mathbb{P}^1(\mathbb{C}_p) \rightarrow \mathbb{P}^1(\mathbb{C}_p)$ be a rational map. Then $\rho_v(\phi(z), \phi(w)) \leq |\text{Res}(\phi)|^{-2} \rho_v(z, w)$ for all $v, w \in \mathbb{P}^1(\mathbb{C}_p)$.*

Let $g(z) = \frac{az+b}{cz+d} \in \text{PSL}(2, \mathbb{C}_p)$. By Lemma 4.5, we have

$$\text{Res}(g(z)) = \text{Res}((a/t)z + b/t, (c/t)z + d/t) = 1 / \|g\|^2 = L(g),$$

where $t = \max\{|a|, |b|, |c|, |d|\} = \|g\|$.

In [33], it is shown that there are two points $x, y \in \mathbb{P}^1(\mathbb{C}_p)$ such that $\sup_{x \neq y} \rho_v(\phi(x), \phi(y)) / \rho_v(x, y) = |\text{Res}(\phi)|^{-2}$. Since $\mathbb{P}^1(\mathbb{C}_p)$ is not compact, the supreme can not be omitted in general cases. However, we get the Lipschitz

constant by other method and show that we can get the supreme when it is the p -adic Möbius map.

Proof of Theorem 1.1

Proof. Since $\rho(g(\zeta_{Gauss}), \zeta_{Gauss}) = 2 \log_p \|g\|$, $\|g\| = 1$ is equivalent to $\rho(g(\zeta_{Gauss}), \zeta_{Gauss}) = 0$ which yields that $L(g) = 1$. The converse is also true. This means that (1), (2), (3) are equivalent.

(2) \Rightarrow (4) If $L(g) = 1$, $L(g^{-1}) = 1$. Hence by Lemma 4.3, we have $\rho_v(z, w) = \rho_v(g^{-1}(g(z)), g^{-1}(g(w))) \leq L(g^{-1})\rho_v(g(z), g(w)) \leq L(g)\rho_v(z, w)$ which yields that $\rho_v(g(z), g(w)) = \rho_v(z, w)$.

(4) \Rightarrow (2) If g is a chordal isometry, then $\rho_v(g(z), g(w)) = \rho_v(z, w)$. By Lemma 4.3, we have $L(g) = 1$.

(1) \Rightarrow (5) Since $\|g\| = 1$, let $g = \frac{az+b}{cz+d}$, we have $\max\{|a|, |b|, |c|, |d|\} = 1$ which yields that $g \in \text{PSL}(2, \mathcal{O}_p)$.

(5) \Rightarrow (1) If $g \in \text{PSL}(2, \mathcal{O}_p)$, let $g = \frac{az+b}{cz+d}$, then $\max\{|a|, |b|, |c|, |d|\} \leq 1$ and $ad-bc = 1$. If $\max\{|a|, |b|, |c|, |d|\} < 1$, then $|ad-bc| < \max\{|ad|, |bc|\} < 1$. This is a contradiction. Hence $\max\{|a|, |b|, |c|, |d|\} = 1$ which yields that $\|g\| = 1$.

(1) \Rightarrow (6) Since $\|g\| = 1$, $\|g^{-1}\| = 1$. Since $\|gh\| \leq \|g\| \|h\| \leq \|h\|$, and $\|h\| = \|g^{-1}gh\| \leq \|g^{-1}\| \|gh\| \leq \|gh\|$, we have $\|gh\| = \|h\|$. Similarly, $\|hg\| = \|h\|$. We can rewrite h as hg^{-1} , and then we have $\|ghg^{-1}\| = \|hg^{-1}\| = \|h\|$, since $g \in \text{PSL}(2, \mathcal{O}_p)$.

(6) \Rightarrow (1) Let $h \in \text{PSL}(2, \mathcal{O}_p)$. Then $\|h\| = \|gh\| = \|g\|$ which yields that $\|g\| = 1$. □

The metric properties of $\text{PSL}(2, \mathcal{O}_p)$ can be used to study the p -adic continued fractions. An *infinite p -adic continued fraction* is a formal expression

$$b_1 + \frac{a_1}{b_2 + \frac{a_2}{b_3 + \frac{a_3}{\dots}}}$$

where $a_i, b_i \in \mathbb{C}_p$ and $a_i \neq 0$. We denote this continued fraction by $\mathbb{K}(a_n|b_n)$. Let $t_n = \frac{a_n}{z+b_n}$ and $T_n = t_1 \circ t_2 \circ t_3 \circ \dots$ for $n = 1, 2, 3, \dots$. The continued fraction is said to be convergent classically if the sequence $\{T_n\}$ converges, else it is said to diverge classically. In the following part, we study the simplest case $\mathbb{K}(1|b_i)$ with $|b_i| \leq 1$. Since $T_{n+1}(\infty) = T_n(0)$, by Theorem 1.1, if the sequence $\{T_n(0)\}$ converges, then $0 = \lim_{n \rightarrow \infty} \rho_v(T_n(0), T_n(\infty)) = \rho_v(0, \infty) = 1$. This is a contradiction. This implies that the continued fraction $\mathbb{K}(1|b_i)$ with $|b_i| \leq 1$ diverge classically.

Example 4.6.

$$\frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}}$$

does not converge classically.

This example shows that the convergence and divergence of p -adic continued fractions are different from those in complex settings.

5. REDUCTION AND p -ADIC MÖBIUS MAPS

We call an elliptic element $f \in \mathrm{PSL}(2, \mathbb{C}_p)$ of order 2 an **involution**. If $g \in \mathrm{PSL}(2, \mathbb{C}_p)$ is a loxodromic element or an elliptic element, g have two fixed points a_g and r_g in $\mathbb{P}^1(\mathbb{C}_p)$. We call the geodesic line A_g which connects a_g and r_g the **axis** of g . Let A be a geodesic line in $\mathbb{P}^1_{\mathrm{Ber}}$. A geodesic line B is **orthogonal** to A if there exists a p -adic Möbius transformation f which is an involution such that endpoints of B are two fixed points of f , and f interchanges endpoints of A .

We introduce a new conception, a **tailed geodesic line**, in order to analyze the geometrical characterization of 2-adic Möbius maps. Let A be a geodesic line and x be the point satisfying $\inf_{y \in A} \rho(x, y) = \log_p 2$. Choosing any point $y \in A$, $A_x = A \cup [x, y]$. Since the Berkovich space is a \mathbb{R} -tree, it is obviously that A_x is independent of the chosen point y , A is the geodesic line associated with A_x , and x is called a **tail**. If A is the axis of an involution f , then there exists a unique point x fixed by f such that A_x is a tailed geodesic line. Hence we call A_x the tailed axis of f .

Lemma 5.1. *Let g be a p -adic Möbius map. Then there exist two involutions $f, h \in \mathrm{PSL}(2, \mathbb{C}_p)$ such that $g = f \circ h$. Furthermore*

- (1) *The axes A of h and B of f are orthogonal to the axis A_g of g .*
- (2) *The endpoints of A are different from the endpoints of B .*
- (3) *The element g is parabolic if and only if A and B share the unique endpoint.*

When $p \geq 3$,

- (4) *the element g is elliptic if and only if $A \cap B \neq \emptyset$;*
- (5) *the element g is loxodromic if and only if $A \cap B = \emptyset$.*

When $p = 2$,

- (6) *the element g is elliptic if and only if the two tailed geodesic lines $A_x \cap B_y \neq \emptyset$;*
- (7) *the element g is loxodromic if and only if the two tailed axes $A_x \cap B_y = \emptyset$.*

Proof. If g is loxodromic or elliptic, without loss of generality, let $g = \lambda^2 z$. The axis A_g is the geodesic line connecting 0 and ∞ . Let $h(z) = -\frac{b^2}{z}$ and $f(z) = g \circ h^{-1}(z) = -\frac{\lambda^2 b^2}{z}$. Thus h has two fixed points $bk, -bk$ and f has two fixed points $\lambda bk, -\lambda bk$, where $k^2 = -1$. It is easy to see that f and

h interchange 0 and ∞ , and $f \circ f = h \circ h = z$, namely f and h are two involutions. Hence A, B are orthogonal to A_g respectively. We prove (1), (2).

In case of $p \geq 3$, if $A \cap B \neq \emptyset$, since the \mathbb{P}_{Ber}^1 is a \mathbb{R} -tree, there exists a point $x \in A \cap B$ corresponding the disc D , which contains endpoints $bk, -bk, \lambda bk, -\lambda bk$ and $|2bk| = |2\lambda bk|$ which implies that $|\lambda| = 1$, i.e. g is elliptic element. If $A \cap B = \emptyset$, either $bk, -bk \notin D(\lambda bk, |\lambda bk|)$ or $\lambda bk, -\lambda bk \notin D(bk, |bk|)$ which implies that $|\lambda bk - bk| > |bk|$ or $|\lambda bk - bk| > |\lambda bk|$, i.e. $|\lambda - 1| > 1$ or $|1 - \frac{1}{\lambda}| > 1$. Hence g is loxodromic. We prove (4),(5).

In case of $p = 2$, if the tailed axes $A_x \cap B_y = \emptyset$, we denote the tails of the tailed axes by $D(bk, |bk|)$ and $D(\lambda bk, |\lambda bk|)$, the two tailed axes A_x and B_y do not intersect, since $D(bk, |bk|)^- \cap D(\lambda bk, |\lambda bk|)^- = \emptyset$, and $D(\lambda bk, |\lambda bk|) \neq D(bk, |bk|)$. Conversely, if two tailed axes do not intersect, then $|\lambda bk| \neq |bk|$, namely $|\lambda| \neq 1$. This implies that g is a loxodromic element. We prove (6),(7)

If g is parabolic, without loss of generality, we can assume that $g(z) = z+1$. Let $f(z) = -z$ and $h(z) = -z-1$. Then $g(z) = f \circ h(z) = -(-z-1) = z+1$. Since the axis A of f is the geodesic line connecting $\{0, \infty\}$ and the axis B of h is the geodesic line connecting $\{-\frac{1}{2}, \infty\}$, we have $\infty \in A \cap B$.

Conversely, if A and B share only one endpoint, without loss of generality, we can assume that A is the geodesic line connecting $\{0, \infty\}$, and B is the geodesic line connecting $\{0, 1\}$. Then $f(z) = -z$ and $h(z) = \frac{z}{2z-1}$ which implies that $g(z) = f \circ h(z) = -\frac{z}{2z-1}$ is a parabolic element, since g has a unique fixed point in $\mathbb{P}^1(\mathbb{C}_p)$. We know that if A and B have different endpoints, then g is either loxodromic or elliptic. □

By the proof of Lemma 5.1, we know that for any p -adic Möbius map g , there exist two half turns f and h such that $g = fh$. Furthermore, following the proof, since f and h are not unique, we can make the axis of f contain the Gauss point ζ_{Gauss} .

We denote the set of fixed points of an element g by $F_g = \{x \in \mathbb{P}_{Ber}^1 | g(x) = x\}$. Let $a, b \in \mathbb{P}^1(\mathbb{C}_p)$, and $x \in \mathbb{H}_{Ber}$.

Proposition 5.2. *Let $g \in \text{PSL}(2, \mathbb{C}_p)$.*

(1) *If g is a loxodromic element, then the set of fixed points of g contains two points in $\mathbb{P}^1(\mathbb{C}_p)$.*

(2) *If g is a tame elliptic element, then the set of fixed points of g is a geodesic line in \mathbb{P}_{Ber}^1 , and $F_g \cap \mathbb{P}^1(\mathbb{C}_p)$ contains two points.*

(3) *Let g be a wild elliptic element. Then the interior of the set of the fixed points of g contains a geodesic line in \mathbb{P}_{Ber}^1 .*

(4) *If g is a parabolic element, then the fixed points of g is an open disc with its boundary with respect to the Berkovich topology.*

Proof. If g is loxodromic or elliptic, we can assume that $g(z) = \lambda^2 z$ with fixed points $0, \infty$. If $|\lambda| > 1$ or $|\lambda| < 1$, then by Lemma 4.2, we know that g can

not fix any point in \mathbb{H}_{Ber} . This proves (1). If $|\lambda| = 1$, then $g \in \text{PSL}(2, \mathcal{O}_p)$ which implies that g fixes every point on the geodesic line connecting $0, \infty$. Furthermore, in case of $|\lambda - 1| = |\lambda + 1| = 1$, for any $x \in \mathbb{H}_{Ber}$ corresponding to the disc $D(a, r)$, we have $g(D(a, r)) = D(\lambda^2 a, |\lambda^2| r) = D(\lambda^2 a, |\lambda^2| r)$, if $gx = x$. Then $|\lambda^2 - 1||a| \leq r$ which implies that $|a| \leq r$, i.e. $0 \in D(a, r)$. Hence x is on the geodesic line connecting 0 and ∞ . This proves (2). When $|\lambda^2 - 1| < 1$, for any $x \in \mathbb{H}_{Ber}$ corresponding to the disc $D(a, r)$, we have $g(D(a, r)) = D(\lambda^2 a, |\lambda^2| r) = D(\lambda^2 a, |\lambda^2| r)$, if $gx = x$. This implies that $|a| \leq \frac{r}{|\lambda^2 - 1|}$. Hence g fixes any point in the hyperbolic disc $\mathcal{B}(x, \frac{1}{|\lambda^2 - 1|})$, i.e. the interior of the set of the fixed points of g contains a geodesic line in \mathbb{P}_{Ber}^1 . This proves (3).

If g is parabolic, we can assume that $g = z + 1$. For any $x \in \mathbb{H}_{Ber}$ corresponding to the disc $D(a, r)$, if $g(x) = x$, then $D(a, r) = D(a + 1, r)$ which implies that $r \geq 1$. This proves (4). □

Lemma 5.3. *If a geodesic line A is orthogonal to the other geodesic line B , then B is also orthogonal to A .*

Proof. Without loss of generality, we can assume that A is a geodesic line with endpoints 0 and ∞ and B is a geodesic line with endpoints -1 and 1 . Then $f(A) = B, f(B) = A$, if $f(z) = \frac{z+1}{z-1}$. □

Lemma 5.4. *Let a geodesic line A be orthogonal to the other geodesic line B .*

- (1) *If $p \geq 3$, then A intersects B at one unique point.*
- (2) *If $p = 2$, then $A \cap B = \emptyset$.*

Proof. Without loss of generality, we may assume that A is the geodesic connecting 0 and ∞ and B is a geodesic line with endpoints $-\alpha, \alpha$. Let the point x correspond to the disc $D(\alpha, |2\alpha|)$ which lies on the geodesic line B and contains the points $-\alpha, \alpha$.

If $p \geq 3$, then $|2\alpha| = |\alpha|$ which implies that $D(\alpha, |2\alpha|)$ contains 0 . Hence B intersects A . Conversely, if B intersects A , then there exists a point x corresponding to a disc $D(0, r)$ containing α or $-\alpha$ which implies that $|\alpha| \leq r$. Hence $D(0, r)$ contains both α and $-\alpha$. If there exist two points $x_1, x_2 \in A \cap B$ corresponding to two discs $D(0, r_1)$ and $D(0, r_2)$ respectively, then either $D(0, r_1) \subset D(0, r_2)$ or $D(0, r_2) \subset D(0, r_1)$. Without loss of generality, we can assume that $D(0, r_1) \subset D(0, r_2)$. Let l_1 be the geodesic segment connecting α and x_2 , and l_2 be the geodesic segment connecting $-\alpha$ and x_2 . Hence $l_1 \cup l_2 \subset B$, but $l_1 \cap l_2$ contains a segment containing x_1 and x_2 . This is a contradiction. Hence A intersects B at a uniquely point.

If $p = 2$, then for any x lying on the geodesic line connecting $\alpha, -\alpha$, we have that the disc corresponding to x must contain α or $-\alpha$. Without loss of generality, let x correspond to the disc $D(\alpha, r)$. If x lies on A , then $D(\alpha, r)$ contains 0 which implies that $|\alpha| \leq r$. Hence $-\alpha \in D(0, r)$. Since ζ corresponds to the disc $D(\alpha, |2\alpha|)$ containing both α and $-\alpha$, the geodesic

line contains the segment which connecting ζ and x . This is a contradiction. Hence $A \cap B = \emptyset$. \square

We say that g keeps a set A invariant if $g(A) = g^{-1}(A) = A$.

Lemma 5.5. *Let A_g be the axis of g . If g is a loxodromic element or an elliptic element, then g keeps the axis A_g invariant. Furthermore, g fixes every point of the axis A_g if and only if g is an elliptic element.*

Proof. Without loss of generality, let $g = \lambda z$. Hence A_g is the geodesic line connecting $0, \infty$. If $|\lambda| > 1$, then g maps each disk $D(0, r)$ to $D(0, |\lambda|r)$ which is also on the geodesic line. If $|\lambda| = 1$, then g maps each disk $D(0, r)$ to $D(0, |\lambda|r)$ which is the disk $D(0, r)$, namely g fixes the point $\zeta_{0,r}$. \square

Lemma 5.6. *If $p \geq 3$, and $g = h \circ f$ is a tame elliptic element, where h, f are two involutions, then two axes of h and f only intersect at a unique point.*

Proof. Without loss of generality, we can assume that $g(z) = \lambda^2(z)$. Thus $f(z) = -\frac{b^2}{z}$ and $h(z) = -\frac{\lambda^2 b^2}{z}$. By Lemma 2.1, we have $|\lambda b - b| = |\lambda - 1||b| = |b|$. This implies that two axes of h and f only intersect at the point $\zeta_{0,|b|}$ which corresponds to the disc $D(0, |b|)$. \square

Lemma 5.7. *If $p \geq 3$, and $g = h \circ f$ is a wild elliptic element, where h, f are two involutions, then two axes of h and f only intersect on a segment, and this segment belongs to the fixed points of g .*

Proof. Without loss of generality, we can assume that $g(z) = \lambda^2 z$, $f(z) = -\frac{1}{z}$ and $h(z) = -\frac{\lambda}{z}$. Let A be the axis of $f(z)$ and B be the axis of $h(z)$. Hence the endpoints of A are $\{-1, 1\}$ and the endpoints of B are $\{-\sqrt{\lambda}, \sqrt{\lambda}\}$. Since $p \geq 3$, then $1 = |-1 - 1| = |-\sqrt{\lambda} - \sqrt{\lambda}|$ which implies that ζ_{Gauss} lies on both the axes A and B . Since $|\lambda - 1| < 1$ and $|\lambda + 1| < 1$, we have $\min\{|\sqrt{\lambda} - 1|, |\sqrt{\lambda} + 1|\} < 1$ which implies that there exists a point $x \in A \cap B \cap \mathbb{H}_{Ber}$ which corresponds to the disc $D(1, |\sqrt{\lambda} - 1|)$ or the disc $D(1, |\sqrt{\lambda} + 1|)$. Hence $A \cap B$ contains either the segment connecting $\zeta_{1,|\sqrt{\lambda}-1|}$ and $\zeta_{0,1}$ or the segment connecting $\zeta_{1,|\sqrt{\lambda}+1|}$ and $\zeta_{0,1}$. This segment belongs to the fixed points of fixed points of f . \square

Lemma 5.8. *If $p \geq 3$, and A and B are two geodesic lines with four distinct endpoints, then there exists a unique geodesic line which is orthogonal to A and B simultaneously.*

Proof. Without loss of generality, we can assume that A is the geodesic with endpoints 0 and ∞ , and B is the other geodesic line with endpoints a and b . If A does not intersect B , then we have $|a - b| < \max\{|a|, |b|\}$. By the ultrametric property, we have $|a| = |b| > |a - b|$. Let C be a geodesic

line with endpoints $\zeta, -\zeta$. By Theorem 5.1, we have that the geodesic line C is orthogonal to the geodesic line A . Let $g = \frac{z-a}{z-b}$. Then the geodesic line B is mapped to the geodesic line A by g . If the geodesic $g(C)$ is also orthogonal to A , then $g(-\zeta) + g(\zeta) = 0$ and $g(C) \cap A \neq \emptyset$. This implies that $\frac{\zeta-a}{\zeta-b} + \frac{-\zeta-a}{-\zeta-b} = 0$, namely $\zeta = \sqrt{ab}$. Then the geodesic line C connecting $-\sqrt{ab}, \sqrt{ab}$ is orthogonal to both A and B simultaneously. \square

Lemma 5.9. *If $p = 2$, and A_x and B_y are two tailed geodesic lines with four distinct endpoints, then there exists a unique geodesic line which is orthogonal to A and B simultaneously.*

Proof. Without loss of generality, we can assume that the endpoints of A are $-1, 1$, and the endpoints of B are t, s . We claim that we can find a p -adic Möbius map $f = \frac{az+b}{cz+d}$ such that $f(-1) = -1, f(1) = -1, f(t) + f(s) = 0$.

Since $f(-1) = -1, f(1) = -1$, we have $a = d, b = c$, and $\frac{at+b}{ct+d} + \frac{as+b}{cs+d} = 0$ which yields that $2abst + (a^2 + b^2)(s+t) + 2ba = 0$. We can lift the solution to the projective space, namely $2ABst + (A^2 + B^2)(s+t) + AB = 0$, and $A^2 - B^2 = C^2$. Since any two curves in the projective space intersect, we have solutions in the projective space. If the solution is $(A : B : 0)$, namely $C = 0$, then $A = B$ or $A = -B$. This implies that $st + s + t + 1 = 0$ or $st - (s+t) + 1 = 0$, namely $s = -1$ or $t = -1$ or $s = 1$ or $t = 1$. This contradicts that two tailed geodesic line have no common endpoints. Hence $C \neq 0$, namely there exists p -adic Möbius map f such that $f(-1) = -1, f(1) = -1, f(t) + f(s) = 0$.

Hence we can assume that the tailed geodesic line A has the endpoints $-1, 1$ and the tailed geodesic line has the endpoints $-\lambda, \lambda$. Then the two tailed geodesic line are orthogonal to the line connecting $0, \infty$ simultaneously. \square

Lemma 5.10. *If $p = 2$, and A_x is a tailed geodesic line with the tail $x \in B$, and $A \cap B = \emptyset$, then there exists a tailed geodesic line B_y such that B_y is orthogonal to A , $y \in A$ and $B \subset B_y$.*

Proof. Let y be the point on the geodesic line A satisfying $\rho(x, y) = \log 2$, and l be the segment connecting x and y such that $B_y = l \cup B$ is the tailed geodesic line satisfying the condition. \square

Lemma 5.11. *If $p = 2$, and $g = h \circ f$ is a tame elliptic element, where h, f are two involutions, then two tailed geodesic lines of h and f only intersect at a unique point.*

Proof. By Lemma 5.9, we can assume that the fixed points of f are $-1, 1$ and the fixed points of h are $-\lambda, \lambda$. Since g is a tame elliptic element, we have $|\lambda - 1| = 1$. Hence the tailed point of the tailed geodesic line of h is

$\zeta_{0,1}$, and the tailed point of the tailed geodesic line of f is also $\zeta_{0,1}$. Since $D(1,1)^- \cap D(\lambda,1)^- = \emptyset$, then the two tailed geodesic lines intersect the unique point $\zeta_{0,1}$. \square

We give the following lemma without proof, which follows from Lemma 5.7 and Lemma 5.9 directly.

Lemma 5.12. *If $p = 2$, and $g = h \circ f$ is a wild elliptic element, where h, f are two involutions, then two tailed axes A_x, B_y of h and f only intersect on a segment, and this segment belongs to the fixed points of g .*

Proof of Theorem 1.2

Proof. If $\bigcap_{g \in G} F_g = \emptyset$, then there exist finitely many elements g_1, \dots, g_n such that $\bigcap_{i=1}^n F_{g_i} = \emptyset$, since the Berkovich space is compact with respect to the weak topology. Hence if we can show that $\bigcap_{i=1}^n F_{g_i} \neq \emptyset$ for any positive integer n , then we prove the theorem.

Let f, g be two elliptic elements, and denote the axes of f, g by A_f, A_g respectively. When $p \geq 3$, by Lemma 5.8, we have that there exists an involution a whose axis A is orthogonal to A_f and A_g simultaneously. By Lemma 5.1, there exist two involutions b, c such that $f = a \circ b$ and $g = a \circ c$. We denote the set of the fixed points of f, g by F_f, F_g respectively, and the axes of a, b, c by A, B, C respectively. By Lemma 5.7, we know that $F_f \supset A \cap B \neq \emptyset$, and $F_g \supset A \cap C \neq \emptyset$, and $B \cap C \neq \emptyset$, since $h = f^{-1} \circ g$ is elliptic. Choosing $x \in A \cap B, y \in A \cap C, z \in B \cap C$, there exists $w \in [x, z] \cap [y, z] \cap [x, y]$, since \mathbb{P}_{Ber}^1 is an \mathbb{R} -tree. This means that $F_f \cap F_g \neq \emptyset$.

By induction, $\bigcap_{i=1, i \neq k}^n F_{g_i} \neq \emptyset$ for $k = 1, \dots, n$, and then we want to show $\bigcap_{i=1}^n F_{g_i} \neq \emptyset$. Since $\bigcap_{i=1, i \neq n-1}^n F_{g_i} \neq \emptyset, \bigcap_{i=1, i \neq n}^n F_{g_i} \neq \emptyset, F_{g_{n-1}} \cap F_{g_n} \neq \emptyset$, choosing $x \in \bigcap_{i=1, i \neq n-1}^n F_{g_i}, y \in \bigcap_{i=1, i \neq n}^n F_{g_i}, z \in F_{g_{n-1}} \cap F_{g_n}$, there exists $w \in [x, z] \cap [y, z] \cap [x, y]$ such that $w \in \bigcap_{i=1}^n F_{g_i}$. This implies that each element in G share at least one fixed point.

When $p = 2$, by Lemma 5.8, we have that there exists an involution a whose axis A is orthogonal to A_f and A_g simultaneously. By Lemma 5.1, there exist two involutions b, c such that $f = a \circ b$ and $g = a \circ c$. We denote the set of the fixed points of f, g by F_f, F_g . Thanks to Lemma 5.9 and Lemma 5.10, there exist two tailed geodesic line A_x, A_y who share the same associated geodesic line A which are orthogonal to two axes A_f and A_g . We denote the tailed axes of b, c by B_x, C_y . By Lemma 5.12, we know that $F_f \supset A_x \cap B_x \neq \emptyset$, and $F_g \supset A_y \cap C_y \neq \emptyset$, and $B_x \cap C_y \neq \emptyset$, since $h = f^{-1} \circ g$ is elliptic. Choosing $u \in A_x \cap B_x, v \in A_y \cap C_y, w \in B_x \cap C_y$, there exists $\omega \in [u, v] \cap [u, w] \cap [v, w]$, since \mathbb{P}_{Ber}^1 is an \mathbb{R} -tree. This means that $F_f \cap F_g \neq \emptyset$.

Following the proof of the case $p = 2$, it is obviously that when $p = 2$, each element in G share at least one fixed point.

By conjugation, we can assume that each element in G shares the unique fixed point ζ_{Gauss} . By Lemma 2.11, we know that each element in G has

good reduction which yields that G has a potentially good reduction. By Lemma 2.10, we know that if g has good reduction, then $g \in \mathrm{PSL}(2, \mathcal{O})$. Since each element $f \in G$ can be written as $\phi f' \phi^{-1}$, where f' has good reduction and $\phi \in \mathrm{PSL}(2, \mathbb{C}_p)$, $\rho_v(f(x), f(y)) = \rho_v(\phi f' \phi^{-1}(x), \phi f' \phi^{-1}(y)) \leq L_1 \rho_v(f' \phi^{-1}(x), f' \phi^{-1}(y)) \leq L_1 \rho_v(\phi^{-1}(x), \phi^{-1}(y)) \leq L_1 L_2 \rho_v(x, y)$, where L_1, L_2 depending only on ϕ . Hence G is equicontinuous on $\mathbb{P}^1(\mathbb{C}_p)$. \square

Theorem 5.13. *If G is a discrete subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$ and the limit set of G is empty, then G has potentially good reduction.*

Proof. We have that G contains no loxodromic element g , since the fixed points of g are in the limit set of G which yields that G contains parabolic elements and elliptic elements only. Since G is a discrete subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$, G contains no parabolic elements. By Theorem 1.2, we know that G has potentially good reduction. \square

Example 5.14. *Let $f_n(z) = z + p^{-n}$ ($n \geq 1$), and the group G is generated by each f_n . Then G contains parabolic elements only and does not have potentially good reduction.*

Proof. For any disc $D(a, r)$ which is fixed by f_n , we have $r \geq p^n$. Since n is arbitrary, the only point fixed by G is the ∞ . Since each generator can commute with each other, we know that each element in G can only fixed the unique point ∞ in \mathbb{P}_{Ber}^1 . \square

Example 5.15. *Let $G \subset \mathrm{PSL}(2, \mathbb{C}_p)$, and ζ_i be the p^i -th primitive root of unity. Suppose that G is generated by*

$$g_i = \begin{pmatrix} \zeta_i & 0 \\ 0 & \zeta_i^{-1} \end{pmatrix},$$

for all the positive integer $i \geq 1$. Then G is discrete, and the limit set $\Lambda(G)$ of G is $\{0, \infty\}$ is a compact set.

Proof. In [30], we have proved that G is a discrete subgroup of $\mathrm{PSL}(2, \mathbb{C}_p)$. Furthermore, the points $\{0, \infty\}$ are the fixed points of all the elements g_i , $i \geq 1$, namely $0, \infty$ are the limit sets of G . \square

6. NORMS OF p -ADIC MÖBIUS MAPS AND ITS APPLICATIONS

Proposition 6.1. *Suppose that $f, g, h \in \mathrm{PSL}(2, \mathbb{C}_p)$. Then*

- (1) $\rho_0(fh, gh) = \rho_0(f, g)$, and $\rho_0(hf, hg) \leq L(h)\rho_0(f, g)$.
- (2) If $h \in \mathrm{PSL}(2, \mathcal{O}_p)$, $\rho_0(h^{-1}fh, h^{-1}gh) = \rho_0(f, g)$.

Proof. Since h is an automorphism on $\mathbb{P}^1(\mathbb{C}_p)$, we have

$$\rho_0(fh, gh) = \sup_{z \in \mathbb{P}^1(\mathbb{C}_p)} \rho_v(fh(z), gh(z)) = \sup_{w=h(z) \in \mathbb{P}^1(\mathbb{C}_p)} \rho_v(f(w), g(w)) = \rho_0(f, g).$$

Since $\rho_v(h(z), h(w)) \leq L(h)\rho_v(z, w)$, we have $\rho_v(hf(z), hg(z)) \leq L(h)\rho_v(f(z), g(z))$ which yields that $\rho_0(hf, hg) \leq L(h)\rho_0(f, g)$.

Since

$$\rho_v(h^{-1}fh, h^{-1}gh) = \rho_v(h^{-1}f, h^{-1}g) \leq L(h)\rho_v(f, g) \leq L(h)L(h^{-1})\rho_v(h^{-1}f, h^{-1}g),$$

we have $\rho_v(h^{-1}f, h^{-1}g) \leq L(h)\rho_v(f, g) \leq \rho_v(h^{-1}f, h^{-1}g)$ which yields that $\rho_0(h^{-1}fh, h^{-1}gh) = \rho_0(f, g)$. \square

Let $m(g) = \|g - g^{-1}\|$ and $M(g) = \frac{\|g - g^{-1}\|}{\|g\|}$.

Proposition 6.2. *Let p be a prime number. Then $p^{-\frac{1}{p-1}} \geq 2^{-1}$.*

Proof. Let $f(x) = x^{-\frac{1}{x-1}}$. Then $f'(x) = f(x)(\ln x - (1 - \frac{1}{x}))/x^2$ which yields that $f'(x) > 0$ if $x \geq 3$. Since $f(3) = 3^{-\frac{1}{2}} \geq 2^{-1} = f(2)$, we have $p^{-\frac{1}{p-1}} \geq 2^{-1}$ if p is a prime number. \square

Theorem 6.3. *Let $p \geq 3$, and g be a p -adic Möbius map. Then $\rho_0(g, I) = M(g)$.*

Proof. By Lemma 4.3, there exists an element $h \in \text{PSL}(2, \mathcal{O}_p)$ such that $hgh^{-1} = \frac{az+b}{d}$ with $ad = 1$. By Proposition 6.1, we have $\rho_0(hgh^{-1}, I) = \rho_0(g, I)$. By Theorem 1.1, we know that $\|g - g^{-1}\| = \|h(g - g^{-1})h^{-1}\| = \|hgh^{-1} - hg^{-1}h^{-1}\|$. Thus we can rewrite hgh^{-1} as g . Hence

$$M(g) = \frac{\max\{|a-d|, |b|\}}{\max\{|a|, |d|, |b|\}}.$$

Moreover

$$\rho_v(g(z), z) = \frac{|a^2z + ab - z|}{\max\{1, |z|\} \max\{1, |a^2z + ab|\}}.$$

If g is parabolic, then

$$\rho_v(g(z), z) = \frac{|b|}{\max\{1, |z|\} \max\{1, |z+b|\}} \leq M(g),$$

and

$$\rho_0(g, I) \geq \rho_v(g(0), 0) = M(g).$$

Thus $\rho_0(g, I) = M(g)$.

If g is loxodromic, we can assume that $|a| > 1$. Since

$$\rho_v(g(0), 0) = \frac{|ab|}{\max\{1, |ab|\}}, \rho_v(g(1), 1) = \frac{|a^2 + ab - 1|}{\max\{1, |a^2 + ab|\}} = \frac{|a^2 + ab|}{\max\{1, |a^2 + ab|\}},$$

$$\max\{|ab|, |a^2 + ab|\} \geq 1,$$

we have $M(g) = 1 \geq \rho_0(g, I) \geq \max\{\rho_v(g(0), 0), \rho_v(g(1), 1)\} = 1 = M(g)$.

If g is elliptic, we have $|a^2 - 1| \leq 1$. If $|ab| > |a^2 - 1|$, then $M(g) = |ab| \geq \rho_0(g, I) \geq \rho_v(g(0), 0) = |ab| = M(g)$. If $|ab| \leq |a^2 - 1|$, there exists a number $\omega \in \mathcal{O}_p$ such that $|(a^2 - 1)\omega + ab| = |a^2 - 1|$ which yields that

$$M(g) = |a^2 - 1| \geq \rho_0(g, I) \geq \rho_v(g(\omega), \omega) = |a^2 - 1| = M(g).$$

□

If $p = 2$, then $2^{-1}M(g) \leq \frac{\max\{|a-d|, |b|\}}{\max\{|a|, |d|, |b|\}} \leq 2M(g)$. Thus we give the following theorem without proof.

Theorem 6.4. *Let $p = 2$, and g be a p -adic Möbius map. Then $2^{-1}M(g) \leq \rho_0(g, I) \leq 2M(g)$.*

Theorem 6.5. *For any p -adic Möbius map g , $\rho_0(g, I) \leq \|g - I\|$.*

Proof. Following Theorem 6.3, we can assume that $g(z) = \frac{az+b}{d}$ with $ad = 1$. Thus $\|g - I\| = \max\{|a - 1|, |b|, |d - 1|\}$. Moreover

$$\rho_v(g(z), z) = \frac{|a^2z + ab - z|}{\max\{1, |z|\} \max\{1, |az + b|\}} \leq 1.$$

If g is loxodromic, we can assume $|a| > 1$. Hence $\max\{|a - 1|, |d - 1|\} > 1$ which yields that $\rho_0(g, I) \leq \|g - I\|$.

If g is parabolic, then $a = 1$. We have

$$\rho_v(g(z), z) = \frac{|b|}{\max\{1, |z|\} \max\{1, |z + b|\}} \leq |b| = \|g - I\|.$$

If g is elliptic, then $|a| = 1$. We have

$$\begin{aligned} \rho_v(g(z), z) &= \frac{|a^2z + ab - z|}{\max\{1, |z|\} \max\{1, |az + b|\}} \leq \frac{|a^2z + ab - z|}{\max\{1, |z|\}} \\ &\leq \frac{\max\{|(a^2 - 1)z|, |ab|\}}{\max\{1, |z|\}} \leq \max\{|a^2 - 1|, |ab|\} \leq \|g - I\|. \end{aligned}$$

□

Let $\varepsilon(g) = \max\{\rho_v(g(z_0), z_0), \rho_v(g(z_1), z_1), \rho_v(g(z_2), z_2)\}$, where z_0, z_1, z_2 are three distinct roots of the equation $z^3 = 1$.

Theorem 6.6. *For any p -adic Möbius map g , we have $2^{-1}\varepsilon(g) \leq M(g) \leq 6\varepsilon(g)$.*

Proof. Since $\varepsilon(g) = \max\{\rho_v(g(z_0), z_0), \rho_v(g(z_1), z_1), \rho_v(g(z_2), z_2)\} \leq \rho_0(g, I)$, by Theorem 6.3 and 6.4, we have $\varepsilon(g) \leq \rho_0(g, I) \leq 2M(g)$.

Since z_0, z_1, z_2 are three distinct roots of the equation $z^3 = 1$, and let $z_0 = 1$, we have $z_1 + z_2 + 1 = 0$, and by Lemma 2.1, Lemma 2.2 and Proposition 6.2, we have $2^{-1} \leq |z_1 - z_2| \leq 1$. This implies that

$$\begin{aligned} |3b| &= |cz_1 + (d - a)z_2 - b + cz_2 + (d - a)z_1 - b + cz_0 + (d - a)z_0 - b| \\ &\leq \max\{|cz_1 + (d - a)z_2 - b|, |cz_2 + (d - a)z_1 - b|, |cz_0 + (d - a)z_0 - b|\}. \end{aligned}$$

We denote $\varepsilon'(g)$ by $\max\{|cz_1 + (d-a)z_2 - b|, |cz_2 + (d-a)z_1 - b|, |cz_0 + (d-a)z_0 - b|\}$. Hence $|b| \leq 3\varepsilon'(g)$ which yields that

$$\max\{|cz_1 + (d-a)z_2|, |cz_2 + (d-a)z_1|, |cz_0 + (d-a)z_0|\} \leq 3\varepsilon'(g).$$

Thus

$\max\{|cz_1 + (d-a)z_2 + cz_2 + (d-a)z_1|, |cz_1 + (d-a)z_2 - cz_2 - (d-a)z_1|\} \leq 3\varepsilon'(g)$, namely $|c + (d-a)| \leq 3\varepsilon'(g)$ and $|c(\frac{z_1}{z_2}) + (d-a)| \leq 3\varepsilon'(g)$ which yields that $|c(\frac{z_1}{z_2} - 1)| \leq 3\varepsilon'(g)$. This implies that $|c| \leq 6\varepsilon'(g)$ and $|(d-a)| \leq 6\varepsilon'(g)$, namely $\max\{|a-d|, |b|, |c|\} \leq 6\varepsilon'(g)$. For any z with $|z| = 1$, we have $\max\{|az + b|, |cz + d|\} \leq \max\{|az|, |b|, |cz|, |d|\} = \|g\|$ which yields that $\max\{|a-d|, |2b|, |2c|\} \leq \max\{|a-d|, |b|, |c|\} \leq 6\varepsilon'(g)$. This implies that $\max\{|a-d|, |2b|, |2c|\} / \|g\| \leq \max\{|a-d|, |b|, |c|\} / \|g\| \leq 6\varepsilon'(g) / \max\{|az + b|, |cz + d|\} \max\{1, |z|\}$ for any z with $|z| = 1$. Since

$$\begin{aligned} \frac{\varepsilon'(g)}{\max\{|az + b|, |cz + d|\} \max\{1, |z|\}} &\leq \max\left\{ \frac{|cz_0^2 + (d-a)z_0 - b|}{\max\{|az_0 + b|, |cz_0 + d|\} \max\{1, |z_0|\}}, \right. \\ &\left. \frac{|cz_1^2 + (d-a)z_1 - b|}{\max\{|az_1 + b|, |cz_1 + d|\} \max\{1, |z_1|\}}, \frac{|cz_2^2 + (d-a)z_2 - b|}{\max\{|az_2 + b|, |cz_2 + d|\} \max\{1, |z_2|\}} \right\} \\ &\leq \varepsilon(g), \end{aligned}$$

we have $M(g) \leq 6\varepsilon(g)$. \square

Corollary 6.7. *For any p -adic Möbius map g , $\frac{1}{4}\varepsilon(g) \leq \rho_0(g, I) \leq \varepsilon(g)$.*

Proof. By Theorem 6.3 and Theorem 6.6, we get the inequality directly. \square

Let $\varepsilon_1(g) = \{\rho_v(g(0), 0), \rho_v(g(1), 1), \rho_v(g(\infty), \infty)\}$.

Theorem 6.8. *For any p -adic Möbius map, $2^{-1}\varepsilon_1(g) \leq M(g) \leq \varepsilon_1(g)$.*

Proof. Since $\varepsilon(g) = \max\{\rho_v(g(0), 0), \rho_v(g(1), 1), \rho_v(g(\infty), \infty)\} \leq \rho_0(g, I)$, by Theorem 6.3, we have $\varepsilon_1(g) \leq \rho_0(g, I) \leq 2M(g)$.

Since

$$\rho_v(g(z), z) = \frac{|cz^2 + (d-a)z - b|}{\max\{|az + b|, |cz + d|\} \max\{1, |z|\}},$$

we have

$$\rho_v(g(0), 0) = \frac{|b|}{\max\{|b|, |d|\}}, \rho_v(g(\infty), \infty) = \frac{|c|}{\max\{|a|, |c|\}},$$

and

$$\rho_v(g(1), 1) = \frac{|c + (d-a) - b|}{\max\{|a + b|, |c + d|\}}.$$

Since $\max\{|a-d|, |2b|, |2c|\} \leq \max\{|b|, |c|, |c + (d-a) - b|\}$ and $\max\{|a + b|, |c + d|\} \leq \max\{|a|, |c|, |b|, |d|\} = \|g\|$, we have

$$M(g) = \frac{\max\{|a-d|, |2b|, |2c|\}}{\|g\|} \leq \frac{\max\{|b|, |c|, |c + (d-a) - b|\}}{\|g\|}$$

$$\leq \max\left\{\frac{|b|}{\max\{|b|, |d|\}}, \frac{|c|}{\max\{|a|, |c|\}}, \frac{|c + (d - a) - b|}{\max\{|a + b|, |c + d|\}}\right\} = \varepsilon_1(g)$$

□

Let $\varepsilon_2(g) = \max\{\rho_v(g(0), 0), \rho_v(g(\infty), \infty)\}$.

Corollary 6.9. *If g is a parabolic element, then $2^{-1}\varepsilon_2(g) \leq M(g) \leq \varepsilon_2(g)$.*

Proof. Since $\varepsilon_2(g) < \varepsilon_1(g)$, we have $2^{-1}\varepsilon_2(g) \leq 2^{-1}\varepsilon_1(g) \leq M(g)$.

By Proposition 3.5, if g is a parabolic element, then $(a + d)^2 = 4$ which yields that $4bc = -(a - d)^2$. Thus it implies that $|a - d| = \sqrt{|4||bc|} \leq \sqrt{|bc|} \leq \max\{|b|, |c|\}$ which yields that $\max\{|a - d|, |2b|, |2c|\} \leq \max\{|b|, |c|\}$. Hence

$$M(g) = \frac{\max\{|a - d|, |2b|, |2c|\}}{\|g\|} \leq \frac{\max\{|a - d|, |b|, |c|\}}{\|g\|}$$

$$\leq \max\left\{\frac{|b|}{\max\{|b|, |d|\}}, \frac{|c|}{\max\{|a|, |c|\}}\right\} = \varepsilon_2(g) = \max\{\rho_v(g(0), 0), \rho_v(g(\infty), \infty)\}.$$

□

As applications of these inequalities, we can get the convergence theorem of p -adic Möbius maps.

Let $\{f_n\}$ be a sequence of p -adic Möbius maps, and \mathbf{U} be the set of points at which the sequence $\{f_n\}$ converges pointwisely, and $f = \lim_{n \rightarrow \infty} f_n$ on \mathbf{U} . Write $f_n \hookrightarrow (\mathbf{U}, f)$ to mean that \mathbf{U} is the set of convergence of f_n and that $f_n \rightarrow f$ on (and only on) \mathbf{U} . In [34], we proved the following theorem.

Theorem 6.10 ([34]). *Suppose that there exists a sequence p -adic Möbius maps f_n such that $f_n \hookrightarrow (\mathbf{U}, f)$ with $\mathbf{U} \neq \emptyset$. Then one of the following possibilities occurs:*

- (a) $\mathbf{U} = \mathbb{P}^1(\mathbb{C}_p)$, and f is a p -adic Möbius map;
- (b) $\mathbf{U} = \mathbb{P}^1(\mathbb{C}_p)$, and f is constant on the complement of one point on \mathbf{U} ;
- (c) $\mathbf{U} = \{z_1, z_2\}$ and $f(z_1) \neq f(z_2)$; or
- (d) f is constant on \mathbf{U} .

We can reprove this theorem by the use of the three-point norms.

If \mathbf{U} contains only one point, it is the case (d), and if \mathbf{U} contains two points only, it is the case (c) or (d). Hence we only need to consider the case when \mathbf{U} contains at least three points.

We prove the following theorem by using different norms of p -adic Möbius maps without using the cross ratios of p -adic Möbius maps.

Theorem 6.11. *Let $\{f_n\}$ be a sequence of p -adic Möbius maps and $z_j, j = 1, 2, 3$ be three distinct points with $f_n(z_j) \rightarrow w_j$, where w_j are also three distinct points. Then a sequence $\{f_n\}$ converge to a p -adic Möbius map f uniformly, where $f(z_j) = w_j, j = 1, 2, 3$.*

Proof. We can find a p -adic Möbius map h such that $h(z_1) = u_1, h(z_2) = u_2, h(z_3) = u_3$, where u_i are the three distinct roots of $z^3 = 1$. Then $hf_n h^{-1}(u_i) \rightarrow w_i$. By Corollary 6.7, we know that $\frac{1}{4}\varepsilon(hf^{-1}f_n h^{-1}) \leq \rho_0(hf^{-1}f_n h^{-1}, I) \leq \varepsilon(hf^{-1}f_n h^{-1})$. This yields that $hg^{-1}f_n h^{-1}$ converges to I uniformly, namely f_n converges to f uniformly. \square

Proposition 6.12 ([34]). *Let $f \in \text{PSL}(2, \mathbb{C}_p)$. Then f preserves the chordal cross ratio, namely*

$$\frac{\rho_v(f(x), f(y))\rho_v(f(z), f(w))}{\rho_v(f(x), f(z))\rho_v(f(y), f(w))} = \frac{\rho_v(x, y)\rho_v(z, w)}{\rho_v(x, z)\rho_v(y, w)}.$$

Theorem 6.13. *Let $\{f_n\}$ be a sequence of p -adic maps, and suppose that there exist three distinct points x_1, x_2, x_3 in $\mathbb{P}^1(\mathbb{C}_p)$ such that $\lim_{n \rightarrow \infty} f_n(x_1) = \infty$, $\lim_{n \rightarrow \infty} f_n(x_2) = \alpha$, $\lim_{n \rightarrow \infty} f_n(x_3) = \beta$, where $\alpha \neq \beta$. Then $f_n \rightarrow \alpha$ on $\mathbb{P}^1(\mathbb{C}_p) \setminus x_3$, namely $\mathbf{U} = \mathbb{P}^1(\mathbb{C}_p)$, and f is constant on the complement of one point on \mathbf{U} .*

Proof. Without loss of generality, we can assume that $x_1 = 0, x_2 = 1, x_3 = \infty$ and $\alpha = 0, \beta = \infty$. Assuming $x_4 \in \mathbb{P}^1(\mathbb{C}_p) \setminus \{0, 1, \infty\}$, if the sequence $\{f_n(x_4)\}$ does not converge to 0, there exists a subsequence $\{f_{n_j}(x_4)\}$ and a fixed positive number δ such that $|f_{n_j}(x_4)| > \delta$.

$$\frac{\rho_v(f_{n_j}(0), f_{n_j}(1))\rho_v(f_{n_j}(\infty), f_{n_j}(x_4))}{\rho_v(f_{n_j}(0), f_{n_j}(\infty))\rho_v(f_{n_j}(1), f_{n_j}(x_4))} = \frac{\rho_v(0, 1)\rho_v(\infty, x_4)}{\rho_v(0, \infty)\rho_v(1, x_4)} \neq 0.$$

Letting n_j tend to ∞ ,

$$0 = \lim_{n_j \rightarrow \infty} \frac{\rho_v(f_{n_j}(0), f_{n_j}(1))\rho_v(f_{n_j}(\infty), f_{n_j}(x_4))}{\rho_v(f_{n_j}(0), f_{n_j}(\infty))\rho_v(f_{n_j}(1), f_{n_j}(x_4))} = \frac{\rho_v(0, 1)\rho_v(\infty, x_4)}{\rho_v(0, \infty)\rho_v(1, x_4)} \neq 0.$$

This is a contradiction. Hence $\lim_{n \rightarrow \infty} f_n(x_4) = 0$. \square

Combing Theorem 6.11 and Theorem 6.13, we prove Theorem 6.10.

7. THE DECOMPOSITION THEOREM OF p -ADIC MÖBIUS MAPS

Two points α, β are called *antipodal* points if there exists an element $u \in \text{PSL}(2, \mathcal{O}_p)$ such that $u(0) = \alpha, u(\infty) = \beta$.

Theorem 7.1. *For any p -adic Möbius map g , there exists an element $u \in \text{PSL}(2, \mathcal{O}_p)$ such that $g = uf$, where either f is a loxodromic element with antipodal fixed points, or $f = I$.*

Proof. If g is a loxodromic element or elliptic element, by Lemma 5.1, there exist two involutions a, b such that $g = ab$, the (tailed) axes of a and b are orthogonal to the axis of g , and the (tailed) axis of a containing ζ_{Gauss} . Let α, β be the fixed points of b . We claim that there exists an element in $h \in \text{PSL}(2, \mathcal{O}_p)$ such that $h(\alpha) + h(\beta) = 0$.

Claim: Without loss of generality, we can assume that $|\alpha| \leq 1$, otherwise we can consider $1/\alpha$, since $1/z \in \text{PSL}(2, \mathcal{O}_p)$. Let $u(z) = z - \alpha \in \text{PSL}(2, \mathcal{O}_p)$ which yields that $u(\alpha) = 0$. Hence we can assume that $\alpha = 0$. If $|\beta| \leq 1$, then let $h(z) = z - \frac{\beta}{2}$ which implies that $h(0) + h(\beta) = 0$. If $|\beta| > 1$, then let $h(z) = \frac{az+b}{cz+d}$ with $h(0) + h(\beta) = 0$. Hence

$$\frac{b}{d} + \frac{a\beta + b}{c\beta + d} = 0$$

which yields that $\beta(ad + bc) + 2bd = 0$. Let $X_1 = ad, X_2 = bc, X_3 = bd$. Hence we have three equations:

$$(1) \quad \beta(X_1 + X_2) + 2X_3 = 0$$

$$(2) \quad X_1 - X_2 = 1$$

$$(3) \quad X_1 + X_2 = \lambda$$

Thus we have $X_1 = \frac{\lambda+1}{2}, X_2 = \frac{\lambda-1}{2}$, and $X_3 = -\frac{\lambda\beta}{2}$. Since $\frac{a}{b} = \frac{X_1}{X_3}$ and $\frac{c}{d} = \frac{X_2}{X_3}$, let $a = \frac{\lambda+1}{2}t, b = -\frac{\lambda\beta}{2}t, c = \frac{\lambda-1}{2}s, d = -\frac{\lambda\beta}{2}s$ which yields that $-\frac{\lambda\beta(\lambda+1)}{4}ts + \frac{\lambda\beta(\lambda-1)}{4}ts = 1$, namely $-\frac{\lambda\beta}{2}ts = 1$. Obviously, $\lambda = 1/\beta, t = 2$, and $s = 1$ is one of the solution of the equations with $\max\{|a|, |b|, |c|, |d|\} \leq 1$, namely $h \in \text{PSL}(2, \mathcal{O}_p)$. Hence we prove the claim.

Since $h(\alpha) + h(\beta) = 0$, the geodesic line connecting $0, \infty$ is orthogonal to the geodesic line connecting $h(\alpha), h(\beta)$, and this line contains ζ_{Gauss} . Let l_1 be the geodesic line which connects the endpoints of involution a . Since all the elements in $\text{PSL}(2, \mathcal{O}_p)$ fix the point ζ_{Gauss} , the point $\zeta_{Gauss} \in h(l_1)$. By Lemma 5.1, we know that there exists an involution c which fixes the point ζ_{Gauss} such that $chbh^{-1}$ is a loxodromic element whose fixed points are $0, \infty$, and $hah^{-1}c$ is an elliptic element in $\text{PSL}(2, \mathcal{O}_p)$, since $hah^{-1}c$ fixes the point ζ_{Gauss} . Thus $hgh^{-1} = hah^{-1}chbh^{-1}$ which yields that $g = h^{-1}hah^{-1}chh^{-1}(chbh^{-1})h$, where $h^{-1}hah^{-1}ch \in \text{PSL}(2, \mathcal{O}_p)$ and $h^{-1}(chbh^{-1})h$ is a loxodromic element with antipodal fixed points.

If g is a parabolic element, then we can assume the fixed point of g is ∞ after conjugating by an element in $\text{PSL}(2, \mathcal{O}_p)$ by the Claim. Thus $g(z) = ab$, where $a(z) = -z, b(z) = -z + b$. Since $a(z) = -z$ contains the point ζ_{Gauss} , by similar discussion above, we can find a involution c which fixes the point ζ_{Gauss} such that $az \in \text{PSL}(2, \mathcal{O}_p)$, and cb is a loxodromic element with antipodal fixed points.

□

Let $\mathcal{U} = \text{PSL}(2, \mathcal{O}_p)$. We define $d(g, \mathcal{U}) = \inf\{\rho_0(g, u) | u \in \mathcal{U}\}$.

Theorem 7.2. *For any p -adic Möbius map, either $d(g, \mathcal{U}) = 0$, if $g \in \mathcal{U}$, or $d(g, \mathcal{U}) = 1$, if $g \notin \mathcal{U}$.*

Proof. If $g \in \mathcal{U}$, then $d(g, \mathcal{U}) = 0$. If $g \notin \mathcal{U}$, then by Theorem 7.1, there exist $u \in \text{PSL}(2, \mathcal{O}_p)$ and a loxodromic element f with antipodal fixed points such that $g = uf$. Hence there exists an element $h \in \text{PSL}(2, \mathcal{O}_p)$ such that $v = hfh^{-1} = \lambda z$. Thus $d(g, \mathcal{U}) = d(uf, \mathcal{U}) = d(v, \mathcal{U})$.

For any $s(z) = \frac{az+b}{cz+d} \in \text{PSL}(2, \mathcal{O}_p)$, since $ad-bc = 1$ and $\max\{|a|, |b|, |c|, |d|\} \leq 1$, we have that $|\frac{b}{d}| = 1$, or $|\frac{a}{c}| = 1$, or $|\frac{b}{d}| > 1, |\frac{a}{c}| < 1$, otherwise, if $|\frac{b}{d}| > 1$ and $|\frac{a}{c}| > 1$, then $|d| < |b| \leq 1$ and $|c| < |a| \leq 1$ which yields that $|ad-bc| \leq \max\{|ad|, |bc|\} < 1$. This contradicts $ad-bc = 1$. Other cases are similar. If $|\frac{b}{d}| = 1$, then $\rho_v(v(0), s(0)) = 1$. If $|\frac{a}{c}| = 1$, then $\rho_v(v(\infty), s(\infty)) = 1$. If $|\frac{b}{d}| > 1, |\frac{a}{c}| < 1$, then $1 \geq |b| > |d|$ and $|a| < |c| \leq 1$. Since $ad-bc = 1$, we have $|b| = |c| = 1 > \max\{|a|, |d|\}$. This implies that $|\frac{d}{c}| < 1$.

$$\rho_v(v(-\frac{d}{c}), s(-\frac{d}{c})) = \frac{|\lambda c(-\frac{d}{c})^2 + (\lambda d - a)(-\frac{d}{c}) - b|}{\max\{1, |-\frac{d}{c}|\} \max\{|a(-\frac{d}{c}) + b|, |c(-\frac{d}{c}) + d|\}} = \frac{|1/c|}{|1/c|} = 1.$$

This yields that for any p -adic Möbius map $g \notin \mathcal{U}$, and any $s \in \mathcal{U}$, $\rho_0(g, s) = 1$, namely $d(g, \mathcal{U}) = 1$. □

Let G be a subgroup of $\text{PSL}(2, \mathbb{C}_p)$. We say that G a discrete subgroup if there exists a positive number ε such that for any non-unit element g with $\|g - I\| > \varepsilon$.

Theorem 7.3. *If G is a subgroup of $\text{PSL}(2, \mathbb{C}_p)$ and $G \cap \mathcal{U} = I$, then G is a discrete subgroup.*

Proof. By Theorem 6.5 and Theorem 7.2, we have $d(f, \mathcal{U}) \leq \rho_0(f, I) \leq \|f - I\|$. Since $G \cap \mathcal{U} = I$, for any nonunit element f , $\|f - I\| \geq 1$. □

Corollary 7.4. *If a subgroup $G \subset \text{PSL}(2, \mathbb{C}_p)$ contains unit element or loxodromic element only, then G is a discrete subgroup.*

Proof. Since each loxodromic element does not belong to \mathcal{U} , by Theorem 7.3, we can get the conclusion directly. □

REFERENCES

- [1] L.V. Ahlfors, *Möbius transformations in several dimensions*, Ordway Professorship Lectures in Mathematics, Univ. of Minnesota, Minneapolis, 1981.
- [2] M.R. Alain, *A course in p -adic analysis*, Graduate Texts in Mathematics 198, Springer-Verlag New York, 2000.
- [3] V. S. Anashin, Uniformly distributed sequences of p -adic integers, (Russian) *Diskret. Mat.* **14** (2002), no. 4, 3-64; translation in *Discrete Math. Appl.* **12** (2002), no. 6, 527-590.
- [4] V. S. Anashin and A. Khrennikov, *Applied algebraic dynamics*, de Gruyter Expositions in Mathematics, 49. Walter de Gruyter Co., Berlin, 2009.

- [5] E. Artin, *Algebraic Numbers and Algebraic Functions*, Gordon and Breach Sci. Publishers, New York, (1967).
- [6] J. V. Armitage and J. R. Parker, Jorgensens Inequality for Non-Archimedean Metric Spaces, *Geometry and Dynamics of Groups and Spaces* **265** (2008), 97-111.
- [7] M. Baker and R. Rumely, *Potential theory and dynamics on the Berkovich projective line*. Mathematical Surveys and Monographs 159, American Mathematical Society, Providence, RI, (2010).
- [8] A. F. Beardon, *The geometry of discrete groups*, Grad. Texts in Math., 91, Springer-Verlag, Berlin, 1983.
- [9] A. F. Beardon, Continued fractions, discrete groups and complex dynamics, *Comput. Methods Funct. Theory* **1** (2001), 535-594.
- [10] A. F. Beardon, The Pointwise convergence of Möbius maps, *Michigan Math. J.* **52**(3) (2004), 483-489.
- [11] A. Chambert-Loir and A. Ducros, Formes différentielles reelles et courants sur les espaces de Berkovich, arXiv:1204.6277.
- [12] A. F. Beardon and I. Short, Norms of Möbius maps, *Bull. London Math. Soc.*, **42** (2010) 499-505.
- [13] R. L. Benedetto, Hyperbolic maps in p -adic dynamics, *Ergodic Theory and Dynamical Systems*, **21** (2001), 1-11.
- [14] R. L. Benedetto, Reduction, dynamics, and Julia sets of rational functions, *J. Number Theory*, **86** (2001), 175-195.
- [15] A. Chambert-Loir *Diophantine Geometry and Analytic Spaces*, in *Tropical and Non-Archimedean Geometry* (Bellairs Workshop in Number Theory, 6-13 Mai 2011), Contemporary Mathematics 605, 2013, p. 161-179. arXiv:1210.0304, PDF.
- [16] A. Ducros, Espaces analytiques p -adiques au sens de Berkovich, talk number 958 of the Bourbaki seminar (march 2006), in *Astrisque* 311 (2007) 137-176.
- [17] P. Erdős and G. Piranian, Sequences of linear fractional transformations, *Michigan Math. J.*, **6** (1959), 205-209.
- [18] C. Favre and M. Jonsson, *The valuative tree*, **1853 Lec. Note. Math., Springer-Verlag, Berlin, 2004.**
- [19] A. H. Fan, M. T. Li, J. Y. Yao and D. Zhou, Strict ergodicity of affine p -adic dynamical systems on \mathbb{Z}_p , *Adv. Math.*, **214**(2) (2007), 666-700.
- [20] C. Favre, J. Kiwi and E. Trucco, A non-Archimedean Montels theorem, *Compositio Mathematica*, **148** (03) (2012), 966-990.
- [21] A. H. Fan, L. M. Liao, Y. F. Wang and D. Zhou, p -adic repellers in \mathbb{Q}_p are subshifts of finite type, *C. R. Math. Acad. Sci. Paris*, **344**(4) (2007), 219-224.
- [22] B. Dragovich, p -Adic and Adelic Quantum Mechanics, *Proc. V. A. Steklov Inst. Math.*, **245** (2004) 72-85.
- [23] B. Dragovich, A. Khrennikov and D. Mihajlović, Linear Fractional p -Adic and Adelic Dynamical Systems; math-ph/0612058.
- [24] F. W. Gehring and G. J. Martin, *The matrix and chordal norms of Möbius transformations*, Complex analysis (ed. J. Hersch and A. Huber; Birkhäuser, Basel, 1988) 51-59.
- [25] F. W. Gehring and G. J. Martin, Inequalities for Möbius transformations and discrete groups, *J. reineangew. Math.*, **418** (1991) 31-76.
- [26] F. Kato, Non-archimedean orbifolds covered by Mumford curves, *J. Algebraic Geom.*, **14** (2005), 1-34.
- [27] P. J. Nicholls, *The ergodic theory of discrete groups*, London Math. Soc. Lecture Note Ser., 143, Cambridge Univ. Press, Cambridge, 1989.
- [28] G. Piranian and W. J. Thron, Convergence properties of sequences of linear fractional transformations, *Michigan Math. J.* **4** (1957), 129-135.
- [29] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*, Grad. Texts in Math., 149, Springer-Verlag, New York, 1994.

- [30] W. Y. Qiu and J. H. Yang, On limit sets of discontinuous subgroups of the Berkovich space, preprint.
- [31] W. Y. Qiu, Y. F. Wang, J. H. Yang, and Y. C. Yin, On metric properties of limit sets of contractive analytic non-Archimedean dynamical systems, *J. Math. Anal. Appl.* **414** (2014) 386-401.
- [32] W. Y. Qiu, J. H. Yang, and Y. C. Yin, The Discrete Subgroups and Jørgensens Inequality for $SL(m, \mathbb{C}_p)$, *Acta Math. Sin., English Series Mar.*, **29**(3) 2013, 417C428.
- [33] J. H. Silverman, *The Arithmetic of Dynamical Systems*, Springer-Verlag-Graduate Texts in Mathematics 241, 2007.
- [34] Y. F. Wang and J. H. Yang, The pointwise convergence of the p -adic Möbius maps, *Sci-China Math.*, **57**(1) 2014, 1-8.

YUEFEI WANG, INSTITUTE OF MATHEMATICS, AMSS, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, P.R.CHINA

E-mail address: wangyf@math.ac.cn

JINGHUA YANG, SHANGHAI UNIV., SHANGHAI, 200444, P.R.CHINA

E-mail address: davidyoung@amss.ac.cn