

# On the structure of the algebra generated by the non-commutative operator graph demonstrating superactivation for a zero-error capacity

G.G. Amosov<sup>\*1</sup> and I.Yu. Zhdanovskiy<sup>†2, 3</sup>

<sup>1</sup>Steklov Mathematical Institute of Russian Academy of Sciences

<sup>2</sup>Moscow Institute of Physics and Technology

<sup>3</sup>National Research University High School of Economics, Laboratory of Algebraic Geometry

February 4, 2021

## Abstract

Recently M.E. Shirokov [1] introduced the non-commutative operator graph depending on the complex parameter  $\theta$  to construct channels with positive quantum zero-error capacity having vanishing n-shot capacity. We study the algebraic structure of this graph. The relations for the algebra generated by the graph are derived. In the limiting case  $\theta = \pm 1$  the graph becomes abelian and degenerates into the direct sum of four one-dimensional irreducible representations of the Klein group.

## 1 Introduction

The superactivation of a quantum channel capacity was discovered in [2]. It appeared that the quantum capacity for a tensor product of two quantum channels may be positive while the quantum capacity of each of these channels is zero. In [3, 4] it was shown that a value of the quantum capacity is closely related to the so-called non-commutative graph of a quantum channel. The same phenomenon for the zero-error classical capacities was established in [5]. In [6, 7] some techniques for studying superactivation by means of non-commutative graphs was introduced. It allows to give low-dimensional examples of superactivation for a quantum capacity. In our paper we shall analyse the algebra generated by the non-commutative graph introduced in [1].

---

\*gramos@mi.ras.ru

†ijdanov@mail.ru

## 2 Preliminaries

Denote  $B(H)$  the algebra of all bounded operators and  $\mathfrak{S}(H)$  the convex set of quantum states (positive unit trace operators) in a finite-dimensional Hilbert state  $H$ . Let  $\Phi : B(H_A) \rightarrow B(H_B)$  be a quantum channel, i.e. a completely positive trace-preserving linear map transmitting  $\mathfrak{S}(H_A)$  to the subset of  $\mathfrak{S}(H_B)$  in the Hilbert spaces  $H_A$  and  $H_B$  [8]. Then the dual map  $\Phi^* : B(H_B) \rightarrow B(H_A)$  is defined by the formula

$$\text{Tr}(\rho\Phi^*(x)) = \text{Tr}(\Phi(\rho)x), \quad \rho \in \mathfrak{S}(H_A), \quad x \in B(H_B).$$

The Stinespring theorem results in the representation of a channel  $\Phi$  in the form

$$\Phi(\rho) = \text{Tr}_{H_E} V\rho V^*, \quad \rho \in \mathfrak{S}(H_A), \quad (1)$$

where  $H_E$  is a Hilbert space of the environment and  $V : H_A \rightarrow H_B \otimes H_E$  is an isometry. The representation (1) allows to define a complementary quantum channel  $\hat{\Phi} : \mathfrak{S}(H_A) \rightarrow \mathfrak{S}(H_E)$  by the formula [8, 9]

$$\hat{\Phi}(\rho) = \text{Tr}_{H_B} V\rho V^*, \quad \rho \in \mathfrak{S}(H_A). \quad (2)$$

Taking into account (1) it is possible to derive the Kraus representation

$$\Phi(\rho) = \sum_{k=1}^{\dim H_E} V_k \rho V_k^*, \quad \rho \in \mathfrak{S}(H_A), \quad (3)$$

where  $\{V_k\}$  is a set of linear operators  $V_k : H_A \rightarrow H_B$ ,  $\sum_j V_k^* V_k = I_{H_A}$ . Using (3) the complementary channel (2) can be represented as follows

$$\hat{\Phi}(\rho) = \sum_{j,k=1}^{\dim H_E} \text{Tr}[V_j \rho V_k^*] |j\rangle\langle k|. \quad (4)$$

The non-commutative graph [3, 5]  $\mathcal{G}(\Phi)$  of a quantum channel  $\Phi$  is a closed linear span of the Kraus operators  $\{V_j^* V_k\}_{j,k=1}^{\dim H_E}$ . It follows from (4) that  $\mathcal{G}(\phi)$  is equal to the subspace  $\hat{\Phi}^*(B(H_E))$ . The operator space  $S$  is a non-commutative graph for some quantum channel iff  $x \in S$  implies  $x^* \in S$  and the identity operator  $I \in S$  [3, 6]. In [1] the following operator graph was introduced

$$\begin{pmatrix} a & b & c\theta & d \\ b & a & d & c/\theta \\ c/\theta & d & a & b \\ d & c\theta & b & a \end{pmatrix} \quad (5)$$

for  $\theta \in \mathbb{C}$ ,  $|\theta| = 1$ . The operator graph (5) is associated with pseudo-diagonal channels [10]

$$\Phi(\rho) = \sum_{j,k} c_{j,k} \langle \psi_j | \rho | \psi_k \rangle |j\rangle\langle k| \quad (6)$$

where  $\{c_{ij}\}$  is a Gram matrix of a collection of unit vectors,  $\{|\psi_i\rangle\}$  is a collection of vectors in  $H$  such that  $\sum_i |\psi_i\rangle\langle\psi_i| = I$ . Here we recover the structure of the algebra  $\mathcal{M}_\theta$  generated by the graph (5) for any complex parameter  $\theta \in \mathbb{C}$ ,  $\theta \neq 0$ .

### 3 The structure of the algebra $\mathcal{M}_\theta$ generated by the graph.

It is straightforward to check that the graph (5) is a linear envelope of the identity  $I$  and the following three matrices:

$$X = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & 0 & \theta & 0 \\ 0 & 0 & 0 & 1/\theta \\ 1/\theta & 0 & 0 & 0 \\ 0 & \theta & 0 & 0 \end{pmatrix}, Z = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad (7)$$

The matrices (7) satisfy the relations

$$X^2 = Y^2 = Z^2 = I, \quad XZ = ZX, \quad YZ = ZY. \quad (8)$$

Let us consider the group  $G$  generated by formal variables  $x, y, z$  satisfying the relations:  $x^2 = y^2 = z^2 = 1, xz = zx, yz = zy$ . Note that adding to the last relations  $xy = yx = z$  we obtain the Klein group. Consider the subalgebra  $\mathcal{M}_\theta \subset Mat_4(\mathbb{C})$  generated by the matrices  $X, Y, Z$ . Thus, we have a well-defined morphism of algebras:  $\phi_\theta : \mathbb{C}G \rightarrow \mathcal{M}_\theta$  defined by the rule:  $x \mapsto X, y \mapsto Y, z \mapsto Z$ . Also, the morphism  $\phi_\theta$  defines a 4-dimensional representation of the group  $G$ .

**Theorem 1.** *We have the following statements:*

- *If  $\theta \neq \pm 1$  then the algebra  $\mathcal{M}_\theta$  is a direct sum of two matrix algebras of size 2. In this case  $\dim_{\mathbb{C}} \mathcal{M}_\theta = 8$ .*
- *If  $\theta = \pm 1$  then the algebra  $\mathcal{M}_\theta$  is isomorphic to the group algebra of the Klein group. In this case  $\dim_{\mathbb{C}} \mathcal{M}_\theta = 4$ .*

*Proof.*

Consider a normal subgroup  $P \triangleleft G$  of index 2 generated by the elements  $g = xy$  and  $z$ . Note that we have the following relation:  $xgx = g^{-1}$ . Using the relations of the group  $G$ , we obtain that the group  $P$  is abelian. Hence, all irreducible representations of  $P$  are 1-dimensional. Any 1-dimensional representation is said to be a character of  $P$ . Thus, a dimension of irreducible representations of  $G$  is less or equal to 2. Let us describe characters of  $P$ . A character  $\chi$  of  $P$  is a morphism:  $\chi : P \rightarrow \mathbb{C} \setminus \{0\}$ ,  $\chi(ab) = \chi(a)\chi(b)$ ,  $a, b \in P$ . The character  $\chi$  is determined by two numbers  $\chi(g) \in \mathbb{C} \setminus \{0\}$  and  $\chi(z)$ ,  $\chi(z)^2 = 1$ . The last condition implies  $\chi(z) = \pm 1$ .

One can describe the standard construction of a  $G$ -representation  $V_\chi$  induced by the character  $\chi$  of  $P$  as follows. Consider a vector  $v$  such that  $av = \chi(a)v$ ,  $a \in P$ . Also, we can consider a "formal" vector  $x \cdot v$ . Thus, the vector space generated by  $v$  and  $x \cdot v$  is a space of  $V_\chi$ . Elements of  $P$  act on  $V_\chi$  by the rule:  $av = \chi(a)v$ . Using the relations  $x^2 = 1$  and  $xax = a^{-1}$ , we get  $a(x \cdot v) = x(xax \cdot v) = xa^{-1} \cdot v = \chi(a^{-1})x \cdot v = \chi^{-1}(a)x \cdot v$ . It means that  $V_\chi$  as representation of  $P$  is a direct sum of 1-dimensional representations corresponding to characters  $\chi$  and  $\chi^{-1}$ . The element  $x$  acts by the formula:  $xv = x \cdot v$  and  $xx \cdot v = x^2v = v$ . Also, one can show that any 2-dimensional irreducible representation of  $G$  is induced by a character of  $P$ . Consider the

representation  $\phi_\theta$  of the group algebra  $\mathbb{C}G$  for  $\theta \neq \pm 1$ . In this case there are two irreducible submodules  $V_\chi$  and  $V_{-\chi}$  of representation  $\phi_\theta$ , where characters  $\chi$  and  $-\chi$  are defined by rule:  $\chi(g) = \theta$  and  $-\chi(g) = -\theta$ . Using the standard arguments, we get that the  $G$ -representation  $\phi_\theta$  is a direct sum  $V_\chi \oplus V_{-\chi}$ . It means that if  $\theta \neq \pm 1$  then algebra  $\mathcal{M}_\theta$  is a direct sum of the matrix algebras of size 2. Thus, we get the first statement. In the case  $\theta = \pm 1$ , one can check that the representation  $\phi$  is a sum of four 1-dimensional representations.

□

We can see that there is a gap in dimension of the algebras  $\mathcal{M}_\theta$  in the case  $\theta = \pm 1$ . We can change this situation as follows. We will construct a family of algebras  $\mathcal{A}_\theta, \theta \in \mathbb{C} \setminus \{0\}$  and a morphism:  $\psi : \mathcal{A}_\theta \rightarrow \mathcal{M}_\theta$  such that  $\dim_{\mathbb{C}} \mathcal{A}_\theta = 8$  and  $\psi : \mathcal{A}_\theta \cong \mathcal{M}_\theta$  for  $\theta \neq \pm 1$ . In the case  $\theta = \pm 1$  morphism  $\psi : \mathcal{A}_\theta \rightarrow \mathcal{M}_\theta$  is a surjective and  $\text{Ker} \psi$  is a 4-dimensional nilpotent ideal of  $\mathcal{A}_\theta$ . Roughly speaking, we construct family of algebras  $\mathcal{A}_\theta$  isomorphic to  $\mathcal{M}_\theta$  for general  $\theta$ . In the case  $\theta = \pm 1$  algebras  $\mathcal{A}_\theta$  "degenerates" to  $\mathcal{M}_\theta$ .

Consider the algebra  $\mathcal{A}_\theta$  generated by "formal" variables  $x, y, z$  and the relations:  $x^2 = y^2 = z^2 = 1, xz = zx, yz = zy, xy + yx = (\theta + \theta^{-1})z$ . We can rewrite the last relation in the following manner:  $g + g^{-1} = (\theta + \theta^{-1})z$ , where  $g = xy$ . One can check that this relation is true for the matrices  $X, Y, Z$ . Thus, we have a surjective morphism:  $\psi : \mathcal{A}_\theta \rightarrow \mathcal{M}_\theta$  defined by the rule:  $\psi : x \mapsto X, y \mapsto Y, z \mapsto Z$ . We can prove the following theorem:

**Theorem 2.**

- If  $\theta \neq \pm 1$  then  $\psi$  is isomorphism.
- If  $\theta = \pm 1$  then the algebra  $\mathcal{A}_\theta$  has the 4-dimensional two-sided nilpotent ideal  $J$  such that  $\psi : \mathcal{A}_\theta/J \cong \mathcal{M}_\theta$ .

Proof.

Let us show that a dimension of the algebra  $\mathcal{A}_\theta$  is equal to 8. Actually, one can show that if  $\theta \neq \pm i$  the algebra  $\mathcal{A}_\theta$  has the following basis:  $1, g, g^2, g^3, x, xg, xg^2, xg^3$ , where  $g = xy$ . In the case  $\theta = \pm i$  one can show that the algebra  $\mathcal{A}_\theta$  has the basis:  $1, g, x, z, xg, xz, gz, xgz$ . Thus, in the case  $\theta \neq \pm 1$ ,  $\psi$  is bijective, hence,  $\psi$  is isomorphism.

Consider the case  $\theta = \pm 1$ . In this case we have the relation:  $g + g^{-1} = \pm 2z$ . Thus,  $(g + g^{-1})^2 = 4z^2 = 4$ . And, hence, we obtain the following relation:

$$(g^2 - 1)^2 = 0. \tag{9}$$

Consider the ideal  $J$  of the algebra  $A$  generated by  $g^2 - 1$ . One can see that  $J$  has the following basis  $g^2 - 1, x(g^2 - 1), g(g^2 - 1), xg(g^2 - 1)$ . Also, consider  $J^2 = \langle t_1 t_2, t_i \in J \rangle$ . Using the relation (9), we get  $J^2 = 0$ . The ideal satisfying this property is said to be a radical. Therefore, the algebra  $\mathcal{A}_\theta$  for  $\theta = \pm 1$  has a 4-dimensional radical. One can check that  $\psi(J) = 0$ .

□

## 4 Conclusion

We have found the general features of the algebra generated by the non-commutative operator graph playing an important role in quantum information theory. We have presented only a

sketch of the theory. We are planning to continue the study in the future to discover the algebraic nature of the quantum superactivation.

## Acknowledgments

The authors are grateful to A.S. Holevo for kind attention to the work and many useful remarks. Theorem 1 was proved by G.G. Amosov, Theorem 2 was proved by I.Yu. Zdanovsky. The work of G.G. Amosov is supported by Russian Science Foundation under grant No 14-21-00162 and performed in Steklov Mathematical Institute of Russian Academy of Sciences. The work of I.Yu. Zhanovskiy is supported by RFBR, research projects 13-01-00234 and 14-01-00416, and was prepared within the framework of a subsidy granted to the HSE by the Government of the Russian Federation for the implementation of the Global Competitiveness Program.

## References

- [1] M.E. Shirokov, Quantum Information Processing, **14:8** 3057-3074 (2015)
- [2] G. Smith, J. Yard, Science, **321**, 1812 (2008); arXiv:0807.4935
- [3] R. Duan, arXiv: 0906.2527 (2009).
- [4] T.S. Cubitt, J. Chen, A.W. Harrow, arXiv:0906.2547 (2009).
- [5] R. Duan, S. Severini, A. Winter, IEEE Trans. Inf. Theory, **59 :2**, 1164-1174 (2013); arXiv:1002.2514.
- [6] M.E. Shirokov, T.V. Shulman, Comm. Math. Phys., **335:3**, 1159-1179 (2015); arXiv:1309.2610.
- [7] M.E.Shirokov, T.V.Shulman, Problems Inform. Transmission. **50:3**, 232-246 (2014); arXiv:1312.3586.
- [8] A.S.Holevo, Quantum systems, channels, information. A mathematical introduction, Berlin, DeGruyter, 2012.
- [9] A.S. Holevo, Probability Theory and Applications, **51:1**, 134-143 (2006); arXiv:quant-ph/0509101.
- [10] T.S. Cubitt, M.B. Ruskai, G. Smith, J. Math. Phys., **49:2** (2008) 102104; arXiv:0802.1360.