

# SCHREIER THEORY OF TRACK CATEGORIES

MARIAM PIRASHVILI

## 1. INTRODUCTION

This paper is a continuation of our study of non-abelian Baues-Wirsching cohomologies. In our previous paper [3], we defined second non-abelian cohomology  $H^2(\mathcal{C}, D)$  of a small category  $\mathcal{C}$  with coefficients in a so-called centralised natural system  $D$ . We proved that  $H^2(\mathcal{C}, D)$  classifies linear extensions of  $\mathcal{C}$  by  $D$ , generalising the corresponding result for abelian natural systems proved in [2].

For an abelian natural system  $D$ , the third cohomology classifies certain abelian track categories [1]. A track category is a 2-category where all 2-morphisms are isomorphisms. A track category is called abelian if for every 1-morphism  $f$ , the group  $Aut(f)$  is abelian.

In a similar fashion to the above, we want to generalise this result for non-abelian track categories. In this paper we solve this problem for the following important case: Given categories  $\mathcal{K}$  and  $\mathcal{C}$  and a functor  $\pi : \mathcal{K} \rightarrow \mathcal{C}$ , which is identity on objects and surjective on morphisms, and  $G$ , a centralised natural system of groups on  $\mathcal{K}$ , we describe the equivalence classes of all track categories  $\mathcal{T}$  for which  $\mathcal{K}$  is the underlying category and  $\mathcal{C}$  is the homotopy category and  $G_f = Aut(f)$ .

## 2. PRELIMINARIES

**2.1. Track categories.** For a small groupoid  $\mathbf{G}$ , the set of its objects will be denoted by  $\mathbf{G}_0$  and the set of morphisms by  $\mathbf{G}_1$ . We have the source and target maps  $\mathbf{G}_1 \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} \mathbf{G}_0$ .

A groupoid is called *abelian* if all automorphism groups are abelian.

**Definition 1.** A category enriched in groupoids  $\mathcal{T}$ , also termed track category, is a 2-category whose all 2-cells are invertible. It is thus a class of objects  $Ob(\mathcal{T})$ , a collection of groupoids  $\mathcal{T}(A, B)$  for  $A, B \in Ob(\mathcal{T})$  called hom-groupoids of  $\mathcal{T}$ , identities  $1_A \in \mathcal{T}(A, A)_0$  and composition functors  $\mathcal{T}(B, C) \times \mathcal{T}(A, B) \rightarrow \mathcal{T}(A, C)$  satisfying the usual equations of associativity and identity morphisms.

Thus  $\mathcal{T}(A, B)_0$  denotes the set of objects of the groupoid  $\mathcal{T}(A, B)$ , while  $\mathcal{T}(A, B)_1$  denotes the set morphisms of the groupoid  $\mathcal{T}(A, B)$ . Elements of  $\mathcal{T}(A, B)_0$  we will call morphisms of  $\mathcal{T}$ , while elements of  $\mathcal{T}(A, B)_1$  are called tracks of  $\mathcal{T}$ . The underlying category of  $\mathcal{T}$  is denoted by  $\mathcal{K}$ .

For  $f, g \in \mathcal{T}(A, B)_1$  we shall write  $f \simeq g$  (and say  $f$  is *homotopic* to  $g$ ) if there exists a morphism  $\alpha$  of  $\mathcal{T}$  from  $f$  to  $g$ . Occasionally this will also be denoted by  $\alpha : f \simeq g$  or  $\alpha : f \Rightarrow g$ . Sometimes  $\alpha$  is called a *homotopy* or a *track* from  $f$  to  $g$ . Homotopy is a natural equivalence relation on morphisms of  $\mathcal{K}$  and determines the homotopy category  $\mathcal{C} = \mathcal{K} / \simeq$ . Objects of  $\mathcal{C}$  are once again the objects in  $Ob(\mathcal{T})$ , while the morphisms of  $\mathcal{C}$  are the homotopy classes of morphisms in  $\mathcal{K}$ . We denote the canonical functor by  $\pi$ . So  $\pi : \mathcal{K} \rightarrow \mathcal{C}$ .

A morphism  $g : B \rightarrow C$  in  $\mathcal{T}$  induces the functors

$$\begin{aligned} g_* : \mathcal{T}(A, B) &\rightarrow \mathcal{T}(A, C), & f &\mapsto gf, & \alpha &\mapsto g_*\alpha, \\ g^* : \mathcal{T}(C, D) &\rightarrow \mathcal{T}(B, D), & h &\mapsto hg, & \beta &\mapsto g^*\beta. \end{aligned}$$

These functors are restrictions of the composition functors. It follows from the definition that the following relations hold:

$$\text{TR 1} \quad (\alpha + \beta) + \gamma = \alpha + (\beta + \gamma),$$

- TR 2  $\alpha + 0 = \alpha = 0 + \alpha,$
- TR 3  $f^*(\alpha + \beta) = f^*(\alpha) + f^*(\beta),$
- TR 4  $g_*(\alpha + \beta) = g_*(\alpha) + g_*(\beta),$
- TR 5  $f^*(0) = 0 = g_*(0),$
- TR 6  $(ff_1)^* = f_1^*f^*, 1^* = 1,$
- TR 7  $(gg_1)_* = g_*g_{1*}, 1_* = 1,$
- TR 8  $g_*f^* = f^*g_*,$
- TR 9  $g_*(\alpha) + f_1^*(\alpha_1) = f^*(\alpha_1) + g_{1*}(\alpha).$

The following diagram explains the 1-morphisms and 2-morphisms in TR 9:

$$A \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{f_1} \end{array} B \begin{array}{c} \xrightarrow{g} \\ \xleftarrow{g_1} \end{array} C, \quad \alpha : f \Rightarrow f_1, \quad \alpha_1 : g \Rightarrow g_1.$$

The equality TR 9 holds in  $\mathcal{T}(gf, g_1f_1)$ .

**Definition 2.** A track functor, or else 2-functor,  $F : \mathcal{T} \rightarrow \mathcal{T}'$  between track categories is a groupoid enriched functor. Thus  $F$  assigns to each  $A \in \text{Ob}(\mathcal{T})$  an object  $F(A) \in \text{Ob}(\mathcal{T}')$ , to each map  $f : A \rightarrow B$  in  $\mathcal{T}$  a map  $F(f) : F(A) \rightarrow F(B)$ , and to each track  $\alpha : f \Rightarrow g$  for  $f, g : A \rightarrow B$  a track  $F(\alpha) : F(f) \Rightarrow F(g)$  in a functorial way, i.e. so that one gets functors  $F_{A,B} : \mathcal{T}(A, B) \rightarrow \mathcal{T}'(F(A), F(B))$ . These assignments are compatible with identities and composition, or equivalently induce a functor  $\mathcal{T}_1 \rightarrow \mathcal{T}'_1$ , that is  $F(1_A) = 1_{F(A)}$  for  $A \in \text{Ob}(\mathcal{T})$ ,  $F(fg) = F(f)F(g)$ , and  $F(\phi \times \psi) = F(\phi) \times (F\psi)$  for any  $\phi : f \Rightarrow f'$ ,  $\psi : g \Rightarrow g'$ ,  $f, f' : B \rightarrow C$ ,  $g, g' : A \rightarrow B$  in  $\mathcal{T}$ .

A track functor  $F : \mathcal{T} \rightarrow \mathcal{T}'$  is called a *weak equivalence* between track categories if the functors  $\mathcal{T}(A, B) \rightarrow \mathcal{T}'(F(A), F(B))$  are equivalences of groupoids for all objects  $A, B$  of  $\mathcal{T}$  and each object  $A'$  of  $\mathcal{T}'$  is homotopy equivalent to some object of the form  $F(A)$ . Such a weak equivalence induces a functor  $F : \mathcal{C} \rightarrow \mathcal{C}'$  between homotopy categories which is an equivalence of categories.

**2.2. Natural systems.** For a category  $\mathbf{I}$ , one denotes by  $\mathbf{FI}$  the *category of factorizations of  $\mathbf{I}$* . Let us recall that objects of the category  $\mathbf{FI}$  are morphisms  $f : i \rightarrow j$  of  $\mathbf{I}$ . A morphism from  $f$  to  $f' : i' \rightarrow j'$  in  $\mathbf{FI}$  is a pair  $(g, h)$ , where  $g : i' \rightarrow i$  and  $h : j \rightarrow j'$  are morphisms in  $\mathbf{I}$  such that

$$f' = h \circ f \circ g.$$

In other words, the following diagram

$$\begin{array}{ccc} j & \xrightarrow{h} & j' \\ f \uparrow & & \uparrow f' \\ i & \xleftarrow{g} & k \end{array}$$

commutes. Composition in  $\mathbf{FI}$  is defined by

$$(g', h')(g, h) = (gg', h'h).$$

It is clear that

$$(g, h) = (g, \text{Id}_{j'}) (\text{Id}_i, h) = (\text{Id}_{i'}, h) (g, \text{Id}_j).$$

Let  $\mathbf{I}$  be a category. A *natural system* on  $\mathbf{I}$  with values in a category  $\mathbf{C}$  is a functor  $D : \mathbf{FI} \rightarrow \mathbf{C}$ . We denote the value of  $D$  on  $f : i \rightarrow j$  by  $D_f$  or  $D(f)$ . If  $f$  is the identity

$\text{Id}_i : i \rightarrow i$  we write  $D_i$  instead of  $D_{\text{Id}_i}$ . We also write  $g^*$  and  $h_*$  instead of  $D(g, \text{Id})$  and  $D(\text{Id}, h)$ . Then

$$D(g, h) = g^*h_* = h_*g^* : D_f \rightarrow D_{hfg}.$$

If  $\mathbf{C}$  is the category of sets; respectively groups, or abelian groups; we say that  $D$  is a natural system of sets; respectively groups, or abelian groups.

Let  $\mathbf{I}$  be a small category. A *centralised natural system of groups* is a natural system  $D$  of groups on  $\mathbf{I}$  such that for any arrows  $i \xrightarrow{f} j \xrightarrow{g} k$  and elements  $x \in D_f, y \in D_g$  one has the equality

$$g_*(x) + f^*(y) = f^*(y) + g_*(x)$$

in the group  $D_{gf}$ . By putting  $i = j = k$  and  $g = f = \text{id}_i$ , it follows that for any object  $i$  the group  $D_i$  is an abelian group.

**2.3. Natural system  $\text{Aut}^{\mathcal{T}}$ .** Let  $\mathcal{T}$  be a track category. Recall that the underlying category  $\mathcal{K}$  has the same objects as  $\mathcal{T}$ , however the set of morphisms  $\text{Hom}_{\mathcal{K}}(A, B)$  is the set of objects of the category  $\mathcal{T}(A, B)$ . For any morphism  $f : A \rightarrow B$  of  $\mathcal{K}$  we let  $\text{Aut}_f^{\mathcal{T}}$  be the collection of all automorphisms of  $f$  in the category  $\mathcal{T}(A, B)$ . Thus, this is the collection of all tracks  $\alpha : f \Rightarrow f$ . It follows from TR 1 and TR 2 that  $D_f$  is a group. Moreover, for any morphism  $g : B \rightarrow C$ , we have maps  $g_* : \text{Aut}_f^{\mathcal{T}} \rightarrow \text{Aut}_{gf}^{\mathcal{T}}$  and  $f^* : \text{Aut}_g^{\mathcal{T}} \rightarrow \text{Aut}_{gf}^{\mathcal{T}}$ , which are group homomorphisms thanks to TR 3 – TR 5. Moreover, in this way one obtains a natural system  $\text{Aut}^{\mathcal{T}}$  of groups on  $\mathcal{K}$ . This follows from the identities TR 6 – TR 8. We claim that  $\text{Aut}^{\mathcal{T}}$  is centralised. To show this it suffices to put  $f = f_1$  and  $g = g_1$  in TR 9 to get:

$$g_*(\alpha) + f^*(\alpha_1) = f^*(\alpha_1) + g_*(\alpha).$$

### 3. COHOMOLOGY OF PRE-TRACK CATEGORIES

**3.1. Pre-track categories.** A pre-track category  $(\pi : \mathcal{K} \rightarrow \mathcal{C}, G)$ , or  $(\pi, G)$  for short, is the following data:

- (1)  $\mathcal{K}$  and  $\mathcal{C}$  are categories and  $\pi$  is a functor which is identity on objects and surjective on morphisms,
- (2)  $G$  is a centralised natural system of groups on  $\mathcal{K}$ .

**Example 3.** *The main example for this is the following. Let  $\mathcal{T}$  be a track category and let  $\mathcal{C}$  be its homotopy category. Then  $(\pi : \mathcal{K} \rightarrow \mathcal{C}, \text{Aut}^{\mathcal{T}})$  is a pre-track category.*

So any track category gives rise to a pre-track category. The question is whether we can say something about the converse.

**Definition 4.** *Let  $(\pi : \mathcal{K} \rightarrow \mathcal{C}, G)$  be a pre-track category. A  $(\pi, G)$ -track category is a pair  $(\mathcal{T}, \sigma)$  where*

- (1)  $\mathcal{T} = (TK \rightrightarrows \mathcal{K})$  is a track category with underlying category  $\mathcal{K}$  satisfying the property: for all  $f, g \in \mathcal{K}(A, B)$  one has  $\pi(f) = \pi(g)$  iff  $\mathcal{T}(f, g) \neq \emptyset$ .
- (2)  $\sigma$  is an isomorphism of natural systems on  $\mathcal{K}$

$$\sigma : \text{Aut}^{\mathcal{T}} \rightarrow G.$$

Two  $(\pi, G)$ -track categories  $(\mathcal{T}, \sigma)$  and  $(\mathcal{T}', \sigma')$  are equivalent if there exists a track functor  $F : \mathcal{T} \rightarrow \mathcal{T}'$  which is identity on objects and the following diagram commutes:

$$\begin{array}{ccc} \text{Aut}^{\mathcal{T}} & \xrightarrow{F} & \text{Aut}^{\mathcal{T}'} \\ & \searrow \sigma & \swarrow \sigma' \\ & & G \end{array}$$

One easily sees that if such an  $F$  exists, then it is an isomorphism of 2-categories. We let  $\text{Tracks}(\pi, G)$  be the collection of equivalence classes of  $(\pi, G)$ -track extensions.

**3.2. Second cohomology of pre-track categories.** Given a pre-track category  $(\pi : \mathcal{K} \rightarrow \mathcal{C}, G)$  we can also define the relative cohomology  $H^2(\pi, G) := Z/\sim$  where  $Z$  is defined as follows.

**Definition 5.**  $Z$  is a collection of triples all  $(\xi, \chi, \phi)$  such that

- (1)  $\xi$  is a function which assigns to all triples of morphisms  $f, g, h : i \rightarrow j$  of  $\mathcal{K}$  with  $\pi(f) = \pi(g) = \pi(h)$ , an element  $\xi(f, g, h) \in G_f$ ,
- (2)  $\chi$  is a function which assigns to each diagram  $i \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} j \begin{array}{c} \xrightarrow{x} \\ \xrightarrow{y} \end{array} k$  with  $\pi(f) = \pi(g)$  and  $\pi(x) = \pi(y)$  an element  $\chi(f, g|x, y) \in G_{xf}$ ,
- (3)  $\phi$  is a function which assigns to each pair of morphisms  $f, g$  in  $\mathcal{K}$  with  $\pi(f) = \pi(g)$  an isomorphism  $\phi_{g,f} : G_g \rightarrow G_f$ .

These functions must satisfy the following equations:

- (i) To simplify notation, first we define  $m_*\phi_{b,a} : G_{mb} \rightarrow G_{ma}$  by

$$(m_*\phi_{b,a})(t) := -\chi(a, b|m, m) + \phi_{mb,ma}(t) + \chi(a, b|m, m)$$

where  $a, b : i \rightarrow j$  and  $m : j \rightarrow k$ . Also, we define  $b^*\phi_{n,m} : G_{nb} \rightarrow G_{mb}$  by

$$(b^*\phi_{n,m})(t) := -\chi(b, b|m, n) + \phi_{nb,mb}(t) + \chi(b, b|m, n),$$

where  $b : i \rightarrow j$  and  $m, n : j \rightarrow k$ . Then we have the following equations:

- (a) For  $f, g, h : i \rightarrow j$ ,  $\phi_{g,f} \circ \phi_{h,g}(t) = \xi(f, g, h) + \phi_{h,f}(t) + \xi(f, g, h)$ .
- (b) For  $a, b : i \rightarrow j$ ,  $m, n : j \rightarrow k$ ,  $\beta \in G_b$  and  $\nu \in G_n$ ,  $m_*\phi_{b,a}(m_*\beta) = m_*(\phi_{b,a}(\beta))$ ,  $a^*\phi_{n,m}(a^*\nu) = a^*(\phi_{n,m}(\nu))$ ,
- (c) and for  $\mu \in G_m$  and  $\alpha \in G_a$ ,  $m_*\phi_{b,a}(b^*\mu) = a^*\mu$ ,  $a^*\phi_{n,m}(n_*\alpha) = m_*\alpha$ .
- (ii) For  $f, g, h, e : i \rightarrow j$  with  $\pi(f) = \pi(g) = \pi(h) = \pi(e)$ , we have to have

$$\xi(f, g, e) + \phi_{g,f}\xi(g, h, e) = \xi(f, h, e) + \xi(f, g, h).$$

- (iii) For  $x, y, z : i \rightarrow j$  with  $\pi(x) = \pi(y) = \pi(z)$  and  $a, b, c : j \rightarrow k$  with  $\pi(a) = \pi(b) = \pi(c)$ , we have to have

$$\xi(ax, by, cz) + \phi_{by,ax}(\chi(y, z|b, c)) + \chi(x, y|a, b) = \chi(x, z|a, c) + x^*\xi(a, b, c) + a_*\xi(x, y, z).$$

- (iv) For  $x, y : i \rightarrow j$  with  $\pi(x) = \pi(y)$ ,  $a, b : j \rightarrow k$  with  $\pi(a) = \pi(b)$  and  $m, n : k \rightarrow l$  with  $\pi(m) = \pi(n)$ , we have to have

$$\chi(ax, by|m, n) + m_*\chi(x, y|a, b) = \chi(x, y|ma, nb) + x^*(\chi(a, b|m, n)).$$

**Definition 6.** We say two such triples  $(\xi, \chi, \phi)$  and  $(\xi', \chi', \phi')$  are equivalent, if there is an element  $\zeta(f, g) \in G_f$  such that

- (1)  $\zeta(f, h) + \xi(f, g, h) - \phi_{g,f}\zeta(g, h) - \zeta(f, g) = \xi'(f, g, h)$ ,
- (2)  $\zeta(ax, by) + \chi(x, y|a, b) - x^*\zeta(a, b) - a_*\zeta(x, y) = \chi'(x, y|a, b)$ , and
- (3)  $\zeta(f, g) + \phi_{g,f}(t) - \zeta(f, g) = \phi'_{g,f}(t)$ .

**Theorem 7.** There is a natural bijection

$$\text{Tracks}(\pi, G) \simeq H^2(\pi, G).$$

*Proof.* Let  $(\mathcal{T}, \sigma)$  be an element of  $\text{Tracks}(\pi, G)$ . Let  $\pi(f) = \pi(g)$ , so  $\mathcal{T}(f, g) \neq \emptyset$ . Choose a track  $H_{f,g} \in \mathcal{T}(f, g)$  so that

$$H_{f,f} = 0, \quad H_{f,g} = -H_{g,f}.$$

Now take  $(f, g, h)$  in such a way that  $\pi(f) = \pi(g) = \pi(h)$ . Then there is a unique element  $\xi(f, g, h) \in G_f$  such that

$$H_{f,g} + H_{g,h} = -\xi(f, g, h) + H_{f,h}.$$

Take  $x, y \in K(i, j)$  and  $a, b \in K(j, k)$  for which  $\pi(x) = \pi(y)$  and  $\pi(a) = \pi(b)$ . Then there is a unique element  $\chi(x, y|a, b) \in G_{ax}$  such that

$$a_*H_{x,y} + y^*H_{a,b} = -\chi(x, y|a, b) + H_{ax,by}.$$

For every  $f$  and  $g$  with  $\pi(f) = \pi(g)$  we define an isomorphism  $\phi_{g,f}(t) := H_{f,g} + t - H_{f,g}$ , where  $t \in \mathcal{T}(g, g)$ .

We claim that the triple  $(\xi, \chi, \phi)$  is an element of  $Z$ .

The Equations (i) of Definition 5 are straightforward to check. First, we check composition:

$$\begin{aligned}\phi_{g,f} \circ \phi_{h,g}(t) &= \phi_{g,f}(H_{g,h} + t - H_{g,h}) = H_{f,g} + H_{g,h} + t - H_{g,h} - H_{f,g} \\ &= -\xi(f, g, h) + \phi_{h,f}(t) + \xi(f, g, h).\end{aligned}$$

Next, we have

$$\begin{aligned}m_*(\phi_{b,a}(\beta)) &= m_*(H_{a,b} + \beta - H_{a,b}) = m_*H_{a,b} + m_*\beta - m_*H_{a,b} \\ &= (m_*\phi_{b,a})(m_*\beta).\end{aligned}$$

Similarly,

$$a^*(\phi_{n,m}(\nu)) = a^*(H_{m,n} + \nu - H_{m,n}) = (a^*\phi_{n,m})(a^*\nu).$$

The next equation

$$m_*\phi_{b,a}(b^*\mu) = m_*H_{a,b} + b^*\mu - m_*H_{a,b} = a^*\mu$$

follows from *TR9*, as shown by the diagram

$$i \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{b} \end{array} j \begin{array}{c} \xrightarrow{m} \\ \xleftarrow{m} \end{array} k, \quad H_{a,b} : a \Rightarrow b, \quad \mu : m \Rightarrow m.$$

The final equation is very similar to this.

To check (ii), we use the associativity property for the expression  $H_{f,g} + H_{g,h} + H_{h,e}$  and the definition of the function  $\xi$  to obtain

$$\begin{aligned}(H_{f,g} + H_{g,h}) + H_{h,e} &= -\xi(f, g, h) + H_{f,h} + H_{h,e} = -\xi(f, g, h) - \xi(f, h, e) + H_{f,e} \\ H_{f,g} + (H_{g,h} + H_{h,e}) &= H_{f,g} - \xi(g, h, e) + H_{g,e} = H_{f,g} - \xi(g, h, e) - H_{f,g} - \xi(f, g, e) + H_{f,e}.\end{aligned}$$

So we have

$$\xi(f, h, e) + \xi(f, g, h) = \xi(f, g, e) + \phi_{g,f}(\xi(g, h, e)).$$

Let  $x, y, z : i \rightarrow j$  and  $a, b, c : j \rightarrow k$  be morphisms in  $K$  such that  $\pi(x) = \pi(y) = \pi(z)$  and  $\pi(a) = \pi(b) = \pi(c)$ . Then we have

$$\begin{aligned}&a_*(H_{x,y} + H_{y,z}) + z^*(H_{a,b} + H_{b,c}) \\ &= a_*H_{x,y} + (a_*H_{y,z} + z^*H_{a,b}) + z^*H_{b,c} \\ &= a_*H_{x,y} + (y^*H_{a,b} + b_*H_{y,z}) + z^*H_{b,c} \\ &= -\chi(x, y|a, b) + H_{ax,by} - \chi(y, z|b, c) + H_{by,cz} \\ &= -\chi(x, y|a, b) + H_{ax,by} - \chi(y, z|b, c) - H_{ax,by} + H_{ax,by} + H_{by,cz} \\ &= -\chi(x, y|a, b) + H_{ax,by} - \chi(y, z|b, c) - H_{ax,by} - \xi(ax, by, cz) + H_{ax,cz}.\end{aligned}$$

On the other hand,

$$\begin{aligned}&a_*(H_{x,y} + H_{y,z}) + z^*(H_{a,b} + H_{b,c}) \\ &= a_*(-\xi(x, y, z) + H_{x,z}) + z^*(-\xi(a, b, c) + H_{a,c}) \\ &= -a_*\xi(x, y, z) + a_*H_{x,z} - z^*\xi(a, b, c) - a_*H_{x,z} + a_*H_{x,z} + z^*H_{a,c} \\ &= -a_*\xi(x, y, z) + a_*H_{x,z} - z^*\xi(a, b, c) - a_*H_{x,z} - \chi(x, z|a, c) + H_{ax,cz} \\ &= -a_*\xi(x, y, z) - a_*\phi_{z,x}(z^*\xi(a, b, c)) - \chi(x, z|a, c) + H_{ax,cz} \\ &= -a_*\xi(x, y, z) - x^*\xi(a, b, c) - \chi(x, z|a, c) + H_{ax,cz} \quad \text{using Equation (i)}.\end{aligned}$$

We therefore obtain Equation (iii).

$$\xi(ax, by, cz) + \phi_{by,ax}(\chi(y, z|b, c)) + \chi(x, y|a, b) = \chi(x, z|a, c) + x^*\xi(a, b, c) + a_*\xi(x, y, z).$$

Let  $x, y : i \rightarrow j$ ,  $a, b : j \rightarrow k$  and  $m, n : k \rightarrow l$  be morphisms in  $K$  such that  $q(x) = q(y)$ ,  $q(a) = q(b)$  and  $q(m) = q(n)$ . By the definition of  $\chi$  we have

$$m_*H_{a,b} + b^*H_{m,n} = -\chi(a, b|m, n) + H_{ma,nb}$$

and

$$m_* a_* H_{x,y} + y^* H_{ma,nb} = -\chi(x, y|ma, nb) + H_{max,nby}.$$

Expressing  $H_{ma,nb}$  from the first equation and substituting in the second, we obtain that the expression  $-\chi(x, y|ma, nb) + H_{max,nby}$  is equal to

$$\begin{aligned} & m_* a_* H_{x,y} + y^* (\chi(a, b|m, n)) + y^* m_* H_{a,b} + y^* b^* H_{m,n} \\ &= m_* a_* H_{x,y} + y^* (\chi(a, b|m, n)) - m_* a_* H_{x,y} + m_* a_* H_{x,y} + y^* m_* H_{a,b} + y^* b^* H_{m,n} \\ &= m_* a_* H_{x,y} + y^* (\chi(a, b|m, n)) - m_* a_* H_{x,y} + m_* (a_* H_{x,y} + y^* H_{a,b}) + y^* b^* H_{m,n} \\ &= m_* a_* H_{x,y} + y^* (\chi(a, b|m, n)) - m_* a_* H_{x,y} - m_* \chi(x, y|a, b) + m_* H_{ax,by} + y^* b^* H_{m,n} \\ &= m_* a_* H_{x,y} + y^* (\chi(a, b|m, n)) - m_* a_* H_{x,y} - m_* \chi(x, y|a, b) - \chi(ax, by|m, n) + H_{max,nby} \\ &= x^* (\chi(a, b|m, n)) - m_* \chi(x, y|a, b) - \chi(ax, by|m, n) + H_{max,nby}, \text{ using Equation (i)}. \end{aligned}$$

Therefore we have

$$\chi(ax, by|m, n) + m^* \chi(x, y|a, b) = \chi(x, y|ma, nb) + x^* (\chi(a, b|m, n)).$$

If we choose another  $H'_{f,g} \in \mathcal{T}(f, g)$ , there is a unique element  $\zeta(f, g) \in G_f$  such that

$$\zeta(f, g) + H_{f,g} = H'_{f,g}.$$

We let  $\xi', \chi'$  and  $\phi'$  be the functions corresponding to  $H'$ . For all  $(f, g, h)$  with  $\pi(f) = \pi(g) = \pi(h)$  we have

$$H'_{f,g} + H'_{g,h} = -\xi'(f, g, h) + H'_{f,h}.$$

Hence,

$$\begin{aligned} \zeta(f, g) + H_{f,g} + \zeta(g, h) + H_{g,h} &= \zeta(f, g) + H_{f,g} + \zeta(g, h) - H_{f,g} + H_{f,g} + H_{g,h} \\ &= -\xi'(f, g, h) + \zeta(f, h) + H_{f,h}. \end{aligned}$$

This gives

$$\zeta(f, h) + \xi(f, g, h) - \phi_{g,f} \zeta(g, h) - \zeta(f, g) = \xi'(f, g, h).$$

For  $(x, y) \in \mathcal{K}(i, j)$  and  $(a, b) \in \mathcal{K}(j, k)$  with  $\pi(x) = \pi(y)$  and  $\pi(a) = \pi(b)$ , we have

$$a_* H'_{x,y} + y^* H'_{a,b} = -\chi'(x, y|a, b) + H'_{ax,by}.$$

So we get

$$\begin{aligned} a_* \zeta(x, y) + a_* H_{x,y} + y^* \zeta(a, b) + y^* H_{a,b} &= a_* \zeta(x, y) + a_* \phi_{y,x} (y^* \zeta(a, b)) - \chi(x, y|a, b) + H_{ax,by} \\ &= -\chi'(x, y|a, b) + \zeta(ax, by) + H_{ax,by}. \end{aligned}$$

Hence

$$\zeta(ax, by) + \chi(x, y|a, b) - x^* \zeta(a, b) - a_* \zeta(x, y) = \chi'(x, y|a, b).$$

We also have for all  $(f, g) \in \mathcal{K}$  with  $\pi(f) = \pi(g)$  and all  $t \in G_f$

$$\zeta(f, g) + \phi_{g,f}(t) - \zeta(f, g) = \phi'_{g,f}(t).$$

The inverse map is constructed in the following way. Let  $(\xi, \chi, \phi) \in H^2(\pi, G)$ .

Let  $\pi(f) = \pi(g)$ . Then we define the set of tracks  $T(f, g)$  to be the set  $G_f \times f, g$ . The track category structure is given by

$$\begin{aligned} (\alpha, f, g) + (\beta, g, h) &= (\alpha + \phi_{g,f}(\beta) - \xi(f, g, h), f, h). \\ a_*(\alpha, f, g) &= (a_* \alpha - \chi(f, g|a, a), af, ag), \\ b^*(\alpha, f, g) &= (b^* \alpha - \chi(b, b|f, g), fb, gb). \end{aligned}$$

Let us check that this indeed gives a track structure.

Now, to verify the relations  $TR1 - TR9$ : First, we check that the addition defined above is associative. We have

$$\begin{aligned} ((\alpha, f, g) + (\beta, g, h)) + (\gamma, h, e) &= (\alpha + \phi_{g,f}(\beta) - \xi(f, g, h), f, h) + (\gamma, h, e) \\ &= (\alpha + \phi_{g,f}(\beta) - \xi(f, g, h) + \phi_{h,f}(\gamma) - \xi(f, h, e), f, e) \\ &= (\alpha + \phi_{g,f}(\beta) + \phi_{g,f} \circ \phi_{h,g}(\gamma) - \xi(f, g, h) - \xi(f, h, e), f, e). \end{aligned}$$

On the other hand,

$$\begin{aligned} (\alpha, f, g) + ((\beta, g, h) + (\gamma, h, e)) &= (\alpha, f, g) + (\beta) + \phi_{h,g}(\gamma) - \xi(g, h, e), g, e) \\ &= (\alpha + \phi_{g,f}(\beta) + \phi_{g,f} \circ \phi_{h,g}(\gamma) - \phi_{g,f}(\xi(g, h, e)) - \xi(f, g, e), f, e). \end{aligned}$$

Using Equation 5 (ii), we can see that we have equality.

Next, for *TR2* we have:

$$(\alpha, f, g) + (0, g, g) = (\alpha + \phi_{g,f}(0) - \xi(f, g, g), f, g) = (\alpha, f, g) = (0, f, f) + (\alpha, f, g)$$

. We also have that

$$\begin{aligned} m^*((\alpha, f, g) + (\beta, g, h)) &= m^*(\alpha + \phi_{g,f}(\beta) - \xi(f, g, h), f, h) \\ &= (m^*(\alpha + \phi_{g,f}(\beta) - \xi(f, g, h)) - \chi(m, m|f, h), fm, hm) \\ &= (m^*\alpha + m^*\phi_{g,f}(m^*\beta) - m^*\xi(f, g, h) - \chi(m, m|f, h), fm, hm) \\ &= (m^*\alpha - \chi(m, m|f, g) + \phi_{gm, fm}(m^*\beta) + \chi(m, m|f, g) - m^*\xi(f, g, h) - \chi(m, m|f, h), fm, hm). \end{aligned}$$

According to Equation 5 (iii),  $-m^*\xi(f, g, h) - \chi(m, m|f, h) = -\chi(m, m|f, g) - \phi_{gm, fm}(\chi(m, m|g, h)) - \xi(fm, gm, hm)$ . Therefore, we have

$$\begin{aligned} &= (m^*\alpha - \chi(m, m|f, g) + \phi_{gm, fm}(m^*\beta) - \phi_{gm, fm}(\chi(m, m|g, h)) - \xi(fm, gm, hm), fm, hm) \\ &= m^*(\alpha, f, g) + m^*(\beta, g, h). \end{aligned}$$

For *TR4*, we have very similarly:

$$\begin{aligned} x_*((\alpha, f, g) + (\beta, g, h)) &= x_*(\alpha + \phi_{g,f}(\beta) - \xi(f, g, h), f, h) \\ &= (x_*(\alpha + \phi_{g,f}(\beta) - \xi(f, g, h)) - \chi(f, h|x, x), xf, xh) \\ &= (x_*\alpha + x_*\phi_{g,f}(x_*\beta) - x_*\xi(f, g, h) - \chi(f, h|x, x), xf, xh) \\ &= (x_*\alpha - \chi(f, g|x, x) + \phi_{xg, xf}(x_*\beta) + \chi(f, g|x, x) - x_*\xi(f, g, h) - \chi(f, h|x, x), xf, xh). \end{aligned}$$

According to Equation 5 (iii),  $-x_*\xi(f, g, h) - \chi(f, h|x, x) = -\chi(f, g|x, x) - \phi_{xg, xf}(\chi(g, h|x, x)) - \xi(xf, xg, xh)$ . Therefore, we have

$$\begin{aligned} &= (x_*\alpha - \chi(f, g|x, x) + \phi_{xg, xf}(x_*\beta) - \phi_{xg, xf}(\chi(g, h|x, x)) - \xi(xf, xg, xh), xf, xh) \\ &= x_*(\alpha, f, g) + x_*(\beta, g, h). \end{aligned}$$

For *TR5*, we have

$$f^*(0, g, g) = (f^*0 - \chi(f, f|g, g), gf, gf) = (0, gf, gf) = (g_*0 - \chi(f, f|g, g), gf, gf) = g_*(0, f, f).$$

Next,

$$m^*n^*(\alpha, f, g) = m^*(n^*\alpha - \chi(n, n|f, g), fn, gn) = (m^*n^*\alpha - m^*\chi(n, n|f, g) - \chi(m, m|fn, gn), fnm, gnm).$$

Using Equation 5 (iv), we get that  $-m^*\chi(n, n|f, g) - \chi(m, m|fn, gn) = -\chi(nm, nm|f, g)$ , therefore giving us

$$= (n^*m^*\alpha - \chi(nm, nm|f, g), fnm, gnm) = (nm)^*(\alpha, f, g).$$

*TR7* can be shown in a very similar manner. For *TR8*, we have

$$\begin{aligned} x_*m^*(\alpha, f, g) &= x_*(m^*\alpha - \chi(m, m|f, g), fm, gm) \\ &= (x_*m^*\alpha - x_*\chi(m, m|f, g) - \chi(fm, gm|x, x), xfm, xgm) \\ &= (x_*m^*\alpha - m^*\chi(f, g|x, x) - \chi(m, m|xf, xg), xfm, xgm), \\ &= m^*x_*(\alpha, f, g), \end{aligned}$$

due to Equation 5 (iv).

We also check *TR9*, the last equation:

$$\begin{aligned}
m_* (\alpha, a, b) + b^* (\mu, m, n) &= (m_* \alpha - \chi(a, b|m, m), ma, mb) + (b^* \mu - \chi(b, b|m, n), mb, nb) \\
&= (m_* \alpha - \chi(a, b|m, m) + \phi_{mb, ma}(b^* \mu - \chi(b, b|m, n)) - \xi(ma, mb, nb), ma, nb) \\
&= (m_* \alpha - \chi(a, b|m, m) + \chi(a, b|m, m) + m_* \phi_{b, a}(b^* \mu) - \chi(a, b|m, m) \\
&\quad - \phi_{mb, ma}(\chi(b, b|m, n)) - \xi(ma, mb, nb), ma, nb).
\end{aligned}$$

According to Equation 5 (iii), this equals to

$$\begin{aligned}
&= (m_* \alpha + m_* \phi_{b, a}(b^* \mu) - m_* \xi(a, b, b) - m_* \phi_{b, a}(b^* \xi(m, m, n)) - \chi(a, b|m, n), ma, nb) \\
&= (m_* \alpha + m_* \phi_{b, a}(b^* \mu) - \chi(a, b|m, n), ma, nb) = (m_* \alpha + a^* \mu - \chi(a, b|m, n), ma, nb).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
a^* (\mu, m, n) + n_* (\alpha, a, b) &= (a^* \mu - \chi(a, a|m, n), ma, na) + (n_* \alpha - \chi(a, b|n, n), na, nb) \\
&= (a^* \mu - \chi(a, a|m, n) + \phi_{na, ma}(n_* \alpha - \chi(a, b|n, n)) - \xi(ma, na, nb), ma, nb) \\
&= (a^* \mu + a^* \phi_{n, m}(n_* \alpha) - \chi(a, a|m, n) - \phi_{na, ma}(\chi(a, b|n, n)) - \xi(ma, na, nb), ma, nb).
\end{aligned}$$

According to Equation 5 (iii), this equals to

$$\begin{aligned}
&= (a^* \mu + a^* \phi_{n, m}(n_* \alpha) - m_* \xi(a, a, z) - m_* \phi_{b, a}(b^* \xi(m, n, n)) - \chi(a, b|m, n), ma, nb) \\
&= (a^* \mu + a^* \phi_{n, m}(n_* \alpha) - \chi(a, b|m, n), ma, nb) = (a^* \mu + m_* \alpha - \chi(a, b|m, n), ma, nb).
\end{aligned}$$

As  $G$  is a centralised natural system, we have  $a^* \mu + m_* \alpha = m_* \alpha + a^* \mu$  in  $G_{ma}$ , therefore giving equality.  $\square$

## REFERENCES

- [1] H.-J. BAUES and M. JIBLADZE. *Classification of abelian track categories*. K-Theory 25(2002) 299-311.
- [2] H.-J. BAUES and G. C. WIRSCHING. *Cohomology of small categories* J. Pure and Appl. Algebra 38(1985), 187-211.
- [3] M. PIRASHVILI. *Second cohomotopy and nonabelian cohomology*. 13(2014), 397-445.