

Wigner matrices, the moments of roots of Hermite polynomials and the semicircle law

M. Kornyik

Eötvös Loránd University

Department of Probability Theory and Statistics

Pázmány Péter sétány 1/C., H-1117, Budapest, Hungary

email:koma@cs.elte.hu

Gy. Michaletzky

Eötvös Loránd University

Department of Probability Theory and Statistics

Pázmány Péter sétány 1/C., H-1117, Budapest, Hungary

email:michaletzky@caesar.elte.hu

April 28, 2019

Abstract

In the present paper we give two alternate proofs of the well known theorem that the empirical distribution of the appropriately normalized roots of the n^{th} monic Hermite polynomial H_n converges weakly to the semicircle law, which is also the weak limit of empirical distribution of the appropriately normalized eigenvalues of a Wigner matrix. In the first proof – based on the recursion satisfied by the Hermite polynomials – we show that the moment generating function of the roots of the H_n is convergent and it satisfies a fixed point equation, which is also satisfied by $c(z^2)$, where $c(z)$ is the generating function of the Catalan numbers C_k . In the second proof we compute the leading term of the k^{th} moments (as a polynomial in n) of H_n and show that it coincides with $C_{k/2}$, the $(k/2)^{\text{th}}$ Catalan number, where k is even. We also mention the known result that the expectation of the characteristic polynomial (p_n) of a Wigner random matrix of size $n \times n$ is exactly the Hermite polynomial H_n , i.e. $Ep_n(x) = H_n(x)$, which suggest the presence of a deep connection between the Hermite polynomials and Wigner matrices.

Keywords: Random matrix, characteristic polynomial, semicircle law, moments of roots of Hermite polynomials.

MSC 2010 : 60B20,15B52,33C45, 15A18

0 Introduction

The first theorem of section 1 presents a short, direct proof of the well known theorem saying that the semicircle law describes the asymptotic distribution of the (normalized) roots of the Hermite polynomials by showing that the generating function (without computing the actual coefficients) of the sum of k^{th} power of the roots converges to $\sum_{k=0}^{\infty} C_k z^{2k} = c(z^2)$, where $c(z)$ denotes the generating function of the Catalan numbers, using a fixed point argument similar to Girko's idea in [8]. In the second theorem we explicitly compute the leading coefficient in n of the sum of k^{th} power of the roots and show that it is equal to $C_{k/2}$ by using the implicit connection between the moments and elementary symmetric polynomials of the roots of an arbitrary polynomial (also known as Newton's identities or Viéta's formulae). This results immediately implies – after proper scaling – the weak convergence of the distribution determined by the scaled roots of H_n , as $n \rightarrow \infty$, to the semicircle law, which is also the limit distribution of the empirical distribution of properly normalized eigenvalues of a Wigner-matrix.

In random matrix theory to analyse the behaviour of the eigenvalues of a random matrix one possibility is to consider the sum of the k^{th} powers of these eigenvalues. This can be done either via analysing the trace of the k^{th} power of the random matrix, or through the k^{th} moments of the roots of its characteristic polynomial. One can find many results of the former type (see [2], [11]), but the latter has not yet been thoroughly investigated ([3], [9]). Since there is an implicit connection between the moments and

elementary symmetric polynomials of the roots (Newton's identities), one can make observations of the moments of roots via examining the coefficients of the characteristic polynomial. In section 2 we show a connection between the Hermite polynomial and the characteristic polynomial of a Wigner matrix. For the sake of completeness we will reprove the result of Diaconis and Gamburd [5] and Forrester and Gamburd [6] claiming that the expectation of the characteristic polynomial of a Wigner matrix is the Hermite polynomial, i.e. if A_n denotes a Wigner matrix of size $n \times n$ (a symmetric matrix with real valued random independent elements such that $Ea_{ij} = 0$ and $Ea_{ij}^2 = 1$ for all $1 \leq i \leq j \leq n$ and $a_{ij} = a_{ji}$ for $i > j$), then

$$Ep_n(x) = E \det(xI - A_n) = H_n(x), \quad (1)$$

where $H_n(x)$ denotes the n^{th} monic Hermite polynomial with leading coefficient 1. These polynomials are known to be orthogonal with respect to the measure $d\mu(x) = \frac{1}{\sqrt{2\pi}}e^{-x^2/2}dx$. The expectation in (1) refers to that of the coefficients of the characteristic polynomial of A_n . To compute this we also present a formula expressing the particular coefficients of this polynomial in terms of the elements of the matrix A_n .

1 Sum of the k^{th} power of the roots of H_n

Let us consider the roots of the Hermite polynomials. Denote by $\xi_1^{(n)}, \dots, \xi_n^{(n)}$ the roots of H_n and denote by $\mu_n = \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j^{(n)}}$ the empirical distribution determined by the normalized roots, where $\lambda_j^{(n)} = \frac{\xi_j^{(n)}}{\sqrt{n}}$, and by $M_n(k) = \sum_{j=1}^n \left(\xi_j^{(n)}\right)^k$ the sum of the k^{th} powers.

The following theorem is known (see e.g. [7]), but we are going to present a short, direct proof of it which is based on the recursive equations satisfied by the Hermite polynomials.

Proposition 1 (See [7]) *The limit distribution of the empirical distribution of the roots of the Hermite polynomial H_n , as $n \rightarrow \infty$, is given by the semicircle distribution. That is*

$$\mu_n \xrightarrow[n \rightarrow \infty]{w} \rho_{sc}(x) dx \quad (2)$$

where ' \xrightarrow{w} ' means weak convergence and $\rho_{sc}(x) = \frac{2}{\pi} \sqrt{1-x^2} \mathbf{1}_{[-1,1]}(x)$.

PROOF: To prove the weak convergence we apply the methods of moments. First for the sake of the reader we present a direct proof of the following known lemma (see [10]) connecting the moments of the empirical distribution of the roots and the corresponding polynomial.

Lemma 1 *Let $p(x) = \sum_{j=0}^n b_j x^j$ be a monic polynomial (i.e. $b_n = 1$) with real coefficients, let $\eta_1 \leq \eta_2 \leq \dots \leq \eta_n$ denote its roots, let $m(k) = \sum_{j=1}^n \eta_j^k$, and let $\mathcal{M}(z) = \sum_{k=0}^{\infty} m(k) z^k$ denote the generating function of the sums of powers of the roots. Then*

$$\mathcal{M}(z) = -\frac{z\widehat{p}'(x)}{\widehat{p}(x)} + n \quad (3)$$

where $\widehat{p}(x) = \sum_{j=0}^n b_{n-j} x^j$ denotes the conjugate polynomial.

PROOF: According to the Newton identities one has

$$\sum_{j=0}^k m(k-j) b_{n-j} = (n-k) b_{n-k} \quad (4)$$

The following computation is straightforward:

$$\begin{aligned} \mathcal{M}(z)\widehat{p}(z) &= \sum_{l=0}^{\infty} m(l) z^l \sum_{j=0}^n b_{n-j} z^j = \sum_{k=0}^{\infty} \sum_{j=0}^k m(k-j) b_{n-j} z^k = \\ &= \sum_{k=0}^{\infty} (n-k) b_{n-k} z^k = n\widehat{p}(z) - z\widehat{p}'(z) \end{aligned}$$

hence

$$\mathcal{M}(z) = -\frac{z\widehat{p}'(z)}{\widehat{p}(z)} + n$$

so the proof is complete. \square

Let us return to the proof of the proposition. Introduce the notation $\mathcal{M}_n(z) := \sum_{k=0}^{\infty} M_n(k)z^k$. We are going to show that

$$\frac{1}{n}\mathcal{M}_n(z/\sqrt{n}) \rightarrow \sum_{k=0}^{\infty} C_k z^{2k}, \quad \text{for } 0 \leq z \leq \frac{1}{3}, \quad (5)$$

where $C_k = \frac{1}{k+1} \binom{2k}{k}$ is the k^{th} Catalan number.

The proof of this claim will be based on the well-known recursive identities of the Hermite polynomials (similar as in [12]):

$$\begin{aligned} H_0(x) &= 1 \\ H_1(x) &= x \\ H_{n+1}(x) &= xH_n(x) - nH_{n-1}(x) \\ H'_n(x) &= nH_{n-1}(x). \end{aligned}$$

Denoting by $\widehat{H}'_n(z) = x^n H_n(\frac{1}{x})$ the conjugate polynomial it can be easily checked that

$$\widehat{H}_0(x) = 1 \quad (6)$$

$$\widehat{H}_1(x) = 1 \quad (7)$$

$$\widehat{H}_{n+1}(x) = \widehat{H}_n(x) - nx^2\widehat{H}_{n-1}(x) \quad (8)$$

$$\widehat{H}'_n(x) = \frac{n}{x} \left(\widehat{H}_n(x) - \widehat{H}_{n-1}(x) \right). \quad (9)$$

Since all the roots of H_n are no greater in absolute value than $2\sqrt{n + \frac{1}{2}}$ we obtain that the conjugate polynomials do not vanish in the interval $[-\frac{1}{3\sqrt{n}}, \frac{1}{3\sqrt{n}}]$. Since $\widehat{H}_n(0) = 1$, they are in fact positive in that interval. This observation combined with equation (8) above implies that in this interval

$$\frac{\widehat{H}_{n-1}(x)}{\widehat{H}_n(x)} \leq \frac{1}{nx^2},$$

consequently

$$\frac{\widehat{H}_{n-1}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})} \leq \frac{1}{z^2}, \quad \text{for } |z| \leq \frac{1}{3}. \quad (10)$$

On the other hand using Lemma 1 equation (9) implies that

$$\mathcal{M}_n(z) = n \left(\frac{\widehat{H}_{n-1}(z)}{\widehat{H}_n(z)} \right).$$

Furthermore $\mathcal{M}_n(z)$ is a monotonically increasing (on $z \geq 0$), convex function (due to its definition and the fact that $M_n(k) = 0$ when k is odd).

Now

$$\frac{\widehat{H}_{n-1}(x)}{\widehat{H}_n(x)} = 1 + \int_0^x \left(\frac{\widehat{H}_{n-1}(y)}{\widehat{H}_n(y)} \right)' dy > 1 + \frac{x}{2} \left(\frac{\widehat{H}_{n-1}(y)}{\widehat{H}_n(y)} \right)' \Big|_{y=x/2},$$

since $\frac{\widehat{H}_{n-1}(x)}{\widehat{H}_n(x)}$ is a positive, convex, monotonically increasing function on $x \geq 0$, hence

$$\left(\frac{\widehat{H}_{n-1}(y)}{\widehat{H}_n(y)} \right)' \Big|_{y=z/\sqrt{n}} \leq \left(\frac{1}{4z^2} - 1 \right) \frac{\sqrt{n}}{z}, \quad \text{for } 0 \leq z \leq \frac{1}{3}$$

which means that $(\widehat{H}_{n-1}/\widehat{H}_n)'(z/\sqrt{n}) = O(\sqrt{n})$. Straightforward computation gives that

$$\left(\frac{\widehat{H}_{n-1}}{\widehat{H}_n}\right)'(x) = \frac{n}{x} \left(\frac{\widehat{H}_{n-1}^2(x)}{\widehat{H}_n^2(x)} - \frac{\widehat{H}_{n-2}(x)}{\widehat{H}_n(x)}\right) + \frac{1}{x} \left(\frac{\widehat{H}_{n-2}(x)}{\widehat{H}_n(x)} - \frac{\widehat{H}_{n-1}(x)}{\widehat{H}_n(x)}\right)$$

hence

$$\begin{aligned} \left(\frac{\widehat{H}_{n-1}}{\widehat{H}_n}\right)' \left(\frac{z}{\sqrt{n}}\right) &= \frac{n\sqrt{n}}{z} \left(\frac{\widehat{H}_{n-1}^2(z/\sqrt{n})}{\widehat{H}_n^2(z/\sqrt{n})} - \frac{\widehat{H}_{n-2}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})}\right) + \\ &+ \frac{\sqrt{n}}{z} \left(\frac{\widehat{H}_{n-2}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})} - \frac{\widehat{H}_{n-1}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})}\right) \end{aligned}$$

and so

$$\frac{\widehat{H}_{n-1}^2(z/\sqrt{n})}{\widehat{H}_n^2(z/\sqrt{n})} - \frac{\widehat{H}_{n-2}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})} = O\left(\frac{1}{n}\right), \quad \text{for } 0 \leq z \leq \frac{1}{3}. \quad (11)$$

Now let $f_n(z) := \frac{\widehat{H}_{n-1}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})}$, then equation (8) implies that

$$1 = f_n(z) - \frac{n-1}{n} z^2 f_n(z) f_{n-1} \left(\sqrt{\frac{n-1}{n}} \cdot z\right)$$

and from (11) it follows that

$$\begin{aligned} f_n^2(z) - f_n(z) f_{n-1} \left(\sqrt{\frac{n-1}{n}} \cdot z\right) &= \frac{\widehat{H}_{n-1}^2(z/\sqrt{n})}{\widehat{H}_n^2(z/\sqrt{n})} - \frac{\widehat{H}_{n-2}(z/\sqrt{n})}{\widehat{H}_n(z/\sqrt{n})} = \\ &= O\left(\frac{1}{n}\right). \end{aligned}$$

Therefore if for some fixed z in the interval above $h(z)$ is a limit point of the sequence $f_n(z)$ then it satisfies the following equation:

$$h(z) = 1 - z^2 h(z)^2. \quad (12)$$

In fact

$$\begin{aligned} 1 &= f_n(z) - \frac{n-1}{n} z^2 f_n^2(z) - \frac{n-1}{n} z^2 \cdot \left[f_n(z) f_{n-1} \left(\sqrt{\frac{n-1}{n}} \cdot z\right) - f_n^2(z) \right] \\ 1 &= f_n(z) - \frac{n-1}{n} z^2 f_n^2(z) + O\left(\frac{1}{n}\right). \end{aligned}$$

It is well-known that the smaller of the two real solutions of (12) is $c(z^2) = \sum_{k=0}^{\infty} C_k z^{2k}$. Since according to (10) the sequence $f_n(z)$ is uniformly bounded in order to prove the convergence of the whole sequence it is enough to prove that

$$f_n(z) \leq c(z^2),$$

for all n , where $0 \leq z \leq \frac{1}{3}$, implying that $c(z^2)$ is the only limit point of this sequence.

For $n = 1$ one has $f_1(z) = 1 \leq c(z^2)$. Equation (8) implies that

$$\begin{aligned} 1 &= \frac{\widehat{H}_{n-1}}{\widehat{H}_n} \left(\frac{z}{\sqrt{n-1}}\right) - z^2 \frac{\widehat{H}_{n-2}}{\widehat{H}_n} \left(\frac{z}{\sqrt{n-1}}\right) = \\ &= f_n \left(\sqrt{\frac{n}{n-1}} \cdot z\right) - z^2 f_n \left(\sqrt{\frac{n}{n-1}} \cdot z\right) f_{n-1}(z). \end{aligned}$$

Let us look at the following map $\xi \mapsto \eta(\xi)$, where

$$1 = \eta(\xi) - z^2 \eta(\xi) \xi,$$

and $\xi \in [0, 1/z^2]$ where $0 \leq z \leq 1/3$ is arbitrary, but fixed. Note that the fixed points of $\xi \rightarrow \eta(\xi)$ are the same as the solutions to (12). Expressing η in term of ξ we get

$$\eta(\xi) = \frac{1}{1 - z^2\xi}.$$

η is a strictly monotonically increasing function on the set $0 \leq \xi < \frac{1}{z^2}$ with $\eta(0) = 1 \leq c(z^2)$, and so $\eta(\xi) \leq c(z^2)$ if $\xi \leq c(z^2)$. Now put $\xi = f_{n-1}(z)$, then $f_n(\sqrt{n/(n-1)} \cdot z) = \eta(\xi) \leq c(z^2)$. Since f_n is monotonically increasing we have

$$f_n(z) \leq f_n\left(\sqrt{\frac{n}{n-1}} \cdot z\right) \leq c(z^2)$$

as well. Thus the only accumulation point of $(f_n(z))_{n \in \mathbb{N}}$ is $c(z^2)$. Remember that $\frac{1}{n}\mathcal{M}_n(z/\sqrt{n}) = f_n(z)$, hence the proof of (5) is complete.

Now the convergence of the power series on the interval $[0, \frac{1}{3}]$ implies the convergence of the coefficients, so

$$\frac{1}{2^k n^{\frac{k}{2}+1}} M_n(k) \rightarrow \begin{cases} \frac{C_{k/2}}{2^k} & k \text{ is even,} \\ 0 & k \text{ is odd.} \end{cases}$$

Shortly, they tend to the corresponding moments of the semicircle distribution since

$$\int_{\mathbb{R}} x^k \rho_{sc}(x) dx = \begin{cases} \frac{C_{k/2}}{2^k} & k \text{ is even,} \\ 0 & k \text{ is odd.} \end{cases}$$

This concludes the proof of the proposition. □

The following theorem shows that instead of the asymptotic property of the moments presented above a stronger statement holds.

Theorem 1 *Let $k, n \in \mathbb{N}$ and $k \geq n$. Then $M_n(k)$ is a polynomial in n , where $M_n(k) = 0$ when k is odd, while*

$$\deg_n M_n(k) = \frac{k}{2} + 1$$

when k is even. In these cases the coefficient of $n^{k/2+1}$ in $M_n(k)$ is given by the Catalan number $C_{k/2}$. In particular

$$M_n(k) = \begin{cases} n^{k/2+1} C_{k/2} + O(n^{k/2}) & k \text{ is even;} \\ 0 & k \text{ is odd,} \end{cases} \quad (13)$$

where $C_{k/2} = \frac{\binom{k}{k/2}}{k/2+1}$.

PROOF: First let us note that $a_{n-k}^{(n)} = 0$ when k is an odd number thus by induction we obtain that $M_n(k) = 0$ for $k = 1, 3, \dots$. Using this fact let us write Newton's identities (4) in the following matrix form:

$$\begin{bmatrix} 1 & 0 & \dots & \dots & 0 \\ a_{n-2}^{(n)} & 1 & 0 & \dots & 0 \\ & & \ddots & & \vdots \\ & & & \ddots & 0 \\ a_{n-2(k-1)}^{(n)} & \dots & \dots & a_{n-2}^{(n)} & 1 \end{bmatrix} \begin{bmatrix} M_n(2) \\ M_n(4) \\ \vdots \\ \vdots \\ M_n(2k) \end{bmatrix} = \begin{bmatrix} -2a_{n-2}^{(n)} \\ -4a_{n-4}^{(n)} \\ \vdots \\ \vdots \\ -2ka_{n-2k}^{(n)} \end{bmatrix}. \quad (14)$$

Since the determinant of the matrix standing on the left hand side is 1 we obtain that

$$M_n(2k) = \det \begin{bmatrix} 1 & 0 & 0 & \dots & -2a_{n-2}^{(n)} \\ a_{n-2}^{(n)} & 1 & 0 & \dots & -4a_{n-4}^{(n)} \\ & & \ddots & & \vdots \\ & & & \ddots & \vdots \\ a_{n-2(k-1)}^{(n)} & a_{n-2(k-2)}^{(n)} & \dots & a_{n-2}^{(n)} & -2ka_{n-2k}^{(n)} \end{bmatrix}. \quad (15)$$

In order to compute the determinant above introduce the notation

$$A(k, l) := \det \begin{bmatrix} 1 & 0 & 0 & \cdots & \binom{n}{l+1} \\ -\frac{n(n-1)}{2} & 1 & 0 & \cdots & -\frac{\binom{n}{l+3}}{2} \\ & \ddots & \ddots & & \\ & & & 1 & \vdots \\ \frac{(-1)^{k-1} \binom{n}{2(k-1)}}{2^{k-1} (k-1)!} & \cdots & \cdots & -\frac{n(n-1)}{2} & \frac{(-1)^{k-1} \binom{n}{l+2(k-1)+1}}{(k-1)! 2^{k-1}} \end{bmatrix} \quad (16)$$

for $k \geq 2, l \geq 1$, where $\binom{n}{l} = n(n-1) \cdots (n-l+1)$. ($\binom{n}{0} = 1$.)

Observe that $M_n(2k) = A(k, 1)$. Multiply the first column by $\binom{n}{l+1}$ and subtract it from the last one, then the j^{th} element ($j \geq 2$) of the last column is given as:

$$\begin{aligned} & \frac{(-1)^{j-1}}{2^{j-1} (j-1)!} \binom{n}{l+2(j-1)+1} - \frac{(-1)^{j-1}}{2^{j-1} (j-1)!} \binom{n}{l+1} \binom{n}{2(j-1)} = \\ & = \frac{(-1)^{j-1}}{2^{j-1} (j-1)!} \binom{n}{2(j-1)} [(n-2j+2)_{l+1} - \binom{n}{l+1}] = \\ & = \frac{(-1)^{j-1}}{2^{j-1} (j-1)!} \binom{n}{2(j-1)} [-2(j-1)] \sum_{h=0}^l [(n-2j+2)_h \times (n-h-1)_{l-h}] \\ & = \frac{(-1)^{j-2}}{2^{j-2} (j-2)!} \sum_{h=0}^l [\binom{n}{2(j-1)+h} \times (n-h-1)_{l-h}]. \end{aligned}$$

due to the fact that

$$\prod_{j=0}^l x_j - \prod_{j=1}^l y_j = \sum_{h=0}^l \left(\prod_{j:j < h} x_j (x_h - y_h) \prod_{j:j > h} y_j \right).$$

Let us observe that now the first element in the last column is zero, while all the other elements can be written as sums with $l+1$ elements, where in the i^{th} summand every element is multiplied by the same factor $(n-h-1)_{l-h}$. Introducing the notation $i = h-1$ we obtain that for $k \geq 3$

$$A(k, l) = \sum_{i=1}^{l+1} A(k-1, i) (n-i)_{l-i+1} \quad (17)$$

For $k = 2$

$$\begin{aligned} A(2, l) &= \det \begin{bmatrix} 1 & n(n-1) \cdots (n-l) \\ -\frac{n(n-1)}{2} & -\frac{1}{2} n(n-1) \cdots (n-l-2) \end{bmatrix} \\ &= \frac{n(n-1) \cdots (n-l)}{2} (n(n-1) - (n-l-1)(n-l-2)) \\ &= \frac{n(n-1) \cdots (n-l)}{2} ((2l+2)n - (l+1)(l+2)). \end{aligned}$$

implying that $\deg_n A(2, l) = l+2$ and the highest degree coefficient is $l+1 > 0$, so for $k = 2$ and arbitrary l we have that $\deg_n A(2, l) = l+2$ with positive highest degree coefficient. Suppose the claim is true for $k-1$ and all l , then

$$\begin{aligned} \deg_n A(k, l) &= \deg_n \left(\sum_{i=1}^{l+1} \prod_{j=i}^l (n-j) A(k-1, i) \right) \\ &\leq \max_i (l-i+1 + \deg_n A(k-1, i)) = k+l. \end{aligned} \quad (18)$$

On the other hand since the highest degree coefficients of $A(k-1, i)$ terms are positive by induction and they are multiplied with factors with highest degree coefficients in n equal to 1, we get that there is no cancellation, that is $\deg_n A(k, l) = k+l$, and its highest degree coefficient remains positive. In particular $\deg_n M_n(2k) = \deg_n A(k, 1) = k+1$.

What remains is to prove that the leading coefficient of $M_n(2k)$ is $C_k = \binom{2k}{k}/(k+1)$. First let us observe that $A(2, l)$ can be written as

$$A(2, l) = \sum_{i=1}^{l+1} \binom{l+1}{i} (n-i)_{l-i+1}.$$

Thus introducing the notation $A(1, l) = (n)_{l+1}$ we obtain that the recursive formula (17) remains true for $k = 2$. Furthermore, with $A(0, 1) = n$, $A(0, l) = 0$ for $l \geq 2$ we might extend the validity of the formula for $k = 1$, as well. Since in the recursive formula for $A(k, l)$ the factors for $A(k-1, i)$ are with leading coefficient one, the leading coefficient of $A(k, l)$ can be obtained as the sum of that of $A(k-1, 1), \dots, A(k-1, l+1)$. Applying now the recursive formula for the elements $A(k-1, i)$ and so on, we obtain that the leading coefficient in n of $A(k, 1)$ is given by the number of $A(1, t)$ terms in the representation of $A(k, 1)$ by these elements. This question can be translated in the following graph theoretical question:

Let us consider the following (directed) graph $((\mathbb{Z}_{\geq 0})^2, E)$: For $a := (i_1, j_1), b := (i_2, j_2) \in (\mathbb{Z}_{\geq 0})^2$ there is an edge from a to b , i.e. $(a, b) \in E$ if and only if $i_2 = i_1 + 1$ and $j_2 = j_1 - 1 + h$ for $h \geq 0$. Let $a(j)$ denote a 's j^{th} coordinate for $j = 1, 2$. The number of simple (directed) paths from the origin to $(k, 0)$ is exactly the coefficient of n^{k+1} in $A(k, 1) = M_n(2k)$. It can be checked easily that for $k = 1$ it is 1, for $k = 2$ it is 2, for $k = 3$ it is 5.

Claim: The number of simple paths from the origin to $(k, 0)$ is exactly C_k .

Proof: By induction on k . Denote by d_{k+1} the number of simple paths from the origin to $(k+1, 0)$, define $d_0 := 1$, denote by o the origin and denote by \mathcal{P}_{k+1} the collection of (directed) simple paths from $(0, 0)$ to $(k+1, 0)$, i.e.

$$\mathcal{P}_{k+1} := \{(o, a_1, a_2, \dots, a_{k+1}) \mid a_i \in (\mathbb{Z}_{\geq 0})^2, \\ a_i(1) = i, (a_{i-1}, a_i) \in E, 1 \leq i \leq k+1, a_{k+1}(2) = 0\},$$

where $a_0 = o$. For any path $P = (o, \dots, a_m)$ set

$$t(P) := \inf\{j \geq 1 \mid a_j = (j, 0)\}$$

and let $\mathcal{P}_{k+1}(i) := \{P \in \mathcal{P}_{k+1} \mid t(P) = i+1\}$. Note that $t(P) = i+1$ means that the first node of the path P whose second coordinate is 0 and differs from the origin is a_{i+1} . It is easy to see that

$\mathcal{P}_{k+1} = \bigcup_{0 \leq i \leq k}^* \mathcal{P}_{k+1}(i)$, hence $|\mathcal{P}_{k+1}| = \sum_{i=0}^k |\mathcal{P}_{k+1}(i)|$. We want to show that

$$|\mathcal{P}_{k+1}(i)| = |\mathcal{P}_i| |\mathcal{P}_{k-i}| = d_i d_{k-i}.$$

Note that $\mathcal{P}_0 = \{(o)\}$ and $\mathcal{P}_1 = \{(o, (1, 0))\}$. Given two paths $P_1 = (o, a_1, \dots, a_i) \in \mathcal{P}_i$ and $P_2 = (o, b_1, \dots, b_{k-i}) \in \mathcal{P}_{k-i}$ one can make a path $P \in \mathcal{P}_{k+1}(i)$ as follows: let us construct $P = (o, c_1, \dots, c_{k+1})$ in such a way that $c_j(1) := a_j(1)$, $c_j(2) := a_j(2) + 1$ for $1 \leq j \leq i$ and $c_j(1) := b_{j-i-1}(1) + i + 1$, $c_j(2) := b_{j-i-1}(2)$ for $i < j \leq k+1$. Remember that $b_0 = o$. It is trivial that $c_{k+1} = (k+1, 0)$ and due to the definition of the graph $(c_j, c_{j+1}) \in E$ for $0 \leq j \leq k+1$. Since $a_j(2) \geq 0$ for $1 \leq j \leq i$, so $c_j(2) \geq 1$ for these indices, while $c_{i+1} = (i+1, 0)$ thus we obtain that $t(P) = i+1$, therefore $P \in \mathcal{P}_{k+1}(i)$. Now take a path $P = (o, c_1, \dots, c_{k+1}) \in \mathcal{P}_{k+1}(i)$. Let $P_1 = (o, a_1, \dots, a_i)$ be defined by $a_j(1) := c_j(1)$ and $a_j(2) := c_j(2) - 1$ for $1 \leq j \leq i$. Since $t(P) = i+1$, one has (for $i \geq 1$) that $c_j \geq 1$ for $1 \leq j \leq i$, therefore $a_j(2) \geq 0$ for $1 \leq j \leq i$. Furthermore, for $i \geq 1$, $c_i = (i, 1)$ – because of $c_{i+1} = (i+1, 0)$ – implying that $a_i = (i, 0)$. Obviously P_1 is a valid path (due to the structure of the graph) and its last node is $(i, 0)$. The number of such paths is given by d_i . Define $P_2 = (o, b_1, \dots, b_{k-i})$ by $b_j(1) = c_{j+i+1}(1) - i - 1$ and $b_j(2) = c_{j+i+1}(2)$. Obviously $b_0 = o = (0, 0)$, and $b_{k-i}(1) = c_{k+1}(1) - i - 1 = k - i$, $b_{k-i}(2) = c_{k+1}(2) = 0$, hence $P_2 \in \mathcal{P}_{k-i}$. The number of such paths is d_{k-i} . (Note that in case of $i = 0$, $\mathcal{P}_0 = \{(o)\}$, $P_1 = (o)$ and $P_2 = (o, b_1, \dots, b_k)$, so one only has to concatenate the two paths in such a way that the coordinates of the nodes of the path are valid, i.e. $c_0 = o$, $c_1 = (1, 0)$, $c_{j+1} = (b_j(1) + 1, b_j(2))$ for $1 \leq j \leq k$. If $P \in \mathcal{P}_{k+1}(0)$ is given, then $P = (o, (1, 0), c_2, \dots, c_{k+1})$ thus $P_1 := (o)$ and $P_2 = (o, b_1, \dots, b_k)$ with $b_j(1) = c_{j+1}(1) - 1$ and $b_j(2) = c_{j+1}(2)$ for $1 \leq j \leq k$.) Now it is easy to see that we found a bijection between $\mathcal{P}_{k+1}(i)$ and $\mathcal{P}_i \times \mathcal{P}_{k-i}$, hence

$$d_{k+1} = \sum_{i=0}^k d_i d_{k-i}. \quad (19)$$

Thus the sequence d_0, d_1, \dots satisfies the same recursion which is valid for the Catalan number. Since as we pointed out above the first two terms of these sequences coincide we obtain by induction that $d_k = C_k = \binom{2k}{k}/(k+1)$ for $k \geq 0$. \square

Remark 1 Let us point out that similar argument shows that the highest degree coefficient in $A(k, l)$ is given by the number of simple paths in the graph defined above starting from the origin ending in $(k, l-1)$.

2 Expectation of the characteristic polynomial

Keeping in mind as also mentioned in the introduction that the limit distribution of the normalized eigenvalues of a Wigner matrix is given by the semicircle law, we now turn to random matrix theory. In the previous section we showed that the limit distribution of the normalized roots of the monic Hermite polynomials is the semicircle distribution. As we will see throughout this section this is not a coincidence, since the expectation of the characteristic polynomial of an $n \times n$ Wigner matrix is exactly the n^{th} (monic) Hermite polynomial. Since the size $n \times n$ of the Wigner matrix will remain unchanged in this section, so for the sake of simplicity we will denote it by A instead of A_n . Diaconis and Gamburd proved in [5] that if A is such that its elements satisfy $P(a_{ij} = \pm 1) = 1/2$ for $1 \leq i \leq j \leq n$, one has

$$E \det[\lambda I - A]^k = H_n^{(k/2)}(x) \quad k \in \mathbb{Z}_{\geq 0},$$

where $H_n^{(k/2)}(x)$ is the generalized Hermite polynomial. These polynomials are orthogonal with respect to the measure $|x|^k e^{-x^2/2} dx$ on \mathbb{R} . Later for $m = 0$ Gamburd and Forrester generalized this theorem in [6], where they proved that if $A = [a_{ij}]$ is a random symmetric matrix, where the diagonal elements are independent and identically distributed with $Ea_{ii} = 0$ for $1 \leq i \leq n$, and the off-diagonal elements are also independent (from the diagonal elements as well) and identically distributed with $Ea_{ij} = 0$ and $D^2 a_{ij} = c^2$ for $1 \leq i < j \leq n$, then one has

$$E \det[\lambda I - A] = c^n H_n(x/c).$$

Their proof goes per definition, i.e. computing

$$E \det[\lambda I - A] = \sum_{\sigma \in S_n} (-1)^{|\sigma|} E \prod_{i=1}^n (\lambda_i \delta_{i\sigma(i)} - a_{i\sigma(i)}).$$

For the sake of completeness we reprove this theorem using a formula for the secular coefficients of the characteristic polynomial in terms of the matrix elements and by showing that the expectation of these coefficients is the same as the appropriate coefficients of the n^{th} monic Hermite polynomial.

Theorem 2 Consider a Wigner matrix A of size $n \times n$, defined as in the introduction, normalized as $Ea_{ij}^2 = 1$, when $i \neq j$. Denote by p its random characteristic polynomial, i.e. $p(x) = \det(xI - A)$. Then

$$Ep(x) = H_n(x), \quad (20)$$

where H_n is the monic Hermite polynomial of degree n .

PROOF: For the proof we will need the following proposition stated without proof (see e.g. [4]).

Proposition 2 Let \mathcal{A} be an arbitrary matrix of size $n \times n$. Then

$$p(x) = \det(xI - \mathcal{A}) = \sum_{k=0}^n (-1)^k \sum_{\substack{I \subset N \\ |I|=k}} \det \mathcal{A}_I x^{n-k},$$

where \mathcal{A}_I denotes the $k \times k$ submatrix of \mathcal{A} spanned by columns and rows indexed by $i_1, \dots, i_k \in I$ and $N = [1, n] \cap \mathbb{N}$.

Based on this result we have that

$$Ep_n(x) = \sum_{k=0}^n (-1)^k x^{n-k} \sum_{\substack{I \subset N \\ |I|=k}} E \det \mathcal{A}_I. \quad (21)$$

On the other hand

$$E \det \mathcal{A}_I = \sum_{\alpha \in S_k(I)} (-1)^{|\alpha|} E \prod_{i \in I} a_{i\alpha(i)},$$

where $S_k(I)$ denotes the set of permutation of the elements in I . Since our matrix A is symmetric with independent elements (i.e. (a_{ij}) , $1 \leq i \leq j \leq n$ are independent random variables and $a_{ij} = a_{ji}$ for $i > j$), we have

$$E[a_{i_1 \alpha_{i_1}} \cdots a_{i_k \alpha_{i_k}}] \neq 0 \Leftrightarrow k \text{ is even and } E[a_{i_1 \alpha_{i_1}} \cdots a_{i_k \alpha_{i_k}}] = E \prod_{j=1}^{k/2} a_{i'_j i''_j}^2$$

for some indices $1 \leq i'_j, i''_j \leq n$. This means that the only permutations that count are those which can be written as the product of $k/2$ disjoint 2-cycles. There is $(k-1)!!$ of these, hence

$$\sum_{\alpha \in S_k(I)} (-1)^{|\alpha|} E[a_{i_1 \alpha_{i_1}} \cdots a_{i_k \alpha_{i_k}}] = (-1)^{k/2} (k-1)!! ,$$

which is independent of the special choice of the subset $I \subset N$ with $|I| = k$, so

$$Ep(x) = \sum_{0 \leq k \leq n, 2|k} (-1)^{k/2} \binom{n}{k} (k-1)!! x^{n-k} = \sum_{0 \leq k \leq n, 2|k} \frac{(-1)^{k/2} \binom{n}{k} k!}{2^{k/2} (k/2)!} x^{n-k}$$

In order to finish the proof of the theorem we have to show that the coefficients above are equal to the coefficients of the monic Hermite polynomial H_n . In the following proposition we give a short proof of this fact. (See Szegő [13].)

Proposition 3 *Let us denote by $H_n(x) = \sum_{j=0}^n a_j^{(n)} x^j$ the Hermite polynomial of degree n . Then*

$$a_{n-k}^{(n)} = \begin{cases} (-1)^{k/2} \frac{\binom{n}{k} k!}{(k/2)! 2^{k/2}} & k \text{ is even;} \\ 0 & k \text{ is odd.} \end{cases} \quad (22)$$

PROOF: The proof is based on the well known fact that

$$H(t, x) = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} = e^{tx - t^2/2}.$$

Since the roots of the Hermite polynomials are lying symmetrically on the real line for odd k one has $\sum_{j=1}^n (\xi_j^{(n)})^k = 0$, where $\xi_j^{(n)}$ denotes the j^{th} largest root of H_n . Instead of $a_k^{(n)}$ we consider $a_{n-k}^{(n)}$, because in this case the parity of k determines whether it will be different from zero, or not. Hence let us consider the exponential generating function G_k of $(a_{n-k}^{(n)})_{n \geq 0}$

$$G_k(z) = \sum_{n=k}^{\infty} a_{n-k}^{(n)} \frac{z^n}{n!}. \quad (23)$$

Note that

$$\sum_{k=0}^{\infty} G_k(z) y^k = \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} a_{n-k}^{(n)} \frac{z^n}{n!} y^k = \sum_{n=0}^{\infty} \widehat{H}_n(y) \frac{z^n}{n!},$$

where $\widehat{H}_n(y) = \sum_{j=0}^n a_{n-j}^{(n)} y^j$. Since $\widehat{H}_n(y) = y^n H_n(y^{-1})$, one has that

$$\sum_{k=0}^{\infty} G_k(z) y^k = \sum_{n=0}^{\infty} H_n(y^{-1}) \frac{(yz)^n}{n!} = H(yz, y^{-1}) = e^{z - (yz)^2/2}, \quad (24)$$

so

$$G_k(z) = \frac{1}{k!} \left. \frac{\partial^k}{\partial y^k} H(yz, \frac{1}{y}) \right|_{y=0} = \frac{1}{k!} e^z \left. \frac{\partial^k}{\partial y^k} e^{-\frac{(yz)^2}{2}} \right|_{y=0}.$$

But since

$$\frac{\partial^k}{\partial y^k} e^{-\frac{(yz)^2}{2}} = \frac{\partial^k}{\partial y^k} \sum_{n=0}^{\infty} \frac{(-1)^n (yz)^{2n}}{n! 2^n} = \sum_{n: n \geq \lceil k/2 \rceil}^{\infty} \frac{(2n)!}{(2n-k)!} \frac{(-1)^n (yz)^{2n-k} z^k}{n! 2^n}$$

we have

$$G_k(z) = \begin{cases} z^k e^z (-1)^{k/2} \frac{1}{(k/2)! 2^{k/2}} & k \text{ is even;} \\ 0 & k \text{ is odd.} \end{cases}$$

Using this formula one gets

$$a_{n-k}^{(n)} = \left. \frac{\partial^n}{\partial z^n} G_k(z) \right|_{z=0} = \begin{cases} (-1)^{k/2} \frac{\binom{n}{k} k!}{(k/2)! 2^{k/2}} & k \text{ is even;} \\ 0 & k \text{ is odd,} \end{cases} \quad (25)$$

hence the proof of this proposition is complete. \square This concludes the proof of the theorem, as well. \square

3 Conclusions

The results show, that in order to get the Hermite polynomials as the expectation of a characteristic polynomial of a Wigner matrix, one only has to assume independence and the existence and finiteness of the second moments. The limiting distribution of the roots of the H_n is the semicircle law as it is shown in Proposition 1. It is also known that the limiting distribution of the eigenvalues of a properly scaled Wigner matrix is given by the semicircle law [1] in the same sense as above, hence there is a deep connection between the Hermite polynomials, random symmetric matrices with independent elements and the semicircle law. This suggests that studying the Hermite polynomials and their roots could give us an insight on the behavior of the eigenvalues of a Wigner random matrix.

References

- [1] L. Arnold. On the asymptotic distribution of the eigenvalues of random matrices. *Journal of Mathematical Analysis and Applications*, 20(2):262–268, 1967.
- [2] Z. D. Bai and Y. Q. Yin. Convergence to the semicircle law. *The Annals of Probability*, pages 863–875, 1988.
- [3] E. Brézin and S. Hikami. Characteristic polynomials of random matrices. *Communications in Mathematical Physics*, 214(1):111–135, 2000.
- [4] R. A. Brualdi and D. Cvetkovic. *A combinatorial approach to matrix theory and its applications*. CRC press, 2008.
- [5] P. Diaconis and A. Gamburd. Random matrices, magic squares and matching polynomials. *Journal of Combinatorics*, 11(4):R2, 2004.
- [6] P. J. Forrester and A. Gamburd. Counting formulas associated with some random matrix averages. *Journal of Combinatorial Theory, Series A*, 113(6):934–951, 2006.
- [7] W. Gawronski. On the asymptotic distribution of the zeros of hermite, laguerre, and jonquiere polynomials. *Journal of Approximation Theory*, 50(3):214–231, 1987.
- [8] V. L. Girko. Spectral theory of random matrices. *Russian Mathematical Surveys*, 40(1):77–120, 1985.
- [9] A. Hardy et al. Average characteristic polynomials of determinantal point processes. In *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, volume 51, pages 283–303. Institut Henri Poincaré, 2015.
- [10] W. Lang. On sums of powers of zeros of polynomials. *Journal of Computational and Applied Mathematics*, 89(2):237–256, 1998.
- [11] V. A. Marchenko and L. A. Pastur. Distribution of eigenvalues for some sets of random matrices. *Matematicheskii Sbornik*, 114(4):507–536, 1967.

- [12] A. Martínez-Finkelshtein, P. Martínez-González, and R. Orive. On asymptotic zero distribution of laguerre and generalized bessel polynomials with varying parameters. *Journal of Computational and Applied Mathematics*, 133(1):477–487, 2001.
- [13] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Colloquium Publications, Voluma XXIII, 1939.