SZEGŐ KERNEL EXPANSION AND EQUIVARIANT EMBEDDING OF CR MANIFOLDS WITH CIRCLE ACTION

HENDRIK HERRMANN, CHIN-YU HSIAO, AND XIAOSHAN LI

ABSTRACT. Let X be a compact strongly pseudoconvex CR manifold with a transversal CR S^1 -action. In this paper, we establish the asymptotic expansion of Szegő kernels of positive Fourier components and by using the asymptotics, we show that X can be equivariant CR embedded into some \mathbb{C}^N equipped with a simple S^1 -action. An equivariant embedding of quasi-regular Sasakian manifold is also derived.

1. Introduction

Let X be a compact strongly pseudoconvex CR manifold. The question of whether or not X admits a CR embedding into a complex Euclidean space has attracted a lot of attention. By a theorem of Boutet de Monvel [6], any compact strongly pseudoconvex CR manifold of dimension greater than or equal to five can be CR embedded into \mathbb{C}^N for some N. The classical example of non-embeddable three dimensional strongly pseudoconvex CR manifold given by Rossi [33] (see also [2]) showed that an arbitrary real analytic deformation of the standard CR structure on the three-sphere may fail to be embeddable. There exists an extensive literature on the embaddability of deformed CR structures. For this subject, we refer the reader to [3, 28, 14, 10, 11] and the references therein. Given a compact strongly pseudoconvex CR manifold equipped with a locally free transversal CR S^1 -action, it was shown in [28, 29] and also [27] that X can always be CR embedded into some complex space.

In this work, we attack the embedding problem for compact strongly pseudoconvex CR manifolds equipped with a transversal CR circle action from a pure analytic point of view. More precisely, we develop an asymptotic expansion for the Szegő kernels (S_m , see Section 2.2) concerning CR functions which lie in the space of positive Fourier components ($H^0_{b,m}(X)$), see Definition 1.8). Our main results involving that features are Theorem 2.6 and Theorem 2.7 which we deduce from a general result about Szegő kernel expansion (see Theorem 2.4) using partially the machinery developed by the second and third named authors in [22, 23]. Theorem 2.6 describes the expansion on the regular part, i.e. the part of the manifold where S^1 acts globally free. Here, the expansion works out well. Difficulties occur on the complement of the regular part, but we can still prove some expansion as

The first-named author was supported by the Graduiertenkolleg 1269: "Global Structures in Geometry and Analysis" and by the Institute of Mathematica, Academia Sinica.

The second-named author was partially supported by Taiwan Ministry of Science of Technology project 104-2628-M-001-003-MY2 and the Golden-Jade fellowship of Kenda Foundation.

The third-named author was supported by NSFC No. 11501422, Postdoctoral Science Foundation of China 2015M570660 and Central university research Fund 2042015kf0049.

stated in Theorem 2.7. Roughly speaking, the problem is that the leading term of the expansion does not change smoothly (even not continuously) from the regular part to its complement.

Inspired by a work of the second named author [19], our second main result in this paper is using Theorem 2.6 and Theorem 2.7 to construct a CR embedding which is equivariant with respect to a simple S^1 -action on \mathbb{C}^N (see Theorem 1.2). Recently, Hsiao-Li-Marinescu [24] established the Kodaira embedding theorem for CR manifolds with circle action which complements the results of this paper with the study of the embedding in the presence of a positive line bundle but without the hypothesis of strict pseudoconvexity.

Before we state the precise embedding result, let us see some examples of compact strongly pseudoconvex CR manifolds in \mathbb{C}^N to get an idea how such simple S^1 -actions could look like.

Example I: Let $X = \{(z_1, z_2, \dots, z_n) \in \mathbb{C}^n : |z_1|^2 + |z_2|^2 + |z_3|^2 + \dots + |z_n|^2 = 1\}$ which is a CR manifold with a transversal CR S^1 -action (see Definition 1.5):

$$e^{i\theta} \circ (z_1, z_2, \dots, z_n) = (e^{im_1\theta} z_1, e^{im_2\theta} z_2, \dots, e^{im_n\theta} z_n),$$

where $(m_1, \ldots, m_n) \in \mathbb{N}^n$.

Example II: $X = \Big\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : \ |z_1|^2 + |z_2|^2 + |z_3|^2 + |z_1^2 + z_2|^4 + |z_2^3 + z_3|^6 = 1 \Big\}.$ Then X admits a transversal CR locally free S^1 -action:

$$e^{i\theta} \circ (z_1, z_2, z_3) = (e^{i\theta} z_1, e^{2i\theta} z_2, e^{6i\theta} z_3).$$

Definition 1.1. We say that an S^1 -action $e^{i\theta}$ on \mathbb{C}^N is simple if

$$e^{i\theta} \circ (z_1, \dots, z_N) = (e^{im_1\theta} z_1, \dots, e^{im_N\theta} z_N), \quad \forall (z_1, \dots, z_N) \in \mathbb{C}^N, \quad \forall \theta \in [0, 2\pi),$$

where $(m_1, \ldots, m_N) \in \mathbb{N}^N$. The minimal weight of a simple S^1 -action on \mathbb{C}^N is given by $\min_{1 \leq j \leq N} m_j$.

Our goal is to prove the following equivariant embedding theorem for compact strongly pseudoconvex CR manifolds which admit a transversal CR S^1 -action.

Theorem 1.2. Let $(X, T^{1,0}X)$ be a compact strongly pseudoconvex CR manifold with a transversal CR locally free S^1 -action $e^{i\theta}$. Then, for any $m_0 \in \mathbb{N}$ we can find $N \in \mathbb{N}$, a simple S^1 -action on \mathbb{C}^N with minimal weight bigger than m_0 and an equivariant CR embedding

$$\Phi: X \to \mathbb{C}^N, \ x \mapsto (\Phi_1(x), \dots, \Phi_N(x)).$$

More precisely, Φ is an embedding, Φ_1, \ldots, Φ_N are CR functions and there exist $m_1, \ldots, m_N \in \mathbb{N}$, $m_j > m_0$ for $j = 1, \ldots, N$, such that

$$\Phi(e^{i\theta} \circ x) = (e^{im_1\theta}\Phi_1(x), \dots, e^{im_N\theta}\Phi_N(x)) = e^{i\theta} \circ \Phi(x), \quad \forall x \in X, \quad \forall \theta \in [0, 2\pi),$$
 where the simple S^1 -action on \mathbb{C}^N is given by $e^{i\theta} \circ (z_1, \dots, z_N) = (e^{im_1\theta}z_1, \dots, e^{im_N\theta}z_N).$

The equivariance, where a lower bound for the minimal weight of the simple S^1 -action on the embedding space can be chosen arbitrarily large, distinguishes Theorem 1.2 from the embedding results mentioned at the beginning of this section.

An application of Theorem 1.2 is an equivariant embedding result for quasiregular Sasakian manifolds. Sasakian manifolds have gained prominence in physics and algebraic geometry, especially in String theory (see [7, 8]). Let X be a compact smooth manifold of dimension $2n-1, n \geq 2$. The triple (X,g,α) where g is a Riemannian metric and α a real 1-form is called Sasakian manifold if the cone $\mathcal{C}(X) = \{(x,t) \in X \times \mathbb{R}_{>0}\}$ is a Kähler manifold with complex struture J and Kähler form $t^2d\alpha + 2tdt \wedge \alpha$ which are compatible with the metric $t^2g + dt \otimes dt$ (see [4], [8], [16], [31], [32]). As a consequence, X is a compact strongly pseudoconvex CR manifold and the Reeb vector field ξ , defined by $\alpha(\cdot) = g(\xi, \cdot)$, induces a transversal CR action on X. If the orbits of ξ are closed, the Sasakian structure is called quasi-regular. In this case, the Reeb vector field generates a locally free transversal CR S^1 -action on X. We thus can identify a compact quasi-regular Sasakian manifold with a compact strongly pseudoconvex CR manifold $(X, T^{1,0}X)$ with a transversal CR locally free S^1 -action where the induced vector field of the S^1 -action coincides with the Reeb vector field on X. From Theorem 1.2, we get

Theorem 1.3. Let X be a quasi-regular Sasakian manifold which admits a transversal CR S^1 -action induced by the Reeb vector field. For any $m_0 \in \mathbb{N}$, there is an equivariant CR embedding of X into some \mathbb{C}^N equipped with a simple S^1 -action with minimal weight bigger than m_0 .

The paper is organized as follows. In the rest of this section we will give an outline of the idea of the proof of Theorem 1.2 and introduce some terminology. Section 2 contains the proofs of the results on Szegő kernel expansion for positive Fourier components. The embedding result (Theorem 1.2) is proven in Section 3.

1.1. The idea of the proof of Theorem 1.2. We refer the reader to Section 1.2, Section 1.3 and Section 1.4 for some notations and terminology used here. Assume that $(X, T^{1,0}X)$ is a compact connected strongly pseudoconvex CR manifold of dimension $2n-1, n \geq 2$, with a transversal CR locally free S^1 -action $e^{i\theta}$. Let T denote the vector field induced by the S^1 -action and let $\overline{\partial}_b$ be the tangential Cauchy-Riemann operator on X. For every $m \in \mathbb{N}$, let $H^0_{b,m}(X) := \{u \in C^\infty(X) : \overline{\partial}_b u = 0, Tu = imu\}$ be the m-th positive Fourier component of the space of global smooth CR functions. The main inspiration of this paper is the following: In [23] the second and third-named author have shown that $\dim H^0_{b,m}(X) \approx m^{n-1}$ as $m \to \infty$. Hence, the space of CR functions which lie in the positive Fourier components is very large and we therefore ask whether X can be CR embedded into some \mathbb{C}^N by CR functions which lie in the positive Fourier components. In this work we give an affirmative answer to this question and as a corollary, we deduce Theorem 1.2. More precisely, we will prove

Theorem 1.4. Let X be a compact connected strongly pseudoconvex CR manifold with a locally free transversal CR S^1 -action. Then X can be CR embedded into some complex space by the CR functions which lie in the positive Fourier components.

Motivated by the second-named author's work on the Kodaira embedding theorem ([19], [20], [21]), we will use the asymptotic expansion of the Szegő kernel with respect to $H_{b,m}^0(X)$ to prove Theorem 1.2. For every $k \in \mathbb{N}$, let X_k and X_{reg} be defined as in (1.3). Let $\{f_j\}_{j=1}^{d_m} \subset H_{b,m}^0(X)$ be an orthonormal basis. The m-th

Szegő kernel $S_m(x,y)$ is given by $S_m(x,y) := \sum_{j=1}^{d_m} f_j(x) \overline{f_j(y)}$. Let us first consider $\Psi^1_m : X \to \mathbb{C}^{d_m}, x \mapsto (f_1(x), \dots, f_{d_m}(x))$.

We notice that $S_m(x,y)=0$ on X_k if $k \nmid m$. From this observation, we see that $X \setminus X_{\text{reg}} \neq \emptyset$ implies that Ψ^1_m can not be an embedding even if m is large. Suppose $X=X_1 \cup X_2 \cup \cdots \cup X_l$. For $1 \leq k \leq l$, let $\{f_j^k\}_{j=1}^{d_{km}}$ be an orthonormal basis of $H^0_{h\,km}(X)$. We next consider the map

$$\Psi_m: X \to \mathbb{C}^{\tilde{N}_m},$$

$$x \mapsto (f_1^1(x), \dots, f_{d_m}^1(x), f_1^2(x), \dots, f_{d_{2m}}^2(x), \dots, f_1^l(x), \dots, f_{d_{l_m}}^l(x)),$$

where $\tilde{N}_m = d_m + d_{2m} + \cdots + d_{lm}$. In Section 2.3, we will show that on a canonical coordinate patch $D \subset X_{\text{reg}}$ with canonical coordinates $x = (z, \theta)$, we have

$$S_m(x,y) \equiv \frac{1}{2\pi} e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \hat{b}(z,w,m) \mod O(m^{-\infty}),$$

$$(1.1) \quad \hat{b}(z,w,m) \sim \sum_{j=0}^{\infty} m^{n-1-j} \hat{b}_j(z,w), \hat{b}_j(z,w) \in C^{\infty}(D \times D), \quad j = 0, 1, 2, \cdots,$$

$$\hat{b}_0(z,z) \neq 0$$

(see Theorem 2.6). Moreover, for fixed $x_0 \in X_k$, k > 1, one finds that $k \nmid m$ implies $S_m(x, x_0) = 0$ and that $k \mid m$ leads to

(1.2)
$$S_m(x,x_0) \equiv \frac{k}{2\pi} e^{im(x_{2n-1} + \Phi(z,0))} \hat{b}(z,0,m) \mod O(m^{-\infty})$$

for some canonical coordinate patch D with canonical coordinates $x=(z,\theta)$, $x_0\in D$, $(z(x_0),\theta(x_0))=(0,0)$ (see Theorem 2.7). It should be mentioned that (1.1) and (1.2) are based on Boutet de Monvel-Sjöstrand's classical result on Szegő kernels [9] (after the seminal work [15] of Fefferman) and the complex stationary phase formula of Melin-Sjöstrand [30]. From (1.1) and (1.2), we can check that Ψ_m is an immersion if m is large. But Ψ_m is not globally injective: in general, assume that m is even, then we can not separate the points $p\in X_k$ and $e^{i\frac{\pi}{k}}\circ p$, where k>1. To overcome this difficulty, let $\{g_j^k\}_{j=1}^{d_{k(m+1)}}$ be an orthonormal basis of $H^0_{b,k(m+1)}(X)$ for $1\leq k\leq l$. For any k, $1\leq k\leq l$, we define a CR map from X to Euclidean space as follows

$$\Phi_m^k: X \to \mathbb{C}^{d_{km}+d_{k(m+1)}}, x \mapsto (f_1^k(x), \cdots, f_{d_{km}}^k(x), g_1^k(x), \cdots, g_{d_{k(m+1)}}^k(x)),$$

and we set

$$\Phi_m: X \to \mathbb{C}^{N_m}, x \to (\Phi_m^1(x), \cdots, \Phi_m^l(x)),$$

where $N_m = \sum_{k=1}^{l} (d_{km} + d_{k(m+1)})$. We thus try to prove that Φ_m is globally injective.

It is not difficult to see that Φ_m can separate the points $p \in X_k$ and $e^{i\theta} \circ p$, where $p \neq e^{i\theta} \circ p$, if m is large enough. But another difficulty comes from the fact that the expansion (1.1) converges only locally uniformly on X_{reg} and on $X \setminus X_{\text{reg}}$, we can only get an expansion of $S_m(x,x_0)$ for fixed $x_0 \in X \setminus X_{\text{reg}}$ and this causes that Φ_m could not be globally injective. To overcome this difficulty, we analyze carefully the behavior of the Szegő kernel $S_m(x,y)$ near the complement of X_{reg} and in

Section 3.2, we can construct many CR functions h_1, \ldots, h_K with large potentials near the complement of X_{reg} which lie in the positive Fourier components such that the map

$$x \in X \to (\Phi_m(x), h_1(x), \dots, h_K(x)) \in \mathbb{C}^{N_m + K}$$

is an embedding if m is large (see Theorem 3.3). This finishes the proof of Theorem 1.4.

1.2. **Set up and terminology.** Let $(X, T^{1,0}X)$ be a compact connected orientable CR manifold of dimension $2n-1, n \geq 2$, where $T^{1,0}X$ is the CR structure of X. That is $T^{1,0}X$ is a subbundle of the complexified tangent bundle $\mathbb{C}TX$ of rank n-1, satisfying $T^{1,0}X \cap T^{0,1}X = \{0\}$, where $T^{0,1}X = \overline{T^{1,0}X}$ and $[\mathcal{V}, \mathcal{V}] \subset \mathcal{V}$, where $\mathcal{V} = C^{\infty}(X, T^{1,0}X)$.

We assume that X admits an S^1 -action: $S^1 \times X \to X, (e^{i\theta}, x) \to e^{i\theta} \circ x$. Here we use $e^{i\theta}$ to denote the S^1 -action. Set $X_{\text{reg}} = \{x \in X : \forall e^{i\theta} \in S^1, \text{ if } e^{i\theta} \circ x = x, \text{ then } e^{i\theta} = \text{id}\}$. For every $k \in \mathbb{N}$, put

(1.3)
$$X_k := \left\{ x \in X : e^{i\theta} \circ x \neq x, \forall \theta \in (0, \frac{2\pi}{k}), e^{i\frac{2\pi}{k}} \circ x = x \right\}.$$

Thus, $X_{\text{reg}} = X_1$. In this paper, for simplicity we always assume that $X_1 \neq \emptyset$. Actually, one can re-normalize the S^1 -action by lifting such that the new S^1 -action satisfies $X_1 \neq \emptyset$, (see [13, Remark 1.14]).

Let $T\in C^\infty(X,TX)$ be the global real vector field induced by the S^1 -action given as follows

(1.4)
$$(Tu)(x) = \frac{\partial}{\partial \theta} \left(u(e^{i\theta} \circ x) \right) \Big|_{\theta=0}, \ u \in C^{\infty}(X).$$

Definition 1.5. We say that the S^1 -action $e^{i\theta}$ ($0 < \theta < 2\pi$) is CR if

$$[T,C^{\infty}(X,T^{1,0}X)]\subset C^{\infty}(X,T^{1,0}X),$$

where $[\ ,\]$ is the Lie bracket between the smooth vector fields on X. Furthermore, we say that the S^1 -action is transversal if for each $x\in X$ one has

$$\mathbb{C}T(x) \oplus T_x^{1,0}(X) \oplus T_x^{0,1}X = \mathbb{C}T_xX.$$

We throughout assume that $(X,T^{1,0}X)$ is a compact connected CR manifold with a transversal CR locally free S^1 -action and we denote by T the global vector field induced by the S^1 -action. Let $\omega_0 \in C^\infty(X,T^*X)$ be the global real one form uniquely determined by $\langle \, \omega_0 \, , \, u \, \rangle = 0$, for every $u \in T^{1,0}X \oplus T^{0,1}X$ and $\langle \, \omega_0 \, , \, T \, \rangle = -1$.

We recall

Definition 1.6. For $x \in X$, the Levi-form \mathcal{L}_x associated with the CR structure is the Hermitian quadratic form on $T_x^{1,0}X$ defined as follows. For any $U, V \in T_x^{1,0}X$, pick $\mathcal{U}, \mathcal{V} \in C^{\infty}(X, T^{1,0}X)$ such that $\mathcal{U}(x) = U, \mathcal{V}(x) = V$. Set

$$\mathcal{L}_x(U,\overline{V}) = \frac{1}{2i} \langle [\mathcal{U},\overline{\mathcal{V}}](x), \omega_0(x) \rangle$$

where $[\ ,\]$ denotes the Lie bracket between smooth vector fields. Note that $\mathcal{L}_x(U,\overline{V})$ does not depend on the choice of \mathcal{U} and \mathcal{V} .

Definition 1.7. The CR structure on X is called pseudoconvex at $x \in X$ if \mathcal{L}_x is semi-positive definite. It is called strongly pseudoconvex at x if \mathcal{L}_x is positive definite. If the CR structure is (strongly) pseudoconvex at every point of X, then X is called a (strongly) pseudoconvex CR manifold.

Denote by $T^{*1,0}X$ and $T^{*0,1}X$ the dual bundles of $T^{1,0}X$ and $T^{0,1}X$, respectively. Define the vector bundle of (0,q)-forms by $\Lambda^q T^{*0,1}X$. Let $D \subset X$ be an open subset. Let $\Omega^{0,q}(D)$ denote the space of smooth sections of $\Lambda^q T^{*0,1}X$ over D.

Fix $\theta_0 \in [0, 2\pi)$. Let

$$de^{i\theta_0}: \mathbb{C}T_xX \to \mathbb{C}T_{e^{i\theta_0}x}X$$

denote the differential map of $e^{i\theta_0}: X \to X$. By the properties of transversal CR S^1 -actions, we can check that

(1.5)
$$de^{i\theta_0}: T_x^{1,0}X \to T_{e^{i\theta_0}x}^{1,0}X, \\ de^{i\theta_0}: T_x^{0,1}X \to T_{e^{i\theta_0}x}^{0,1}X, \\ de^{i\theta_0}(T(x)) = T(e^{i\theta_0}x).$$

Let $(e^{i\theta_0})^*: \Lambda^q(\mathbb{C}T^*X) \to \Lambda^q(\mathbb{C}T^*X)$ be the pull back of $e^{i\theta_0}, q = 0, 1, \dots, n-1$. From (1.5), we can check that for every $q = 0, 1, \dots, n-1$

(1.6)
$$(e^{i\theta_0})^* : \Lambda^q T_{e^{i\theta_0}x}^{*0,1} X \to \Lambda^q T_x^{*0,1} X.$$

Let $u \in \Omega^{0,q}(X)$ be a section. The Lie derivative of u along the direction T is denoted by Tu. From (1.6) we have $Tu \in \Omega^{0,q}(X)$ for all $u \in \Omega^{0,q}(X)$.

Let $\overline{\partial}_b:\Omega^{0,q}(X)\to\Omega^{0,q+1}(X)$ be the tangential Cauchy-Riemann operator. From (1.5), it is straightforward to deduce

(1.7)
$$T\overline{\partial}_b = \overline{\partial}_b T \text{ on } \Omega^{0,q}(X).$$

For every $m \in \mathbb{Z}$, put $\Omega_m^{0,q}(X) := \{u \in \Omega^{0,q}(X) : Tu = imu\}$. We denote by $\overline{\partial}_{b,m}$ the restriction of $\overline{\partial}_b$ to $\Omega_m^{0,q}(X)$. From (1.7) we have the $\overline{\partial}_{b,m}$ -complex for every $m \in \mathbb{Z}$:

(1.8)
$$\overline{\partial}_{b,m}: \cdots \to \Omega_m^{0,q-1}(X) \to \Omega_m^{0,q}(X) \to \Omega_m^{0,q+1}(X) \to \cdots$$

For $m \in \mathbb{Z}$, the q-th $\overline{\partial}_{b,m}$ -cohomology is given by

(1.9)
$$H_{b,m}^{q}(X) := \frac{\operatorname{Ker} \overline{\partial}_{b} : \Omega_{m}^{0,q}(X) \to \Omega_{m}^{0,q+1}(X)}{\operatorname{Im} \overline{\partial}_{b} : \Omega_{m}^{0,q-1}(X) \to \Omega_{m}^{0,q}(X)}.$$

Moreover, we have (see Theorem 1.13 in [23])

(1.10)
$$\dim H^q_{b,m}(X) < \infty$$
, for all $q = 0, ..., n - 1$.

Definition 1.8. A function $u \in C^{\infty}(X)$ is a Cauchy-Riemann function (CR function for short) if $\overline{\partial}_b u = 0$, that is $\overline{Z}u = 0$ for all $Z \in C^{\infty}(X, T^{1,0}X)$. For $m \in \mathbb{N}$, $H^0_{b,m}(X)$ is called the m-th positive Fourier component of the space of CR functions.

1.3. Hermitian CR geometry.

Definition 1.9. Let D be an open set and let $V \in C^{\infty}(D, \mathbb{C}TX)$ be a vector field over D. We say that V is rigid if

$$de^{i\theta_0}(V(x)) = V(e^{i\theta_0}x)$$

holds for any $x \in D$ and $\theta_0 \in [0, 2\pi)$ satisfying $e^{i\theta_0} \circ x \in D$.

Definition 1.10. Let $\langle \cdot | \cdot \rangle$ be a Hermitian metric on $\mathbb{C}TX$. We say that $\langle \cdot | \cdot \rangle$ is rigid if for all rigid vector fields V, W on D, where D is any open set, we have

$$\langle V(x)|W(x)\rangle = \langle (de^{i\theta_0}V)(e^{i\theta_0}\circ x)|(de^{i\theta_0}W)(e^{i\theta_0}\circ x)\rangle, \forall x\in D, \theta_0\in [0,2\pi).$$

Lemma 1.11 (Theorem 9.2 in [21]). Let X be a compact CR manifold with a transversal CR S^1 -action. There always exists a rigid Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathbb{C}TX$ such that $T^{1,0}X \perp T^{0,1}X$, $T \perp (T^{1,0}X \oplus T^{0,1}X)$, $\langle T|T \rangle = 1$ and $\langle u|v \rangle$ is real if u,v are real tangent vectors.

From now on, we fix a rigid Hermitian metric $\langle\cdot|\cdot\rangle$ on $\mathbb{C}TX$ satisfying all the properties in Lemma 1.11. The Hermitian metric $\langle\cdot|\cdot\rangle$ on $\mathbb{C}TX$ induces by duality a Hermitian metric on $\mathbb{C}T^*X$ and also on the bundles of (0,q)-forms $\Lambda^qT^{*0,1}X,q=0,1\cdots,n-1$. We will denote all these induced metrics by $\langle\cdot|\cdot\rangle$. For every $v\in\Lambda^qT^{*0,1}X$, we write $|v|^2:=\langle v|v\rangle$. We have the pointwise orthogonal decompositions:

$$\mathbb{C}T^*X = T^{*1,0}X \oplus T^{*0,1}X \oplus \{\lambda\omega_0 : \lambda \in \mathbb{C}\},$$

$$\mathbb{C}TX = T^{1,0}X \oplus T^{0,1}X \oplus \{\lambda T : \lambda \in \mathbb{C}\}.$$

For $p \in X$, locally there is an orthonormal frame $\{U_1, \ldots, U_{n-1}\}$ of $T^{1,0}X$ such that the Levi-form \mathcal{L}_p is diagonal with respect to this frame. That is, $\mathcal{L}_p(U_i, \overline{U_j}) = \lambda_j \delta_{ij}$, where $\delta_{ij} = 1$ if i = j, $\delta_{ij} = 0$ if $i \neq j$. The entries $\{\lambda_1, \ldots, \lambda_{n-1}\}$ are called the eigenvalues of the Levi-form at p with respect to the rigid Hermitian metric $\langle \cdot | \cdot \rangle$. Moreover, the determinant of \mathcal{L}_p is defined by $\det \mathcal{L}_p = \lambda_1(p) \cdots \lambda_{n-1}(p)$.

1.4. **Canonical local coordinates.** In this work, we need the following result due to Baouendi-Rothschild-Treves, (see [1]).

Theorem 1.12. Let X be a compact CR manifold of $\dim X = 2n-1, n \geq 2$ with a transversal CR S^1 -action. Let $\langle \cdot | \cdot \rangle$ be a rigid Hermitian metric on X as in Lemma 1.11. For $x_0 \in X$, there exists a local patch D and local coordinates $(x_1, \dots, x_{2n-1}) = (z, \theta) = (z_1, \dots, z_{n-1}, \theta), z_j = x_{2j-1} + ix_{2j}, 1 \leq j \leq n-1, x_{2n-1} = \theta$, centered at x_0 . In terms of these coordinates, $D = \{(z, \theta) \in \mathbb{C}^{n-1} \times \mathbb{R} : |z| < \varepsilon, |\theta| < \delta\}$ and on D

(1.11)
$$T = \frac{\partial}{\partial \theta}$$

$$Z_{j} = \frac{\partial}{\partial z_{j}} + i \frac{\partial \varphi(z)}{\partial z_{j}} \frac{\partial}{\partial \theta}, j = 1, \dots, n - 1,$$

where $\{Z_j(x)\}_{j=1}^{n-1}$ form a basis of $T_x^{1,0}X$, for each $x \in D$ and $\varphi(z) \in C^\infty(D,\mathbb{R})$ is independent of θ . Moreover, on D, $\varphi(z) = \sum_{j=1}^{n-1} \lambda_j |z_j|^2 + O(|z|^3), \forall (z,\theta) \in D$, where $\{\lambda_j\}_{j=1}^{n-1}$ are the eigenvalues of the Levi-form of X at x_0 with respect to the given rigid

Hermitian metric on X. We call D a canonical local patch and (z, θ, φ) canonical coordinates centered at x_0 .

In order to make use of the S^1 -action locally, we need coordinates which contains nearly whole orbits. On the regular part, this is possible without any restrictions while on the irregular part we just find such coordinates approximately. This issue is stated in the following two lemmas.

Lemma 1.13 ([23], Lemma 1.17). Fix $x_0 \in X_{reg}$. Then we can find canonical coordinates (z, θ, φ) centered at x_0 and defined on a canonical local patch $D = \{(z, \theta) : |z| < \varepsilon_0, |\theta| < \pi\}$.

Lemma 1.14 ([23], Lemma 1.18). Let $x_0 \in X_k$ be a point, where $k \in \mathbb{N}$, k > 1. For every $\epsilon > 0$, ϵ small, we can find canonical coordinates (z, θ, φ) centered at x_0 and defined on a canonical local patch $D_{\epsilon} = \{(z, \theta) : |z| < \varepsilon_0, |\theta| < \frac{\pi}{k} - \epsilon\}$.

Lemma 1.15 ([23], Lemma 1.19). Fix $x_0 \in X$. Let (z, θ, φ) be canonical coordinates centered at x_0 and defined on a canonical chart $D = \tilde{D} \times (-\delta, \delta)$. We denote by dv_X the volume form associated with the rigid Hermitian metric. Then on D one has $dv_X = \lambda(z)dv(z)d\theta$ with $\lambda(z) \in C^{\infty}(\tilde{D}, \mathbb{R})$ which does not depend on θ and $dv(z) = 2^{n-1}dx_1 \cdots dx_{2n-2}$.

2. Szegő kernel expansion

2.1. **Some standard notations.** First, we introduce some standard notations and definitions. We shall use the following notations: $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. An element $\alpha = (\alpha_1, \cdots, \alpha_n) \in \mathbb{N}_0^n$ will be called a multiindex and the size of α is $|\alpha| = \alpha_1 + \cdots + \alpha_n$. We write $x^{\alpha} = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, $x = (x_1, \cdots, x_n)$, $\partial_x^{\alpha} = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_n}^{\alpha_n}$, $\partial_x^{\alpha} = \frac{\partial^{|\alpha|}}{\partial x^{\alpha}}$. Let $z = (z_1, \cdots, z_n)$, $z_j = x_{2j-1} + ix_{2j}$, $j = 1, \cdots, n$ be the coordinates of \mathbb{C}^n . We write $z^{\alpha} = z_1^{\alpha_1} \cdots z_n^{\alpha_n}$, $\overline{z}^{\alpha} = \overline{z}_1^{\alpha_1} \cdots \overline{z}_n^{\alpha_n}$, $\frac{\partial^{|\alpha|}}{\partial z^{\alpha}} = \partial_{z_1}^{\alpha_1} \cdots \partial_{z_n}^{\alpha_n}$, $\frac{\partial^{|\alpha|}}{\partial \overline{z}^{\alpha}} = \partial_{\overline{z}_1}^{\alpha_1} \cdots \partial_{\overline{z}_n}^{\alpha_n}$. In this section, we will study the semi-classical asymptotic expansion of the

In this section, we will study the semi-classical asymptotic expansion of the Szegő kernel of positive Fourier components. We recall some notations in semi-classical analysis.

Definition 2.1. Let W be an open subset of \mathbb{R}^N . Let S(1;W)=S(1) be the set of $a\in C^\infty(W)$ such that for every $\alpha\in\mathbb{N}_0^N$, there exists constant C_α such that $|\partial_x^\alpha a(x)|\leq C_\alpha$ on W. If a=a(x,k) depends on $k\in(1,\infty)$, we say that $a(x,k)\in S_{\mathrm{loc}}(1;W)=S_{\mathrm{loc}}(1)$ if $\chi(x)a(x,k)$ is uniformly bounded in S(1) when k varies in $(1,\infty)$ for every $\chi(x)\in C_0^\infty(W)$. For $m\in\mathbb{R}$, we put $S_{\mathrm{loc}}^m(1;W)=S_{\mathrm{loc}}^m(1)=k^mS_{\mathrm{loc}}(1)$. If $a\in S_{\mathrm{loc}}^{m_0}(1)$, $a_j\in S_{\mathrm{loc}}^{m_j}(1)$, $m_j\searrow -\infty$, we say that $a\sim \sum_{j=0}^\infty a_j$ in $S_{\mathrm{loc}}^{m_0}(1)$ if $a-\sum_{j=0}^{N_0}a_j\in S_{\mathrm{loc}}^{m_{N_0+1}}(1)$ for every N_0 .

Let W_1, W_2 be two open subsets of \mathbb{R}^N . If $A: C_0^\infty(W_1) \to \mathcal{D}'(W_2)$ is continuous, by the Schwartz kernel theorem (Theorem 5.2.1 in [17]) we write $K_A(x,y)$ or A(x,y) to denote the distribution kernel of A. The following two statements are equivalent

- (a) A can be extended to an continuous operator : $\mathcal{E}'(W_1) \to C^{\infty}(W_2)$,
- (b) $A(x,y) \in C^{\infty}(W_1 \times W_2)$.

If A satisfies (a) or (b), we say that A is smoothing.

A k-dependent continuous operator $A_k: C_0^\infty(W_1) \to \mathcal{D}'(W_2)$ is called k-negligible if A_k is smoothing and the kernel $A_k(x,y)$ of A_k satisfies $|\partial_x^\alpha \partial_y^\beta A_k(x,y)| = O(k^{-m})$ locally uniformly on every compact set in $W_1 \times W_2$, for all multi-indices $\alpha, \beta \in \mathbb{N}_0^N$ and all $m \in \mathbb{N}_0$. Let $C_k: C_0^\infty(W_1) \to \mathcal{D}'(W_2)$ be another k-dependent continuous operator. We write $A_k \equiv C_k \mod O(k^{-\infty})$ or $A_k(x,y) \equiv C_k(x,y) \mod O(k^{-\infty})$ if $A_k - C_k$ is k-negligible. We write $A_k = C_k + O(k^{-\infty})$ if $A_k \equiv C_k \mod O(k^{-\infty})$. Similarly, we write $B_k(x) \equiv 0 \mod O(k^{-\infty})$ for any k-dependent smooth function $B_k(x) \in C^\infty(W)$ if $|\partial_x^\alpha B_k(x)| = O(k^{-m})$ locally uniformly on every compact subset of W for all $\alpha \in \mathbb{N}_0^N$ and all $m \in \mathbb{N}_0$.

2.2. Asymptotic Szegő kernel expansion. Let $(\cdot|\cdot)$ be the inner product on $\Omega^{0,0}(X)$ induced by dv_X . Let $L^2(X)$ (resp. $L^2_m(X)$) be the completions of $\Omega^{0,0}(X)$ (resp. $\Omega^{0,0}_m(X)$) with respect to $(\cdot|\cdot)$. By elementary Fourier analysis one has $L^2_m(X) \perp L^2_{m'}(X)$ for $m \neq m', m, m' \in \mathbb{Z}$. For $m \in \mathbb{Z}$, let $Q_m : L^2(X) \to L^2_m(X)$ be the orthogonal projection with respect to $(\cdot|\cdot)$.

From now on we assume $m \in \mathbb{N}$. Let $S_m: L^2(X) \to H^0_{b,m}(X)$ be the orthogonal projection with respect to $(\cdot | \cdot)$. We call S_m the m-th Szegő projection. From (1.10), one finds $\dim H^0_{b,m}(X) < \infty$. Let $\{f_j\}_{j=1}^{d_m}$ be an orthonormal basis of $H^0_{b,m}(X)$. Then the m-th Szegő kernel function is given by $S_m(x) = \sum_{j=1}^{d_m} |f_j(x)|^2$. Let $S_m(x,y)$ be the distribution kernel with respect to the operator S_m which is given by $S_m(x,y) = \sum_{j=1}^{d_m} f_j(x) \overline{f_j(y)}$. The goal of this section is to study the semiclassical asymptotic expansion of $S_m(x,y)$.

We extend $\overline{\partial}_b$ to $L^2(X)$ in the sense of distribution and denote its kernel by $\operatorname{Ker}(\overline{\partial}_b) = \{u \in L^2(X) : \overline{\partial}_b u = 0\}$ which is a closed subspace of $L^2(X)$. Let $S: L^2(X) \to \operatorname{Ker}(\overline{\partial}_b)$ be the usual Szegő projection. We denote by S(x,y) the distribution kernel of the Szegő projection.

Lemma 2.2. With the notations above, we have

(2.1)
$$H_{b,m}^0(X) = \operatorname{Ker}(\overline{\partial}_b) \cap L_m^2(X)$$

and

(2.2)
$$S_m u = SQ_m u = Q_m Su, \ \forall u \in C^{\infty}(X).$$

Proof. It is obvious that $H_{b,m}^0(X) \subset \operatorname{Ker}(\overline{\partial}_b) \cap L_m^2(X)$. The converse is a direct corollary from following subelliptic estimate (see theorem 1.12 in [22])

(2.3)
$$||u||_s \le C_{s,m}(||\overline{\partial}_b u||_{s-1} + ||u||), \forall u \in H^s(X) \cap L^2_m(X), s \ge 1,$$

where $H^s(X)$ is the usual Sobolev space on X, $||u||_s$ is the usual Sobolev norm of order s and $C_{s,m}$ is a constant.

For any $u \in C^{\infty}(X)$, write $u = u_1 + u_2$, $u_1 \in H^0_{b,m}(X)$, $u_2 \in H^0_{b,m}(X)^{\perp}$. For any $v \in H^0_{b,m}(X)$, we have

$$(S_m u|v) = (u_1|v) = (u|v) = (Q_m u|v) = (SQ_m u|v).$$

For any $v \in L^2(X) \cap H^0_{b,m}(X)^{\perp}$, we have

$$(S_m u|v) = 0 = (SQ_m u|v)$$

since $S_m u, SQ_m u \in H^0_{b,m}(X)$. This implies $S_m u = SQ_m u$ for all $u \in C^\infty(X)$. Similarly, we have $S_m u = Q_m Su$ for all $u \in C^\infty(X)$.

Fix $x_0 \in X$. Let (z, θ, φ) be canonical coordinates centered at x_0 and defined on a canonical local patch $D_1 = \{(z, \theta) : |z| < \varepsilon_1, |\theta| < \delta_1\}$. Choose $D = \{(z, \theta) : |z| < \varepsilon, |\theta| < \delta\} \subseteq D_1$.

Choose two cut-off functions $\chi, \chi_1 \in C_0^{\infty}(D_1)$ such that $\chi = 1$ in some small neighborhood of \overline{D} and $\chi_1 = 1$ in some small neighborhood of $\sup \chi$. By Lemma 2.2, we have $S_m = SQ_m$ and hence

$$\chi S_m = \chi S Q_m = \chi S \chi_1 Q_m + \chi S (1 - \chi_1) Q_m.$$

We write $F=\chi S(1-\chi_1)$ and $F_m=\chi S(1-\chi_1)Q_m$ and denote by F(x,y) and $F_m(x,y)$ the distribution kernels of F and F_m , respectively. We will show

Lemma 2.3. $F_m: C_0^{\infty}(D) \to \mathcal{E}'(D_1)$ is m-negligible.

Proof. Since supp $\chi \cap \text{supp } (1-\chi_1)=\emptyset$, by a result of Boutet de Monvel-Sjöstrand [9] (see also [18] and [26]) we know that F is smoothing. Let $\cup_{j=1}^{n_0} U_j$ be a finite covering of X. We assume that all the $U_j s, 1 \leq j \leq n_0$ are canonical local patches. Choose a partition of unity $\{\rho_j\}_{j=1}^{n_0}$ with $\text{supp} \rho_j \in U_j, 1 \leq j \leq n_0$, and $\sum_{j=1}^{n_0} \rho_j = 1$ on X. Then for all $u \in C_0^\infty(D)$ we have

(2.4)
$$F_m u = FQ_m u = F\left(\sum_{j=1}^{n_0} \rho_j Q_m u\right) = \sum_{j=1}^{n_0} F(\rho_j Q_m u).$$

For $1 \leq j \leq n_0$, let $y = (w, y_{2n-1})$ be canonical coordinates in U_j . Then on U_j one finds

$$\rho_j Q_m u = \rho_j(y)(Q_m u)(y) = \rho_j(y)\hat{u}_m(w)e^{imy_{2n-1}}.$$

Set $F_j(x,y) = F(x,y)\rho_j(y)$ for $x \in D, y \in U_j$. Then on D, by a direct calculation we have

(2.5)

$$F(\rho_j Q_m u)(x) = -\frac{1}{2\pi mi} \int_{U_j} \left(\int_0^{2\pi} \frac{\partial F_j}{\partial y_{2n-1}} (x, e^{i\theta} \circ y) e^{im\theta} d\theta \right) u(y) \lambda(w) dv(w) dy_{2n-1}.$$

By (2.4), (2.5) and the induction method, we have $F_m(x,y) = O(m^{-N})$ locally uniformly for all $N \in \mathbb{N}$ and similarly for the derivatives. Thus, the lemma follows.

Set $G = \chi S \chi_1$ and $G_m = \chi S \chi_1 Q_m$. Write $D_1 = \tilde{D}_1 \times (-\delta_1, \delta_1)$ and $D = \tilde{D} \times (-\delta, \delta)$ with $\tilde{D}_1 = \{z \in \mathbb{C}^{n-1} : |z| < \varepsilon_1\}$ and $\tilde{D} = \{z \in \mathbb{C}^{n-1} : |z| < \varepsilon\}$. Assume that on D_1 , $\chi_1(y) = \tilde{\chi}_1(w)\tilde{\chi}_2(y_{2n-1})$ holds with $\tilde{\chi}_1(w) \in \underline{C_0^{\infty}}(\tilde{D}_1), \tilde{\chi}_2(y_{2n-1}) \in C_0^{\infty}(-\delta_1, \delta_1)$ and $\tilde{\chi}_1(w) = 1$ in some small neighborhood of \tilde{D} and $\tilde{\chi}_2 = 1$ in some small neighborhood of $[-\delta, \delta]$. Let $u \in C_0^{\infty}(D)$ be a smooth function. On D_1 , we write $(Q_m u)(y) = \hat{u}_m(w)e^{imy_{2n-1}}, \hat{u}_m(w) \in C^{\infty}(\tilde{D}_1)$. Then on D we have (2.6)

$$G_m u(x) = \int_{\tilde{D}_1} \tilde{\chi}_1(w) \hat{u}_m(w) \lambda(w) \left(\int_{-\delta_1}^{\delta_1} \chi(x) S(x, w, y_{2n-1}) \tilde{\chi}_2(y_{2n-1}) e^{imy_{2n-1}} dy_{2n-1} \right) dv(w).$$

In order to calculate the integral with respect to dy_{2n-1} in (2.6), we need the following result due to Boutet de Monvel and Sjöstrand [9], [18] and Hsiao-Marinescu [26].

Theorem 2.4. Let X be a compact strongly pseudoconvex CR manifold with a transversal CR S^1 -action. For any $x_0 \in X$, let D_1 be the canonical local patch defined as in Theorem 1.12 with canonical coordinates (z, θ, φ) centered at x_0 . Then on $D_1 \times D_1$ the distribution kernel S(x, y) of the Szegő projection $S: L^2(X) \to Ker(\overline{\partial}_b)$ satisfies

(2.7)
$$S(x,y) = \int_0^\infty e^{i\Psi(x,y)t} b(x,y,t) dt$$

in the sense of oscillatory integrals, where

(2.8)

$$\Psi(x,y) \in C^{\infty}(D_1 \times D_1), \Psi(x,y) = x_{2n-1} - y_{2n-1} + \Phi(z,w),
\Phi(z,w) = -\overline{\Phi}(w,z), \exists c > 0 : \operatorname{Im}\Phi \ge c|z-w|^2, \Phi(z,w) = 0 \Leftrightarrow z = w,
\Phi(z,w) = i(\varphi(z) + \varphi(w)) - 2i \sum_{|\alpha|+|\beta| \le N} \frac{\partial^{|\alpha|+|\beta|}\varphi}{\partial z^{\alpha}\partial \overline{z}^{\beta}}(0) \frac{z^{\alpha}}{\alpha!} \frac{\overline{w}^{\beta}}{\beta!} + O(|(z,w)|^{N+1}), \forall N \in \mathbb{N}_0,
h(x,y,t) \ge \sum_{\alpha=0}^{\infty} h_{\alpha}(x,y)t^{n-1-k} \text{ in } S^{n-1}(1;D_{\alpha} \times D_{\alpha})$$

$$b(x, y, t) \sim \sum_{k=0}^{\infty} b_k(x, y) t^{n-1-k} \text{ in } S_{\text{loc}}^{n-1}(1; D_1 \times D_1),$$

$$b_j(x,y) \in C^{\infty}(D_1 \times D_1), j = 0, 1, \cdots,$$

$$b_0(x,x) = \frac{1}{2}\pi^{-n} \left| \det \mathcal{L}_x \right|, \quad \forall x \in D_1.$$

By Theorem 2.4, the integral with respect to dy_{2n-1} in (2.6) can be computed by making use of the stationary phase formula due to Melin-Sjörstrand [30]. Substituting (2.7) and (2.8) to (2.6) and changing coordinates $t = m\sigma$, where $m \in \mathbb{N}$ and $\sigma \in \mathbb{R}_+$, we have

(2.9)

$$\int_{-\delta_{1}}^{\delta_{1}} \chi(x) S(x, w, y_{2n-1}) \tilde{\chi}_{2}(y_{2n-1}) e^{imy_{2n-1}} dy_{2n-1}$$

$$= m \int_{-\delta_{1}}^{\delta_{1}} \int_{0}^{\infty} e^{im[(x_{2n-1} - y_{2n-1})\sigma + \Phi(z, w)\sigma + y_{2n-1}]} \chi(x) b(x, y, m\sigma) \tilde{\chi}_{2}(y_{2n-1}) d\sigma dy_{2n-1}.$$

Set

$$\tilde{\Psi}(x, w, y_{2n-1}, \sigma) = (x_{2n-1} - y_{2n-1})\sigma + \Phi(z, w)\sigma + y_{2n-1}.$$

Then one finds

$$\frac{\partial \tilde{\Psi}}{\partial \sigma} = x_{2n-1} - y_{2n-1} + \Phi(z, w), \frac{\partial \tilde{\Psi}}{\partial y_{2n-1}} = -\sigma + 1.$$

For any fixed (x,w) the critical point of $\tilde{\Psi}$ is denoted by $x_c=(y_{2n-1},\sigma)=(x_{2n-1}+\Phi(z,w),1)$ which is the solution of the equations $\frac{\partial \tilde{\Psi}}{\partial \sigma}=0, \frac{\partial \tilde{\Psi}}{\partial y_{2n-1}}=0$. Moreover, the Hessian of $\tilde{\Psi}$ with respect to variables (y_{2n-1},σ) at the critical point x_c is

$$\begin{pmatrix}
\frac{\partial^2 \tilde{\Psi}}{\partial \sigma \partial \sigma} & \frac{\partial^2 \tilde{\Psi}}{\partial \sigma \partial y_{2n-1}} \\
\frac{\partial^2 \tilde{\Psi}}{\partial y_{2n-1} \partial \sigma} & \frac{\partial^2 \tilde{\Psi}}{\partial y_{2n-1} \partial y_{2n-1}}
\end{pmatrix}\Big|_{x_c} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

which implies that $\tilde{\Psi}(x, w, y_{2n-1}, \sigma)$ is a non-degenerate complex valued phase function for any fixed (x, w) in the sense of Melin and Sjöstrand [30]. Hence,

one can apply the stationary phase formula of Melin and Sjöstrand [30] to carry out the $d\sigma dy_{2n-1}$ integration in (2.9):

$$(2.10) \qquad m \int_{-\delta_{1}}^{\delta_{1}} \int_{0}^{\infty} e^{im\tilde{\Psi}(x,w,y_{2n-1},\sigma)} \chi(x)b(x,y,m\sigma)\tilde{\chi}_{2}(y_{2n-1})d\sigma dy_{2n-1}$$

$$= m \int_{-\delta_{1}}^{\delta_{1}} \int e^{im\tilde{\Psi}} \tau(\sigma)\chi(x)b(x,y,m\sigma)\tilde{\chi}_{2}(y_{2n-1})d\sigma dy_{2n-1}$$

$$+ m \int_{-\delta_{1}}^{\delta_{1}} \int e^{im\tilde{\Psi}} (1-\tau(\sigma))\chi(x)b(x,y,m\sigma)\tilde{\chi}_{2}(y_{2n-1})d\sigma dy_{2n-1},$$

where $\tau(\sigma) \in C_0^{\infty}(\mathbb{R})$ with $\operatorname{supp} \tau \in (\frac{1}{2}, \frac{3}{2})$ and $\tau = 1$ near $\sigma = 1$.

First we show that on $D_1 \times \tilde{D}_1$, the second term on the right-hand side of (2.10) satisfies the following identity

(2.11)
$$m \int_{-\delta_1}^{\delta_1} \int e^{im\tilde{\Psi}(x,w,y_{2n-1},\sigma)} (1-\tau(\sigma))\chi(x)b(x,y,m\sigma)\tilde{\chi}_2(y_{2n-1})d\sigma dy_{2n-1} \equiv 0 \mod O(m^{-\infty}).$$

This is a direct corollary of the following formula

$$e^{im\tilde{\Psi}} = \frac{1}{im(1-\sigma)} \frac{\partial}{\partial y_{2n-1}} e^{im\tilde{\Psi}}$$

and the integration by parts with respect to the variable y_{2n-1} . For convenience we denote by $H_m(x, w)$ the left-hand side of (2.11).

Making use of Melin-Sjöstrand's stationary phase formula [30], the first term on the right-hand side of (2.10) becomes

(2.12)
$$m \int_{-\delta_1}^{\delta_1} \int e^{im\tilde{\Psi}} \tau(\sigma) \chi(x) b(x, y, m\sigma) \tilde{\chi}_2(y_{2n-1}) d\sigma dy_{2n-1}$$
$$\equiv e^{im(x_{2n-1} + \Phi(z, w))} \chi(x) \hat{b}(x, w, m) \mod O(m^{-\infty}),$$

where

(2.13)
$$\hat{b}(x, w, m) \sim \sum_{j=0}^{\infty} \hat{b}_j(x, w) m^{n-1-j} \text{ in } S_{\text{loc}}^{n-1}(1; D_1 \times \tilde{D}_1),$$
$$\hat{b}_j(x, w) \in C^{\infty}(D_1 \times \tilde{D}_1), j = 0, 1, 2, \cdots.$$

In particular, one has

(2.14)
$$\hat{b}_0(x,w) = (2\pi)\tilde{b}_0(x,w,x_{2n-1} + \Phi(z,w)), \ \hat{b}_0(x,z) = \pi^{1-n} \left| \det \mathcal{L}_x \right|,$$

where \tilde{b}_0 denotes an almost analytic extension of b_0 , that is $\tilde{b}_0(\widetilde{x},\widetilde{y}) \in C^\infty(U_1 \times U_1)$ with $\tilde{b}_0|_{D_1 \times D_1} = b_0$ and $\left|\overline{\partial}_{\widetilde{x}} \tilde{b}_0(\widetilde{x},\widetilde{y})\right| + \left|\overline{\partial}_{\widetilde{y}} \tilde{b}_0(\widetilde{x},\widetilde{y})\right| \leq C_N(|\operatorname{Im} \widetilde{x}|^N + |\operatorname{Im} \widetilde{y}|^N)$, for every N > 0 where $C_N > 0$ is a constant. Here U_1 is an open set in \mathbb{C}^{2n-1} with $U_1 \cap \mathbb{R}^{2n-1} = D_1$ (we identify D_1 with an open set in \mathbb{R}^{2n-1}) and \widetilde{x} , \widetilde{y} are complex coordinates of \mathbb{C}^{2n-1} . Substituting (2.11) and (2.12) to (2.6) one has

(2.15)
$$G_{m}u = \int_{\tilde{D}_{1}} \tilde{\chi}_{1}(w)\hat{u}_{m}(w)e^{im(x_{2n-1}+\Phi(z,w))}\chi(x)\hat{b}(x,w,m)\lambda(w)dv(w) + \int_{\tilde{D}_{1}} \tilde{\chi}_{1}(w)\hat{u}_{m}(w)H_{m}(x,w)\lambda(w)dv(w)$$

with $H_m(x, w) \equiv 0 \mod O(m^{-\infty})$ on $D_1 \times \tilde{D}_1$.

Choose $\eta(y_{2n-1}) \in C_0^{\infty}(-\delta_1, \delta_1)$ such that $\int_{-\delta_1}^{\delta_1} \eta(y_{2n-1}) dy_{2n-1} = 1$. Then the first term on the right-hand side of (2.15) is equal to

(2.16)

$$\int_{D_1} (Q_m u)(y) \tilde{\chi}_1(w) \eta(y_{2n-1}) e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \chi(x) \hat{b}(x,w,m) \lambda(w) dw dy_{2n-1}$$

$$= \chi(x) \int_{D_1} (Q_{-m} B_m)(x,y) u(y) \lambda(w) dy = \chi(x) \int_{D} (Q_{-m} B_m)(x,y) u(y) \lambda(w) dy.$$

Here, we have set

(2.17)
$$B_m(x,y) = e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \hat{b}(x,w,m) \tilde{\chi}_1(w) \eta(y_{2n-1})$$

and $(Q_{-m}B_m)(x,y)$ denotes that Q_{-m} acts on $B_m(x,y)$ seen as a function in the variable y. Combining (2.15), (2.16), (2.17) and Lemma 2.3, we have

$$S_m(x,y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} B_m(x,e^{i\theta} \circ y) e^{im\theta} d\theta + A_m(x,y), \forall x,y \in D \times D,$$

where $A_m(x,y) \equiv 0 \mod O(m^{-\infty})$. On the other hand, we have $S_m(x,y) = \sum_{j=1}^{d_m} f_j(x) \overline{f_j(y)}$. On D, we can write $f_j(x) = \hat{f_j}(z) e^{imx_{2n-1}}$ which leads to $S_m(x,y) = \sum_{j=1}^{d_m} \hat{f_j}(z) \overline{\hat{f_j}(w)} e^{im(x_{2n-1}-y_{2n-1})}$. Thus,

(2.18)
$$e^{-imx_{2n-1}}S_m(x,y) = \sum_{j=1}^{d_m} \hat{f}_j(z)\overline{\hat{f}_j(w)}e^{im(-y_{2n-1})}$$

does not depend on x_{2n-1} . We get

(2.19)

$$e^{-imx_{2n-1}}S_m(x,y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-imx_{2n-1}}B_m(x,e^{i\theta} \circ y)e^{im\theta}d\theta + e^{-imx_{2n-1}}A_m(x,y).$$

Choose $\chi_0(x_{2n-1}) \in C_0^{\infty}(-\delta, \delta)$ such that $\int_{-\delta}^{\delta} \chi_0(x_{2n-1}) dx_{2n-1} = 1$. From (2.18) and (2.19) we have

(2.20)

$$e^{-imx_{2n-1}}S_m(x,y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\delta}^{\delta} \chi_0(x_{2n-1})e^{-imx_{2n-1}}B_m(x,e^{i\theta} \circ y)e^{im\theta}dx_{2n-1}d\theta + C_m(z,y),$$

where $C_m(z,y) = \int_{-\delta}^{\delta} A_m(x,y) e^{-imx_{2n-1}} \chi_0(x_{2n-1}) dx_{2n-1}$, $C_m(z,y) \equiv 0 \mod O(m^{-\infty})$. Set

(2.21)
$$\hat{S}_m(x,y) = e^{imx_{2n-1}} \int_{\delta_n}^{\delta_1} \chi_0(x_{2n-1}) e^{-imx_{2n-1}} B_m(x,y) dx_{2n-1}.$$

From (2.17),(2.19), (2.20) and (2.21) we have

Theorem 2.5. Let X be as in Theorem 2.4. Consider the orthogonal projection S_m : $L^2(X) \to H^0_{b,m}(X)$. We denote by $S_m(x,y)$ the distribution kernel of S_m . For $x_0 \in X$, let (z,θ,φ) be canonical coordinates centered at x_0 and defined on a canonical local

patch $D_1 = \{(z, \theta) : |z| < \varepsilon_1, |\theta| < \delta_1\}$. For any $D = \{(z, \theta) : |z| < \varepsilon, |\theta| < \delta\} \subseteq D_1$, we have

$$S_m(x,y) \equiv \frac{1}{2\pi} \int_{-\pi}^{\pi} \hat{S}_m(x, e^{i\theta} \circ y) e^{im\theta} d\theta \mod O(m^{-\infty})$$

on $D \times D$, where

(2.22)

$$\hat{S}_{m}(x,y) = e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \hat{b}(z,w,m) \tilde{\chi}_{1}(w) \eta(y_{2n-1}),$$

$$\Phi(z,w) = i(\varphi(z) + \varphi(w)) - 2i \sum_{|\alpha| + |\beta| \le N} \frac{\partial^{|\alpha| + |\beta|} \varphi}{\partial z^{\alpha} \partial \overline{z}^{\beta}}(0) \frac{z^{\alpha}}{\alpha!} \frac{\overline{w}^{\beta}}{\beta!} + O(|(z,w)|^{N+1}),$$

$$\hat{b}(z,w,m) \sim \sum_{k=0}^{\infty} m^{n-1-k} \hat{b}_{k}(z,w) \text{ in } S_{\text{loc}}^{n-1}(1; \tilde{D} \times \tilde{D}), \tilde{D} = \{z \in \mathbb{C}^{n-1} : |z| < \varepsilon\},$$

$$\hat{b}_{0}(z,w) = (2\pi) \int_{-\delta}^{\delta} \tilde{b}_{0}(z,x_{2n-1},w,x_{2n-1} + \Phi(z,w)) \chi_{0}(x_{2n-1}) dx_{2n-1},$$

$$\hat{b}_{0}(z,z) = \pi^{-(n-1)} |\det \mathcal{L}_{x}|, \quad x = (z,0), \quad \forall z \in \tilde{D},$$

and

$$\begin{split} \hat{b}_{j}(z,w) &\in C^{\infty}(\tilde{D} \times \tilde{D}), \forall j; \chi_{0}(x_{2n-1}) \in C^{\infty}_{0}(-\delta,\delta), \int_{-\delta}^{\delta} \chi_{0}(x_{2n-1}) dx_{2n-1} = 1; \\ \chi_{1}(w) &\in C^{\infty}_{0}(\tilde{D}_{1}), \quad \chi_{1} = 1 \text{ in a neighborhood of } \overline{\tilde{D}}; \\ \eta(y_{2n-1}) &\in C^{\infty}_{0}(-\delta_{1},\delta_{1}), \int_{-\delta_{1}}^{\delta_{1}} \eta(y_{2n-1}) dy_{2n-1} = 1. \end{split}$$

Here, \tilde{b}_0 is as in (2.14).

2.3. Asymptotic Szegő kernel expansion on X_{reg} . Recall that X_{reg} is the regular part of X, that is $X_{\text{reg}} = \{x \in X : \forall e^{i\theta} \in S^1, \text{ if } e^{i\theta} \circ x = x, \text{ then } e^{i\theta} = \text{id}\}$, and that we assume $X_{\text{reg}} \neq \emptyset$. Fix $x_0 \in X_{\text{reg}}$. Let (z, θ, φ) be canonical coordinates centered at x_0 and defined on a canonical patch $D_1 = \{(z, \theta) : |z| < \varepsilon_1, |\theta| < \pi\}$. Set $D = \{(z, \theta) \in \mathbb{C}^{n-1} \times \mathbb{R} : |z| < \varepsilon, |\theta| < \frac{\pi}{2}\}$ with $\varepsilon < \varepsilon_1$. From Theorem 2.5 it follows that

(2.23)
$$S_m(x,y) \equiv e^{-imy_{2n-1}} \frac{1}{2\pi} \int_{-\pi}^{\pi} \hat{S}_m(x,(w,\theta)) e^{im\theta} d\theta \mod O(m^{-\infty})$$

holds on $D \times D$. Substituting (2.22) to (2.23), we have

(2.24)
$$S_m(x,y) \equiv \frac{1}{2\pi} e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \hat{b}(z,w,m) \mod O(m^{-\infty}),$$
$$S_m(x,x) \equiv \frac{1}{2\pi} \hat{b}(z,z,m) \mod O(m^{-\infty}).$$

Thus, from (2.24) we have

Theorem 2.6. Let X be as in Theorem 2.4. For $x_0 \in X_{reg}$, let (z, θ, φ) be canonical coordinates centered at x_0 and defined on a canonical patch $D_1 = \{(z, \theta) : |z| < 0\}$

 $\varepsilon_1, |\theta| < \pi$ }. Set $D = \{(z, \theta) \in \mathbb{C}^{n-1} \times \mathbb{R} : |z| < \varepsilon, |\theta| < \frac{\pi}{2}\} \in D_1$. Then on $D \times D$, we have

$$S_m(x,y) \equiv \frac{1}{2\pi} e^{im(x_{2n-1} - y_{2n-1} + \Phi(z,w))} \hat{b}(z,w,m) \mod O(m^{-\infty}),$$

where

(2.25)
$$\hat{b}(z, w, m) \sim \sum_{j=0}^{\infty} m^{n-1-j} \hat{b}_{j}(z, w) \text{ in } S_{\text{loc}}^{n-1}(1, \tilde{D} \times \tilde{D}),$$

$$\hat{b}_{j}(z, w) \in C^{\infty}(\tilde{D} \times \tilde{D}), \quad j = 0, 1, 2, \cdots,$$

$$\hat{b}_{0}(z, z) = \pi^{-(n-1)} |\det \mathcal{L}_{x}|, \quad x = (z, 0), \quad \forall z \in \tilde{D}.$$

Here, we set $\tilde{D} = \{z \in \mathbb{C}^{n-1} : |z| < \varepsilon\}$. In particular, we have

(2.26)
$$S_m(x,x) \equiv \frac{1}{2\pi} \hat{b}(z,z,m) \mod O(m^{-\infty}).$$

2.4. Asymptotic Szegő kernel expansion on the complement of X_{reg} . In this section, we try to get the asymptotic expansion of the Szegő kernel on the complement of X_{reg} . Fix $x_0 \in X_k$ for some k > 1, where X_k is defined in (1.3). Let (z,θ,φ) be canonical coordinates centered at x_0 and defined on a canonical chart $D_1 = \{(z,\theta): |z| < \varepsilon_1, |\theta| < \frac{\pi}{k} - \epsilon\}$. It is straightforward to see that there is a small neighborhood $D = \{(z,\theta): |z| < \varepsilon, |\theta| < \delta\} \in D_1$ of x_0 such that

(2.27)
$$e^{i\theta} \circ (0,0) \neq (z,\hat{\theta}), \ \forall \theta \in [0,2\pi), \ (z,\hat{\theta}) \in D, \ z \neq 0.$$

From Theorem 2.5, we have

(2.28)
$$S_{m}(x,x_{0}) \equiv \frac{1}{2\pi} \sum_{s=1}^{k} \int_{\frac{2\pi}{k}(s-1)}^{\frac{2\pi}{k}s} \hat{S}_{m}(x,e^{i\theta} \circ (0,0)) e^{im\theta} d\theta \mod O(m^{-\infty})$$

$$\equiv \frac{1}{2\pi} \sum_{s=1}^{k} e^{i\frac{2\pi}{k}(s-1)m} \int_{0}^{\frac{2\pi}{k}} \hat{S}_{m}(x,e^{i\theta} \circ (0,0)) e^{im\theta} d\theta \mod O(m^{-\infty})$$

for any $x \in D$. By a direct calculation, we find

(2.29)
$$\sum_{s=1}^{k} e^{i\frac{2\pi}{k}(s-1)m} = \begin{cases} k, & \text{if } k \mid m \\ 0, & \text{if } k \nmid m \end{cases} ;$$

From (2.27), we can check that

(2.30)
$$\frac{k}{2\pi} \int_{-\frac{\pi}{k}}^{\frac{\pi}{k}} \hat{S}_m(x, e^{i\theta} \circ (0, 0)) e^{im\theta} d\theta = \frac{k}{2\pi} \int_{-\frac{\pi}{k}}^{\frac{\pi}{k}} \hat{S}_m(x, (0, \theta)) e^{im\theta} d\theta$$

holds. Substituting (2.29) to (2.28) for $k \mid m$ and using (2.30), we have

(2.31)
$$S_m(x,x_0) \equiv \frac{k}{2\pi} \int_{-\frac{\pi}{k}}^{\frac{\pi}{k}} \hat{S}_m(x,(0,\theta)) e^{im\theta} d\theta \mod O(m^{-\infty}).$$

Substituting (2.22) to (2.31) yields

$$S_m(x, x_0) \equiv \frac{k}{2\pi} e^{im(x_{2n-1} + \Phi(z,0))} \hat{b}(z, 0, m) \mod O(m^{-\infty}).$$

Summing up, we obtain

Theorem 2.7. Let X be as in Theorem 2.4. Assume $x_0 \in X_k$, k > 1. Let D_1 , D and (z, θ, φ) be as above with $\frac{\pi}{k} - \epsilon$ replaced by δ_1 . For $k \nmid m$ we have $S_m(x, x_0) = 0$ for all $x \in D$. For $k \mid m$ we have

(2.32)
$$S_m(x,x_0) \equiv \frac{k}{2\pi} e^{im(x_{2n-1} + \Phi(z,0))} \hat{b}(z,0,m) \mod O(m^{-\infty})$$

on D. In particular, given $k \mid m$ and $x = x_0$, we have

$$S_m(x_0, x_0) = \frac{k}{2\pi} \hat{b}(0, 0, m) + O(m^{-\infty})$$

and

$$\hat{b}(0,0,m) \sim \hat{b}_0(0,0)m^{n-1} + \hat{b}_1(0,0)m^{n-2} + \cdots$$

in the sense that for any $N \in \mathbb{N}_0$ there exists $C_N > 0$ independent of m such that

$$\left| \hat{b}(0,0,m) - \sum_{j=0}^{N} \hat{b}_{j}(0,0)m^{n-1-j} \right| \le C_{N} m^{n-2-N}$$

holds for all $m \in \mathbb{N}$.

3. Equivariant embedding of CR manifolds

Let X be a compact strongly pseudoconvex CR manifolds with a locally free transversal CR S^1 -action. Now we use the Szegő kernel expansion we have established in Section 2 to get the equivariant embedding of X.

3.1. Immersion of CR manifold. We assume $X = X_1 \cup X_2 \cup \cdots \cup X_l$, $X_1 \neq \emptyset$, where X_k , $1 \leq k \leq l$, is defined in (1.3). For $1 \leq k \leq l$ let $\{f_j^k\}_{j=1}^{d_{km}}$ and $\{g_j^k\}_{j=1}^{d_{k(m+1)}}$ be orthonormal bases of $H_{b,km}^0(X)$ and $H_{b,k(m+1)}^0(X)$, respectively. Now for $1 \leq k \leq l$ we can define a CR map from X to Euclidean space as follows

$$\Phi_m^k: X \to \mathbb{C}^{d_{km}+d_{k(m+1)}}, x \mapsto (f_1^k(x), \cdots, f_{d_{km}}^k(x), g_1^k(x), \cdots, g_{d_{k(m+1)}}^k(x)).$$

Combining the $\Phi_m^k s, 1 \leq k \leq l$, we define a CR map

$$\Phi_m: X \to \mathbb{C}^{N_m}, x \to (\Phi_m^1(x), \cdots, \Phi_m^l(x)),$$

where $N_m = \sum_{k=1}^l (d_{km} + d_{k(m+1)})$. If the transversal CR S^1 -action on X is globally free, then $X = X_1 = X_{\text{reg}}$ and Epstein [14] showed that Φ^1_m is an CR embedding when m is large. However, if the transversal CR S^1 -action on X is just locally free the CR functions in $H^0_{b,m}(X) \bigoplus H^0_{b,m+1}(X)$ are not enough for the embedding. The reason is that the space $H^0_{b,m}(X) \bigoplus H^0_{b,m+1}(X)$ will be not enough to separate the points in $X \setminus X_{\text{reg}}$.

Now we use the asymptotic Szegő kernel expansion in Section 2 to establish

Lemma 3.1. The map $\Phi_m: X \to \mathbb{C}^{N_m}$ is an immersion when m is large.

Proof. For $x_0 \in X_k$, let (z,θ,φ) be canonical coordinates centered at x_0 and defined on a canonical local patch $D=\{(z,\theta): |z|<\varepsilon, |\theta|<\delta\}=\tilde{D}\times(-\delta,\delta)$. Assume that k|m. Let $\{f_j\}_{j=1}^{d_m}\subset H^0_{b,m}(X)$ be an orthonormal basis. Since $S_m(x,y)=$

 $\sum_{j=1}^{d_m} f_j(x) \overline{f_j(y)}$, we have that $\overline{S_m(x,y)} = S_m(y,x)$. Furthermore, for any $u \in C_0^\infty(D)$ we have $S_m u(x) = \int_D S_m(x,y) u(y) dv_X(y)$ and hence

(3.1)
$$\overline{S_m u} = \int_D \overline{S_m(x,y)u(y)} dv = \int_D S_m(y,x) \overline{u(y)} dv_X.$$

Choose cut-off functions $\chi \in C_0^\infty(\mathbb{C}^{n-1}), \chi_2 \in C_0^\infty(-\delta, \delta)$ such that $\mathrm{supp}\chi \Subset \{w \in \mathbb{C}^{n-1}: |w|<1\}$ and $\int_{-\delta}^\delta \chi_2(y_{n-1})dy_{2n-1}=1$. For $j=1,\cdots,n-1$, set

(3.2)
$$u_j(y) = w_j \chi \left(\frac{\sqrt{m}w}{\log m} \right) \chi_2(y_{2n-1}) e^{imy_{2n-1}} e^{im\text{Re}\Phi(w,0)},$$

where Φ is as in Theorem 2.5. Then $u_j \in C_0^{\infty}(X)$ has its support in D if m is sufficiently large. Define $v_j = S_m u_j, j = 1, \dots, n-1$. Then from Theorem 2.5 and (3.1) we have

$$\overline{S_m u_j(x)} = \frac{1}{2\pi} \int_D \int_{-\pi}^{\pi} \hat{S}_m(y, e^{i\theta} \circ x) e^{im\theta} d\theta \overline{u_j(y)} dv_X + \int_D \overline{R_m(x, y) u_j(y)} dv_X,$$

where $R_m(x,y) \equiv 0 \mod O(m^{-\infty})$. With respect to the canonical local coordinates, one notes that

(3.3)
$$\frac{\partial \hat{S}_m(y, e^{i\theta} \circ x)}{\partial \overline{z}_j} \Big|_{x=x_0} = \frac{\partial \hat{S}_m}{\partial \overline{z}_j} (y, e^{i\theta} \circ x_0).$$

Then by (3.3) and a direct calculation we have

(3.4)
$$\frac{\partial \overline{S_m u_j}}{\partial \overline{z_j}}(x_0) \equiv \frac{k}{2\pi} \int_D \int_{-\frac{\pi}{L}}^{\frac{\pi}{L}} \frac{\partial \hat{S}_m}{\partial \overline{z_j}}(y, (0, \theta)) e^{im\theta} d\theta \overline{u_j(y)} dv_X + O(m^{-\infty})$$

which leads to

Substituting (3.5) to (3.4), we have

$$\frac{\partial \overline{S_m u_j}}{\partial \overline{z_j}}(x_0) = \frac{k}{2\pi} \int_D e^{im(y_{2n-1} + \Phi(w,0))} \left[2m(\lambda_j w_j + O(|w|^2)) \hat{b}(w,0,m) + \frac{\partial \hat{b}(w,0,m)}{\partial \overline{z}_j} \right] \times \overline{u_j(y)} dv_X + O(m^{-\infty}).$$

Substituting (3.2) to (3.6) and taking the coordinate transformation $w \to \frac{w}{\sqrt{m}}$, we have

$$(3.7) \frac{\partial \overline{S_m u_j}}{\partial \overline{z_j}}(x_0) = \frac{k}{2\pi} \int_{|w| \le \log m} e^{-m \operatorname{Im} \Phi(\frac{w}{\sqrt{m}}, 0)} m^{-(n-1)} \times \left[2(\lambda_j |w_j|^2 + \frac{1}{\sqrt{m}} O(|w|^3)) \hat{b}(\frac{w}{\sqrt{m}}, 0, m) + \frac{1}{\sqrt{m}} \frac{\partial \hat{b}(\frac{w}{\sqrt{m}}, 0, m)}{\partial \overline{z}_j} \overline{w_j} \right] \times \chi(\frac{w}{\log m}) \lambda(\frac{w}{\sqrt{m}}) dv(w) + O(m^{-\infty}),$$

and hence

(3.8)
$$\lim_{m \to \infty} \frac{\partial \overline{S_m u_j}}{\partial \overline{z_j}}(x_0) = \frac{k}{2\pi} \int_{\mathbb{C}^{n-1}} e^{-\lambda |w|^2} 2\lambda_j |w_j|^2 \hat{b}_0(0,0) dv(w) = c_j \neq 0,$$

where $\lambda |w|^2 = \sum_{j=1}^{n-1} \lambda_j |w_j|^2$ and c_j is a non-zero real number.

For $j \neq k$, we can repeat the procedure above and get

(3.9)
$$\frac{\partial \overline{S}_{m} u_{j}}{\partial \overline{z}_{k}}(x_{0}) = \frac{k}{2\pi} \int_{|w| \leq \log m} e^{-m \operatorname{Im}\Phi(\frac{w}{\sqrt{m}}, 0)} m^{-(n-1)} \times \left[(2\lambda_{k} w_{k} \overline{w}_{j} + \frac{1}{\sqrt{m}} O(|w|^{3})) \hat{b}(\frac{w}{\sqrt{m}}, 0, m) + \frac{1}{\sqrt{m}} \frac{\partial \hat{b}(\frac{w}{\sqrt{m}}, 0, m)}{\partial \overline{z}_{k}} \overline{w}_{j} \right] \times \left[(\frac{w}{\log m}) \lambda(\frac{w}{\sqrt{m}}) dw + O(m^{-\infty}). \right]$$

Letting $m \to \infty$, we get

(3.10)
$$\lim_{m \to \infty} \frac{\partial \overline{S_m u_j}}{\partial \overline{z}_k}(x_0) = \frac{k}{2\pi} \int_{\mathbb{C}^{n-1}} e^{-\lambda |w|^2} 2\lambda_k w_k \overline{w}_j \hat{b}_0(0,0) dv(w) = 0.$$

Similarly, one computes

(3.11)
$$\frac{\partial S_m u_j}{\partial z_k}(x_0) = \frac{k}{2\pi} \int_{|w| \le \log m} e^{-m \operatorname{Im}\Phi(\frac{w}{\sqrt{m}},0)} m^{-(n-1)} \times \left[(2\lambda_k \overline{w}_k \overline{w}_j + \frac{1}{\sqrt{m}} O(|w|^3)) \hat{b}(\frac{w}{\sqrt{m}},0,m) + \frac{1}{\sqrt{m}} \frac{\partial \hat{b}(\frac{w}{\sqrt{m}},0,m)}{\partial z_k} \overline{w}_j \right] \times \chi(\frac{w}{\log m}) \lambda(\frac{w}{\sqrt{m}}) dw + O(m^{-\infty}).$$

Letting $m \to \infty$, we have

(3.12)
$$\lim_{m \to \infty} \frac{\partial \overline{S_m u_j}}{\partial z_k}(x_0) = \frac{k}{2\pi} \int_{\mathbb{C}^{n-1}} e^{-\lambda |w|^2} 2\lambda_k \overline{w}_k \overline{w}_j \hat{b}_0(0,0) dv(w) = 0.$$

Given j=n, choose $\chi_3(y_{2n-1})\in C_0^\infty(-\delta_1,\delta_1)$ satisfying $\int_{-\delta_1}^{\delta_1}y_{2n-1}\chi_3(y_{2n-1})=1$. Set

$$u_n = m y_{2n-1} \chi_3(m y_{2n-1}) e^{i m y_{2n-1}} \chi\left(\frac{\sqrt{m}w}{\log m}\right) e^{i m \operatorname{Re}\Phi(w,0)}, \quad v_n = S_m u_n.$$

Then by (3.3) and the same argument as in (2.28) we have

$$(3.13) \quad \frac{\partial \overline{S_m u_n}}{\partial x_{2n-1}}(x_0) = \frac{k}{2\pi} \int_D \int_{-\frac{\pi}{k}}^{\frac{\pi}{k}} \frac{\partial \hat{S}_m}{\partial x_{2n-1}}(y, e^{i\theta} \circ x_0) e^{im\theta} d\theta \overline{u_n(y)} dv_X + O(m^{-\infty}).$$

By a direct calculation, we have

(3.14)
$$\frac{\partial \hat{S}_m}{\partial x_{2n-1}}(y,0,\theta) = e^{im(y_{2n-1}-\theta+\Phi(w,0))}\hat{b}(w,0,m) \left[-im\eta(\theta) + \frac{\partial \eta(\theta)}{\partial \theta} \right].$$

Substituting (3.14) to (3.13) and using the fact that $\int_{-\frac{\pi}{h}}^{\frac{\pi}{h}} \frac{\partial \eta(\theta)}{\partial \theta} d\theta = 0$, we find

(3.15)
$$\frac{k}{2\pi} \int_{D} \int_{-\frac{\pi}{k}}^{\frac{\pi}{k}} \frac{\partial \hat{S}_{m}}{\partial x_{2n-1}} (y, e^{i\theta} \circ x_{0}) e^{im\theta} d\theta \overline{u_{n}(y)} dv_{X}$$

$$= \frac{-ik}{2\pi} \int_{|w| < \log m} m^{-(n-1)} \hat{b}(\frac{w}{\sqrt{m}}, 0, m) e^{-m\operatorname{Im}\Phi(\frac{w}{\sqrt{m}}, 0)} \lambda(w) dv(w).$$

Substituting (3.15) to (3.13) and letting $m \to \infty$, we have

(3.16)
$$\lim_{m \to \infty} \frac{\partial \overline{S_m u_n}(x_0)}{\partial x_{2n-1}} = \frac{-ik}{2\pi} \hat{b}_0(0,0) \int_{\mathbb{C}^{n-1}} e^{-\lambda |w|^2} dv(w) = ic_n \neq 0,$$

where c_n is a nonzero real number.

On the other hand, for $j = 1, \dots, n-1$ by a similar calculation we have

(3.17)
$$\frac{\partial \overline{S_m u_n}}{\partial \overline{z}_j}(x_0) = \frac{k}{2\pi} \int_{|w| \le \log m} e^{-m \operatorname{Im} \Phi(\frac{w}{\sqrt{m}}, 0)} \left[2(\lambda_j \frac{w_j}{\sqrt{m}} + \frac{1}{m} O(|w|^2)) \hat{b}(\frac{w}{\sqrt{m}}, 0, m) + \frac{1}{m} \frac{\partial \hat{b}}{\partial \overline{z}_j} (\frac{w}{\sqrt{m}}, 0, m) \right] \chi(\frac{w}{\log m}) \lambda(\frac{w}{\sqrt{m}}) m^{-(n-1)} dv(w).$$

By (3.17) we get

(3.18)
$$\left| \frac{\partial \overline{S_m u_n}}{\partial \overline{z}_i} (x_0) \right| \le C \frac{1}{\sqrt{m}},$$

where C is a constant which does not depend on x_0 and m. Similarly, we find

(3.19)
$$\left| \frac{\partial \overline{S_m u_n}}{\partial z_j} (x_0) \right| \le C \frac{1}{\sqrt{m}}.$$

Set $v_j=\alpha_{2j-1}+i\alpha_{2j},\ j=1,\cdots,n$. Then combining the above arguments there are positive constants c,C independent of x_0 and m and a sequence $\{\varepsilon_m\}$ which does not depend on $x_0\in X$ with $\varepsilon_m\to 0$ as $m\to\infty$ such that the following estimates hold

(3.20)
$$\left| \frac{\partial \alpha_{j}}{\partial x_{j}}(x_{0}) \right| \geq c; \left| \frac{\partial \alpha_{2n}}{\partial x_{2n-1}}(x_{0}) \right| \geq c, j = 1, \dots, 2n - 2,$$

$$\left| \frac{\partial \alpha_{j}}{\partial x_{k}}(x_{0}) \right| \leq \varepsilon_{m}, j \neq k, j, k = 1, \dots, 2n - 2,$$

$$\left| \frac{\partial \alpha_{2n}}{\partial x_{j}}(x_{0}) \right| \leq C \frac{1}{\sqrt{m}}, j = 1, \dots, 2n - 2.$$

From (3.20) the real Jacobian matrix of Φ_m is non-degenerate at any $x_0 \in X$ when m is large enough which implies that Φ_m is an immersion. Thus, we get the conclusion of the lemma.

3.2. Analysis near the complement of X_{reg} . In order to get the global embedding of CR manifolds by CR functions which lie in the positive Fourier components we need the following

Proposition 3.2. Fix $x_0 \in X \setminus X_{reg}$. Without loss of generality, we assume $x_0 \in X_{k_0}$ for some $k_0 > 1$. We have

- (1) There exist a positive integer m_0 and a neighborhood $U(x_0)$ of x_0 such that $\Phi_{m_0}^{k_0}: U(x_0) \to \mathbb{C}^{d_{k_0m_0}+d_{k_0(m_0+1)}}$ is an embedding and $S_{k_0m_0}(x,x_0) \neq 0$, $S_{k_0(m_0+1)}(x,x_0) \neq 0$, for all $x \in U(x_0)$.
- (2) There exist positive constants ε_0 , δ_0 and a neighborhood $V(x_0)$ of x_0 with $V(x_0) \subseteq U(x_0)$ such that

$$e^{i\theta} \circ V(x_0) \subset U(x_0), \forall \theta \in I(x_0, \varepsilon_0),$$

 $-1 \le \cos k_0 \theta \le 1 - \delta_0, \forall \theta \notin I(x_0, \varepsilon_0), 0 \le \theta < 2\pi$

holds, where $I(x_0, \varepsilon_0)$ is given by

(3.21)

$$I(x_0, \varepsilon_0) = \{\theta : 0 \le \theta < \varepsilon_0\} \cup \{\theta : |\theta - \frac{2\pi}{k_0}| < \varepsilon_0\} \cup \{\theta : |\theta - \frac{4\pi}{k_0}| < \varepsilon_0\} \cup \cdots \cup \{\theta : |\theta - \frac{2(k_0 - 1)\pi}{k_0}| < \varepsilon_0\} \cup \{\theta : 2\pi - \varepsilon_0 < \theta < 2\pi\}.$$

(3) Fix $0 < \sigma < \frac{\delta_0}{100}$, where $\delta_0 > 0$ is as in (2). There exist a positive integer m_1 and a neighborhood $W(x_0)$ of x_0 with $W(x_0) \in V(x_0)$ such that $S_{k_0m_1}(x,x_0) \neq 0$ for all $x \in W(x_0)$ and the real part of $\frac{S_{k_0(m_1+1)}(x,x_0)}{S_{k_0m_1}(x,x_0)}$ denoted by $\mathcal{R}_{k_0m_1}(x)$ satisfies

$$|1 - \mathcal{R}_{k_0 m_1}(x)| < \sigma, \forall x \in W(x_0).$$

The imaginary part of $\frac{S_{k_0(m_1+1)}(x,x_0)}{S_{k_0m_1}(x,x_0)}$ denoted by $\mathcal{I}_{k_0m_1}(x)$ satisfies the following inequality

$$|\mathcal{I}_{k_0m_1}(x)| < \frac{\sigma}{8}, \forall x \in W(x_0).$$

(4) For any positive constant c > 0, there exist a positive integer m_2 and a neighborhood $\hat{W}(x_0) \subseteq W(x_0)$ of x_0 such that

$$|S_{k_0m_2}(x,x_0)| > \frac{c}{2}, \forall x \in \hat{W}(x_0)$$

and

$$|S_{k_0m_2}(y,x_0)| < \frac{c}{8}, \forall y \notin \bigcup_{0 \le \theta < 2\pi} e^{i\theta} \circ W(x_0)$$

hold.

Proof. Fix $x_0 \in X_{k_0}$ and let D be the canonical local patch given in Theorem 1.12. From (2.32), we have for any $D' \in D$ and $N \in \mathbb{N}$, there exists a constant $C_{D',N}$ such that

(3.22)
$$|S_{k_0m}(x,x_0)| \ge \frac{k_0}{2\pi} \left| \hat{b}(z,0,k_0m) \right| e^{-k_0m\operatorname{Im}\Phi(z,0)} - C_{D',N}m^{-N}, m >> 1$$

holds. Given $x=(z,\theta)$ with $|z|\leq \frac{1}{m}, |\theta|\leq \frac{1}{m}$, one has $|S_{k_0m}(x,x_0)|>0$ when $m\gg 1$. Thus, there is a $\lambda_0>0$ such that for all $m\geq \lambda_0$ we have $|S_{k_0m}(x,x_0)|>0$ for all $x\in U_m(x_0)$, where $U_m(x_0)=\{(z,\theta):|z|<\frac{1}{m},|\theta|<\frac{1}{m}\}$. Moreover, from the proof of Lemma 3.1, we see that there is a $\lambda_1>0$ such that for all $m\geq \lambda_1$, there is a small neighborhood $\widetilde{U}_m(x_0)$ of x_0 such that $\Phi_m^{k_0}:\widetilde{U}_m(x_0)\to \mathbb{C}^{d_{k_0m_0}+d_{k_0(m_0+1)}}$ is an embedding. Taking $m_0\geq \lambda_0+\lambda_1$ and setting $U(x_0)=U_{m_0}(x_0)\cap U_{m_0+1}(x_0)\cap \widetilde{U}_{m_0}(x_0)$, we get (1).

Since $x_0 \in X_{k_0}$, we have $e^{i\frac{2\pi}{k_0}j} \circ x_0 = x_0$ for $0 \le j \le k_0, j \in \mathbb{Z}$. Then for any ε_0 we define $I(x_0, \varepsilon_0)$ as in (3.21). When ε_0 is sufficiently small there exists a small neighborhood of x_0 denoted by $V(x_0) \in U(x_0)$ such that $e^{i\theta} \circ V(x_0) \subset U(x_0)$ for $\theta \in I(x_0, \varepsilon_0)$. For $\theta \notin I(x_0, \varepsilon_0)$, $0 \le \theta < 2\pi$, we have $|k_0\theta - 2\pi j| \ge \varepsilon_0 k_0$ for every $j = 0, 1, \ldots, k_0$ which implies that there exists a constant δ_0 depending on ε_0 such that $-1 \le \cos k_0 \theta \le 1 - \delta_0$ for $\theta \notin I(x_0, \varepsilon_0)$. Thus we get the conclusion of (2) in this proposition.

From the proof of (1), there is an $\widetilde{m}_1 > 0$ such that for every $m \geq \widetilde{m}_1$, there is a neighborhood $W_m(x_0)$ of x_0 such that $S_{k_0m}(x,x_0) \neq 0$ and $S_{k_0(m+1)}(x,x_0) \neq 0$. We assume that $m \geq \widetilde{m}_1$ and $x \in W_m(x_0)$. By (2.32), we have

$$S_{k_0m}(x,x_0) \equiv \frac{k_0}{2\pi} e^{ik_0 m(x_{2n-1} + \Phi(z,0))} \hat{b}(z,0,m) \mod O(m^{-\infty}),$$

$$(3.23) \quad S_{k_0(m+1)}(x,x_0) \equiv \frac{k_0}{2\pi} e^{ik_0(m+1)(x_{2n-1} + \Phi(z,0))} \hat{b}(z,0,m+1) \mod O(m^{-\infty}),$$

$$\hat{b}(z,0,m) \sim \sum_{j=0}^{\infty} \hat{b}_j(z,0) m^{n-1-j} \text{ in } S_{\text{loc}}^{n-1}(1;D).$$

Write

$$\frac{S_{k_0(m+1)}(x,x_0)}{S_{k_0m}(x,x_0)} = \mathcal{R}_{k_0m}(x) + i\mathcal{I}_{k_0m}(x).$$

Since $\hat{b}_0(0,0) \neq 0$ (see Theorem 2.6), we have $\hat{b}(0,0,m) \neq 0$ for m large and this implies that $\hat{b}(z,0,m) \neq 0$ when |z| is sufficiently small. We assume that $\hat{b}(z,0,m) \neq 0$ for every $m \geq \widetilde{m}_1$ and every $(z,0) \in W_m(x_0)$. Set

$$a_m(x) = \frac{k_0}{2\pi} e^{ik_0 m(x_{2n-1} + \Phi(z,0))} \hat{b}(z,0,m), b_m(x) = S_{k_0 m}(x,x_0) - a_m(x).$$

From (3.23), for any $D' \in V(x_0) \in D$ and any $N \in \mathbb{N}$ there exists a positive constant $C_{D',N}$ such that

$$\sup_{x \in D'} |S_{k_0 m}(x) - a_m(x)| \le C_{D', N} m^{-N}, m >> 1,$$

holds. For any $m \geq \widetilde{m}_1$, define $V_m(x_0) = \{x = (z, \theta) \in D, |z| < \frac{1}{m}, |\theta| < \frac{1}{m}\} \cap W_m(x_0)$, then $V_m(x_0) \in D'$ when m is sufficiently large. Then on $V_m(x_0)$,

we have

(3.24)
$$|b_{m+1}(x)| \le C_{D',N} \frac{1}{(m+1)^N}, |b_m(x)| \le C_{D',N} \frac{1}{m^N}.$$

On the other hand, we have $|a_m(x)|=\frac{k_0}{2\pi}e^{-k_0m{\rm Im}\Phi(z,0)}\hat{b}(z,0,m)$. From (2.8), by a direct calculation we get ${\rm Im}\Phi(z,0)=\lambda|z|^2+O(|z|^3)$. So we assume D' to be sufficiently small such that on D' we have

$$|c_1|z|^2 \le \text{Im}\Phi(z,0) \le |c_2|z|^2$$

for some constants c_1, c_2 . Then

(3.25)
$$|a_m(x)| \ge \hat{c}m^{n-1}, \forall x \in V_m(x_0), \quad \frac{a_{m+1}(x)}{a_m(x)} \approx 1, \quad \forall x \in V_m(x_0),$$

holds for some positive constant \hat{c} when m is sufficiently large. Since

$$\frac{S_{k_0(m+1)}(x,x_0)}{S_{k_0m}(x,x_0)} = \frac{b_{m+1} + a_{m+1}}{b_m + a_m} = \frac{\frac{b_{m+1}}{a_m} + \frac{a_{m+1}}{a_m}}{\frac{b_m}{a_m} + 1},$$

(3.24) and (3.25) imply

$$\frac{S_{k_0(m+1)}(x, x_0)}{S_{k_0m}(x, x_0)} \approx 1, \forall x \in V_m(x_0)$$

for m >> 1. Then for any fixed $0 < \sigma < \frac{\delta_0}{100}$, we can choose m_1 sufficiently large such that $W(x_0) = \{(z,\theta): |z| < \frac{1}{m_1}, |\theta| < \frac{1}{m_1}\}$ satisfies $W(x_0) \in V(x_0)$ and on $W(x_0)$ we have

(3.26)
$$|1 - \mathcal{R}_{k_0 m_1}(x)| < \sigma, |\mathcal{I}_{k_0 m_1}(x)| < \frac{\sigma}{8}.$$

Thus, we get the conclusion of (3) in the proposition.

Choose a neighborhood $W_1(x_0)$ of x_0 such that $W_1(x_0) \in W(x_0)$ holds. Following the same arguments as in the proof of Lemma 2.3, we have

(3.27)
$$S_{k_0m}(x_0, y) \equiv 0 \mod O(m^{-\infty}), \forall y \notin \bigcup_{0 < \theta < 2\pi} e^{i\theta} \circ \overline{W_1(x_0)}.$$

Since $X \setminus \bigcup_{0 \le \theta < 2\pi} e^{i\theta} \circ W(x_0) \subseteq X \setminus \bigcup_{0 \le \theta < 2\pi} e^{i\theta} \circ \overline{W_1(x_0)}$, (3.27) implies that for any N > 0 there exists a constant C_N satisfying

$$|S_{k_0m}(x_0,y)| \le C_N m^{-N} \ \forall m >> 1, \forall y \in X \setminus \bigcup_{0 \le \theta < 2\pi} e^{i\theta} \circ W(x_0).$$

Thus, for any c>0, there exists n_0 such that for any $m>n_0$ we have $|S_{k_0m}(x_0,y)|<\frac{c}{8}$ for all $y\not\in\bigcup_{0\leq\theta<2\pi}e^{i\theta}\circ W(x_0)$. Then following the same arguments as in the proof

of (1) in the proposition, there exists a positive integer m_2 and a neighborhood $\hat{W}(x_0) \in W_1(x_0) \in W(x_0)$ such that $|S_{k_0m_2}(x,x_0)| > \frac{c}{2}$ holds for all $x \in \hat{W}(x_0)$ and moreover $|S_{k_0m_2}(x_0,y)| < \frac{c}{8}$ holds for all $y \notin X \setminus \bigcup_{0 \le \theta < 2\pi} e^{i\theta} \circ W(x_0)$. Thus, we get

the conclusion of (4) in this proposition.

3.3. Embedding of CR manifold by positive Fourier components. Now, we are going to establish the global embedding of the CR manifolds, which have a locally free transversal CR S^1 -action, by positive Fourier components.

Since $X \setminus X_{\text{reg}} \in X$, there exist finite $\hat{W}(x_i) \in W(x_i) \in V(x_i) \in U(x_i)$ and positive constants $m_0(x_i), m_1(x_i), m_2(x_i)$ with respect to the points $x_i, 0 \le i \le n_0$ satisfying the properties in Proposition 3.2 and moreover $X \setminus X_{\text{reg}} = \bigcup_{i=1}^{n_0} \hat{W}(x_i)$. Without loss of generality, we assume that $x_i \in X_{k_i}, 0 \le i \le n_0$. For every $i = 0, 1, \ldots, n_0$, set

$$H_{x_{i}} = \bigoplus_{j=0}^{2} \left(H_{b,k_{i}m_{j}(x_{i})}^{0}(X) \bigoplus H_{b,k_{i}(m_{j}(x_{i})+1)}^{0}(X) \right),$$

$$H_{m} = \bigoplus_{k=1}^{l} \left(H_{b,km}^{0}(X) \bigoplus H_{b,k(m+1)}^{0}(X) \right) \bigoplus_{i=0}^{n_{0}} H_{x_{i}}.$$

Recall that $X=X_1\cup X_2\cup\cdots\cup X_l$, $X_{\text{reg}}=X_1\neq\emptyset$. Put $N_m=\dim H_m$ and let $\{f_j\}_{j=1}^{N_m}$ be an orthonormal basis of H_m with respect to its decomposition. Define a map

$$\Phi_m: X \to \mathbb{C}^{N_m}, x \mapsto (f_1(x), \cdots, f_{N_m}(x)).$$

We will prove the following

Theorem 3.3. Let X be a compact connected strongly pseudoconvex CR manifold with a locally free transversal CR S^1 -action. Then Φ_m is an embedding when m is large.

Proof. By Lemma 3.1, we know that Φ_m is an immersion when m is large. Now we show that Φ_m is injective when m is large by seeking a contradiction. We assume that there exist two sequences $\{\hat{y}_m\}, \{\hat{z}_m\} \subset X, \ \hat{y}_m \neq \hat{z}_m$ such that $\Phi_m(\hat{y}_m) = \Phi_m(\hat{z}_m)$. Since X is compact, there exist subsequences of $\{\hat{y}_m\}, \{\hat{z}_m\}$ which are also denoted by $\{\hat{y}_m\}, \{\hat{z}_m\}$ such that $\hat{y}_m \to \hat{y}, \hat{z}_m \to \hat{z}$ for $m \to \infty$.

First we assume that $\hat{y}, \hat{z} \in X \setminus X_{\text{reg}}$.

Case I: $\hat{y} = e^{i\theta_0} \circ \hat{z}, \hat{z} \in X_k$ for some k > 1 and $\hat{z} \in U(x_i)$ for some i. By assumption of \hat{y}_m, \hat{z}_m we have that

(3.28)
$$S_{k_i m_0(x_i)}(\hat{y}, x_i) = S_{k_i m_0(x_i)}(\hat{z}, x_i), \\ S_{k_i (m_0(x_i)+1)}(\hat{y}, x_i) = S_{k_i (m_0(x_i)+1)}(\hat{z}, x_i).$$

In the following context, we will omit x_i in $m_j(x_i)$, j = 0, 1, 2 for brevity if it makes no confusion. Then (3.28) implies

$$e^{ik_i m_0 \theta_0} S_{k_i m_0}(\hat{z}, x_i) = S_{k_i m_0}(\hat{z}, x_i),$$

$$e^{ik_0 (m_0 + 1)\theta_0} S_{k_i (m_0 + 1)}(\hat{z}, x_i) = S_{k_i (m_0 + 1)}(\hat{z}, x_i).$$

By (1) in Proposition 3.2, we have $e^{ik_i\theta_0}=1$. Then $\theta_0=\frac{2\pi}{k_i}m$ holds for some $m\in\mathbb{Z}$. The rigid Hermitian metric on X implies that $e^{i\theta}:X\to X$ is an isometric map for each θ . Thus, we have

(3.29)
$$\operatorname{dist}(\hat{y}, x_i) = \operatorname{dist}(e^{i\frac{2\pi}{k_i}m} \circ \hat{z}, x_i) = \operatorname{dist}(e^{i\frac{2\pi}{k_i}m} \circ \hat{z}, e^{i\frac{2\pi}{k_i}m} \circ x_i) = \operatorname{dist}(\hat{z}, x_i).$$

This implies $\hat{y} \in U(x_i)$ if the $U(x_i)$ we chose is a geodesic ball centered at x_i . This is a contradiction since Φ_m is an embedding on $U(x_i)$.

case II: $\hat{y} \neq e^{i\theta} \circ \hat{z}, \forall 0 < \theta < 2\pi$. We assume $\hat{z} \in \hat{W}(x_i)$. Since Φ_m is an embedding on $U(x_i)$, we must have $\hat{y} \notin U(x_i)$. Now we have a claim as follows Claim: $\hat{y} \notin \bigcup_{0 \leq \theta \leq 2\pi} e^{i\theta} \circ W(x_i)$.

We prove the claim by seeking a contradiction. If it is not true, there exists a $\hat{z}_1 \in W(x_i)$ such that $\hat{y} = e^{i\hat{\theta}} \circ \hat{z}_1$ holds for some $\hat{\theta} \in [0, 2\pi)$. By (2) in Proposition 3.2 we have $\hat{\theta} \notin I(x_i, \varepsilon_i)$ and $-1 \le \cos k_i \hat{\theta} \le 1 - \delta_i$. Since

$$S_{k_i m_1}(\hat{y}, x_i) = S_{k_i m_1}(\hat{z}, x_i), \ S_{k_i (m_1 + 1)}(\hat{y}, x_i) = S_{k_i (m_1 + 1)}(\hat{z}, x_i),$$

we find

(3.30)
$$\frac{S_{k_i(m_1+1)}(\hat{z}, x_i)}{S_{k_i m_1}(\hat{z}, x_i)} = e^{ik_i \hat{\theta}} \frac{S_{k_i(m_1+1)}(\hat{z}_1, x_i)}{S_{k_i m_1}(\hat{z}_1, x_i)}.$$

From (3.30) we have

$$\mathcal{R}_{k_i m_1}(\hat{z}) + i \mathcal{I}_{k_i m_1}(\hat{z}) = (\cos k_i \hat{\theta} + i \sin k_i \hat{\theta}) (\mathcal{R}_{k_i m_1}(\hat{z}_1) + i \mathcal{I}_{k_i m_1}(\hat{z}_1))$$

which leads to

$$\mathcal{R}_{k_i m_1}(\hat{z}) = \mathcal{R}_{k_i m_1}(\hat{z}_1) \cos k_i \hat{\theta} - \mathcal{I}_{k_i m_1}(\hat{z}_1) \sin k_i \hat{\theta}.$$

Then we get

$$(3.31) 1 - \mathcal{R}_{k_i m_1}(\hat{z}) = 1 + (1 - \mathcal{R}_{k_i m_1}(\hat{z}_1)) \cos k_i \hat{\theta} - \cos k_i \hat{\theta} + \mathcal{I}_{k_i m_1}(\hat{z}_1) \sin k_i \hat{\theta}.$$

From (3.31) we have

$$|1 - \mathcal{R}_{k_i m_1}(\hat{z})| \ge 1 - \cos k_i \hat{\theta} - |1 - \mathcal{R}_{k_i m_1}(\hat{z}_1)| - |\mathcal{I}_{k_i m_1}(\hat{z}_1)|.$$

By (3) in Proposition 3.2 we have

$$\sigma \ge |1 - \mathcal{R}_{k_i m_1}(\hat{z})| \ge 1 - (1 - \delta_0) - \sigma - \frac{\sigma}{8},$$

that is

$$(2 + \frac{1}{8})\sigma \ge \delta_0.$$

This is contradiction with $0 < \sigma < \frac{\delta_0}{100}$. Thus we get the conclusion of the claim. From the above claim and by (4) in Proposition 3.2, we have

$$|S_{k_i m_2}(\hat{z}, x_i)| > \frac{c}{2}, |S_{k_i m_2}(\hat{y}, x_i)| < \frac{c}{8}.$$

This is a contradiction with

$$S_{k_i m_2}(\hat{z}, x_i) = S_{k_i m_2}(\hat{y}, x_i).$$

Next, we assume $\hat{y}, \hat{z} \in X_{\text{reg}}$.

Case III: $\hat{y}, \hat{z} \in X_{\text{reg}}$ and $\hat{y} = e^{i\hat{\theta}} \circ \hat{z}$ for some $\hat{\theta} \in [0, 2\pi)$. Choose canonical coordinates (z, θ, φ) centered at some $\hat{z}_0 \in X$ and defined on a canonical local patch $D = \{(z, \theta) : |z| < \varepsilon, |\theta| < \pi\}$ such that in terms of the canonical coordinates, $\hat{z} = (0, \theta_1), \ \hat{y} = (0, \theta_2)$. Then we have $\theta_2 - \theta_1 = \hat{\theta}$. Let $\{f_j\}_{j=1}^{d_m}$ and $\{g_j\}_{j=1}^{d_{m+1}}$ be orthonormal bases of $H^0_{b,m}(X)$ and $H^0_{b,m+1}(X)$, respectively. Then by the assumptions on \hat{y}_m and \hat{z}_m we have

$$S_m(\hat{z}_m, \hat{y}_m) = S_m(\hat{z}_m, \hat{z}_m), S_{m+1}(\hat{z}_m, \hat{y}_m) = S_{m+1}(\hat{z}_m, \hat{z}_m).$$

Without loss of generality, we assume $\hat{z}_m, \hat{y}_m \in D$ for each m. Then in terms of canonical local coordinates, we write $\hat{z}_m = (z_m, \theta_m)$ and $\hat{y}_m = (w_m, \eta_m)$. By Theorem 2.6 we have

$$S_{m}(\hat{z}_{m}, \hat{y}_{m}) = \frac{1}{2\pi} e^{im(\theta_{m} - \eta_{m} + \Phi(z_{m}, w_{m}))} \hat{b}(z_{m}, w_{m}, m) + O(m^{-\infty}),$$

$$S_{m+1}(\hat{z}_{m}, \hat{y}_{m}) = \frac{1}{2\pi} e^{i(m+1)(\theta_{m} - \eta_{m} + \Phi(z_{m}, w_{m}))} \hat{b}(z_{m}, w_{m}, m+1) + O(m^{-\infty}),$$

$$S_{m}(\hat{z}_{m}, \hat{z}_{m}) = \frac{1}{2\pi} \hat{b}(z_{m}, z_{m}, m) + O(m^{-\infty}),$$

$$S_{m+1}(\hat{z}_{m}, \hat{z}_{m}) = \frac{1}{2\pi} \hat{b}(z_{m}, z_{m}, m+1) + O((m+1)^{-\infty}).$$

We assume $\lim_{m\to\infty} m \mathrm{Im} \Phi(z_m,w_m) = M$ (M can be ∞).

(a): Assume

$$\lim_{m \to \infty} m \operatorname{Im} \Phi(z_m, w_m) = M \in (0, \infty].$$

From $S_m(\hat{z}_m, \hat{y}_m) = S_m(\hat{z}_m, \hat{z}_m)$ and (3.32) we have

$$e^{im(\theta_m - \eta_m + \Phi(z_m, w_m))} \hat{b}(z_m, w_m, m) = \hat{b}(z_m, z_m, m) + O(m^{-\infty}).$$

Then we get

$$m^{-(n-1)}|\hat{b}(z_m, w_m, m)|e^{-m\operatorname{Im}\Phi(z_m, w_m)} = m^{-(n-1)}|\hat{b}(z_m, z_m, m) + O(m^{-\infty})|.$$

Letting $m \to \infty$, we have

$$\hat{b}(0,0) = e^{-M}\hat{b}(0,0),$$

that is $\hat{b}(0,0) = 0$. Thus, we get a contradiction.

(b): Assume

(3.33)
$$\lim_{m \to \infty} m \operatorname{Im} \Phi(z_m, w_m) = 0.$$

From $S_{m+1}(\hat{z}_m, \hat{y}_m) - S_m(\hat{z}_m, \hat{y}_m) = S_{m+1}(\hat{z}_m, \hat{z}_m) - S_m(\hat{z}_m, \hat{z}_m)$ combined with (3.32) we have

$$m^{-(n-1)} \left| e^{im(\theta_m - \eta_m + \Phi(z_m, w_m))} \left[e^{i(\theta_m - \eta_m + \Phi(z_m, w_m))} \hat{b}(z_m, w_m, m + 1) - \hat{b}(z_m, w_m, m) \right] \right|$$

$$= m^{-(n-1)} \left| \hat{b}(z_m, z_m, m + 1) - \hat{b}(z_m, z_m, m) \right| + O(m^{-\infty}).$$

Letting $m \to \infty$ and using (3.33), we have

$$|e^{i\hat{\theta}}\hat{b}(0,0) - \hat{b}(0,0)| = 0.$$

Hence, $\hat{\theta} = 0$ and $\hat{z} = \hat{y}$. Put

$$f_m(t) = \frac{|S_m(t\hat{z}_m + (1-t)\hat{y}_m, \hat{y}_m)|^2}{S_m(t\hat{z}_m + (1-t)\hat{y}_m, t\hat{z}_m + (1-t)\hat{y}_m)S_m(\hat{y}_m, \hat{y}_m)}.$$

We have

(3.34)
$$f_m(0) = \frac{S_m(\hat{y}_m, \hat{y}_m)^2}{S_m(\hat{y}_m, \hat{y}_m)^2} = 1,$$

$$f_m(1) = \frac{|S_m(\hat{z}_m, \hat{y}_m)|^2}{S_m(\hat{z}_m, \hat{z}_m)S_m(\hat{y}_m, \hat{y}_m)} = \frac{S_m(\hat{y}_m, \hat{y}_m)^2}{S_m(\hat{y}_m, \hat{y}_m)S_m(\hat{y}_m, \hat{y}_m)} = 1.$$

By the Schwartz inequality, we get $0 \le f_m(t) \le 1$. Then (3.34) implies that there is a $t_m \in (0,1)$ such that $f_m'(t_m) = 0$ and $f_m''(t_m) \ge 0$ holds. Hence, we get

(3.35)
$$\liminf_{m \to \infty} \frac{f_m''(t_m)}{|z_m - w_m|^2 m} \ge 0.$$

Then, making use of the same arguments as in [19]((4.22)) in Theorem 4.7), we have that (3.35) is impossible under the assumption (3.33).

Case IV: $\hat{z}, \hat{y} \in X_{\text{reg}}, \hat{y} \neq e^{i\theta} \circ \hat{z}$ for any $\theta \in [0, 2\pi)$. Choose a canonical local patch $D(\hat{z})$ with canonical coordinates (z, θ, φ) centered at \hat{z} . Since $\hat{z} \in X_{\text{reg}}$, we can apply Lemma 1.13 and have that $D(\hat{z})$ can be chosen such that $D(\hat{z}) = \{(z, \theta) : |z| < \varepsilon, |\theta| < \pi\}$ holds in terms of canonical coordinates. Let ε be sufficiently small such that $\overline{D(\hat{z})}$ is S^1 -invariant. Since $\hat{y} \neq e^{i\theta} \circ \hat{z}$ for all $\theta \in [0, 2\pi)$, for ε small enough we can choose a canonical local patch $D(\hat{y})$ such that $\overline{D(\hat{y})} \cap \overline{D(\hat{z})} = \emptyset$ holds. Choose two functions $\chi, \chi_1 \in C_0^\infty(X)$ satisfying $\chi = 1$ in a small neighborhood of $\overline{D(\hat{z})}$ and $\chi_1 = 1$ in a small neighborhood of supp χ and supp $\chi \cap \overline{D(\hat{y})} = \sup \chi_1 \cap \overline{D(\hat{y})} = \emptyset$. Choose $\chi_0(w) \in C_0^\infty(\mathbb{C}^{n-1})$ such that $\sup \chi_0(w) \in \{w : |w| < 1\}$ and $\int_{\mathbb{C}^{n-1}} \chi_0(w) dv(w) = 1$ hold. Furthermore, choose $\eta_0(y_{2n-1}) \in C_0^\infty(-\pi,\pi)$ with $\int_{-\pi}^{\pi} \eta_0(y_{2n-1}) dy_{2n-1} = 1$. For any $m \in \mathbb{N}$, set

(3.36)
$$u_m(y) = m^{n-1} e^{im(y_{2n-1} - \theta_m - \operatorname{Re}\Phi(z_m, w))} \eta_0(y_{2n-1}) \chi_0(m(w - z_m)) \in C_0^{\infty}(D(\hat{z})).$$

Then we have

(3.37)
$$S_m u_m(\hat{y}_m) = \chi S_m u_m(\hat{y}_m) + (1 - \chi) S_m u_m(\hat{y}_m) = (1 - \chi) S_m u_m(\hat{y}_m)$$
 and

(3.38)
$$(1-\chi)S_m u_m(\hat{y}_m) = (1-\chi)S\chi_1 Q_m u_m(\hat{y}_m) + (1-\chi)S(1-\chi_1)Q_m u_m(\hat{y}_m).$$

Since $\overline{D(\hat{z})}$ is an S^1 -invariant subset and $\operatorname{supp} u_m \in D(\hat{z})$, we have $\operatorname{supp} Q_m u_m \in \overline{D(\hat{z})}$. This implies

(3.39)
$$(1 - \chi)S(1 - \chi_1)Q_m u_m(\hat{y}_m) = 0.$$

Then by the same arguments as in the proof of Lemma 2.3, we get

(3.40)
$$(1 - \chi) S \chi_1 Q_m u_m(\hat{y}_m) = O(m^{-\infty}).$$

Combining (3.37), (3.38), (3.39) and (3.40), we conclude

$$S_m u_m(\hat{y}_m) = O(m^{-\infty}).$$

On the other hand, we have

$$S_{m}u_{m}(\hat{z}_{m}) = \frac{m^{n-1}}{2\pi} \int_{X} e^{-m\operatorname{Im}\Phi(z_{m},w)} \hat{b}(z_{m},w,m) \chi_{0}(m(w-z_{m}))\lambda(w)dv(w) + O(m^{-\infty})$$

$$= \frac{1}{2\pi} \int_{\{w \in \mathbb{C}^{n-1}: |w| < 1\}} e^{-m\operatorname{Im}\Phi(z_{m},\frac{w}{m}+z_{m})} \hat{b}(z_{m},\frac{w}{m}+z_{m},m) \times$$

$$\chi_{0}(w)\lambda(\frac{w}{m}+z_{m})m^{-(n-1)}dv(w) + O(m^{-\infty}).$$

Since $\operatorname{Im}\Phi(z_m,\frac{w}{m}+z_m)\geq c_0|\frac{w}{m}|^2$ for some constant c_0 , we have $-m\operatorname{Im}\Phi(z_m,\frac{w}{m}+z_m)\to 0$ uniformly on $\{w\in\mathbb{C}^{n-1}:|w|<1\}$ as $m\to\infty$. Letting $m\to\infty$ we have

$$\lim_{m \to \infty} S_m u_m(\hat{z}_m) = \frac{1}{2\pi} \hat{b}(0,0) \neq 0.$$

This is a contradiction with the assumption $S_m u_m(\hat{z}_m) = S_m u_m(\hat{y}_m)$.

Case V: $\hat{z} \in X_{\text{reg}}, \hat{y} \notin X_{\text{reg}}$. We have that $\hat{y} \neq e^{i\theta} \circ \hat{z}$ for all $\theta \in [0, 2\pi)$. Following the same arguments as in Case IV, we find that this is impossible.

Thus, we get the conclusion of Theorem 3.3.

Acknowledgement

The authors would like to thank Professor Paul Yang for his interest in this work. The third-named author would like to thank the Institute of Mathematics, University of Cologne for the hospitality during his visit. We also thank the referee for many detailed remarks that have helped to improve the presentation.

REFERENCES

- [1] M.-S.Baouendi and L.-P. Rothschild and F. Treves, CR structures with group action and extendability of CR functions, Invent. Math., 83 (1985), 359-396.
- [2] D. M. Burns, Global behavior of some tangential Cauchy-Riemann equations, Partial differential equations and geometry (Proc. Conf., Park City, Utah, 1977), pp. 51-56, Lecture Notes in Pure and Appl. Math., 48, Dekker, New York, 1979.
- [3] D.-M. Burns and C.-L. Epstein, *Embeddability for three-dimensional CR manifolds*, J. Amer. Math. Soc. **3** (4) (1990), 809-841.
- [4] D.-E. Blair, *Contact Manifolds in Riemannian Geometry*, Lecture Notes in Mathematics, Springer-Verlag, **509** (1976).
- [5] J. Bland, Contact geometry and CR structures on \mathbb{S}^3 , Acta Math., 172 (1994), 1-49.
- [6] L. Boutet de Monvel, *Intégration des équations de Cauchy-Riemann induites formelles*, Séminaire Goulaouic-Lions-Schwartz 1974–1975; Équations aux derivées partielles linéaires et non linéaires, Centre Math., École Polytech., Paris, 1975, Exp. no. 9, pp. 13.
- [7] C.P. Boyer and K. Galicki, *Sasakian geometry*, Oxford mathematical monographs, Oxford Univ. Press, 2006.
- [8] C.P. Boyer, K. Galicki, S. Simanca, *Canonical Sasakian metrics*, Commun. Math. Phys. **279** (2008), 705C733.
- [9] L. Boutet de Monvel and J. Sjöstrand, *Sur la singularité des noyaux de Bergman et de Szegő*, Astérisque, 34-35 (1976), 123-164.
- [10] S. Chanillo, H-L. Chiu and P. Yang, *Embeddability for three-dimensional Cauchy-Riemann manifolds and CR Yamabi invariants*, Duke Math. J., **161**(15) (2012), 2909-2921.
- [11] S. Chanillo, H-L. Chiu and P. Yang, *Embedded three-dimensional CR manifolds and the non-negativity of Paneitz operators*, Contemporary Mathematics, **599** (2013), 65-82.
- [12] S. C. Chen, M. C. Shaw, *Partial differential equations in several complex variables*, AMS/IP Studies in Advanced Mathematics, 19, American Mathematical Society, Providence, RI; International Press, Boston, MA, 2001.
- [13] J-H. Cheng, C-Y. Hsiao and I-H. Tsai, Index theorems on CR manifolds with S^1 actions, arXiv:1511.00063.
- [14] C. L. Epstein, *CR-structures on three dimensional circle bundles*, Invent. Math., **109**(1992), 351-403.
- [15] C. Fefferman, *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. 26 (1974), 1-65.
- [16] H. Geiges, Normal Contact Structures on 3-Manifolds, Tohoku Math. J., 49 (1997), 415-422.
- [17] L. Hörmander, *The analysis of linear partial differential operators*, I, Classics in Mathematics, Springer-Verlag, Berlin, 2003.
- [18] C-Y. Hsiao, *Projections in several complex variables*, Mem. Soc. Math. France Nouv. Ser. **123** (2010).

- [19] C.-Y. Hsiao, Bergman kernel asymptotics and a pure analytic proof of Kodaira embedding theorem, Springer Proceedings in Mathematics and Statistics, Volume 144, (2015), 161-173.
- [20] C.-Y. Hsiao, Existence of CR sections for high power of semi-positive generalized Sasakian CR line bundles over generalized Sasakian CR manifolds, Ann. Glob. Anal. Geom., 47(1) (2015), 13–62. DOI 10.1007/s10455-014-9434-0.
- [21] C-Y. Hsiao, Szegő kernel asymptotics for high power of CR line bundles and Kodaira embedding theorems on CR manifolds, to appear in Memories of American Mathematical Society, arXiv:1401.6647.
- [22] C-Y. Hsiao and X. Li, Szegö kernel asymptotics and Morse inequalities on CR manifolds with S^1 action, arXiv:1502.02365.
- [23] C-Y. Hsiao and X. Li, Morse inequalities for Fourier components of Kohn Rossi cohomology of CR manifolds with S^1 action, Math. Z., 284 (2016), no. 1-2, 441-468.
- [24] C-Y. Hsiao, X. Li and G. Marinescu, Equivariant Kodaira embedding of CR manifolds with circle action, arXiv:1603.08872.
- [25] C-Y. Hsiao and G. Marinescu, Szegö kernel asymptotics and Morse inequalties on CR manifolds, Math.Z., **271** (2012), 509-573.
- [26] C-Y. Hsiao and G. Marinescu, Asymptotics of spectral function of lower energy forms and Bergman kernel of semi-positive and big line bundles, Comm. Anal. Geom. **22**(1) (2014), 1-108.
- [27] C-Y. Hsiao and G. Marinescu, *On the singularities of the Szegő projections on lower energy forms*, 57 pages, to appear in Journal of differential geometry, arXiv:1407.6305.
- [28] L. Lempert, *On three dimensional Cauchy-Riemann manifolds*, Jour. Amer. Math. Soc., **5** (4) (1992), 923-969.
- [29] G. Marinescu and N. Yeganefar, *Embeddability of some strongly pseudoconvex manifolds*, Trans. Amer. Math. Soc. 359 (2007), 4757-4771.
- [30] A. Melin and J. Sjöstrand, Fourier integral operators with complex-valued phase functions, Springer Lecture Notes in Math., 459, (1975), 120-223.
- [31] L. Ornea and M. Verbitsky, *Embeddings of compact Sasakian manifolds*, Mathematical Research Letters, **14**(4), (2007), 703-710.
- [32] L. Ornea and M. Verbitsky, *Sasakian structures on CR-manifolds*, Geometriae Dedicata, **125**(1), (2007), 159-173.
- [33] H. Rossi, Attaching analytic spaces to an analytic space along a pseudoconvex boundary, Proc. Conf. on Complex Manifolds (A. Aeppli, E. Calabi, R. Rohrl, eds.), Springer-Verlag,

Mathematical Institute, University of Cologne, Weyertal 86-90, 50931 Cologne, Germany

E-mail address: post@hendrik-herrmann.de or heherrma@math.uni-koeln.de

Institute of Mathematics, Academia Sinica, 6F, Astronomy-Mathematics Building, No.1, Sec.4, Roosevelt Road, Taipei 10617, Taiwan

E-mail address: chsiao@math.sinica.edu.tw or chinyu.hsiao@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, HUBEI 430072, CHINA & INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, 6F, ASTRONOMY-MATHEMATICS BUILDING, No.1, SEC.4, ROOSEVELT ROAD, TAIPEI 10617, TAIWAN

E-mail address: xiaoshanli@whu.edu.cn or xiaoshanli@math.sinica.edu.tw