# OPTIMAL SURVIVING STRATEGY FOR DRIFTED BROWNIAN MOTIONS WITH ABSORPTION

WENPIN TANG AND LI-CHENG TSAI

ABSTRACT. We study the 'Up the River' problem formulated by Aldous [Ald02], where a unit drift is distributed among a finite collection of Brownian particles on  $\mathbb{R}_+$ , which are annihilated once they reach the origin. Starting K particles at x = 1, we prove Aldous' conjecture [Ald02] that the 'push-the-laggard' strategy of distributing the drift asymptotically (as  $K \to \infty$ ) maximizes the total number of surviving particles, with approximately  $\frac{4}{\sqrt{\pi}}K^{1/2}$  surviving particles. We further establish the hydrodynamic limit of the particle density, in terms of a two-phase Partial Differential Equation (PDE) with a moving boundary, by utilizing certain integral identities and coupling techniques.

#### 1. INTRODUCTION

In this paper we study the 'Up the River' problem formulated by Aldous [Ald02]. That is, we consider K independent Brownian particles, which all start at x = 1, and are absorbed (annihilated) once they hit x = 0. Granted a unit drift, we ask what is the optimal strategy of dividing and allocating the drift among all surviving particles in order to maximize the number of particles that survive forever. More precisely, letting  $B_i(t)$ ,  $i = 1, \ldots, K$ , denote independent standard Brownian motions, we define the model as a  $\mathbb{R}^K_+$ -valued diffusion  $(X_i(t); t \ge 0)_{1 \le i \le K}$ , satisfying

$$X_{i}(t) = 1 + B_{i}(t \wedge \tau_{i}) + \int_{0}^{\tau_{i}} \phi_{i}(s) ds.$$
(1.1)

Here  $\tau_i := \inf\{t > 0 : X_i(t) = 0\}$  denotes the absorption time of the *i*-th particle, and the strategy is any  $[0,1]^K$ -valued,  $\{B_i(t)\}_{i=1}^K$ -progressively measurable function  $(\phi_i(t); t \ge 0)_{1 \le i \le K}$  such that  $\sum_{i=1}^K \phi_i(t) \le 1$ ,  $\forall t \ge 0$ . Our goal is to maximize  $\widetilde{U}(\infty)$ , where

$$\widetilde{U}(\infty) := \lim_{t \to \infty} \widetilde{U}(t), \quad \widetilde{U}(t) := |\{i : X_i(t) > 0\}|.$$

(Here  $\tilde{U}(t)$  actually depends on K, but we suppress the dependence in this notation, and reserve notations like  $\tilde{U}_K(t)$  for scaled quantities). Inspired by the 'Up the River: Race to the Harbor' board game, this simple model serves as a natural optimization problem for a random environment with limited resources. For K = 2, [MS06] obtains an explicit expression of the law of  $\tilde{U}(t)$ , and for large K, numerical results are obtained in [Han13] for the discrete analog of (1.1).

Focusing on the asymptotic behavior as  $K \to \infty$ , we prove that the optimal strategy is the naïve **push-the-laggard strategy** 

$$\phi_i(t) := \mathbf{1}_{\{X_i(t) = Z(t)\}}, \quad \text{where } Z(t) := \min\{X_i(t) : X_i(t) > 0\}, \tag{1.2}$$

<sup>2010</sup> Mathematics Subject Classification. Primary 60K35; Secondary 35Q70, 82C22.

Key words and phrases. Atlas model, competing Brownian particles, hydrodynamic limit, Stefan problems, moving boundary.

which allocates all the unit drift on the laggard Z(t).

**Remark 1.1.** Ties may occur in (1.2), namely  $|\{i : X_i(t) = Z(t)\}| > 1$ . Here we break the ties in an *arbitrarily* fixed manner. That is, any  $(\phi_i(t))_i$  satisfying  $\sum_{i:X_i(t)=Z(t)} \phi_i(t) = 1$ is regarding as a push-the-laggard strategy. As bounds in this paper is independent of the exact choice of breaking the ties, hereafter we fix some arbitrary way of breaking the ties and refer to (1.2) as *the* push-the-laggard strategy.

Furthermore, we prove that, due to self-averaging,  $\widetilde{U}(\infty)$  is in fact deterministic to the leading order under the push-the-laggard strategy, more explicitly  $\widetilde{U}(\infty) \approx \frac{4}{\sqrt{\pi}} K^{1/2}$ . Let  $\widetilde{U}_K(t) := K^{-1/2} \widetilde{U}(t)$ . The following is our main result:

# Theorem 1.2.

(a) Regardless of the strategy, for any fixed  $n < \infty$  and  $\gamma \in (0, \frac{1}{4})$ , we have

$$\mathbf{P}\left(\widetilde{U}_{K}(\infty) \leq \frac{4}{\sqrt{\pi}} + K^{-\gamma}\right) \geq 1 - CK^{-n}, \quad \forall K < \infty,$$
(1.3)

where  $C = C(n, \gamma) < \infty$  depends only on n and  $\gamma$ , not on the strategy.

(b) Under the push-the-laggard strategy, for any fixed  $\alpha \in (0, \frac{1}{96})$  and  $n < \infty$ , we have

$$\mathbf{P}\left(|\widetilde{U}_{K}(\infty) - \frac{4}{\sqrt{\pi}}| \le K^{-\alpha}\right) \ge 1 - CK^{-n}, \quad \forall K < \infty,$$
(1.4)

where  $C = C(\alpha, n) < \infty$  depends only on  $\alpha$  and n.

**Remark 1.3.** While the exponent  $-\frac{1}{4}^+$  of the error term in Theorem 1.2(a) (originating from the control on the relevant martingales) is optimal, the choice of exponent  $\alpha \in (0, \frac{1}{96})$  in Theorem 1.2(b) is purely technical. The later may be improved by establishing sharper estimates, which we do not pursue in this paper.

Theorem 1.2 resolves Aldous' conjecture [Ald02, Conjecture 2] in a slightly different form. The intuition leading to such a theorem, as well as the the main ingredient of proving it, is the hydrodynamic limit picture given in [Ald02]. To be more precise, we consider the diffusively scaled process  $X_i^K(t) := K^{-1/2}X_i(tK)$  under the push-the-laggard strategy, with  $Z_K(t) := \min\{X_i^K(t) : X_i^K(t) > 0\} = K^{-1/2}Z(tK)$  and the scaled complementary distribution function

$$\widetilde{U}_{K}(t,x) := K^{-1/2} \sum_{i=1}^{K} \mathbf{1}_{(x,\infty)}(X_{i}^{K}(t)).$$
(1.5)

Let p(t,x) denote the standard heat kernel, with  $p^{N}(t,y,x) := p(t,y-x) + p(t,y+x)$ denoting the Neumann heat kernel. We expect  $(\widetilde{U}_{K}(t,x), Z_{K}(t))$  to be well-approximated by  $(\widetilde{U}_{\star}(t,x), z_{\star}(t))$ . Here  $\widetilde{U}_{\star}(t,x)$  and  $z_{\star}(t)$  are deterministic functions, which are defined in *two* separated phases as follows. For  $t \leq \frac{1}{2}$ , the **absorption phase**, we define

$$\widetilde{U}_{\star}(t,x) := 2p(t,x) + \int_0^t p^{\mathsf{N}}(t-s, z_{\star}(s), x) ds, \quad \forall t \le \frac{1}{2}, \ x \ge 0,$$
(1.6)

$$z_{\star}(t) := 0, \quad \forall t \le \frac{1}{2}; \tag{1.7}$$

and for  $t > \frac{1}{2}$ , the **moving boundary phase**, we define

$$\widetilde{U}_{\star}(t,x) := 2p(t,x) + \int_0^t p^{N}(t-s, z_{\star}(s), x) ds, \quad \forall t > \frac{1}{2}, \ x \ge z_{\star}(t),$$
(1.8)

where  $z_{\star}(t)$  is characterized by the following integral equation

$$\begin{cases} z_{\star}(\cdot) \in \mathcal{C}(\mathbb{R}_{+}), \text{ nondecreasing,} \\ \int_{0}^{\infty} p(t - \frac{1}{2}, z_{\star}(t) - y) U_{\star}(\frac{1}{2}, y) dy = \int_{\frac{1}{2}}^{t} p(t - s, z_{\star}(t) - z_{\star}(s)) ds, \quad \forall t \in (\frac{1}{2}, \infty), \end{cases}$$
(1.9)

for  $U_{\star}(\frac{1}{2}, x) := \widetilde{U}_{\star}(\frac{1}{2}, 0) - \widetilde{U}_{\star}(\frac{1}{2}, x)$ . We prove in Section 4 that (1.9) admits a unique solution.

The solution  $(\tilde{U}_{\star}, z_{\star})$  to (1.6)-(1.9) is closely related to certain PDE problems, as follows. Let  $\tilde{\Phi}(t, y) := \mathbf{P}(B(t) \geq y)$  denote the complementary error function. For  $t \leq \frac{1}{2}$ , with  $z_{\star}(t) = 0$ , we use integration by parts in (1.6) to get  $2\int_{0}^{t} p(t-s,x)ds = 2\int_{0}^{t} p(s,x)ds = 4tp(t,x) + 4\int_{0}^{t} s\partial_{s}p(s,x)ds$ , followed by the change of variable  $s \to \frac{x}{\sqrt{s}}$  applied to the last integral, thereby obtaining  $\tilde{U}_{\star}(t,x) = 2p(t,x) + 4tp(t,x) - 4x\tilde{\Phi}(t,x)$ . This expression is further rewritten as

$$\widetilde{U}_{\star}(t,x) = \int_{x}^{\infty} u_1(t,y) dy, \qquad (1.10)$$

where 
$$u_1(t,x) := -2\partial_x p(t,x) + 4\widetilde{\Phi}(t,x).$$
 (1.11)

That is,  $\widetilde{U}_{\star}(t, \cdot)$  is the complementary distribution function of  $u_1(t, \cdot)$  as in (1.11), which solves the following PDE with a fixed boundary at x = 0:

$$\partial_t u_1 = \frac{1}{2} \partial_{xx} u_1 \quad \forall 0 < t < \frac{1}{2}, \ x > 0,$$
 (1.12a)

$$u_1(t,0) = 2, \quad \forall 0 < t < \frac{1}{2},$$
 (1.12b)

$$\lim_{t \to 0} (u_1(t, x) + 2\partial_x p(t, x)) = 0, \quad \forall x \ge 0.$$
(1.12c)

For  $t > \frac{1}{2}$ , we consider the following **Stefan problem**, a PDE with a moving boundary:

$$z_2(t)$$
 nondecreasing,  $z_2(\frac{1}{2}) = 0$ , (1.13a)

$$\partial_t u_2 = \frac{1}{2} \partial_{xx} u_2, \quad \forall t > \frac{1}{2}, \ x > z_2(t) \tag{1.13b}$$

$$u_2(\frac{1}{2}, x) = u_1(\frac{1}{2}, x), \quad \forall x \ge 0,$$
 (1.13c)

$$u_2(t, z_2(t)) = 2, \quad \forall t \ge \frac{1}{2},$$
 (1.13d)

$$2\frac{d}{dt}z_2(t) + \frac{1}{2}\partial_x u_2(t, z_2(t)) = 0, \quad \forall t > \frac{1}{2}.$$
(1.13e)

As we show in Lemma 4.1, for each sufficiently smooth solution  $(u_2, z_2)$  to (1.13),  $\widetilde{U}_{\star}(t, x) := \int_x^\infty u_2(t, y) \mathbf{1}_{\{y \ge z_{\star}(t)\}} dy$  and  $z_{\star}(t) := z_2(t) \mathbf{1}_{\{t>1/2\}}$  satisfy (1.8)–(1.9).

**Remark 1.4.** Here we take the integral representations (1.8)-(1.9) as the *definition* of the hydrodynamic limit equation. This formulation is more convenient for our purpose, and in particular it does not require showing the smoothness of the solution. We note that, however, it should be possible to establish classical solutions to (1.13) with sufficient smoothness, by converting (1.13) to the parabolic variational inequality. See, for example, [Fri10]. We do not pursue this direction here.

Before stating the precise result on hydrodynamic limit, we explain the intuition of how (1.12)-(1.13) arise from the behavior of the particle system. Indeed, the heat equations (1.12a) and (1.13b) model the diffusive behavior of  $(X_i^K(t))_i$  away from  $Z_K(t)$ . In view of the equilibrium measure of gaps of the infinite Atlas model [PP08], near  $Z_K(t)$  we expect

the particle density to be 2 to balance the drift exerted on  $Z_K(t)$ , yielding the boundary conditions (1.12b) and (1.13d). The function  $-2\partial_x p(t, x)$  is the average density of the system without the drift. (The singularity of  $-2\partial_x p(t, x)$  at t = 0 captures the overabundance of particles at t = 0 compared to the scaling  $K^{1/2}$ .) As the drift affects little of the particle density near t = 0, we expect the initial condition (1.12c). The absorption phase ( $t \leq \frac{1}{2}$ ) describes the initial state of the particle system with high density, where particles are constantly absorbed, yielding a fixed boundary  $Z_K(t) \approx 0$ . Under the push-the-laggard strategy, the system enters a new phase at  $t \approx \frac{1}{2}$ , where the density of particles is low enough ( $\leq 2$  everywhere) so that the drift carries all remaining particles away from 0. This results in a moving boundary  $Z_K(t)$ , with an additional boundary condition (1.13e), which simply paraphrases the conservation of particles  $\frac{d}{dt} \int_{z_2(t)}^{\infty} u_2(t, y) dy = 0$ .

Hereafter we use  $|f|_{\mathcal{C}(D)} := \sup_{x \in D} |f(x)|$  to denote the uniform norm over the designated domain D. The following is our result on the hydrodynamic limit of  $(\widetilde{U}_K(t,x), Z_K(t))$ :

**Theorem 1.5** (hydrodynamic limit). Under the push-the-laggard strategy, for any fixed  $\alpha \in (0, \frac{1}{96})$  and  $T, n < \infty$ , there exists  $C = C(T, L, \alpha, n) < \infty$  such that

$$\mathbf{P}\left(\sup_{t\in[0,T],x\in\mathbb{R}}\left\{|\widetilde{U}_{K}(t,x)-\widetilde{U}_{\star}(t,x)|t^{\frac{3}{4}}\right\}\leq CK^{-\alpha}\right)\geq1-CK^{-n},\quad\forall K<\infty,\qquad(1.14)$$

$$\mathbf{P}\left(|Z_K - z_\star|_{\mathcal{C}([0,T])} \le CK^{-\alpha}\right) \ge 1 - CK^{-n}, \quad \forall K < \infty.$$
(1.15)

Under the push-the-laggard strategy (1.2), the process  $(X_i^K(t))_i$  is closely related to the Atlas model [Fer02]. The latter is a simple special case of diffusions with rank-dependent drift: see [BFK05, CP10, CP11, IPB<sup>+</sup>11, IK10, IKS13], for their ergodicity and sample path properties, and [DSVZ12, PS14] for their large deviations properties as the dimension tends to infinity. In particular, the hydrodynamic limit and fluctuations of the Atlas-type model have been analyzed in [CDSS, DT15, HJV15].

Here we take one step further and analyze the combined effect of rank-dependent drift and absorption, whereby demonstrating the two-phase behavior. With the absorption at x = 0, previous methods of analyzing the large scale behaviors of diffusions with rank-dependent drift do not apply. In particular, the challenge of proving Theorem 1.5 originates from the lack of invariant measure (for the absorption phase) and the singularity at t = 0, where a rapid transition from K particles to an order of  $K^{1/2}$  particles occurs. Here we solve the problem by adopting a *new* method of exploiting certain integral identities of the particle system that mimic (1.6)–(1.9). Even though here we mainly focus on the push-the-laggard strategy under the initial condition  $X_i(0) = 1$ ,  $\forall i$ , the integral identities apply to general rank-dependent drifts and initial conditions, and may be used for analyzing for general models with both rank-dependent drifts and absorption.

Acknowledgment. We thank David Aldous for suggesting this problem for research. WT thanks Jim Pitman for helpful discussion throughout this work, and Craig Evans for pointing out the relation between (1.13) and parabolic variational inequalities. LCT thanks Amir Dembo for enlightening discussion at the early stage of this work. LCT was partially supported by the NSF through DMS-0709248.

#### 2. Proof of Theorem 1.2 and 1.5

Throughout this paper we use  $C = C(\alpha, \beta, ...) < \infty$  to denote a generic deterministic finite constant depends only on the designated variables.

With  $z_{\star}(t)$  as in (1.7) and (1.9), we unify the integral representations (1.6) and (1.8) into a single one as

$$\widetilde{U}_{\star}(t,x) = 2p(t,x) + \int_0^t p^{N}(t-s, z_{\star}(s), x)ds, \quad \forall t > 0, \ x \ge z_{\star}(t).$$
(2.1)

Adopting the notations  $\phi_i^K(t) := \phi(tK), \ \tau_i^K := K^{-1}\tau_i$ , and

$$\Psi(t, y, x) := \Phi(t, y - x) - \widetilde{\Phi}(t, y + x),$$
(2.2)

$$\widetilde{G}_{K}(t,x) := K^{\frac{1}{2}}\Psi(t,K^{-\frac{1}{2}},x) = K^{\frac{1}{2}} \int_{0}^{K^{-\frac{1}{2}}} p^{N}(t,y,x)dy, \qquad (2.3)$$

in Section 3, we establish the following microscopic analog of (2.1).

**Proposition 2.1.** Let  $(\phi_i^K(t); t \ge 0)_{1 \le i \le K}$  be any given strategy. For all  $t < \infty$  and  $x \ge 0$ ,

$$\widetilde{U}_{K}(t,x) = \widetilde{G}_{K}(t,x) + \sum_{i=1}^{K} \int_{0}^{t \wedge \tau_{i}^{K}} \phi_{i}^{K}(s) p^{N}(t-s, X_{i}^{K}(s), x) ds + R_{K}(t,x).$$
(2.4)

Here  $R_K(t,x)$  is some remainder term such that, given any  $T, n < \infty$  and  $\gamma \in (0, \frac{1}{4})$ 

$$\mathbf{P}\left(\sup_{x\in\mathbb{R}}|R_{K}(t,x)| \le K^{-\frac{1}{4}+\gamma}t^{-\frac{3}{4}}, \ \forall t\le T\right) \ge 1 - CK^{-n}, \quad \forall K<\infty,$$
(2.5)

where  $C = C(T, \gamma, n) < \infty$  is independent of the strategy.

Now, with  $\widetilde{U}_K(t) = \widetilde{U}_K(t, 0)$ , specializing (2.4) at  $t = \frac{1}{2}$  yields

$$\widetilde{U}_{K}(\frac{1}{2}) = K^{\frac{1}{2}} \int_{0}^{K^{-1/2}} 2p(\frac{1}{2}, x) dx + \sum_{i=1}^{K} \int_{0}^{\frac{1}{2}} \phi_{i}^{K}(s) 2p(\frac{1}{2} - s, X_{i}^{K}(s)) ds + R_{K}(\frac{1}{2}, 0)$$

$$\leq 2p(\frac{1}{2}, 0) + \int_{0}^{\frac{1}{2}} 2p(\frac{1}{2} - s, 0) ds + R_{K}(\frac{1}{2}, 0) = \frac{4}{\sqrt{\pi}} + R_{K}(\frac{1}{2}, 0).$$
(2.6)

With  $U_K(\infty) \leq U_K(\frac{1}{2})$ , combining (2.6) with (2.5) yields Theorem 1.2(a).

Postponing the proof of Theorem 1.2(b), we proceed to proving Theorem 1.5. Specialize (2.4) to the push-the-laggard strategy (1.2) to obtain

$$\widetilde{U}_K(t,x) = \widetilde{G}_K(t,x) + \int_0^t p^{\mathcal{N}}(t-s, Z_K(s), x)ds + R_K(t,x), \quad \forall t < \tau_{\text{ext}},$$
(2.7)

where  $\tau_{\text{ext}} := \max\{\tau_i^K : i\}$  is the extinction time. In Section 5, using a simple comparison argument we show that

**Lemma 2.2.** For any fixed  $T, n < \infty$ , there exists  $C = C(T, n) < \infty$  such that  $\mathbf{P}(\tau_{ext} > T) \ge 1 - CK^{-n}, \forall K < \infty$ .

Next, for the heat kernel p(t, x) it is standard to show that

$$|p(t,x) - p(t,x')| \le C(\alpha)|x - x'|^{\alpha} t^{-\frac{1+\alpha}{2}}, \qquad \forall \alpha \in (0,1),$$
(2.8)

$$|p(t,x) - p(t',x)| \le C(\alpha)|t - t'|^{\frac{\alpha}{2}}(t')^{-\frac{1+\alpha}{2}}, \qquad \forall \alpha \in (0,1), \ t' < t < \infty.$$
(2.9)

From (2.8) we obtain

$$|\widetilde{G}_K(t,x) - 2p(t,x)| \le C(\alpha) K^{-\frac{\alpha}{2}} t^{-\frac{1+\alpha}{2}}, \qquad \qquad \forall \alpha \in (0,1), \qquad (2.10)$$

$$\int_{0}^{s} |p^{N}(t-s,x,z) - p^{N}(t-s,x,z')| ds \le C(\alpha')|z-z'|^{\alpha'} t^{\frac{1-\alpha'}{2}}, \quad \forall \alpha' \in (0,1).$$
(2.11)

With these estimates and Lemma 2.2, comparing (2.1) and (2.7), we find that (1.14) follows immediately form (1.15). The later is established for *i*) the absorption phase; and *ii*) the moving boundary phase, as follows.

(i) In Section 5 adapting a stopping argument to analyze (2.7), we prove

**Proposition 2.3.** For any fixed  $\beta \in (0, \frac{1}{24})$  and  $n < \infty$ , there exists  $C = C(\beta, n) < \infty$  such that

$$\mathbf{P}\left(\sup_{s\leq\frac{1}{2}+\frac{1}{7}K^{-2\beta}}Z_{K}(s)\leq K^{-\beta}\right)\geq 1-CK^{-n}, \ \forall K<\infty.$$
(2.12)

Further, as shown in Section 4,  $z_{\star}(s)$  grows quadratically near  $s = \frac{1}{2}$ ,

**Lemma 2.4.** We have  $\lim_{t\downarrow 0} \{t^{-2}z_{\star}(t+\frac{1}{2})\} = \frac{2}{\sqrt{\pi}}$ .

With this and  $z_{\star}(s) = 0$ ,  $\forall s \leq \frac{1}{2}$ , Proposition 2.3 implies (1.15) for  $s \leq \frac{1}{2} + \frac{1}{7}K^{-2\beta}$ , for any fixed  $\beta \in (0, \frac{1}{96})$ .

(ii) In Section 6.3 we establish the hydrodynamic lower bound as follows

**Proposition 2.5.** For any fixed  $\alpha \in (0, \frac{1}{24})$  and  $T, n < \infty$ , there exists  $C = C(T, \alpha, n) < \infty$  such that  $\mathbf{P}(\inf_{s \in [\frac{1}{2},T]}(Z_K(s) - z_{\star}(s)) \geq -K^{-\alpha}) \geq 1 - CK^{-n}, \forall K < \infty$ .

Combining this with Lemma 2.4, we then conclude that for any fixed  $\beta \in (0, \frac{1}{96}), T, n < \infty$ ,

$$\mathbf{P}\left(\inf_{s\in[\frac{1}{2}+\frac{1}{7}K^{-2\beta},T]}Z_K(s)>0\right)\geq 1-C(T,\beta,n)K^{-n},\quad\forall K<\infty.$$

As (1.15) is proven for  $s \leq \frac{1}{2} + \frac{1}{7}K^{-2\beta}$  in (*i*), to prove the hydrodynamic upper bound, we consider only  $s > \frac{1}{2} + \frac{1}{7}K^{-2\beta}$ , and, without loss of generality, *turn off* the absorption for  $s > \frac{1}{2} + K^{-2\beta}$ . That is, we fix  $\beta \in (0, \frac{1}{96})$  and consider the modified process  $(X_i^{*,K}(s))_i$ , which is the same as  $(X_i^K(s))_i$  except that no particle is absorbed for  $s \geq \frac{1}{2} + K^{-2\beta}$ , with  $Z_K^*(s) := \min\{X_i^{*,K}(s) : X_i^{*,K}(s) > 0\}$ . Obtaining the hydrodynamic limit of  $Z_K^*(s)$  requires the following analog of (2.4) for Atlas models. More precisely, for  $\lambda > 0$ , we say that  $(Y_i(t); t)_{1 \leq i \leq n}$  is a  $\lambda$ -Atlas model (with *n* particles) if it evolves according to the following stochastic differential equations:

$$dY_i(t) = \mathbf{1}_{\{X_i(t) = Z^Y(t)\}} \,\lambda dt + dB_i(t) \quad \text{for } 1 \le i \le n, \quad Z^Y(t) := \min\{Y_i(t)\}, \tag{2.13}$$

and call  $(Y_i(t))_i$  a **unit Atlas model** if  $\lambda = 1$ . Let  $Y_i^K(t) := K^{-1/2}Y_i(tK)$  and  $Z_K^Y(t) := K^{-1/2}Z^Y(tK)$  denote the scaled processes. (Here K is just a scaling parameter, not necessarily related to the number of particles). Define the corresponding empirical measure

$$\mu_{t,K}^{Y}(\bullet) := K^{-1/2} \sum_{i} \delta_{Y_{i}^{K}(t)}(\bullet)$$
(2.14)

and distribution function  $U_K^Y(t, x) := \langle \mu_{t,K}^Y, \mathbf{1}_{(0,x)} \rangle$ , and let  $||X||_p := (\mathbf{E} |X|^p)^{1/p}$  denote the  $L^p$  norm of a random variable. In Section 3 we prove

**Proposition 2.6.** Let  $t_{in} < \infty$  and  $(Y_i(t); t \ge Kt_{in})_i$  be a unit Atlas model. We let  $Y^K := (Y_i^K(t); t \ge t_{in})_i$ ,  $U_K^Y(t, x)$  and  $\mu_{t,K}^Y$  to be as in the preceding. Assume  $(Y_i^K(t_{in}))_i$  satisfies the following initial condition: given any  $\alpha \in (0, 1)$  and  $n < \infty$ , there exist  $D_*, D_{\alpha,n} < \infty$  such that

$$\mathbf{P}(|\{Y_i^K(t_{in})\}_i| \le K) \ge 1 - \exp(-\frac{1}{D_*}K^{1/2}),$$
(2.15)

$$\|\mu_{t_{in},K}^{Y}([a,b])\|_{n} \le D_{\alpha,n}|b-a|^{\alpha}, \quad \forall |b-a| \ge K^{-1/2}.$$
(2.16)

Then, for all  $t > t_{in}$ , we have

$$U_K^Y(t,x) = \int_0^\infty p(t-t_{in}, x-y) U_K^Y(t_{in}, y) dy - \int_{t_{in}}^t p(t-s, Z_K^Y(s), x) ds + R_K^Y(t, x).$$
(2.17)

Here  $R_K^Y(t,x)$  is a remainder term such that, given any  $T, n < \infty$  and  $\gamma \in (0, \frac{1}{4})$ ,

$$\mathbf{P}\left(|R_K^Y|_{\mathcal{C}([t_{in},T]\times\mathbb{R})} \le K^{-\frac{1}{4}+\gamma}\right) \ge 1 - C_D K^{-n}, \ \forall K < \infty,$$
(2.18)

where  $C_D < \infty$  depends only on T, n and  $D_*, D_{\alpha,j}, j = 1, \ldots, n$ .

By using Proposition 2.6, in Section 6.2 we prove

**Proposition 2.7.** For any fixed  $\alpha < \beta \in (0, \frac{1}{96})$  and  $T, n < \infty$ , there exists  $C = C(T, \alpha, n) < \infty$  such that  $\mathbf{P}(\sup_{s \in [\frac{1}{2} + \frac{1}{7}K^{-2\beta}, T]}(Z_K^*(s) - z_\star(s)) \leq K^{-\alpha}) \geq 1 - CK^{-n}, \forall K < \infty.$ 

Now we turn to proving Theorem 1.2(b). In Section 6.3, we prove the following proposition via a simple scaling argument.

**Proposition 2.8.** For any fixed  $n < \infty$ , there exists  $C = C(n) < \infty$  such that

$$\mathbf{P}\left(\inf_{t\geq 1} Z_K(\frac{1}{2}+t) > 0\right) \ge 1 - CK^{-n}, \ \forall K < \infty.$$
(2.19)

As shown in Lemma 4.2,  $\widetilde{U}_{\star}(s, z(s)) = \widetilde{U}_{\star}(\frac{1}{2}, 0) = \frac{4}{\sqrt{\pi}}, \forall s \geq \frac{1}{2}$ . With this, Theorem 1.2(b) readily follows from Theorem 1.5 and Proposition 2.8.

The rest of the article is organized as follows. In Section 3 we establish Propositions 2.1 and 2.6 concerning the microscopic integral representations. In Section 4, we settle the uniqueness and stability of (1.9), as well as Lemma 2.4. Based on this, in Section 5 we prove Proposition 2.3. In Section 6, we construct a solution to (1.9) as the hydrodynamic limit of a certain Atlas process, and prove Propositions 2.5-2.8.

# 3. Proof Propositions 2.1 and 2.6

Recall that the scaled empirical measure of a  $\lambda$ -Atlas model is defined in (2.14). We now define the analogous scaled empirical measure of  $(X_i^K(t))_i$ :

$$\mu_{t,K}(\bullet) := K^{-1/2} \sum_{\{i:X_i^K(t)>0\}} \delta_{X_i^K(t)}(\bullet).$$
(3.1)

The first step is to derive (2.4) and (2.17), via Ito calculus, with explicit remainder terms  $R_K(t,x)$  and  $R_K^Y(t_{\rm in},t,x)$ . To this end, we note that under the scaling  $X_i^K(s) := K^{-1/2}X_i(sK)$ , the SDE (1.1) transforms into

$$dX_{i}^{K}(s) = \phi_{i}^{K}(s)K^{\frac{1}{2}}d(s \wedge \tau_{i}^{K}) + dB_{i}^{K}(s \wedge \tau_{i}^{K}).$$
(3.2)

Recall the definition of  $\Psi(t, y, x)$  from (2.2). To derive (2.4), fixing arbitrary  $t < \infty, x \ge 0$ and  $\varepsilon > 0$ , with  $\Psi(t + \varepsilon - s, y)$  solving the backward heat equation  $\partial_s \Psi + \frac{1}{2} \partial_{yy} \Psi = 0$ , we apply Itô's formula to  $F_i(s) := \Psi(t + \varepsilon - s, X_i^K(s), x)$  using (3.2) to obtain

$$F_{i}(t \wedge \tau_{i}^{K}) - F_{i}(0) - K^{\frac{1}{2}} \int_{0}^{t \wedge \tau_{i}^{K}} \phi_{i}^{K}(s) p^{N}(t + \varepsilon - s, X_{i}^{K}(s), x) ds = M_{i,K}^{N,\varepsilon}(t, x), \qquad (3.3)$$

where  $M_{i,K}^{N,\varepsilon}(t,x) := \int_0^{t\wedge\tau_i^K} p^N(t+\varepsilon-s, X_i^K(s), x) dB_i^K(s)$ . With  $\Psi(s,0,x) = 0$ , we have  $F_i(t\wedge\tau_i^K) = \Psi(\varepsilon, X_i^K(t), x)$ . Using this in (3.3), summing the result over *i*, and dividing both sides by  $K^{1/2}$ , we arrive at

$$\langle \mu_{t,K}, \Psi(\varepsilon, \bullet, x) \rangle = \tilde{G}_K(t + \varepsilon, x)$$
  
 
$$+ \sum_{i=1}^K \int_0^t \phi_i^K(s) p^{\mathcal{N}}(t + \varepsilon - s, X_i^K(s), x) ds + M_K^{\mathcal{N},\varepsilon}(t, x),$$
 (3.4)

where  $\widetilde{G}_K(t,x)$  is defined as in (2.3) and  $M_K^{N,\varepsilon}(t,x) := K^{-1/2} \sum_{i=1}^K M_{i,K}^{N,\varepsilon}(t,x)$ .

With  $\lim_{\varepsilon \to 0} \langle \mu_{t,K}, \Psi(\varepsilon, \cdot, x) \rangle = \widetilde{U}_K(t, x)$ , it is natural to let  $\varepsilon \to 0$  here, which, however, introduces undesired discontinuity in  $M_K^{N,\varepsilon}(t, x)$ . Instead, we simply fix  $\varepsilon := K^{-1}$  and, adopting hereafter the notation  $t_K := t + \frac{1}{K}$ , define the discrepancies introduced by the shift of  $\varepsilon = K^{-1}$  as

$$E_K(t,x) := \langle \mu_{t,K}, (\Psi(K^{-1}, \bullet, x) - \Psi(0, \bullet, x)) \rangle, \qquad (3.5)$$

$$Q_K^{\rm N}(t,x) := \sum_{i=1}^{K} \int_0^t \phi_i^K(s) \left( p^{\rm N}(t_K - s, X_i^K(s), x) - p^{\rm N}(t - s, X_i^K(s), x) \right) ds, \qquad (3.6)$$

with the convention  $\Psi(0, y, x) := \mathbf{1}_{[x,\infty)}(y) - \mathbf{1}_{(-\infty,-x]}(y)$ . Defining further the corresponding martingale term

$$M_K^{\rm N}(t,x) := K^{-\frac{1}{2}} \sum_{i=1}^K \int_0^{t \wedge \tau_i^K} p^{\rm N}(t_K - s, X_i^K(s), x) dB_i^K(s), \qquad (3.7)$$

from (3.4) we thus obtain (2.4) for

$$R_{K}(t,x) := (\widetilde{G}_{K}(t_{K},x) - \widetilde{G}_{K}(t,x)) - E_{K}(t,x) + Q_{K}^{N}(t,x) + M_{K}^{N}(t,x).$$
(3.8)

Similarly, for (2.17), applying the preceding argument to the test function  $\Phi(t-s+\varepsilon, y-x)$ , we have

$$\langle \mu_{t,K}^{Y}, \Phi(\varepsilon, x - \bullet) \rangle = \langle \mu_{t_{\text{in}},K}^{Y}, \Phi(t - t_{\text{in}} + \varepsilon, x - \bullet) \rangle - \int_{t_{\text{in}}}^{t} p(t + \varepsilon - s, Z_{K}^{Y}(s), x) ds - M_{K}^{\varepsilon}(t, x), \quad \forall x \in \mathbb{R},$$
(3.9)

where  $M_K^{\varepsilon}(t,x) := K^{-1/2} \sum_i \int_{t_{in}}^t p(t-s+\varepsilon, Y_i^K(s)-x) dB_i^K(s)$ . Setting  $\varepsilon = K^{-1}$ , we thus obtain (2.17) for

$$R_K^Y(t,x) := E_K^Y(0,t,x) - E_K^Y(t-t_{\rm in},t_{\rm in},x) - Q_K(t_{\rm in},t,x) - M_K(t,x), \qquad (3.10)$$

where

$$E_K^Y(t,s,x) := \langle \mu_{s,K}^Y, \Phi(t_K,x-\bullet) - \Phi(t,x-\bullet) \rangle, \qquad (3.11)$$

$$Q_K(t_{\rm in}, t, x) := \int_{t_{\rm in}}^t \left( p(t_K - s, x - Z_K^Y(s)) - p(t - s, x - Z_K^Y(s)) \right) ds, \tag{3.12}$$

$$M_K(t,x) := K^{-\frac{1}{2}} \sum_{i=1}^K \int_{t_{\rm in}}^t p(t_K - s, x - Y_i^K(s)) dB_i^K(s).$$
(3.13)

It now remains only to show

**Lemma 3.1.** The remainder terms,  $R_K(t, x)$  and  $R_K^Y(t, x)$ , given as in (3.8) and (3.10), satisfy the respective bounds (2.5) and (2.18).

Proving Lemma 3.1 requires bounds on the empirical measures. Recall the definition of  $\mu_{t,K}^{Y}$  from (2.14).

**Lemma 3.2.** Fixing  $n, T < \infty$  and  $\alpha \in (0, 1)$ , we let  $\mu_{t,K}^Y$  and  $t_{in} < \infty$  be as in Proposition 2.6, we have

$$\|\mu_{t_{in}+s,K}^{Y}([a,b])\|_{n} \le C_{D}|b-a|^{\alpha} \left( \left(\frac{|b-a|}{\sqrt{s_{K}}}\right)^{1-\alpha} + 1 \right), \qquad \forall K^{-\frac{1}{2}} \le |b-a| \le 1, \ s \le T, \quad (3.14)$$

$$\|\langle \mu_{t_{in}+s,K}^{Y}, p(t_{K}, \cdot -x) \rangle\|_{n} \le C_{D} t_{K}^{\frac{\alpha-1}{2}} \left( \left(\frac{t_{K}}{s_{K}}\right)^{\frac{1-\alpha}{2}} + 1 \right), \quad \forall x \in \mathbb{R}, \ s, t < T,$$
(3.15)

where  $C_D < \infty$  dependence only on  $T, \alpha, n, D_*$  and  $D_{\alpha,n}$  as in (2.15)–(2.16).

Proof. Fixing such  $T, n, \alpha$  and [a, b], we let  $C_D = C(T, n, \alpha, D_*, D_{\alpha,1}, \ldots, D_{\alpha,n}) < \infty$  denote a generic constant as in the preceding. Defining the extend interval  $[a', b'] \supset [a, b]$ , where  $a' := a - K^{-1/2}$  and  $b' := b + K^{-1/2}$ , with  $|b-a| \ge K^{-1/2}$  we have  $\Phi(K^{-1}, b'-y) - \Phi(K^{-1}, a'-y) \ge \frac{1}{C} \mathbf{1}_{[a,b]}(y), \forall y \ge 0$ . With this, we set  $\varepsilon := K^{-1}$  in (3.9) and take the difference of the resulting equation for x = a' and for x = b' to obtain

$$\mu_{t_{\text{in}}+s,K}^{Y}([a,b]) \le C(|J_1| + |J_2| + |J_3|), \tag{3.16}$$

where  $J_1 := \int_0^s \left( p(s_K - s', a' - Z_K^Y(t_{\rm in} + s')) - p(s_K - s', b' - Z_K^Y(t_{\rm in} + s')) \right) ds',$ 

$$J_2 := \langle \mu_{t_{\text{in},K}}^Y, \Phi(s_K, b' - \bullet) - \Phi(s_K, a' - \bullet) \rangle = \int_{a'}^{b'} \langle \mu_{t_{\text{in},K}}^Y, p(s_K, x - \bullet) \rangle dx, \qquad (3.17)$$

and  $J_3 := M_K(t_{in} + s, b') - M_K(t_{in} + s, a')$ . Define  $h(a, b; s) := |b - a|^{\alpha} ((|b - a|/\sqrt{s_K})^{1-\alpha} + 1)$ . For  $J_1$ , by (2.8) we have

$$|J_1| \le C|b - a|^{\alpha} \le Ch(a, b; s).$$
(3.18)

To bound  $J_2$  and  $J_3$ , we first derive bounds on  $f_n(s,t) := \sup_{x \in \mathbb{R}} \|\langle \mu_{t_{in}+s,K}^Y, p(t_K, \cdot -x) \rangle\|_n$ . To this end, letting

$$\nu_{*,n} := \sup\left\{ \|\mu_{s+t_{\text{in}},K}^{Y}([a,b])\|_{n}(h(a,b;s))^{-1} : |b-a| \ge K^{-\frac{1}{2}}, \ s \le T \right\},$$
(3.19)

we fix  $x \in \mathbb{R}$  and partition  $\mathbb{R}$  into intervals  $I_j := x + [\sqrt{t_K j}, \sqrt{t_K (j+1)}), j \in \mathbb{Z}$ , each of length  $\sqrt{t_K} \ge K^{-1/2}$ . As the heat kernel  $p(t_K, y - x) = t_K^{-1/2} p(1, (y - x)/\sqrt{t_K})$  decreases in |y - x|, we have

$$\langle \mu_{t_{\text{in}}+s,K}^{Y}, p(t_{K}, \cdot - x) \rangle \leq \sum_{j \in \mathbb{Z}} \frac{1}{\sqrt{t_{K}}} p(1, j_{*}) \mu_{t_{\text{in}}+s,K}^{Y}(I_{j}), \quad j_{*} := |j| \land |j+1|.$$

Taking the  $L^n$ -norm of both sides using the triangle inequality, with  $\nu_{*,n}$  as in (3.19),  $D_{\alpha,n}$  as in (2.16), and  $\sum_{j} p(1, j_*) < \infty$ , we have

$$f_n(s,t) \le C\nu_{*,n} t_K^{\frac{\alpha-1}{2}} \big( (t_K/s_K)^{\frac{1-\alpha}{2}} + 1 \big), \tag{3.20}$$

$$f_n(0,t) \le CD_{\alpha,n} t_K^{\frac{\alpha-1}{2}}.$$
 (3.21)

Now, with  $J_2$  as in (3.17), by (3.21) we clearly have

$$||J_2||_n \le C|b-a|s_K^{\frac{\alpha-1}{2}} \le Ch(a,b;s).$$
(3.22)

As for  $J_3$ , with  $M_K(t_{in} + s, x)$  defined as in the proceeding,  $J_3$  is a martingale integral of quadratic variation  $K^{-1/2} \int_0^s \langle \mu_{s,K}^Y, q(s', \cdot) \rangle ds'$ , where  $q(s', y) := p(s_K - s', a' - y) - p(s_K - s', b' - y)$ . Hence, by the Burkholder–Davis–Gundy (BDG) inequality we have

$$\|J_3\|_j^2 \le C_D K^{-\frac{1}{2}} \int_0^s \|\langle \mu_{t_{\text{in}}+s',K}^Y, q^2(s', \cdot) \rangle\|_{j/2} ds',$$
  
$$\le C_D K^{-\frac{1}{2}} \int_0^s |q(s', \cdot)|_{\mathcal{C}(\mathbb{R})} f_{j-1}(s', s-s') ds', \quad j=2,\dots,n,$$
(3.23)

where in the last inequality we used  $f_{j/2}(s,t) \leq f_{j-1}(s,t)$ . Fixing  $\beta := ((2\alpha - 1)_+, \alpha)$ , by (2.8) we have  $|q(s', \cdot)|_{\mathcal{C}(\mathbb{R})} \leq C|b-a|^{\beta}(s_K-s')^{-\frac{1+\beta}{2}}$ . Using this and (3.20) in (3.23), we thus arrive at

$$\|J_3\|_j^2 \le CK^{-\frac{1}{2}} |b'-a'|^\beta \nu_{*,j-1} \int_0^s \left( (s_K - s')^{-\frac{1+\beta}{2}} s_K'^{\frac{\alpha-1}{2}} + (s_K - s')^{-1+\frac{\alpha-\beta}{2}} \right) ds'.$$

With  $-1 + \frac{\alpha - \beta}{2} > -1$  and  $K^{-1/2} \le K^{-(2\alpha - \beta)/2} \le |b - a|^{2\alpha - \beta}$ , we further obtain

$$\|J_3\|_j^2 \le C|b-a|^{2\alpha} s_K^{\frac{\alpha-\beta}{2}} \nu_{*,j-1} \le Ch(a,b;s)^2 \nu_{*,j-1}.$$
(3.24)

Now, combining the preceding bounds (3.18), (3.22) and (3.24) on  $J_1, J_2, J_3$  with (3.16), we obtain

$$\|\mu_{s,K}^{Y}([a,b])\|_{j} \le C_{D}h(a,b;s)(1+(\nu_{*,j-1})^{\frac{1}{2}}),$$
(3.25)

for j = 2, ..., n. As this holds for all  $|b - a| \ge K^{-1/2}$ , we further obtain  $\nu_{*,j} \le C_D(1 + (\nu_{*,j-1})^{\frac{1}{2}})$ , for j = 2, ..., n. Consequently (3.14) follows once we prove it for n = 1. The latter, with  $\mathbf{E}(J_3) = 0$  and the bounds (3.22) and (3.18), follows by taking expectation in (3.16).

Next we establish bounds on  $\mu_{t,K}$ , defined as in (3.1).

**Lemma 3.3.** Let  $n, T < \infty$  and  $\alpha \in (0, 1)$ . Given any strategy,

$$\|\mu_{s,K}([a,b])\|_{n} \le C|b-a|^{\alpha} s_{K}^{-\frac{1+\alpha}{2}}, \qquad \forall |b-a| \ge K^{-\frac{1}{2}}, \ s \le T,$$
(3.26)

$$\|\langle \mu_{s,K}, p(t_K, \cdot - x) \rangle\|_n \le C t_K^{-\frac{1-\alpha}{2}} s_K^{-\frac{1+\alpha}{2}}, \qquad \forall x \in \mathbb{R}, \ s, t \le T,$$
(3.27)

where  $C = C(T, \alpha, n) < \infty$  is independent of the strategy.

*Proof.* The proof is similar to that of Lemma 3.2, by using (3.4) for  $\varepsilon = K^{-1}$ . More precisely, letting

$$\mu_{*,n} := \sup \left\{ \|\mu_s([a,b])\|_n |b-a|^{-\alpha} s_K^{\frac{1+\alpha}{2}} : |b-a| \ge K^{-\frac{1}{2}}, s \le T \right\},\$$

following the same argument as we obtained (3.20) and (3.25), here we obtain that

$$\|\langle \mu_{s,K}, p(t_K, \bullet - x) \rangle\|_n \le C \mu_{*,n} t_K^{\frac{\alpha - 1}{2}} s_K^{-\frac{1 + \alpha}{2}},$$
(3.28)

$$\|\mu_{s,K}([a,b])\|_{j} \le C(\Psi(K^{-1}, \bullet, b') - \Psi(K^{-1}, \bullet, a')) \le C|b-a|^{\alpha} s_{K}^{-\frac{1+\alpha}{2}} (1 + (\mu_{*,j-1})^{\frac{1}{2}}), \quad (3.29)$$

where  $a' := a - K^{-1/2}$  and  $b' := b + K^{-1/2}$ . However, unlike (3.25), the inequality (3.29) holds true only for  $[a', b'] \subset \mathbb{R}_+$ . Consequently, intervals [a, b] with  $a \in [0, K^{-1/2})$  are left out, and hence (3.29) cannot be closed. To circumvent this problem, for  $j = 1, \ldots, n$ , we consider the modified process  $(X_i^{K,j}(t))_i$ , where particles are absorbed at level  $x = (j - n)K^{-1/2}$ , instead of x = 0. Let  $\mu_{s,K}^j$  denote the corresponding empirical measure of  $(X_i^{K,j}(t))_i$ . Under the natural coupling, we clearly have  $\mu_{s,K}^1(\cdot) \ge \mu_{s,K}^2(\cdot) \ge \ldots \ge \mu_{s,K}^n(\cdot) = \mu_{s,K}(\cdot)$ . With this, letting

$$\mu_{*,j'}^j := \sup \left\{ \|\mu_s^j([a,b])\|_{j'} |b-a|^{-\alpha} s_K^{\frac{1+\alpha}{2}} : |b-a| \ge K^{-\frac{1}{2}}, s \le T \right\},\$$

similar to (3.29) we have

$$\begin{aligned} \|\mu_{s,K}^{j}([a,b])\|_{j} &\leq C|b-a|^{\alpha} s_{K}^{-\frac{1+\alpha}{2}} (1+(\mu_{*,j-1}^{j-1})^{\frac{1}{2}}), \quad j=2,\dots,n, \end{aligned}$$
(3.30)  
$$\mathbf{E} \left(\mu_{s,K}^{1}([a,b])\right) &\leq C|b-a|^{\alpha} s_{K}^{-\frac{1+\alpha}{2}}, \end{aligned}$$

for all  $[a,b] \subset \mathbb{R}_+ + (j-n)K^{-1/2}$  with  $|b-a| \geq K^{-1/2}$ . The difference between (3.30) and (3.28) is that here, for each  $j = 2, \ldots, n$ , we bound the relevant terms in terms of  $\mu_{t,K}^{j-1}$ , not in terms of  $\mu_{t,K}^{j}$  itself. Consequently, (3.30) holds even for  $a \in [(j-n)K^{-1/2}, (j-n+1)K^{-1/2})$ , and hence

$$\mu_{*,j}^{j} \leq C(1 + (\mu_{*,j-1}^{j-1})^{\frac{1}{2}}), \ j = 2, \dots, n, \quad \mu_{*,1}^{1} \leq C.$$

From this the desired results follow.

**Lemma 3.4.** Let  $T, L, n < \infty$  and

$$I_{x,\alpha} := [-K^{-\alpha} + x, x + K^{-\alpha}].$$
(3.31)

(a) For any given  $\gamma \in (0, \frac{1}{4})$ ,  $\alpha \in (\frac{1}{2} - 2\gamma, \frac{1}{2}]$  and any strategy,

$$\mathbf{P}\left(\mu_{t,K}(I_{x,\alpha}) \le t^{-\frac{3}{4}} K^{-\frac{1}{4}+\gamma}, \ \forall t \le T, |x| \le L\right) \ge 1 - CK^{-n},\tag{3.32}$$

where  $C = C(T, \alpha, \gamma, n) < \infty$  is independent of the strategy.

(b) Letting  $\mu_{t,K}^Y$  and  $t_{in} < \infty$  be as in Proposition 2.6, we have, for any  $\alpha \in (\frac{1}{4}, \frac{1}{2}]$ ,

$$\mathbf{P}\left(\mu_{t,K}^{Y}(I_{x,\alpha}) \le K^{-\frac{1}{4}}, \ \forall t_{in} \le t \le T, |x| \le L\right) \ge 1 - C_D K^{-n}, \tag{3.33}$$

where  $C_D < \infty$  dependence only on  $T, \alpha, n$  and  $D_*, D_{\alpha,j}, j \in \mathbb{Z}$ , as in (2.15)-(2.16).

Proof. We first prove Part (b). Fixing  $L, T, n < \infty$ ,  $t_{\text{in}} \leq T$  and  $\alpha \in (\frac{1}{4}, \frac{1}{2}]$ , we use C(k) to denote generic finite constants that may depend on  $L, T, \alpha, n, D_*, D_{\alpha,j}$  and the designated variable k. To the end of proving (3.33), we cover [-L, L] by intervals  $I_j := [jK^{-\alpha}, (j + 1)K^{-\alpha})$  of length  $K^{-\alpha}$ ,  $|j| \leq LK^{\alpha}$ . Since each  $I_{x,\alpha}$  is contained in the union of three consecutive such intervals  $I_j$ , it suffices to prove

$$\mathbf{P}(\mu_{t,K}^{Y}(I_{j}) \leq \frac{1}{3}K^{-\frac{1}{4}}, \ \forall |j| \leq LK^{\alpha}, t_{\text{in}} \leq t \leq T) \geq 1 - CK^{-n}.$$
(3.34)

By (3.14) we have, for all  $t \in [t_{in} + K^{-1}, T]$ ,

$$\|\mu_{t,K}^{Y}(I_{j})\|_{k} \leq C(k,\beta)|I_{j}|^{\beta} \left( (|I_{j}|K^{1/2})^{1-\beta} + 1 \right) \leq C(k,\beta)(K^{-\alpha + \frac{1-\beta}{2}} + K^{-\alpha\beta}),$$
(3.35)

for any fixed  $\beta \in (0,1)$  and  $k < \infty$ . With  $\alpha > \frac{1}{4}$ , fixing  $\beta$  close enough to 1 we have  $\|\mu_{t,K}^Y(I_j)\|_k \leq C(k)K^{-\frac{1}{4}-\varepsilon}$ , for some fixed  $\varepsilon > 0$ . With this, applying Markov's inequality we obtain

$$\mathbf{P}(\mu_{t,K}^{Y}(I_j) \ge \frac{1}{9}K^{-\frac{1}{4}}) \le C(k)K^{-k\varepsilon}.$$
(3.36)

Now, fixing  $k \ge (n + \alpha + 2)\varepsilon^{-1}$  and taking the union bound of (3.36) over  $|j| \le LK^{\alpha}$  and  $t = t_{\ell} := \ell K^{-2} + t_{\text{in}}, 1 \le \ell \le TK^2$ , we arrive at

$$\mathbf{P}(\mu_{t_{\ell},K}^{Y}(I_{j}) \leq \frac{1}{9}K^{-\frac{1}{4}}, \ \forall |j| \leq LK^{\alpha}, 1 \leq \ell \leq TK^{2}) \geq 1 - CK^{-n}.$$
(3.37)

To move from the 'discrete time'  $t_{\ell}$  to 'continuous time'  $t \in [t_{\text{in}}, T]$ , we need to control  $\mu_{s,K}^{Y}(I_j)$  for  $s \in [t_{\ell-1}, t_{\ell}] := J_{\ell}$ . Within each  $J_{\ell}$ , since each  $Y_i^K(s)$  particle evolves as a drifted Brownian motion (with drift  $\leq K^{1/2}$ ), we have that

$$\mathbf{P}(|Y_i^K(s) - Y_i^K(t_\ell)| \le K^{-\alpha}, \forall s \in J_\ell) \ge 1 - \exp(-\frac{1}{C}K^{1-\alpha}) \ge 1 - CK^{-n-3}.$$
(3.38)

Further taking the union bound of this over  $\ell \leq TK^{-2}$  and over all particles  $i = 1, \ldots$ , with (2.15), we obtain

$$\mathbf{P}\left(\sup_{s\in J_{\ell}}|Y_{i}^{K}(s)-Y_{i}^{K}(t_{\ell})|\leq K^{-\alpha}, \ \forall i,\forall 1\leq\ell\leq TK^{2}\right)\geq1-CK^{-n}$$

That is, with high probability, no particle travels farther than distance  $|I_j|$  within each time interval  $J_{\ell}$ , and therefore  $\mu_{s,K}^Y(I_j) \leq \mu_{t_{\ell},K}^Y(I_{j-1} \cup I_j \cup I_{j+1}), \forall s \in J_{\ell}$ . We thus conclude (3.34) from (3.37).

Part (a) is proven by similar argument as in the preceding. The only difference is that, instead of a moment bound of the form (3.35), we have from (3.26) the moment bound

$$\|\mu_{t,K}(I_j)\|_k \le C(k)|I_j|^{\frac{1}{2}}t^{-\frac{3}{4}} \le C(k)K^{-\frac{\alpha}{2}}t^{-\frac{3}{4}},\tag{3.39}$$

for all  $k < \infty$ . With  $\frac{\alpha}{2} > \frac{1}{4} - \gamma$ , (3.39) yields (3.32) by same argument we obtained (3.35).  $\Box$ 

Proof of Lemma 3.1. Fixing arbitrary  $t_{in} \leq T < \infty$  and  $\gamma \in (0, \frac{1}{4})$ , we let  $C(k) < \infty$  denote a generic constant depending only on  $T, n, \alpha, \gamma$  and the designated variable k, and  $C_D(k) < \infty$  denote a constant that may further depend  $D_*, D_{\alpha,j}, j = 1, \ldots, n$ .

We first claim that, to the end of proving (2.5) and (2.18), it suffices to prove

$$\mathbf{P}\left(|R_K(t,x)| \le K^{-\frac{1}{4}+\gamma} t^{-\frac{3}{4}}, \ \forall t \le T, x \in [a,a+1]\right) \ge 1 - CK^{-n},\tag{3.40}$$

$$\mathbf{P}\left(|R_{K}^{Y}(t_{\rm in}, t, x)| \le K^{-\frac{1}{4}+\gamma}, \ \forall t_{\rm in} \le t \le T, x \in [a, a+1]\right) \ge 1 - C_{D}K^{-n},\tag{3.41}$$

for arbitrary  $a \in \mathbb{R}$ . Indeed, taking the union bound of (3.40). over  $a \in \mathbb{Z} \cap [-K, K]$  yields

$$\mathbf{P}\left(|R_K(t,x)| \le K^{-\frac{1}{4}+\gamma}t^{-\frac{3}{4}}, \ \forall t \le T, |x| \le K\right) \ge 1 - CK^{-n+1}.$$
(3.42)

Further, since each  $X_i^K(t)$  evolves as a Brownian motion with drift at most  $K^{-1/2}$  and absorption, the event  $\Omega_K := \{|X_i^K(t)| \leq \frac{1}{2}K, \forall t \leq T, \forall i\}$  holds with probability  $1 - CK^{-n}$ . On  $\Omega_K$  we have

$$\widetilde{U}_K(t,x) = \widetilde{U}_K(t,K), \quad \forall x \ge K, \qquad \widetilde{U}_K(t,-x) = \widetilde{U}_K(t,-x) = 0, \quad \forall x < 0,$$

$$\left|\int_0^t p^{\mathcal{N}}(t-s, Z_K(s), \pm x)ds - \int_0^t p^{\mathcal{N}}(t-s, Z_K(s), \pm K)ds\right| \le CK^{-1}, \quad \forall x \ge K.$$

Now, expressing R(t,x) as  $\widetilde{U}_K(t,x) - \widetilde{G}_K(t,x) - \int_0^t p^N(t-s, Z_K(s), x) ds$ , we then conclude that  $\sup_{x \in \mathbb{R}} |R_K(t,x)| \leq \sup_{|x| \leq K} |R_K(t,x)| + CK^{-1}$ . Combining this with (3.42) yields the desired bound (2.5). A similar argument shows that (3.41) implies (2.18).

We now proceed to proving (3.40)–(3.41), which amounts to bounding each term on the r.h.s. of (3.8) and (3.10). To this end, fixing  $t \leq T$  and  $x \in [a, a + 1]$ , we next establish the desired bounds on

- i)  $|\widetilde{G}_{K}(t_{K}, x) \widetilde{G}_{K}(t, x)|;$ ii)  $Q_{K}^{N}(t, x)$  and  $Q_{K}(t_{\text{in}}, t, x);$ iii)  $E_{K}(t, x), E_{K}^{Y}(0, t, x)$  and  $E_{K}^{Y}(t t_{\text{in}}, t, x);$  and iv)  $M_{K}(t, x)$  and  $M_{K}^{N}(t_{\text{in}}, t, x).$
- (i) With  $\widetilde{G}_K(t,x)$  defined as in (2.3), by (2.9) for  $\alpha' = \frac{1}{4}$  we have

$$|\tilde{G}_K(t_K, x) - \tilde{G}_K(t, x)| \le CK^{-\frac{1}{4}}t^{-\frac{3}{4}}.$$
(3.43)

(*ii*) Applying (2.9) for  $\alpha = \frac{1}{2}$  in (3.6) and in (3.12) yields

$$|Q_K^{\rm N}(t,x)|, \ |Q_K(t_{\rm in},t,x)| \le CK^{-\frac{1}{4}}.$$
 (3.44)

(*iii*) Let  $I_{x,\alpha}$  be as in (3.31). As  $t \mapsto |\Phi(t_K, y) - \Phi(t, y)|$  decreases in t and  $\Phi(K^{-1}, -|y|) \leq 1$  $C \exp(-K^{1/2}|y|)$ , we have

$$|\Phi(t_K, y - x) - \Phi(t, y - x)| \le C(\mathbf{1}_{I_{x,\alpha}}(y) + \exp(-K^{\alpha - \frac{1}{2}})),$$
(3.45)

$$|\Psi(\frac{1}{K}, y, x) - \Psi(0, y, x)| \le C(\mathbf{1}_{I_{x,\alpha} \cup I_{-x,\alpha}}(y) + \exp(-K^{\alpha - \frac{1}{2}})),$$
(3.46)

for fixed  $\alpha \in (\frac{1}{2} - 2\gamma, \frac{1}{2})$ . Now, cover  $[-a - 1, -a] \cup [a, a + 1]$  by intervals of the form  $I_{x',\alpha}$ , with  $x' \in \mathcal{X} := \{\pm (a + 2jK^{-\alpha}) : 0 \le j \le \frac{1}{2}K^{\alpha}\}$ . We then have

$$\mathbf{1}_{I_{x,\alpha}}(y), \ \mathbf{1}_{I_{x,\alpha}\cup I_{-x,\alpha}}(y) \le 4 \max_{x'\in\mathcal{X}} \left\{ \mathbf{1}_{I_{x',\alpha}}(y) \right\}.$$

Combining this with (3.45)-(3.46), and inserting the result into (3.5) and (3.11) using  $|\{X_i^K(t)\}_i| \leq K$  and (2.15), we arrive at

$$|E_{K}(t,x)| \le C \max_{x' \in \mathcal{X}} \left\{ \mu_{t,K}(\mathbf{1}_{I_{x',\alpha}}) \right\} + CK \exp(-K^{\alpha - \frac{1}{2}})),$$

 $|E_K^Y(t - t_{\rm in}, t, x)| \le C \max_{x' \in \mathcal{X}} \left\{ \mu_{t,K}^Y(\mathbf{1}_{I_{x',\alpha}}) \right\} + CK \exp(-K^{\alpha - \frac{1}{2}})), \text{ with probability } 1 - C_D K^{-n}.$ 

Further using Lemma 3.4 to bound the first terms on the r.h.s., we thus obtain

$$\mathbf{P}(|E_K(t,x)| \le CK^{-\frac{1}{4}+\gamma}t^{-\frac{3}{4}}, \ \forall t \le T, x \in [a,a+1]) \ge 1 - CK^{-n},$$
(3.47)

$$\mathbf{P}(|E_K^Y(t - t_{\rm in}, t, x)| \le C_D K^{-\frac{1}{4}}, \ \forall t_{\rm in} \le t \le T, x \in [a, a+1]) \ge 1 - C_D K^{-n}.$$
(3.48)

(*iv*) We recall from [Kun97] the following Kolmogorov-type estimate.

**Lemma 3.5** ([Kun97, Theorem 1.4.1]). Let  $t_{in} < T < \infty$ ,  $a \in \mathbb{R}$ , and let F be a  $\mathcal{C}([t_{in}, \infty) \times$  $\mathbb{R}$ )-valued process. If, for some  $\alpha_1, \alpha_2, k \in \mathbb{N}$  and  $C_1 < \infty$  with  $\frac{1}{k\alpha_1} + \frac{1}{k\alpha_2} < 1$ ,

$$\|F(t_{in},0)\|_k \le C_1,\tag{3.49}$$

$$||F(t,x) - F(t',x')||_k \le C_1(|t - t'|^{\alpha_1} + |x - x'|^{\alpha_2}),$$
(3.50)

$$\forall t, t' \in [t_{in}, T], \ x, x' \in [a, a+1], \ then \ \left\| |F|_{\mathcal{C}([t_{in}, T] \times [a, a+1])} \right\|_k \le C_2 = C_2(C_1, T, \alpha_1, \alpha_2) < \infty.$$

Note that, although the dependence of  $C_2$  is not explicitly designated in [Kun97, Theorem 1.4.1], under the present setting, it is clear from the proof of [Kun97, Lemma 1.4.2, Lemma 1.4.3] that  $C_2 = C_2(C_1, T, \alpha_1, \alpha_2, k)$ .

Our goal is to apply Lemma 3.5 for  $F(t,x) := K^{1/4}M_K(t,x)$ . For such F we have  $F(t_{\rm in}, 0) = 0$ , whereby (3.49) follows. Turning to verifying (3.50), we fix t < t' and  $x, x' \in \mathbb{R}$ . With  $M_K(t,x)$  defined as in (3.13), we telescope F(t,x) - F(t',x') into  $F_1 + F_2 - F_3$ , where

$$\begin{split} F_1 &:= K^{-1/4} \sum_i \int_{t_{\rm in}}^t f_1(s, Y_i^K(s)) dB_i^K(s), \qquad F_2 := K^{-1/4} \sum_i \int_{t_{\rm in}}^t f_2(s, Y_i^K(s)) dB_i^K(s), \\ F_3 &:= K^{-1/4} \sum_i \int_t^{t'} f_3(s, Y_i^K(s)) dB_i^K(s), \end{split}$$

 $f_1(s,y) := p(t_K - s, y - x) - p(t_K - s, y - x'), f_2(s,y) := p(t_K - s, y - x') - p(t'_K - s, y - x')$ and  $f_3(s'', y) := p(t'_K - s, y - x')$ . Similar to the way we obtained (3.23), here by the BDG inequality we have

$$\begin{split} \|F_1\|_k^2 &\leq C(k) \int_{t_{\rm in}}^t \|\langle \mu_{s,K}^Y, f_1(s, \cdot)^2 \rangle\|_{k/2} ds, \quad \|F_2\|_k^2 \leq C(k) \int_{t_{\rm in}}^t \|\langle \mu_{s,K}^Y, f_2(s, \cdot)^2 \rangle\|_{k/2} ds, \\ \|F_3\|_k^2 &\leq C(k) \int_t^{t'} \|\langle \mu_{s,K}^Y, f_3(s, \cdot)^2 \rangle\|_{k/2} ds, \end{split}$$

for all k > 1. On the r.h.s., the kernel functions  $f_1, f_2, f_3$  appear in square (i.e. power of two). We use (2.8)–(2.9) to replace 'one power' of them with  $C(t-s)^{-\frac{3}{4}}|x-x'|^{\frac{1}{2}}$ ,  $C(t-s)^{-\frac{3}{4}}|t-t'|^{\frac{1}{4}}$ and  $C(t-s)^{-\frac{1}{2}}$ , respectively, and then use (3.15) for  $\alpha = \frac{3}{4}$  to bound the remaining expression  $\|\langle \mu_{s,K}^Y, f_j(s, \cdot) \rangle\|_{n/2}$ , j = 1, 2, 3, whereby obtaining

$$\begin{aligned} \|F_1\|_k^2 &\leq C_D(k) \int_{t_{\rm in}}^t |x - x'|^{\frac{1}{2}} ((t - s)^{-\frac{7}{8}} + (t - s)^{-\frac{3}{4}} s^{-\frac{1}{8}}) ds \leq C_D(k) |x - x'|^{\frac{1}{2}}, \\ \|F_2\|_k^2 &\leq C_D(k) \int_{t_{\rm in}}^t |t - t'|^{\frac{1}{4}} ((t - s)^{-\frac{7}{8}} + (t - s)^{-\frac{3}{4}} s^{-\frac{1}{8}}) ds \leq C_D(k) |t - t'|^{\frac{1}{4}}, \\ \|F_3\|_k^2 &\leq C_D(k) \int_t^{t'} ((t - s)^{-\frac{5}{8}} + (t - s)^{-\frac{1}{2}} s^{-\frac{1}{8}}) ds \leq C_D(k) |t - t'|^{\frac{3}{8}}. \end{aligned}$$

We thus verify (3.50) for  $(\alpha_1, \alpha_2) = (\frac{1}{8}, \frac{1}{4})$ , whereby  $|||M_K|_{\mathcal{C}([0,T]\times[a,a+1])}||_n \leq C_D(n)K^{-\frac{1}{4}}$ . From this and Markov's inequality, we conclude

$$\mathbf{P}\left(|M_K|_{\mathcal{C}([0,T]\times[a,a+1])} \le K^{-\frac{1}{4}+\gamma}\right) \ge 1 - C_D K^{-n\gamma}.$$
(3.51)

The term  $M_K^N(t, x)$  is bounded by similar procedures as in the preceding. The only difference is that the estimate (3.27), unlike (3.15), introduces a singularity of  $M_K^N(t, x)$  as  $t \to 0$ , so we set  $F(t, x) := t^{\frac{3}{4}} K^{1/4} M_K^N(t, x)$  (instead of  $F(t, x) := K^{1/4} M_K^N(t, x)$ ). The extra prefactor  $t^{\frac{3}{4}}$  preserves the moment estimate (3.50) since  $t^{\frac{3}{4}}$  is  $\alpha$ -Hölder continuous for all  $\alpha < \frac{3}{4}$ . Consequently, following the preceding argument we obtain

$$\mathbf{P}\left(\sup_{t\in[0,T],x\in[a,a+1]}|M_{K}^{N}(t,x)| \le t^{-\frac{3}{4}}K^{-\frac{1}{4}+\gamma}\right) \ge 1 - CK^{-n\gamma}.$$
(3.52)

Now, combining the bounds (3.43), (3.47), (3.51)–(3.52) from (i)–(iv), we conclude the desired results (2.5) and (2.18).

### 4. The Stefan Problem

In this section, we develop the necessary PDE tools for the Stefan problem. Throughout this section, we fix  $u_{in} \in \mathcal{C}(\mathbb{R}_+) \cap L^1(\mathbb{R}_+)$  with  $0 < u_{in}(x) \leq 2, \forall x \geq 0$ , and let  $U_{in}(x) := \int_0^x u_{in}(y) dy$ . Note that the function  $U_{\star}(\frac{1}{2}, x) := \widetilde{U}_{\star}(\frac{1}{2}, 0) - \widetilde{U}_{\star}(\frac{1}{2}, x)$  in (1.9) indeed satisfies this condition (see (1.10)).

As mentioned in Remark 1.4, we take the integral equation (1.9), instead of (1.13), as the definition of the Stefan problem. To motivate such a definition, we first prove the following:

**Lemma 4.1.** For any classical solution (u, z) to the following PDE,

$$\partial_t u = \frac{1}{2} \partial_{xx} u, \quad \forall \ 0 < t < T, \ x > z(t), \tag{4.1a}$$

$$u(t, z(t)) = 2, \quad \forall t \ge 0, \tag{4.1b}$$

$$\frac{d}{dt}z(t) + \frac{1}{4}\partial_x u(t, z(t)) = 0, \quad \forall t > 0,$$

$$(4.1c)$$

$$u(0,x) = u_{in}(x), \quad \forall x \ge 0, \tag{4.1d}$$

$$z \in \mathcal{C}^1((0,\infty)) \cap \mathcal{C}([0,\infty)), \text{ nondecreasing, } z(0) = 0,$$

$$u \in \mathcal{C}^{2}(\mathcal{D}) \cap \mathcal{C}(\overline{\mathcal{D}}) \cap L^{\infty}(\overline{\mathcal{D}}), \text{ where } \mathcal{D} := \{(t, x) : t > 0, x \ge z(t)\},$$

 $U(t,x) := \int_{z(t)}^{x} u(t,y) dy$  satisfies

$$U(t,x) = \int_0^\infty p(t,x-y)U_{in}(y)dy - \int_0^t p(t-s,x-z(s))ds,$$
(4.2)

In particular, with U(t, z(t)) = 0, we have

$$\int_0^\infty p(t, z(t) - y) U_{in}(y) dy = \int_0^t p(t - s, z(t) - z(s)) ds.$$
(4.3)

*Proof.* By (4.1a), (4.1b)–(4.1c), U(s, y) solves the heat equation in  $\{(s, y) : s > 0, x > z(s)\}$ . With this, for any fixed t > 0 and  $x \in \mathbb{R}$ , we integrate Green's identity

$$\frac{1}{2}\partial_y((\partial_y p)U - p(\partial_y U)) + \partial_s(pU) = 0, \text{ where } p = p(t - s, x - y), \ U = U(s, y),$$

over  $\{(s, y) : \varepsilon < s < t - \varepsilon, y > z(s) + \varepsilon\}$ . Letting  $\varepsilon \to 0$ , and combining the result with U(s, z(s)) = 0 and (1.13d), we conclude (4.2).

Next, we establish the connection between the integral representations (4.2) and (1.8). Lemma 4.2. Given any solution  $z_{\star}$  to (1.9), for  $t \in (0, \infty)$  we define

$$U_{\star}(t+\frac{1}{2},x) := \int_{0}^{\infty} p(t,x-y)U_{\star}(\frac{1}{2},y)dy - \int_{0}^{t} p(t-s,x-z_{\star}(\frac{1}{2}+s))ds, \qquad (4.4)$$

where  $U_{\star}(\frac{1}{2}, x) := \widetilde{U}_{\star}(\frac{1}{2}, 0) - \widetilde{U}_{\star}(\frac{1}{2}, x)$  is as in the preceding. We then have that

$$U_{\star}(\frac{1}{2}+t,x) = \widetilde{U}_{\star}(\frac{1}{2}+t,0) - \widetilde{U}_{\star}(\frac{1}{2}+t,x), \ \forall x \ge 0, \ t \in \mathbb{R}_{+}.$$
(4.5)

In particular, since by (1.9)  $U_{\star}(s, z_{\star}(s)) = 0$ , we have  $\widetilde{U}_{\star}(s, z_{\star}(s)) = \widetilde{U}_{\star}(\frac{1}{2}, 0) = \frac{4}{\sqrt{\pi}}, \forall s \geq \frac{1}{2}$ .

Proof. From (4.4), we clearly have that  $U_{\star} \in \mathcal{C}([\frac{1}{2}, \infty) \times \mathbb{R}) \cap L^{\infty}([\frac{1}{2}, \infty) \times \mathbb{R})$  solves the heat equation in  $\{(s, x) : s > \frac{1}{2}, x < z_{\star}(s)\}$ . Combining this with  $U_{\star}(s, z_{\star}(s)) = 0, \forall s \ge \frac{1}{2}$  (by (1.9)) and  $U_{\star}(\frac{1}{2}, x) = 0, \forall x \le 0$ , we then conclude from the uniqueness of the heat equation that  $U_{\star}(s, x) = 0, \forall s \ge \frac{1}{2}, x \le z_{\star}(s)$ . In particular,

$$U_{\star}(\frac{1}{2}+t,x) = U_{\star}(\frac{1}{2}+t,x) + U_{\star}(\frac{1}{2}+t,-x)$$
  
=  $\int_{0}^{\infty} p^{N}(\frac{1}{2}+t,x-y)U_{\star}(\frac{1}{2},y)dy - \int_{0}^{t} p^{N}(t-s,z_{\star}(\frac{1}{2}+t),x)ds, \quad \forall x \in \mathbb{R}_{+}.$  (4.6)

The desired identity (4.5) now follows by comparing (4.6) and (2.1), using the semigroup property  $p^{N}(t+s, y, x) = \int_{0}^{\infty} p^{N}(t, y, z) p^{N}(t, z, x) dz$  of  $p^{N}$ .

Next we turn to the well-posedness of (4.3). The following lemma settles the existence of (4.3).

**Lemma 4.3.** Fixing  $T < \infty$  and  $U_{in}(x)$  as in the proceeding, there exists a nondecreasing  $z \in \mathcal{C}([0,T])$  satisfying (4.3) for all  $t \in [0,T]$ .

Postponing the proof of Lemma 4.3 until Section 6.1, we proceed to establishing the uniqueness, or more generally, the stability, of (4.3). More precisely, we consider  $w \in \mathcal{C}([0,T])$  that satisfies w(0) = 0 and

$$\int_{0}^{\infty} p(t, w(t) - y) U_{\rm in}(y) dy = f(t, w(t)) + \int_{0}^{t} p(t - s, w(t) - w(s)) ds, \tag{4.7}$$

where  $f \in L^{\infty}([0,T] \times \mathbb{R})$  is a generic perturbation. Define a seminorm

$$|w|'_{[0,T]} := \sup\{w(t) - w(t'), \ 0 \le t \le t' \le T\}$$
(4.8)

measuring how nondecreasing the given function is.

**Lemma 4.4.** Let  $U_{in}(x) = \int_0^x u_{in}(y) dy$  be as in the preceding. Fixing  $T < \infty$  and  $f_1(t, x), f_2(t, x) \in L^{\infty}([0, T] \times \mathbb{R})$ , we consider  $w_1$  and  $w_2$  satisfying (4.7) for  $f = f_1$  and  $f = f_2$ , respectively. Let  $L := \sup\{|w_1(t)|, |w_2(t)| : t \leq T\} + 1$ . There exists  $C_1 = C_1(T, L) < \infty$  such that

$$\sup_{0 \le t \le T} (w_1(t) - w_2(t)) \le C_1(|f_1 - f_2|_{\mathcal{C}([0,T] \times \mathbb{R})} + |w_2|'_{[0,T]}).$$

for all  $f_1, f_2 \in L^{\infty}([0,T] \times \mathbb{R})$  such that  $C_1(|f_1 - f_2|_{\mathcal{C}([0,T] \times \mathbb{R})} + |w_2|_{[0,T]}) \leq 1$ .

Indeed, when  $f_1 = f_2 = 0$ , Lemma 4.4 yields

**Corollary 4.5.** For any  $U_{in}$  as in the preceding, the solution to (4.3) is unique.

Proof of Lemma 4.4. Without loss of generality we assume  $T \ge 1$ . To simplify notations, let  $\varepsilon := |f_1 - f_2|_{\mathcal{C}([0,T]\times\mathbb{R})}$  and  $\varepsilon' := |w|'_{[0,T]}$ .

Let  $F(t,z) := \int_0^\infty p(t,z-y)U_{\rm in}(y)dy$  denote the r.h.s. of (4.2). With  $\partial_x U_{\rm in}(y) = u_{\rm in}(y) > 0$ ,  $\forall y \ge 0$ , we have  $\partial_z F(t,z) > 0$ ,  $\forall z \ge 0$ , so there exists  $c_1 = c_1(T,L) > 0$  such that

$$\partial_z F(t,z) \ge c_1 > 0, \quad \forall 0 \le z \le L, \ 0 \le t \le T.$$

$$(4.9)$$

Fixing  $C_1 := 4/c_1$ , we let  $\delta := C_1(\varepsilon + \varepsilon') \leq 1$ ,  $w_2^{\delta}(t) := w_2(t) + \delta$ , and  $t^* := \inf\{t \leq T : w_1(t) \geq w_2^{\delta}(t)\}$  be the first hitting time. Taking the difference of (4.7) for  $(t, f) = (t^*, f_1)$  and for  $(t, f) = (t^*, f_2)$ , we obtain

$$F(t^*, w_2^{\delta}(t^*)) - F(t^*, w_2(t^*)) \le \varepsilon + \int_0^{t^*} g^*(s) ds, \qquad (4.10)$$

where  $g^*(s) := p(t^* - s, w_1(t^*) - w_1(s)) - p(t^* - s, w_2(t^*) - w_2(s))$ . With  $w_1(t^*) = w_2(t^*) + \delta$ and  $w_1(s) - \delta \le w_2(s) \le w_2(t^*) + \varepsilon'$ ,  $\forall s \le t^*$ , we have  $w_1(t^*) - w_1(s) \ge w_2(t^*) - w_2(s) \ge -\varepsilon'$ , whereby

$$g_*(s) \le 1_{\{w_2(t^*) - w_2(s) < \varepsilon'\}} (p(t^* - s, w_1(t^*) - w_1(s)) - p(t^* - s, \varepsilon')) \le p(t^* - s, 0) - p(t^* - s, \varepsilon').$$
(4.11)

Integrating (4.11) into (4.10) and then applying the following readily verified identity

$$\int_{0}^{t} p(s,x)ds = 2tp(t,x) - 2|x|\widetilde{\Phi}(t,|x|), \qquad (4.12)$$

we obtain

$$\int_{0}^{t^{*}} g_{*}(s)ds \leq \sqrt{\frac{2T}{\pi}} (1 - \exp(-\varepsilon'^{2}/2T)) + 2\varepsilon' < 4\varepsilon',$$
(4.13)

for all  $\varepsilon' \leq 1$  and  $T \geq 1$ . Now, if  $t^* \leq T$ , combining (4.13) with (4.10) and (4.9) yields  $\delta c_1 < \varepsilon + 4\varepsilon'$ , leading to a contradiction. Consequently, we must have  $t^* > T$ .

We now turn to the proof of Lemma 2.4. For sufficiently smooth solutions to the PDE (1.13), one can easily calculate  $\frac{d^2}{dt^2} z_{\star}(\frac{1}{2}) = -\frac{1}{8} \partial_{xxx} u_1(\frac{1}{2}, 0) = \frac{2}{\sqrt{\pi}}$  by differentiating (1.13e) and (1.13b). Here, as we take the integral equation (1.9) as the definition of the Stefan problem, we prove Lemma 2.4 by a different argument, without assuming the smoothness of of  $z_{\star}$ .

Proof of Lemma 2.4. To simplify notations, throughout this proof we let  $w(t) := z_{\star}(\frac{1}{2} + t)$ , and use r(t, x) to denote a generic bounded function of (t, x).

Let  $F(t,x) := \int_0^\infty p(t,x-y)U_\star(\frac{1}{2},y)dy$  denote the l.h.s. of (1.9) (with a  $\frac{1}{2}$  shift in time) and let  $u_\star(x) := \partial_x U_\star(\frac{1}{2},x) = -2\partial p(\frac{1}{2},x) + 4\widetilde{\Phi}(\frac{1}{2},x)$  (see (1.10)). Integrating by parts followed by a change of variable  $y \mapsto \frac{y}{\sqrt{t}}$  yields

$$F(t,x) := \sqrt{t} \int_0^\infty u_\star(\sqrt{t}y) \Phi(1, \frac{x}{\sqrt{t}} - y) dy.$$

Further, with  $u_{\star} \in \mathcal{C}^{\infty}(\mathbb{R}_{+}) \cap L^{\infty}(\mathbb{R}_{+}), u_{\star}(0) = 2$  and  $\frac{d}{dx}u_{\star}(0) = \frac{d^{2}}{dx^{2}}u_{\star}(0) = 0$ , we have  $F(t,x) = t^{1/2}F_{0}(\frac{x}{\sqrt{t}}) - t^{2}F_{3}(\frac{x}{\sqrt{t}}) + t^{5/2}r(t,x)$ , where

$$F_0(x') := 2 \int_0^\infty \Phi(1, x' - y) dy, \quad F_3(x') := \frac{a_3}{6} \int_0^\infty y^3 \Phi(1, x' - y) dy,$$

where  $a_3 := -\frac{d^3}{dx^3} u_{\star}(0) = \frac{16}{\sqrt{\pi}}$ .

Insert the preceding expression of F into (1.9) to obtain

$$t^{1/2}F_0(\frac{w(t)}{\sqrt{t}}) - t^2F_3(\frac{w(t)}{\sqrt{t}}) + t^{5/2}r(t,x) = \int_0^t p(t-s,w(t)-w(s))ds \le t^{1/2}F_0(0), \quad (4.14)$$

where the last inequality follows by replacing p(t - s, w(t) - w(s)) with p(t - s, 0). As  $x' \mapsto F_0(x')$  is strictly increasing and continuous, letting  $t \downarrow 0$  we conclude that  $\lim_{t\downarrow 0} \frac{w(t)}{\sqrt{t}} = 0$ . Now, dividing both sides of (4.14) by  $t^2$ , and letting  $t \downarrow 0$  using  $\lim_{t\downarrow 0} \frac{w(t)}{\sqrt{t}} = 0$ , we further deduce that  $\lim_{t\downarrow 0} t^{-3/2}(F_0(\frac{w(t)}{\sqrt{t}}) - F_0(0)) - F_3(0) \leq 0$ . With  $\frac{d}{dx'}F_0(0) = 1$ , Taylor-expanding  $F_0$  in the preceding inequality yields  $\limsup_{t\downarrow 0} \frac{w(t)}{t^2} \leq F_3(0) = \frac{a_3}{8} = \frac{2}{\sqrt{\pi}}$ . From this, we further obtain that

$$\int_0^t p(t-s, w(t) - w(s)) ds \ge \int_0^t p(t-s, \frac{1}{C}t^2) ds \ge t^{1/2} F_0(0) - t^{5/2} r(t, x).$$

Combining this with (4.14) yields  $\liminf_{t\downarrow 0} \frac{w(t)}{t^2} \ge F_3(0) = \frac{2}{\sqrt{\pi}}$ .

## 5. The Absorption Phase

We begin by settling Lemma 2.2.

Proof of Lemma 2.2. Consider the modified process  $(X'_i(t); t \ge 0)_{1 \le i \le K}$  consisting of K independent Brownian motions starting at x = 1 and absorbed once they reach x = 0, and let  $\tau'_{\text{ext}} := \inf\{K^{-1}t : X_i(t) = 0, \forall i\}$  denote the corresponding scaled extinction time. Under the natural coupling of  $(X'_i(t))_i$  and  $(X_i(t))_i$  (by letting them sharing the underlying Brownian motion), we clearly have  $\tau_{\text{ext}} \ge \tau'_{\text{ext}}$ . For the later, it is straightforward to verify that

$$\mathbf{P}(\tau'_{\text{ext}} \le T) = \left(\mathbf{P}\left(\inf_{t \le T} (B(Kt) + 1) \le 0\right)\right)^K \le \exp(-\frac{1}{C}K^{1/2}).$$

where  $B(\cdot)$  is a standard Brownian motion and  $C = C(T) < \infty$ . From this the desired result follows.

We proceed to proving Proposition 2.3. To this end, we first establish the following short time estimate.

**Lemma 5.1.** For any fixed  $\alpha \in (0, \frac{1}{2})$  and  $n < \infty$ , there exists  $C = C(\alpha, n) < \infty$  such that

$$\mathbf{P}\left(\sup_{t\in[0,K^{-2\alpha}]}Z_K(t)\leq K^{-\alpha}\right)\geq 1-CK^{-n},\quad\forall K<\infty.$$
(5.1)

Proving Lemma 5.1 requires the following classical result from [Fel71].

**Lemma 5.2** ([Fel71, Chapter X.5, Example (c)]). Let  $(B(t); t \ge 0)$  be a standard Brownian motion (starting from 0), and let  $0 < a < b < \infty$ . Defining  $\rho(t, a, b) := \mathbf{P}(0 < B(s) + a < b, \forall s \le t)$ , we have

$$\rho(t,a,b) = \sum_{n=0}^{\infty} \frac{4}{(2n+1)\pi} \sin\left(\frac{(2n+1)\pi a}{b}\right) \exp\left(-\frac{(2n+1)^2 \pi^2}{2b^2}t\right).$$
 (5.2)

Proof of Lemma 5.1. Fixing  $\alpha \in (0, \frac{1}{2})$  and  $n < \infty$ , we let  $C := C(\alpha, n) < \infty$  denote a generic constant depending only on  $\alpha$  and n.

We consider first the modified process  $(\widehat{X}_i^K(t); t \ge 0)_{1 \le i \le K}$ , which consists of K independent Brownian motions starting at  $x = K^{-\frac{1}{2}}$ , and absorbed at x = 0 and  $x = \frac{1}{2}K^{-\alpha}$ . We begin by deriving a lower bound on

$$\widehat{N} := \left| \left\{ i : 0 < \widehat{X}_i^K(s) < \frac{1}{2} K^{-\alpha}, \forall s \le K^{-2\alpha} \right\} \right|,$$

the number of surviving particles of up to time  $K^{-2\alpha}$ . As  $\widehat{N}$  is the sum of i.i.d. Bernoulli $(\rho_K^*)$  random variables, where  $\rho_K^* := \rho(K^{-2\alpha}, K^{-\frac{1}{2}}, \frac{1}{2}K^{-\alpha})$ , by the Chernoff bound we have

$$\mathbf{P}(\widehat{N} \ge \frac{1}{2}K\rho_K^*) \ge 1 - \exp\left(-\frac{1}{8}K\rho_K^*\right).$$
(5.3)

Now, specializing (5.2) at  $(t, a, b) = (K^{-2\alpha}, K^{-\frac{1}{2}}, \frac{1}{2}K^{-\alpha})$ , we obtain

$$\rho_K^* = \sum_{n=0}^{\infty} \rho_{K,n}' \exp\left(-2(2n+1)^2 \pi^2\right), \text{ where } \rho_{K,n}' := \frac{4}{(2n+1)\pi} \sin(2(2n+1)\pi K^{\alpha-\frac{1}{2}}).$$

With  $\lim_{K\to\infty} (K^{\frac{1}{2}-\alpha}\rho'_{K,n}) = 8$  and  $|\rho'_{K,n}| \le 8K^{\alpha-\frac{1}{2}}$ , it is straightforward to show that

$$\lim_{K \to \infty} (K^{\frac{1}{2} - \alpha} \rho_K^*) = 8 \sum_{n=0}^{\infty} \exp\left(-2(2n+1)^2 \pi^2\right) > 0$$

Consequently,  $\rho_K^* \geq \frac{1}{C} K^{\alpha - \frac{1}{2}}$ . Inserting this into (5.3), we arrive at

$$\mathbf{P}(\widehat{N} \ge \frac{1}{C}K^{\alpha + \frac{1}{2}}) \ge 1 - \exp(-\frac{1}{C}K^{\alpha + \frac{1}{2}}) \ge 1 - CK^{-n}.$$

Next, we consider the process  $(\widehat{X}_i^{\prime K}(t); t \geq 0)_{1 \leq i \leq K}$ , consisting of K independent Brownian motions starting at  $x = K^{-\frac{1}{2}}$  and absorbed only at x = 0, coupled to  $(\widehat{X}_i^K(t))_i$  by the aforementioned natural coupling. Let  $\Gamma := \{\widehat{X}_i^{\prime K}(K^{-2\alpha}) : 0 < \widehat{X}_i^{\prime K}(s) < \frac{1}{2}K^{-\alpha}, \forall s \leq K^{-2\alpha}\}$ denote the set of all  $\widehat{X}_i^{\prime K}$  particles that stay within  $(0, \frac{1}{2}K^{-\alpha})$  for all  $s \leq K^{-2\alpha}$ , and let  $\widehat{N}' := |\Gamma|$ . We clearly have  $\widehat{N}' \geq \widehat{N}$ , whereby

$$\mathbf{P}(\widehat{N}'(K^{-2\alpha}) \ge \frac{1}{C}K^{\alpha + \frac{1}{2}}) \ge 1 - CK^{-n}.$$
(5.4)

Now, couple  $(\widehat{X}_i^{\prime K}(t))_i$  and  $(X_i^K(t))_i$  (under the push-the-laggard strategy) by the natural coupling. On the event  $\{\widehat{N}' \geq \frac{1}{C}K^{\alpha+\frac{1}{2}}\}$ , to move all particles in  $\Gamma$  to the level  $x = K^{-\alpha}$  requires at least a drift of  $\widehat{N}'\frac{K^{-\alpha}}{2} \geq \frac{1}{C}K^{\frac{1}{2}}$ , while the total amount of (scaled) drift at disposal is  $K^{-2\alpha+\frac{1}{2}}$ , which is less than  $\frac{1}{C}K^{\frac{1}{2}}$  for all large enough K. Consequently, the desired result (5.1) follows from (5.4).

Proof of Proposition 2.3. Fixing  $\beta \in (0, \frac{1}{24})$ ,  $\alpha \in (4\beta, \frac{1}{2} - 8\beta)$  and  $n < \infty$ , we  $C = C(\beta, n) < \infty$  denote a generic constant. We consider the threshold function

$$z^{*}(t) = K^{-\alpha} \mathbf{1}_{\{t \le K^{-2\alpha}\}} + (\sqrt{t}K^{-\beta}) \mathbf{1}_{\{t > K^{-2\alpha}\}},$$
(5.5)

with the corresponding hitting time  $\tau := \inf\{t \in \mathbb{R}_+ : Z_K(t) \ge z^*(t)\}$ . It suffices to show  $\mathbf{P}(\tau > \frac{1}{2} + \frac{1}{7}K^{-2\beta}) \ge 1 - CK^{-n}$ . To this end, by Lemma 5.1, without loss of generality we assume  $\tau \in (K^{-2\alpha}, 1)$ . As the trajectory of  $Z_K$  is continuous except when it hits 0, we have  $Z_K(\tau) \ge z^*(\tau)$ . Hence at time  $\tau$ , no particle exists between 0 and  $z^*(\tau)$ , or equivalently  $\widetilde{U}_K(\tau, z^*(\tau)) = \widetilde{U}_K(\tau, 0)$ . With this, taking the difference of (2.7) at  $x = z^*(\tau)$  and at x = 0, and multiplying the result by  $\sqrt{\frac{\pi\tau}{2}}$ , we obtain

$$h_{1} = h_{2} + \sqrt{\frac{\pi\tau}{2}} (R_{K}(\tau, z^{*}(\tau)) - R_{K}(\tau, 0)),$$
  
where  $h_{1} := \sqrt{\frac{\pi\tau}{2}} (\widetilde{G}_{K}(\tau, 0) - \widetilde{G}_{K}(\tau, z^{*}(\tau))), h_{2} := \sqrt{\frac{\pi\tau}{2}} \int_{0}^{\tau} f_{2}(s, Z_{K}(s), z^{*}(\tau)) ds,$  and  
 $f_{2}(s, z, z') := p^{N}(\tau - s, z, z') - p^{N}(\tau - s, z, 0).$  (5.6)

Further using (2.5), for fixed  $\gamma \in (0, \frac{1-2\alpha}{4} - 4\beta)$ , to control the remainder term  $(R_K(\tau, z^*) - R_K(\tau, 0))$ , with  $\tau \geq K^{-2\alpha}$ , we have

$$h_1 \le h_2 + CK^{\frac{2\alpha-1}{4}+\gamma},$$
 (5.7)

with probability  $1 - CK^{-n}$ . To show  $\tau \ge \frac{1}{2} + \frac{1}{7}K^{-2\beta}$ , we next derive a lower bound on  $h_1$  and an upper bound on  $h_2$ .

With  $G_K(t, x)$  defined as in (2.3), we have

$$h_1 = h_1(K^{-\beta}), \text{ where } h_1(a) = \sqrt{K\tau} \int_0^{\frac{1}{\sqrt{K\tau}}} \left(e^{-\frac{y^2}{2}} - \frac{1}{2}e^{-\frac{(y+a)^2}{2}} - \frac{1}{2}e^{-\frac{(y-a)^2}{2}}\right) dy.$$

Taylor-expanding  $h_1(a)$  to the fifth order, we further obtain that  $h_1 \ge a^2 h_{12} + a^4 h_{14} - Ca^6$ , for  $a = K^{-\beta}$ , where  $h_{12} := \sqrt{K\tau} \int_0^{\frac{1}{\sqrt{K\tau}}} e^{-y^2/2} (\frac{1}{2} - \frac{1}{2}y^2) dy$  and  $h_{14} := \sqrt{K\tau} \int_0^{\frac{1}{\sqrt{K\tau}}} e^{-y^2/2} (-\frac{1}{8} + \frac{1}{4}y^2 - \frac{1}{24}y^4) dy$ . Further Taylor-expanding  $h_{12}$  and  $h_{14}$  in  $\frac{1}{\sqrt{K\tau}}$  yields  $h_{12} \ge \frac{1}{2} - \frac{C}{K\tau}$  and  $h_{14} \ge -\frac{1}{8} - \frac{C}{K\tau}$ , and therefore

$$h_1 \ge \frac{1}{2}a^2 - \frac{1}{8}a^4 - Ca^2(\frac{1}{K\tau} + a^4) \ge \frac{1}{2}a^2 - \frac{1}{8}a^4 - Ca^2(K^{2\alpha - 1} + a^4), \text{ for } a = K^{-\beta}.$$
 (5.8)

Turning to estimating  $h_2$ , we first observe that the function  $f_2(s, z, z')$  as in (5.6) increases in  $z, \forall z \leq z'$ , as is readily verified by taking derivative as follows:

$$\sqrt{2\pi(\tau-s)^3} \partial_z f_2(s,z,z')$$

$$= z'(e^{-(z-z')^2/2} - e^{-(z+z')^2/2}) - z(e^{-(z-z')^2/2} + e^{-(z+z')^2/2}) + 2ze^{-z^2/2}$$

$$\geq z(e^{-(z-z')^2/2} - e^{-(z+z')^2/2}) - z(e^{-(z-z')^2/2} + e^{-(z+z')^2/2}) + 2ze^{-z^2/2} \ge 0.$$

Now, since  $t \mapsto z^*(t)$  is increasing for all  $t \ge K^{-2\alpha}$ , to obtain a upper bound on  $h_2$  we replace  $Z_K(s)$  with  $z^*(\tau)$  for  $s \ge K^{-2\alpha}$ . Further, with  $\int_0^{K^{-2\alpha}} p^N(\tau - s, z, z') ds \le CK^{-\alpha}$ , we obtain  $h_2 \le CK^{-\alpha} + \sqrt{\frac{\pi\tau}{2}} \int_0^{\tau} f_2(s, z^*(\tau), z^*(\tau)) ds$ . With  $z^*(\tau) = K^{-\beta} \sqrt{\tau}$ , the last integral is evaluated explicitly by using (4.12), yielding

$$h_2 \le \tau h_2(K^{-\beta}) + CK^{-\alpha}$$
, where  $h_2(a) = 1 + e^{-2a^2} - 2e^{-a^2/2} + 2a \int_a^{2a} e^{-y^2/2} dy$ .

Taylor-expanding  $h_2(a)$  to the fifth order, we further obtain

$$h_2 \le \tau (a^2 - \frac{7}{12}a^4 + Ca^6) + CK^{-\alpha}, \quad \text{for } a = K^{-\beta}.$$
 (5.9)

Now, combining (5.7)–(5.9), we arrive at

$$\tau \ge \frac{\frac{1}{2} - \frac{1}{8}a^2 - C(K^{2\alpha - 1} + a^4)}{1 - \frac{7}{12}a^2 + C(a^4 + a^{-2}K^{-\alpha} + a^{-2}K^{\frac{2\alpha - 1}{4} + \gamma})}, \text{ for } a = K^{-\beta}.$$
 (5.10)

With  $\alpha$  and  $\gamma$  chosen as in the preceding, it is now easy to check that  $\tau \geq \frac{1}{2} + \frac{1}{6}K^{-2\gamma} + (\text{higher order terms})$ , from which the desired result follows.

## 6. The Moving Boundary Phase

6.1. **Proof of Lemma 4.3.** Recall the definition of  $\lambda$ -Atlas model from (2.13). We further define the **infinite**  $\lambda$ -Atlas model to be the analog of (2.13) with semi-infinitely many particles, i.e.

$$dY_i(t) = \mathbf{1}_{\{X_i(t) = Z^Y(t)\}} \,\lambda dt + dB_i(t) \quad \text{for } i = 1, 2, \dots,$$
(6.1)

where  $Z^{Y}(t) := \inf\{X_{i}(t) : i \in \mathbb{N}\}$  again denotes the lowest ranked particle. General conditions on  $(X_{i}(0))_{i\in\mathbb{N}}$  for (6.1) to be well-posed are studied in [IKS13, Shk11]. Here we consider only a particular initial condition, i.e.  $(Y_{i}(0))_{i\in\mathbb{N}} \sim \text{PPP}_{+}(2\lambda)$ , the Poisson Point

Process on  $\mathbb{R}_+$  with rate  $2\lambda$ , which is in fact the invariance measure of the gap process  $(Y_{i+1}(t) - Y_i(t))_{i \in \mathbb{N}}$  [PP08].

We begin by recalling some comparison techniques of the Atlas models. Given two random vectors  $X = (X_i), Y = (Y_i) \in \mathbb{R}^n$ , we denote by  $X \succeq Y$  that Y is stochastically bounded above by X, namely there exists a coupling of X, Y under which  $X_i \ge Y_i, i = 1, ..., n$ .

**Lemma 6.1.** Fixing  $\lambda > 0$ , we let  $(Y_i^1(t))_i$  and  $(Y_i^2(t))_i$  be either finite  $\lambda$ -Atlas models or infinite  $\lambda$ -Atlas models starting from PPP<sub>+</sub>(2 $\lambda$ ), with  $n_j := |\{Y_i^j(0)\}|$  particles, respectively.

- (a) If  $n_1 \ge n_2$  and  $Y_i^1(0) \le Y_i^2(0)$  for  $1 \le i \le n_2$ , then there exists a coupling of  $(Y_i^1(t))_i$ and  $(Y_i^2(t))_i$  such that  $Y_i^1(t) \le Y_i^2(t)$ , for all  $1 \le i \le n_2$  and t > 0.
- (b) Consider the case  $n_2 < \infty$ . Let  $G^2(t) := (Y_{i+1}^2(t) Y_i^2(t))_{1 \le i \le n_2 1}$  denote the gap process of  $Y^2(t)$ . If initially  $G^2(0) \succeq \bigotimes_{i=1}^{n_2 1} \operatorname{Exp}(2\lambda)$ , then  $G^2(t) \succeq \bigotimes_{i=1}^{n_2 1} \operatorname{Exp}(2\lambda)$ , for all t > 0.

Proof. Part (a) follows readily from [Sar15, Corollary 3.9, Remark 9]. As for (b), consider the infinite  $\lambda$ -Atlas model  $(Y'_i(t))_{i\in\mathbb{N}}$  started from PPP<sub>+</sub>(2 $\lambda$ ). According to [PP08, Theorem 1],  $\otimes_{i\in\mathbb{N}} \operatorname{Exp}(2\lambda)$  is an invariance measure of the gap process, i.e.  $(Y'_{i+1}(t) - Y'_i(t))_i \sim \otimes_{i\in\mathbb{N}} \operatorname{Exp}(2\lambda)$ , for all t > 0. With this, Part (b) follows by coupling  $(Y'_i(t))_i$  and  $(Y'_i(t))_i$  using Part (a).

Fix  $T < \infty$ ,  $u_{\text{in}} \in \mathcal{C}(\mathbb{R}_+) \cap L^1(\mathbb{R}_+)$  with  $0 < u_{\text{in}}(x) \leq 2$ ,  $\forall x \geq 0$ , and let  $U_{\text{in}}(x) := \int_0^x u_{\text{in}}(y) dy$  as in the preceding. Proving Lemma 4.3 amounts to constructing a nondecreasing solution  $z \in \mathcal{C}([0,T])$  to (4.3), which is obtained as the hydrodynamic limit of a certain process as follows. Setting  $u_{\text{in}}^K(x) := u_{\text{in}}(K^{-1/2}x)$ , we let  $\hat{Y}(t) := (\hat{Y}_i(t); t \geq 0)_i$  be a unit-Atlas model with the initial condition  $(\hat{Y}_i(0))_i \sim \text{PPP}_+(u_{\text{in}}^K)$ , the Poisson Point Process on  $\mathbb{R}_+$  with density  $u_{\text{in}}^K$ . Scale the process  $Y_i^K(t) := K^{-1/2}Y_i(K^{-1}t)$  as in the preceding, and let  $Z_K^Y(t) := \inf\{Y_i^K(t)\}$ . We begin by showing that  $Z_K^Y(t)$  is approximately nondecreasing. Recall the definition of  $|\cdot|_{[0,T]}$  from (4.8).

**Lemma 6.2.** For any fixed  $\beta \in (0, \frac{1}{4})$  and  $n < \infty$ , there exists  $C = C(T, \beta, n) < \infty$  such that  $\mathbf{P}(|Z_K^Y|'_{[0,T]} \leq K^{-\beta}) \geq 1 - CK^{-n}, \forall K < \infty$ .

*Proof.* Fixing such  $T, \beta, n$ , we let  $C = C(T, \beta, n) < \infty$  denote a generic finite constant depending only on  $T, \beta, n$ .

Let  $G_i^K(t) := Y_{i+1}^K(t) - Y_i^K(t)$  denote the corresponding gap process. With  $u_{in}(x) \le 2, \forall x \ge 0$ , we have that  $(G_i^K(0))_i \succeq \otimes_i \operatorname{Exp}(2\sqrt{K})$ , so by Lemma 6.1(b),  $(G_i^K(t))_i \succeq \otimes_i \operatorname{Exp}(2\sqrt{K})$ . Fixing  $s_* \in [0, T]$ , we start an *infinite*  $2\sqrt{K}$ -Atlas model  $(\widehat{Y}_i^K(t); t \ge s_*)_i$  with the initial condition

$$Z_{K}^{Y}(s_{*}) = \widehat{Y}_{1}^{K}(s_{*}) < \widehat{Y}_{2}^{K}(s_{*}) < \ldots \to \infty, \quad (\widehat{Y}_{i+1}^{K}(s_{*}) - \widehat{Y}_{i}^{K}(s_{*}))_{i} \sim \bigotimes_{i \in \mathbb{N}} \operatorname{Exp}(2\sqrt{K}). \quad (6.2)$$

Let  $Y_{(1)}^{K}(s_{*}) \leq Y_{(2)}^{K}(s_{*}) \leq \ldots$  denote the ordered  $Y_{i}^{K}$  particles at time  $s_{*}$ . With (6.2), we then couple  $(\widehat{Y}_{i}^{K}(s_{*}))_{i}$  and  $(Y_{(i)}^{K}(s_{*}))_{i}$  in such a way that  $Y_{(i)}^{K}(s_{*}) \geq \widehat{Y}_{i}^{K}(s_{*}), \forall 1 \leq i \leq |\{Y_{i}^{K}(t)\}|$ . Let  $\widehat{Z}_{K}(t) := \min\{\widehat{Y}_{i}^{K}(t)\}$ . By Lemma 6.1(a), we then have a coupling of  $(Y_{i}^{K}(t))_{i}$  and  $(\widehat{Y}_{i}^{K}(t))_{i}$  such that  $Z_{K}^{Y}(t) \geq \widehat{Z}_{K}(t), \forall t \geq s_{*}$ . Furthermore, by [DT15, Proposition 2.3, Remark 2.4], with  $\beta < \frac{1}{4}$ , we have  $\mathbf{P}(\sup_{t \in [s_{*}, s_{*}+T]} |\widehat{Z}_{K}(t) - \widehat{Z}_{K}(s_{*})| \leq \frac{1}{2}K^{-\beta}) \geq 1 - CK^{-n-2}$ ,  $\forall K < \infty$ . Consequently, by the preceding coupling of we obtain

$$\mathbf{P}\left(\sup_{t\in[s_*,T]} (Z_K^Y(s_*) - Z_K^Y(t)) \le \frac{1}{2}K^{-\beta}\right) \ge 1 - CK^{-n-2}, \quad \forall K < \infty.$$
(6.3)

Further taking the union bound of (6.3) over  $s_* = s_\ell := K^{-2}T\ell$ ,  $1 \le \ell \le K^2$ , we have

$$\mathbf{P}\left(\sup_{t\in[s_{\ell},T]} (Z_K^Y(t_{\ell}) - Z_K^Y(t)) \le \frac{1}{2}K^{-\beta}, \ \forall 1 \le \ell \le TK^2\right) \ge 1 - CK^{-n}.$$
(6.4)

Within each  $s \in [s_{\ell}, s_{\ell+1}]$ , by the same argument we obtained (3.38), we obtain here the following continuity estimate:

$$\mathbf{P}\left(\sup_{s\in[s_{\ell},s_{\ell+1}]}|Z_{K}^{Y}(s)-Z_{K}^{Y}(t_{\ell})|\leq\frac{1}{2}K^{-\beta}\right)\geq1-CK^{-n-2}.$$
(6.5)

Taking the union bound of (6.5) over  $1 \le \ell \le K^2$ , and then combining the result with (6.4), we obtain the desired result.

With  $(Y_i^K(0))_i \sim \text{PPP}_+(K^{1/2}u_{\text{in}})$ , one easily verifies that (2.15)–(2.16) hold. With this, we now apply Proposition 2.6 to obtain

$$\begin{aligned} U_K^Y(t,x) &= \int_0^\infty p(t,x-y) U_K^Y(0,y) dy - \int_0^t p(t-s,x-Z_K^Y(s)) ds + R_K^Y(t,x), \\ &= \int_0^\infty p(t,x-y) U_{\rm in}(y) dy - \int_0^t p(t-s,x-Z_K^Y(s)) ds + f_K'(t,x) + R_K^Y(t,x), \end{aligned}$$

where  $R_K^Y(0, t, x)$  satisfies (2.18) and  $f'_K(t, x) := \int_0^\infty p(t, x - y)(U_K^Y(0, y) - U_{\text{in}}(y))dy$ . Further, with  $U_K^Y(t, Z_K^Y(t)) = 0$ , setting  $x = Z_K^Y(t)$  yields

$$\int_{0}^{\infty} p(t, Z_{K}^{Y}(t) - y) U_{\text{in}}(y) dy = \int_{0}^{t} p(t - s, Z_{K}^{Y}(t) - Z_{K}^{Y}(s)) ds - f_{K}'(t, Z_{K}^{Y}(t)) - R_{K}^{Y}(t, Z_{K}^{Y}(t)).$$
(6.6)

Next we establish a uniform upper bound on  $\{Z_K^Y\}_K$ .

**Lemma 6.3.** For any fixed  $T, n < \infty$ , there exists  $C = C(T, n) < \infty$  and  $L = L(T, U_{in}) < \infty$  such that  $\mathbf{P}(\sup_{t \leq T} Z_K^Y(t) \leq L) \geq 1 - CK^{-n}, \forall K < \infty$ .

*Proof.* With the bound (2.18) on  $R_K^Y(t, x)$ , we begin by establishing a bound on  $f'_K$ . Indeed,  $|f'_K(t, x)| \leq \sup_{y \in \mathbb{R}} |U_K^Y(0, y) - U_{in}(y)|$ . With  $u_{in} \in L^1(\mathbb{R})$ , we have  $U_{in}(\infty) < \infty$ , and  $(U_K^Y(0, y) - U_{in}(y); y \geq 0)$  is a compensated Poisson process, so by Doob's  $L^p$  maximal inequality we have

$$\mathbf{E}\left(\sup_{y\in\mathbb{R}}\left|U_{K}^{Y}(0,y)-U_{\mathrm{in}}(y)\right|\right)^{k} \leq C \mathbf{E}\left(\left|U_{K}^{Y}(0,\infty)-U_{\mathrm{in}}(\infty)\right|\right)^{k} \leq C K^{-\frac{k}{4}},\tag{6.7}$$

for any fixed  $k < \infty$ , where  $C = C(k) < \infty$ . From this we conclude that

$$\mathbf{P}(\|f'_K\|_{\mathcal{C}([0,T]\times\mathbb{R})} \le K^{-\beta}) \ge 1 - C(\beta,k)K^{-k},$$
(6.8)

for any fixed  $\beta \in (0, \frac{1}{4})$  and  $k < \infty$ .

Now, we set  $z^*(t) := 1 + at$ , for some  $a \in \mathbb{R}_+$  to be specified, and consider the first hitting time  $\tau := \inf\{t : Z_K^Y(t) \ge z^*(t)\}$ . Indeed, as  $z \mapsto \int_0^\infty p(\tau, z - y)U_{\text{in}}(y)dy$  is increasing, we have

$$\int_{0}^{\infty} p(\tau, Z_{K}^{Y}(\tau) - y) U_{\text{in}}(y) dy \ge \int_{0}^{\infty} p(\tau, 1 - y) U_{\text{in}}(y) dy := f_{1}(\tau)$$
(6.9)

$$\int_0^\tau p(\tau - s, Z_K^Y(\tau) - Z_K^Y(s)) ds \le \int_0^\infty p(s, as) ds := f_2(a).$$
(6.10)

Fixing  $T < \infty$ , with  $u_{\text{in}} > 0$ , we clearly have  $\inf_{t \leq T} f_1(t) := f_* > 0$  and  $\lim_{a \to \infty} f_2(a) = 0$ . With this, fixing some large enough a with  $f_2(a) < \frac{1}{2}f_*$ , we insert the bounds (6.8)–(6.10), (2.18) into (6.6) to obtain

$$\mathbf{P}\left(\{f_* \le \frac{1}{2}f_* + K^{-\beta}\} \cap \{\tau \ge T\}\right) \ge 1 - C(\beta, n)K^{-n}.$$

With  $f_* > 0$ ,  $\{f_* \leq \frac{1}{2}f_* + K^{-\beta}\}$  is impossible for all large enough K, so we obtain the desired result for L = aT + 1.

We are now ready to construct the solution z to (4.3).

**Lemma 6.4.** There exists nondecreasing  $z \in C([0,T])$  which solves (4.3). Such z is in fact the uniform limit of  $Z_K$ : for any fixed  $\beta \in (0, \frac{1}{4})$  and  $n < \infty$ , there exists  $C = C(\beta, n) < \infty$ such that

$$\mathbf{P}(|Z_K^Y - z|_{\mathcal{C}([0,T])} \le K^{-\beta}) \ge 1 - CK^{-n}, \quad \forall K < \infty.$$
(6.11)

Proof. Fixing  $K_1 < K_2 < \infty$ ,  $1 \le n < \infty$  and  $\beta \in (0, \frac{1}{4})$ , we begin by estimating  $\sup_{t \in [0,T]} |Z_{K_1}(t) - Z_{K_2}(t)|$ . To this end, we apply Lemma 6.3 and Lemma 4.4 for  $(w_1, w_2) = (Z_{K_1}^Y, Z_{K_2}^Y)$  and for  $(w_1, w_2) = (Z_{K_2}^Y, Z_{K_1}^Y)$  to obtain

$$|Z_{K_1}^Y - Z_{K_2}^Y|_{\mathcal{C}([0,T])} \le C_1 \sum_{j=1,2} \left( |Z_{K_j}^Y|_{[0,T]}' + |R_{K_j}^Y|_{\mathcal{C}([0,T] \times \mathbb{R})} + |f_{K_j}'|_{\mathcal{C}([0,T] \times \mathbb{R})} \right), \tag{6.12}$$

where  $C_1 < \infty$  is as in Lemma 4.4. Next, using (6.8), (2.18) and Lemma 6.2 to bound the r.h.s. of (6.12), we further obtain

$$\mathbf{P}\left(|Z_{K_1}^Y - Z_{K_2}^Y|_{\mathcal{C}([0,T])} \le (1 - 2^{-\beta})K_1^{-\beta}\right) \ge 1 - C(\beta, n)K_1^{-n}.$$
(6.13)

Consider the subsequence  $K'_m := 2^m$ . Letting  $(K_1, K_2) = (K'_{m+j}, K'_{m+j+1})$  in (6.13), and taking the union bound over all  $j \in \mathbb{Z}_{>0}$ , we then arrive at

$$\mathbf{P}\left(|Z_{K'^m}^Y - Z_{K'_{m+j}}^Y|_{\mathcal{C}([0,T])} \le (K'_m)^{-\beta}, \forall j < \infty\right) \ge 1 - C(\beta, n)(K'_m)^{-n}.$$
(6.14)

From this, we conclude that  $\{Z_{K'_m}^Y\}_m$  is almost surely Cauchy in  $\mathcal{C}([0,T])$ , and hence converges to a (possibly random) limit  $Z \in \mathcal{C}([0,T])$ . Now, letting  $K \to \infty$  in (6.6), with (2.18) and (6.8), we find that Z is a solution to (4.3), and  $t \mapsto Z(t)$  is nondecreasing by Lemma 6.2. As the solution to (4.3) is unique (by Corollary 4.5), Z = z must in fact be deterministic. Finally, from (6.13)–(6.14), we have that actually the whole sequence  $\{Z_K^Y\}_K$  converges (in probability) to z, and that (6.11) holds.

6.2. **Proof of Proposition 2.7.** Fix  $\beta \in (0, \frac{1}{96})$  and let  $\sigma_K := \frac{1}{2} + \frac{1}{7}K^{-2\beta}$ . Recall that  $(X_i^{*,K}(s); s \ge 0)_i$  denote the modified process where the absorption is *turned off* for  $s \ge \sigma_K$ . The proof is similar to that of Lemma 6.4. Indeed, with  $|\{X_i^K(\sigma_K) : X_i^K(\sigma_K) > 0\}| \le K$  and (3.26), we have that  $(X_i^{*,K}(\sigma_K))_i$  satisfies (2.15)–(2.16), so, by Proposition 2.6, we have

$$U_{K}^{*}(t+\sigma_{K},x) = \int_{0}^{\infty} p(t,x-y)U_{K}(\sigma_{K},y)dy - \int_{0}^{t} p(t-s,x-Z(s+\sigma_{K}))ds + R_{K}^{Y}(t,x),$$

where  $U_K(s, x) := K^{-1/2} \sum_{i=1}^K \mathbf{1}_{(0,x)}(X_i^K(s))$  and  $U_K^*(s, x) := K^{-1/2} \sum_{i=1}^K \mathbf{1}_{(0,x)}(X^{*,K}(s))$ denote the scaled distribution functions for  $(X_i^K(s))_i$  and  $(X_i^{*,K}(s))_i$ , respectively. Now, let  $Z_K^*(s) := \inf\{X_i^{*,K}(s)\}$  and  $W_K(t) := Z_K^*(t + \sigma_K) - Z_K^*(\sigma_K)$ . Similar to (6.6), with  $U_K^*(s, Z_K^*(s)) = 0$ , we have

$$\int_0^\infty p(t, W_K(t) - y) U_\star(\sigma_K, y + z_\star(\sigma_K)) dy = \int_0^t p(t - s, W_K(t) - W_K(s)) ds - f_K^*(t, W_K(t)) - R_K^Y(t, W_K(t) + Z_K^*(\sigma_K)),$$

where  $f_K^*(t,x) = \int_0^\infty p(t,x-y)(U_K(\sigma_K,y+Z_K^*(\sigma_K)) - U_\star(\sigma_K,y+z_\star(\sigma_K)))dy$ . With  $U_\star(s,x)$  and  $z_\star(s)$  as in (4.4) and (1.9), respectively, it is straightforward to verify that

$$\int_0^\infty p(t, z_\star(\sigma_K + t) - y) U_\star(\sigma_K, y) dy = \int_0^t p(t - s, z_\star(\sigma_K + t) - z_\star(\sigma_K + s)) ds$$

Hence, letting  $w(t) := z_{\star}(t + \sigma_K) - z_{\star}(\sigma_K)$ , we have

$$\int_0^\infty p(t, w(t) - y) U_{\star}(\sigma_K, y + z_{\star}(\sigma_K)) dy = \int_0^t p(t - s, w(t) - w(s)) ds.$$

Now, fixing  $T < \infty$ , we apply Lemma 4.4 for  $(w_1, w_2) = (W_K, w)$  to obtain

$$\sup_{\in [0,T]} \left( Z_K^*(\sigma_K + t) - z_\star(\sigma_K + t) \right) \le C_1(|f_K^*|_{\mathcal{C}([0,T]\times\mathbb{R})} + |R_K^Y|_{\mathcal{C}([\sigma_K,\sigma_K + T]\times\mathbb{R})}).$$

The terms  $|R_K^Y|_{\mathcal{C}([\sigma_K,\sigma_K+T]\times\mathbb{R})}$  is bounded by (2.18). As for  $|f_K^*|_{\mathcal{C}([0,T]\times\mathbb{R})}$ , with  $\int_0^\infty p(t, x - y)dy \leq 1$ , we have

$$|f_{K}^{*}(t,x)| \leq |U_{\star}(\sigma_{K}, \cdot - z_{\star}(\sigma_{K})) - U_{\star}(\sigma_{K}, \cdot - Z_{K}^{*}(\sigma_{K}))|_{\mathcal{C}(\mathbb{R})},$$
  
$$\leq |U_{\star}(\sigma_{K}, \cdot - z_{\star}(\sigma_{K})) - U_{\star}(\sigma_{K}, \cdot - Z_{K}^{*}(\sigma_{K}))|_{\mathcal{C}(\mathbb{R})} + |U_{K}(\sigma_{K}, \cdot) - U_{\star}(\sigma_{K}, \cdot)|_{\mathcal{C}(\mathbb{R})}.$$
(6.15)

With  $|\partial_x U_{\star}(\sigma_K, x)| \leq 2$ , The first term on (6.15) is at most  $2|Z_K^*(\sigma_K) - z_{\star}(\sigma_K)| \leq 2|Z_K^*(\sigma_K)| + 2|z_{\star}(\sigma_K)|$ . This, by (2.12) and Lemma 2.4, is bounded by  $CK^{-\beta}$  with probability  $1 - CK^{-n}$ . To complete the proof, we bound the last term in (6.15) as follows.

**Lemma 6.5.** Given any fixed  $n < \infty$  and  $\beta \in (0, \frac{1}{24})$ , there exists  $C = C(T, n, \beta) < \infty$  such that

$$\mathbf{P}\left(|U_K(s, \cdot) - U_{\star}(s, \cdot)|_{\mathcal{C}(\mathbb{R})} \le K^{-\beta}\right) \ge 1 - CK^{-n}, \quad \forall s \in [\frac{1}{2}, \sigma_K].$$
(6.16)

Proof. Fixing such n and  $\beta$ , we let  $C = C(n,\beta) < \infty$  denote a generic finite constant depending only on these variables. Fixing  $s \in [\frac{1}{2}, \sigma_K]$ , with  $U_K(s,x) = \widetilde{U}_K(s,0) - \widetilde{U}_K(s,x)$  and  $U_{\star}(s,x) = \widetilde{U}_{\star}(s,0) - \widetilde{U}_{\star}(s,x)$  (by Lemma 4.2), we clearly have  $|U_K(s,x) - U_{\star}(s,x)| \leq 2 \sup_{y \in \mathbb{R}} |\widetilde{U}_K(s,y) - \widetilde{U}_{\star}(s,y)|$ . To further bound the r.h.s., we take the difference of the integral

representations (2.7) and (2.1), and use (2.10)–(2.11) to bound the corresponding terms, thereby obtaining

$$|\widetilde{U}_K(s,y) - \widetilde{U}_\star(s,y)| \le C(\varepsilon)K^{1-\varepsilon} + C(\varepsilon)|Z_K|^{1-\varepsilon}_{\mathcal{C}([0,\sigma_K])} + |R_K(s,y)|,$$
(6.17)

for any fixed  $\varepsilon \in (0, 1)$ . Further applying (2.12) and (2.5) for bounding the last two terms, we obtain the desired bound (6.16).

6.3. **Proof of Propositions 2.5 and 2.8.** Fix  $\alpha \in (0, \frac{1}{24})$  and  $n, T < \infty$ , and let  $(Y_i^K(t))_i$ and  $Z_K^Y(t)$  be as in Section 6.1, for the *specific* initial condition  $U_{in}(y) = V(y) = U_\star(\frac{1}{2}, y - K^{-\alpha}) + 2(y \wedge K^{-\alpha})_+$ . With  $|V(\cdot) - U_\star(\frac{1}{2}, \cdot)|_{\mathcal{C}(\mathbb{R})} \leq CK^{-\alpha}$ , applying the same argument as in the proof of Lemma 6.4 one readily obtains

$$\mathbf{P}\left(|Z_K^Y(\boldsymbol{\cdot}) - z_\star(\frac{1}{2} + \boldsymbol{\cdot})|_{\mathcal{C}([0,T])} \le K^{-\alpha}\right) \ge 1 - C(T,\alpha,n)K^{-n}, \quad \forall K < \infty.$$
(6.18)

With this, Proposition 2.5 clearly follows from the following coupling result.

**Lemma 6.6.** There exist a coupling of  $Z_K(\frac{1}{2}+t)$  and  $Z_K^Y(t)$  such that

$$\mathbf{P}\left(Z_{K}^{Y}(t) - K^{-\alpha} \le Z_{K}(\frac{1}{2} + t), \ \forall t \le \tau_{ext}\right) \ge 1 - CK^{-n},\tag{6.19}$$

where  $C = C(\alpha, n) < \infty$  and  $\tau_{ext} := \inf\{t : X_i^K(\frac{1}{2} + t) = 0, \forall i\}$  is the extinction time for  $(X_i^K(t))_i$ .

Proof. Fixing such  $\alpha, n$ , we let  $C = C(\alpha, n) < \infty$  denote a generic finite constant that depend only on  $\alpha, n$ . Ordering the particles at t = 0:  $Y_1^K(0) < Y_2^K(0) < \cdots$ , and  $X_1^K(\frac{1}{2}) < X_2^K(\frac{1}{2}) < \cdots$ , we first claim that, regardless of the coupling, with probability  $1 - CK^{-n}$ , we have

$$|\{Y_i^K(0)\}| \ge |\{X_i^K(\frac{1}{2})\}|, \text{ and } Y_i^K(0) - K^{-\alpha} \le X_i^K(\frac{1}{2}), \text{ for } 1 \le i \le |\{X_i^K(\frac{1}{2})\}|, \quad (6.20)$$

or equivalently, with probability  $1 - CK^{-n}$ 

$$U_K^Y(0, x + K^{-\alpha}) \ge U_K(\frac{1}{2}, x), \quad \forall x \in \mathbb{R}_+.$$
 (6.21)

Indeed, by Lemma 6.5 and (6.7), we have  $U_K(\frac{1}{2}, x) \leq U_\star(\frac{1}{2}, x) + K^{-\alpha}$  and  $V(x + K^{-\alpha}) \leq U_K^Y(0, x + K^{-\alpha}) - K^{\alpha}, \forall x \in \mathbb{R}_+$ , with probability  $1 - CK^{-n}$ . Combining these with  $V(x + K^{-\alpha}) \geq U(x) + 2K^{-\alpha}, \forall x \geq 0$ , we see that (6.21) follows.

Now, with (6.20), we proceed to constructing the coupling for t > 0. Let  $\tau_1 := \inf\{t \ge 0 : Z_K(\frac{1}{2}+t) = 0\}$  be the first hitting time of  $Z_K(\frac{1}{2}+\cdot)$  at x = 0. For  $t \in [0, \tau_1)$ , both processes  $(Y_i^K(t))_i$  and  $(X_i^K(t))_i$  evolve as  $\sqrt{K}$ -Atlas models, so by Lemma 6.1(a), with probability  $1 - CK^{-n}$ ,

$$Y_i^K(t) - K^{-\alpha} \le X_i^K(\frac{1}{2} + t)$$
 for  $1 \le i \le |\{X_i^K(\frac{1}{2})\}|$  and  $0 \le t < \tau_1$ .

As  $(X_i^K(\frac{1}{2}+t))_i$  loses a particle at time  $t = \tau_1$ , by ordering  $\{Y_i^K(\tau_1)\}$  and  $\{X_i^K(\frac{1}{2}+\tau_1)\}$ , we retain the type of ordering as in (6.20). As  $Z_K(\frac{1}{2}+t)$  hits 0 at most  $|\{X_i^K(\frac{1}{2})\}| < \infty$  times, iterating the prescribed procedure yields the desired coupling.

Turning to the proof of Proposition 2.8, we fix  $n < \infty$ . By Lemma 6.6, it suffices to show

$$\mathbf{P}(\inf_{t \ge 1} Z_K^Y(t) > 0) \ge 1 - C(n)K^{-n}, \quad \forall K < \infty.$$
(6.22)

We prove this by a scaling argument as follows. For  $m \in \mathbb{N}$ , we let  $K_m := 2^m K$  and  $Y_i^{K,m}(t) := 2^{\frac{m}{2}} Y_i^{K_m}(\frac{t}{2^m})$ . The scaling here is chosen so that each  $Y^{K,m}(t)$  is a  $\sqrt{K}$ -Atlas model. With V(x) defined as in the preceding, by (1.10)–(1.11) we have that

$$\frac{d}{dx}V(x) := v(x) = \begin{cases} 2 , & \text{for } x \in (0, K^{-\alpha}), \\ -2\partial_x p(\frac{1}{2}, x - K^{-\alpha}) + 4\widetilde{\Phi}(\frac{1}{2}, x - K^{-\alpha}), & \text{for } x \in (K^{-\alpha}, \infty). \end{cases}$$
(6.23)

Now, with  $(Y_i^K(0))_i \sim \text{PPP}_+(\sqrt{K}v)$ , we have  $(Y_i^{K,m}(0))_i \sim \text{PPP}_+(\sqrt{K}v_m)$ , where  $v_m(x) := v(2^{-m/2}x)$ . As v(x) is decreasing for all  $x \in \mathbb{R}_+$  (which is verified by differentiating v(x), as in (6.23)), we have  $v \leq v_1 \leq v_2 \leq \cdots$ , so there exists a coupling of  $(Y_i^{K,m}(t))_i, m \geq 1$ , such that

$$\begin{aligned} |\{Y_i^{K,1}(0)\}_i| &\leq |\{Y_i^{K,2}(0)\}_i| \leq \dots, \\ Y_i^{K,m}(0) &\geq Y_i^{K,m+1}(0), \quad \forall 1 \leq i \leq |\{Y_i^{K,m}(0)\}_i|, \ m \geq 1. \end{aligned}$$

Letting  $Z_K^m(t) := \min\{Y_i^{K,m}(t)\}$ , by Lemma 6.1(b), we then have a coupling of  $Z_K^Y(t)$  and  $Z_{K,m}^Y(t)$  such that  $Z_K^Y(t) \ge Z_{K,m}^Y(t) = 2^{\frac{m}{2}} Z_{K_m}^Y(\frac{t}{2^m}), \forall t \ge 0$ . Consequently,

$$\inf_{2^m \le t \le 2^{m+1}} Z_K^Y(t) \ge \inf_{1 \le t \le 2} \{ 2^{\frac{m}{2}} Z_{K_m}^Y(t) \}.$$
(6.24)

By Lemma 2.4, we have  $z_{\star}(\frac{1}{2}+t) > K^{-\alpha} > 0$ ,  $\forall t \ge 1$ ,  $\alpha > 0$  and K large enough. Hence by (6.18), we have  $\mathbf{P}(\inf_{1\le t\le 2} Z_{K_m}^Y(t) > 0) \ge 1 - C(n)K_m^{-n} = 1 - C(n)2^{-mn}K^{-n}$ . Combining this with (6.24), and applying the union bound over  $m \in \mathbb{N}$ , we conclude (6.22).

#### References

- [Ald02] D. Aldous. Unpublished, available at http://www.stat.berkeley.edu/~aldous/Research/OP/river.pdf. 2002.
- [BFK05] A. D. Banner, R. Fernholz, and I. Karatzas. Atlas models of equity markets. Ann. Appl. Probab., 15(4):2296–2330, 2005.
- [CDSS] M. Cabezas, A. Dembo, A. Sarantsev, and V. Sidoravicius. Brownian particles of rank-dependent drifts: out of equilibrium behavior. *In preperation.*
- [CP10] S. Chatterjee and S. Pal. A phase transition behavior for Brownian motions interacting through their ranks. *Probab. Theory Related Fields*, 147(1-2):123–159, 2010.
- [CP11] S. Chatterjee and S. Pal. A combinatorial analysis of interacting diffusions. J. Theoret. Probab., 24(4):939–968, 2011.
- [DSVZ12] A. Dembo, M. Shkolnikov, S. Varadhan, and O. Zeitouni. Large deviations for diffusions interacting through their ranks. To appear Comm. Pure Appl. Math., 2012.
- [DT15] A. Dembo and L.-C. Tsai. Equilibrium fluctuation of the Atlas model. 2015. arXiv:1503.03581.
- [Fel71] W. Feller. An introduction to probability theory and its applications, volume II of Second edition. John Wiley & Sons, Inc., New York-London-Sydney, 1971.
- [Fer02] E. R. Fernholz. Stochastic portfolio theory. Springer, 2002.
- [Fri10] A. Friedman. Variational principles and free-boundary problems. Courier Corporation, 2010.
- [Han13] W. Han. http://hanweijian.com/research/2013-research-projects/random-particle-motion/. 2013.
- [HJV15] F. Hernández, M. Jara, and F. J. Valentim. Equilibrium fluctuations for a discrete Atlas model. arXiv preprint arXiv:1507.04786, 2015.
- [IK10] T. Ichiba and I. Karatzas. On collisions of Brownian particles. The Annals of Applied Probability, 20(3):951–977, 2010.
- [IKS13] T. Ichiba, I. Karatzas, and M. Shkolnikov. Strong solutions of stochastic equations with rankbased coefficients. Probab. Theory Related Fields, 156(1-2):229–248, 2013.
- [IPB<sup>+</sup>11] T. Ichiba, V. Papathanakos, A. Banner, I. Karatzas, and R. Fernholz. Hybrid Atlas models. Ann. Appl. Probab., 21(2):609–644, 2011.

- [Kun97] H. Kunita. Stochastic flows and stochastic differential equations, volume 24. Cambridge University press, 1997.
- [MS06] H. McKean and L. A. Shepp. The advantage of capitalism vs. socialism depends on the criterion. Journal of Mathematical Sciences, 139(3):6589–6594, 2006.
- [PP08] S. Pal and J. Pitman. One-dimensional Brownian particle systems with rank-dependent drifts. Ann. Appl. Probab., pages 2179–2207, 2008.
- [PS14] S. Pal and M. Shkolnikov. Concentration of measure for Brownian particle systems interacting through their ranks. *Ann. Appl. Probab.*, 24(4):1482–1508, 2014.
- [Sar15] A. Sarantsev. Comparison techniques for competing Brownian particles. 2015. arXiv:1305.1653.
- [Shk11] M. Shkolnikov. Competing particle systems evolving by interacting Levy processes. Ann. Appl. Probab., 21(5):1911–1932, 2011.

STATISTICS DEPARTMENT, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720 *E-mail address*: wenpintang@stat.berkeley.edu

L.-C. TSAI, DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, CA 94305 *E-mail address*: lctsai@stanford.edu