

THE SORTED EFFECTS METHOD: DISCOVERING HETEROGENEOUS EFFECTS BEYOND THEIR AVERAGES

VICTOR CHERNOZHUKOV, IVAN FERNANDEZ-VAL, YE LUO

ABSTRACT. The partial (*ceteris paribus*) effects of interest in nonlinear and interactive linear models are heterogeneous as they can vary dramatically with the underlying observed or unobserved covariates. Despite the apparent importance of heterogeneity, a common practice in modern empirical work is to largely *ignore it* by reporting average partial effects (or, at best, average effects for some groups). While average effects provide very convenient scalar summaries of typical effects, by definition they fail to reflect the entire variety of the heterogeneous effects. In order to discover these effects much more fully, we propose to estimate and report *sorted effects* – a collection of estimated partial effects sorted in increasing order and indexed by percentiles. By construction the sorted effect curves *completely represent* and help visualize the range of the heterogeneous effects in one plot. They are as convenient and easy to report in practice as the conventional average partial effects. They also serve as a basis for *classification analysis*, where we divide the observational units into most or least affected groups and summarize their characteristics. We provide a quantification of uncertainty (standard errors and confidence bands) for the estimated sorted effects and related classification analysis, and provide confidence sets for the most and least affected groups. The derived statistical results rely on establishing key, new mathematical results on Hadamard differentiability of a multivariate sorting operator and a related classification operator, which are of independent interest.

We apply the *sorted effects method* and classification analysis to demonstrate several striking patterns in the gender wage gap. We find that this gap is particularly strong for married women, ranging from -60% to 0% between the 2% and 98% percentiles, as a function of observed and unobserved characteristics; while the gap for never married women ranges from -40% to $+20\%$. The most adversely affected women tend to be married, do not have college degrees, work in sales, and have high levels of potential experience.

Keywords: Sorting, Partial Effect, Marginal Effect, Sorted Effect, Classification Analysis, Nonlinear Model, Functional Analysis, Differential Geometry, Gender Wage Gap

1. INTRODUCTION

In nonlinear and interactive linear models the partial (*ceteris paribus*) effects of interest often vary with respect to the underlying covariates. For example, consider a binary response model with conditional choice probability $P(Y = 1 \mid X) = F(X^\top \beta)$, where Y is a binary response variable, X is a vector of covariates, F is a distribution function such as the standard normal or logistic, and β is a vector of coefficients. The partial or predictive effect (PE) of a marginal

Date: May 29, 2018.

MIT, BU and University of Florida. First version of October, 2011. We thank the coeditor, three anonymous referees, Alberto Abadie, Daron Acemoglu, Isaiah Andrews, Joshua Angrist, Debopam Bhattacharya, Jaroslav Borovicka, Kasey Buckles, Denis Chetverikov, Jerry Hausman, Tetsuya Kaji, Pat Kline, Roger Koenker, Siyi Luo, Maddie McKelway, Anna Mikusheva, Whitney Newey, Claudia Olivetti, Andres Santos, Camille Terrier, Frank Windmeijer, Abigail Wozniak, and seminar participants at multiple institutions for helpful discussions. We gratefully acknowledge research support from the National Science Foundation.

change in a continuous covariate X_j with coefficient β_j on the conditional choice probability is

$$\Delta(X) = f(X^\top \beta) \beta_j, \quad f(v) = \partial F(v) / \partial v,$$

which generally varies in the population of interest with the covariate vector X , as X varies according to some distribution, say μ . A common empirical practice is to report the average partial effect (APE),

$$E[\Delta(X)] = \int \Delta(x) d\mu(x),$$

as a single summary measure of the PE (e.g., Wooldridge (2010, Chap. 2)), or to report effects for some groups (e.g., Angrist and Pischke (2008)). However, the APE completely disregards the heterogeneity of the PE and may give a very incomplete picture of the impact of the covariates.

In this paper we propose complementing the APE by reporting the entire set of PEs sorted in increasing order and indexed by a ranking with respect to the distribution of the covariates in the population of interest. These sorted effects correspond to percentiles of the PE,

$$\Delta_\mu^*(u) = u^{\text{th}}\text{-quantile of } \Delta(X), \quad X \sim \mu,$$

and provide a more complete representation of the heterogeneity of $\Delta(X)$. We shall call these effects as sorted predictive or partial effects (SPE) by default, as most models are predictive.¹ We also show how to use the SPEs to carry out classifications analysis (CA). This analysis consists of classifying the observational units into most or least affected depending on whether their PEs are above or below some tail SPE, and then comparing the moments or distribution of the covariates of the most and least affected groups.

Heterogeneous effects also arise in the most basic linear models with interactions (Oaxaca, 1973; Cox, 1984). Consider a conditional mean model for the Mincer earnings function:

$$Y = P(T, W)^\top \beta + \epsilon, \quad E[\epsilon | T, W] = 0, \quad X = (T, W),$$

where Y is log wage, T is an indicator of gender (or race, treatment, or program participation), and W is a vector of labor market characteristics. The vector $P(T, W)$ is a collection of transformations of T and W , involving some interaction between T and W . For example, Oaxaca (1973) used the specification $P(T, W) = (TW, (1 - T)W)$. Then, the PE of changing $T = 0$ to $T = 1$ is

$$\Delta(X) = P(1, W)^\top \beta - P(0, W)^\top \beta,$$

which is a measure of the gender wage gap conditional on worker characteristics. The function $u \mapsto \Delta_\mu^*(u)$ provides again a complete summary of the entire range of PEs. The left panel of Figure 1 illustrates the SPE of the conditional gender wage gap for women. The SPE varies sharply from around -40 to 6.5% , and does not coincide with the average PE of -20% . The PE is especially (negatively) large for women who have any of the following characteristics: married, low educated, high experience, and working on sales occupations – this follows from the classification analysis,

¹When the underlying model has a structural or causal interpretation, we may use the name sorted structural effects or sorted treatment effects.




FIGURE 1. APE and SPE (introduced in this paper) of the gender wage gap for women. Estimates and 90% bootstrap uniform confidence bands (derived in this paper) based on a linear model with interactions for the conditional expectation (left) and quantile (right) functions.

where we compare the average characteristics of the subpopulations of women with covariate values X such that $\Delta(X)$ is above the 90% percentile and below the 10% percentile. We refer the reader to Section 3 for a detailed discussion of this example.

The general settings that we deal with in this paper as well as the specific results we obtain are as follows: Let X denote a covariate vector, $\Delta(X)$ denote a generic PE of interest, μ denote the distribution of X in the population of interest, and \mathcal{X} denote the interior of the support of X in this population. The SPE is obtained by sorting the multivariate function $x \mapsto \Delta(x)$ in increasing order with respect to μ . Using tools from differential geometry, we prove that this multivariate sorting operator is Hadamard differentiable with respect to the PE function Δ and the distribution μ at the regular values of $x \mapsto \Delta(x)$ on \mathcal{X} . This key and new mathematical result allows us to derive the large sample properties of the empirical SPE, which replace Δ and μ by sample analogs, obtained from parametric or semi-parametric estimators, using the functional delta method. In particular, we derive a functional central limit theorem and a bootstrap functional central limit theorem for the empirical SPE. The main requirement of these theorems is that the empirical Δ and μ also satisfy functional central limit theorems, which hold for many estimators used in empirical economics under general sampling conditions. We use the properties of the empirical SPE to construct confidence sets for the SPE that hold uniformly over quantile indices. We also show under the same conditions that the empirical version of the objects in the classification analysis follow functional central limit theorems and bootstrap functional central limit theorems.

We derive these result by establishing the Hadamard differentiability of a classification operator related to the multivariate sorting operator.

Related technical literature: Previously, Chernozhukov, Fernández-Val, and Galichon (2010) derived the properties of the rearrangement (sorting) operator in the univariate case with known μ (standard uniform distribution). Those results were motivated by a completely different problem – namely, restoration of monotonicity in conditional quantile estimation – rather than the problem of summarizing heterogeneous effects by the SPEs. These prior technical results are not applicable to our case as soon as the dimension of X is greater than one, which is the case in all modern applications where effects are of interest. Moreover, the previous results are not applicable even in the univariate case since the measure μ is not known in all envisioned applications. The properties of the sorting operator are different in the multivariate case and require tools from differential geometry: computation of functional (Hadamard) derivatives of the sorting operator with respect to perturbations of Δ require us to work with integration on $(d_x - 1)$ -dimensional manifolds of the type $\{\Delta(x) = \delta\}$, where $d_x = \dim X$. Moreover, we also need to compute functional derivatives with respect to suitable perturbations of the measure μ . In econometrics or statistics, Sasaki (2015) also used differential geometry to characterize the structural properties of derivatives of conditional quantile functions in nonseparable models; and Kim and Pollard (1990) used tools from differential geometry to derive the large sample properties of the maximum score and other cube root consistent estimators. Relative to these papers, we share the use of differential geometry tools as a general proof strategy, but we apply these tools to establish the analytical properties of different functionals – namely, the SPEs. Moreover, our results on the functional differentiability of the sorting and classification operators in the multivariate case constitute new mathematical results, which are of interest in their own right.

Organization of the paper: In Section 2 we discuss the quantities of interest in nonlinear and interactive linear models with examples; introduce the SPE and related CA, along with their empirical counterparts; and outline the main inferential results. In Section 3 we provide an empirical application to the gender wage gap in the U.S. in 2015. We derive the properties of the empirical SPE and CA in large samples and show how to use these properties to make inference in Section 4. Appendix A provides some key mathematical results on the differentiability of the multivariate sorting and classification operators and Appendix B contains the proof of the main results. All other proofs are given in the online appendix with supplementary material (SM), which also contains additional technical material, and results from Monte Carlo simulations and an empirical application to mortgage denials using binary response models (Chernozhukov, Fernandez-Val, and Luo, 2017).

2. SORTED EFFECTS AND CLASSIFICATION ANALYSIS

We start by discussing the objects of interest in nonlinear and interactive linear models.

2.1. Effects of Interest. We consider a general model characterized by a predictive function $g(X)$, where X is a d_x -vector of covariates that may contain unobserved components, as in quantile regression models. The function g usually arises from a model for a response variable Y , which can be discrete or continuous. We call the function g predictive because the underlying model can be either predictive or causal under additional assumptions, but we do not insist on estimands having a causal interpretation. For example, in a mean regression model, $g(X) = E[Y | X]$ corresponds to the expectation function of Y conditional on X ; in a binary response model, $g(X) = P[Y = 1 | X]$ corresponds to the choice probability of $Y = 1$ conditional on X ; in a quantile regression model, $g(X) = Q_Y[\epsilon | Z]$, where the covariate $X = (\epsilon, Z)$ consists of the unobservable rank variable ϵ with a uniform distribution, $\epsilon | Z \sim U(0, 1)$, and the observed covariate vector Z , and where $Q_Y[\tau | Z]$ is the conditional τ^{th} -quantile of Y given Z .

Let $X = (T, W)$, where T is the key covariate or treatment of interest, and W is a vector of control variables. We are interested in the effects of changes in T on the function g holding W constant. These effects are usually called partial effects, marginal effects, or treatment effects. We call them predictive effects (PE) throughout the paper, as such a name most accurately describes the meaning of the estimand (especially when a causal interpretation is not available). If T is discrete, the PE is

$$\Delta(x) = \Delta(t, w) = g(t_1, w) - g(t_0, w), \quad (2.1)$$

where t_1 and t_0 are two values of T that might depend on t (e.g., $t_0 = 0$ and $t_1 = 1$, or $t_0 = t$ and $t_1 = t + 1$). This PE measures the effect of changing T from t_0 to t_1 holding W constant at w . If T is continuous and $t \mapsto g(t, w)$ is differentiable, the PE is

$$\Delta(x) = \Delta(t, w) = \partial_t g(t, w), \quad (2.2)$$

where ∂_t denotes $\partial/\partial t$, the partial derivative with respect to t . This PE measures the effect of a marginal change of T from the level t holding W constant at w .²

We consider the following examples in the empirical applications of Section 3 and SM.

Example 1 (Binary response model). Let Y be a binary response variable such as an indicator for mortgage denial, and X be a vector of covariates related to Y . The predictive function of the probit or logit model takes the form:

$$g(X) = P(Y = 1 | X) = F(P(X)^\top \beta),$$

where $P(X)$ is a vector of known transformations of X , β is a parameter vector, and F is a known distribution function (the standard normal distribution function in the probit model or standard logistic distribution function in the logit model). If T is a binary variable such as an indicator

²We can also consider high-order and crossed effects. For example, $\Delta(x) = \partial_{t_2}^2 g(t, w)$ gives the second-order PE of the continuous treatment T if $t \mapsto g(t, w)$ is twice differentiable; and, letting $X = (T, S, W)$ where T and S are discrete, $\Delta(x) = g(t_1, s_1, w) - g(t_0, s_1, w) - g(t_1, s_0, w) + g(t_0, s_0, w)$ gives the crossed effect or interaction of T and S .

for black applicant and W is a vector of controls such as the applicant characteristics relevant for the bank decision, the PE,

$$\Delta(x) = F(P(1, w)^\top \beta) - F(P(0, w)^\top \beta),$$

describes the difference in predicted probability of mortgage denial for a black applicant and a white applicant, conditional on a specific value w of the observable characteristics W . \square

Example 2 (Interactive linear model with additive error). Let Y be the logarithm of the wage. Suppose $X = (T, W)$, where T is an indicator for female worker and W are other worker characteristics. We can model the conditional expectation function of log wage using the linear interactive model:

$$Y = g(X) + \epsilon = P(T, W)^\top \beta + \epsilon, \quad E[\epsilon | T, W] = 0, \quad X = (T, W),$$

where $P(T, W)$ is a collection of transformations of T and W , involving some interaction between T and W . For example, $P(T, W) = (TW, (1 - T)W)$. Then the PE

$$\Delta(x) = P(1, w)^\top \beta - P(0, w)^\top \beta$$

is the (average) gender wage gap or difference between the expected log wage of a woman and a man, conditional on a specific value w of the characteristics W . \square

Example 3 (Linear model with non-additive error, or QR model). Let Y be log wage, T be an indicator for female worker, and W be a vector of worker characteristics as in the previous example. Suppose we model the conditional quantile function of log wage using the linear interactive model:

$$Y = g(X) = P(T, W)^\top \beta(\epsilon), \quad \epsilon | T, W \sim U(0, 1), \quad X = (T, W, \epsilon),$$

where $P(T, W)^\top \beta(\tau)$ is the conditional τ^{th} -quantile of Y given T and W . Thus the covariate vector $X = (T, W, \epsilon)$ includes the observed covariates (T, W) as well as the rank variable ϵ , which is an unobserved factor (e.g., ‘‘ability rank’’). Here $P(T, W)$ is a collection of transformations of T and W , e.g., $P(T, W) = (TW, (1 - T)W)$. Then the PE

$$\Delta(x) = P(1, w)^\top \beta(\tau) - P(0, w)^\top \beta(\tau), \quad x = (t, w, \tau),$$

is the (τ^{th} -quantile) gender wage gap or difference between the conditional τ^{th} -quantile of log-wage of a woman and a man, conditional on a specific value w of the characteristics W .

Note that in this case,

$$X \sim \mu, \quad \mu = F_{T,W} \times F_\epsilon,$$

where F_ϵ is the distribution function of the standard uniform random variable, and $F_{T,W}$ is the distribution of (T, W) . For estimation purposes, we will have to exclude the tail quantile indices, so F_ϵ will be redefined to have support on a set of the form $[\ell, 1 - \ell]$, where $0 < \ell < 0.5$ is a small positive number. \square

The set of examples listed above are the most basic, leading cases, arising mostly in predictive analysis and program evaluation. Our theoretical results are rather general and are not limited to these cases. Thus, they allow for both Δ and μ to originate from causal or structural models and to be estimated by structural methods. For example, in treatment effects models with selection on observables (Rosenbaum and Rubin, 1983), the PE is the conditional average treatment effect $\Delta(x) = E(Y_1 - Y_0 \mid X = x)$, where Y_1 and Y_0 are potential outcomes in the treated and non-treated statuses and X is a vector of covariates. The standard approach is to aggregate the conditional average treatment effects by integration with respect to the distribution of the covariates in the population of interest μ . This yields the average treatment effect if μ is the distribution in the entire population or the average treatment effect on the treated if μ is the distribution in the treated subpopulation. The SPE can be used to complement the analysis by reporting the entire range of conditional average treatment effects, and also to determine the optimal treatment allocation with budget constraints. Thus, Bhattacharya and Dupas (2012) showed that under some conditions this optimal allocation has a cutoff determined by a tail percentile of the conditional average treatment effects, i.e. by a SPE. Another example is the welfare analysis described in Hausman and Newey (2017), where $\Delta(x)$ is the compensating or equivalent variation of a price change conditional on covariates such as income and demographic characteristics, and μ is the distribution of covariates in the population of interest.

In all the previous examples, the PE $\Delta(x)$ is a function of x and therefore can be different for each observational unit. To summarize this effect in a single measure, a common practice in empirical economics is to average the PEs. Averaging, however, masks most of the heterogeneity in the PE allowed by nonlinear or interactive linear models. We propose reporting the entire set of values of the PE sorted in increasing order and indexed by a ranking $u \in [0, 1]$ with respect to the population of interest. These sorted effects provide a more complete representation of the heterogeneity in the PE than the average effects.

Definition 2.1 (*u-SPE*). The u^{th} -sorted predictive effect with respect to μ is

$$\Delta_\mu^*(u) := \inf\{\delta \in \mathbb{R} : F_{\Delta, \mu}(\delta) \geq u\}, \quad F_{\Delta, \mu}(\delta) := E_\mu[1\{\Delta(X) \leq \delta\}],$$

where E_μ denotes expectation with respect to μ .

The u -SPE is the u^{th} -quantile of $\Delta(X)$ when X is distributed according to μ . As for the average effect, μ can be chosen to select a target subpopulation from the entire population. For example, when T is a treatment indicator:

- If μ is set to the marginal distribution of X in the entire population, then $\Delta_\mu^*(u)$ is the *population u-SPE*.
- If μ is set to the distribution of X conditional on $T = 1$, then $\Delta_\mu^*(u)$ is the *u-SPE on the treated or exposed*.

By considering $\Delta_\mu^*(u)$ at multiple quantile indices, we obtain a one-dimensional representation of the heterogeneity of the PE. Accordingly, our object of interest is the *SPE-function*

$$\{u \mapsto \Delta_\mu^*(u) : u \in \mathcal{U}\}, \quad \mathcal{U} \subseteq [0, 1],$$

where \mathcal{U} is the set of quantile indices of interest.

We also show how to use the u -SPE for classification analysis. Let $u \in \mathcal{U}$, with $u < 1/2$, and Z be a d_z -dimensional random vector that includes X and possibly other variables such as Y in Examples 1–3. By abuse of notation, we also denote the distribution of Z over its support \mathcal{Z} as μ .

Definition 2.2 (u -CA). The u^{th} -classification analysis consists of 2 steps: (i) Assign all Z with $\Delta(X) < \Delta_\mu^*(u)$ to the u -least affected subpopulation, and all Z with $\Delta(X) > \Delta_\mu^*(1 - u)$ to the u -most affected subpopulation. (ii) Obtain the moments and distribution of Z in the least and most affected subpopulations. We denote by $\Lambda_{\Delta, \mu}^{-u}(t)$ and $\Lambda_{\Delta, \mu}^{+u}(t)$ generic objects indexed by $t \in \mathbb{R}^{d_z}$ in the least and most affected subpopulations, respectively. For example, $\Lambda_{\Delta, \mu}^{-u}(t) = \mathbb{E}_\mu[Z^t \mid \Delta(X) < \Delta_\mu^*(u)]$ corresponds to the t -moment of Z in the u -least affected subpopulation, and $\Lambda_{\Delta, \mu}^{-u}(t) = \mathbb{E}_\mu[1(Z \leq t) \mid \Delta(X) < \Delta_\mu^*(u)]$ to the distribution of Z at t in the u -least affected subpopulation.³ We define the same quantities in the u -most affected subpopulation replacing $\Delta(X) < \Delta_\mu^*(u)$ by $\Delta(X) > \Delta_\mu^*(1 - u)$ in the conditioning set.

2.2. Empirical SPE. In practice, we replace the PE Δ and the distribution μ by sample analogs to construct plug-in estimators of the SPE. Let $\widehat{\Delta}(x)$ and $\widehat{\mu}(x)$ be estimators of $\Delta(x)$ and $\mu(x)$ obtained from $\{(Y_i, T_i, W_i) : 1 \leq i \leq n\}$, an independent and identically distributed sample of size n from (Y, T, W) .

Definition 2.3 (Empirical u -SPE). The estimator of Δ_μ^* is

$$\widehat{\Delta}_\mu^*(u) := \widehat{\Delta}_{\widehat{\mu}}^*(u) = \inf\{\delta \in \mathbb{R} : F_{\widehat{\Delta}, \widehat{\mu}}(\delta) \geq u\}, \quad F_{\widehat{\Delta}, \widehat{\mu}}(\delta) = \mathbb{E}_{\widehat{\mu}}[1\{\widehat{\Delta}(X) \leq \delta\}] =: \widehat{F}_{\widehat{\Delta}, \widehat{\mu}}(\delta).$$

Then the empirical SPE-function is

$$\{u \mapsto \widehat{\Delta}_\mu^*(u) : u \in \mathcal{U}\}, \quad \mathcal{U} \subseteq [0, 1],$$

where \mathcal{U} is the set of indices of interest that typically excludes tail indices and satisfies other technical conditions stated in Section 4.

Example 1 (Binary response model, cont.) The estimator of the PE is

$$\widehat{\Delta}(x) = F\left(P(1, w)^\top \widehat{\beta}\right) - F\left(P(0, w)^\top \widehat{\beta}\right),$$

³For a d_z -dimensional random variable $Z = (Z_1, \dots, Z_{d_z})$ and $t = (t_1, \dots, t_{d_z}) \in \mathbb{R}^{d_z}$, we denote $Z^t := \prod_{j=1}^{d_z} Z_j^{t_j}$ and $\{Z \leq t\} := \{Z_1 \leq t_1, \dots, Z_{d_z} \leq t_{d_z}\}$.

where $\widehat{\beta}$ is the maximum likelihood (ML) estimator of β ,

$$\widehat{\beta} \in \arg \max_{b \in \mathbb{R}^{d_p}} \sum_{i=1}^n [Y_i \log F(P(T_i, W_i)^\top b) + (1 - Y_i) \log \{1 - F(P(T_i, W_i)^\top b)\}], \quad d_p = \dim P(T, W).$$

□

Example 2 (Interactive linear model with additive error, cont.) The estimator of the PE is

$$\widehat{\Delta}(x) = P(1, w)^\top \widehat{\beta} - P(0, w)^\top \widehat{\beta},$$

where $\widehat{\beta}$ is the ordinary least squares (OLS) estimator of β ,

$$\widehat{\beta} \in \arg \min_{b \in \mathbb{R}^{d_p}} \sum_{i=1}^n [Y_i - P(T_i, W_i)^\top b]^2, \quad d_p = \dim P(T, W).$$

□

Example 3 (Linear model with non-additive error, cont.) The estimator of the PE is

$$\widehat{\Delta}(x) = P(1, w)^\top \widehat{\beta}(\tau) - P(0, w)^\top \widehat{\beta}(\tau),$$

where $\widehat{\beta}(\tau)$ is the Koenker and Basset (1978) quantile regression (QR) estimator of $\beta(\tau)$,

$$\widehat{\beta}(\tau) \in \arg \min_{b \in \mathbb{R}^{d_p}} \sum_{i=1}^n \rho_\tau(Y_i - P(T_i, W_i)^\top b), \quad d_p = \dim P(T, W), \quad \rho_\tau(v) = (\tau - 1\{v < 0\})v.$$

□

Remark 2.1 (Estimation of μ). Let S denote the indicator for an observational unit belonging to the subpopulation of interest. For example, if $S = T$, then $S = 1$ indicates the unit is in the subpopulation of the treated and $S = 0$ indicates the unit is in the subpopulation of the untreated. The indicator S can also incorporate other restrictions, for example $S = 1\{X \in \mathcal{X}\}$ restricts the support of covariate X to the region \mathcal{X} . Finally, if S is always 1, then this means that we work with the entire population. Estimation of μ can be done using the empirical distribution:

$$\widehat{\mu}(x) = \sum_{i=1}^n S_i 1\{X_i \leq x\} / \sum_{i=1}^n S_i,$$

provided that $\sum_{i=1}^n S_i > 0$. An alternative would be to use the smoothed empirical distribution.

If μ can be decomposed into known and unknown parts, then we only need to estimate the unknown parts. Thus, $\mu = F_{T,W} \times F_\epsilon$ in Example 3, where F_ϵ is known to be the uniform distribution and $F_{T,W}$ is unknown, but can be estimated by the empirical distribution of (T, W) in the part of the population of interest. □

2.3. Empirical CA. The empirical version of the u -CA classifies the observations in the sample using the empirical PEs and u -SPE, and computes the moments and distributions in the resulting most and least affected subsamples.

Definition 2.4 (Empirical u -CA). The empirical u^{th} -classification analysis consists of 2 steps: (1) Assign all Z_i with $\widehat{\Delta}(X_i) < \widehat{\Delta}_\mu^*(u)$ to the u -least affected subsample, and all Z_i with $\widehat{\Delta}(X_i) > \widehat{\Delta}_\mu^*(1-u)$ to the u -most affected subsample. (2) Estimate the moments and distribution of Z in the least and most affected subpopulations by the empirical analogs in the least and most affected subsamples, i.e. $\widehat{\Lambda}_{\Delta,\mu}^{-u}(t) = \Lambda_{\widehat{\Delta},\widehat{\mu}}^{-u}(t)$ and $\widehat{\Lambda}_{\Delta,\mu}^{+u}(t) = \Lambda_{\widehat{\Delta},\widehat{\mu}}^{+u}(t)$. For example, $\widehat{\Lambda}_{\Delta,\mu}^{-u}(t) = \mathbb{E}_{\widehat{\mu}} \left[Z^t \mid \widehat{\Delta}(X) < \widehat{\Delta}_\mu^*(u) \right]$ estimates the t -moment of Z in the u -least affected subpopulation and $\widehat{\Lambda}_{\Delta,\mu}^{-u}(t) = \mathbb{E}_{\widehat{\mu}} \left[1(Z \leq t) \mid \widehat{\Delta}(X) < \widehat{\Delta}_\mu^*(u) \right]$ the distribution of Z at t in the u -least affected subpopulation. The corresponding estimators in the u -most affected subpopulation are constructed replacing $\widehat{\Delta}(X) < \widehat{\Delta}_\mu^*(u)$ by $\widehat{\Delta}(X) > \widehat{\Delta}_\mu^*(1-u)$ in the conditioning set. Here we use the same notation as in Definition 2.2.

2.4. Inference on SPE. The main inferential result for the SPE can be previewed as follows. Assume that the PE function $x \mapsto \Delta(X)$ is not locally flat in the sense that the norm of its gradient does not vanish anywhere over the support, and other regularity conditions stated in Section 4. Then, the empirical SPE-process is \sqrt{n} -consistent and converges in distribution to a centered Gaussian process, namely

$$\sqrt{n}(\widehat{\Delta}_\mu^*(u) - \Delta_\mu^*(u)) \rightsquigarrow Z_\infty(u) \text{ in } \ell^\infty(\mathcal{U}),$$

the metric space of bounded functions on \mathcal{U} , as a stochastic process indexed by $u \in \mathcal{U}$, where \mathcal{U} is a compact subset of $(0, 1)$. Moreover, the exchangeable bootstrap algorithm specified in Algorithm 2.1 estimates consistently the law of $Z_\infty(u)$.

The next corollary to Theorem 4.1 in Section 4 provides uniform bands that cover the SPE-function simultaneously over a region of values of u with prespecified probability in large samples. It does cover pointwise confidence bands for the SPE-function at a specific quantile index u as a special case by simply taking \mathcal{U} to be the singleton set $\{u\}$.

Corollary 2.1 (Inference on SPE-function using Limit Theory and Bootstrap). *Under the assumptions of Theorem 4.1, for any $0 < \alpha < 1$,*

$$\mathbb{P} \left\{ \Delta_\mu^*(u) \in \left[\widehat{\Delta}_\mu^*(u) - \widehat{t}_{1-\alpha}(\mathcal{U}) \widehat{\Sigma}(u)^{1/2} / \sqrt{n}, \widehat{\Delta}_\mu^*(u) + \widehat{t}_{1-\alpha}(\mathcal{U}) \widehat{\Sigma}(u)^{1/2} / \sqrt{n} \right] : u \in \mathcal{U} \right\} \rightarrow 1 - \alpha,$$

where $\widehat{t}_{1-\alpha}(\mathcal{U})$ is any consistent estimator of $t_{1-\alpha}(\mathcal{U})$, the $(1-\alpha)$ -quantile of

$$t(\mathcal{U}) := \sup_{u \in \mathcal{U}} |Z_\infty(u)| \Sigma(u)^{-1/2},$$

and $u \mapsto \widehat{\Sigma}(u)$ is a uniformly consistent estimator of $u \mapsto \Sigma(u)$, the variance function of $u \mapsto Z_\infty(u)$. We provide consistent estimators of $t_{1-\alpha}(\mathcal{U})$ and $u \mapsto \Sigma(u)$ in Algorithm 2.1.

We now describe a practical bootstrap algorithm to estimate the quantiles of $t(\mathcal{U})$. Let $(\omega_1, \dots, \omega_n)$ denote the bootstrap weights, which are nonnegative random variables independent of the data obeying the conditions stated in van der Vaart and Wellner (1996). For example, $(\omega_1, \dots, \omega_n)$ is a multinomial vector with dimension n and probabilities $(1/n, \dots, 1/n)$ in the empirical bootstrap. In what follows B is the number of bootstrap draws, such that $B \rightarrow \infty$. In our experience, setting $B \geq 500$ suffices for good accuracy.

Algorithm 2.1 (Bootstrap law of $t(\mathcal{U})$ and its quantiles). **1)** Draw a realization of the bootstrap weights $(\omega_1, \dots, \omega_n)$. **2)** For each $u \in \mathcal{U}$, compute $\widehat{\Delta}_\mu^*(u) = \widetilde{\Delta}_\mu^*(u)$, a bootstrap draw of $\widehat{\Delta}_\mu^*(u) = \widehat{\Delta}_{\widetilde{\mu}}^*(u)$, where $\widetilde{\Delta}$ and $\widetilde{\mu}$ are the bootstrap versions of $\widehat{\Delta}$ and $\widehat{\mu}$ that use $(\omega_1, \dots, \omega_n)$ as sampling weights in the computation of the estimators. Construct a bootstrap draw of $Z_\infty(u)$ as $\widetilde{Z}_\infty(u) = \sqrt{n}(\widetilde{\Delta}_\mu^*(u) - \widehat{\Delta}_\mu^*(u))$. **3)** Repeat steps (1)-(2) B times. **4)** For each $u \in \mathcal{U}$, compute a bootstrap estimator of $\Sigma(u)^{1/2}$ such as the bootstrap interquartile range rescaled with the normal distribution, $\widehat{\Sigma}(u)^{1/2} = (q_{0.75}(u) - q_{0.25}(u))/(z_{0.75} - z_{0.25})$, where $q_p(u)$ is the p th sample quantile of $\widetilde{Z}_\infty(u)$ in the B draws and z_p is the p th quantile of $N(0, 1)$. **5)** Use the empirical distribution of $\tilde{t}(\mathcal{U}) = \sup_{u \in \mathcal{U}} |\widetilde{Z}_\infty(u)| \widehat{\Sigma}(u)^{-1/2}$ across the B draws to approximate the distribution of $t(\mathcal{U}) = \sup_{u \in \mathcal{U}} |Z_\infty(u)| \Sigma(u)^{-1/2}$. In particular, construct $\widehat{t}_{1-\alpha}(\mathcal{U})$, an estimator of $t_{1-\alpha}(\mathcal{U})$, as the $(1 - \alpha)$ -quantile of the B draws of $\tilde{t}(\mathcal{U})$.

Remark 2.2 (Monotonization of the bands). While the SPE-function $u \mapsto \Delta_\mu^*(u)$ is increasing by definition, the end functions of the confidence band $u \mapsto \widehat{\Delta}_\mu^*(u) \pm \widehat{t}_{1-\alpha}(\mathcal{U}) \widehat{\Sigma}(u)^{1/2} / \sqrt{n}$ might not be increasing. Chernozhukov, Fernández-Val, and Galichon (2009) showed that monotonizing the end functions via rearrangement reduces the width of the band in uniform norm, while increases coverage in finite-samples. We use this refinement in the empirical examples.⁴ \square

Remark 2.3 (Finite-Sample Bias Corrections). The empirical u -SPE might be biased in small samples, specially at the tails. Bootstrap is also useful to improve the estimator and confidence bands. Thus, a corrected estimator can be formed as $2\widehat{\Delta}_\mu^* - \overline{\Delta}_\mu^*$, and a corrected $(1 - \alpha)$ -confidence band as $\left[2\widehat{\Delta}_\mu^* - \overline{\Delta}_\mu^* \pm \widehat{t}_{1-\alpha}(\mathcal{U}) \widehat{\Sigma}(u)^{1/2} / \sqrt{n}\right]$, where $\overline{\Delta}_\mu^*$ is the mean of the bootstrap draw of the estimator. In Appendix H of the SM, we show that this correction reduces the bias of the estimator and increases the coverage of the confidence bands in a simulation calibrated to the gender wage gap application. \square

2.5. Inference on CA. Let $\Lambda_{\Delta, \mu}^u(t) := [\Lambda_{\Delta, \mu}^{-u}(t), \Lambda_{\Delta, \mu}^{+u}(t)]$ and $\widehat{\Lambda}_{\Delta, \mu}^u(t) := [\widehat{\Lambda}_{\Delta, \mu}^{-u}(t), \widehat{\Lambda}_{\Delta, \mu}^{+u}(t)]$. The main inferential result for CA can be previewed as follows: the empirical CA-process converges in distribution to a centered bivariate Gaussian process, namely

$$\sqrt{n}(\widehat{\Lambda}_{\Delta, \mu}^u(t) - \Lambda_{\Delta, \mu}^u(t)) \rightsquigarrow Z_\infty^u(t) \text{ in } \ell^\infty(\mathbb{R}^{d_z})^2, \quad (2.3)$$

⁴In practice, the rearrangement simply consists in sorting the two vectors containing the discretized version of the end-functions in increasing order; see Chernozhukov, Fernández-Val, and Galichon (2009) for more details.

as a stochastic process indexed by $t \in \mathbb{R}^{d_z}$. Moreover, exchangeable bootstrap estimates consistently the law of $Z_\infty^u(t)$.

The next corollary to Theorem 4.2 in Section 4 provides uniform bands that cover L linear combinations of the 2-dimensional vector $\Lambda_{\Delta, \mu}^u(t)$ with coefficients c_1, \dots, c_L simultaneously over $t \in \mathcal{T}$ with prespecified probability in large samples. It covers pointwise confidence intervals for the mean of the k^{th} component of Z for least affected as a special case with $L = 1$ linear combination, $c_1 = (1, 0)'$, and $\mathcal{T} = \{e_k\}$, where e_k is a unit vector with a one in the k^{th} position. Joint confidence intervals for s differences of means of the $k_1^{\text{th}}, \dots, k_s^{\text{th}}$ components of Z between most and least affected are a special case with $L = 1$ linear combination, $c_1 = (-1, 1)'$, and $\mathcal{T} = \{e_{k_1}, \dots, e_{k_s}\}$. Joint uniform bands for the distribution of the k^{th} component of Z for most and least affected are also a special case with $L = 2$ linear combinations, $c_1 = (1, 0)$, $c_2 = (0, 1)$, and $\mathcal{T} = \{t \in \mathbb{R}^{d_z} : t_j = \bar{T}, j \neq k\}$, where \bar{T} is an arbitrarily large number. By appropriate choice of the linear combinations and the index set T , we can therefore conduct multiple tests while preserving the significance level from simultaneous inference problems (Romano, Shaikh, and Wolf, 2010a; Romano, Shaikh, and Wolf, 2010b; List, Shaikh, and Xu, 2016). We show examples in the empirical application of Section 3.

Corollary 2.2 (Inference on CA-function using Limit Theory and Bootstrap). *Under the assumptions of Theorem 4.2, for any $0 < \alpha < 1$,*

$$\mathbb{P} \left\{ c'_\ell \Lambda_{\Delta, \mu}^u(t) \in c'_\ell \hat{\Lambda}_{\Delta, \mu}^u(t) \pm \hat{t}_{1-\alpha}^u(\mathcal{T}, L) [c'_\ell \hat{\Sigma}^u(t) c_\ell]^{1/2} / \sqrt{n} : t \in \mathcal{T}, \ell = 1, \dots, L \right\} \rightarrow 1 - \alpha,$$

where $\hat{t}_{1-\alpha}^u(\mathcal{T}, L)$ is any consistent estimator of $t_{1-\alpha}^u(\mathcal{T}, L)$, the $(1 - \alpha)$ -quantile of

$$t^u(\mathcal{T}, L) := \sup_{t \in \mathcal{T}, \ell = 1, \dots, L} |c'_\ell Z_\infty^u(t)| [c'_\ell \Sigma^u(t) c_\ell]^{-1/2},$$

and $t \mapsto \hat{\Sigma}^u(t)$ is a uniformly consistent estimator of $t \mapsto \Sigma^u(t)$, the variance function of $t \mapsto Z_\infty^u(t)$. A p -value of the null hypothesis $c'_\ell \Lambda_{\Delta, \mu}^u(t) = r_\ell(t)$ for all $t \in \mathcal{T}$ and $\ell = 1, \dots, L$ of the realization of the statistic $\sup_{t \in \mathcal{T}, \ell = 1, \dots, L} |c'_\ell \hat{\Lambda}_{\Delta, \mu}^u(t) - r_\ell(t)| [c'_\ell \hat{\Sigma}^u(t) c_\ell]^{-1/2} = s$ is

$$S_{t^u(\mathcal{T}, L)}(s) = \mathbb{P} (t_{1-\alpha}^u(\mathcal{T}, L) > s).$$

We provide consistent estimators of $t_{1-\alpha}^u(\mathcal{T}, L)$, $u \mapsto \Sigma^u(t)$ and $S_{t^u(\mathcal{T}, L)}(t)$ in Algorithm 2.2.

Algorithm 2.2 (Bootstrap law of $t(\mathcal{T}, L)$, quantiles and p-values). **1)** Draw a realization of the bootstrap weights $(\omega_1, \dots, \omega_n)$. **2)** For each $t \in \mathcal{T}$, compute $\tilde{\Lambda}_{\Delta, \mu}^{-u}(t) = \Lambda_{\tilde{\Delta}, \tilde{\mu}}^{-u}(t)$ and $\tilde{\Lambda}_{\Delta, \mu}^{+u}(t) = \Lambda_{\tilde{\Delta}, \tilde{\mu}}^{+u}(t)$, a bootstrap draw of $\hat{\Lambda}_{\Delta, \mu}^{-u}(t) = \Lambda_{\hat{\Delta}, \hat{\mu}}^{-u}(t)$ and $\hat{\Lambda}_{\Delta, \mu}^{+u}(t) = \Lambda_{\hat{\Delta}, \hat{\mu}}^{+u}(t)$, where $\tilde{\Delta}$ and $\tilde{\mu}$ are the bootstrap versions of $\hat{\Delta}$ and $\hat{\mu}$ that use $(\omega_1, \dots, \omega_n)$ as sampling weights in the computation of the estimators. Construct a bootstrap draw of $Z_\infty^u(t)$ as $\tilde{Z}_\infty^u(t) = \sqrt{n}(\tilde{\Lambda}_{\Delta, \mu}^u(t) - \hat{\Lambda}_{\Delta, \mu}^u(t))$, where $\tilde{\Lambda}_{\Delta, \mu}^u(t) = [\tilde{\Lambda}_{\Delta, \mu}^{-u}(t), \tilde{\Lambda}_{\Delta, \mu}^{+u}(t)]$. **3)** Repeat steps (1)-(2) B times. **4)** For each $t \in \mathcal{T}$ and $\ell = 1, \dots, L$, compute a bootstrap estimator of $[c'_\ell \Sigma^u(t) c_\ell]^{1/2}$ such as the bootstrap interquartile range rescaled with the normal distribution $[c'_\ell \tilde{\Sigma}^u(t) c_\ell]^{1/2} = (q_{0.75}^u(t, \ell) - q_{0.25}^u(t, \ell)) / (z_{0.75} - z_{0.25})$, where $q_p^u(t, \ell)$ is the p th sample quantile of $c'_\ell \tilde{Z}_\infty^u(t)$ in the B draws and z_p is the p th quantile of $N(0, 1)$. **5)** Use

the empirical distribution of $\tilde{t}(\mathcal{T}, L) = \sup_{t \in \mathcal{T}, \ell=1, \dots, L} |c'_\ell \tilde{Z}_\infty^u(t)| [c'_\ell \widehat{\Sigma}^u(t) c_\ell]^{-1/2}$ across the B draws to approximate the distribution of $t(\mathcal{T}, L) = \sup_{t \in \mathcal{T}, \ell=1, \dots, L} |c'_\ell Z_\infty^u(t)| [c'_\ell \Sigma^u(t) c_\ell]^{-1/2}$. In particular, construct $\widehat{t}_{1-\alpha}(\mathcal{T}, L)$, an estimator of $t_{1-\alpha}(\mathcal{T}, L)$, as the $(1-\alpha)$ -quantile of the B draws of $\tilde{t}(\mathcal{T}, L)$, and an estimation of the p -value $S_{\tilde{t}(\mathcal{T}, L)}(s)$ as the proportion of the B draws of $\tilde{t}(\mathcal{T}, L)$ that are greater than s .

2.6. Inference on Most and Least Affected Subpopulations. In addition to moments and distributions, we can conduct inference on the subpopulations of most and least affected.⁵ Let

$$\mathcal{M}^{-u} := \{(x, y) \in \mathcal{Z} : \Delta(x) \leq \Delta_\mu^*(u)\}, \quad \mathcal{M}^{+u} := \{(x, y) \in \mathcal{Z} : \Delta(x) \geq \Delta_\mu^*(1-u)\},$$

be the sets representing the u -least and u -most affected subpopulation, respectively. Here we assume that \mathcal{Z} is compact or that the support of (X, Y) has been intersected with a compact set to form \mathcal{Z} . We can construct an outer $(1-\alpha)$ -confidence set for \mathcal{M}^{-u} as⁶

$$\mathcal{CM}^{-u}(1-\alpha) = \{(x, y) \in \mathcal{Z} : \widehat{\Sigma}^{-1/2}(x, u) \sqrt{n} [\widehat{\Delta}(x) - \widehat{\Delta}_\mu^*(u)] \leq \widehat{c}(1-\alpha)\},$$

where $\widehat{c}(1-\alpha)$ is a consistent estimator of $c(1-\alpha)$, the $(1-\alpha)$ -quantile of the random variable

$$V_\infty = \sup_{\{x \in \mathcal{X} : \Delta(x) = \Delta_\mu^*(u)\}} \Sigma^{-1/2}(x, u) [G_\infty(x) - Z_\infty(u)],$$

and $x \mapsto \widehat{\Sigma}(x, u)$ is a uniformly consistent estimator of $x \mapsto \Sigma(x, u)$, the variance function of the process $G_\infty(x) - Z_\infty(u)$ defined in Section 4. The estimator $\widehat{c}(1-\alpha)$ can be obtained as the $(1-\alpha)$ -quantile of the bootstrap version of V_∞ ,

$$\widetilde{V}_\infty^* = \sup_{\{x \in \mathcal{X} : \widehat{\Delta}(x) = \widehat{\Delta}_\mu^*(u)\}} \widehat{\Sigma}^{-1/2}(x, u) \sqrt{n} \left([\widetilde{\Delta}(x) - \widetilde{\Delta}_\mu^*(u)] - [\widehat{\Delta}(x) - \widehat{\Delta}_\mu^*(u)] \right),$$

where $\widetilde{\Delta}(x)$ and $\widetilde{\Delta}_\mu^*(u)$ are defined as in Algorithm 2.1. A similar $(1-\alpha)$ -confidence set, $\mathcal{CM}^{+u}(1-\alpha)$, can be constructed for \mathcal{M}^{+u} . These sets can be visualized by plotting all 2 or 3 dimensional projections of their elements. We provide an example of such plots in Section 3. An immediate consequence of the set inference results in Chernozhukov, Kocatulum, and Menzel (2015) and the results of this paper is the following corollary:

Corollary 2.3 (Inference on Most and Least Affected Subpopulations). *The sets $\mathcal{CM}^{-u}(1-\alpha)$ and $\mathcal{CM}^{+u}(1-\alpha)$ cover \mathcal{M}^{-u} and \mathcal{M}^{+u} with probability approaching $1-\alpha$, and $\mathcal{CM}^{-u}(1-\alpha)$ and $\mathcal{CM}^{+u}(1-\alpha)$ are consistent in the sense that they approach to \mathcal{M}^{-u} and \mathcal{M}^{+u} at a \sqrt{n} -rate with respect to the Hausdorff distance.*

⁵Here we follow the set inference approach described in Chernozhukov, Kocatulum, and Menzel (2015), which builds on Chernozhukov, Hong, and Tamer (2007). In addition our results justify the use of subsampling-based methods as in Chernozhukov, Hong, and Tamer (2007) and Romano and Shaikh (2010).

⁶Note that we can also similarly construct an inner confidence region, which is the complement of the outer confidence region of $\mathcal{X} \setminus \mathcal{M}^{-u}$, see Chernozhukov, Kocatulum, and Menzel (2015) for relevant discussion.

3. EMPIRICAL ANALYSIS OF THE GENDER WAGE GAP

We report the main results of the application to the gender wage gap using data from the U.S. March Supplement of the Current Population Survey (CPS) in 2015. In Appendix H of the SM, we complement the analysis with supporting results from a simulation calibrated to this application. There, we find that our estimation and inference methods perform well in finite samples that closely mimic the characteristics of the CPS data. This exercise serves to indirectly verify the plausibility of the main regularity conditions mentioned in Section 2 and formally stated in Section 4.

Our sample consists of white, non-hispanic individuals who are aged 25 to 64 years and work more than 35 hours per week during at least 50 weeks of the year. We exclude self-employed workers; individuals living in group quarters; individuals in the military, agricultural or private household sectors; individuals with inconsistent reports on earnings and employment status; individuals with allocated or missing information in any of the variables used in the analysis; and individuals with hourly wage rate below \$3. The resulting sample contains 32,523 workers including 18,137 men and 14,382 of women.

We estimate interactive linear models with additive and non-additive errors, using mean and quantile regressions, respectively. The outcome variable Y is the logarithm of the hourly wage rate constructed as the ratio of the annual earnings to the total number of hours worked, which is constructed in turn as the product of number of weeks worked and the usual number of hours worked per week. The key covariate T is an indicator for female worker, and the control variables W include 5 marital status indicators (widowed, divorced, separated, never married, and married); 5 educational attainment indicators (less than high school graduate, high school graduate, some college, college graduate, and advanced degree); 4 region indicators (midwest, south, west, and northeast); a quartic in potential experience constructed as the maximum of age minus years of schooling minus 7 and zero, i.e., $experience = \max(age - education - 7, 0)$; 5 occupation indicators (management, professional and related; service; sales and office; natural resources, construction and maintenance; and production, transportation and material moving); 12 industry indicators (mining, quarrying, and oil and gas extraction; construction; manufacturing; wholesale and retail trade; transportation and utilities; information; financial services; professional and business services; education and health services; leisure and hospitality; other services; and public administration); and all the two-way interactions between the education, experience, occupation and industry variables except for the occupation-industry interactions.⁷ All calculations use the CPS sampling weights to account for nonrandom sampling in the March CPS.

⁷The sample selection criteria and the variable construction follow Mulligan and Rubinstein (2008). The occupation and industry categories follow the 2010 Census Occupational Classification and 2012 Census Industry Classification, respectively.

TABLE 1. Descriptive Statistics of Workers

	All	Women	Men		All	Women	Men
Log wage	3.15	3.02	3.25	O.manager	0.48	0.55	0.43
Female	0.44	1.00	0.00	O.service	0.10	0.10	0.09
MS.married	0.65	0.61	0.68	O.sales	0.23	0.31	0.16
MS.widowed	0.01	0.02	0.01	O.construction	0.09	0.01	0.15
MS.separated	0.02	0.02	0.02	O.production	0.11	0.04	0.17
MS.divorced	0.13	0.16	0.10	I.minery	0.03	0.01	0.04
MS.Nevermarried	0.19	0.18	0.20	I.construction	0.06	0.01	0.09
E.lhs	0.02	0.02	0.03	I.manufacture	0.14	0.08	0.18
E.hsg	0.25	0.21	0.28	I.retail	0.13	0.11	0.14
E.sc	0.28	0.29	0.27	I.transport	0.04	0.02	0.06
E.cg	0.28	0.30	0.27	I.information	0.02	0.02	0.03
E.ad	0.16	0.18	0.15	I.finance	0.08	0.10	0.07
R.northeast	0.19	0.19	0.19	I.professional	0.11	0.10	0.13
R.midwest	0.27	0.28	0.27	I.education	0.24	0.40	0.11
R.south	0.35	0.35	0.35	I.leisure	0.05	0.05	0.04
R.west	0.18	0.18	0.19	I.services	0.03	0.03	0.04
Experience	21.68	21.72	21.65	I.public	0.07	0.06	0.07

Source: March Supplement CPS 2015.

Table 1 reports sample means of the variables used in the analysis. Working women are more highly educated than working men, have about the same potential experience, and are less likely to be married and more likely to be divorced. They work relatively more often in managerial and sales occupations and in the industries providing education and health services. Working men are relatively more likely to work in construction and production occupations within non-service industries. The unconditional gender wage gap is 23%.

Figure 1 of Section 1 plots estimates and 90% confidence bands for the APE and SPE-function on the treated (women) of the conditional gender wage gap using additive and non-additive error models. The PEs are obtained as described in Examples 2 and 3 with $P(T, W) = (TW, (1 - T)W)$. In this case $\dim P(T, W) = 332$, which makes it very difficult to identify any pattern about the gender wage gap just by looking at the regression coefficients. The distribution $F_{T,W}$ is estimated by the empirical distribution of (T, W) for women, and F_ϵ is approximated by a uniform distribution over the grid $\{.02, .03, \dots, .98\}$. The confidence bands are constructed using Algorithm 2.1 with standard exponential weights (weighted bootstrap) and $B = 500$, and are uniform for the SPE-function over the grid $\mathcal{U} = \{.01, .02, \dots, .98\}$. We monotoneize the bands using the rearrangement method described in Remark 2.2, and implement the finite sample corrections described in Remark 2.3. After controlling for worker characteristics, the gender wage gap for women remains on average around 20%. More importantly, we uncover a striking amount of

heterogeneity, with the PE ranging between -6.5 and 40% in the additive error model and between -14 and 54% in the non-additive error model.⁸

TABLE 2. Classification Table – Average Characteristics of the 10% Least and Most Affected Women by Gender Wage Gap

	10% Least		10% Most			10% Least		10% Most	
	Est.	S.E.	Est.	S.E.		Est.	S.E.	Est.	S.E.
Log wage	3.08	0.03	2.97	0.03	O.manager	0.67	0.04	0.38	0.04
M.married	0.28	0.03	0.87	0.02	O.service	0.08	0.02	0.10	0.02
M.widowed	0.02	0.01	0.01	0.01	O.sales	0.19	0.03	0.42	0.04
M.separated	0.02	0.01	0.01	0.00	O.construction	0.02	0.01	0.01	0.00
M.divorced	0.15	0.02	0.07	0.02	O.production	0.03	0.01	0.09	0.02
M.nevermarried	0.52	0.03	0.04	0.01	I.minery	0.01	0.01	0.02	0.01
E.lhs	0.01	0.01	0.06	0.01	I.construction	0.02	0.01	0.02	0.01
E.hsg	0.08	0.02	0.30	0.04	I.manufacture	0.02	0.01	0.11	0.02
E.sc	0.15	0.03	0.23	0.04	I.retail	0.06	0.02	0.19	0.03
E.cg	0.37	0.04	0.17	0.03	I.transport	0.01	0.01	0.04	0.01
E.ad	0.39	0.04	0.24	0.03	I.information	0.04	0.01	0.05	0.01
R.ne	0.24	0.02	0.18	0.02	I.finance	0.03	0.02	0.09	0.03
R.mw	0.26	0.02	0.28	0.02	I.professional	0.06	0.02	0.04	0.02
R.so	0.31	0.02	0.39	0.03	I.education	0.46	0.04	0.33	0.04
R.we	0.19	0.02	0.15	0.02	I.leisure	0.10	0.03	0.01	0.01
Experience	13.05	1.03	26.32	0.75	I.services	0.09	0.02	0.01	0.01
					I.public	0.09	0.02	0.08	0.02

PE estimated from a linear conditional quantile model with interactions.

Standard Errors obtained by weighted bootstrap with 500 repetitions.

Table 8 shows the results of a classification analysis, exhibiting characteristics of women that are most and least affected by the gender wage gap together with standard errors obtained by weighted bootstrap. We focus here on the non-additive model, but the results from the additive model are similar. Since the PE are predominantly negative, we define the most affected as $\Delta(X) < \Delta_{\mu}^*(u)$ and the least affected as $\Delta(X) > \Delta_{\mu}^*(1 - u)$ to facilitate the interpretation. According to this model the 10% of the women *most affected* by the gender wage gap on average earn lower wages, are much more likely to be married, much less likely to be never married, have lower education, live in the South, possess much more potential experience, are more likely to have sales and non managerial occupations, and work more often in manufacture and retail and less often in education industries than the 10% least affected women.

Table 3 tests if the differences found in table 8 are statistically significant. It reports p-values for the test of equality of means for most and least affected women. The first p-value accounts for simultaneous inference on all variables within a given category. For example, it accounts that

⁸In the 2016 version of the paper we found similar patterns of heterogeneity using CPS 2012 data with a specification that did not include occupation and industry indicators.

TABLE 3. Classification Table – Difference in the Average Characteristics of the 10% Most and Least Affected Women by Gender Wage Gap

	Est.	S.E.	P-val. ¹	JP-val. ²		Est.	S.E.	P-val. ¹	JP-val. ²
Log wage	-0.10	0.04	0.03	0.70	O.manager	-0.29	0.06	0.00	0.00
M.married	0.59	0.04	0.00	0.00	O.service	0.02	0.03	0.99	1.00
M.widowed	-0.02	0.02	0.93	1.00	O.sales	0.22	0.06	0.00	0.03
M.separated	-0.01	0.01	0.89	1.00	O.construction	-0.01	0.01	0.67	1.00
M.divorced	-0.08	0.04	0.46	0.86	O.production	0.06	0.02	0.08	0.42
M.nevermarried	-0.48	0.04	0.00	0.00	I.minery	0.01	0.01	0.97	1.00
E.lhs	0.05	0.01	0.01	0.17	I.construction	-0.00	0.01	1.00	1.00
E.hsg	0.22	0.05	0.00	0.00	I.manufacture	0.08	0.02	0.02	0.15
E.sc	0.08	0.06	0.70	1.00	I.retail	0.12	0.04	0.06	0.32
E.cg	-0.19	0.06	0.01	0.16	I.transport	0.04	0.01	0.08	0.39
E.ad	-0.15	0.06	0.07	0.46	I.information	0.01	0.02	1.00	1.00
R.ne	-0.06	0.04	0.35	0.99	I.finance	0.06	0.04	0.78	0.99
R.mw	0.02	0.04	0.95	1.00	I.professional	-0.01	0.03	1.00	1.00
R.so	0.08	0.04	0.23	0.97	I.education	-0.13	0.06	0.29	0.74
R.we	-0.04	0.03	0.69	1.00	I.leisure	-0.09	0.03	0.04	0.22
Experience	13.27	1.54	0.00	0.00	I.services	-0.07	0.02	0.02	0.15
					I.public	-0.01	0.03	1.00	1.00

PE estimated from a linear conditional quantile model with interactions.

Standard Errors and p-values obtained by weighted bootstrap with 500 repetitions.

¹ These p-values are adjusted for multiplicity to account for joint testing of zero coefficients on for all variables within a category: M E, R, O, or I.

² These p-values are adjusted for multiplicity to account for joint testing of zero coefficients on all the variables in the table.

we are conducting five tests corresponding to the five categories of marital status. For the non categorical variables log wage and experience the p-values are for one test. The second p-value accounts for simultaneous inference of all the differences displayed in the table.⁹ These p-values are obtained by Algorithm 2.2 with the appropriate choice of vectors of linear combinations and set \mathcal{T} , and 500 weighted bootstrap repetitions. The p-values show that most of the differences from table 8 are statistically significant at conventional significant levels after controlling for simultaneous inference. In particular, the most affected women are significantly more likely to be married, high-school graduates, more experienced, and in sales occupations, and less likely to be never married and in managerial occupations under the most strict simultaneous inference correction. Blau and Kahn (2017) have recently documented the importance of differences in occupation and industry to explain the gender wage gap using data from the Panel Study of Income Dynamics (PSID) 1980-2010 and a different methodology based on wage decompositions. Consistent with our findings,

⁹We employ the so called "single-step" methods for controlling the family-wise error rate. To generate a (somewhat) higher power, we recommend to employ the p-values generated via "step-down" methods, such as those reported in Romano and Wolf (2016) and List, Shaikh, and Xu (2016).

they argue that this importance might be due to compensating differentials. Unlike Blau and Kahn (2017) and previous studies in the literature, our analysis uncovers significant heterogeneity in the extent of the gender wage gap and relates this heterogeneity to human capital, occupation, industry and other characteristics.




FIGURE 2. APE and SPE of the gender wage gap for women by marital status. Estimates and 90% bootstrap uniform confidence bands for the conditional quantile function.

We further explore these findings by analyzing the APE and SPE on the treated *conditional* on marital status and unobserved rank in the non-additive error model. Figures 2 and 3 show estimates and 90% confidence bands of the APE and SPE-function of the gender wage gap for 2 subpopulations defined by marital status (married and never married) and 3 subpopulations defined by unobserved rank (first decile, median and ninth decile, where the unobserved rank is .1, .5 and .9, respectively). The confidence bands are constructed as in fig. 1. We find significant heterogeneity in the gender gap within each subpopulation, and also between subpopulations defined by marital status and unobserved rank. The SPE-function is more negative for married women and at the tails of the conditional distribution. Married women at the top decile suffer from the highest gender wage gaps. This pattern is consistent with “glass-ceiling” effects behind the gender wage gap (Albrecht, Bjorklund, and Vroman, 2003).

Figure 4 plots simultaneous 90% confidence bands for the distribution of experience and log wage for the most and least affected women. They are obtained by Algorithm 2.2 with 500 weighted bootstrap replications. The estimated distribution of experience for the most affected first-order stochastically dominates the same estimated distribution for the least affected women.




FIGURE 3. APE and SPE of the gender wage gap for women by unobserved ranking in the conditional distribution. Estimates and 90% bootstrap uniform confidence bands for the conditional quantile function.




FIGURE 4. Estimates and 90% weighted bootstrap joint uniform confidence bands for the distributions of experience and log wages of the 10% most and least affected women by gender wage gap.

Moreover, the uniform bands confirm that this dominance is statistically significant at the 90% confidence level for the underlying distributions. The estimated (marginal) distribution of log wage for the least affected first-order dominates the same estimated distribution for most affected, but we cannot reject that the underlying distributions are equal at the 10% significance level. The results of the classification analysis are consistent with preferences that make never married highly educated young women working on managerial occupations be more career-oriented.¹⁰




FIGURE 5. Estimates and 90% weighted bootstrap confidence bands for projections of the confidence sets for characteristics of 10% most and least affected women by gender wage gap.

Finally, Figure 5 plots two dimensional projections of experience-log wage and experience-marital status of the confidence sets for the 10% most and least affected subpopulations. We show the results from the additive error model for the conditional expectation. Here we use a simplified specification that excludes the two-way interactions from W to get more precise estimates of all the PEs. We obtain 90% confidence sets for the most and least affected subpopulations by weighted bootstrap with standard exponential weights and 500 repetitions. The sets $\mathcal{CM}^{-0.1}(0.90)$ and $\mathcal{CM}^{+0.1}(0.90)$ include 23% and 19% of the women in the sample, respectively.¹¹ The projections show that there are relatively more least affected women with low experience at all wage levels, more high affected women with high wages with between 15 and 25 years of experience, and more least affected women which are not married at all experience levels.

¹⁰We find similar results using the additive error model. We do not report these results for the sake of brevity.

¹¹Recall that in this application the set $\mathcal{CM}^{-0.1}(0.90)$ corresponds to most affected women and $\mathcal{CM}^{+0.1}(0.90)$ to least affected women. We drop one woman that is included in both sets.

4. DETAILED LARGE SAMPLE THEORY

4.1. Detailed Large Sample Theory for SPE. For an open set \mathcal{K} , let the class \mathcal{C}^1 on \mathcal{K} denote the set of continuously differentiable real valued functions on \mathcal{K} . We make the following technical assumptions about the PE function $\Delta : \mathbb{R}^{d_x} \mapsto \mathbb{R}$ and the distribution of the covariates:

S.1. The part of the domain of the PE function $x \mapsto \Delta(x)$ of interest, \mathcal{X} , is open and its closure $\overline{\mathcal{X}}$ is compact. The distribution μ is absolutely continuous with respect to the Lebesgue measure with density μ' . There exists an open set $B(\mathcal{X})$ containing $\overline{\mathcal{X}}$ such that $x \mapsto \Delta(x)$ is \mathcal{C}^1 on $B(\mathcal{X})$, and $x \mapsto \mu'(x)$ is continuous on $B(\mathcal{X})$ and is zero outside the domain of interest, i.e. $\mu'(x) = 0$ for any $x \in B(\mathcal{X}) \setminus \mathcal{X}$.

S.2. Let $\mathcal{M}_\Delta(\delta) := \{x \in \mathcal{X} : \Delta(x) = \delta\}$. For any regular value δ of Δ on $\overline{\mathcal{X}}$, we assume that the closure of $\mathcal{M}_\Delta(\delta)$ has a finite number of connected branches.

The following property of the set $\mathcal{M}_\Delta(\delta)$ is a useful implication of Assumptions S.1 and S.2 that we will exploit in the analysis.

Remark 4.1 (Properties of $\mathcal{M}_\Delta(\delta)$). By Theorem 5-1 in Spivak (1965, p. 111), S.1 and S.2 imply that $\mathcal{M}_\Delta(\delta)$ is a $(d_x - 1)$ -manifold without boundary in \mathbb{R}^{d_x} of class \mathcal{C}^1 for any δ that is a regular value of $x \mapsto \Delta(x)$ on $\overline{\mathcal{X}}$.

Assumption S.1 imposes mild smoothness conditions on the PE function $x \mapsto \Delta(x)$. It also requires that all the components of the covariate X are continuous random variables. We defer the treatment of the case where X has both continuous and discrete components to the SM. As a matter of generalization, our theoretical analysis allows us to replace that $x \mapsto \mu'(x)$ vanishes on $\partial\mathcal{X}$, by the weaker condition that the intersection of $\mathcal{M}_\Delta(\delta)$ and the boundary of \mathcal{X} have zero volume with respect to μ , namely

$$\int_{\mathcal{M}_\Delta(\delta)} 1_{\{x \in \partial\mathcal{X}\}} \frac{\mu'(x)}{\|\partial\Delta(x)\|} d\text{Vol} = 0, \quad (4.4)$$

where $\partial\mathcal{X}$ denotes the boundary of $\overline{\mathcal{X}}$, $\partial\Delta(x)$ is the gradient of $x \mapsto \Delta(x)$, and $\int_{\mathcal{M}} f(x) d\text{Vol}$ denotes the integral of the function f on the manifold \mathcal{M} with respect to volume; see Appendix D in the SM for a brief review on Differential Geometry. This relaxation is relevant to cover the case where X includes an uniformly distributed component such as the unobserved rank in Example 3.¹²

Assumption S.2 imposes shape restrictions on $x \mapsto \Delta(x)$ that rule out cases such as infinite cyclical oscillations or flat areas. A simple sufficient condition for S.2 is that the map $x \mapsto \Delta(x)$ does not have critical points on $\overline{\mathcal{X}}$. This means that $x \mapsto \Delta(x)$ is not locally flat anywhere on $\overline{\mathcal{X}}$, which we define to mean that the norm of the gradient, $\|\partial\Delta(x)\|$, does not vanish on $x \in \overline{\mathcal{X}}$.

¹² In the numerical examples of Section H in the SM, the first two designs only satisfy this relaxed condition.

In this case, any δ in the image of $\bar{\mathcal{X}}$ under Δ is regular. This condition is probably the most relevant for practice and can be verified in applications, at least informally.

Remark 4.2 (Verification of Regularity Conditions in Practice). The main regularity condition is that PE function $x \mapsto \Delta(x)$ be smooth and not locally flat, namely $\|\partial\Delta(x)\|$ does not vanish. Our inferential results are developed under this assumption, and they do not apply otherwise. To verify if these results apply in practice, we strongly recommend to conduct a Monte Carlo experiment using a data generating process that mimics the application at hand.¹³ Indeed, failure of the inference method in the simulation experiment implies failure of the regularity conditions. We provide an application of this supporting analysis to the gender wage gap example in Appendix H. Looking forward, it would be useful to develop further an inference method with good robustness properties with respect to the regularity conditions, i.e. that remains uniformly valid when the PE function is (close to being) locally flat. We delegate this line of research to future work.¹⁴

We make the following assumptions about the estimator of the PE. Let $\ell^\infty(\mathcal{T})$ denote the set of bounded and measurable functions $g : \mathcal{T} \rightarrow \mathbb{R}$ and \mathcal{F} a fixed subset of continuous functions on $B(\mathcal{X})$. Let $\ell^\infty(B(\mathcal{X}))$ be the set of bounded and measurable functions on $B(\mathcal{X})$ and \rightsquigarrow denote weak convergence (convergence in distribution).

S.3. $\widehat{\Delta}$, the estimator of Δ , belongs to \mathcal{F} with probability approaching 1 and obeys a functional central limit theorem, namely,

$$a_n(\widehat{\Delta} - \Delta) \rightsquigarrow G_\infty \text{ in } \ell^\infty(B(\mathcal{X})),$$

where a_n is a sequence such that $a_n \rightarrow \infty$ as $n \rightarrow \infty$, and $x \mapsto G_\infty(x)$ is a tight process that has almost surely uniformly continuous sample paths on $B(\mathcal{X})$.

In the parametric and semiparametric models of Examples 1–3, S.3 holds under weak conditions that guarantee asymptotic normality of the ML, OLS and QR estimators. For the QR estimator in Example 3 where the unobserved rank is one of the covariates, these conditions include that the density of Y conditional on X be bounded away from zero (Koenker, 2005), which is facilitated by excluding tail quantile indexes.

Let $\widehat{\mu}$ be the estimator of the distribution μ . It is convenient to identify μ and $\widehat{\mu}$ with the operators:

$$g \mapsto \mu(g) = \int g(x)d\mu(x), \quad g \mapsto \widehat{\mu}(g) = \int g(x)d\widehat{\mu}(x),$$

mapping from the set $\mathcal{G} := \{x \mapsto 1(f(x) \leq \delta) : f \in \mathcal{F}, \delta \in \mathcal{V}\}$ to \mathbb{R} , where \mathcal{F} is the fixed subset of continuous functions on $B(\mathcal{X})$ containing Δ , and \mathcal{V} is any compact set of \mathbb{R} . We require \mathcal{G} to be

¹³In fact, we recommend doing this for every econometric method.

¹⁴For instance, it is of interest to determine whether the use of subsampling instead of bootstrap can deliver a more robust inference method when the PE function is close to being flat; see, e.g. Romano and Shaikh (2012).

totally bounded under the $L^2(\mu)$ norm. Define \mathbb{H} as the set of all bounded linear operators H on \mathcal{G} of the form

$$g \mapsto H(g),$$

which are uniformly continuous on $g \in \mathcal{G}$ under the $L^2(\mu)$ norm. We define the boundedness of these operators with respect to the norm:

$$\|H\|_{\mathcal{G}} = \sup_{g \in \mathcal{G}} |H(g)|,$$

and define the corresponding distance between two operators H and \tilde{H} in \mathbb{H} as $\|H - \tilde{H}\|_{\mathcal{G}} = \sup_{g \in \mathcal{G}} |H(g) - \tilde{H}(g)|$. Clearly, $\mu \in \mathbb{H}$.

We make the following assumption about $\hat{\mu}$.

S.4. The function $x \mapsto \hat{\mu}(x)$ is a distribution over $B(\mathcal{X})$ obeying in \mathbb{H} ,

$$b_n(\hat{\mu} - \mu) \rightsquigarrow H_{\infty}, \quad (4.5)$$

where $g \mapsto H_{\infty}(g)$ is a.s. an element of \mathbb{H} (i.e. it has almost surely uniformly continuous sample paths on \mathcal{G} with respect to the $L^2(\mu)$ metric) and b_n is a sequence such that $b_n \rightarrow \infty$ as $n \rightarrow \infty$.

When $\hat{\mu}$ is the empirical distribution based on a random sample from the population with distribution μ , then $b_n = \sqrt{n}$ and $H_{\infty} = B_{\mu}$, where B_{μ} is a μ -Brownian Bridge, i.e. a Gaussian process with zero mean and covariance function $(g_1, g_2) \mapsto \mu(g_1 g_2) - \mu(g_1)\mu(g_2)$. In this case condition S.4 imposes that the function class

$$\mathcal{G} = \{x \mapsto 1(f(x) \leq \delta) : f \in \mathcal{F}, \delta \in \mathcal{V}\}$$

is μ -Donsker. Note that \mathcal{F} is the parameter space that contains $\Delta(x)$ as well as $\hat{\Delta}(x)$ in S.3. In parametric models for the PE where $\mathcal{F} = \{f(x, \theta) : \theta \in \Theta\}$, f is known, $\theta \subseteq \mathbb{R}^{d_{\theta}}$ with $d_{\theta} < \infty$, and $x \mapsto f(x, \theta)$ is \mathcal{C}^1 on \mathcal{X} for all $\theta \in \Theta$, the class \mathcal{G} is μ -Donsker under mild conditions specified for example in van der Vaart (1998, Chap. 19). Examples 1 and 2 specify the PE parametrically. Lemma F.1 in the SM gives other sufficient conditions for the Donsker property.

The following result is derived as a consequence of the new mathematical results on the Hadamard differentiability of the sorting operator, stated in Lemma A.2 in the Appendix (proof given in SM due to space constraints), in conjunction with the functional delta method. It shows that the empirical SPE-function follows a FCLT over sets of quantiles corresponding to Δ_{μ}^* pre-images of compact sets of \mathbb{R} .

Define \mathcal{D} as a compact set consisting of regular values of $x \mapsto \Delta(x)$ on $\overline{\mathcal{X}}$, and $\mathcal{U} := \{\tilde{u} \in [0, 1] : \Delta_{\mu}^*(\tilde{u}) \in \mathcal{D}, f_{\Delta, \mu}(\Delta_{\mu}^*(\tilde{u})) > \varepsilon\}$, for a fixed $\varepsilon > 0$, where $f_{\Delta, \mu}(\Delta_{\mu}^*(\tilde{u}))$ is the density of $\Delta(X)$ defined in Lemma A.1(a). Let $r_n := a_n \wedge b_n$, the slowest of the rates of convergence of $\hat{\Delta}$ and $\hat{\mu}$. Assume $r_n/a_n \rightarrow s_{\Delta} \in [0, 1]$ and $r_n/b_n \rightarrow s_{\mu} \in [0, 1]$, where $s_{\Delta} = 0$ when $b_n = o(a_n)$ and $s_{\mu} = 0$ when $a_n = o(b_n)$. For example, $s_{\mu} = 0$ if μ is treated as known.

Theorem 4.1 (FCLT for $F_{\widehat{\Delta}, \widehat{\mu}}$ and $\widehat{\Delta}_\mu^*$). *Suppose that S.1-S.4 hold, and the convergence in S.3 and S.4 holds jointly. Then, as $n \rightarrow \infty$,*

(a) *The estimator of the distribution of PE obeys a functional central limit theorem, namely, in $\ell^\infty(\mathcal{D})$,*

$$r_n(F_{\widehat{\Delta}, \widehat{\mu}}(\delta) - F_{\Delta, \mu}(\delta)) \rightsquigarrow s_\Delta T_\infty(\delta) + s_\mu H_\infty(g_{\Delta, \delta}),$$

as a stochastic process indexed by $\delta \in \mathcal{D}$, where

$$T_\infty(\delta) := - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{G_\infty(x) \mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol}.$$

(b) *The empirical SPE-process obeys a functional central limit theorem, namely in $\ell^\infty(\mathcal{U})$,*

$$r_n(\widehat{\Delta}_\mu^*(u) - \Delta_\mu^*(u)) \rightsquigarrow - \frac{s_\Delta T_\infty(\Delta_\mu^*(u)) + s_\mu H_\infty(g_{\Delta, \Delta_\mu^*(u)})}{\int \frac{\mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol}} =: Z_\infty(u), \quad (4.6)$$

as a stochastic process indexed by $u \in \mathcal{U}$.

Remark 4.3 (Critical values). (a) Theorem 4.1 shows that $\delta \mapsto F_{\widehat{\Delta}, \widehat{\mu}}(\delta)$ ($u \mapsto \widehat{\Delta}_\mu^*(u)$) follows a FCLT over any compact set \mathcal{D} (the Δ_μ^* pre-image of \mathcal{D}), where \mathcal{D} excludes the critical values of $x \mapsto \Delta(x)$ on $\overline{\mathcal{X}}$. Thus, we can set $\mathcal{D} = \Delta(\overline{\mathcal{X}}) := \{\Delta(x) : x \in \overline{\mathcal{X}}\}$ when the map $x \mapsto \Delta(x)$ does not have critical points on $\overline{\mathcal{X}}$. This case is nice because it allows us not to worry about critical values when performing inference, and practically relevant as it occurs very naturally in many applications. For instance, it arises whenever $\Delta(x)$ is strictly locally monotonic in some direction. (b) In numerical examples reported in the SM, we find that the bootstrap inference method proposed performs well even in models where $x \mapsto \Delta(x)$ has critical points, without excluding the corresponding critical values from \mathcal{D} . This evidence suggests that the exclusion of critical values might not be necessary for inference. \square

4.2. Detailed Large Sample Theory for CA. It is convenient to modify the notation for the u -CA separating the dependence on $\Delta_\mu^*(u)$ from Δ and μ and specifying the characteristic of interest as φ_t . Moreover, when $Z = (X, Y)$ we remove the dependence on Y by taking expectations conditional on X . Let $\Lambda_{\widehat{\Delta}, \widehat{\mu}, \widehat{\Delta}_\mu^*(u)}(\varphi_t) := \widehat{\Lambda}_{\Delta, \mu}^u(t)$ and $\Lambda_{\Delta, \mu, \Delta_\mu^*(u)}(\varphi_t) := \Lambda_{\Delta, \mu}^u(t)$, where $\varphi_t \in \mathcal{F}_M \cup \mathcal{F}_I$, $t = (t_1, \dots, t_{d_z}) \in \mathbb{R}^{d_z}$, $u \in \mathcal{U}$, $\mathcal{F}_M := \{\int z_1^{t_1} \dots z_{d_z}^{t_{d_z}} d\mu(y | x) : t_1, \dots, t_{d_z} \in \{0, 1, 2, \dots\}, \int |z_1^{t_1} \dots z_{d_z}^{t_{d_z}}| d\mu(z) < \infty, t_1 + \dots + t_{d_z} \leq M\}$, M is some fixed integer, $\mu(y | x)$ is the distribution of Y at y conditional on $X = x$, and $\mathcal{F}_I := \{\int 1(z_1 \leq t_1, \dots, z_{d_z} \leq t_{d_z}) d\mu(y | x) : t_1, \dots, t_{d_z} \in \mathbb{R}\}$. For example, $\varphi_t(x) = x^{t_x} \mathbb{E}[Y^{t_y} | X = x]$ or $\varphi_t(x) = 1(x \leq t_x) \mu(t_y | x)$ for $t = (t_x, t_y)$. To derive the properties of $\Lambda_{\widehat{\Delta}, \widehat{\mu}, \widehat{\Delta}_\mu^*(u)}(\varphi_t)$, we use that the class of functions $\widetilde{\mathcal{G}} = \{1(f \leq \delta) \varphi : \varphi \in \mathcal{F}_M \cup \mathcal{F}_I, \delta \in \mathcal{V}, f \in \mathcal{F}\}$ is μ -Donsker. When $x \mapsto \mu(y | x)$ is continuous, this property holds by assumption S.4 when $\widehat{\mu}$ is the empirical distribution.¹⁵

¹⁵Lemma F.2 in the SM gives other sufficient conditions for the Donsker property.

The following result is derived as a consequence of the new mathematical results on the Hadamard differentiability of the classification operator, stated in Lemma A.3 in the Appendix (proof given in SM due to space constraints), in conjunction with the functional delta method.

Theorem 4.2 (FCLT for $\Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t)$). *Suppose that S.1-S.4 hold, the convergence in S.3 and S.4 holds jointly, and $u \in \mathcal{U}$. If $Z = (X, Y)$, then assume that \mathcal{Y} is compact and $x \mapsto \mu(y | x)$ is continuous on $B(\mathcal{X})$ for all $y \in \mathcal{Y}$. Then, as $n \rightarrow \infty$, (a) $\Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t)$ obeys a FCLT with respect to $t \mapsto \varphi_t \in \mathcal{F}_M$, namely, in $\ell^\infty(\mathbb{R}^{d_z})^2$,*

$$\begin{aligned} r_n \left(\Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t) - \Lambda_{\Delta, \mu, \Delta_\mu^*(u)}(\varphi_t) \right) \\ \rightsquigarrow \int_{\mathcal{M}_\Delta(\delta)} \tilde{\varphi}_t(x) \frac{Z_\infty(u) - s_\Delta G_\infty(x)}{\|\partial \Delta(x)\|} \mu'(x) d\text{Vol} + s_\mu H_\infty(h_{\Delta, \delta, \varphi_t}) =: Z_\infty^u(t), \end{aligned}$$

as a stochastic process indexed by $t \in \mathbb{R}^{d_z}$, where $\tilde{\varphi}_t(x) = [\varphi_t(x) - \Lambda_{\Delta, \mu, \delta}(\varphi_t)]/F_{\Delta, \mu}(\delta)$, $\tilde{h}_{\Delta, \delta, \varphi_t} := \tilde{\varphi}_t(x) 1\{\Delta(x) \leq \delta\}$, and $Z_\infty(u)$ is the limit process of Theorem 4.1; and (b) if in addition Assumption AS.1 holds, then $\Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t)$ obeys the same FCLT with respect to $t \mapsto \varphi_t \in \mathcal{F}_I$.

Assumption AS.1 is a technical condition stated in Appendix E of the SM to deal with the discontinuity of the indicator functions when $\varphi_t \in \mathcal{F}_I$. A sufficient condition for AS.1 is that

$$\int_{\mathcal{M}_\Delta(\delta) \cap \{x: x_k = t_k\}} d\text{Vol} = 0$$

holds uniformly over all $\delta \in \mathcal{V}$, $t_k \in \mathbb{R}$ and $k = 1, 2, \dots, d_x$. In other words, the manifold $\mathcal{M}_\Delta(\delta)$ and the set of points $\{x : x_k = t_k\}$ can not have an intersection with positive volume of $(d_x - 1)$ -dimension.

4.3. Bootstrap Inference for SPE and CA. Corollaries 2.1 and 2.2 use critical values of statistics related to the limit processes Z_∞ and Z_∞^u to construct confidence bands and p-values. These critical values can be hard to obtain in practice. In principle one can use simulation, but it might be difficult to numerically locate and parametrize the manifold $\mathcal{M}_\Delta(\delta)$, and to evaluate the integrals on $\mathcal{M}_\Delta(\delta)$ needed to compute the realizations of $Z_\infty(u)$ and $Z_\infty^u(t)$. This creates a real challenge to implement our inference methods. To deal with this challenge we employ (exchangeable) bootstrap to compute critical values (Præstgaard and Wellner, 1993; van der Vaart and Wellner, 1996) instead of simulation. We show that the bootstrap law is consistent to approximate the distribution of the limit processes of Theorems 4.1 and 4.2.

To state the bootstrap validity result formally, we follow the notation and definitions in van der Vaart and Wellner (1996). Let D_n denote the data vector and let $B_n = (\omega_1, \dots, \omega_n)$ be the vector of bootstrap weights. Consider a random element $\tilde{Z}_n = Z_n(D_n, B_n)$ in a normed space \mathbb{D} . We say that the bootstrap law of \tilde{Z}_n consistently estimates the law of some tight random element Z_∞

and write $\tilde{Z}_n \rightsquigarrow_{\mathbb{P}} Z_\infty$ if

$$\sup_{h \in \text{BL}_1(\mathbb{D})} |\mathbb{E}_{B_n} h(\tilde{Z}_n) - \mathbb{E}_{\mathbb{P}} h(Z_\infty)| \rightarrow_{\mathbb{P}} 0,$$

where $\text{BL}_1(\mathbb{D})$ denotes the space of functions with Lipschitz norm at most 1; \mathbb{E}_{B_n} denotes the conditional expectation with respect to B_n given the data D_n ; $\mathbb{E}_{\mathbb{P}}$ denotes the expectation with respect to \mathbb{P} , the distribution of the data D_n ; and $\rightarrow_{\mathbb{P}}$ denotes convergence in (outer) probability.

The next result is a consequence of the functional delta method for the exchangeable bootstrap. Let $\Lambda_{\tilde{\Delta}, \tilde{\mu}, \tilde{\Delta}_\mu^*(u)}(\varphi_t) := \tilde{\Lambda}_{\tilde{\Delta}, \mu}^u(t)$, the bootstrap draw of $\hat{\Lambda}_{\tilde{\Delta}, \mu}^u(t)$ defined in Algorithm 2.2.

Theorem 4.3 (Bootstrap FCLT for $\hat{\Delta}_\mu^*$ and $\Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t)$). *Suppose that the bootstrap is consistent for the law of the estimator of the PE, namely $a_n(\tilde{\Delta} - \hat{\Delta}) \rightsquigarrow_{\mathbb{P}} G_\infty$ in $\ell^\infty(B(\mathcal{X}))$, and for the law of the estimated measure, namely $b_n(\tilde{\mu} - \hat{\mu}) \rightsquigarrow_{\mathbb{P}} H_\infty$ in \mathbb{H} . Then, (1) under the assumptions of Theorem 4.1, the bootstrap is consistent for the law of the empirical SPE-process, namely*

$$r_n(\tilde{\Delta}_\mu^*(u) - \hat{\Delta}_\mu^*(u)) \rightsquigarrow_{\mathbb{P}} Z_\infty(u) \text{ in } \ell^\infty(\mathcal{U});$$

and (2) under the assumptions of Theorem 4.2, the bootstrap is consistent for the law of the empirical CA-process, namely

$$r_n \left(\Lambda_{\tilde{\Delta}, \tilde{\mu}, \tilde{\Delta}_\mu^*(u)}(\varphi_t) - \Lambda_{\hat{\Delta}, \hat{\mu}, \hat{\Delta}_\mu^*(u)}(\varphi_t) \right) \rightsquigarrow Z_\infty^u(t) \text{ in } \ell^\infty(\mathbb{R}^{d_z})^2.$$

Theorem 4.3 employs the high-level condition that the bootstrap can approximate consistently the laws of $\hat{\Delta}$ and $\hat{\mu}$, after suitable rescaling. In Examples 1-3 when $\hat{\mu}$ is the empirical measure based on the random sample of size n , the exchangeable bootstrap method entails randomly reweighing the sample using the weights $(\omega_1, \dots, \omega_n)$, which include empirical bootstrap and i.i.d. exponential weights, for example. In this case the high level condition holds if the weights satisfy the conditions stated in equation (3.6.8) of van der Vaart and Wellner (1996). We refer to van der Vaart and Wellner (1996) and Chernozhukov, Fernández-Val, and Melly (2013) for bootstrap FCLT for parametric and semi parametric estimators of Δ including least squares, quantile regression, and distribution regression, as well as nonparametric estimators of μ including the empirical distribution function.

APPENDIX A. KEY NEW MATHEMATICAL RESULTS: HADAMARD DIFFERENTIABILITY OF SORTING AND CLASSIFICATION OPERATORS

A.1. Notation. We denote the PE as $\Delta(x)$, the empirical PE as $\hat{\Delta}(x)$, and $\partial\Delta(x) := \partial\Delta(x)/\partial x$, the gradient of $x \mapsto \Delta(x)$. For a vector $v = (v_1, \dots, v_{d_v}) \in \mathbb{R}^{d_v}$, $\|v\|$ denotes the Euclidian norm of v , that is $\|v\| = \sqrt{v^\top v}$, where the superscript \top denotes transpose.

A.2. Basic Analytical Properties of Sorted Functions. The following lemma establishes the properties of the distribution function $\delta \mapsto F_{\Delta,\mu}(\delta)$ and the SPE-function $u \mapsto \Delta_\mu^*(u)$.

Define \mathcal{D} as a compact set consisting of regular values of $x \mapsto \Delta(x)$ on $\overline{\mathcal{X}}$.

Lemma A.1 (Basic Properties of $F_{\Delta,\mu}$ and Δ_μ^*). *Under conditions S.1 and S.2:*

1. *For any $\delta \in \mathcal{D}$, the derivative of $F_{\Delta,\mu}(\delta)$ with respect to δ is:*

$$f_{\Delta,\mu}(\delta) := \partial_\delta F_{\Delta,\mu}(\delta) = \int_{\mathcal{M}_\Delta(\delta)} \frac{\mu'(x)}{\|\partial\Delta(x)\|} d\text{Vol}. \quad (\text{A.7})$$

This integral is well-defined because the gradient $x \mapsto \partial\Delta(x)$ is finite, continuous, and bounded away from 0 on $\mathcal{M}_\Delta(\delta) \subseteq \overline{\mathcal{X}}$. The map $\delta \mapsto f_{\Delta,\mu}(\delta)$ is uniformly continuous on \mathcal{D} .

2. *Fix $\varepsilon > 0$, then for any $u \in \mathcal{U} := \{\tilde{u} \in [0, 1] : \Delta_\mu^*(\tilde{u}) \in \mathcal{D}, f_{\Delta,\mu}(\Delta_\mu^*(\tilde{u})) > \varepsilon\}$, the derivative of $\Delta_\mu^*(u)$ respect to u is:*

$$\partial_u \Delta_\mu^*(u) = \frac{1}{f_{\Delta,\mu}(\Delta_\mu^*(u))}. \quad (\text{A.8})$$

Moreover, the derivative map $u \mapsto \partial_u \Delta_\mu^(u)$ is uniformly continuous on \mathcal{U} .*

A.3. Functional Derivatives of Sorting-Related Operators. We consider the properties of the distribution function and the SPE-function as functional operators $(\Delta, \mu) \mapsto F_{\Delta,\mu}$ and $(\Delta, \mu) \mapsto \Delta_\mu^*$. We show that these operators are Hadamard differentiable with respect to (Δ, μ) . These results are critical ingredients to deriving the large sample distributions of the empirical versions of $F_{\Delta,\mu}$ and Δ_μ^* in Section 4.

We now recall the definition of uniform Hadamard differentiability from van der Vaart and Wellner (1996).

Definition A.1 (Hadamard Derivative Uniformly in an Index). Suppose the linear spaces \mathbb{D} and \mathbb{E} are equipped with the norms $\|\cdot\|_{\mathbb{D}}$ and $\|\cdot\|_{\mathbb{E}}$, and Θ is a compact subset of a metric space. A map $\phi_\theta : \mathbb{D}_\phi \subseteq \mathbb{D} \rightarrow \mathbb{E}$ is called Hadamard-differentiable uniformly in $\theta \in \Theta$ at $f \in \mathbb{D}_\phi$ tangentially to a subspace $\mathbb{D}_0 \subseteq \mathbb{D}$ if there is a continuous linear map $\partial_f \phi_\theta : \mathbb{D}_0 \rightarrow \mathbb{E}$ such that uniformly in $\theta \in \Theta$:

$$\frac{\phi_\theta(f + t_n h_n) - \phi_\theta(f)}{t_n} - \partial_f \phi_\theta[h] \rightarrow 0, \quad n \rightarrow \infty, \quad (\text{A.9})$$

for all converging real sequences $t_n \rightarrow 0$ and $\|h_n - h\|_{\mathbb{D}} \rightarrow 0$ such that $f + t_n h_n \in \mathbb{D}_\phi$ for every n , and $h \in \mathbb{D}_0$; moreover, the map $(\theta, h) \mapsto \partial_f \phi_\theta[h]$ is continuous on $\Theta \times \mathbb{D}_0$.

In what follows, we let \mathbb{F} denote the space of continuous functions on $B(\mathcal{X})$ equipped with the sup-norm, and \mathbb{F}_0 denote a subset of \mathbb{F} that contains uniformly continuous functions.

Lemma A.2 (Hadamard differentiability of $(\Delta, \mu) \mapsto F_{\Delta,\mu}$ and $(\Delta, \mu) \mapsto \Delta_\mu^*$). *Let $\mathbb{D} := \mathbb{F} \times \mathbb{H}$ and $\mathbb{D}_0 := \mathbb{F}_0 \times \mathbb{H}$. Assume that S.1-S.2 hold. Then,*

(a) The map $(\Delta, \mu) \mapsto F_{\Delta, \mu}(\delta)$, mapping $\mathbb{D} \rightarrow \mathbb{R}$, is Hadamard differentiable uniformly in $\delta \in \mathcal{D}$ at (Δ, μ) tangentially to \mathbb{D}_0 with the derivative map $\partial_{\Delta, \mu} F_{\Delta, \mu}(\delta) : \mathbb{D}_0 \rightarrow \mathbb{R}$ defined by

$$(G, H) \mapsto \partial_{\Delta, \mu} F_{\Delta, \mu}(\delta)[G, H] := - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{G(x)\mu'(x)}{\|\partial\Delta(x)\|} d\text{Vol} + H(g_{\Delta, \delta}).$$

(b) The map $(\Delta, \mu) \mapsto \Delta_{\mu}^*(u)$, mapping $\mathbb{D} \rightarrow \mathbb{R}$ is Hadamard differentiable uniformly in $u \in \mathcal{U}$ at (Δ, μ) tangentially to \mathbb{D}_0 with the derivative map, $\partial_{\Delta, \mu} \Delta_{\mu}^*(u) : \mathbb{D}_0 \rightarrow \mathbb{R}$, defined by

$$(G, H) \mapsto \partial_{\Delta, \mu} \Delta_{\mu}^*(u)[G, H] := - \frac{\partial_{\Delta, \mu} F_{\Delta, \mu}(\Delta_{\mu}^*(u))[G, H]}{f_{\Delta, \mu}(\Delta_{\mu}^*(u))}.$$

A.4. Functional Derivatives of Classification Operators. Let $\tilde{\mathbb{D}} := \mathbb{F} \times \tilde{\mathbb{H}} \times \mathbb{R}$ and $\tilde{\mathbb{D}}_0 := \mathbb{F}_0 \times \tilde{\mathbb{H}} \times \mathbb{R}$, where \mathbb{F} and \mathbb{F}_0 are defined as before; $\tilde{\mathbb{H}}$ is the set of bounded linear operators mapping from the set $\tilde{\mathcal{G}} := \{\varphi 1(\Delta \leq \delta) : f \in \mathcal{F}, \varphi \in \mathcal{F}_I \cup \mathcal{F}_M, \delta \in \mathcal{V}\}$ to \mathbb{R} , with norm

$$\|H\|_{\tilde{\mathcal{G}}} = \sup_{g \in \tilde{\mathcal{G}}} |H(g)|,$$

where the map $g \mapsto H(g)$ is uniformly continuous on $g \in \tilde{\mathcal{G}}$ under the $L^2(\mu)$ norm. We derive the properties of the least affected classification operator $\Lambda_{\Delta, \mu, \delta}^- : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ defined by

$$\Lambda_{\Delta, \mu, \delta}^-(\varphi_t) := \int \varphi_t(x) 1\{\Delta(x) \leq \delta\} d\mu(x) / \int 1\{\Delta(x) \leq \delta\} d\mu(x),$$

where $\varphi_t \in \mathcal{F}_M$ for moments and $\varphi_t \in \mathcal{F}_I$ for distributions of the components of Z , and $\delta = \Delta_{\mu}^*(u)$ for some $u \in \mathcal{U}$. The properties of the most affected operator $\Lambda_{\Delta, \mu, \delta}^+ : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ can be derived using similar arguments, which are omitted for brevity.

Lemma A.3 (Hadamard differentiability of $(\Delta, \mu, \delta) \mapsto \Lambda_{\Delta, \mu, \delta}^-$). *Assume that Assumptions S.1 and S.2 hold, $\delta \in \mathcal{D}$, and $F_{\Delta, \mu}(\delta) > 0$. Then,*

(a) *The map $\Lambda_{\Delta, \mu, \delta}^-(\varphi_t) : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $\varphi_t \in \mathcal{F}_M$ at (Δ, μ, δ) tangentially to $\tilde{\mathbb{D}}_0$.*

(b) *If in addition Assumption AS.1 stated in Appendix E of the SM holds, the map $\Lambda_{\Delta, \mu, \delta}^-(\varphi_t) : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $\varphi_t \in \mathcal{F}_I$ at (Δ, μ, δ) tangentially to $\tilde{\mathbb{D}}_0$.*

(c) *The derivative map $\partial_{\Delta, \mu, \delta} \Lambda_{\Delta, \mu, \delta}^-(\varphi_t) : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ is defined by:*

$$(G, H, K) \mapsto \partial_{\Delta, \mu, \delta} \Lambda_{\Delta, \mu, \delta}^-(\varphi_t)[G, H, K] := \int_{\mathcal{M}_{\Delta}(\delta)} \tilde{\varphi}_t(x) \frac{K - G(x)}{\|\partial\Delta(x)\|} d\text{Vol} + H(\tilde{h}_{\Delta, \delta, \varphi_t}),$$

where $\tilde{\varphi}_t(x) = [\varphi_t(x) - \Lambda_{\Delta, \mu, \delta}^-(\varphi_t)] / \int 1(\Delta(x) \leq \delta) d\mu(x)$ and $\tilde{h}_{\Delta, \delta, \varphi_t} := \tilde{\varphi}_t(x) 1\{\Delta(x) \leq \delta\}$.

APPENDIX B. PROOFS OF SECTION 4

We first recall Theorem 3.9.4 of van der Vaart and Wellner (1996).

Lemma B.1 (Delta-method). *Let \mathbb{D} and \mathbb{E} be metrizable topological vector spaces, and Θ is a compact subset of a metric space. Let $\phi_\theta : \mathbb{D}_\phi \subseteq \mathbb{D} \rightarrow \mathbb{E}$ be a Hadamard differentiable mapping uniformly in $\theta \in \Theta$ at $f \in \mathbb{D}$ tangentially to $\mathbb{D}_0 \subseteq \mathbb{D}$, with derivative $\partial_f \phi_\theta$. Let $\widehat{f}_n : \Omega_n \rightarrow \mathbb{D}_\phi$ be stochastic maps taking values in \mathbb{D}_ϕ such that $r_n(\widehat{f}_n - f) \rightsquigarrow J_\infty$ for some sequence of constants $r_n \rightarrow \infty$, where J_∞ is separable and takes values in \mathbb{D}_0 . Then $r_n(\phi_\theta(\widehat{f}_n) - \phi_\theta(f)) \rightsquigarrow \partial_f \phi_\theta[J_\infty]$, as a stochastic process indexed by $\theta \in \Theta$.*

Proof of Theorem 4.1. The statements follow directly from Lemma A.2, and Lemma B.1, by setting $\phi_\theta = F_{\Delta, \mu}(\delta)$ with $\theta = \delta$ or $\phi_\theta = \Delta_\mu^*(u)$ with $\theta = u$, $\mathbb{D}_\phi = \mathbb{D} = \mathbb{F} \times \mathbb{H}$, $\mathbb{E} = \mathbb{R}$, $\mathbb{D}_0 = \mathbb{F}_0 \times \mathbb{H}$, $f = (\Delta, \mu)$, $\widehat{f}_n = (\widehat{\Delta}, \widehat{\mu})$, and $J_\infty = (s_\Delta G_\infty, s_\mu H_\infty)$. The expression of $\partial_f \phi_\theta$ for each statement is the Hadamard derivative in the corresponding statement of Lemma A.2. \square

Proof of Theorem 4.2. The statements follow directly from Lemma A.4, and Lemma B.1, by setting $\phi_\theta = \Lambda_{\Delta, \mu, \delta}^-$, $\theta = t$, $\mathbb{D}_\phi = \mathbb{D} = \mathbb{F} \times \widetilde{\mathbb{H}} \times \mathbb{R}$, $\mathbb{E} = \mathbb{R}$, $\mathbb{D}_0 = \mathbb{F}_0 \times \widetilde{\mathbb{H}} \times \mathbb{R}$, $f = (\Delta, \mu, \Delta_\mu^*(u))$, $\widehat{f}_n = (\widehat{\Delta}, \widehat{\mu}, \widehat{\Delta}_\mu^*(u))$, and $J_\infty = (s_\Delta G_\infty, s_\mu H_\infty, Z_\infty)$. The expression of $\partial_f \phi_\theta$ for each statement is the Hadamard derivative in the corresponding statement of Lemma A.4. \square

To prove Theorem 4.3, we recall Theorem 3.9.11 of van der Vaart (1998). Here we use the notation for bootstrap convergence \rightsquigarrow_P defined in Section 4.3.

Lemma B.2 (Delta-method for bootstrap in probability). *Let \mathbb{D} and \mathbb{E} be metrizable topological vector spaces, and Θ is a compact subset of a metric space. Let $\phi_\theta : \mathbb{D}_\phi \subseteq \mathbb{D} \mapsto \mathbb{E}$ be a Hadamard-differentiable mapping uniformly in $\theta \in \Theta$ at f tangentially to \mathbb{D}_0 with derivative $\partial_f \phi_\theta$. Let \widehat{f}_n be a random element such that $r_n(\widehat{f}_n - f) \rightsquigarrow J_\infty$. Let \widetilde{f}_n be a stochastic map in \mathbb{D} , produced by a bootstrap method, such that $r_n(\widetilde{f}_n - \widehat{f}_n) \rightsquigarrow_P J_\infty$. Then, $r_n(\phi_\theta(\widetilde{f}_n) - \phi_\theta(\widehat{f}_n)) \rightsquigarrow_P \partial_f \phi_\theta[J_\infty]$, as a stochastic process indexed by $\theta \in \Theta$.*

Proof of Theorem 4.3. The statement (1) follows directly from Lemma A.2, and Lemma B.2, by setting $\phi_\theta = \Delta_\mu^*(u)$, $\theta = u$, $\mathbb{D}_\phi = \mathbb{D} = \mathbb{F} \times \mathbb{H}$, $\mathbb{E} = \mathbb{R}$, $\mathbb{D}_0 = \mathbb{F}_0 \times \mathbb{H}$, $f = (\Delta, \mu)$, $\widehat{f}_n = (\widehat{\Delta}, \widehat{\mu})$, and $J_\infty = (s_\Delta G_\infty, s_\mu H_\infty)$. The expression of $\partial_f \phi_\theta$ is the Hadamard derivative in statement (b) of Lemma A.2. The statement (2) follows directly from Lemma A.3, and Lemma B.2, by setting $\phi_\theta = \Lambda_{\Delta, \mu, \delta}^-$, $\theta = t$, $\mathbb{D}_\phi = \mathbb{D} = \mathbb{F} \times \widetilde{\mathbb{H}} \times \mathbb{R}$, $\mathbb{E} = \mathbb{R}$, $\mathbb{D}_0 = \mathbb{F}_0 \times \widetilde{\mathbb{H}} \times \mathbb{R}$, $f = (\Delta, \mu, \Delta_\mu^*(u))$, $\widehat{f}_n = (\widehat{\Delta}, \widehat{\mu}, \widehat{\Delta}_\mu^*(u))$, and $J_\infty = (s_\Delta G_\infty, s_\mu H_\infty, Z_\infty)$. The expression of $\partial_f \phi_\theta$ is the Hadamard derivative in statement (c) of Lemma A.3. \square

Supplement to “The Sorted Effects Method: Discovering Heterogeneous Effects Beyond Their Averages”

Victor Chernozhukov, Iván Fernández-Val, Ye Yuo

ABSTRACT. The supplementary material contains 7 appendices with additional results and some omitted proofs. Appendix C introduces some notation. Appendix D includes a brief review of differential geometry. Appendix E gathers the proofs of the key mathematical results in Appendix A. Appendix F provides sufficient conditions for the μ -Donsker properties in Section 4. Appendix G extends the theoretical analysis to include discrete covariates. Appendices H and I report the results of 3 numerical simulations and an empirical application to the effect of race on mortgage denials, respectively.

APPENDIX C. NOTATION

For a possibly multivariate random variable X , \mathcal{X} denotes the interior of the support of X in the part of the population of interest, μ denotes the distribution of X over \mathcal{X} , and $\hat{\mu}$ denotes an estimator of μ . We denote the expectation with respect to the distribution $\tilde{\mu}$ by $E_{\tilde{\mu}}$. We denote the PE as $\Delta(x)$, the empirical PE as $\hat{\Delta}(x)$, and $\partial\Delta(x) := \partial\Delta(x)/\partial x$, the gradient of $x \mapsto \Delta(x)$. We also use $a \wedge b$ to denote the minimum of a and b . For a vector $v = (v_1, \dots, v_{d_v}) \in \mathbb{R}^{d_v}$, $\|v\|$ denotes the Euclidian norm of v , that is $\|v\| = \sqrt{v^\top v}$, where the superscript \top denotes transpose. For a non-negative integer r and an open set \mathcal{K} , the class \mathcal{C}^r on \mathcal{K} includes the set of r times continuously differentiable real valued functions on \mathcal{K} . The symbol \rightsquigarrow denotes weak convergence (convergence in distribution), and \rightarrow_P denotes convergence in (outer) probability.

APPENDIX D. BACKGROUND ON DIFFERENTIAL GEOMETRY

We recall some definitions from differential geometry that are used in the analysis. For a continuously differentiable function $\Delta : B(\mathcal{X}) \rightarrow \mathbb{R}$ defined on an open set $B(\mathcal{X}) \subseteq \mathbb{R}^{d_x}$ containing the set \mathcal{X} , $x \in \mathcal{X}$ is a *critical point* of Δ on \mathcal{X} , if

$$\partial\Delta(x) = 0, \tag{D.10}$$

where $\partial\Delta(x)$ is the gradient of $\Delta(x)$; otherwise x is a *regular point* of Δ on \mathcal{X} . A value δ is a *critical value* of Δ on \mathcal{X} if the set $\{x \in \mathcal{X} : \Delta(x) = \delta\}$ contains at least one critical point; otherwise δ is a *regular value* of Δ on \mathcal{X} .

In the multi-dimensional space, $d_x > 1$, a function Δ can have continuums of critical points. For example, the function $\Delta(x_1, x_2) = \cos(x_1^2 + x_2^2)$ has continuums of critical points on the circles $x_1^2 + x_2^2 = k\pi$ for each positive integer k .

We recall now several core concepts related to manifolds from Spivak (1965) and Munkres (1991).

Definition D.1 (Manifold). Let d_k , d_x and r be positive integers such that $d_x \geq d_k$. Suppose that \mathcal{M} is a subspace of \mathbb{R}^{d_x} that satisfies the following property: for each point $m \in \mathcal{M}$, there is a set \mathcal{V} containing m that is open in \mathcal{M} , a set \mathcal{K} that is open in \mathbb{R}^{d_k} , and a continuous map $\alpha_m : \mathcal{K} \rightarrow \mathcal{V}$ carrying \mathcal{K} onto \mathcal{V} in a one-to-one fashion, such that: (1) α_m is of class \mathcal{C}^r on \mathcal{K} , (2) $\alpha_m^{-1} : \mathcal{V} \rightarrow \mathcal{K}$ is continuous, and (3) the Jacobian matrix of α_m , $D\alpha_m(k)$, has rank d_k for each $k \in \mathcal{K}$. Then \mathcal{M} is called a d_k -manifold without boundary in \mathbb{R}^{d_x} of class \mathcal{C}^r . The map α_m is called a *coordinate patch* on \mathcal{M} about m . A set of coordinate patches that covers \mathcal{M} is called an *atlas*.

Definition D.2 (Connected Branch). For any subset \mathcal{M} of a topological space, if any two points m_1 and m_2 cannot be connected via path in \mathcal{M} , then we say that m_1 and m_2 are not connected. Otherwise, we say that m_1 and m_2 are connected. We say that $\mathcal{V} \subseteq \mathcal{M}$ is a *connected branch* of \mathcal{M} if all points of \mathcal{V} are connected to each other and do not connect to any points in $\mathcal{M} \setminus \mathcal{V}$.

Definition D.3 (Volume). For a $d_x \times d_k$ matrix $A = (x_1, x_2, \dots, x_{d_k})$ with $x_i \in \mathbb{R}^{d_x}$, $1 \leq i \leq d_k \leq d_x$, let $\text{Vol}(A) = \sqrt{\det(A^T A)}$, which is the *volume* of the parallelepiped $P(A)$ with edges given by the columns of A , $P(A) = \{c_1 x_1 + \dots + c_{d_k} x_{d_k} : 0 \leq c_i \leq 1, i = 1, \dots, d_k\}$.

The volume measures the amount of mass in \mathbb{R}^{d_k} of a d_k -dimensional parallelepiped in \mathbb{R}^{d_x} , $d_k \leq d_x$. This concept is essential for integration on manifolds, which we will discuss shortly. First we recall the concept of integration on parameterized manifolds:

Definition D.4 (Integration on a parametrized manifold). Let \mathcal{K} be open in \mathbb{R}^{d_k} , and let $\alpha : \mathcal{K} \rightarrow \mathbb{R}^{d_x}$ be of class \mathcal{C}^r on \mathcal{K} , $r \geq 1$. The set $\mathcal{M} = \alpha(\mathcal{K})$ together with the map α constitute a *parametrized d_k -manifold* in \mathbb{R}^{d_x} of class \mathcal{C}^r . Let g be a real-valued continuous function defined at each point of \mathcal{M} . The *integral of g over \mathcal{M} with respect to volume* is defined by

$$\int_{\mathcal{M}} g(m) d\text{Vol} := \int_{\mathcal{K}} (g \circ \alpha)(k) \text{Vol}(D\alpha(k)) dk, \quad (\text{D.11})$$

provided that the right side integral exists. Here $D\alpha(k)$ is the Jacobian matrix of the mapping $k \mapsto \alpha(k)$, and $\text{Vol}(D\alpha(k))$ is the volume of matrix $D\alpha(k)$ as defined in Definition D.3.

The above definition coincides with the usual interpretation of integration. The integral can be extended to manifolds that do not admit a global parametrization α using the notion of partition of unity. This partition is a set of smooth local functions defined in a neighborhood of the manifold. The following Lemma shows the existence of the partition of unity and is proven in Lemma 25.2 in Munkres (1991).

Lemma D.1 (Partition of Unity on \mathcal{M} of class \mathcal{C}^∞). *Let \mathcal{M} be a d_k -manifold without boundary in \mathbb{R}^{d_x} of class \mathcal{C}^r , $r \geq 1$, and let ϑ be an open cover of \mathcal{M} . Then, there is a collection $\mathcal{P} = \{p_i \in$*

$\mathcal{C}^\infty : i \in \mathcal{I}\}$, where p_i is defined on an open set containing \mathcal{M} for all $i \in \mathcal{I}$, with the following properties: (1) For each $m \in \mathcal{M}$ and $i \in \mathcal{I}$, $0 \leq p_i(m) \leq 1$, (2) for each $m \in \mathcal{M}$ there is an open set $\mathcal{V} \in \vartheta$ containing m such that all but finitely many $p_i \in \mathcal{P}$ are 0 on \mathcal{V} , (3) for each $m \in \mathcal{M}$, $\sum_{p_i \in \mathcal{P}} p_i(m) = 1$, and (4) for each $p_i \in \mathcal{P}$ there is an open set $\mathcal{U} \in \vartheta$, such that $\text{supp}(p_i) \subseteq \mathcal{U}$.

Now we are ready to recall the definition of integration on a manifold.

Definition D.5 (Integration on a manifold with partition of unity). Let $\vartheta := \{\vartheta_j : j \in \mathcal{J}\}$ be an open cover of a d_k -manifold without boundary \mathcal{M} in \mathbb{R}^{d_x} of class \mathcal{C}^r , $r \geq 1$. Suppose there is an coordinate patch $\alpha_j : \mathcal{V}_j \subseteq \mathbb{R}^{d_k} \rightarrow \vartheta_j$, that is one-to-one and of class \mathcal{C}^r on \mathcal{V}_j for each $j \in \mathcal{J}$. Denote $\mathcal{K}_j = \alpha_j^{-1}(\mathcal{M} \cap \vartheta_j)$. Then for a real-valued continuous function g defined on an open set that contains \mathcal{M} , the *integral of g over \mathcal{M} with respect to volume* is defined by:

$$\int_{\mathcal{M}} g(m) d\text{Vol} := \sum_{j \in \mathcal{J}} \sum_{i \in \mathcal{I}} \int_{\mathcal{K}_j} [(p_i g) \circ \alpha_j](k) \text{Vol}(D\alpha_j(k)) dk, \quad (\text{D.12})$$

provided that the right side integrals exist, where $\{p_i \in \mathcal{C}^\infty : i \in \mathcal{I}\}$ is a partition of unity on \mathcal{M} of class \mathcal{C}^∞ that satisfies the conditions of Lemma D.1. Munkres (1991, p. 212) shows that the integral does not depend on the choice of cover and partition of unity.

APPENDIX E. PROOFS OF APPENDIX A

To analyze the analytical properties of the SPE-function, it is convenient to treat the PE as a multivariate real-valued function

$$\Delta : B(\mathcal{X}) \rightarrow \mathbb{R},$$

where $B(\mathcal{X}) \subseteq \mathbb{R}^{d_x}$ contains the set \mathcal{X} . Let μ be a distribution function. The distribution of Δ with respect to μ is the function $F_{\Delta, \mu} : \mathbb{R} \rightarrow [0, 1]$ with

$$F_{\Delta, \mu}(\delta) = \int 1\{\Delta(x) \leq \delta\} d\mu(x). \quad (\text{E.13})$$

The SPE-function is the map

$$\Delta_\mu^* : \mathcal{U} \subseteq [0, 1] \rightarrow \mathbb{R},$$

defined at each point as the left-inverse function of $F_{\Delta, \mu}$, i.e.,

$$\Delta_\mu^*(u) := F_{\Delta, \mu}^{\leftarrow}(u) := \inf_{\delta \in \mathbb{R}} \{F_{\Delta, \mu}(\delta) \geq u\}. \quad (\text{E.14})$$

From this functional perspective, the map $u \mapsto \Delta_\mu^*(u)$ is the result of applying a sorting operator to the map $x \mapsto \Delta(x)$ that sorts the values of Δ in increasing order weighted by μ . The next subsections provide the proofs of 3 results:

- 1) Lemma A.1, which characterizes some analytical properties of the distribution function $\delta \mapsto F_{\Delta, \mu}(\delta)$ and the sorted function $u \mapsto \Delta_\mu^*(u)$,
- 2) Lemma A.2, which derives the functional derivatives of $F_{\Delta, \mu}$ and Δ_μ^* with respect to Δ and μ , and

- 3) Lemma A.3, which derives the functional derivatives of the related classification operator $\Lambda_{\Delta, \mu, \delta}^-$ with respect to Δ , μ and δ .

E.1. Proof of Lemma A.1. We use the following results in the proof of Lemma A.1.

Lemma E.1. *If $\Delta : B(\mathcal{X}) \rightarrow \mathbb{R}$ is \mathcal{C}^1 on an open set $B(\mathcal{X}) \subseteq \mathbb{R}^{d_x}$, then for any compact subset $\overline{\mathcal{X}}$ of $B(\mathcal{X})$, the sets of critical points and critical values of $x \mapsto \Delta(x)$ on $\overline{\mathcal{X}}$ are closed.*

Proof. (1) Critical points: since $x \mapsto \partial\Delta(x)$ is continuous on $\overline{\mathcal{X}}$ and $\overline{\mathcal{X}}$ is compact, the set of points $x \in \overline{\mathcal{X}}$ such that $\partial\Delta(x) = 0$ is closed.

(2) Critical values: since $x \mapsto \Delta(x)$ is continuous and $\overline{\mathcal{X}}$ is compact, the image set $\Delta(\overline{\mathcal{X}})$ is a compact set in \mathbb{R} . For any sequence of critical values $\{\delta_i\}_{i \geq 1}^\infty$ in $\Delta(\overline{\mathcal{X}})$, there is a corresponding sequence $\{x_i\}_{i \geq 1}^\infty$ in $\overline{\mathcal{X}}$ such that $\Delta(x_i) = \delta_i$. Suppose $\{\delta_i\}_{i \geq 1}^\infty$ converges to $\delta_0 \in \Delta(\overline{\mathcal{X}})$. By compactness of $\overline{\mathcal{X}}$, we can find a converging subsequence of $\{x_i\}_{i \geq 1}^\infty$ with limit $x_0 \in \overline{\mathcal{X}}$ such that $\Delta(x_i) = \delta_i$. Then by continuity of $x \mapsto \partial\Delta(x)$, $\partial\Delta(x_0) = 0$. By continuity of $x \mapsto \Delta(x)$, $\Delta(x_0) = \delta_0$, and therefore $\delta_0 = \Delta(x_0)$ is a critical value of $\Delta(x)$. Hence the set of critical values is closed. \square

Lemma E.2. *For a compact set \mathcal{V} in a metric space \mathbb{D} , suppose there is an open cover $\{\theta_i : i \in I\}$ of \mathcal{V} . Then there exists a finite open sub-cover of \mathcal{V} and $\eta > 0$, such that for every point $x \in \mathcal{V}$, the η -ball around x is contained in the finite sub-cover.*

Proof of Lemma E.2. Since \mathcal{V} is a compact set in the metric space \mathbb{D} (with metric $\|\cdot\|_{\mathbb{D}}$), then any open cover $\{\theta_i : i \in I\}$ of \mathcal{V} has a finite open subcover $\{\tilde{\theta}_i : i = 1, 2, \dots, m\}$ which covers \mathcal{V} .

Let $\Theta = \cup_{i=1}^m \tilde{\theta}_i$. We prove the statement of the lemma by contradiction. Suppose for any $i > 0$, there exists some point $x_i \in \mathbb{D}$ such that $d(x_i, \mathcal{V}) := \inf_{v \in \mathcal{V}} \|x_i - v\|_{\mathbb{D}} < i^{-1}$ and $x_i \notin \Theta$. Then, by compactness of \mathcal{V} there exists $v_i \in \mathcal{V}$ such that $d(x_i, \mathcal{V}) = d(x_i, v_i) < i^{-1}$. Let v_0 be the limit of $\{v_i : i \geq 1\}$. By compactness of \mathcal{V} , $v_0 \in \mathcal{V}$. Since $d(x_i, v_0) \rightarrow 0$ as $i \rightarrow \infty$ and Θ is an open cover of \mathcal{V} , there must be a open ball $B(v_0)$ around v_0 such that $B(v_0) \subseteq \Theta$, which contradicts with $x_i \notin \Theta$, for i large enough. Therefore there must be an η such that the η -ball around any $x \in \mathcal{V}$ is covered by Θ . \square

Proof of Lemma A.1. The proof of statement (2) follows directly from the inverse function theorem.

The proof of statement (1) is divided in two steps. Step 1 constructs a finite set of open rectangles that covers the set $\mathcal{M}_{\Delta}(\delta)$ and has certain properties that allow us to apply a change of variable to the derivative of $\delta \mapsto F_{\Delta, \mu}(\delta)$. Step 2 expresses the derivative as an integral on a manifold.

For a subset $\mathcal{S} \subseteq \mathbb{R}^{d_x}$ and $\eta > 0$, define $B_\eta(\mathcal{S}) := \{x \in \mathbb{R}^{d_x} : d(x, \mathcal{S}) = \inf_{s \in \mathcal{S}} \|x - s\| < \eta\}$. Similarly, for any $\delta \in \mathbb{R}$ and $\eta > 0$, define $B_\eta(\delta) := (\delta - \eta, \delta + \eta)$. Without loss of generality, we assume that $\mathcal{M}_\Delta(\delta)$ only has one connected branch. We will discuss the case where $\mathcal{M}_\Delta(\delta)$ has multiple connected branches at the end of the proof of this lemma.

Step 1. For any regular value $\delta \in \mathcal{D}$, the set $\mathcal{M}_\Delta(\delta)$ is a $(d_x - 1)$ -manifold in \mathbb{R}^{d_x} of class \mathcal{C}^1 by Theorem 5-1 in Spivak (1965, p. 111). Denote $\widetilde{\mathcal{M}}_\Delta(\delta) := \{x \in B(\mathcal{X}) : \Delta(x) = \delta\}$ and $\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta)) := \cup_{\delta' \in B_\eta(\delta)} \widetilde{\mathcal{M}}_\Delta(\delta')$ for $\eta > 0$. These enlargements of the set $\mathcal{M}_\Delta(\delta)$ are used to apply a change of variable technique to integrals on $\mathcal{M}_\Delta(\delta)$.

By assumptions S.1-S.2, there exists $\eta_1 > 0$ small enough and $C > c > 0$ such that:

(1) $\overline{B_{\eta_1}(\delta)} := [\delta - \eta_1, \delta + \eta_1] \subseteq \Delta(\overline{\mathcal{X}}) := \{\Delta(x) : x \in \overline{\mathcal{X}}\}$ and contains no critical values of Δ on $\overline{\mathcal{X}}$, and $B_{\eta_1}(\overline{\mathcal{X}}) \subseteq B(\mathcal{X})$.

(2) $\inf_{x \in \overline{\widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\overline{\mathcal{X}})}} \|\partial\Delta(x)\| > c$.

(3) $\sup_{x \in \overline{\widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\overline{\mathcal{X}})}} \|\partial\Delta(x)\| < C$.

(4) For any $\eta < \eta_1$, $\widetilde{\mathcal{M}}_\Delta(\delta) \cap B_\eta(\overline{\mathcal{X}})$ is a $(d_x - 1)$ -manifold in \mathbb{R}^{d_x} of class \mathcal{C}^1 .

Indeed, by Lemma E.1, the set of regular values is open. Therefore, there exists a small neighborhood $B_\eta(\delta)$ with $\eta > 0$ such that there exists no critical value of Δ on $\overline{\mathcal{X}}$ in $B_\eta(\delta)$. Then any $\eta_1 < \eta$ satisfies statement (1). Statements (2) and (3) follow by the compactness of $\overline{\mathcal{X}}$, the continuity of mapping $x \mapsto \partial\Delta(x)$, and assumptions S.1 and S.2. Statement (4) is implied by Theorem 5-1 in Spivak (1965, p. 111).

Next, we establish a finite cover of $\widetilde{\mathcal{M}}_\Delta(B_{\eta_2}(\delta)) \cap B_{\eta_2}(\overline{\mathcal{X}})$ with certain good properties, for some $\eta_2 < \eta_1$.

For any $\eta_3 < \eta_1$, $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta)) \cap B_{\eta_3}(\overline{\mathcal{X}})$ satisfies the properties (2)–(4) stated above. Consider the rectangles $\theta(x) := X_1(x) \times \dots \times X_{d_x}(x)$ centered at $x = (x_1, \dots, x_{d_x})$ where $X_k(x) := (x_k - a_k(x), x_k + a_k(x))$, with $a_k(x) > 0$, $k = 1, 2, \dots, d_x$. Let $A(x) := \sup_{1 \leq k \leq d_x} a_k(x)$ be such that:

$$\overline{\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta)) \cap B_{\eta_3}(\overline{\mathcal{X}})} \subseteq \cup_{x \in \widetilde{\mathcal{M}}_\Delta(\delta) \cap B_{\eta_3}(\overline{\mathcal{X}})} \theta(x) \subseteq \widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\overline{\mathcal{X}}),$$

which can be fulfilled by using small enough η_3 .

By continuity of $x \mapsto \partial\Delta(x)$, for small enough $A(x)$ and any $x' \in \theta(x)$, there always exists an index $i(x) \in \{1, 2, \dots, d_x\}$ such that $|\partial_{x_{i(x)}} \Delta(x')| \geq \frac{c}{2\sqrt{d_x}}$ since $\|\partial\Delta(x')\| \geq c$ for all $x' \in \theta(x)$ by the property (2) above, where $\partial_x := \partial/\partial x$. Also we can find a finite set of $\theta(x)$'s, denoted as $\Theta := \{\theta(x^i)\}_{i=1}^m$, such that Θ forms a finite open cover of $\overline{\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta)) \cap B_{\eta_3}(\overline{\mathcal{X}})}$. We rename these open rectangles as $\theta_i := \theta(x^i)$, $i \in \{1, 2, \dots, m\}$, where $\theta_i = X_{i1} \times \dots \times X_{id_x}$ and $X_{ik} := X_k(x^i)$, $k \in \{1, \dots, d_x\}$.

For a given $i \in \{1, 2, \dots, m\}$, consider the center of θ_i , denoted as x^i . Without loss of generality, we can assume that $i(x^i) = d_x$. Then, for all $x' \in \theta(x^i)$, $|\partial_{x_{d_x}} \Delta(x')| \geq c/2\sqrt{d_x}$. This means that $\Delta(x)$ is partially monotonic in x_{d_x} on $\theta(x^i)$. By the implicit function theorem, there exists g such that $g(x'_1, x'_2, \dots, x'_{d_x-1}, \delta') = x'_{d_x}$, for any $x' = (x'_1, x'_2, \dots, x'_{d_x}) \in \widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta)) \cap \theta(x^i)$ and $\delta' = \Delta(x')$. Also by the implicit function theorem,

$$\partial g(x'_1, \dots, x'_{d_x-1}, \delta') = \frac{-(\partial_{x_1} \Delta(x'), \partial_{x_2} \Delta(x'), \dots, \partial_{x_{d_x-1}} \Delta(x'), -1)}{\partial_{d_x} \Delta(x')}.$$

So $\|\partial g(x'_1, \dots, x'_{d_x-1}, \delta')\| \leq \frac{\|\partial \Delta(x')\|}{|\partial_{d_x} \Delta(x')|} \leq \frac{2(C+1)\sqrt{d_x}}{c} := \Lambda$ because $|\partial_{x_{d_x}} \Delta(x')| \geq c/2\sqrt{d_x}$ and $\|\partial \Delta(x')\| \leq C$. Therefore,

$$\begin{aligned} |g(x'_1, x'_2, \dots, x'_{d_x-1}, \delta') - x^i_{d_x}| &= |g(x'_1, x'_2, \dots, x'_{d_x-1}, \delta') - g(x^i_1, x^i_2, \dots, x^i_{d_x-1}, \delta)| \\ &\leq \sup_{x' \in \theta(x), \delta' = \Delta(x')} \|\partial g(x_1, x_2, \dots, x_{d_x-1}, \delta')\| \cdot \|(x_1 - x^i_1, x_2 - x^i_2, \dots, x_{d_x-1} - x^i_{d_x-1}, \delta' - \delta)\| \\ &\leq \Lambda(\sqrt{a_1^2(x^i) + \dots + a_{d_x-1}^2(x^i)} + \eta_3), \end{aligned}$$

since $\|(x_1 - x^i_1, \dots, x_{d_x-1} - x^i_{d_x-1}, \delta' - \delta)\| \leq \|(x_1 - x^i_1, \dots, x_{d_x-1} - x^i_{d_x-1})\| + |\delta' - \delta|$, with $\|(x_1 - x^i_1, \dots, x_{d_x-1} - x^i_{d_x-1})\| \leq \sqrt{a_1^2(x^i) + \dots + a_{d_x-1}^2(x^i)}$ and $|\delta' - \delta| < \eta_3$.

We can choose $a_1(x^i) = a_2(x^i) = \dots = a_{d_x-1}(x^i) = \eta_4$ and $a_{d_x}(x^i) = 2(1 + \eta_3)\Lambda(\sqrt{d_x} - 1\eta_4 + \eta_3)$, using η_4 small enough in order to fulfill the following property of θ_i : with η_4 small enough,

$$\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta)) \cap \theta_i \subseteq X_{i1} \times \dots \times X_{i,d_x-1} \times \left(x^i_{d_x} - \frac{a_{d_x}(x^i)}{2(1 + \eta_3)}, x^i_{d_x} + \frac{a_{d_x}(x^i)}{2(1 + \eta_3)} \right),$$

or geometrically, the tube $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta))$ does not intersect θ_i 's faces except at the ones which are *parallel* to the vector $(0, \dots, 0, 1) \in \mathbb{R}^{d_x}$. In such a case, we say that $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta))$ intersects θ_i at the axis x_{d_x} . More generally, for all $i \in \{1, 2, \dots, m\}$, $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta))$ intersects θ_i at axis $i(x^i)$, where x^i is the center of θ_i . This property implies that g is a well-defined injection from $X_{i1} \times \dots \times X_{i,d_x-1} \times B_{\eta_3}(\delta)$ to $X_{i1} \times \dots \times X_{i,d_x}$, for $i \in \{1, \dots, m\}$, which will allow us to perform a change of variable in the equation (E.16). Such a property holds for any $\eta_2 < \eta_3$.

Step 2. Let η_2 be such that $0 < \eta_2 < \eta_3$. We first apply partition of unity to the open cover $\Theta = \{\theta_i\}_{i=1}^m$ of $\widetilde{\mathcal{M}}_\Delta(B_{\eta_2}(\delta)) \cap B_{\eta_2}(\overline{\mathcal{X}})$ of Step 1.

By Lemma D.1, for the finite open cover Θ of the manifold $\widetilde{\mathcal{M}}_\Delta(B_{\eta_2}(\delta)) \cap B_{\eta_2}(\overline{\mathcal{X}})$, we can find a set of \mathcal{C}^∞ partition of unity p_j , $1 \leq j \leq J$ on Θ with the properties given in the lemma.

Our main goal is to compute

$$\partial_\delta F_{\Delta, \mu}(\delta) = \lim_{h \rightarrow 0} \frac{F_{\Delta, \mu}(\delta + h) - F_{\Delta, \mu}(\delta)}{h}.$$

Denote $B_\eta^+(\delta) = [\delta, \delta + \eta]$, for any $\delta \in \mathbb{R}$ and $\eta > 0$. Denote $\mathcal{M}_\Delta(B_\eta^+(\delta)) = \cup_{\delta' \in B_\eta^+(\delta)} \mathcal{M}_\Delta(\delta')$, and $\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) = \cup_{\delta' \in B_\eta^+(\delta)} \widetilde{\mathcal{M}}_\Delta(\delta')$.

For any $0 < \eta < \eta_2$, $\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \subseteq \widetilde{\mathcal{M}}_\Delta(B_\eta(\delta))$. Therefore, the properties (1) to (4) stated in Step 1 are satisfied when we replace $\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta))$ by $\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta))$. Note that,

$$\begin{aligned} F_{\Delta,\mu}(\delta + \eta) - F_{\Delta,\mu}(\delta) &= \int_{x \in \mathcal{X}} 1(\delta \leq \Delta(x) \leq \delta + \eta) \mu'(x) dx \\ &= \int_{\mathcal{M}_\Delta(B_\eta^+(\delta))} \mu'(x) dx = \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta))} \mu'(x) dx = \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \Theta} \mu'(x) dx \\ &= \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap (\cup_{i=1}^m \theta_i)} \mu'(x) \sum_{j=1}^J p_j(x) dx = \sum_{1 \leq i \leq m, 1 \leq j \leq J} \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \theta_i} p_j(x) \mu'(x) dx. \end{aligned} \quad (\text{E.15})$$

This third and fourth equalities hold because $\mu'(x) = 0$ for any $x \in \widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \setminus \mathcal{M}_\Delta(B_\eta^+(\delta))$ and $x \in \widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \setminus \Theta$, respectively.

For any $i \in \{1, 2, \dots, m\}$, without loss of generality, suppose that $\mathcal{M}_\Delta(B_\eta^+(\delta))$ intersects $\theta_i = X_{i1} \times \dots \times X_{id_x}$ at the x_{d_x} axis. Then, $|\partial_{x_{d_x}} \Delta(x)| \geq c/\sqrt{d_x}$ on θ_i , and we can apply the implicit function theorem to show existence of the \mathcal{C}^1 implicit function $g : X_{i1} \times \dots \times X_{i(d_x-1)} \times B_\eta^+(\delta) \rightarrow X_{id_x}$, such that $\Delta(x_1, \dots, x_{d_x-1}, g(x_1, \dots, x_{d_x-1}, \delta')) = \delta'$ for all $(x_1, \dots, x_{d_x-1}, \delta') \in X_{i1} \times \dots \times X_{i(d_x-1)} \times B_\eta^+(\delta)$. Define the injective mapping ψ_{d_x} as:

$$\begin{aligned} \psi_{d_x} : X_{i1} \times \dots \times X_{i(d_x-1)} \times B_\eta^+(\delta) &\rightarrow X_{i1} \times \dots \times X_{i(d_x-1)} \times X_{id_x}, \\ \psi_{d_x}(x_{-d_x}, \delta') &= (x_{-d_x}, g(x_{-d_x}, \delta')) \text{ for } x_{-d_x} := (x_1, x_2, \dots, x_{d_x-1}). \end{aligned}$$

In equation (E.15), we apply a change of variable defined by the map ψ_{d_x} to the (i, j) -th element of the sum:

$$\begin{aligned} \int_{\theta_i \cap \widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta))} p_j(x) \mu'(x) dx &= \int_{X_{i1} \times \dots \times X_{i(d_x-1)} \times B_\delta^+(\eta)} (p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x}) |\det(D\psi_{d_x})| d\delta' dx_{-d_x} \\ &= \int_{X_{i1} \times \dots \times X_{i(d_x-1)}} \int_{B_\delta^+(\eta)} \frac{(p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x})}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}|} d\delta' dx_{-d_x} \\ &= \eta \int_{X_{i1} \times \dots \times X_{i(d_x-1)}} \frac{(p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x})}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}|} dx_{-d_x} + o(\eta). \end{aligned} \quad (\text{E.16})$$

The second equality follows because

$$D\psi_{d_x}(x_{-d}, \delta) = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \partial_\delta g(x_{-d_x}, \delta) \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 1/\partial_{x_{d_x}} \Delta(\tilde{x}) \end{bmatrix},$$

where $\tilde{x} = \psi_{d_x}(x_{-d}, \delta)$.

The last equality follows as $\eta \rightarrow 0$, because by the uniform continuity of

$$(x_{-d_x}, \delta') \mapsto (p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x}) / |\partial_{x_{d_x}} \Delta \circ \psi_{d_x}| \Big|_{(x_{-d_x}, \delta')}$$

over $(x_{-d_x}, \delta') \in X_{i_1} \times \dots \times X_{i_{(d_x-1)}} \times B_\eta^+(\delta)$. In (E.16), the last component of ψ_{d_x} is fixed to be δ without being specified for simplicity. We will maintain this convention in the rest of the proof whenever the variable of integration is x_{-d_x} (excluding x_{d_x}).

Next, we write the first term of (E.16) as an integral on a manifold, which is

$$\eta \int_{X_{i_1} \times \dots \times X_{i_{(d_x-1)}}} \frac{(p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x})}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}|} dx_{-d_x} = \eta \int_{\widetilde{\mathcal{M}}_\Delta(\delta) \cap \theta_i} \frac{p_j(x) \mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol}. \quad (\text{E.17})$$

Summing up over i and j in (E.16) and using Definition 5.5,

$$\sum_{1 \leq i \leq m, 1 \leq j \leq J} \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \theta_i} p_j(x) \mu'(x) dx = \eta \int_{\widetilde{\mathcal{M}}_\Delta(\delta) \cap \Theta} \frac{\mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol} + o(\eta). \quad (\text{E.18})$$

Let us explain (E.17). Equation (E.17) is calculated using the following fact: The mapping $\alpha : X_{i_1} \times \dots \times X_{i_{d_x-1}} \rightarrow X_{i_1} \times \dots \times X_{i_{d_x}}$ such that $\alpha(x_1, \dots, x_{d_x-1}) = (x_1, \dots, x_{d_x-1}, g(x_1, \dots, x_{d_x-1}, \delta))$ has Jacobian matrix

$$D\alpha^\top(x_{-d_x}) = \begin{bmatrix} 1 & 0 & \dots & 0 & \partial_{x_1} g(x_{-d_x}) \\ 0 & 1 & \dots & 0 & \partial_{x_2} g(x_{-d_x}) \\ \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 1 & \partial_{x_{d_x-1}} g(x_{-d_x}) \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & 0 & (\partial_{x_1} \Delta / \partial_{x_{d_x}} \Delta)(\tilde{x}) \\ 0 & 1 & \dots & 0 & (\partial_{x_2} \Delta / \partial_{x_{d_x}} \Delta)(\tilde{x}) \\ \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 1 & (\partial_{x_{d_x-1}} \Delta / \partial_{x_{d_x}} \Delta)(\tilde{x}) \end{bmatrix},$$

where $\tilde{x} = (x_1, \dots, x_{d_x-1}, g(x_1, \dots, x_{d_x-1}, \delta))$. The volume of $D\alpha$ is $\text{Vol}(D\alpha) = \sqrt{\det(D\alpha^\top D\alpha)}$, where $D\alpha^\top D\alpha = I_{d_x-1} + \partial g \partial g^\top$. By the Matrix Determinant Lemma,

$$\text{Vol}(D\alpha)(x_{-d_x}) = \sqrt{1 + \partial g^\top \partial g} = \|\partial \Delta\| / \|\partial_{x_{d_x}} \Delta\| \Big|_{x=\tilde{x}}.$$

Hence, the left hand side of equation (E.17) is:

$$\eta \int_{X_{i_1} \times \dots \times X_{i_{(d_x-1)}}} \frac{(p_j \circ \psi_{d_x}) \cdot (\mu' \circ \psi_{d_x})}{\|\partial \Delta \circ \psi_{d_x}\|} \text{Vol}(D\alpha) dx_{-d_x},$$

and it can be further re-expressed as the right side of (E.17) using Definition 5.4.

By equations (E.15) and (E.18),

$$\frac{F_{\Delta, \mu}(\delta + \eta) - F_{\Delta, \mu}(\delta)}{\eta} = \int_{\mathcal{M}_\Delta(\delta)} \frac{\mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol} + o(1), \quad (\text{E.19})$$

where we use that $\mu'(x) = 0$ for all $x \in \widetilde{\mathcal{M}}_\Delta(\delta) \setminus \mathcal{M}_\Delta(\delta)$. Similarly, we can show that

$$\frac{F_{\Delta, \mu}(\delta) - F_{\Delta, \mu}(\delta - \eta)}{\eta} = \int_{\mathcal{M}_\Delta(\delta)} \frac{\mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol} + o(1).$$

Thus, we conclude that $F_{\Delta, \mu}(\delta)$ is differentiable at $\delta \in \mathcal{D}$ with derivative

$$f_{\Delta, \mu}(\delta) := \partial_\delta F_{\Delta, \mu}(\delta) = \int_{\mathcal{M}_\Delta(\delta)} \frac{\mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol}.$$

Finally, if $\mathcal{M}_\Delta(\delta)$ has multiple branches but a finite number of them, we can repeat Step 1 and 2 in the proof above for each individual branch. Since the number of connected branches is finite, the remainders in equation (E.19) converge to 0 uniformly. Thus, adding up the results for all connected branches in equation (E.19), the statements of Lemma A.1 hold. \square

E.2. Proof of Lemma A.2. We use the following results in the proof of Lemma A.2.

Lemma E.3 (Continuity). *Let f be a measurable function defined on $B_\eta(\mathcal{X}) \subset B(\mathcal{X})$ which vanishes outside \mathcal{X} , where $\eta > 0$ is a constant. Let δ be a regular value of Δ on $\overline{\mathcal{X}}$. Suppose f is continuous on $\widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\overline{\mathcal{X}})$ for any $\delta \in \mathcal{D}$ and some small η_1 such that $0 < \eta_1 < \eta$. Then, $\delta \mapsto \int_{\mathcal{M}_\Delta(\delta)} f d\text{Vol}$ is continuous on \mathcal{D} .*

Proof. First, we follow Step 1 in the Proof of Lemma A.1. Suppose we have a set of open rectangles $\Theta = \{\theta_1, \dots, \theta_m\}$ such that $\widetilde{\mathcal{M}}_\Delta(B_{\eta_2}(\delta)) \cap B_{\eta_2}(\overline{\mathcal{X}}) \subset \cup_{i=1}^m \theta_i \subset \overline{\cup_{i=1}^m \theta_i} \subset \widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\overline{\mathcal{X}})$ for any $\eta_2 < \eta_3$, where η_3 is a small enough positive number, $\eta_3 < \eta_1$. Moreover, let η_3 be small enough such that all $\delta' \in B_{\eta_3}(\delta)$ are regular values. By compactness of $\overline{\cup_{i=1}^m \theta_i}$, f is bounded and uniformly continuous on $\cup_{i=1}^m \theta_i$.

By construction, θ_i , $i = 1, 2, \dots, m$, satisfies that $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3})$ intersects θ_i at axis $i(\theta_i)$, for any $\eta_2 < \eta_3$.

Then, following Step 2 in the Proof of Lemma A.1, there exists a set of \mathcal{C}^∞ partition of unity functions $x \mapsto p_j(x)$ of Θ , $j = 1, 2, \dots, J$.

Then, for any $\delta' \in B_{\eta_3}(\delta)$, by the definition of partition of unity,

$$\int_{\mathcal{M}_\Delta(\delta')} f d\text{Vol} = \sum_{1 \leq i \leq m, 1 \leq j \leq J} \int_{\widetilde{\mathcal{M}}_\Delta(\delta') \cap \theta_i} p_j(x) f(x) d\text{Vol}. \quad (\text{E.20})$$

The equation (E.20) holds since $f(x) = 0$ for all $x \notin \mathcal{X}$.

To show that $\int_{\mathcal{M}_\Delta(\delta')} f d\text{Vol}$ converges to $\int_{\mathcal{M}_\Delta(\delta)} f d\text{Vol}$ as δ' converges to δ , it suffices to show that $\int_{\widetilde{\mathcal{M}}_\Delta(\delta') \cap \theta_i} p_j(x) f(x) d\text{Vol}$ converges to $\int_{\widetilde{\mathcal{M}}_\Delta(\delta) \cap \theta_i} p_j(x) f(x) d\text{Vol}$ as δ' converges to δ , for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, J$.

Without loss of generality, assume that $\widetilde{\mathcal{M}}_\Delta(B_{\eta_3}(\delta))$ intersects θ_i at axis $i(\theta_i) = d_x$. Then, there exists constants $c > 0$ and $C > 0$ such that $\partial_{x_{d_x}} \Delta(x) > c$ and $\|\partial \Delta(x)\| < C$ for all $x \in \theta_i$, $i = 1, 2, \dots, m$.

We can apply the implicit function theorem to establish existence of the \mathcal{C}^1 function $g : X_{i_1} \times \dots \times X_{i_{(d_x-1)}} \times B_\eta^+(\delta) \rightarrow X_{i_{d_x}}$, such that $\Delta(x_1, \dots, x_{d_x-1}, g(x_1, \dots, x_{d_x-1}, \delta')) = \delta'$ for all $(x_1, \dots, x_{d_x-1}, \delta') \in X_{i_1} \times \dots \times X_{i_{(d_x-1)}} \times B_\eta(\delta)$. Define the one-to-one mapping ψ_{d_x} as:

$$\psi_{d_x} : X_{i_1} \times \dots \times X_{i_{(d_x-1)}} \times B_\eta^+(\delta) \rightarrow X_{i_1} \times \dots \times X_{i_{(d_x-1)}} \times X_{i_{(d_x)}},$$

where $\psi_{d_x}(x_{-d_x}, \delta') = (x_{-d_x}, g(x_{-d_x}, \delta'))$ for $x_{-d_x} := (x_1, x_2, \dots, x_{d_x-1})$. Note that ψ_{d_x} and g are both \mathcal{C}^1 functions.

For any δ' such that $|\delta' - \delta| < \eta_3$, by the change of variables we have:

$$\int_{\widetilde{\mathcal{M}}_{\Delta}(\delta') \cap \theta_i} p_j(x) f(x) d\text{Vol} = \int_{X_1 \times X_2 \times \dots \times X_{d_x-1}} (p_j f) \circ \psi_{d_x}(x_{-d_x}, \delta') \frac{\|\partial \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')\|}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} dx_{-d_x}. \quad (\text{E.21})$$

Since $|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')| = |\partial_{x_{d_x}} \Delta|_{x=\psi_{d_x}(x_1, \dots, x_{d_x-1}, \delta')} > c$ for all $\delta' \in B_{\eta_3}(\delta)$ and $x_{-d_x} \in X_1 \times X_2 \times \dots \times X_{d_x-1}$ and $p_j, f, \partial \Delta$ and $\partial_{x_{d_x}} \Delta$ are uniformly continuous functions on $\widetilde{\mathcal{M}}_{\Delta}(B_{\eta_3}(\delta)) \cap B_i$, conclude that the map

$$(p_j f) \circ \psi_{d_x} \frac{\|\partial \Delta \circ \psi_{d_x}\|}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}|} \text{ is uniformly continuous on } X_1 \times \dots \times X_{d_x-1} \times B_{\eta_3}(\delta).$$

Since $X_1 \times \dots \times X_{d_x-1}$ and is bounded, it immediately follows that $\delta' \mapsto \int_{\widetilde{\mathcal{M}}_{\Delta}(\delta') \cap \theta_i} p_j(x) f(x) d\text{Vol}$ is continuous at $\delta' = \delta$, and hence

$$\delta' \mapsto \int_{\mathcal{M}_{\Delta}(\delta')} f d\text{Vol} = \sum_{1 \leq i \leq m, 1 \leq j \leq J} \int_{\widetilde{\mathcal{M}}_{\Delta}(\delta') \cap \theta_i} p_j(x) f(x) d\text{Vol}$$

is continuous at $\delta' = \delta$.

This argument applies to every $\delta \in \mathcal{D}$, and by compactness of \mathcal{D} the continuity claim extends to the entire \mathcal{D} . \square

Lemma E.4 (Hadamard differentiability of $\Delta \mapsto F_{\Delta, \mu}$ and $\Delta \mapsto \Delta_{\mu}^*$). *Suppose that S.1-S.2 hold. Then:*

(a) *The map $F_{\Delta, \mu}(\delta) : \mathbb{F} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $\delta \in \mathcal{D}$ at Δ tangentially to \mathbb{F}_0 , with the derivative map $\partial_{\Delta} F_{\Delta, \mu}(\delta) : \mathbb{F}_0 \rightarrow \mathbb{R}$ defined by*

$$G \mapsto \partial_{\Delta} F_{\Delta, \mu}(\delta)[G] := - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{G(x) \mu'(x)}{\|\partial \Delta(x)\|} d\text{Vol}.$$

(b) *The map $\Delta_{\mu}^*(u) : \mathbb{F} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $u \in \mathcal{U}$ at Δ tangentially to \mathbb{F}_0 , with the derivative map $\partial_{\Delta} \Delta_{\mu}^*(u) : \mathbb{F}_0 \rightarrow \mathbb{R}$ defined by:*

$$G \mapsto \partial_{\Delta} \Delta_{\mu}^*(u)[G] := - \frac{\partial_{\Delta} F_{\Delta, \mu}(\Delta_{\mu}^*(u))[G]}{f_{\Delta, \mu}(\Delta_{\mu}^*(u))}.$$

Proof of Lemma E.4. To shows statement (a), for any $G_n \rightarrow G \in \mathbb{F}_0$ under sup-norm such that $\Delta + t_n G_n \in \mathbb{F}$, and $t_n \rightarrow 0$, we consider

$$\frac{F_{\Delta + t_n G_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)}{t_n}.$$

By assumption, any function $G \in \mathbb{F}_0$ is bounded and uniformly continuous on $B(\mathcal{X})$. Hence, G_n is uniformly bounded for $n \geq N$, since $G_n \rightarrow G$ in sup-norm.

For any $\delta \in \mathcal{D}$ we consider a procedure similar to Lemma A.1. We use the same notation as in Step 1 of the proof of Lemma A.1. Suppose for $\eta_1 > 0$ small enough, we have a rectangle cover $\Theta = \cup_{i=1}^m \theta_i \subseteq B(\mathcal{X})$ of $\widetilde{\mathcal{M}}_\Delta(B_{\eta_1}(\delta)) \cap B_{\eta_1}(\mathcal{X})$ such that for all $\eta < \eta_1$, $\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta))$ intersects each θ_i at some axis $i(\theta_i)$, $1 \leq i \leq m$. As before, there is a partition of unity $\{p_j\}_{j=1}^J$ on the cover sets $\Theta = \{\theta_i\}_{i=1}^m$. As in the proof of Lemma A.1, we can rewrite

$$\begin{aligned} & \frac{\int_{\mathcal{X}} [1\{\Delta(x) + t_n G_n(x) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x) dx}{t_n} \\ &= \sum_{1 \leq i \leq m, 1 \leq j \leq J} \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \theta_i} p_j(x) \frac{[1\{\Delta(x) + t_n G_n(x) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x)}{t_n} dx. \end{aligned}$$

Then, for any fixed positive number $|\zeta|$, there exist N large enough such that $\sup_{x \in B(\mathcal{X}), n \geq N} |G_n - G| < |\zeta|$. Moreover, for any $x \in B(\mathcal{X})$, and large enough n ,

$$1\{\Delta(x) + t_n G_n(x) \leq \delta\} \leq 1\{\Delta(x) + t_n(G(x) - \zeta) \leq \delta\}.$$

As in Step 2 of the proof of Lemma A.1, suppose $\theta_i = X_{i1} \times \dots \times X_{id_x}$ intersects $\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta))$ at $i(\theta_i) = x_{d_x}$. Define the parametrization

$$\psi_{d_x} : X_{i1} \times \dots \times X_{i,d_x-1} \times B_\eta(\delta) \mapsto \theta_i,$$

$$\psi_{d_x}(x_{-d_x}, \delta') = (x_{-d_x}, g(x_{-d_x}, \delta')),$$

where $g(x_{-d_x}, \delta')$ is the implicit function derived from equation $\Delta(x) = \delta'$, for any $\delta' \in B_\eta(\delta)$. Therefore, for large enough n ,

$$\begin{aligned} & \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \theta_i} p_j(x) \frac{[1\{\Delta(x) + t_n G_n(x) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x)}{t_n} dx \\ & \leq \frac{\int_{\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta)) \cap \theta_i} [1\{\Delta(x) + t_n(G(x) - \zeta) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x) dx}{t_n}. \end{aligned}$$

Next, by a change of variables $\psi_{d_x}^{-1}$ from θ_i to $X_{i1} \times \dots \times X_{i,d_x-1} \times B_\eta(\delta)$,

$$\begin{aligned} & \int_{\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta)) \cap \theta_i} p_j(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - t_n(G(x) - \zeta)\} \mu'(x)}{t_n} dx \\ &= \int_{X_{i1} \times \dots \times X_{i,d_x-1}} \int_{B_\eta(\delta)} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \frac{1\{\delta \leq \delta' \leq \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta)\}}{t_n} d\delta' dx_{-d_x} \\ &= \int_{X_{i1} \times \dots \times X_{i,d_x-1}} \int_{B_\eta(\delta) \cap [\delta, \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta)]} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')| t_n} d\delta' dx_{-d_x} \\ & \leq - \int_{X_{i1} \times \dots \times X_{i,d_x-1}} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta) \end{aligned}$$

$$\begin{aligned}
&= - \int_{\theta_i \cap \widetilde{\mathcal{M}}_\Delta(\delta)} p_j(x) \mu'(x) \frac{G(x) - \zeta}{\|\partial\Delta(x)\|} d\text{Vol} + o(\eta) \\
&= - \int_{\theta_i \cap \mathcal{M}_\Delta(\delta)} p_j(x) \mu'(x) \frac{G(x) - \zeta}{\|\partial\Delta(x)\|} d\text{Vol} + o(\eta),
\end{aligned}$$

where the inequality in the above equation holds by continuity of $(x_{-d_x}, \delta') \mapsto (p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta') / |\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|$. More specifically, fixing $\eta > 0$ and x_{-d_x} , for $t_n \rightarrow 0$,

$$B_\eta(\delta) \cap [\delta, \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta)] = [\delta, \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta)]$$

and

$$\frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \rightarrow \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|}$$

as $\delta' \rightarrow \delta$. The last equality above holds because $\mu'(x) = 0$ for all $x \in \widetilde{\mathcal{M}}_\Delta(\delta) \setminus \mathcal{M}_\Delta(\delta)$.

Since m and J are fixed for any $n \geq N$, and $|G \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta|$ is bounded by some absolute constant, $\sum_j p_j(x) = 1$ and $p_j(x) \geq 0$, we can let $\zeta \rightarrow 0$ to conclude that:

$$\lim_{n \rightarrow \infty} \frac{F_{\Delta+t_n G_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} \leq \sum_{i=1}^m \sum_{j=1}^J - \int_{\theta_i \cap \mathcal{M}_\Delta(\delta)} p_j(x) \mu'(x) \frac{G(x)}{\|\partial\Delta(x)\|} d\text{Vol}.$$

The right side is given by:

$$- \int_{\mathcal{M}_\Delta(\delta)} \frac{\mu'(x) G(x)}{\|\partial\Delta(x)\|} d\text{Vol}.$$

On the other hand,

$$1(\Delta(x) + t_n G_n(x) \leq \delta) \geq 1(\Delta(x) + t_n(G(x) + \zeta) \leq \delta)$$

for some $\zeta > 0$. So,

$$\begin{aligned}
&\int_{\widetilde{\mathcal{M}}_\Delta(B_\eta^+(\delta)) \cap \theta_i} p_j(x) \frac{[1\{\Delta(x) + t_n G_n(x) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x)}{t_n} dx \\
&\geq \frac{\int_{\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta)) \cap \theta_i} [1\{\Delta(x) + t_n(G(x) + \zeta) \leq \delta\} - 1\{\Delta(x) \leq \delta\}] \mu'(x) dx}{t_n}.
\end{aligned}$$

And, by a change of variables $\psi_{d_x}^{-1}$ from θ_i to $X_{i1} \times \dots \times X_{i, d_x-1} \times B_\eta(\delta)$,

$$\begin{aligned}
&\int_{\widetilde{\mathcal{M}}_\Delta(B_\eta(\delta)) \cap \theta_i} p_j(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - t_n(G(x) + \zeta)\} \mu'(x)}{t_n} dx \\
&= \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \int_{B_\eta(\delta)} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \frac{1\{\delta \leq \delta' \leq \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) + \zeta)\}}{t_n} d\delta' dx_{-d_x} \\
&= \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \int_{B_\eta(\delta) \cap [\delta, \delta - t_n(G \circ \psi_{d_x}(x_{-d_x}, \delta) + \zeta)]} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')| t_n} d\delta' dx_{-d_x} \\
&\geq - \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \frac{(p_j \cdot \mu') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (G \circ \psi_{d_x}(x_{-d_x}, \delta) + \zeta) dx_{-d_x} - o(\eta)
\end{aligned}$$

$$\begin{aligned}
&= - \int_{\theta_i \cap \widetilde{\mathcal{M}}_{\Delta}(\delta)} p_j(x) \mu'(x) \frac{G(x) + \zeta}{\|\partial\Delta(x)\|} d\text{Vol} - o(\eta) \\
&= - \int_{\theta_i \cap \mathcal{M}_{\Delta}(\delta)} p_j(x) \mu'(x) \frac{G(x) + \zeta}{\|\partial\Delta(x)\|} d\text{Vol} - o(\eta).
\end{aligned}$$

Let $\zeta \rightarrow 0$ and $\eta \rightarrow 0$, it follows that

$$\lim_{n \rightarrow \infty} \frac{F_{\Delta + t_n G_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} \geq - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{\mu'(x) G(x)}{\|\partial\Delta(x)\|} d\text{Vol}.$$

Combining the two inequalities, we conclude that $F_{\Delta, \mu}(\delta)$ is Hadamard-differentiable at Δ tangentially to \mathbb{F}_0 with derivative

$$\partial_{\Delta} F_{\Delta, \mu}(\delta)[G] = - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{\mu'(x) G(x)}{\|\partial\Delta(x)\|} d\text{Vol}.$$

To show that the result holds uniformly in $\delta \in \mathcal{D}$, we use the equivalence between uniform convergence and continuous convergence (e.g., Resnick (1987, p.2)). Take a sequence δ_n in \mathcal{D} that converges to $\delta \in \mathcal{D}$. Then, the preceding argument applies to this sequence and $\partial_{\Delta} F_{\Delta, \mu}(\delta_n)[G] \rightarrow \partial_{\Delta} F_{\Delta, \mu}(\delta)[G]$ by uniform continuity of $\delta \mapsto \partial_{\Delta} F_{\Delta, \mu}(\delta)[G]$ on \mathcal{D} , which holds by Lemma E.3 because G , μ' , and $\|\partial\Delta\|$ are continuous on $\overline{\mathcal{X}}$ and \mathcal{D} excludes neighborhoods of the critical values of Δ in $\overline{\mathcal{X}}$.

To show statement (b), note that by statement (a), Hadamard differentiability of the quantile map, see e.g., Lemma 3.9.20 in van der Vaart and Wellner (1996), and the chain rule for Hadamard differentiation, the inverse map $\Delta_{\mu}^*(u)$ is Hadamard differentiable at Δ tangentially to \mathbb{F}_0 with the derivative map

$$\partial_{\Delta} \Delta_{\mu}^*(u)[G] = - \left. \frac{\partial_{\Delta} F_{\Delta, \mu}(\delta)[G]}{\partial_{\delta} F_{\Delta, \mu}(\delta)} \right|_{\delta = \Delta_{\mu}^*(u)} = \frac{\partial_{\Delta} F_{\Delta, \mu}(\Delta_{\mu}^*(u))[G]}{f_{\Delta, \mu}(\Delta_{\mu}^*(u))},$$

uniformly in the index $u \in \mathcal{U} = \{u \in (0, 1) : \Delta_{\mu}^*(u) \in \mathcal{D}, f_{\Delta, \mu}(\Delta_{\mu}^*(u)) \geq \varepsilon\}$. \square

Proof of Lemma A.2 . To show Statement (a), Consider $t_n \rightarrow 0$ and $(G_n, H_n) \rightarrow (G, H) \in \mathbb{D}_0 := \mathbb{F}_0 \times \mathbb{H}$ as $n \rightarrow \infty$, such that $(\Delta + t_n G_n, \mu + t_n H_n) \in \mathbb{D}$. Let $\Delta_n := \Delta + t_n G_n$ and $\mu_n := \mu + t_n H_n$. Then, we can decompose

$$F_{\Delta_n, \mu_n}(\delta) - F_{\Delta, \mu}(\delta) = [F_{\Delta_n, \mu_n}(\delta) - F_{\Delta_n, \mu}(\delta)] + [F_{\Delta_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)].$$

By Lemma E.4,

$$\frac{F_{\Delta_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} = - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{G(x) \mu'(x)}{\|\partial\Delta(x)\|} d\text{Vol} + o(1).$$

Let $g_{\Delta, \delta} := 1(\Delta(x) \leq \delta)$. By definition of $F_{\Delta_n, \mu_n}(\delta)$,

$$\frac{F_{\Delta_n, \mu_n}(\delta) - F_{\Delta_n, \mu}(\delta)}{t_n} = H_n(g_{\Delta_n, \delta}).$$

Note that

$$H_n(g_{\Delta_n, \delta}) - H(g_{\Delta, \delta}) = [H_n(g_{\Delta_n, \delta}) - H_n(g_{\Delta, \delta})] + [H_n - H](g_{\Delta, \delta}).$$

The second term goes to 0 by the assumption $H_n \rightarrow H$ in \mathbb{H} . For the first term, we further decompose

$$|H_n(g_{\Delta_n, \delta}) - H_n(g_{\Delta, \delta})| \leq |H_n(g_{\Delta_n, \delta}) - H(g_{\Delta_n, \delta})| + |H_n(g_{\Delta, \delta}) - H(g_{\Delta, \delta})| + |H(g_{\Delta_n, \delta}) - H(g_{\Delta, \delta})|.$$

The first two terms go to 0 by $\|H_n - H\|_{\mathcal{G}} \rightarrow 0$. Moreover, $H(g_{\Delta_n, \delta}) \rightarrow H(g_{\Delta, \delta})$ because $g_{\Delta_n, \delta}(X) = 1(\Delta_n(X) \leq \delta) \rightarrow g_{\Delta, \delta}(X) = 1(\Delta(X) \leq \delta)$ in the $L^2(\mu)$ norm, since $\Delta_n \rightarrow \Delta$ in the sup norm and $\Delta(X)$ has an absolutely continuous distribution, and since we require the operator H to be continuous under the $L^2(\mu)$ norm.

We conclude that for any $\delta \in \mathcal{D}$,

$$\frac{F_{\Delta_n, \mu_n}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} \rightarrow - \int_{\mathcal{M}_{\Delta}(\delta)} \frac{G(x)\mu'(x)}{\|\partial\Delta(x)\|} d\text{Vol} + H(g_{\Delta, \delta}) = \partial_{\Delta, \mu} F_{\Delta, \mu}(\delta)[G, H].$$

By an argument similar to the proof of Lemma E.4, it can be shown that the convergence is uniform in $\delta \in \mathcal{D}$.

Statement (b) follows by statement (a) and the Hadamard differentiability of the quantile map uniformly in the quantile index, see, e.g., Lemma 3.9.20 in van der Vaart and Wellner (1996). \square

E.3. Proof of Lemma A.3. We will denote the functions in the classes \mathcal{F}_M and \mathcal{F}_I by $\varphi_t(x)$ whenever we want to distinguish $x = (x_1, \dots, x_{d_x})$, the argument of the function, from $t := (t_1, \dots, t_{d_z})$, the index of the function in the class. Otherwise, we will use $\varphi(x)$. To analyze $\Lambda_{\Delta, \mu, \delta}^-$ it is convenient to introduce the operator $\Upsilon_{\Delta, \mu, \delta} : \mathbb{D} \rightarrow \mathbb{R}$ defined by

$$\Upsilon_{\Delta, \mu, \delta}(\varphi) := \int \varphi(x) 1\{\Delta(x) \leq \delta\} d\mu(x),$$

since $\Lambda_{\Delta, \mu, \delta}^-(\varphi) = \Upsilon_{\Delta, \mu, \delta}(\varphi) / \Upsilon_{\Delta, \mu, \delta}(1)$.

Let $\widetilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)) := \cup_{\delta' \in B_{\eta}(\delta)} \widetilde{\mathcal{M}}_{\Delta}(\delta')$, where $\widetilde{\mathcal{M}}_{\Delta}(\delta) := \{x \in B(\mathcal{X}) : \Delta(x) = \delta\}$ and $B_{\eta}(\delta) := (\delta - \eta, \delta + \eta)$ for any $\delta \in \mathcal{V}$ and $\eta > 0$. When $\varphi_t \in \mathcal{F}_I$ we make the following technical assumption to deal with the discontinuity of the indicator functions:

AS.1. Define the set $\widetilde{\mathcal{Z}}_{k, \eta}(\delta, t_k) := \{x_{-k} : (x_k, x_{-k}) \in \widetilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)), x_k = t_k\}$ for any $\eta > 0$, $\delta \in \mathcal{V}$, $k = 1, 2, \dots, d_x$, and $t_k \in \mathbb{R}$. Then, for any $\epsilon > 0$, there exist $\eta_0 > 0$ such that for any $\eta < \eta_0$, $\int_{\widetilde{\mathcal{Z}}_{k, \eta}(\delta, t_k)} d\mu(x_{-k}) \leq \epsilon$ holds uniformly over all $\delta \in \mathcal{V}$, $t_k \in \mathbb{R}$ and $k = 1, 2, \dots, d_x$.

The next result shows that $(\Delta, \mu, \delta) \mapsto \Upsilon_{\Delta, \mu, \delta}$ is Hadamard differentiable.

Lemma E.5 (Hadamard differentiability of $(\Delta, \mu, \delta) \mapsto \Upsilon_{\Delta, \mu, \delta}$). *Assume that Assumptions S.1 and S.2 hold and $\delta \in \mathcal{D}$. Then,*

(a) *The map $\Upsilon_{\Delta, \mu, \delta}(\varphi) : \mathbb{D} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $\varphi \in \mathcal{F}_M$ at (Δ, μ, δ) tangentially to \mathbb{D}_0 .*

(b) If in addition Assumption AS.1 holds, the map $\Upsilon_{\Delta,\mu,\delta}(\varphi) : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ is Hadamard-differentiable uniformly in $\varphi \in \mathcal{F}_I$ at (Δ, μ, δ) tangentially to $\tilde{\mathbb{D}}_0$.

(c) The derivative map $\partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(\varphi) : \tilde{\mathbb{D}} \rightarrow \mathbb{R}$ is defined by:

$$(G, H, K) \mapsto \partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(\varphi)[G, H, K] := \int_{\mathcal{M}_{\Delta}(\delta)} \varphi(x) \frac{K - G(x)}{\|\partial\Delta(x)\|} d\text{Vol} + H(h_{\Delta,\delta,\varphi}),$$

where $h_{\Delta,\delta,\varphi} := \varphi(x)1\{\Delta(x) \leq \delta\}$.

Proof of Lemma E.5. Statements (a) and (b) follow by similar arguments. For brevity, we focus on the proof of Statement (b) and mention the changes needed for the proof of Statement (a), which is simpler.

To show Statement (b), consider $s_n \rightarrow 0$ and $(G_n, H_n, K_n) \rightarrow (G, H, K) \in \tilde{\mathbb{D}}_0$ as $n \rightarrow \infty$, such that $(\Delta + s_n G_n, \mu + s_n H_n, \delta + s_n K_n) \in \tilde{\mathbb{D}}$. Let $\Delta_n := \Delta + s_n G_n$, $\mu_n := \mu + s_n H_n$, and $\delta_n := \delta + s_n K_n$. Then, we can decompose

$$\Upsilon_{\Delta_n, \mu_n, \delta_n}(\varphi) - \Upsilon_{\Delta, \mu, \delta}(\varphi) = [\Upsilon_{\Delta_n, \mu_n, \delta_n}(\varphi) - \Upsilon_{\Delta_n, \mu, \delta_n}(\varphi)] + [\Upsilon_{\Delta_n, \mu, \delta_n}(\varphi) - \Upsilon_{\Delta, \mu, \delta}(\varphi)]. \quad (\text{E.22})$$

The first term of (E.22) satisfies

$$\frac{\Upsilon_{\Delta_n, \mu_n, \delta_n}(\varphi) - \Upsilon_{\Delta_n, \mu, \delta_n}(\varphi)}{s_n} = H_n(h_{\Delta_n, \delta_n, \varphi}) = H(h_{\Delta, \delta, \varphi}) + o(1).$$

The first equality follows from linearity of $\mu \mapsto \Upsilon_{\Delta_n, \mu, \delta_n}(\varphi)$ and $h_{\Delta_n, \delta_n, \varphi} = \varphi(x)1\{\Delta_n(x) \leq \delta_n\}$. To show the second equality note that

$$H_n(h_{\Delta_n, \delta_n, \varphi}) - H(h_{\Delta, \delta, \varphi}) = H_n(h_{\Delta_n, \delta_n, \varphi}) - H_n(h_{\Delta, \delta, \varphi}) + [H_n - H](h_{\Delta, \delta, \varphi}),$$

where the second term goes to zero by the assumption $H_n \rightarrow H$ in $\tilde{\mathbb{H}}$. For the first term, we further decompose

$$\begin{aligned} |H_n(h_{\Delta_n, \delta_n, \varphi}) - H_n(h_{\Delta, \delta, \varphi})| &\leq |H_n(h_{\Delta_n, \delta_n, \varphi}) - H(h_{\Delta_n, \delta_n, \varphi})| \\ &\quad + |H_n(h_{\Delta, \delta, \varphi}) - H(h_{\Delta, \delta, \varphi})| + |H(h_{\Delta_n, \delta_n, \varphi}) - H(h_{\Delta, \delta, \varphi})|. \end{aligned}$$

By definition of the space $\tilde{\mathbb{H}}$, the first two terms go to 0 by $\|H_n - H\|_{\tilde{\mathcal{G}}} \rightarrow 0$.

Moreover, $H(h_{\Delta_n, \delta_n, \varphi}) \rightarrow H(h_{\Delta, \delta, \varphi})$ because

$$h_{\Delta_n, \delta_n, \varphi}(X) = \varphi(X)1(\Delta_n(X) \leq \delta_n) \rightarrow h_{\Delta, \delta, \varphi}(X) = \varphi(X)1(\Delta(X) \leq \delta)$$

in the $L^2(\mu)$ norm, since $\Delta_n \rightarrow \Delta$ in the sup norm and $\Delta(X)$ has an absolutely continuous distribution, and since we require the operator H to be continuous under the $L^2(\mu)$ norm.

Next we show that the second term of (E.22) satisfies

$$\frac{\Upsilon_{\Delta_n, \mu, \delta_n}(\varphi) - \Upsilon_{\Delta, \mu, \delta}(\varphi)}{s_n} = \int_{\mathcal{M}_{\Delta}(\delta)} \varphi(x) \frac{K - G(x)}{\|\partial\Delta(x)\|} \mu'(x) d\text{Vol} + o(1).$$

The proof follows the same steps as the proof of Lemma E.4 after noticing that we can write

$$\Upsilon_{\Delta_n, \mu, \delta_n}(\varphi) = \Upsilon_{\tilde{\Delta}_n, \mu, \delta}(\varphi),$$

where $\tilde{\Delta}_n = \Delta + s_n \tilde{G}_n$ with $\tilde{G}_n = G_n - K_n$, and replacing $\mu'(x)$ by $\tilde{\mu}'(x) = \varphi(x)\mu'(x)$.

Specifically, following the notation in the proof of Lemma A.1,

$$\Upsilon_{\tilde{\Delta}_n, \mu, \delta}(\varphi) = \sum_{i=1}^m \sum_{j=1}^J \int_{\tilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)) \cap \theta_i} p_j(x) \varphi(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - s_n \tilde{G}_n(x)\}}{s_n} \tilde{\mu}'(x) dx.$$

Without loss of generality, assume that θ_i intersects with $\tilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta))$ at the axis $x_{k_i} = x_{d_x}$. When $\varphi(x) \in \mathcal{F}_I$, each component in the above summation satisfies:

$$\begin{aligned} & \int_{\tilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)) \cap \theta_i} p_j(x) \varphi(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - s_n \tilde{G}_n(x)\} \tilde{\mu}'(x)}{s_n} dx \\ &= \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \int_{B_{\eta}(\delta)} \frac{(p_j \cdot \varphi \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \times \frac{1\{\delta \leq \delta' \leq \delta - s_n \tilde{G}_n \circ \psi_{d_x}(x_{-d_x}, \delta')\}}{s_n} d\delta' dx_{-d_x} \\ &= \int_{\tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x})} \int_{B_{\eta}(\delta)} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \times \frac{1\{\delta \leq \delta' \leq \delta - s_n \tilde{G}_n \circ \psi_{d_x}(x_{-d_x}, \delta')\}}{s_n} d\delta' dx_{-d_x} \\ &+ \int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \int_{B_{\eta}(\delta)} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \times \frac{1\{\delta \leq \delta' \leq \delta - s_n \tilde{G}_n \circ \psi_{d_x}(x_{-d_x}, \delta')\}}{s_n} d\delta' dx_{-d_x}, \end{aligned}$$

where $\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x}) := [X_{i1} \times \dots \times X_{i, d_x-1}] \cap \tilde{\mathcal{Z}}_{d_x, \eta}(\delta, t_{d_x})$ and $\tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x}) := X_{i1} \times \dots \times X_{i, d_x-1} \setminus \tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})$. When $\varphi(x) \in \mathcal{F}_M$, then we could simply let $\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x}) = \emptyset$ in the rest of the proof.

Partition $t = (t_x, t_y)$ corresponding to $Z = (X, Y)$. Although $x \mapsto \varphi(x) = 1(x \leq t_x)\mu(t_y | x)$ is a discontinuous function, $\delta \mapsto \varphi(x) \circ \psi_{d_x}(x_{-d_x}, \delta)$ is continuous for those x such that $x_{-d_x} \in \tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x})$ and $\delta = \Delta(x)$. Accordingly, we partition the integral in two regions because the integrand is not necessarily continuous on $\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x}) \times B_{\eta}(\delta)$. We use Assumption AS.1 to bound the integral in this region. Thus, for any $\epsilon > 0$, for η being small enough, the area of $\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})$, defined as $\int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \mu'(x_{-d_x}) dx_{-d_x}$, is less than or equal to ϵ by AS.1 uniformly over δ and t_{d_x} . Then, for large enough n , $k = 1, 2, \dots, d_x$ and some arbitrarily small $\zeta > 0$, by

continuity of the integrand,

$$\begin{aligned}
& \int_{\tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x})} \int_{B_\eta(\delta)} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta')}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|} \times \frac{1\{\delta \leq \delta' \leq \delta - s_n \tilde{G}_n \circ \psi_{d_x}(x_{-d_x}, \delta')\}}{s_n} d\delta' dx_{-d_x} \\
& \leq - \int_{\tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x})} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\tilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta) \\
& = - \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\tilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} \\
& \quad + \int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\tilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta),
\end{aligned}$$

where $\tilde{G} = G - K$.

The inequality above holds by continuity of the integrand $(x_{-d_x}, \delta') \mapsto (p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta') / |\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta')|$ on $\tilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x}) \times B_\eta(\delta)$, and

$$\left| \int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\tilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} \right| \leq \int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} C dx_{-d_x} \leq C\epsilon,$$

for

$$C := \sup_{x_{-d_x} \in X_{i1} \times \dots \times X_{i, d_x-1}} \left| \frac{(p_j \cdot \tilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\tilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) \right|,$$

which is bounded from above, because all components in C are bounded from above and $|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|$ is bounded away from zero. Similarly, for s_n large enough,

$$\left| \int_{\tilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \int_{B_\eta(\delta)} (p_j \cdot \psi_{d_x}) \times \frac{(1\{\delta \leq \Delta \leq \delta - s_n \tilde{G}_n(x)\} \tilde{\mu}') \circ \psi_{d_x}}{s_n |\partial_{x_{d_x}} \Delta \circ \psi_{d_x}|} d\delta' dx_{-d_x} \right| \leq C\epsilon.$$

Therefore, combining the previous results

$$\begin{aligned}
& \int_{\widetilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)) \cap \theta_i} p_j(x) \varphi(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - s_n \widetilde{G}_n(x)\}}{s_n} \widetilde{\mu}'(x) dx \\
& \leq - \int_{\widetilde{\mathcal{X}}_{d_x, \eta}^c(\delta, t_{d_x})} \frac{(p_j \cdot \widetilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\widetilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta) + C\epsilon \\
& = - \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \frac{(p_j \cdot \widetilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\widetilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} \\
& + \int_{\widetilde{\mathcal{X}}_{d_x, \eta}(\delta, t_{d_x})} \frac{(p_j \cdot \widetilde{\mu}') \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\widetilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta) + C\epsilon \\
& \leq - \int_{X_{i1} \times \dots \times X_{i, d_x-1}} \frac{(p_j \cdot \widetilde{\mu}) \circ \psi_{d_x}(x_{-d_x}, \delta)}{|\partial_{x_{d_x}} \Delta \circ \psi_{d_x}(x_{-d_x}, \delta)|} (\widetilde{G} \circ \psi_{d_x}(x_{-d_x}, \delta) - \zeta) dx_{-d_x} + o(\eta) + 2C\epsilon \\
& = - \int_{\theta_i \cap \widetilde{\mathcal{M}}_{\Delta}(\delta)} p_j(x) \varphi(x) \cdot \widetilde{\mu}'(x) \frac{\widetilde{G}(x) - \zeta}{\|\partial \Delta(x)\|} d\text{Vol} + o(\eta) + 2C\epsilon \\
& = - \int_{\theta_i \cap \mathcal{M}_{\Delta}(\delta)} p_j(x) \varphi(x) \cdot \widetilde{\mu}'(x) \frac{\widetilde{G}(x) - \zeta}{\|\partial \Delta(x)\|} d\text{Vol} + o(\eta) + 2C\epsilon,
\end{aligned}$$

where ζ , η and ϵ can be arbitrarily small for large enough n .

Similarly, we can show that

$$\begin{aligned}
& \int_{\widetilde{\mathcal{M}}_{\Delta}(B_{\eta}(\delta)) \cap \theta_i} p_j(x) \varphi(x) \frac{1\{\delta \leq \Delta(x) \leq \delta - s_n \widetilde{G}_n(x)\}}{s_n} \widetilde{\mu}'(x) dx \\
& \geq - \int_{\theta_i \cap \mathcal{M}_{\Delta}(\delta)} p_j(x) \varphi(x) \cdot \widetilde{\mu}'(x) \frac{\widetilde{G}(x) - \zeta}{\|\partial \Delta(x)\|} d\text{Vol} - o(\eta) - 2C\epsilon
\end{aligned}$$

Since we can choose η and ϵ to be arbitrarily small, we conclude that for any $\varphi \in \mathcal{F}_I$,

$$\frac{\Upsilon_{\Delta_n, \mu_n, \delta_n}(\varphi) - \Upsilon_{\Delta, \mu, \delta}(\varphi)}{s_n} \rightarrow \int_{\mathcal{M}_{\Delta}(\delta)} \varphi(x) \frac{K - G(x)}{\|\partial \Delta(x)\|} \mu'(x) d\text{Vol} + H(\varphi(x) 1\{\Delta(x) \leq \delta\}).$$

To show that the result holds uniformly in $\varphi \in \mathcal{F}_I$, we use the equivalence between uniform convergence and continuous convergence (e.g., Resnick (1987, p.2)). Take a sequence $\varphi^n \in \mathcal{F}_I$ that converges to $\varphi \in \mathcal{F}_I$ in the $L^1(\mu)$ norm, i.e., $\int_{\mathcal{X}} |\varphi^n - \varphi| d\mu \rightarrow 0$ as $n \rightarrow \infty$. Then, the preceding argument applies to this sequence and $\partial_{\Delta, \mu, \delta} \Upsilon_{\Delta, \mu, \delta}(\varphi^n)[K, G, H] \rightarrow \partial_{\Delta, \mu, \delta} \Upsilon_{\Delta, \mu, \delta}(\varphi)[K, G, H]$ by linearity of the map $\varphi \mapsto \partial_{\Delta, \mu, \delta} \Upsilon_{\Delta, \mu, \delta}(\varphi)[K, G]$.

□

Proof of Lemma A.3. Note that $\Lambda_{\Delta, \mu, \delta}^-(\varphi) = \Upsilon_{\Delta, \mu, \delta}(\varphi) / \Upsilon_{\Delta, \mu, \delta}(1)$, where $\Upsilon_{\Delta, \mu, \delta}(1) = \int 1(\Delta(x) \leq \delta) d\mu(x) = F_{\Delta, \mu}(\delta)$.

By Lemma E.5, $\Upsilon_{\Delta,\mu,\delta}(\varphi)$ and $\Upsilon_{\Delta,\mu,\delta}(1)$ are Hadamard-differentiable at (Δ, μ, δ) tangentially to $\tilde{\mathbb{D}}_0$. Then, by the chain rule for Hadamard-differentiable mappings, $\Lambda_{\tilde{\Delta},\mu,\delta}^-(\varphi)$ is Hadamard-differentiable at (Δ, μ, δ) tangentially to $\tilde{\mathbb{D}}_0$ since $\Upsilon_{\Delta,\mu,\delta}(1) > 0$. The derivative map is obtained from

$$\partial_{\Delta,\mu,\delta}\Lambda_{\tilde{\Delta},\mu,\delta}^-(\varphi) = \frac{\partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(\varphi)}{F_{\Delta,\mu}(\delta)} - \Lambda_{\Delta,\mu,\delta}(\varphi) \frac{\partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(1)}{F_{\Delta,\mu}(\delta)},$$

after replacing the expressions of $\partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(\varphi)$ and $\partial_{\Delta,\mu,\delta}\Upsilon_{\Delta,\mu,\delta}(1)$ from Lemma E.5 and grouping terms. □

APPENDIX F. SUFFICIENT CONDITIONS FOR μ -DONSKER PROPERTIES IN SECTION 4

Lemma F.1 (Sufficient conditions for \mathcal{G} being μ -Donsker). *Suppose S.1-S.2 hold, and \mathcal{V} is the union of a finite number of compact intervals. Suppose that \mathcal{F} satisfies:*

$$\sup_{\tilde{\Delta} \in \mathcal{F}} \sup_{x \in B(\mathcal{X})} \|\partial \tilde{\Delta}(x) - \partial \Delta(x)\| + \sup_{\tilde{\Delta} \in \mathcal{F}} \sup_{x \in B(\mathcal{X})} |\tilde{\Delta}(x) - \Delta(x)| < c_0.$$

Let $N(\epsilon, \mathcal{F}, \|\cdot\|_\infty)$ be the ϵ -covering number of the class \mathcal{F} under L_∞ norm. Suppose that $\int_0^1 \sqrt{\log N(\epsilon^2, \mathcal{F}, \|\cdot\|_\infty)} d\epsilon < \infty$. If c_0 is small enough, then \mathcal{G} is μ -Donsker.

Proof of Lemma F.1. Since \mathcal{V} is a union of finite number of closed intervals, for any $\zeta > 0$, we can construct a collection of closed intervals $\mathcal{I} := \{[a_i, b_i] : i = 1, 2, \dots, r\}$ such that: (1) $|b_i - a_i| < \zeta$, (2) $[a_i, b_i] \subset \mathcal{V}$, (3) $\cup_{i=1}^r [a_i, b_i] = \mathcal{V}$, (4) $a_i \leq b_i \leq a_{i+1} \leq b_{i+1}$, for all $i = 1, 2, \dots, r-1$, and (5) $r \leq \frac{C_0}{\zeta}$, where C_0 is a constant.

Using S.1 and S.2 and the assumptions of the Lemma, there exists $\eta > 0$ small enough such that the following conditions hold:

(1) There exist constants c and C such that $\|\partial \Delta(x)\| \leq C$ for all $x \in \bar{\mathcal{X}}$ and $\|\partial \Delta(x)\| \geq c$ in $\tilde{\mathcal{M}}_\Delta(B_\eta(\delta))$ for some small $\eta > 0$ and all $\delta \in \mathcal{D}$.

(2) Uniformly in $\tilde{\Delta} \in \mathcal{F}$,

$$\frac{c}{2} \leq \inf_{x \in \tilde{\mathcal{M}}_\Delta(B_\eta(\delta))} \|\partial \tilde{\Delta}(x)\| \leq \sup_{x \in \tilde{\mathcal{M}}_\Delta(B_\eta(\delta))} \|\partial \tilde{\Delta}(x)\| \leq \frac{c}{2} + C.$$

Moreover, using arguments similar to those used to show Lemma A.1, we can verify that:

(3) Uniformly in $\tilde{\Delta} \in \mathcal{F}$, uniformly in $\delta \in \mathcal{V}$,

$$f_{\tilde{\Delta},\mu}(\delta) = \int_{\mathcal{M}_{\tilde{\Delta}}(\delta)} \frac{\mu'(x)}{\|\partial \tilde{\Delta}(x)\|} d\text{Vol} < K_1,$$

for some finite constant K_1 .

Define the norm $\|g\|_{2,\mu}^2 := \int_{\mathcal{X}} g(x)^2 \mu'(x) dx$. For $\eta > 0$ small enough, for any $\delta \in \mathcal{V}$ and $\tilde{\Delta} \in \mathcal{F}$,

$$\|1(\tilde{\Delta} \leq \delta) - 1(\tilde{\Delta} \leq \delta + \eta)\|_{2,\mu}^2 = \int 1(\delta \leq \tilde{\Delta}(x) \leq \delta + \eta) \mu'(x) dx = \int_{\delta' \in B_{\eta}^+(\delta)} f_{\tilde{\Delta},\mu}(\delta') d\delta' \leq K_1 \eta.$$

Similarly, $\|1(\tilde{\Delta} \leq \delta) - 1(\tilde{\Delta} \leq \delta - \eta)\|_{2,\mu}^2 \leq K_1 \eta$.

Let $B_{\zeta,\infty}(\Delta_1), \dots, B_{\zeta,\infty}(\Delta_{q_\zeta})$ be a set of ζ -balls centered at $\Delta_1, \dots, \Delta_{q_\zeta}$ under sup norm that covers \mathcal{F} , where $q_\zeta = N(\zeta, \mathcal{F}, \|\cdot\|_\infty)$. Then, $[\Delta_j - \zeta, \Delta_j + \zeta]$ are covering brackets of \mathcal{F} , $j = 1, 2, \dots, q_\zeta$. For any $\tilde{\Delta} \in [\Delta_j - \zeta, \Delta_j + \zeta]$ and $\delta \in [a_i, b_i]$, $i = 1, 2, \dots, r$, then the bracket $[1(\Delta_j + \zeta \leq a_i), 1(\Delta_j - \zeta \leq b_i)]$ covers $1(\tilde{\Delta} \leq \delta)$. For ζ small enough, the size of the bracket $[1(\Delta_j + \zeta \leq a_i), 1(\Delta_j - \zeta \leq b_i)]$ under the norm $\|\cdot\|_{2,\mu}$ is:

$$\|1(\Delta_j + \zeta \leq a_i) - 1(\Delta_j - \zeta \leq b_i)\|_{2,\mu}^2 = \|1(\Delta_j \leq b_i + \zeta) - 1(\Delta_j \leq a_i - \zeta)\|_{2,\mu}^2 \leq 3K_1 \zeta,$$

since $|b_i - a_i| < \zeta$ by construction. Therefore, for ζ small enough, $\{[1(\Delta_j + \zeta \leq a_i), 1(\Delta_j - \zeta \leq b_i)] : j = 1, 2, \dots, q_\zeta, i = 1, 2, \dots, r\}$, form a set of $\sqrt{3K_1 \zeta}$ -brackets under the norm $\|\cdot\|_{2,\mu}$ that covers \mathcal{G} . The total number of brackets is $r q_\zeta \leq \frac{C_0}{\zeta} N(\zeta, \mathcal{F}, \|\cdot\|_\infty)$. Or equivalently, for ζ small enough,

$$N_{\square}(\zeta, \mathcal{G}, \|\cdot\|_{2,\mu}) \leq \frac{3K_1 C_0}{\zeta^2} N(\zeta^2/(3K_1), \mathcal{F}, \|\cdot\|_\infty).$$

Then by assumption,

$$\begin{aligned} \int_0^1 \sqrt{\log(N_{\square}(\zeta, \mathcal{G}, \|\cdot\|_{2,\mu}))} d\zeta &\leq \int_0^1 \sqrt{\log\left(\frac{3K_1 C_0}{\zeta^2} N(\zeta^2/(3K_1), \mathcal{F}, \|\cdot\|_\infty)\right)} d\zeta \\ &\lesssim \int_0^1 \sqrt{\log\left(\frac{3K_1 C_0}{\zeta^2}\right)} d\zeta + \int_0^1 \sqrt{\log(N(\zeta^2/(3K_1), \mathcal{F}, \|\cdot\|_\infty))} d\zeta < \infty. \end{aligned}$$

We conclude that \mathcal{G} is μ -Donsker by Donsker theorem (van der Vaart, 1998, Theorem 19.5). \square

Lemma F.2 (Sufficient conditions for $\tilde{\mathcal{G}}$ being μ -Donsker). *Suppose S.1-S.2 hold, and \mathcal{V} is the union of a finite number of compact intervals. Suppose that \mathcal{F} satisfies:*

$$\sup_{\tilde{\Delta} \in \mathcal{F}} \sup_{x \in B(\mathcal{X})} \|\partial \tilde{\Delta}(x) - \partial \Delta(x)\| + \sup_{\tilde{\Delta} \in \mathcal{F}} \sup_{x \in B(\mathcal{X})} |\tilde{\Delta}(x) - \Delta(x)| < c_0.$$

Let $N(\epsilon, \mathcal{F}, \|\cdot\|_\infty)$ be the ϵ -covering number of the class \mathcal{F} under L_∞ norm. Suppose that $\int_0^1 \sqrt{\log N(\epsilon^2, \mathcal{F}, \|\cdot\|_\infty)} d\epsilon < \infty$. If c_0 is small enough, then $\tilde{\mathcal{G}}$ is μ -Donsker.

Proof of Lemma F.2. First, \mathcal{F}_I and \mathcal{F}_M are both μ -Donsker. By Lemma F.1, the class \mathcal{F} is μ -Donsker. Since the class of the product of two functions from Donsker classes is Donsker, $\tilde{\mathcal{G}}$ is μ -Donsker. \square

APPENDIX G. EXTENSION OF THEORETICAL ANALYSIS TO DISCRETE VARIABLES

We consider the case where the covariate X includes discrete components. Without loss of generality we assume that the first component of X is discrete and the rest are continuous. Accordingly, we consider the partition $X = (D, C)$. Let $\mathcal{X}_{c|d}$ denote the interior of the support of C conditional on $D = d$, \mathcal{X}_d denote the support of D , $\mu_{c|d}$ denote the distribution of C conditional on $D = d$, μ_d denote the distribution of D , and $\pi_d(d) = \mathbb{P}(D = d)$. As above, $d_x = \dim(X)$, and \mathcal{D} is a compact set consisting of regular values of Δ on $\overline{\mathcal{X}} := \cup_{d \in \mathcal{X}_d} \{d\} \times \overline{\mathcal{X}_{c|d}}$, where $\overline{\mathcal{X}_{c|d}}$ is the closure of $\mathcal{X}_{c|d}$.

We adjust S.1-S.4 to hold conditionally at each value of the discrete covariate.

S.1'. The set \mathcal{X}_d is finite. For any $d \in \mathcal{X}_d$: the set $\mathcal{X}_{c|d}$ is open and its closure $\overline{\mathcal{X}_{c|d}}$ is compact; the distribution $\mu_{c|d}$ is absolutely continuous with respect to the Lebesgue measure with density $\mu'_{c|d}$; and there exists an open set $B(\mathcal{X}_{c|d})$ containing $\overline{\mathcal{X}_{c|d}}$ such that $c \mapsto \Delta(d, c)$ is \mathcal{C}^1 on $B(\mathcal{X}_{c|d})$, and $c \mapsto \mu'_{c|d}(c)$ is continuous on $B(\mathcal{X}_{c|d})$ and is zero outside $\mathcal{X}_{c|d}$, i.e. $\mu'(x) = 0$ for any $x \in B(\mathcal{X}_{c|d}) \setminus \mathcal{X}_{c|d}$.

S.2'. For any $d \in \mathcal{X}_d$ and any regular value δ of Δ on $\overline{\mathcal{X}_{c|d}}$, $\mathcal{M}_{\Delta|d}(\delta) := \{c \in \overline{\mathcal{X}_{c|d}} : \Delta(d, c) = \delta\}$ is either a $(d_x - 2)$ -manifold without boundary on \mathbb{R}^{d_x-1} of class \mathcal{C}^1 with finite number of connected branches, or an empty set.

S.3'. $\widehat{\Delta}$, the estimator of Δ , obeys a functional central limit theorem, namely,

$$a_n(\widehat{\Delta} - \Delta) \rightsquigarrow G_\infty \text{ in } \ell^\infty(B(\mathcal{X})),$$

where a_n is a sequence such that $a_n \rightarrow \infty$ as $n \rightarrow \infty$, and $c \mapsto G_\infty(d, c)$ is a tight process that has almost surely uniformly continuous sample paths on $B(\mathcal{X}_{c|d})$ for all $d \in \mathcal{X}_d$.

Let $B(\mathcal{X}) := \cup_{d \in \mathcal{X}_d} \{d\} \times B(\mathcal{X}_{c|d})$; \mathcal{F} denote a set of continuous functions on $B(\mathcal{X})$ equipped with the sup-norm; \mathcal{V} be any compact subset of \mathbb{R} ; \mathbb{H} be the set of all bounded operators $H : g \mapsto H(g)$ uniformly continuous on $\mathcal{G} = \{1(f \leq \delta) : f \in \mathcal{F}, \delta \in \mathcal{V}\}$ with respect to the $L^2(\mu)$ norm, which are represented as:

$$H(g) = \sum_{d \in \mathcal{X}_d} H_d(d) \int g(c, d) d\mu_{c|d}(c) + \sum_{d \in \mathcal{X}_d} \pi_d(d) H_{c|d}(g(\cdot, d)),$$

where $d \mapsto H_d(d)$ is a function that takes on finitely many values and $g \mapsto H_{c|d}(g)$ is a bounded linear operator on \mathcal{G} . Equip the space \mathbb{H} with the sup norm $\|\cdot\|_{\mathcal{G}}$: $\|H\|_{\mathcal{G}} = \sup_{g \in \mathcal{G}} |H(g)|$. Let $\mu(x) = \mu_d(d)\mu_{c|d}(c)$ and $\widehat{\mu}(x) = \widehat{\mu}_d(d)\widehat{\mu}_{c|d}(c)$.

S.4'. The function $x \mapsto \widehat{\mu}(x)$ is a distribution over $B(\mathcal{X})$ obeying in \mathbb{H} ,

$$b_n(\widehat{\mu} - \mu) \rightsquigarrow H_\infty,$$

where $H_\infty \in \mathbb{H}$ a.s., b_n is a sequence such that $b_n \rightarrow \infty$ as $n \rightarrow \infty$, and H_∞ can be represented as:

$$H_\infty(g) = \sum_{d \in \mathcal{X}_d} H_{d,\infty}(d) \int g(c, d) d\mu_{c|d}(c) + \sum_{d \in \mathcal{X}_d} \pi_d(d) H_{c|d,\infty}(g(\cdot, d)).$$

We generalize Lemmas A.1 and A.2 to the case where X includes discrete components.

Define $\mathbb{D} := \mathbb{F} \times \mathbb{H}$ and $\mathbb{D}_0 := \mathbb{F}_0 \times \mathbb{H}$, where \mathbb{F} is the set of continuous functions on $B(\mathcal{X})$ and \mathbb{F}_0 is a subset of \mathbb{F} containing uniformly continuous functions.

Lemma G.1 (Properties of $F_{\Delta,\mu}$ and Δ_μ^* with discrete X). *Suppose that S.1' and S.2' hold. Then, $\delta \mapsto F_{\Delta,\mu}(\delta)$ is differentiable at any $\delta \in \mathcal{D}$, with derivative function $f_{\Delta,\mu}(\delta)$ defined as:*

$$f_{\Delta,\mu}(\delta) := \partial_\delta F_{\Delta,\mu}(\delta) = \sum_{d \in \mathcal{X}_d} \pi_d(d) \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{\mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol}.$$

The map $\delta \mapsto f_{\Delta,\mu}(\delta)$ is uniformly continuous on \mathcal{D} .

(1) The map $F_{\Delta,\mu}(\delta) : \mathbb{D} \rightarrow \mathbb{R}$ is Hadamard differentiable uniformly in $d \in \mathcal{D}$ at (Δ, μ) tangentially to \mathbb{D}_0 , with derivative map $\partial_{\Delta,\mu} F_{\Delta,\mu}(\delta) : \mathbb{D}_0 \rightarrow \mathbb{R}$ defined by:

$$\begin{aligned} (G, H) \mapsto \partial_{\Delta,\mu} F_{\Delta,\mu}(\delta)[G, H] := & - \sum_{d \in \mathcal{X}_d} \pi_d(d) \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{G(d, c) \mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol}(c) \\ & + \sum_{d \in \mathcal{X}_d} H_d(d) \int \mathbf{1}\{\Delta(d, c) \leq \delta\} \mu'_{c|d}(c) dc \\ & + \sum_{d \in \mathcal{X}_d} \pi_d(d) H_{c|d}(\mathbf{1}\{\Delta(\cdot, d) \leq \delta\}). \end{aligned}$$

(2) The map $\Delta_\mu^*(u) : \mathbb{D} \rightarrow \mathbb{R}$ is Hadamard differentiable uniformly in $u \in \mathcal{U}$ at (Δ, μ) tangentially to \mathbb{D}_0 , with derivative map $\partial_{\Delta,\mu} \Delta_\mu^*(u) : \mathbb{D}_0 \rightarrow \mathbb{R}$ defined by:

$$(G, H) \mapsto \partial_{\Delta,\mu} \Delta_\mu^*(u)[G, H] := - \frac{\partial F_{\Delta,\mu}(\Delta_\mu^*(u))[G, H]}{f_{\Delta,\mu}(\Delta_\mu^*(u))},$$

where $\mathcal{U} = \{\tilde{u} \in [0, 1] : \Delta_\mu^*(\tilde{u}) \in \mathcal{D}, f_{\Delta,\mu}(\Delta_\mu^*(\tilde{u})) > \varepsilon\}$ for fixed $\varepsilon > 0$.

Proof of Lemma G.1. Note that $F_{\Delta,\mu}(\delta) = \sum_{d \in \mathcal{X}_d} \pi_d(d) \int_{c \in \mathcal{X}_d} \mathbf{1}(\Delta(d, c) \leq \delta) \mu'_{c|d}(c) dc$. Given the results of Lemma A.1, for each d ,

$$\partial_\delta \int_{\mathcal{X}_{c|d}} \mathbf{1}(\Delta(d, c) \leq \delta) \mu'_{c|d}(c) dc = \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{\mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol}.$$

Therefore, averaging over $d \in \mathcal{X}_d$,

$$f_{\Delta,\mu}(\delta) := \partial_\delta F_{\Delta,\mu}(\delta) = \sum_{d \in \mathcal{X}_d} \pi_d(d) \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{\mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol},$$

where we use that \mathcal{X}_d is a finite set.

Next we prove the statements (1) and (2). Let $G_n \in \mathbb{F}$ and $H_n \in \mathbb{H}$ such that $G_n \rightarrow G \in \mathbb{F}_0$ and $H_n \rightarrow H \in \mathbb{H}$. Let $\Delta_n = \Delta + t_n G_n$ and $\mu_n = \mu + t_n H_n$, where $t_n \rightarrow 0$ as $n \rightarrow \infty$.

As in the proof of Lemma A.2, we decompose

$$F_{\Delta_n, \mu_n}(\delta) - F_{\Delta, \mu}(\delta) = [F_{\Delta_n, \mu_n}(\delta) - F_{\Delta_n, \mu}(\delta)] + [F_{\Delta_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)].$$

Applying the same argument as in the proof of Lemma A.2 to each d and averaging over $d \in \mathcal{X}_d$, for any $\delta \in \mathcal{D}$

$$\frac{F_{\Delta_n, \mu}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} = - \sum_{d \in \mathcal{X}_d} \mu_d(d) \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{G(d, c) \mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol} + o(1),$$

where we use that \mathcal{X}_d is a finite set. By assumption S.4' and a similar argument to the proof of Lemma A.2,

$$\frac{F_{\Delta_n, \mu_n}(\delta) - F_{\Delta_n, \mu}(\delta)}{t_n} = H(g_{\Delta, \delta}) + o(1), \quad g_{\Delta, \delta}(c, d) = 1\{\Delta(c, d) \leq \delta\}$$

We conclude that for any $\delta \in \mathcal{D}$,

$$\frac{F_{\Delta_n, \mu_n}(\delta) - F_{\Delta, \mu}(\delta)}{t_n} \rightarrow - \sum_{d \in \mathcal{X}_d} \mu_d(d) \int_{\mathcal{M}_{\Delta|d}(\delta)} \frac{G(d, c) \mu'_{c|d}(c)}{\|\partial_c \Delta(d, c)\|} d\text{Vol} + H(g_{\Delta, \delta}) = \partial_{\Delta, \mu} F_{\Delta, \mu}(\delta)[G, H].$$

By an argument similar to the proof of Lemma A.2, it can be shown that the convergence is uniform in $\delta \in \mathcal{D}$. This shows statement (1).

Statement (2) follows from statement (1) and Theorem 3.9.20 of van der Vaart and Wellner (1996) for inverse maps, using an argument analogous to the proof of statement (b) in Lemma A.2. \square

We are now ready to derive a functional central limit theorem for the empirical SPE-function. As in Theorem 4.1, let $r_n := a_n \wedge b_n$, the slowest of the rates of convergence of $\widehat{\Delta}$ and $\widehat{\mu}$, where $r_n/a_n \rightarrow s_\Delta \in [0, 1]$ and $r_n/b_n \rightarrow s_\mu \in [0, 1]$.

Theorem G.1 (FCLT for $\widehat{\Delta}_\mu^*(u)$ with discrete X). *Suppose that S.1'-S.4' hold, the convergence in S.3' and S.4' holds jointly, and $\widehat{\Delta} \in \mathcal{F}$ with probability approaching 1. Then, the empirical SPE-process obeys a functional central limit theorem, namely in $\ell^\infty(\mathcal{U})$,*

$$r_n(\widehat{\Delta}_\mu^*(u) - \Delta_\mu^*(u)) \rightsquigarrow \partial_{\Delta, \mu} \Delta_\mu^*(u)[s_\Delta G_\infty, s_\mu H_\infty], \quad (\text{G.23})$$

as a stochastic process indexed by $u \in \mathcal{U}$, where \mathcal{U} is defined in Lemma G.1.

Remark G.1 (Bootstrap FCLT for $\widehat{\Delta}_\mu^*(u)$ with discrete X). The exchangeable bootstrap is consistent to approximate the distribution of the limit process in (G.23) under the same conditions as in Theorem 4.3, replacing S.1-S.4 by S.1'-S.4'. Accordingly, we do not repeat the statement here. \square

Remark G.2 (CA with discrete covariates). The results of the classification analysis can also be extended to the case where X contains discrete components following analogous arguments as for the SPE. We omit the details for the sake of brevity. \square

Proof of Theorem G.1. The result follows from Lemma G.1 and Lemma B.1. \square

APPENDIX H. SOME NUMERICAL ILLUSTRATIONS

We evaluate the accuracy of the asymptotic approximations to the distribution of the empirical SPE in small samples using numerical simulations. In particular, we compare pointwise 95% confidence intervals for the SPE based on the asymptotic and exact distributions of the empirical SPE. The exact distribution is approximated numerically by simulation. The asymptotic distribution is obtained analytically from the FCLT of Theorem 4.1, and approximated by bootstrap using Theorem 4.3. We first consider two simulation designs where the limit process in Theorem 4.1 has a convenient closed-form analytical expression. The designs differ on whether the PE-function $x \mapsto \Delta(x)$ has critical points or not. We hold fix the values of the covariate vector X in all the calculations, and accordingly we treat the distribution μ as known. For the bootstrap inference, we use empirical bootstrap with $B = 3,000$ repetitions. All the results are based on 3,000 simulations. The last design is calibrated to mimic the gender wage gap application.

Design 1 (No critical points). We consider the PE-function

$$\Delta(x) = x_1 + x_2, \quad x = (x_1, x_2),$$

with the covariate vector X uniformly distributed in $\mathcal{X} = (-1, 1) \times (-1, 1)$. The corresponding SPE is

$$\Delta_\mu^*(u) = 2(\sqrt{2u} - 1)1(u \leq 1/2) + 2(1 - \sqrt{2(1-u)})1(u > 1/2),$$

where we use that $\Delta(X)$ has a triangular distribution with parameters $(-2, 0, 2)$. The sample size is $n = 441$ and the values of X are held fixed in the grid $\{-1, -0.9, \dots, 1\} \times \{-1, -0.9, \dots, 1\}$. Figure 6 plots $x \mapsto \Delta(x)$ on \mathcal{X} , and $u \mapsto \Delta_\mu^*(u)$ on $(0, 1)$. Here we see that $x \mapsto \Delta(x)$ does not have critical values, and that $u \mapsto \Delta_\mu^*(u)$ is a smooth function.

To obtain an analytical expression of the limit $Z_\infty(u)$ of Theorem 4.1, we make the following assumption on the estimator of the PE:

$$\sqrt{n}(\widehat{\Delta}(x) - \Delta(x)) = \exp[\Delta(x)] \sum_{i=1}^n Z_i / \sqrt{n},$$

where Z_1, \dots, Z_n is an i.i.d. sequence of standard normal random variables. Hence

$$Z_\infty(u) \sim N(0, \exp[2\Delta_\mu^*(u)]),$$

so that $\widehat{\Delta}_\mu^*(u) \overset{a}{\sim} N(\Delta_\mu^*(u), \exp[2\Delta_\mu^*(u)]/n)$, where $\overset{a}{\sim}$ denotes asymptotic approximation to the distribution.




FIGURE 6. PE-function and SPE-function in Design 1. Left: PE function $x \mapsto \Delta(x)$. Right: SPE function $u \mapsto \Delta_\mu^*(u)$.

Table 4 reports biases and compares the standard deviations of the empirical SPE with the asymptotic standard deviations, $\exp[\Delta_\mu^*(u)]/\sqrt{n}$, at the quantile indices $u \in \{0.1, 0.2, \dots, 0.9\}$. The biases are small relative to dispersions and the asymptotic approximations are very close to the exact standard deviations. We also find that 95% confidence intervals constructed using the asymptotic approximations, $\hat{\Delta}_\mu^*(u) \pm 1.96 \exp[\Delta_\mu^*(u)]/\sqrt{n}$, have coverage probabilities close to their nominal levels at all indices. These asymptotic confidence intervals are not feasible in general, either because $\Delta_\mu^*(u)$ are unknown or more generally because it is not possible to characterize analytically the distribution of $Z_\infty(u)$. In practice we propose approximating this distribution by bootstrap. In this case the empirical bootstrap version of the empirical SPE is constructed from the bootstrap PE

$$\tilde{\Delta}(x) = \Delta(x) + \exp[\Delta(x)] \sum_{i=1}^n \omega_i Z_i/n,$$

where $(\omega_1, \dots, \omega_n)$ is a multinomial vector with dimension n and probabilities $(1/n, \dots, 1/n)$ independent of Z_1, \dots, Z_n . The last column of the table shows that the empirical coverages of bootstrap 95% confidence intervals are close to their nominal levels at all quantile indices.

Design 2 (Critical points). We consider the PE-function

$$\Delta(x) = x^3 - 3x,$$

TABLE 4. Properties of Empirical SPE in Design 1

u	Bias	Std. Dev.		Pointwise Coverage (%)	
	($\times 100$)	Exact	Asymptotic	Asymptotic	Bootstrap [†]
0.1	0.016	0.014	0.014	95.10	95.03
0.2	0.024	0.021	0.021	95.10	95.03
0.3	0.032	0.029	0.029	95.10	95.03
0.4	0.044	0.039	0.039	95.10	95.03
0.5	0.053	0.047	0.048	95.10	95.03
0.6	0.065	0.058	0.058	95.10	95.03
0.7	0.088	0.078	0.079	95.10	95.03
0.8	0.119	0.105	0.106	95.10	95.03
0.9	0.177	0.157	0.158	95.10	95.03

Notes: 3,000 simulations with sample size $n = 441$.

[†]3,000 bootstrap repetitions. Nominal level is 95%.

with covariate X uniformly distributed on $\mathcal{X} = (-3, 3)$. Figure 7 plots $x \mapsto \Delta(x)$ on \mathcal{X} , and $u \mapsto \Delta_\mu^*(u)$ on $(0, 1)$.¹⁶ Here we see that $x \mapsto \Delta(x)$ has two critical points at $x = -1$ and $x = 1$ with corresponding critical values at $\delta = 2$ and $\delta = -2$. The SPE-function $u \mapsto \Delta_\mu^*(u)$ has two kinks at $u = 1/6$ and $u = 5/6$, the Δ_μ^* pre-images of the critical values.

To obtain an analytical expression of the limit $Z_\infty(u)$ of Theorem 4.1, we make the following assumption on the estimator of the PE:

$$\sqrt{n}(\widehat{\Delta}(x) - \Delta(x)) = (x/2)^2 \sum_{i=1}^n Z_i / \sqrt{n},$$

where Z_1, \dots, Z_n is an i.i.d. sequence of standard normal variables. This assumption is analytically convenient because after some calculations we find that for $u \notin \{1/6, 5/6\}$,

$$Z_\infty(u) \sim N(0, S(\Delta_\mu^*(u))^2 / (4n)),$$

where

$$S(\delta) = 1(\delta < -2)\check{\Delta}_1(\delta)^2 + 1(-2 < \delta < 2) \sum_{k=1}^3 \frac{\check{\Delta}_k(\delta)^2 |\check{\Delta}_k(\delta)^2 - 1|^{-1}}{\sum_{j=1}^3 |\check{\Delta}_j(\delta)^2 - 1|^{-1}} + 1(\delta > 2)\check{\Delta}_1(\delta)^2,$$

and $\check{\Delta}_1(\delta)$, $\check{\Delta}_2(\delta)$ and $\check{\Delta}_3(\delta)$ are real roots of $\Delta(x) - \delta = 0$ sorted in increasing order.¹⁷ Hence, $\widehat{\Delta}_\mu^*(u) \stackrel{a}{\sim} N(\Delta_\mu^*(u), S(\Delta_\mu^*(u))^2 / (4n))$.

¹⁶We obtain $u \mapsto \Delta_\mu^*(u)$ analytically using the characterization of Chernozhukov, Fernández-Val, and Galichon (2010) for the univariate case.

¹⁷The equation $\Delta(x) - \delta = x^3 - 3x - \delta = 0$ has three real roots when $\delta \in (-2, 2)$, and one real root when $\delta < -2$ or $\delta > 2$.




FIGURE 7. PE-function and SPE-function in Design 2. Left: PE function $x \mapsto \Delta(x)$. Right: SPE function $u \mapsto \Delta_\mu^*(u)$.

Table 5 reports biases and compares the standard deviations of the empirical SPE in samples of size $n = 601$ with the asymptotic standard deviations at the quantile indices $u \in \{1/12, 2/12, \dots, 11/12\}$, where the values of X are held fixed in the grid $\{-3, -2.99, \dots, 3\}$. The biases are small relative to dispersion except at the kinks $u = 1/6$ and $u = 5/6$. The asymptotic approximation is close to the exact standard deviation, except for the quantiles at the kinks where the asymptotic standard deviations are not well-defined because $\check{\Delta}_k(\delta)^2 - 1 = 0$. We also find that pointwise 95% confidence intervals constructed using the asymptotic distribution and empirical bootstrap have coverage probabilities close to their nominal levels. Interestingly, the bootstrap provides coverages close to the nominal levels even at the kinks.

Design 3 (Calibration to CPS data). This design is calibrated to the interactive linear model with additive error for the conditional expectation in the gender wage gap application of Section 3. More specifically, we generate log wages as

$$Y_i = P(T_i, W_i)' \beta + \sigma \varepsilon_i, \quad i = 1, \dots, n,$$

where the covariates $X_i = (T_i, W_i)$ are fixed to the values in the 2015 CPS data set, $P(T, W) = (TW, (1 - T)W)$, β and σ^2 are the least squares estimates of the regression coefficients and residual variance in the data set, $(\varepsilon_1, \dots, \varepsilon_n)$ is a sequence of i.i.d. standard normal random variables independent of X_i , and $n = 32,523$, the sample size in the application. For each

TABLE 5. Properties of Empirical SPE in Design 2

u	Bias	Std. Dev.		Pointwise Coverage (%)	
	($\times 100$)	Exact	Asymptotic	Asymptotic	Bootstrap [†]
1/12	0.068	0.126	0.127	95.67	95.80
1/6	-2.393	0.054	–	–	95.67
1/4	-0.005	0.025	0.025	95.83	95.77
1/3	-0.016	0.028	0.028	95.80	95.90
5/12	0.045	0.030	0.030	95.63	95.47
1/2	0.023	0.030	0.031	92.73	97.53
7/12	-0.020	0.030	0.030	95.20	95.80
2/3	0.049	0.028	0.028	95.53	95.67
3/4	0.039	0.025	0.025	95.53	95.73
5/6	2.447	0.053	–	–	95.73
11/12	0.068	0.126	0.127	95.67	95.80

Notes: 3,000 simulations with sample size $n = 601$.

[†]3,000 bootstrap repetitions. Nominal level is 95%.




FIGURE 8. Bias, standard deviation and root mean square error of empirical and bias corrected SPE functions. Results obtained from 500 repetitions of a design calibrated to the CPS 2015 data.

simulated sample $\{(Y_i, X_i) : 1 \leq i \leq n\}$, we reestimate the model by least squares, obtain the empirical SPE-function on the treated over a grid of percentile indexes $\mathcal{U} = \{0.02, 0.03, \dots, 0.98\}$, and construct a 90% uniform confidence band for the SPE-function using Algorithm 2.1 with standard exponential weights and $B = 200$. We repeat this procedure 500 times.

Figure 8 reports bias, standard deviation (Std. Dev.) and root mean squared error (RMSE) of the empirical and bias corrected SPE functions, see Remark 2.3. We find that the empirical SPE displays negative bias in the lower tail and positive bias in the upper tail, which are reduced by the bootstrap bias correction. The correction slightly increases dispersion, but reduces overall rmse for most percentiles, specially at the tails. Table 8 reports the empirical coverage of 90% confidence bands constructed around the empirical and bias corrected SPE functions. Here we find that the uncorrected bands undercover the entire SPE function, whereas the corrected bands have coverage above the nominal level. One possible reason for the overcoverage of the bootstrap corrected bands is that we keep the covariates fixed across samples, which is not accounted by the bootstrap procedure. To sum up, we find that the bootstrap corrections of Remark 2.3 reduce the bias of the empirical SPE and improve the coverage of the confidence bands in finite samples.

TABLE 6. Coverage of 90% Confidence Bands

	Uncorrected	Bootstrap Bias Corrected
Coverage	0.80	0.98

Notes: 500 simulations and 200 bootstrap repetitions.

DGP calibrated to CPS 2015.

APPENDIX I. EFFECT OF RACE ON MORTGAGE DENIALS

To study the effect of race in the bank decisions of mortgage denials or racial mortgage denial gap, we use data on mortgage applications in Boston from 1990 (see Munnell, Tootell, Browne, and McEneaney (1996)). The Federal Reserve Bank of Boston collected these data in relation to the Home Mortgage Disclosure Act (HMDA), which was passed to monitor minority access to the mortgage market. Providing better access to credit markets can arguably help the disadvantaged groups escape poverty traps. Following Stock and Watson (2011, Chap 11), we focus on white and black applicants for single-family residences. The sample includes 2,380 observations corresponding to 2,041 white applicants and 339 black applicants.

We estimate a binary response model where the outcome variable Y is an indicator for mortgage denial, the key covariate T is an indicator for the applicant being black, and the controls W contain financial and other characteristics of the applicant that banks take into account in the mortgage decisions. These include the monthly debt to income ratio; monthly housing expenses to income ratio; a categorical variable for “bad” consumer credit score with 6 categories (1 if no slow payments or delinquencies, 2 if one or two slow payments or delinquencies, 3 if more than two slow payments or delinquencies, 4 if insufficient credit history for determination, 5 if delinquent credit history with payments 60 days overdue, and 6 if delinquent credit history with payments 90 days overdue); a categorical variable for “bad” mortgage credit score with 4 categories (1 if no late mortgage payments, 2 if no mortgage payment history, 3 if one or two late mortgage payments,

and 4 if more than two late mortgage payments); an indicator for public record of credit problems including bankruptcy, charge-offs, and collective actions; an indicator for denial of application for mortgage insurance; two indicators for medium and high loan to property value ratio, where medium is between .80 and .95 and high is above .95; and three indicators for self-employed, single, and high school graduate.

TABLE 7. Descriptive Statistics of Mortgage Applicants

	All	Black	White
Deny	0.12	0.28	0.09
Black	0.14	1.00	0.00
Debt-to-income ratio	0.33	0.35	0.33
Expenses-to-income ratio	0.26	0.27	0.25
Bad consumer credit	2.12	3.02	1.97
Bad mortgage credit	1.72	1.88	1.69
Credit problems	0.07	0.18	0.06
Denied mortgage insurance	0.02	0.05	0.02
Medium loan-to-value ratio	0.37	0.56	0.34
High loan-to-value ratio	0.03	0.07	0.03
Self-employed	0.12	0.07	0.12
Single	0.39	0.52	0.37
High school graduate	0.98	0.97	0.99
number of observations	2,380	339	2,041

Table 7 reports the sample means of the variables used in the analysis. The probability of having the mortgage denied is 19% higher for black applicants than for white applicants. However, black applicants are more likely to have socio-economic characteristics linked to a denial of the mortgage.

Figure 9 plots estimates and 90% confidence sets of the population APE and SPE-function of being black. The PEs are obtained as described in Example 1 of the main text using a logit model with $P(X) = X = (T, W)$ and $\hat{\mu}$ equal to the empirical distribution of X in the whole sample. The confidence bands are constructed using Algorithm 2.1 with multinomial weights (empirical bootstrap) and $B = 500$, and are uniform for the SPE-function over the grid $\mathcal{U} = \{.02, .03, \dots, .98\}$. We monotone the bands using the rearrangement method of Chernozhukov, Fernández-Val, and Galichon (2009). After controlling for applicant characteristics, black applicants are still on average 5.3% more likely to have the mortgage denied than white applicants. Moreover, the SPE-function shows significant heterogeneity, with the PE ranging between 0 and 15%. Thus, there exists a subgroup of applicants that is 15% more likely to be denied a mortgage if they were black, and there is a subgroup of applicants that is not affected by racial mortgage denial gap. Table 8 shows the results of the classification analysis, answering the question “who is affected the most and who the least?” The table shows that the 10% of the applicants *most affected* by




FIGURE 9. APE and SPE (introduced in this paper) of being black on the probability of mortgage denial. Estimates and 90% bootstrap uniform confidence bands (derived in this paper) based on a logit model are shown.

racial mortgage denial gap are *more likely* to have either of the following characteristics relative to the 10% of the *least affected* applicants: self employed, single, black, high debt to income ratio, high expense to income ratio, high loan to value ratio, medium or high loan-to-income ratio, bad consumer or credit scores, and credit problems.

REFERENCES

- ALBRECHT, J., A. BJORKLUND, AND S. VROMAN (2003): “Is There a Glass Ceiling in Sweden?,” *Journal of Labor Economics*, 21(1), 145–177.
- ANGRIST, J. D., AND J.-S. PISCHKE (2008): *Mostly harmless econometrics: An empiricist’s companion*. Princeton university press.
- BHATTACHARYA, D., AND P. DUPAS (2012): “Inferring welfare maximizing treatment assignment under budget constraints,” *J. Econometrics*, 167(1), 168–196.
- BLAU, F. D., AND L. M. KAHN (2017): “The Gender Wage Gap: Extent, Trends, and Explanations,” *Journal of Economic Literature*, 55(3), 789–865.
- CHERNOZHUKOV, V., I. FERNÁNDEZ-VAL, AND A. GALICHON (2009): “Improving point and interval estimators of monotone functions by rearrangement,” *Biometrika*, 96(3), 559–575.
- CHERNOZHUKOV, V., I. FERNÁNDEZ-VAL, AND A. GALICHON (2010): “Quantile and probability curves without crossing,” *Econometrica*, 78(3), 1093–1125.
- CHERNOZHUKOV, V., I. FERNANDEZ-VAL, AND Y. LUO (2017): “Supplement to “The Sorted Effects Method: Discovering Heterogenous Effects Beyond Their Averages”,” Discussion paper, ArXiv.
- CHERNOZHUKOV, V., I. FERNÁNDEZ-VAL, AND B. MELLY (2013): “Inference on counterfactual distributions,” *Econometrica*, 81(6), 2205–2268.
- CHERNOZHUKOV, V., H. HONG, AND E. TAMER (2007): “Estimation and Inference on Identified Parameter Sets in Econometric Models,” *Econometrica*.

TABLE 8. Who is affected the most and who the least? Classification Analysis – Averages of Characteristics of the Mortgage Applicants Least and Most Affected by Racial Discrimination

Characteristics of the Group	10% Most Affected		10% Least Affected	
	PE	> .11	PE	< .018
Deny	0.44	(0.03)	0.11	(0.04)
Black	0.37	(0.04)	0.07	(0.02)
Debt-to-income	0.39	(0.01)	0.25	(0.02)
Expenses-to-income	0.28	(0.01)	0.21	(0.02)
Bad consumer credit	4.64	(0.25)	1.31	(0.09)
Bad mortgage credit	1.99	(0.07)	1.37	(0.12)
Credit problems	0.45	(0.05)	0.05	(0.02)
Denied mortgage insurance	0.01	(0.01)	0.06	(0.04)
Medium loan-to-house	0.58	(0.06)	0.07	(0.04)
High loan-to-house	0.13	(0.03)	0.02	(0.01)
Self employed	0.18	(0.05)	0.05	(0.03)
Single	0.59	(0.05)	0.11	(0.06)
High school grad	0.93	(0.03)	1.00	(0.01)

Std. errors in parentheses obtained by bootstrap with 200 repetitions.

- CHERNOZHUKOV, V., E. KOCATULUM, AND K. MENZEL (2015): “Inference on sets in finance,” *Quant. Econ.*, 6(2), 309–358.
- COX, D. R. (1984): “Interaction,” *Internat. Statist. Rev.*, 52(1), 1–31, With discussion and a reply by the author.
- HAUSMAN, J. A., AND W. K. NEWEY (2017): “Nonparametric Welfare Analysis,” *Annual Review of Economics*, 9(1), 521–546.
- KIM, J., AND D. POLLARD (1990): “Cube Root Asymptotics,” *Ann. Statist.*, 18(1), 191–219.
- KOENKER, R. (2005): *Quantile regression*. Cambridge university press.
- KOENKER, R., AND G. BASSET (1978): “Regression Quantiles,” *Econometrica*, 46(1), 33–50.
- LIST, J. A., A. M. SHAIKH, AND Y. XU (2016): “Multiple hypothesis testing in experimental economics,” Discussion paper, National Bureau of Economic Research.
- MULLIGAN, C. B., AND Y. RUBINSTEIN (2008): “Selection, Investment, and Women’s Relative Wages Over Time,” *The Quarterly Journal of Economics*, 123(3), 1061–1110.
- MUNKRES, J. R. (1991): *Analysis on manifolds*. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA.
- MUNNELL, A. H., G. M. B. TOOTELL, L. E. BROWNE, AND J. MCEANEY (1996): “Mortgage Lending in Boston: Interpreting HMDA Data,” *The American Economic Review*, 86(1), pp. 25–53.
- OAXACA, R. (1973): “Male-Female Wage Differentials in Urban Labor Markets,” *International Economic Review*, 14(3), 693–709.
- PRÆSTGAARD, J., AND J. A. WELLNER (1993): “Exchangeably weighted bootstraps of the general empirical process,” *Ann. Probab.*, 21(4), 2053–2086.
- RESNICK, S. I. (1987): *Extreme values, regular variation, and point processes*, vol. 4 of *Applied Probability. A Series of the Applied Probability Trust*. Springer-Verlag, New York.

- ROMANO, J. P., AND A. M. SHAIKH (2010): "Inference for the identified set in partially identified econometric models," *Econometrica*, 78(1), 169–211.
- (2012): "On the uniform asymptotic validity of subsampling and the bootstrap," *The Annals of Statistics*, 40(6), 2798–2822.
- ROMANO, J. P., A. M. SHAIKH, AND M. WOLF (2010a): "Hypothesis Testing in Econometrics," *Annual Review of Economics*, 2(1), 75–104.
- (2010b): "multiple testing," in *The New Palgrave Dictionary of Economics*, ed. by S. N. Durlauf, and L. E. Blume. Palgrave Macmillan, Basingstoke.
- ROMANO, J. P., AND M. WOLF (2016): "Efficient computation of adjusted p-values for resampling-based stepdown multiple testing," *Statistics & Probability Letters*, 113, 38–40.
- ROSENBAUM, P. R., AND D. B. RUBIN (1983): "The central role of the propensity score in observational studies for causal effects," *Biometrika*, 70(1), 41–55.
- SASAKI, Y. (2015): "What do quantile regressions identify for general structural functions?," *Econometric Theory*, 31(5), 1102–1116.
- SPIVAK, M. (1965): *Calculus on manifolds. A modern approach to classical theorems of advanced calculus*. W. A. Benjamin, Inc., New York-Amsterdam.
- STOCK, J., AND M. WATSON (2011): *Introduction to Econometrics (3rd edition)*. Addison Wesley Longman.
- VAN DER VAART, A. W. (1998): *Asymptotic Statistics*. Cambridge University Press.
- VAN DER VAART, A. W., AND J. A. WELLNER (1996): *Weak Convergence and Empirical Processes*. Springer Series in Statistics.
- WOOLDRIDGE, J. M. (2010): *Econometric Analysis of Cross Section and Panel Data*. Cambridge, Massachusetts: The MIT Press, second edn.