Entanglement and topological interfaces

E. Brehm, I. Brunner, D. Jaud and C. Schmidt-Colinet

Arnold Sommerfeld Center, Ludwig-Maximilians-Universität Theresienstrae 37, 80333 München, Germany E-mail: E.Brehm@physik.uni-muenchen.de, Ilka.Brunner@physik.uni-muenchen.de, Daniel.Jaud@physik.uni-muenchen.de,

schmidt.co@physik.uni-muenchen.de

ABSTRACT: In this paper we consider entanglement entropies in two-dimensional conformal field theories in the presence of topological interfaces. Tracing over one side of the interface, the leading term of the entropy remains unchanged. The interface however adds a subleading contribution, which can be interpreted as a relative (Kullback-Leibler) entropy with respect to the situation with no defect inserted. Reinterpreting boundaries as topological interfaces of a chiral half of the full theory, we rederive the left/right entanglement entropy in analogy with the interface case. We discuss WZW models and toroidal bosonic theories as examples.

Contents

1	Introduction	1
2	Topological Interfaces	4
3	Entanglement Entropy	7
	3.1 The replica trick and conformal interfaces	8
4	Entanglement Entropy through Topological Interfaces	9
5	Left/Right Entanglement Entropy	17
6	Results for Bosonic Tori	22
	6.1 EE through topological defects	22
	6.2 LREE of bosonic tori	24
7	Conclusion and Outlook	26
A	Entanglement entropy of a fusion product	28

1 Introduction

In the study of entanglement entropy we are often interested in universal terms, in particular if these terms depend only on a limited number of parameters in the quantum theory. For two-dimensional conformal field theories, formulae for such terms were found in the seminal work of [1] and [2]. Concretely, consider a critical system with a subsystem of dimensionless length L in some units. For large L, the entanglement entropy of this subsystem has the expansion

$$S_L = \frac{c}{3} \log L + S_{\rm sub} \,.$$
 (1.1)

Here, c is the central charge of the CFT, and the term S_{sub} comprises subleading terms, including in particular constant terms. In the context of statistical mechanics, the subleading terms will depend on the information of the full statistical system, not just its RG fixed point. In this sense, the subleading terms are non-universal, whereas the leading term depends only on basic data of the underlying CFT.

Besides systems defined on Riemann surfaces without boundaries, one area of investigation is centered around the entanglement in systems with boundaries or interfaces. There are several possibilities to specify subsystems, leading to different entanglement entropies. One possibility is to single out a spacial interval terminating at the boundary. If the subsystem of length L ends on a boundary specified by some boundary condition b, the expression for the entanglement entropy becomes [2]

$$S_L^{(b)} = \frac{c}{6} \log L + \log g_b + \frac{S_{\text{sub}}}{2}.$$
 (1.2)

Comparing with (1.1), the factor 1/2 in the overall coefficient of the leading term reflects the area law. The quantity g_b is the universal ground-state degeneracy [3] of the boundary condition b. In string theory, g_b defines the mass of the D-brane [4]. The important observation [2] is that $\log g_b$ in (1.2) is a universal contribution of the boundary. The other subleading terms are non-universal, where S_{sub} denotes the same terms as in the bulk case (1.1).

Via the folding trick, this result can also be applied to interfaces between two CFTs if the interface splits the system symmetrically. In [5–7], both boundary and interface entanglement entropy in this real-space case were investigated by AdS/CFT methods based on the Ryu-Takayanagi area law [8, 9].

Naturally, it is of interest to generalise the results on defect and boundary entropies further. For the case of interfaces, one would like to consider situations not constrained by the requirement of geometric reflection symmetry. For the case of boundaries, one would like to consider subsystems that are not specified by the geometry of the system, but by decomposing the Hilbert space into left- and right-movers [10, 11]. In this paper, we will discuss the entanglement entropy through the special class of topological interfaces, and show how the same techniques can be employed to determine the left/right entanglement entropy for boundaries.

The problem of entanglement through interfaces has been approached before in special examples, in particular for the case of free bosons in [12]. The interface splits the system into two parts, and one is interested in the entanglement entropy of the subspace on one side of the interface. The same method of computation was used in [13] to determine the entanglement across defects of the Ising model, giving in particular a CFT computation of results obtained earlier in [14, 15]. A recent investigation from the AdS/CFT point of view of these setups can be found in [16], see [17, 18] for further results in this direction.

In these examples, it was observed that the entanglement entropy in the presence of interfaces receives two kinds of modifications to the pure bulk expression (1.1). The first one is a reduction of the coefficient of the universal logarithmic term of the bulk theory. This correction depends on the "strength" of the interface, which can be defined using its transmissivity or reflectivity [19]. The other modification is a universal contribution to the constant part of the bulk term. This shift was observed to be independent of the transmissivity, and indeed only depends on the topological information contained in the interface operator.

In this paper we will study the constant shifts of the entanglement entropy in the presence of interfaces in some more detail and generality. Our discussion is restricted to the case of topological interfaces, preserving the full conformal invariance of the bulk theory. Topological interfaces are tensionless domain walls of two-dimensional CFTs and can be moved freely on the two-dimensional space-time as long as no other interfaces or operator insertions are crossed. They add interesting structure to any CFT, as they can for example merge smoothly, defining a product structure on the space of interfaces. Topological interfaces define maps between the Hilbert spaces of the CFTs on the two sides of the interface. These maps in particular intertwine the action of the Virasoro algebra, as a consequence of the fact that the interface is topological. We will consider the case where the interface also intertwines the action of higher symmetry generators.

Of particular interest are cases where the higher symmetry renders the theory rational. Tracing over one side of the topological interface, we show that one obtains a natural reduced density matrix. Strictly speaking, the associated probability distribution is determined by the fusion product of the interface with its conjugate. This combination can in particular be regarded as a map from a single CFT to itself. In the case of a rational theory, the probability distribution acts on the space of irreducible representations appearing in the decomposition of the bulk Hilbert space of that theory with respect to the symmetry algebra. We compute the entanglement entropy through the interface and show that it is given by the negative of the Kullback-Leibler entropy relative to the probability distribution associated to the trivial (invisible) interface.

The Kullback-Leibler divergence of one probability distribution with respect to another is always nonnegative, and vanishes only if the two probability distributions agree. We discuss what this means in terms of the interface. The map between interfaces and probability distributions is not one-to-one, there is in general more than one interface corresponding to a single probability distribution. In particular, not only the trivial interface leads to a vanishing subleading contribution to the entanglement entropy, but so do all interfaces corresponding to symmetries. These interfaces map states to symmetry-transformed states, and therefore do not lead to any information loss. This minimum of the information loss can only be saturated if the partition functions of the CFTs on the two sides of the interface agree and the fusion product of the interface with its dual is the identity. For the other cases, we will derive a bound on the subleading term of the entanglement entropy.

We apply our result to the $su(2)_k$ WZW models in the limit of large levels k, where the theory approaches a sigma-model on S^3 with H-flux. Here, the probability distributions associated to topological defects become continuous. We compute the shift in entanglement entropy for elementary defects in the large-k limit. For generic elementary defects, we obtain rational numbers approaching 1. In certain distinguished cases however the Kullback-Leibler divergence develops sharp peaks that can be attributed to divisibility properties involving the level k.

Furthermore, we slightly extend the framework of rational CFTs and discuss also toroidal compactifications of free bosons. Here, the situation is particularly simple, as all topological interfaces fall into the class of "duality interfaces" introduced in [20]. The defining property of this class of interfaces is that fusion with the orientiation reversed interface yields a superposition of symmetry defects. Thus, the insertion of this superposition amounts to an orbifold-like projection of the initial theory. When one associates a probability, non-invariant states are assigned probability 0, whereas all invariant states are distributed equally. This amounts to a shift in the entanglement entropy by the logarithm of the number of symmetry defects appearing in the fusion. For the free boson theory, this contribution can be identified as the index of a sublattice of the winding and momentum lattice to which the interface couples. This index appears also in the g-factor of the interface, such that in the special case of bosonic theories the shift in the entanglement entropy agrees with $\log g^2$. Note that the shift vanishes for duality defects implementing T-duality transformations, which is as expected since there is no information loss due to T-duality.

Besides the entanglement across topological defects we consider entanglement entropies between non-geometrical subsectors for boundaries. As mentioned before, in the presence of a boundary a suggestive subsector of the space of states is provided by the chirality of the symmetry algebra. Since the boundary gluing conditions couple the left- and the rightmoving degrees of freedom, the idea is to trace out the holomorphic or antiholomorphic sector of the space of states. The authors of [10] studied this left/right entanglement entropy for a free boson CFT. Results for general rational CFTs and an interpretation in a gravitational context were obtained in [11].

In this paper, we give an alternative derivation of the results of [11]. For this, we reinterpret the conformal boundary as a defect-like object on the full plane. To do so, we start with a theory on the upper half-plane with a boundary condition along the real line. We employ the doubling trick to fold the dependence on anti-holomorphic coordinates to the lower half plane. The boundary condition on the boundary state can then be interpreted as an intertwining property for an interface operator associated to the boundary condition. The computation of the entanglement entropy then resembles the one for the interface case. However, the interpretation in terms of a Kullback-Leibler divergence is lost, as there is no natural trivial boundary condition that could serve as a reference point.

Again, we consider su(2) WZW models and toroidal theories as examples. In the special case of tori, the left/right entanglement entropy is given in terms of the g-factor.

The plan of the paper is as follows. In section 2 we recall some basics of topological interfaces in two-dimensional CFTs, and fix some notation. In section 3 we briefly review the method of [1, 12] for the computation of the entanglement entropy through conformal interfaces. We then turn to the case where the interface is topological in section 4. Here, we work out the relation between entanglement entropy and Kullback-Leibler divergence, and discuss the rational case in detail. The relevance of our formulae to the left/right entanglement entropy (LREE) is discussed in section 5. Finally, we turn to the case of free bosons compactified on tori in section 6. We determine the entanglement entropy through topological interfaces as well as the left/right entanglement entropy in this case, which slightly extends the rational framework in a simple example.

2 Topological Interfaces

The study of one-dimensional interfaces between two-dimensional CFTs has a long history [21, 22]. Locally, an interface sets gluing conditions for all pairs of local fields separated by the interface. In that sense an interface defines a map between the algebras of local fields on the two sides. Similar to a boundary condition, such an interface condition admits local excitations and constitutes a one-dimensional subsector of the full quantum field theory. At a conformal fixed point the interface preserves at least one half of the bulk conformal charges. If the interface runs along the real axis of the complex plane and separates CFT1 from CFT2, the condition reads

$$\lim_{y \searrow 0} (T^{(1)}(x+iy) - \tilde{T}^{(1)}(x-iy)) = \lim_{y \nearrow 0} (T^{(2)}(x+iy) - \tilde{T}^{(2)}(x-iy)), \qquad (2.1)$$

where $T^{(n)}$ and $\tilde{T}^{(n)}$ are the holomorphic and antiholomorphic components of the energymomentum tensor of CFT*n*. The requirement (2.1) is a necessary local condition. Similarly as in the case of boundary conditions, further local conditions follow from sewing relations, and global conditions arise from modular constraints on the torus.

In some sense, conformal interfaces generalise the notion of conformal boundary conditions, which are the special solutions where both sides of (2.1) are equal to zero, or occur if one side of the interface is trivial. Another set of special solutions to (2.1) is obtained when the interface commutes with both the left- and the right-moving Virasoro algebra, such that (2.1) is solved separately for the holomorphic and the antiholomorphic component of the energy-momentum tensor. Evidently, this can only happen when the theories on the two sides have equal left- and right-moving central charges. An interface corresponding to such a solution can be freely deformed and moved on the Riemann surface, as long as it does not cross any operator insertions. These interfaces were dubbed topological in [23]. Initially introduced in [24], topological interfaces have been studied in particular in rational CFTs (see *e.g.* [20, 25] and references therein).

A topological interface can be regarded as an operator on the space of states, acting as a constant map between (left-right pairs of) isomorphic Virasoro representations. In the case where the conformal symmetry is enhanced to a larger chiral symmetry algebra, the topological interface condition may or may not respect the additional symmetry. Topological interfaces between CFT1 and CFT2 thus naturally fall into classes corresponding to the preserved common symmetry subalgebra. For a given topological interface, we consider the decomposition of the space of states of CFTn (n = 1, 2) with respect to this common subalgebra,

$$\mathcal{H}^{n} = \bigoplus_{(i\bar{i})} M^{n}_{i\bar{i}} \mathcal{H}_{i} \otimes \mathcal{H}_{\bar{i}}.$$
(2.2)

The indices i and $\bar{\imath}$ label (generally different) irreducible highest weight representations of the two chiral parts of the common subalgebra. The non-negative integers $M_{i\bar{\imath}}^n$ give the multiplicities of the pair of representations $(i, \bar{\imath})$. We will assume that our theories are unitary and have a discrete spectrum of highest weight states of the chiral subalgebra, and that there is a unique vacuum state. For the following discussion we will also assume that the modular S transformation of representation characters is given by a discrete matrix.

An operator corresponding to a general topological interface will then be denoted

$$\mathcal{I}_A = \sum_{\mathbf{i}} d_{A\mathbf{i}} \|\mathbf{i}\|.$$
(2.3)

Generally, we will use capital indices to label the interfaces. We use bold-face indices **i** to refer to a pair of left-right products of irreducible representations in the two adjacent CFTs,

$$\mathbf{i} \equiv (i, \bar{\imath}; \alpha, \beta) \,. \tag{2.4}$$

Here, (i, \bar{i}) labels the transmitted pair of representations. The indices $\alpha = 1, 2, \ldots, M_{i\bar{i}}^1$ and $\beta = 1, 2, \ldots, M_{i\bar{i}}^2$ are the multiplicity labels of this pair on the two sides of the interface. The symbol $\|\mathbf{i}\|$ in (2.3) denotes the Ishibashi-type projector which acts as an intertwiner between the two pairs of representations, *i.e.*

$$\|\mathbf{i}\| : \mathcal{H}_i \otimes \mathcal{H}_{\bar{i}}^{(\alpha)} \to \mathcal{H}_i \otimes \mathcal{H}_{\bar{i}}^{(\beta)}$$
(2.5)

and

$$J_n \|\mathbf{i}\| = \|\mathbf{i}\| J_n , \qquad (2.6)$$

where J_n denotes any symmetry generator. Note that in general one can include an automorphism of the extended symmetry algebra into the action of the interface; this generalisation is straightforward.

One important property of topological interfaces is that they admit a fusion product. The fusion product has the geometric interpretation of moving the interface lines on top of each other, and interpreting the result as a topological interface between the two remaining CFTs. While fusion may also be defined for the more general conformal interfaces, it is particularly straightforward in the topological case, where it basically consists of map composition [24]. When writing the coefficients d_{ABi} of the fusion product $\mathcal{I}_{AB} = \mathcal{I}_A \mathcal{I}_B$ we will suppress the summation over multiplicity labels,

$$d_{AB\mathbf{i}} = \sum_{\gamma} d_{A(i\bar{\imath};\alpha,\gamma)} d_{B(i\bar{\imath};\gamma,\beta)} \equiv d_{A\mathbf{i}} d_{B\mathbf{i}} \,. \tag{2.7}$$

The fusion product extends to fusion of a topological interface with (non-topological) conformal interfaces, in particular with boundary conditions.

Due to invariance under modular S transformation, the coefficients d_{Ai} must satisfy [24]

$$\sum_{\mathbf{i}} S_{ij} S_{\bar{\imath}\bar{\jmath}} \operatorname{Tr} d_{A^* \mathbf{i}} d_{A\mathbf{i}} = \mathcal{N}_{j\bar{\jmath}A}^{\ A} \in \mathbb{N}_0, \qquad (2.8)$$

where A^* labels the orientation reversed interface with corresponding defect operator $\mathcal{D}_{A^*} = \mathcal{D}_A^{\dagger}$, S_{ij} is an element of the modular S matrix, and the trace is over multiplicity labels. This is the analogue of the Cardy condition for conformal boundary states. The $\mathcal{N}_{j\bar{j}}{}^B_A$ count the multiplicity of the pair of representations (j,\bar{j}) in a system where the topological interfaces intersect a spatial slice, *i.e.* for a time evolution parallel to the interfaces. The condition (2.8) restricts the possible values of coefficients d_{Ai} , and it also requires that linear superpositions of interfaces must have integer coefficients. We refer to interfaces which cannot be decomposed into a superposition of other interfaces with positive coefficients as 'elementary'. The set of elementary interfaces forms a basis for all topological interfaces of the same class. Obviously any interface for which at least one of the $\mathcal{N}_{j\bar{j}A}$ is equal to 1 is elementary. In fact, due to the operator-state correspondence in the theory on the interface any elementary interface has at least $\mathcal{N}_{00}{}^A_A = 1$, *i.e.* the vacuum in parallel time evolution occurs with multiplicity 1.

Consider a set of topological interfaces \mathcal{I}_A for which $d_{A\mathbf{i}}$ provides a unitary transformation from projectors $\|\mathbf{i}\|$ to the \mathcal{I}_A . It can be shown [24] that the corresponding $\mathcal{N}_{i\bar{i}A}^{\ B}$ form a representation of a tensor product of fusion algebras,

$$\sum_{B} \mathcal{N}_{i\bar{\imath}A}{}^{B} \mathcal{N}_{j\bar{\jmath}B}{}^{C} = \sum_{\mathbf{k}} N_{ij}{}^{k} N_{\bar{\imath}\bar{\jmath}}{}^{\bar{k}} \mathcal{N}_{k\bar{k}}{}^{C}_{A}.$$
(2.9)

In the last formula, the N_{ij}^{k} are the fusion rules of the chiral algebra. It is easy to see that a topological interface \mathcal{I}_{A} in such a set is elementary.

A particular instance where we know a set of d_{Ai} that provides a change of basis occurs in rational CFT, *i.e.* in theories where the index set $\{i\}$ in (2.2) is finite. The simplest case are the diagonal theories — theories which are charge conjugation invariant $(i = \bar{i})$, and where the multiplicities for all chiral algebra representations are 1. In such a theory there are topological defects¹ of the form

$$\mathcal{D}_{\mathbf{a}} = \sum_{i} \frac{S_{ai}}{S_{0i}} \left\| i \right\|.$$
(2.10)

These defects have $\mathcal{N}_{0a}^{\ a} = N_{0a}^{\ a} = 1$ and are therefore elementary. They provide a basis for the set of topological defects which respect the chiral symmetry.

In cases where the chiral algebra admits a global symmetry G, we find among the topological interfaces the so-called symmetry defects. Each element $g \in G$ can be associated to a topological defect \mathcal{D}_g . By definition, these interfaces glue any field to its image under the symmetry operation. Hence, they implement an action of G through

$$\mathcal{D}_{q}^{\dagger} = \mathcal{D}_{q^{-1}}, \qquad \mathcal{D}_{g}\mathcal{D}_{h} = \mathcal{D}_{gh} \qquad \forall h, g \in G.$$
(2.11)

A broader class of interfaces are the duality interfaces introduced in [20]. Their defining property is that

$$\mathcal{I}\mathcal{I}^{\dagger} = \bigoplus_{g \in G} \mathcal{D}_g \,, \tag{2.12}$$

where G is a finite symmetry group of the CFT. The fusion product of a duality interface with its adjoint contains a superposition of group-like defects corresponding to a symmetry (sub-)group. Duality defects were first introduced in the context of RCFT, where they can be used to relate CFTs with the same chiral algebra but different modular invariants. However, the definition can be extended also to the non-rational context. Prominent examples for duality interfaces implement dualities such as T-duality in free field theories, or the Kramers-Wannier duality in the Ising model [26].

3 Entanglement Entropy

Entanglement entropy measures quantum correlation between subsystems. Let $\rho = |\psi\rangle\langle\psi|$ be the density matrix of a system in a pure quantum state $|\psi\rangle$. Let the Hilbert space be a direct product $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$ where A and B are the subsystems. The reduced density matrix of A is $\rho_A = \text{Tr}_B \rho$. The *entanglement entropy* is the corresponding von Neumann entropy

$$S_A = -\operatorname{Tr} \rho_A \log \rho_A \,. \tag{3.1}$$

¹In this paper we usually refer to interfaces as defects if the CFTs on the two sides are identical.

 $S_{\rm B}$ is defined analogously. For the density matrix of a pure quantum state one always has $S_{\rm A} = S_{\rm B}$. In the simplest case the pure state is the ground state $|0\rangle$ of the system. We then refer to the corresponding quantity (3.1) as the ground state entanglement entropy.

One way to compute (3.1) makes use of the replica trick [2]. The theory is considered on K copies of the original Riemann surface, glued together along the subsystem A in a cyclic fashion. Tracing over all copies of the subsystem B, the reduced density matrix becomes ρ_A^K . Its trace in the K-sheeted Riemann surface can be written as

$$\operatorname{Tr}_{\mathcal{A}} \rho_{\mathcal{A}}^{K} = \frac{Z(K)}{Z(1)^{K}}, \qquad (3.2)$$

where Z(K) is the full partition function on the K-sheeted Riemann surface. One analytically continues this expression to complex values of K, and obtains the entanglement entropy (3.1) from

$$S_{\mathcal{A}} = (1 - \partial_K) \log Z(K)|_{K \to 1}.$$

$$(3.3)$$

3.1 The replica trick and conformal interfaces

In the following, we will briefly review a construction of Z(K) due to [1, 12], which in principle allows to derive the entanglement entropy through general conformal interfaces connecting two conformal field theories. The same construction was also used in [13].

Consider a conformal interface \mathcal{I} along the imaginary axis of the complex plane, with CFT1 on Re w > 0 and CFT2 on Re w < 0. With time flowing along the defect line, the subsystems A and B consist of the positive and negative real axis, respectively. Following the replica trick, the corresponding K-sheeted Riemann surface consists of K copies of the complex plane, glued together cyclically along a branch cut on the positive real axis, as illustrated on the left of Figure 1.

In order to evaluate the partition function Z(K) we introduce the cutoffs $|w| = \epsilon$ and |w| = L and change coordinates to $z = \log w$. Observe that this transformation is compatible with (2.1). The resulting cylinder is illustrated on the right of Figure 1. As in [12] we regularise the partition function by imposing periodicity in Re z and choosing $\epsilon = \frac{1}{L}$. Periodicity can be imposed since the cut-offs L and ϵ are very large and very small, respectively. In these limits the result does not depend on the specific choice of boundary condition. The identification $\epsilon = 1/L$ is somewhat arbitrary. It will lead to a factor of two in the final result for the leading term of the entanglement entropy. This will have the benefit that the bulk term will have the familiar form $c/3 \log L$, even though since L is only an IR cut-off, the 'interval' is actually physically a half line with only one end point, in which case the entanglement entropy should be reduced by a factor 1/2 due to the area law. On the resulting torus one observes that the shape of the defects is unaltered under the global conformal transformation that changes the time evolution parallel to the interfaces to the one flowing orthogonally to the interfaces. We conclude that Z(K) is given by a torus partition function with 2K interfaces inserted,

$$Z(K) = \operatorname{Tr}_{1} \left(\mathcal{I}^{\dagger} e^{-\delta H_{2}} \mathcal{I} e^{-\delta H_{1}} \cdots \mathcal{I} e^{-\delta H_{1}} \right)$$

= $\operatorname{Tr}_{1} \left(\mathcal{I}^{\dagger} e^{-\delta H_{2}} \mathcal{I} e^{-\delta H_{1}} \right)^{K},$ (3.4)

where H_1 and H_2 are the Hamilton operators in the respective CFT, and

$$\delta = \frac{2\pi^2}{\log L/\epsilon} = \frac{\pi^2}{\log L} \,. \tag{3.5}$$

Obviously the evaluation and analytic continuation of (3.4) depends heavily on \mathcal{I} . For non-topological conformal defects, Z(K) is in general very hard to compute. An explicit expression which permitted the computation of the entanglement entropy was obtained in [12] for the case of a single free boson, and in [13] for conformal defects of the free fermion and the Ising model. However, for topological defects the expression for Z(K) simplifies considerably, as we will see in the following.

There is one feature of the entanglement entropy as we define it here which is rather obvious already at this stage. From (3.4) it is easy to see that (3.3) is invariant under any rescalings of the interface. While interfaces generically have a standard normalisation derived from their properties under modular transformations, this means in particular that superpositions $M\mathcal{I}$ of identical interfaces \mathcal{I} yield the same entanglement as a single \mathcal{I} .



Figure 1. Sketch of the K-sheeted Riemann surface we use in the replica trick. After imposing an UV cutoff ϵ , a IR cutoff L, the surface corresponds to the cylinder on the upper right. To derive Z(K) we impose periodicity also in the direction of the real part of z to obtain a torus.

4 Entanglement Entropy through Topological Interfaces

In the limit of a large IR cutoff L, or equivalently $\delta \ll 1$, the entanglement entropy through a topological interface \mathcal{I} follows from the torus partition function (3.4) in a rather straightforward way. The torus partition function includes K insertions of \mathcal{I} and of its adjoint \mathcal{I}^{\dagger} , which commute with both Virasoro algebras and therefore in particular with the Hamiltonian $H \propto L_0 + \tilde{L}_0$. Thus we can write

$$Z(K) = \operatorname{Tr}\left(\mathcal{I}e^{-\delta H}\mathcal{I}^{\dagger}e^{-\delta H}\right)^{K} = \operatorname{Tr}\left((\mathcal{I}\mathcal{I}^{\dagger})^{K}e^{-2\delta HK}\right).$$
(4.1)

For the general topological interface (2.3) we find

$$Z(K) = \sum_{(i,\bar{i})} \operatorname{Tr} \left(d_{A\mathbf{i}} \, d_{A^*\mathbf{i}} \right)^K \chi_i \left(e^{-2\delta K} \right) \chi_{\bar{i}} \left(e^{-2\delta K} \right) \,, \tag{4.2}$$

where $\chi_i(q)$ is the character of the representation *i*. In (4.2) and in the remainder of this section, Tr denotes the trace over multiplicity indices. Applying a modular *S* transformation we obtain

$$Z(K) = \sum_{(i,\bar{i})} \sum_{j,\bar{j}} \operatorname{Tr} \left(d_{A\mathbf{i}} \, d_{A^*\mathbf{i}} \right)^K S_{ij} S_{\bar{\imath}\bar{\jmath}} \, \chi_j \left(e^{-\frac{2\pi^2}{\delta K}} \right) \chi_{\bar{\jmath}} \left(e^{-\frac{2\pi^2}{\delta K}} \right) \,. \tag{4.3}$$

In the limit $\delta \ll 1$ only the vacuum with the energy $E_0 = -\frac{c}{12}$ will contribute to the sum. The contribution of every other state in the theory is exponentially suppressed. Therefore the partition function is approximately given by

$$Z(K) \approx \underbrace{\sum_{(i,\bar{i})} \operatorname{Tr} \left(d_{A^* \mathbf{i}} \, d_{A \mathbf{i}} \right)^K S_{i0} S_{\bar{i}0} \, e^{\frac{\pi^2 c}{6\delta K}} \,. \tag{4.4}$$
$$\underbrace{=}_{\equiv A(K)}$$

The factor A(K) contains the information about the topological interface. For the entanglement entropy we then obtain

$$S = (1 - \partial_K) \log Z(K) \big|_{K \to 1}$$

$$\approx (1 - \partial_K) \left(\frac{\pi^2 c}{6\delta K} + \log A(K) \right) \Big|_{K \to 1}$$

$$= \frac{c}{3} \log L + \left[\log A(1) - \frac{A'(1)}{A(1)} \right].$$
(4.5)

In the last line we have used (3.5), and a prime denotes the derivative with respect to K. Note that time in the channel described in (4.3) runs parallel to the interface. From (2.8) we therefore find that

$$A(1) = \mathcal{N}_{\mathbf{0}A}^{\ A} \tag{4.6}$$

is a non-negative integer. It is the multiplicity of the vacuum representation in the twisted torus partition function in the channel where time evolves along the interface and its conjugate. If the interface is elementary we have A(1) = 1. For the derivative of A(K) one obtains

$$A'(1) = \sum_{(i,\bar{\imath})} S_{i0} S_{\bar{\imath}0} \operatorname{Tr} \left(d_{A^* \mathbf{i}} d_{A \mathbf{i}} \right) \log \left(d_{A^* \mathbf{i}} d_{A \mathbf{i}} \right) \,.$$

Inserting this in (4.5), the entanglement entropy becomes

$$S = \frac{c}{3} \log L + \left[\log \mathcal{N}_{\mathbf{0}A}^{\ A} - \frac{1}{\mathcal{N}_{\mathbf{0}A}^{\ A}} \sum_{(i,\bar{\imath})} S_{i0} S_{\bar{\imath}0} \operatorname{Tr} \left(d_{A^* \mathbf{i}} \, d_{A\mathbf{i}} \right) \log \left(d_{A^* \mathbf{i}} \, d_{A\mathbf{i}} \right) \right] \,. \tag{4.7}$$

Within CFT1 we now define

$$p_{(i\bar{\imath},\alpha\alpha')}^{A} = \frac{d_{A^{*}\mathbf{i}} d_{A\mathbf{i}} S_{i0} S_{\bar{\imath}0}}{\mathcal{N}_{\mathbf{0}A}^{A}}, \qquad (4.8)$$

where the multiplicity labels α and α' both run from 1 to $M_{i\bar{\imath}}^1$. For every pair $(i, \bar{\imath})$, the matrix $p_{\mathbf{i}}^A \equiv p_{(i\bar{\imath},\alpha\alpha')}^A$ is a positive-semidefinite Hermitian matrix², *i.e.* the eigenvalues of the $p_{\mathbf{i}}^A$ are real and positive. Moreover, by (2.8) we have

$$\sum_{(i,\bar{i})} \operatorname{Tr} p_{\mathbf{i}}^{A} = 1.$$
(4.9)

The set of all eigenvalues therefore forms a probability distribution. In a quantisation where time runs orthogonally to the interface, the value of $\text{Tr } p_i^A$ is the probability of finding the system CFT1 in the Ishibashi-type state associated to the sector (i, i), after tracing out CFT2.³ Such a state is thermal within its sector, and the set of p_i^A should therefore be understood as defining a reduced density matrix. Observe that the distribution corresponding to the identity defect in CFT1 is given by

$$p_{\mathbf{i}}^{id} = S_{i0} S_{\bar{\imath}0} \,\delta_{\alpha\alpha'} \,, \qquad \alpha, \alpha' = 1, \, 2, \dots, M_{i\bar{\imath}}^1 \,.$$

$$(4.10)$$

Equation (4.7) can now be written as

$$S = \frac{c}{3} \log L - \sum_{(i,\bar{i})} \operatorname{Tr} p_{\mathbf{i}}^{A} \log \frac{p_{\mathbf{i}}^{A}}{p_{\mathbf{i}}^{id}}.$$
 (4.11)

This is our main result of this section. The quantity

$$s(\mathcal{I}_A) := -\sum_{(i,\vec{\imath})} \operatorname{Tr} p_{\mathbf{i}}^A \log \frac{p_{\mathbf{i}}^A}{p_{\mathbf{i}}^{id}}$$
(4.12)

is the negative of the relative entropy — the Kullback-Leibler divergence [28] — of the probability distribution associated to \mathcal{I}_A on the CFT1 side, measured with respect to the probability distribution associated to the identity defect \mathcal{D}_{id} of CFT1. One interpretation of this quantity is the amount of information lost when the probability distribution is wrongly assumed to be given by \mathcal{D}_{id} , while it is in reality given by \mathcal{I}_A .⁴

The relative entropy is always non-negative, and vanishes only if the compared probability distributions agree.⁵ Therefore we have $s(\mathcal{I}_A) \leq 0$, which corresponds to the intuition that an interface cannot enhance the transmissivity beyond the one of the identity defect in CFT1. We have $s(\mathcal{I}_A) = 0$ if and only if $p^A = p^{\text{id}}$. This is the case precisely if $d_{A^*i}d_{Ai}$ is the identity matrix for all pairs of representations (i, i) which appear in CFT1. A necessary requirement for the existence of an interface with this property is that the representation multiplicities of CFT2 must not be smaller than those of CFT1. Since both CFTs are

²Recall that in unitary theories $S_{i0} > 0$

 $^{^{3}}$ For an interpretation along these lines in terms of a three-dimensional topological field theory see [27].

⁴The fact that we find a relative entropy ties in nicely with the results [29]. There it was observed that in higher dimensions, a perturbation of the shape of the entangled region leads to a shift in the entanglement entropy given by the Kullback-Leibler divergence of the probability distributions defined by the reduced density matrices before and after the perturbation. While the shape does not play a role in our setup, the shift in the entanglement entropy is again associated with the difference of the involved density matrices.

⁵Continuous distributions have to agree almost everywhere.

unitary and have a single vacuum state, modular invariance in fact forces CFT1 and CFT2 to have identical spectra. Since the necessary condition $d_{A^*i}d_{Ai} = 1$ then means that the fusion product of the defect and its conjugate is the identity, we have

$$\exists \mathcal{I} : s(\mathcal{I}) = 0 \quad \Leftrightarrow \quad Z_{\text{CFT1}} = Z_{\text{CFT2}} \quad \text{and} \quad \mathcal{I}^{\dagger} \mathcal{I} = \mathcal{D}_{id} \quad \text{in CFT1}.$$
(4.13)

For general CFT1 and CFT2 we may give a simple upper bound for s, based on the restricted data we have been employing so far. Without loss of generality every interface \mathcal{I} between CFT1 and CFT2 can be associated with a set of diagonal matrices p_i . Each of these matrices p_i has at most $T_{i\bar{\imath}} = \min(M_{i\bar{\imath}}^1, M_{i\bar{\imath}}^2)$ eigenvalues different from 0. Varying the remaining eigenvalues we look for the maximal value of s under the linear constraint (4.9). This is only one constraint out of the set (2.8), such that this calculation will obviously lead to an upper bound. A maximal value of s would be achieved for the distribution

$$p_{\mathbf{i}} = \operatorname{diag}(p_{(i\bar{\imath},1)}, \dots, p_{(i\bar{\imath},T_{i\bar{\imath}})}, 0, \dots, 0) \quad \text{with} \quad p_{(i\bar{\imath},\alpha)} = \frac{S_{i0}S_{\bar{\imath}0}}{\sum_{(j,\bar{\jmath})} T_{j\bar{\jmath}}S_{j0}S_{\bar{\jmath}0}} \,. \tag{4.14}$$

This distribution yields the upper bound

$$s \le \log\left(\sum_{(i,\bar{\imath})} T_{i\bar{\imath}} S_{i0} S_{\bar{\imath}0}\right) \,. \tag{4.15}$$

The bound is strictly smaller than zero if there is at least one (i, i) with $T_{i\bar{\imath}} < M_{i\bar{\imath}}^1$. As we have seen above, this is equivalent to having at least one pair (i, i) where $M_{i\bar{\imath}}^1 \neq M_{i\bar{\imath}}^2$. The bound (4.15) is zero if and only if the theories CFT1 and CFT2 have the same spectrum. In cases where the CFTs on the two sides are identical, the distribution (4.14) is in particular obtained from the identity defect.

We emphasise that different interfaces can lead to the same distribution (4.8), and thus to the same entanglement entropy. In particular, fusing any interface on either side with a symmetry defect of the respective theory will leave the distribution unaltered. The reference distribution p^{id} of (4.10) is therefore also obtained from any symmetry defect in CFT1. On the other hand, every defect whose fusion product with a particular topological interface leaves the probability distribution of the interface unaltered is a symmetry defect.

The distributions also do not change if we superpose the same interface multiple times. This is obvious from the interpretation of the probability distribution mentioned above. In agreement to our remark in section 3.1, an interface \mathcal{I} formally has the same probability distribution as $M\mathcal{I}$ for any rescaling $M \in \mathbb{C}^*$, and therefore in particular for superpositions of the same interface. However, the change in the entanglement entropy is difficult to compute for general superposition and fusion. This is so because it is in general difficult to see how closely the probability distribution of the resulting interface follows p_i^{id} .

For concreteness, let us now consider the defects (2.10) in a rational theory with diagonal modular invariant. By (4.8), the interfaces (2.10) of diagonal RCFTs lead to a probability distribution

$$p_i^a = |S_{ia}|^2 \,. \tag{4.16}$$

From our result (4.11) we therefore obtain the entanglement entropy

$$S = \frac{c}{3} \log L - \sum_{i} |S_{ia}|^2 \log \left| \frac{S_{ia}}{S_{i0}} \right|^2.$$
(4.17)

Example 1: Duality interfaces

As a class of examples we consider the duality interfaces (2.12). Here \mathcal{II}^{\dagger} projects the theory onto a sector invariant under a symmetry group G. Invariant states pick up a constant prefactor of |G|, the order of the group. On the level of equation (4.1) this means that

$$Z(K) = \operatorname{Tr}\left((\mathcal{II}^{\dagger})^{K} e^{-2\delta HK}\right) = \operatorname{Tr}\left((\bigoplus_{g \in G} \mathcal{D}_{g})^{K} e^{-2\delta HK}\right) = |G|^{K} \operatorname{Tr}_{inv}\left(e^{-2\delta HK}\right),$$
(4.18)

where in the last line the trace is taken only over the invariant subsector of the initial Hilbert space. This partition function is a projection of an initial partition function, which is in line with the fact that correlators of invariant fields in orbifold theories are obtained by projection from the initial theory. The prefactor $|G|^K$ will drop out in the calculation of the entanglement entropy, so that effectively we consider the entanglement of an initial system with a projected system. However, in comparison with the system with only the trivial defect inserted, the projection contains a factor of $|G|^{-1}$ that leads to a shift in the entanglement entropy for duality interfaces. The entanglement entropy in the presence of such a duality interface is therefore

$$S = \frac{c}{3} \log L - \log |G|.$$
 (4.19)

In terms of the probability distributions introduced earlier we find for the duality defects

$$p_{\mathbf{i}}^{duality} = p_{\mathbf{i}}^{id}|G|$$
 for **i** invariant, $p_{\mathbf{i}}^{duality} = 0$ otherwise . (4.20)

The shift in the entanglement entropy encodes the information loss under a projection.

Example 2: Ising model

The critical Ising model is described by three primaries id, ϵ , σ . It is an example of a diagonal rational theory. The S matrix of the Ising model is given by

$$S_{ij} = \frac{1}{2} \begin{pmatrix} 1 & 1 & \sqrt{2} \\ 1 & 1 & -\sqrt{2} \\ \sqrt{2} & -\sqrt{2} & 0 \end{pmatrix}, \quad \text{with } i, j \in \{id, \epsilon, \sigma\}.$$
(4.21)

The three elementary topological defects of the Ising model are therefore

$$\mathcal{D}_{id} = \|id\| + \|\epsilon\| + \|\sigma\|,$$

$$\mathcal{D}_{\epsilon} = \|id\| + \|\epsilon\| - \|\sigma\|,$$

$$\mathcal{D}_{\sigma} = \sqrt{2}\|id\| - \sqrt{2}\|\epsilon\|.$$

The defect corresponding to the vacuum *id* is the identity defect. The defect \mathcal{D}_{ϵ} is a symmetry defect implementing the \mathbb{Z}_2 symmetry of the Ising model. The presence of these two defects does not result in a shift of the entanglement entropy. The third defect \mathcal{D}_{σ} implements Kramers-Wannier duality. It satisfies the fusion rules

$$\mathcal{D}_{\sigma}\mathcal{D}_{\sigma} = \mathcal{D}_{id} + \mathcal{D}_{\epsilon} \,. \tag{4.22}$$

From our formula (4.17) we deduce that the entanglement entropy of \mathcal{D}_{σ} is

$$S(\sigma) = \frac{c}{3} \log L - \log 2,$$
 (4.23)

which also agrees with the result (4.19) for duality interfaces where the order of the group is 2. The result also reproduces the constant shift in the entanglement entropy observed in [13].

Example 3: $su(2)_k$ interfaces and the large k limit

The diagonal WZW model based on the chiral algebra su(2) at level k has irreducible representations labelled by half-integer spins s. Using the index convention i = 2s, the integer label i runs from 0 to k. The modular S matrix is given by

$$S_{ij} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{\pi(i+1)(j+1)}{k+2}\right) \,. \tag{4.24}$$

By (4.17), the entanglement entropy in the presence of an elementary defect \mathcal{D}_a of the form (2.10) reads

$$S(\mathcal{D}_a) = \frac{c}{3} \log L - \frac{2}{k+2} \sum_{i=0}^{k} \sin^2 \left(\frac{\pi(a+1)(i+1)}{k+2} \right) \log \frac{\sin^2 \left(\frac{\pi(a+1)(i+1)}{k+2} \right)}{\sin^2 \left(\frac{\pi(i+1)}{k+2} \right)}.$$
 (4.25)

Note that the defect \mathcal{D}_k does not change the entanglement entropy, since this defect simply implements the \mathbb{Z}_2 -symmetry acting on the representation labels as $a \to k - a$.

At large k one obtains the WZW model based on su(2). The central charge is c = 3, and the model can be presented in terms of three bosons on a target space S^3 with nonvanishing H-flux at large radius. The Z₂-symmetry corresponds to the reflection symmetry of the three-sphere. At any k, the theory contains elementary defects \mathcal{D}_a for non- negative integers $a \leq k$. To find their geometric interpretation, we recall a few facts on the interpretation of symmetry preserving boundary states. Quite generally, symmetry preserving D-branes on group manifolds wrap conjugacy classes [30, 31], which can be automorphismtwisted. In particular, the symmetry preserving (Cardy-)states of a WZW model wrap ordinary conjugacy classes of the underlying group G. To give an interpretation to defects, we first use the folding trick to map defects to permutation boundary conditions for the WZW model based on $G \times G$. Geometrically, these branes wrap twisted conjugacy classes where the automorphism is the permutation of the two factors, and the conjugacy class of $(g_1, g_2) \in G \times G$ takes the form [32]

$$C_{\omega}(g_1, g_2) = \left\{ (h_1^{-1}g_1h_2, h_2^{-1}g_2h_1) \, | \, h_1 \in G_1, h_2 \in G_2 \right\}.$$
(4.26)

The multiplication map $m: G \times G \to G$ maps these conjugacy classes to the conjugacy classes in the diagonal G. Indeed, the twisted conjugacy classes of $G \times G$ correspond precisely to the pre-images of the conjugacy classes of G under the multiplication map [32]. In the case of SU(2) they take the form $S^3 \times S^2$, as the regular untwisted conjugacy classes of SU(2) are generically isomorphic to S^2 . The conjugacy classes of ± 1 are special and correspond to points. This gives a geometric interpretation to the fact that the defects \mathcal{D}_a carry the same labels as Cardy boundary states. Indeed, the label a corresponds to a polar angle distinguishing the different 2-spheres $S^2 \subset S^3$ of a single SU(2).

We first compute the entanglement entropy in the large k limit while keeping the label a fixed. In the limit $k \to \infty$, the correction $s(\mathcal{D}_a)$ to the universal bulk entanglement entropy log L obtained from the defect \mathcal{D}_a becomes an integral,

$$s(\mathcal{D}_a) = -\lim_{k \to \infty} \frac{2}{k+2} \sum_{i=0}^k \sin^2\left(\frac{\pi(a+1)(i+1)}{k+2}\right) \log\frac{\sin^2\left(\frac{\pi(a+1)(i+1)}{k+2}\right)}{\sin^2\left(\frac{\pi(i+1)}{k+2}\right)}$$
$$= -2\int_0^1 dx \, \sin^2(\pi(a+1)x) \log(\frac{\sin^2(\pi(a+1)x)}{\sin^2(\pi x)}) \,. \tag{4.27}$$

In particular, we see that in the large k limit, the probability distribution of the interface is a continuous sine-square distribution

$$p_a(x) = 2\sin^2 \pi (a+1)x, \quad x \in [0,1].$$
 (4.28)

The distributions of sphere-like conjugacy classes are related to a conjugacy class corresponding to a point.

The integration can be performed by elementary methods. We first split the logarithmic term. The first of the two resulting summands,

$$\int_0^1 dx \, \sin^2(\pi(a+1)x) \log(\sin^2(\pi(a+1)x)) = \frac{1}{\pi} \int_0^\pi dy \, \sin^2 y \log(\sin^2 y) = \frac{1}{2} - \log 2 \quad (4.29)$$

is independent of a. In the other summand we use $2\sin^2 x = 1 - \cos(2x)$ to obtain

$$-\int_{0}^{1} dx \sin^{2}(\pi(a+1)x) \log(\sin^{2}(\pi x)) =$$

= $-\frac{1}{2\pi} \int_{0}^{\pi} dy \log(\sin^{2} y) + \frac{1}{2\pi} \int_{0}^{\pi} dy \cos(2(a+1)y) \log(\sin^{2} y).$ (4.30)

The first integral on the right-hand side of (4.30) yields $-\int_0^{\pi} dy \log \sin^2 y = 2\pi \log 2$. In the second integral we use partial integration to obtain

$$\int_0^{\pi} dy \, \cos(2(a+1)y) \log(\sin^2 y) = = -\frac{1}{a+1} \int_0^{\pi} dy \, \sin(2(a+1)y) \cot(y) = -\frac{\pi}{a+1} \,.$$
(4.31)

Using (4.29) - (4.31), (4.27) becomes

$$s(\mathcal{D}_a) = -\frac{a}{a+1}, \qquad a \ll k.$$
(4.32)

In particular, the contribution to the entanglement entropy from such an elementary defect \mathcal{D}_a is given by a rational number.

However, there is a second class of defects, for which the approximations made in the calculation leading to the result (4.32) do not hold. This is in particular the case if we pick a such that a + 1 divides k + 2 and take the limit keeping the ratio (a + 1)/(k + 2) fixed. Let us for example consider the case a = k/2 (k even), geometrically corresponding to the equatorial two-sphere, which is the fixed point under the involution $a \to k - a$. In this case the probabilities p_i^a vanish for i odd, and take the value 2/(k + 2) for i even. Using similar methods as above, the entanglement entropy in the limit $k \to \infty$ becomes

$$s(\mathcal{D}_{\underline{k}}) = -\log 4, \qquad k \to \infty.$$
(4.33)

Since $\log 4 > 1$ one observes that $-s(\mathcal{D}_{\frac{k}{2}})$ deviates substantially from the value (4.32). In fact, plotting of $-s(\mathcal{D}_a)$ at finite even k one observes a peak in the entanglement at a = k/2. Similar, less pronounced peaks are obtained at other values where a + 1 divides k + 2.

For generic defects, a + 1 does not divide k + 2, but of course (a + 1)/(k + 2) is still a rational number that we denote l/n, where l, n are coprime. It is natural to ask what happens if instead of a (as in the computation leading to (4.32)) we keep l/n fixed when taking the large k limit. In this case we find from (4.25) the expression

$$s(\mathcal{D}_{\frac{l(k+2)}{n}-1}) = -\log(2n) - H(n), \qquad k \to \infty,$$
 (4.34)

where H(n) is the entropy of a probability distribution $p_m = \frac{2}{n} \sin^2(\frac{\pi m}{n})$ for m = 1, 2, ..., n,

$$H(n) = \sum_{m=1}^{n} \frac{2}{n} \sin^2(\frac{\pi m}{n}) \log\left(\frac{2}{n} \sin(\frac{\pi m}{n})^2\right).$$
(4.35)

Note that the values of s in (4.34) are multiply degenerate, as the right-hand side does not depend on l. The entropies (4.34) are bounded from below by $s(D_{\frac{k}{2}})$, showing again that the defect corresponding to the equatorial two-sphere has minimum entanglement entropy. On the other hand, for $n \gg l$ they quickly approach the value -1 from below, such that this asymptotic expression in fact comes rather close to the approximation (4.32).

We will not go much further into details, and instead plot the entanglement entropy correction $-s(\mathcal{D}_a)$ at a finite value of k together with the approximation (4.32) in figure 2. The plot illustrates that the values of $s(\mathcal{D}_a)$ approach the asymptotic values (4.32) rather well for generic values of a. It also illustrates the peaks of the values at the special points where (4.34) deviates strongly from (4.32).

A nice pattern arises when we consider the fusion product of elementary defects at fixed labels a and b for $k \to \infty$. For finite k, the product $\mathcal{D}_{a \times b} = \mathcal{D}_a \mathcal{D}_b$ has the decomposition

$$\mathcal{D}_{a \times b} = \sum_{c} N_{ab}^{\ c} \mathcal{D}_{c} \tag{4.36}$$

in terms of elementary defects, where $N_{ab}^{\ c}$ are the fusion rules. Using the fact that in the large k limit the number of vacua on the defects contained in the fusion product is

$$\mathcal{N}_{0\,a\times b}^{a\times b} = \min(a,b) + 1\,,\tag{4.37}$$



Figure 2. Plots of $-s(\mathcal{D}_a)$ for large values of k, together with the asymptotic values (4.32). The peaks in the plots are captured by the asymptotic expression (4.34).

the probability distribution for $\mathcal{D}_{a \times b}$ is given by

$$p_i^{a \times b} = \frac{2}{\min(a,b)+1} \frac{\sin^2(\frac{\pi(a+1)(i+1)}{k+2})\sin^2(\frac{\pi(b+1)(i+1)}{k+2})}{(k+2)\sin^2(\frac{\pi(i+1)}{k+2})}.$$
(4.38)

The expression for $s(\mathcal{D}_{a \times b})$ in the large-k limit can be written as two summands, by splitting off the part involving the logarithm of the factor $\min(a, b) + 1$ in (4.38). As explained in appendix A, using elementary methods one can show that in all cases

$$s(\mathcal{D}_{a \times b}) = -\frac{p}{q} + \log(\min(a, b) + 1),$$
 (4.39)

where p and q are natural numbers depending on the labels a and b. Note that the argument in the logarithm is the number of elementary defects in the decomposition of the fusion product. However, the fact that this logarithm directly reflects the number of elementary defects in the decomposition is true only if each of these elementary defects appears with multiplicity 1.

5 Left/Right Entanglement Entropy

In this section we consider a system with a boundary. As mentioned in the introduction, the real-space entanglement entropy of a system with boundary receives a correction by the boundary entropy $s = \log g$, where g is the universal non-integer ground-state degeneracy of [3]. The entanglement entropy we are interested in is the left/right entanglement entropy (LREE) considered before in [10] for the free boson and in [11, 33] for generic CFTs. The two subsystems consist of the left- and right-moving part of the Hilbert space.

As mentioned in section 2, a conformal boundary condition is the maximally reflective solution to the interface conformality condition (2.1). For a CFT in the upper half-plane, the components of the bulk energy-momentum tensor satisfy

$$T = \tilde{T}|_{\mathbb{R}} \,. \tag{5.1}$$

To each boundary condition one can associate a boundary state. If we map the upper half plane to the unit disc, the boundary condition becomes such a boundary state in standard radial quantisation. The defining property for a conformal boundary state $|B\rangle$ is the gluing condition

$$(L_n - \bar{L}_{-n})|B\rangle = 0$$
. (5.2)

This means that the boundary state breaks one half of the conformal charges. Such a boundary state is coherent, *i.e.* it belongs to an extension of the closed string Hilbert space, and it is in particular not normalisable. As before, we decompose the Hilbert space as in (2.2). If the boundary state preserves the full chiral algebra, the gluing condition (5.2) is supplemented by similar conditions for the additional generators. Together these gluing conditions can only be solved in a sector $\mathcal{H}_i \otimes \mathcal{H}_{\bar{i}}$ where the two representations in the product are isomorphic. In this way one obtains the Ishibashi states $|i\rangle\rangle$ [34]. For a pair of a boundary state and a dual boundary state imposed on the edges of an annulus, the global constraint of consistency under the analog of the modular S transformation is referred to as the Cardy constraint [35], or as the open-closed string duality. As in the case of topological defects it reduces the linear space of solutions of the local gluing conditions to the positive cone of a lattice. The remaining consistent boundary states

$$|B\rangle = \sum_{\mathbf{i}} b_{B\mathbf{i}} |\mathbf{i}\rangle\rangle \tag{5.3}$$

are linear combinations of Ishibashi states. In this section, bold-faced indices

$$\mathbf{i} = (i, \alpha, \beta) \tag{5.4}$$

only contain one representation label. The sum in (5.3) and in the rest of this section only runs over representations of the bulk space of states with $i = \bar{i}$. The multiplicity labels α and β distinguish the different instances where the representation i appears in the holomorphic and antiholomorphic part of the space of states, respectively.

In our original setup, correlators and fields in the BCFT will depend on holomorphic and anti-holomorphic coordinates restricted to the upper half plane. Using the doubling trick, we regard the dependence on anti-holomorphic coordinates \bar{z} on the upper half plane as a dependence on holomorphic coordinate $z^* = \bar{z}$ for mirror fields on the lower half plane. This means that we consider a chiral construction on the full plane, where the stress tensor is continuous everywhere. The boundary condition is then a topological interface in this chiral part of a CFT, located on the real line.

Unfolding the Ishibashi states one obtains interface-like operators $\|\mathbf{i}\|$ that project onto a specific pair of representations \mathbf{i} . We therefore associate to the boundary state (5.3) an interface operator

$$\mathcal{I}(B) = \sum_{\mathbf{i}} b_{B\mathbf{i}} \|\mathbf{i}\|.$$
(5.5)

Our computation of the entanglement entropy now proceeds in analogy with the previous sections. The characters in the expression

$$Z(K) = \sum_{i} \operatorname{Tr} \left(b_{B^* \mathbf{i}} b_{B \mathbf{i}} \right)^K \operatorname{Tr}_{\mathcal{H}_i} \left(e^{-2\delta H} \right) = \sum_{i} \operatorname{Tr} \left(b_{B^* \mathbf{i}} b_{B \mathbf{i}} \right)^K \chi_i \left(e^{-2\delta K} \right)$$
(5.6)

can be written by means of the modular S matrix as

$$Z(K) = \sum_{i} \operatorname{Tr} \left(b_{B^* \mathbf{i}} b_{B \mathbf{i}} \right)^K S_{ij} \chi_j \left(e^{-\frac{2\pi^2}{\delta K}} \right) \approx \sum_{i} \operatorname{Tr} \left(b_{B^* \mathbf{i}} b_{B \mathbf{i}} \right)^K S_{i0} e^{\frac{2\pi^2 c}{\delta K 24}}.$$
 (5.7)

For the general boundary state with open string vacuum multiplicity \mathcal{N}_{0B}^{B} we write the entanglement entropy again in terms of a probability distribution. The distribution is defined by the traces of the matrices

$$p_{\mathbf{i}}^{B} = \frac{b_{B^{*}\mathbf{i}}b_{B\mathbf{i}}\,S_{i0}}{\mathcal{N}_{0B}^{B}}\,.$$
(5.8)

In (5.8) we abuse the index notation in the same way as in the previous chapters — while the indices **i** on the right-hand side contain one multiplicity label for holomorphic and one for antiholomorphic representations (and we again suppress the summation over interior labels), the index **i** on the left-hand side includes two multiplicity labels of the same kind. The LREE attributed to a system with boundary condition B then reads

$$S = \frac{c}{6} \log L - \sum_{i} \operatorname{Tr} p_{\mathbf{i}}^{B} \log \frac{p_{\mathbf{i}}^{B}}{S_{i0}}.$$
(5.9)

Note that the prefactor of the logarithmic term is one half of the prefactor in the case of a full theory with a topological interface. This reproduces the area law result mentioned in [2].

A natural question is whether it is again possible to interpret the result in terms of a Kullback-Leibler divergence. However, for interfaces there is a generic "neutral" interface (the identity defect) with respect to which one can compute the relative entropy. This is no longer the case for boundaries, as there is no "neutral" boundary on the full plane that could serve as a reference point. There will generically always be "information loss" when left movers are scattered by the boundary into right movers. Exceptional cases occur when the boundary condition is a permutation boundary condition obtained by folding an identity or symmetry defect to a boundary condition for a tensor product of identical CFTs.

Technically, one can try to interpret the denominator in the logarithm of (5.9) as a distribution corresponding to the entries S_{i0} times appropriate identity matrices. However, the sum over the traces of these matrices is in general not equal to 1, and therefore not a probility distribution. In the cases where it is, we indeed obtain the relative entropy with respect to a permutation boundary state, where each b_i is a permutation matrix. However, in general we conclude that the interpretation as a relative entropy fails in the case of the LREE boundary states.

An immediate consequence of loosing the interpretation of the LREE as a relative entropy is that the contribution

$$s = -\sum_{i} \operatorname{Tr} p_{\mathbf{i}}^{B} \log \frac{p_{\mathbf{i}}^{B}}{S_{i0}}$$
(5.10)

is not necessarily negative any more. The technique we applied in the case of interfaces in section 4 now yields the upper bound

$$s \le \log\left(\sum_{i} S_{i0}\right). \tag{5.11}$$

This bound does not need to be negative. As an example consider boundary states in diagonal rational models. Elementary boundary states which preserve the rational symmetry are labelled by irreducible representations b of the symmetry algebra. The coefficients b_{bi} of the elementary boundary states in this case are

$$b_{bi} = \frac{S_{bi}}{\sqrt{S_{i0}}} \,. \tag{5.12}$$

From (5.8) and (5.10) we obtain the LREE

$$S = \frac{c}{6} \log L - \sum_{i} |S_{bi}|^2 \log \frac{|S_{bi}|^2}{S_{i0}}.$$
(5.13)

This reproduces the result obtained previously in [11]. It seems plausible that all symmetrypreserving boundary states in a diagonal model have the LREE of the Cardy brane associated to the identity as an upper bound,

$$s \le -\sum_{i} S_{0i}^2 \log S_{0i} \,. \tag{5.14}$$

The right-hand side is stricter than the bound (5.11), and it is always positive.

Example: Ising model

The LREE for boundary states of the Ising model has been discussed in [11]. We quote the results here for illustration. The Cardy states in the Ising model are explicitly given in terms of Ishibashi states by

$$|id\rangle = \frac{1}{\sqrt{2}} \left(|id\rangle\rangle + |\epsilon\rangle\rangle + 2^{\frac{1}{4}} |\sigma\rangle\rangle \right),$$

$$|\epsilon\rangle = \frac{1}{\sqrt{2}} \left(|id\rangle\rangle + |\epsilon\rangle\rangle - 2^{\frac{1}{4}} |\sigma\rangle\rangle \right),$$

$$|\sigma\rangle = |id\rangle\rangle - |\epsilon\rangle\rangle.$$

(5.15)

The contributions to the LREE we obtain from (5.10) are

$$s = \frac{3\log 2}{4}$$
 for $|id\rangle, |\epsilon\rangle$, and $s = 0$ for $|\sigma\rangle$. (5.16)

Example: $su(2)_k$ boundary states and the $k \to \infty$ limit

Analogously to the example of $su(2)_k$ defects in section 4 we consider the LREE of boundary states in the WZW models $su(2)_k$ in the limit $k \to \infty$. For finite k, the theory is diagonal and rational, and the formulae of [11] apply (see [36] for a discussion of the LREE in WZW models at finite k). The Cardy states (5.12) are again labelled by spins s = b/2 for b = 0, 1, ..., k. From (5.13), the universal contribution to the LREE by the state B_b is

$$s(b) = -\sum_{i=0}^{k} \frac{2}{k+2} \sin^2(\pi(b+1)\frac{(i+1)}{k+2}) \log \frac{\sin^2(\pi(b+1)\frac{i+1}{k+2})}{\sin(\pi\frac{i+1}{k+2})} + \log \sqrt{\frac{2}{k+2}}.$$
 (5.17)

Here we have split off a factor depending only on k from the argument of the logarithm, and used that $\sum_i p_i^b = 1$. Observe that the shift term $-\log(k+2)$ in (5.17) has the right form to be identified with (the logarithm of) the radius of the target space. The target space of the $su(2)_k$ WZW model is the (fuzzy) sphere S^3 at radius $R = \sqrt{k}$. As in the defect case, the sum in (5.17) becomes an integral in the $k \to \infty$ limit. By the same methods as in section 4 we obtain

$$s(b) = -\frac{2b+1}{2b+2} + \frac{1}{2}\log 2 + \log R, \qquad k \to \infty, \,. \tag{5.18}$$

The positive (and infinite) contribution from the radius is similar to the radius contribution to the LREE of Dirichlet branes of the compactified boson [10], see (6.26).

Example: Fusion of defect and boundary in the $su(2)_k$ WZW model

To extend the result of the last example we consider the fusion product of an elementary defect operator \mathcal{D}_a with a Cardy boundary state $|b\rangle$. This yields a new boundary state $|B\rangle = \mathcal{D}_a|b\rangle$. From (2.10) and (5.12) we see that the coefficients of $|B\rangle$ are given by

$$b_{Bi} = \frac{S_{ai}^2 S_{bi}^2}{S_{0i}^{3/2}}.$$
(5.19)

The number of open-string vacua in the self-spectrum of $|B\rangle$ is

$$\mathcal{N}_{0B}^{\ B} = \sum |b_{Bi}|^2 S_{0i} = \min(a, b) + 1, \qquad (5.20)$$

as in the case of the fusion products of two elementary defects in example 3 of section 4. The subleading contribution to the LREE can be written as

$$s(a \times b) = \log(\min(a, b) + 1) - \frac{\sum_{i} |b_{Bi}|^2 S_{0i} \log |b_{Bi}|^2}{\min(a, b) + 1}.$$
 (5.21)

Observe that $\min(a, b) + 1$ is also again the number of elementary branes in the decomposition

$$|B\rangle = \sum_{c} N_{ab}^{\ c} |c\rangle \tag{5.22}$$

of the fusion product.

In the large k limit of the $su(2)_k$ WZW model, the LREE of the fusion product differs from the entropy of the original boundary state $|b\rangle$ again by a rational term and the logarithm of the number of elementary branes in the decomposition. Indeed, in the limit of large k the numerator in the second term of the right-hand side of (5.21) becomes

$$\sum_{i} |b_{Bi}|^2 S_{0i} \log |b_{Bi}|^2 \xrightarrow{k \to \infty} (5.23)$$
$$-\log \sqrt{\frac{2}{k+2}} + \frac{2}{\pi} \int_0^{\pi} \frac{\sin^2((a+1)x)\sin^2((b+1)x)}{\sin^2(x)} \log \left(\frac{\sin^2((a+1)x)\sin^2((b+1)x)}{\sin^3(x)}\right) dx \,.$$

A similar calculation as in example 3 of section 4 leads to

$$s(a \times b) = \log \min(a+1, b+1) + \frac{1}{2}\log 2 + \log R - \frac{p}{q}, \qquad k \to \infty,$$
 (5.24)

for some $p, q \in \mathbb{N}$. The difference between the entanglement entropy of the boundary state after fusion (5.24) and the original boundary state (5.18) for $k = \infty$ is therefore

$$s(a \times b) - s(b) = \log(\min(a, b) + 1) - \frac{p}{q}.$$
(5.25)

6 Results for Bosonic Tori

6.1 EE through topological defects

For the case of d free bosons compactified on a torus the interface operators are explicitly known [37]. The ground states of the theory form an even, self dual lattice $\Gamma \subset \mathbb{R}^{d,d}$. The lattice vectors are of the form $\gamma = (p, \bar{p})$, where the d-dimensional vectors p and \bar{p} denote left- and right-moving momenta. We will consider topological interfaces that also preserve the full $u(1)^d$ symmetry. These interfaces are specified by a gluing matrix $\Lambda \in O(d|\mathbb{R}) \times$ $O(d|\mathbb{R})$. Similarly to the rational case discussed earlier, the interface operators can be written as linear combinations of operators between $u(1)^d$ highest weight representations:

$$\mathcal{I}_{12}(\Lambda) = \sum_{\gamma \in \Gamma_{12}^{\Lambda}} d_{\Lambda\gamma} ||\gamma|| \,. \tag{6.1}$$

As before, $||\gamma||$ is an intertwiner of the representation space specified by the lattice vector γ , and $d_{\Lambda\gamma}$ are prefactors constrained by consistency under modular S transformation. The range of the summation is restricted to a sublattice, given in terms of a gluing condition Λ for the lattices Γ_1 and Γ_2 on the two sides of the interface,

$$\Gamma_{12}^{\Lambda} = \{ \gamma \in \Gamma_1 \, | \, \Lambda \gamma \in \Gamma_2 \} = \Gamma_1 \cap \Lambda^{-1} \Gamma_2 \subset \Gamma_1.$$
(6.2)

For admissible gluing conditions Λ , the sublattice Γ_{12}^{Λ} has full rank. Consistency under modular S transformation then demands that $d_{\Lambda\gamma} = g_{12}^{\Lambda} \exp(2\pi i \varphi(\gamma))$, where $\varphi \in (\Gamma_{12}^{\Lambda})^*$ and

$$(g_{12}^{\Lambda})^2 = |\Gamma_1 / \Gamma_{12}^{\Lambda}| \tag{6.3}$$

is the index of the sublattice Γ_{12}^{Λ} inside the lattice Γ_1 . The topological interface operator splits into a lattice and an oscillator part,

$$\mathcal{I}_{12} = \mathcal{I}_{12}^0(\Lambda) \prod_{n>0} \mathcal{I}_{12}^n(\Lambda) , \qquad (6.4)$$

where

$$\mathcal{I}_{12}^{0} = g_{12}^{\Lambda} \sum_{\gamma \in \Gamma_{12}^{\Lambda}} e^{2\pi i \varphi(\gamma)} |\Lambda \gamma \rangle \langle \gamma |$$
(6.5)

gives the map for the zero modes and the

$$\mathcal{I}_{12}^{n} = \exp\left(-\frac{1}{n}\left(a_{-n}^{2}\Lambda_{11}a_{n}^{1} + \tilde{a}_{-n}^{2}\Lambda_{22}\tilde{a}_{n}^{1}\right)\right)$$
(6.6)

for n > 0 give the contribution of the higher modes. It is implicitly understood that modes of CFT1 act from the right and modes of CFT2 from the left of \mathcal{I}_{12}^0 .

In order to determine the entanglement entropy we proceed as before. The partition function of the K-sheeted Riemann surface for the topological defect (4.1) is

$$Z(K) = \operatorname{Tr}\left(\left(\mathcal{I}\mathcal{I}^{\dagger}\right)^{K} e^{-2\delta KH}\right)$$

= $(g_{12}^{\Lambda})^{2K} \sum_{(p,\bar{p})=\gamma\in\Gamma_{12}^{\Lambda}} \chi_{p}(i\frac{\delta}{\pi}K)\bar{\chi}_{\bar{p}}(i\frac{\delta}{\pi}K),$ (6.7)

where the χ_p are the u(1) characters. We perform a modular S transformation and express Z(K) in terms of characters depending on the variable $i\pi/K\delta$. This leads to a summation over lattice vectors in the dual lattice Γ_{12}^{\vee} . In the limit $\delta \ll 1$ we approximate the lattice sum by the dominant contribution of the vacuum $p = \bar{p} = 0$,

$$Z(K) = (g_{12}^{\Lambda})^{2K} \sum_{(q,\bar{q})\in\Gamma_{12}^{\vee}} a_{(q,\bar{q})}\chi_q(i\frac{\pi}{K\delta})\bar{\chi}_{\bar{q}}(-i\frac{\pi}{K\delta})$$

$$\approx (g_{12}^{\Lambda})^{2K} a_{(0,0)}e^{\frac{\pi^2 d}{6\delta K}}.$$
(6.8)

We now use that the interfaces with the normalization (6.3) are elementary. For K = 1, we obtain the ordinary defect partition function, where the multiplicity of the vacuum propagating in the dual channel is 1. We therefore conclude that $a_{(0,0)} = 1/(g_{12}^{\Lambda})^2$.

Using $\delta = \pi^2 / \log(L)$, c = d for the central charge of d bosons, and (6.3), the entanglement entropy is given by

$$S = (1 - \partial_K) \log(Z(K)) \Big|_{K=1} = \frac{c}{3} \log(L) - \log |\Gamma_1 / \Gamma_{12}^{\Lambda}|.$$
(6.9)

In the special case d = 1, *i.e.* for a free boson compactified on a circle, conformal interfaces between CFT1 and CFT2 are classified by two winding numbers k_1 and k_2 . For generic compactification radii these interfaces are not topological, but they become so by choosing radii to satisfy the relation $R_1/R_2 = k_2/k_1$ [38]. In this case the index of the sublattice is [37, 38]

$$|\Gamma_1 / \Gamma_{12}^{\Lambda}| = |k_1 k_2|, \qquad (6.10)$$

such that the entanglement entropy through the topological interface is given by

$$S = \frac{1}{3}\log(L) - \log|k_1k_2| , \qquad (6.11)$$

in agreement with the result of [12].

All topological toroidal interfaces are duality interfaces according to our previous definitions. A subclass of them are symmetry interfaces and describe automorphisms of the toroidal CFT. They are associated to gluing matrices in the T-duality group $O(d, d, \mathbb{Z})$. In particular, for those matrices we get $\Gamma_{12}^{\Lambda} = \Gamma_1$, which means that the defect couples to the full momentum lattice and no ground states are projected out. In this case, there is no contribution to the entanglement entropy from the interface. The broader class of duality interfaces is specified by matrices in $O(d, d, \mathbb{Q})$. These interfaces can in particular be related to orbifold constructions. In the case of a single circle, where $R_1/R_2 = k_2/k_1$ [37, 38], the theory with radius R_1 can be obtained from the theory with radius R_2 by orbifolding with respect to the shift symmetry

$$X \mapsto X + 2\pi R_1 \,. \tag{6.12}$$

The orbifold group generated by this symmetry is $\mathbb{Z}_{|k_1k_2|}$, and hence of order $|k_1k_2|$. It is clear that \mathcal{II}^{\dagger} projects the theory with R_2 onto the sector invariant under the orbifold group. We see that for circle theories the contribution to the sub-leading term of the entanglement entropy is set by the order of this orbifold group,

$$S = \frac{1}{3}\log(L) - \log|G|, \qquad (6.13)$$

in agreement with the general result (4.19).

6.2 LREE of bosonic tori

We consider the LREE for d free bosons compactified on a torus. The gluing conditions can be written as [37, 39]

$$(a_n + O\tilde{a}_{-n})|B\rangle = 0, \qquad (6.14)$$

where $O \in O(d|\mathbb{R})$. The ground states solving the zero mode condition are given by

$$\Gamma^{O} = \left\{ \begin{pmatrix} -Ox\\ x \end{pmatrix} \cap \Gamma \mid x \in \mathbb{R}^{d} \right\}, \qquad (6.15)$$

where Γ is the charge lattice of the torus model. The boundary state is a superposition of Ishibashi states $|p, \bar{p}\rangle$ built on ground states $(p, \bar{p}) \in \Gamma^O$,

$$|B\rangle = g \sum_{(p,\bar{p})\in\Gamma^O} e^{i\varphi(p,\bar{p})} |p,\bar{p}\rangle\rangle, \qquad (6.16)$$

where the function $\varphi \in (\Gamma^O)^*$ specifies the *D*-brane moduli and the *g*-factor is fixed by the condition that the open-string vacuum appears with multiplicity 1. As before, we unfold the boundary state and associate an interface between left- and right-movers of the free boson theory. For this, we introduce the projections $\pi(\Gamma^O)$ and $\bar{\pi}(\Gamma^O)$ of the lattice Γ^O to the left- and right-moving parts respectively. On the level of ground states, the interfaces therefore maps $\pi(\Gamma^O) \ni p \to -Op \in \bar{\pi}(\Gamma^O)$. The *g*-factor is given by

$$g = vol(\pi(\Gamma^O)), \tag{6.17}$$

the volume of the unit cell of $\pi(\Gamma^O)$. The computation of the entanglement entropy now proceeds in analogy to the case of topological interfaces. The partition functions on the K-sheeted torus are approximated by

$$Z(K) = g^{2K} \sum_{(p,\bar{p})\in\Gamma^O} \chi_p(i\frac{\delta}{\pi}K) \xrightarrow{S \operatorname{trsf}, \delta \ll 1} g^{2K-2} e^{\frac{\pi^2}{12\delta K}}, \qquad (6.18)$$

where we used again that the vacuum in the open string channel has multiplicity 1 for K = 1. For $c = d, \delta = \pi^2 / \log(L)$ we obtain from this the LREE

$$S = \frac{c}{6}\log(L) - \log \operatorname{vol}(\pi(\Gamma^O)).$$
(6.19)

The subleading part of the LREE is determined by the g factor of the boundary state. To relate this quantity to the torus geometry, let us recall that Γ is a Narain lattice given by

$$\Gamma = \left\{ \begin{pmatrix} \frac{1}{2}E^{-1}N + E^{T}(1+B)M \\ -\frac{1}{2}E^{-1}N + E^{T}(1-B)M \end{pmatrix} \mid M, N \in \mathbb{Z}^{d} \right\} ,$$
(6.20)

where $G = EE^T$ is the metric and B the Kalb-Ramond field on the target space, and Nand M are the momentum and winding quantum numbers, respectively. Let us consider a D1 brane in d = 2 dimensions for the geometric case where B is zero. If the D1 brane were located in infinite flat space, we would specify the direction of the brane by specifying the momenta perpendicular to it; a localisation of the brane to its world-volume direction would then be achieved by integration over these momenta. On a torus, the momenta are part of a lattice. We can fix our brane by choosing the elementary generator of transverse momenta that couple to the brane to be given by

$$N^0 = \begin{pmatrix} N_1^0 \\ N_2^0 \end{pmatrix}, \tag{6.21}$$

with two integers N_i^0 that we assume to be relatively prime. This choice determines the winding modes our D1 brane can couple to. The elementary winding generator $M^0 = (M_1^0, M_2^0)$ is again specified by two coprime integers M_1^0, M_2^0 , and have to satisfy the orthogonality constraint

$$N_1^0 M_1^0 + N_2^0 M_2^0 = 0. (6.22)$$

The equation is solved by $M_1^0 = -N_2^0$, $M_2^0 = N_1^0$. By this we have fixed a D1 brane for which the lattice Γ^O is precisely spanned by the two generators N^0 and M^0 for N and M in (6.20), respectively. It can be checked explicitly that these lattice vectors solve (6.14) with

$$O = \begin{pmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{pmatrix} \in O(2, \mathbb{R}), \qquad (6.23)$$

where $\theta = \arctan\left(-(E^{-1}N^0)_1/(E^{-1}N^0)_2\right)$ [38]. To compute the *g*-factor we now have to compute the volume of (the unit cell of) this lattice, projected to the left-movers. After some algebra one obtains

$$g^{2} = \frac{1}{2 \det E} \left((N_{2}^{0})^{2} G_{11} + (N_{1}^{0})^{2} G_{22} - 2G_{12} N_{2}^{0} N_{1}^{0} \right) .$$
 (6.24)

Mapping N_i^0 to the winding numbers of the brane $N_1^0 = k_2, N_2^0 = -k_1$, we see that

$$g^2 = \frac{length^2}{2vol},\tag{6.25}$$

where *length* refers to the length of the brane and *vol* to the volume of the torus. This is in agreement with geometrical expectations. Note also, that in the special case of a rectangular torus with diagonal metric where the radii are related by a rational number the above result agrees with the one for topological interfaces of the previous section.

The left- right-entanglement entropy has been computed before for the case of a free boson in [10]. To compare the results, note that in one dimension the left-moving momenta are given by $a_0 = N/2R + MR$ and the right-moving momenta by $\bar{a}_0 = N/2R - MR$. The matrix O in the gluing condition reduces to a choice of sign. For Dirichlet branes we have O = 1, and only ground states without winding contribute to the boundary state. We therefore have $\Gamma^O = \{(N/2R, N/2R)\}$, and the volume of the projected unit cell is 1/2R. Similar considerations also hold for Neumann branes. Our result (6.19) for the LREE of a single boson compactified on a circle thus gives

$$S = \frac{1}{6} \log L - \begin{cases} \log R & \text{for } O = -1 \text{ (Neumann b.c.)} \\ \log \frac{1}{2R} & \text{for } O = 1 \text{ (Dirichlet b.c.)} \end{cases}$$
(6.26)

which in particular reproduces the results of [10].

7 Conclusion and Outlook

In this paper we have discussed the entanglement entropy through topological interfaces, and the left/right entanglement entropy at conformal boundaries. Our focus has been on unitary CFTs with discrete spectrum. For an interface between CFT1 and CFT2 we trace out the half-plane on the CFT2 side of the interface. The eigenvalues of the resulting reduced density matrix (4.8) give the probability of finding the reduced system CFT1 in a thermal state in the representation (i, \bar{i}) . The general result for the entanglement entropy through a topological interface is obtained in (4.11). The universal bulk term proportional to log L is not affected, as correlation functions do not depend on the position or shape of the interface insertion. To subleading order the topological interface contributes a universal term to the entanglement entropy. The contribution is the negative of the relative entropy (Kullback-Leibler divergence) of the distribution associated to the interface, compared with the situation when there was no interface to start with.

By unfolding a theory with a boundary we obtain a topological interface in a chiral theory, allowing a derivation of the left/right entanglement entropy (5.9) in analogy to the interface case. In this situation we lose the interpretation of the entanglement entropy in terms of a relative entropy. However, the derivation of the entanglement entropy proceeds in analogy to the interface case, and the resulting formulas have a similar structure.

The Kullback-Leibler divergence has many appealing features. In this paper, we have interpreted only some of them in the context of defects and interfaces. In particular, we have given a meaning to its positivity – the interface will never increase the entanglement. We have also connected the vanishing of the Kullback-Leibler divergence to properties of interfaces that do not lead to information loss.

One particular issue that we did not touch in this paper is that the Kullback-Leibler divergence measures the difference between probability distributions, and in this sense shares some properties with a distance (for an application to distances between quantum field theories see e.g. [40]). In the present paper we only obtained Kullback-Leibler divergences of interfaces with respect to the identity defect. In order to explore the distance property, it would be interesting to study a concrete physical realisation for Kullback-Leibler divergences with respect to arbitrary interfaces as reference points.

The Kullback-Leibler divergence is evidently not symmetric, and even after a suitable symmetrisation fails to satisfy the triangle inequality. The distance measure it might yield for topological defects would therefore share the same features.

We recall that interfaces have been used to define distances before. In particular, in [41] the interface entropy $\log g$ of deformation interfaces⁶ was identified with Calabi's Diastasis function. Ultimately, the proposal was that the *g*-factor between interfaces can be used to define a distance between different CFTs. However, also in this case it was observed that the triangle inequality will not be satisfied.

A further observation for the $\log g$ distance is that it gives rise to a metric at the infinitesimal level, which coincides with the Zamolodchikov metric on the moduli space. This is again similar in the case of the Kullback-Leibler divergence, where the infinitesimal limit yields the Fisher information metric. At the moment, for our purposes this property is at the moment rather formal, as the interfaces we have studied here are generally labelled by discrete numbers.

Note that the deformation interfaces considered in [41] are in particular not topological. In fact, all of them are in the deformation class of the identity defect. The entanglement entropy through such interfaces has been investigated only in examples. One particular case is the theory of a compactified free boson [12]. There the subleading term of the entanglement entropy vanishes, and instead the prefactor of the log term changes. In fact, it is suggestive that the subleading terms we discussed in this paper vanish more generally for such deformations of the identity. It would be interesting to prove a general statement about the form of the entanglement entropy along these lines.

Finally, let us comment on N = 2 supersymmetric theories. Such theories can be topologically twisted, and one can formulate boundary as well as interface gluing conditions that are compatible with the topological twist. On the level of the topological theory, all interfaces can be moved freely. It would be interesting to consider entanglement entropy through topological interfaces in the supersymmetric situation, where entanglement should have a topological interpretation. For results on the supersymmetric case without interfaces see [42].

Acknowledgements

The authors would like to thank Charles Melby-Thompson and Erik Plauschinn for useful discussions. This work was partially supported by the DFG Transregio Grant TRR33 and the Excellence Cluster Universe.

⁶Deformation interfaces relate different CFTs in the same moduli space in a minimal way.

A Entanglement entropy of a fusion product

In this appendix we show that the pattern (4.39) holds. This follows from the fact that in the large k limit, the contribution

$$-2\int_{0}^{1} \frac{\sin^{2}(\pi(a+1)x)\sin^{2}(\pi(b+1)x)}{\min(a+1,b+1)\cdot\sin^{2}(\pi x)}\log\left(\frac{\sin^{2}(\pi(a+1)x)\sin^{2}(\pi(b+1)x)}{\sin^{4}(\pi x)}\right) dx \quad (A.1)$$

is a rational number. In order to see this, the basic integral that has to be evaluated is

$$-\frac{1}{\pi} \int_0^\pi \frac{\sin^2((a+1)x)\sin^2((b+1)x)}{\sin^2(x)} \log\left(\sin^2((l+1)x)\right) dx.$$
(A.2)

By the two representations of Clausen's function Cl_1 we can express the logarithmic factor in terms of the sum

$$\log(\sin^2(y)) = -\log 4 - \sum_{k=1}^{\infty} \frac{2\cos(2ky)}{k}.$$
 (A.3)

We note that the log 2 terms cancel out in (A.1), so it is enough to keep only the sum over cosines from the right-hand side of (A.3) for further calculations. One might be concerned that while the individual terms in the summation over k give rational results, resummation may yet yield something non-rational. In order to see that this is not the case we eliminate the sine functions in the denominator of (A.2) by writing the remaining sine functions in the numerator in terms of spread polynomials [43]

$$\sin^2(nx) = \sum_{p=0}^{n-1} \frac{n}{n-p} \binom{2n-1-p}{p} (-4)^{n-1-p} \sin^{2(n-p)}(x) =: \sum_{\{p\}} \sin^{2p}(x) \,. \tag{A.4}$$

Since we are not interested in the precise value of the finite result, the only relevant property for us is that the spread polynomials have rational coefficients and finite order. We define the summation symbol on the right-hand side to indicate a finite sum of trigonometric functions with rational coefficients. We reduce the right-hand side further by the identity

$$\sin^{2p}(x) = \frac{1}{2^{2p}} \binom{2p}{p} + \frac{2}{2^{2p}} \sum_{r=0}^{p-1} (-1)^{p-r} \binom{2p}{r} \cos(2(n-r)x) =: \sum_{\{r\}} \cos(2rx) \,. \tag{A.5}$$

The purpose of writing (A.4) and (A.5) is to demonstrate that (A.1) can indeed be written in the form

$$\frac{1}{\pi} \int_0^{\pi} \sum_{\{s\}} \cos(2sx) \sum_{k=1}^{\infty} \cos(2k(l+1)x) \frac{1}{k} \, dx \,. \tag{A.6}$$

By the integral identity $\int_0^{\pi} \cos(nx) \cos(mx) dx = \frac{\pi}{2} \delta_{n,m}$, the finite sum over s reduces the infinite sum over k to a rational result, and we obtain (4.39) as proposed.

References

 C. Holzhey, F. Larsen, and F. Wilczek, Geometric and renormalized entropy in conformal field theory, Nucl. Phys. B424 (1994) 443-467, [hep-th/9403108].

- [2] P. Calabrese and J. Cardy, Entanglement entropy and conformal field theory, J.Phys. A42 (2009) 504005, [arXiv:0905.4013].
- [3] I. Affleck and A. W. W. Ludwig, Universal noninteger "ground-state degeneracy" in critical quantum systems, Phys. Rev. Lett. 67 (Jul, 1991) 161–164.
- [4] J. A. Harvey, S. Kachru, G. W. Moore, and E. Silverstein, Tension is dimension, JHEP 03 (2000) 001, [hep-th/9909072].
- [5] T. Azeyanagi, A. Karch, T. Takayanagi, and E. G. Thompson, Holographic calculation of boundary entropy, JHEP 03 (2008) 054–054, [arXiv:0712.1850].
- [6] T. Takayanagi, Holographic boundary entropy and Janus solutions, Prog. Theor. Phys. Suppl. 177 (2009) 191–202.
- [7] J. Estes, K. Jensen, A. O'Bannon, E. Tsatis, and T. Wrase, On holographic defect entropy, JHEP 05 (2014) 084, [arXiv:1403.6475].
- [8] S. Ryu and T. Takayanagi, Holographic derivation of entanglement entropy from AdS/CFT, Phys. Rev. Lett. 96 (2006) 181602, [hep-th/0603001].
- [9] S. Ryu and T. Takayanagi, Aspects of holographic entanglement entropy, JHEP 08 (2006) 045, [hep-th/0605073].
- [10] L. A. Pando Zayas and N. Quiroz, Left-right entanglement entropy of boundary states, JHEP 01 (2015) 110, [arXiv:1407.7057].
- [11] D. Das and S. Datta, Universal features of left-right entanglement entropy, arXiv:1504.0247.
- [12] K. Sakai and Y. Satoh, Entanglement through conformal interfaces, JHEP 0812 (2008) 001, [arXiv:0809.4548].
- [13] E. M. Brehm and I. Brunner, Entanglement entropy through conformal interfaces in the 2d Ising model, arXiv:1505.0264.
- [14] I. Peschel, Entanglement entropy with interface defects, Journal of Physics A Mathematical General 38 (May, 2005) 4327–4335, [cond-mat/0502034].
- [15] V. Eisler and I. Peschel, Solution to the fermionic entanglement problem with interface defects, Ann. phys 522 (May, 2010) 679, [arXiv:1005.2144].
- [16] M. Gutperle and J. D. Miller, Entanglement entropy at holographic interfaces, arXiv:1511.08955.
- [17] D. Bak, M. Gutperle, and R. A. Janik, Janus black holes, JHEP 10 (2011) 056, [arXiv:1109.2736].
- [18] D. Bak and H. Min, Multi-faced black Janus and entanglement, JHEP 03 (2014) 046, [arXiv:1311.5259].
- [19] T. Quella, I. Runkel, and G. M. Watts, Reflection and transmission for conformal defects, JHEP 0704 (2007) 095, [hep-th/0611296].
- [20] J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, Duality and defects in rational conformal field theory, Nucl. Phys. B763 (2007) 354–430, [hep-th/0607247].
- [21] M. Oshikawa and I. Affleck, Defect lines in the Ising model and boundary states on orbifolds, Phys. Rev. Lett. 77 (1996) 2604–2607, [hep-th/9606177].

- [22] C. Bachas, J. de Boer, R. Dijkgraaf, and H. Ooguri, Permeable conformal walls and holography, JHEP 06 (2002) 027, [hep-th/0111210].
- [23] C. Bachas and M. Gaberdiel, Loop operators and the Kondo problem, JHEP 11 (2004) 065, [hep-th/0411067].
- [24] V. B. Petkova and J. B. Zuber, Generalized twisted partition functions, Phys. Lett. B504 (2001) 157-164, [hep-th/0011021].
- [25] J. Fuchs, M. R. Gaberdiel, I. Runkel, and C. Schweigert, Topological defects for the free boson cft, J. Phys. A40 (2007) 11403, [arXiv:0705.3129].
- [26] J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, Kramers-Wannier duality from conformal defects, Phys. Rev. Lett. 93 (2004) 070601, [cond-mat/0404051].
- [27] X.-L. Qi, H. Katsura, and A. W. W. Ludwig, General relationship between the entanglement spectrum and the edge state spectrum of topological quantum states, Physical Review Letters 108 (May, 2012) 196402, [arXiv:1103.5437].
- [28] S. Kullback and R. Leibler, On information and sufficiency, Annals of Mathematical Statistics 22 (1951) 79–86.
- [29] T. Faulkner, R. G. Leigh, and O. Parrikar, Shape dependence of entanglement entropy in conformal field theories, arXiv:1511.0517.
- [30] A. Yu. Alekseev and V. Schomerus, *D-branes in the WZW model*, *Phys. Rev.* D60 (1999) 061901, [hep-th/9812193].
- [31] C. Bachas, M. R. Douglas, and C. Schweigert, Flux stabilization of D-branes, JHEP 05 (2000) 048, [hep-th/0003037].
- [32] J. M. Figueroa-O'Farrill and S. Stanciu, *D*-branes in $AdS_3 \times S^3 \times S^3 \times S^1$, *JHEP* **04** (2000) 005, [hep-th/0001199].
- [33] I. Affleck, N. Laflorencie, and E. S. Sørensen, Entanglement entropy in quantum impurity systems and systems with boundaries, J.Phys. A42 (2009) 504009.
- [34] N. Ishibashi, The boundary and crosscap states in conformal field theories, Modern Physics Letters A 4 (1989) 251–264.
- [35] J. L. Cardy, Boundary conditions, fusion rules and the Verlinde formula, Nuclear Physics B 324 (1989), no. 3 581 – 596.
- [36] H. J. Schnitzer, Left-right entanglement entropy, D-branes, and level-rank duality, arXiv:1505.07070.
- [37] C. Bachas, I. Brunner, and D. Roggenkamp, A worldsheet extension of O(d, d; z), JHEP 1210 (2012) 039, [arXiv:1205.4647].
- [38] C. Bachas and I. Brunner, Fusion of conformal interfaces, JHEP 0802 (2008) 085, [arXiv:0712.0076].
- [39] R. Blumenhagen, D. Lüst, and S. Theisen, *Basic Concepts of String Theory*. Theoretical and Mathematical Physics. Springer Berlin Heidelberg, 2013.
- [40] V. Balasubramanian, J. J. Heckman, and A. Maloney, *Relative entropy and proximity of quantum field theories*, JHEP 05 (2015) 104, [arXiv:1410.6809].
- [41] C. P. Bachas, I. Brunner, M. R. Douglas, and L. Rastelli, Calabi's diastasis as interface entropy, Phys. Rev. D90 (2014), no. 4 045004, [arXiv:1311.2202].

- [42] A. Giveon and D. Kutasov, Supersymmetric Renyi entropy in CFT₂ and AdS₃, JHEP 01 (2016) 042, [arXiv:1510.08872].
- [43] S. Goh, Chebyshev polynomials and spread polynomials, Honours Thesis, School of Mathematics and Statistics, UNSW (2005).