# Borell's formula on a Riemannian manifold and applications

Joseph Lehec

September 28, 2018

#### 1 Introduction

Throughout the article  $(\Omega, \mathcal{A}, P)$  is a probability space, equipped with a filtration  $(\mathcal{F}_t)_{t\leq T}$  and carrying a standard n–dimensional Brownian motion  $(B_t)_{t\leq T}$ . The time horizon  $T$  shall vary along the article, most of the time it will be finite. By standard we mean that  $(B_t)$  starts from 0 and has quadratic covariation given by  $[B]_t = tI_n$  for all  $t \leq T$ . Let  $\mathbb H$  be the Cameron-Martin space, namely the space of absolutely continuous paths  $u: [0, T] \to \mathbb{R}^n$ , starting from 0 and equipped with the norm

$$
||u||_{\mathbb{H}} = \left(\int_0^T |\dot{u}_s|^2 ds\right)^{1/2},
$$

where  $|\dot{u}_s|$  denotes the Euclidean norm of the derivative of u at time s. In this context a *drift* is a process which is adapted to the filtration  $(\mathcal{F}_t)$  and which belongs to  $\mathbb H$  almost surely. Let  $\gamma_n$  be the standard Gaussian measure on  $\mathbb{R}^n$ . The starting point of the present article is the so called Borell formula: If  $f: \mathbb{R}^n \to \mathbb{R}$  is measurable and bounded from below then

<span id="page-0-0"></span>
$$
\log\left(\int_{\mathbb{R}^n} e^f d\gamma_n\right) = \sup_U \left\{ \mathsf{E}\left[f(B_1 + U_1) - \frac{1}{2} ||U||_{\mathbb{H}}^2\right] \right\} \tag{1}
$$

where the supremum is taken over all drifts U (here the time horizon is  $T = 1$ ). Actually a more general formula holds true, where the function  $f$  is allowed to depend on the whole trajectory of  $(B_t)$  rather than just  $B_1$ . More precisely, let  $(W, \mathcal{B}, \gamma)$  be the *n*-dimensional Wiener space: W is the space of continuous paths from  $[0, T]$  to  $\mathbb{R}^n$ ,  $\mathcal{B}$  is the Borel  $\sigma$ -field associated to the topology given by the uniform convergence (uniform convergence on compact sets if  $T = +\infty$ ) and  $\gamma$  is the law of the standard Brownian motion. If  $F: \mathbb{W} \to \mathbb{R}$  is measurable and bounded from below then

<span id="page-0-1"></span>
$$
\log\left(\int_{\mathbb{W}} \mathbf{e}^F \, d\gamma\right) = \sup_{U} \left\{ \mathsf{E} \left[ F(B+U) - \frac{1}{2} ||U||_{\mathbb{H}}^2 \right] \right\} \tag{2}
$$

Of course  $(1)$  is retrieved by applying the latter formula to a functional F of the form  $F(w) = f(w_1)$ . Formula [\(2\)](#page-0-1) is due to Boué and Dupuis [\[6\]](#page-14-0), we refer to our previous work [\[13\]](#page-14-1) for more historical comments and an alternate proof of the formula. We are interested in the use of such formulas to prove functional inequalities. This was initiated by Borell in [\[5\]](#page-14-2), in which he proved [\(1\)](#page-0-0) and showed that it yields the Prékopa–Leindler inequality very easily. This was further developed in the author's works [\[13,](#page-14-1) [14\]](#page-14-3) where many other functional inequalities were derived from [\(1\)](#page-0-0) and [\(2\)](#page-0-1). The main purpose of the present article is to establish a version of Borell's formula [\(1\)](#page-0-0) for the Brownian motion on a Riemannian manifold and to give a couple of applications, including a new proof of the Brascamp–Lieb inequality on the sphere of Carlen, Lieb and Loss.

### <span id="page-1-1"></span>2 Borell's formula for a diffusion

Let  $\sigma: \mathbb{R}^n \to M_n(\mathbb{R})$ , let  $b: \mathbb{R}^n \to \mathbb{R}^n$  and assume that the stochastic differential equation

<span id="page-1-0"></span>
$$
dX_t = \sigma(X_t) dB_t + b(X_t) dt \tag{3}
$$

has a unique strong solution. We also assume for simplicity that the explosion time is  $+\infty$ . Then there exists a measurable functional

$$
G\colon\mathbb{R}^n\times\mathbb{W}\to\mathbb{W}
$$

(it is probably safer to complete the  $\sigma$ -field  $\beta$  at this stage) such that for every  $x \in \mathbb{R}^n$  the process  $X = G(x, B)$  is the unique solution of [\(3\)](#page-1-0) starting from x. This hypothesis is satisfied in particular if  $\sigma$  and b are locally Lipschitz and grow at most linearly, see for instance [\[11,](#page-14-4) Chapter IV]. The process  $(X_t)$  is then a diffusion with generator  $L$  given by

$$
Lf = \frac{1}{2} \langle \sigma \sigma^T, \nabla^2 f \rangle + \langle b, \nabla f \rangle.
$$

for every  $\mathcal{C}^2$ -smooth function f. We denote the associated semigroup by  $(P_t)$ : for any test function  $f$ 

$$
P_t f(x) = \mathsf{E}_x \left[ f(X_t) \right],
$$

where as usual the subscript x denotes the starting point of  $(X_t)$ . Fix a finite time horizon T. Fix  $x \in \mathbb{R}^n$ , let  $f: R^n \to \mathbb{R}$  and assume that f is bounded from below. Applying the representation formula [\(2\)](#page-0-1) to the functional

$$
F \colon w \in \mathbb{W} \mapsto f(G(x, w)_T)
$$

we get

$$
\log \int_\mathbb{W} \mathrm{e}^{f(G(x,w)_{T})} \, \gamma(dw) = \sup_{U} \left\{ \mathsf{E}\left[f\left(G(x,B+U)_{T}\right) - \frac{1}{2} \|U\|_{\mathbb{H}}^{2}\right] \right\}
$$

where the supremum is taken on all drifts  $U$ . By definition of the semigroup  $(P_t)$  we have

$$
\log \int_{\mathbb{W}} e^{f(G(x,w)_T)} \gamma(dw) = \log P_T(e^f)(x).
$$

Also, we have the following lemma.

<span id="page-2-1"></span>**Lemma 1.** Let  $(U_t)_{t\leq T}$  be a drift. The process  $X^U = G(x, B+U)$  is the unique process satisfying

<span id="page-2-0"></span>
$$
X_t^U = x + \int_0^t \sigma(X_s^U) \, (dB_s + dU_s) + \int_0^t b(X_s^U) \, ds, \quad t \le T \tag{4}
$$

almost surely.

*Proof.* Assume first that  $||U||_{\mathbb{H}}$  is bounded. Then by Novikov's criterion the process  $(D_t)$  given by

$$
D_t = \exp\left(-\int_0^t \langle \dot{U}_s, dB_s \rangle - \frac{1}{2} \int_0^t |\dot{U}_s|^2 ds\right)
$$

is a uniformly integrable martingale. Moreover, according to Girsanov's formula, under the measure Q given by  $dQ = D_T dP$ , the process  $B + U$  is a standard Brownian motion on  $[0, T]$ , see for instance [\[12,](#page-14-5) section 3.5] for more details. Now since the stochastic differential equation [\(3\)](#page-1-0) is assumed to have a unique strong solution and by definition of  $G$ , almost surely for  $Q$ , the unique process satisfying [\(4\)](#page-2-0) is  $X^U = G(x, B + U)$ . Since Q and P are equivalent this is the result. For general  $U$ , the result follows by applying the bounded case to  $(U_t) = (U_{t \wedge T_n})$  where  $T_n$  is the stopping time

$$
T_n = \inf \left\{ t \ge 0 : \int_0^t |\dot{U}_s|^2 ds \ge n \right\},\,
$$

 $\Box$ 

and letting *n* tend to  $+\infty$ .

To sum up, we have established the following result.

<span id="page-2-2"></span>**Theorem 2.** For any function  $f: \mathbb{R}^n \to \mathbb{R}$  bounded from below we have

$$
\log P_T(e^f)(x) = \sup_U \left\{ \mathsf{E}\left[f(X_T^U) - \frac{1}{2} ||U||_{\mathbb{H}}^2\right] \right\},\,
$$

where the supremum is taken over all drifts U and the process  $X^U$  is the unique solution of [\(4\)](#page-2-0).

Remarks. This direct consequence of the representation formula [\(2\)](#page-0-1) was already noted by Boué and Dupuis. They used it to recover Freidlin and Wentzell's large deviation principle as the diffusion coefficient is sent to 0. Let us also note that the non explosion hypothesis is not essential. One can consider  $\mathbb{R}^n \cup \{\infty\}$ , the one point compactification of  $\mathbb{R}^n$ , set  $X_t = \infty$  after explosion time, and restrict to functions  $f$  that tend to  $0$  at infinity. In the same way we could also deal with a Dirichlet boundary condition.

#### <span id="page-3-2"></span>3 Borell's formula on a Riemannian manifold

Let  $(M, g)$  be a complete Riemannian manifold of dimension n. In this section we wish to establish a Borell type formula for the Brownian motion on  $M$ . To do so we need to recall first the intrinsic construction of the Brownian motion on M.

Let us start with some definitions from Riemannian geometry. Recall that the orthonormal frame bundle  $\mathcal{O}(M)$  is the set of  $(n + 1)$ –tuples

$$
\phi = (x, \phi^1, \dots, \phi^n)
$$

where x is in M and  $(\phi^1, \ldots, \phi^n)$  is an orthonormal basis of  $T_x(M)$ . Given an element  $\phi = (x, \phi^1, \dots, \phi^n)$  of  $\mathcal{O}(M)$  and a vector  $v \in T_x(M)$ , the *horizontal* lift of v at  $\phi$ , denoted  $\mathcal{H}(v)$ , is an element of  $T_{\phi}(\mathcal{O}(M))$  defined as follows: Choose a curve  $(x_t)$  starting from x with speed v and for  $i \leq n$  let  $\phi_t^i$  be the parallel translation of  $\phi^i$  along  $(x_t)$ . Since parallel translation preserves the inner product, we thus obtain a smooth curve  $(\phi_t)$  on  $\mathcal{O}(M)$  and we can set

$$
\mathcal{H}(v)=\dot{\phi}_0.
$$

This is a lift of  $v$  in the sense that for any smooth  $f$  on  $M$ 

$$
\mathcal{H}(v)(f\circ\pi)=v(f),
$$

where  $\pi: \mathcal{O}(M) \to M$  is the canonical projection. Now for  $i \leq n$  we define a vector field on  $\mathcal{O}(M)$  by setting

$$
\mathcal{H}^i(x,\phi^1,\ldots,\phi^n)=\mathcal{H}(\phi^i).
$$

The operator

$$
\Delta_{\mathcal{H}} = \sum_{i=1}^n (\mathcal{H}^i)^2
$$

is called the horizontal Laplacian. It is related to the Laplace–Beltrami operator on M, denoted  $\Delta$ , through the following commutation property: for any smooth  $f$  on  $M$  we have

<span id="page-3-1"></span>
$$
\Delta_{\mathcal{H}}(f \circ \pi) = \Delta(f) \circ \pi.
$$
 (5)

Note that the horizontal Laplacian is by definition a sum of squares of vector fields, and that this is typically not the case for the Laplace–Beltrami operator. We are now in a position to define the horizontal Brownian motion on  $\mathcal{O}(M)$ . Let

$$
B_t = (B_t^1, \ldots, B_t^n)
$$

be a standard *n*-dimensional Brownian motion. We consider the following stochastic differential equation on  $\mathcal{O}(M)$ 

<span id="page-3-0"></span>
$$
d\Phi_t = \sum_{i=1}^d \mathcal{H}^i(\Phi_t) \circ dB_t^i.
$$
 (6)

Throughout the rest of the article, the notation  $H \circ dM$  denotes the Stratonovitch integral. The equation [\(6\)](#page-3-0) is a short way of saying that for any smooth function g on  $\mathcal{O}(M)$  we have

$$
g(\Phi_t) = g(\Phi_0) + \sum_{i=1}^n \int_0^t \mathcal{H}^i(g)(\Phi_t) \circ dB_t^i.
$$

This always has a strong solution, see [\[11,](#page-14-4) Theorem V.1.1.]. Let us assume additionally that it does not explode in finite time. This is the case in partic-ular if the Ricci curvature of M is bounded from below, see for instance [\[10,](#page-14-6) section 4.2], where a more precise criterion is given. Translating the equation above in terms of Itô increments we easily get

$$
dg(\Phi_t) = \sum_{i=1}^n \mathcal{H}^i(g)(\Phi_t) dB_t^i + \frac{1}{2} \Delta_{\mathcal{H}} g(\Phi_t) dt.
$$

Let  $(X_t)$  be the process given by  $X_t = \pi(\Phi_t)$ . Applying the previous formula and  $(5)$  we see that for any smooth f on M

<span id="page-4-0"></span>
$$
df(X_t) = \sum_{i=1}^{n} \Phi_t^i(f)(X_t) dB_t^i + \frac{1}{2} \Delta f(X_t) dt.
$$
 (7)

In particular

$$
f(X_t) - \frac{1}{2} \int_0^t \Delta f(X_s) ds, \quad t \ge 0
$$

is a local martingale. This shows that  $(X_t)$  is a Brownian motion on M. The process  $(X_t)$  is called the *stochastic development* of  $(B_t)$ . In the sequel, it will be convenient to identify the orthogonal basis  $\Phi_t^1, \ldots, \Phi_t^n$  with the orthogonal map

$$
x \in \mathbb{R}^n \to \sum_{i=1}^n x_i \Phi_t^i \in T_{X_t}(M).
$$

Then the equation [\(7\)](#page-4-0) can be rewritten

$$
df(X_t) = \langle \nabla f(X_t), \Phi_t dB_t \rangle + \frac{1}{2} \Delta f(X_t) dt.
$$

Similarly the equation [\(6\)](#page-3-0) can be rewritten in a more concise form

<span id="page-4-1"></span>
$$
d\Phi_t = \mathcal{H}(\Phi_t) \circ dB_t.
$$
 (8)

To sum up, the process  $(\Phi_t)$  is an orthonormal basis above  $(X_t)$  which is used to map the Brownian increment  $dB_t$  from  $\mathbb{R}^n$  to the tangent space of M at  $X_t$ . Now we establish a Borell type formula for the process  $(X_t)$ . We know that there exists a measurable functional

$$
G\colon \mathcal{O}(M)\times\mathbb{W}\to \mathcal{C}(\mathbb{R},\mathcal{O}(M))
$$

such that the process  $\Phi = G(\phi, B)$  is the unique solution of [\(8\)](#page-4-1) starting from  $\phi$ . Let  $\phi \in \mathcal{O}(M)$ , let  $T > 0$ , let  $f : M \to \mathbb{R}$  and assume that f is bounded from below. Applying the representation formula [\(2\)](#page-0-1) to the functional

$$
F \colon w \in \mathbb{W} \mapsto f \circ \pi \left( G(\phi, w)_T \right)
$$

we get

$$
\log \left( \int_{\mathbb{W}} e^{f \circ \pi (G(\phi, B)_{T})} d\gamma \right) = \sup_{U} \left\{ \mathsf{E} \left[ f \circ \pi (G(\phi, B + U)_{T}) - \frac{1}{2} ||U||_{\mathbb{H}}^{2} \right] \right\}.
$$

Let  $x = \pi(\phi)$ . Since  $\pi(G(\phi, B))$  is a Brownian motion on M starting from x we have

$$
\log \left( \int_{\mathbb{W}} e^{f \circ \pi (G(\phi, B)_{T})} d\gamma \right) = \log P_{T}(e^{f})(x),
$$

where  $(P_t)$  is the heat semigroup on M. Also, letting  $\Phi^U = G(\phi, B + U)$  and reasoning along the same lines as in the proof of Lemma [1,](#page-2-1) we obtain that  $\Phi^U$ is the only solution to

$$
d\Phi_t^U = \mathcal{H}(\Phi_t^U) \circ (dB_t + dU_t)
$$

starting from  $\phi$ . We also let  $X^U = \pi(\Phi^U)$  and call this process the stochastic development of  $B + U$  starting from  $\phi$ . To sum up, we have established the following result.

<span id="page-5-0"></span>**Theorem 3.** Let  $f: M \to \mathbb{R}$ , let  $\phi \in \mathcal{O}(M)$ , let  $x = \pi(\phi)$  and let  $T > 0$ . If f is bounded from below then

$$
\log P_T\left(e^f\right)(x) = \sup_U \left\{ \mathsf{E}\left[f(X_T^U) - \frac{1}{2} ||U||_{\mathbb{H}}^2\right] \right\},\,
$$

where the supremum is taken on all drifts U and where given a drift U, the process  $X^U$  is the stochastic development of  $B+U$  starting from  $\phi$ .

## 4 Brascamp–Lieb inequality on the sphere

In the article [\[14\]](#page-14-3), we explained how to derive the Brascamp–Lieb inequality and its reversed version from Borell's formula. In this section we extend this to the sphere, and give a proof based on Theorem [3](#page-5-0) of the spherical version of the Brascamp–Lieb inequality, due to Carlen, Lieb and Loss in [\[7\]](#page-14-7).

<span id="page-5-1"></span>**Theorem 4.** Let  $g_1, \ldots, g_{n+1}$  be non–negative functions on the interval  $[-1, 1]$ . Let  $\sigma_n$  be the Haar measure on the sphere  $\mathbb{S}^n$ , normalized to be a probability measure. We have

$$
\int_{\mathbb{S}^n} \prod_{i=1}^{n+1} g_i(x_i) \, \sigma_n(dx) \le \prod_{i=1}^{n+1} \left( \int_{\mathbb{S}^n} g_i(x_i)^2 \, \sigma_n(dx) \right)^{1/2}
$$

Remark. The inequality does not hold if we replace the  $L^2$  norm in the righthand side by a smaller  $L^p$  norm, like the  $L^1$  norm. Somehow this 2 accounts for the fact that the coordinates of a uniform random vector on  $\mathbb{S}^n$  are not independent. We refer to the introduction of [\[7\]](#page-14-7) for a deeper insight on this inequality.

In addition to the Borell type formulas established in the previous two sections, our proof relies on a sole inequality, spelled out in the lemma below. Let  $P_i: \mathbb{S}^n \to [-1,1]$  be the application that maps x to its i–th coordinate  $x_i$ . The spherical gradient of  $P_i$  at x is the projection of the coordinate vector  $e_i$  onto  $x^{\perp}$ :

$$
\nabla P_i(x) = e_i - x_i x.
$$

<span id="page-6-1"></span>**Lemma 5.** Let  $x \in \mathbb{S}^n$  and let  $y \in x^{\perp}$ . For  $i \leq n+1$ , if  $\nabla P_i(x) \neq 0$  let

$$
\theta^i = \frac{\nabla P_i(x)}{|\nabla P_i(x)|}
$$

and let  $\theta_i$  be an arbitrary unit vector of  $x^{\perp}$  otherwise. Then for any  $y \in x^{\perp}$  we have

$$
\sum_{i=1}^{n+1} \langle \theta^i, y \rangle^2 \le 2|y|^2.
$$

*Proof.* Assume first that  $\nabla P_i(x) \neq 0$  for every i. Since  $\nabla P_i(x) = e_i - x_i x$  and  $y$  is orthogonal to  $x$  we then have

$$
\langle \theta^i,y\rangle^2=y_i^2+x_i^2\langle \theta^i,y\rangle^2\leq y_i^2+x_i^2|y|^2.
$$

Summing this over  $i \leq n+1$  yields the result. On the other hand, if there exists i such that  $\nabla P_i(x) = 0$  then  $x = \pm e_i$  and it is almost immediate to check that the desired inequality holds true.  $\Box$ 

*Proof of Theorem [4.](#page-5-1)* Let us start by describing the behaviour of a given coordinate of a Brownian motion on  $\mathbb{S}^n$ . Let  $(B_t)$  be a standard Brownian motion on  $\mathbb{R}^n$ , let  $\phi$  be a fixed element of  $\mathcal{O}(\mathbb{S}^n)$  and let  $(\Phi_t)$  be the horizontal Brownian motion given by

$$
\Phi_0 = \phi
$$
 and  $d\Phi_t = \mathcal{H}(\Phi_t) \circ dB_t$ .

We also let  $X_t = \pi(\Phi_t)$  be the stochastic development of  $(B_t)$  and  $X_t^i = P_i(X_t)$ , for every  $i \leq n + 1$ . We have

<span id="page-6-0"></span>
$$
dX_t^i = \langle \nabla P_i(X_t), \Phi_t dB_t \rangle + \frac{1}{2} \Delta P_i(X_t) dt.
$$
 (9)

Let  $\theta$  be an arbitrary unit vector of  $\mathbb{R}^n$  and define a process  $(\theta_t^i)$  by

$$
\theta_t^i = \begin{cases} \Phi_t^* \left( \frac{\nabla P_i(X_t)}{|\nabla P_i(X_t)|} \right), & \text{if } \nabla P_i(X_t) \neq 0, \\ \theta & \text{otherwise.} \end{cases}
$$

Since  $\Phi_t$  is an orthogonal map  $\theta_t$  belongs to the unit sphere of  $\mathbb{R}^n$ . Consequently the process  $(W_t^i)$  defined by  $dW_t^i = \langle \theta_t^i, dW_t \rangle$  is a one dimensional standard Brownian motion. Observe that  $|\nabla P_i| = (1 - P_i^2)^{1/2}$  and recall that  $P_i$  is an eigenfunction for the spherical Laplacian:  $\Delta P_i = -nP_i$ . Equality [\(9\)](#page-6-0) becomes

$$
dX_t^i = \left(1 - (X_t^i)^2\right)^{1/2} dW_t^i - \frac{n}{2} X_t^i dt.
$$

This stochastic differential equation is usually referred to as the Jacobi diffusion in the literature, see for instance [\[2,](#page-14-8) section 2.7.4]. What matters for us is that it does possess a unique strong solution. Indeed the drift term is linear and although the diffusion factor  $(1 - x^2)^{1/2}$  is not locally Lipschitz, it is Hölder continuous with exponent  $1/2$ , which is sufficient to insure strong uniqueness in dimension 1, see for instance [\[15,](#page-15-0) section V.40]. Let  $(Q_t)$  be the semigroup associated to the process  $(X_t^i)$ . The stationary measure  $\nu_n$  is easily seen to be given by

$$
\nu_n(dt) = c_n 1\!\!1_{[-1,1]}(t)(1-t^2)^{\frac{n}{2}-1} dt,
$$

where  $c_n$  is the normalization constant. Obviously  $\nu_n$  coincides with the pushforward of  $\sigma_n$  by  $P_i$ .

We now turn to the actual proof of the theorem. Let  $g_1, \ldots, g_{n+1}$  be non negative functions on  $[-1, 1]$  and assume (without loss of generality) that they are bounded away from 0. Let  $f_i = \log(g_i)$  for all i and let

$$
f \colon x \in \mathbb{S}^n \mapsto \sum_{i=1}^{n+1} f_i(x_i).
$$

The functions  $f_i, f$  are bounded from below. Fix a time horizon T, let U be a drift and let  $(\Phi_t^U)$  be the process given by

$$
\begin{cases} \Phi_0^U = \phi \\ d\Phi_t^U = \mathcal{H}(\Phi_t^U) \circ (dB_t + dU_t), \quad t \leq T. \end{cases}
$$

We also let  $X_t^U = \pi(\Phi_t^U)$  be the stochastic development of  $B + U$ . These processes are well defined by the results of the previous section. We want to bound  $f(X_T^U) - \frac{1}{2} ||U||_{\mathbb{H}}^2$  from above. By definition

$$
f(X_T^U) = \sum_{i=1}^{n+1} f_i(P_i X_T^U)).
$$

Let  $(\theta_t^i)$  be the process given by

$$
\theta_t^i = \Phi_t^*\left(\frac{\nabla P_i(X_t^U)}{|\nabla P_i(X_t^U)|}\right)
$$

(again replace this by an arbitrary fixed unit vector if  $\nabla P_i(X_t^U) = 0$ ). Then let  $(W_t^i)$  be the one dimensional Brownian motion given by  $dW_t^i = \langle \theta_t^i, dW_t \rangle$  and let  $(U_t^i)$  be the one dimensional drift given by  $dU_t^i = \langle \theta_t^i, dU_t \rangle$ . The process  $(P_i(X_t^U))$  then satisfies

<span id="page-8-0"></span>
$$
dP_i(X_t^U) = \left(1 - P_i(X_t^U)^2\right)^{1/2} \left(dW_t^i + dU_t^i\right) - \frac{n}{2} P_i(X_t^U) dt. \tag{10}
$$

Applying Lemma [5,](#page-6-1) we easily get

$$
\sum_{i=1}^{n+1} \|U^i\|_{\mathbb{H}}^2 \le 2\|U\|_{\mathbb{H}}^2,
$$

almost surely (note that in the left hand side of the inequality  $\mathbb H$  is the Cameron– Martin space of  $\mathbb R$  rather than  $\mathbb R^n$ ). Therefore

<span id="page-8-1"></span>
$$
f(X_T^U) - \frac{1}{2} ||U||_{\mathbb{H}}^2 \le \sum_{i=1}^{n+1} \left( f_i(P_i(X_T^U)) - \frac{1}{4} ||U^i||_{\mathbb{H}}^2 \right).
$$
 (11)

Recall [\(10\)](#page-8-0) and apply Theorem [2](#page-2-2) to the semigroup  $(Q_t)$  and to the function  $2f_i$ rather than  $f_i$ . This gives

$$
\mathsf{E}\left[f_i(P_i(X_t^U)) - \frac{1}{4} ||U^i||_{\mathbb{H}}^2\right] \le \frac{1}{2}\log Q_T(e^{2f_i})(x_i),
$$

for every  $i \leq n+1$ . Taking expectation in [\(11\)](#page-8-1) thus yields

$$
\mathsf{E}\left[f(X_T^U) - \frac{1}{2} ||U||_{\mathbb{H}}^2\right] \leq \frac{1}{2} \sum_{i=1}^{n+1} \log Q_T\left(e^{2f_i}\right)(x_i).
$$

Taking the supremum over all drifts  $U$  and using Theorem [3](#page-5-0) we finally obtain

$$
P_T(e^f)(x) \leq \prod_{i=1}^{n+1} (Q_T(e^{2f_i})(x_i))^{1/2}.
$$

The semigroup  $(P_t)$  is ergodic and converges to  $\sigma_n$  as t tends to  $+\infty$ . Similarly  $(Q_t)$  converges to  $\nu_n$ . So letting T tend to  $+\infty$  in the previous inequality gives

$$
\int_{\mathbb{S}^n} e^f d\sigma_n \le \prod_{i=1}^{n+1} \left( \int_{[-1,1]} e^{2f_i} d\nu_n \right)^{1/2},
$$

 $1/2$ 

which is the result.

Remark. Barthe, Cordero–Erausquin and Maurey in [\[3\]](#page-14-9) and together with Ledoux in [\[4\]](#page-14-10) gave several extensions of Theorem [4.](#page-5-1) The method exposed here also allows to recover most of their results. We chose to stick to the original statement of Carlen, Lieb and Loss for simplicity.

 $\Box$ 

#### 5 The dual formula

In [\[13\]](#page-14-1), we established a dual version of Borell's formula [\(1\)](#page-0-0). It states as follows: If  $\mu$  is an absolutely continuous measure on  $\mathbb{R}^n$  satisfying some (reasonable) technical assumptions, the relative entropy of  $\mu$  with respect to the Gaussian measure is given by the following formula

<span id="page-9-0"></span>
$$
H(\mu \mid \gamma_n) = \inf \left\{ \frac{1}{2} \mathsf{E} \left[ \|U\|_{\mathbb{H}}^2 \right] \right\} \tag{12}
$$

where the infimum is taken over all drifts U such that  $B_1 + U_1$  has law  $\mu$ . Informally this says that the minimal energy needed to constrain the Brownian motion to have a prescribed law at time 1 coincides with the relative entropy of this law with respect to  $\gamma_n$ . The infimum is actually a minimum and the optimal drift can be described as follows. We let f be the density of  $\mu$  with respect to  $\gamma_n$  and  $(P_t)$  be the heat semigroup on  $\mathbb{R}^n$ . The following stochastic differential equation

$$
\begin{cases}\nX_0 = 0 \\
dX_t = dB_t + \nabla \log P_{1-t} f(X_t) dt, \quad t \le 1.\n\end{cases}
$$
\n(13)

has a unique strong solution. The solution satisfies  $X_1 = \mu$  in law and is optimal in the sense that there is equality in  $(12)$  for the drift U given by  $\tilde{U}_t = \nabla \log P_{1-t} f(X_t)$ . The purpose of this section is to describe the Riemannian counterpart of [\(12\)](#page-9-0). There is also a version for diffusions such as the ones considered in section [2](#page-1-1) but we shall omit it for the sake of brevity.

The setting of this section is the same as that of section [3:](#page-3-2)  $(M, q)$  is a complete Riemannian manifold of dimension  $n$  whose Ricci curvature is bounded from below and  $(B_t)$  is a standard Brownian motion on  $\mathbb{R}^n$ . We denote the heat semigroup on M by  $(P_t)$ .

**Theorem 6.** Fix  $x \in M$  and a time horizon T. Let  $\mu$  be a probability measure on M, assume that  $\mu$  is absolutely continuous with respect to  $\delta_x P_T$  and let f be its density. If f is Lipschitz and bounded away from 0 then

$$
H(\mu \mid \delta_x P_T) = \inf \left\{ \frac{1}{2} \mathsf{E} \left[ ||U||_{\mathbb{H}}^2 \right] \right\}
$$

where the infimum is taken all drifts U such that the stochastic development of  $B + U$  starting from x has law  $\mu$  at time T.

Proving that any drift satisfying the constraint has energy at least as large as the relative entropy of  $\mu$  is a straightforward adaptation of Proposition 1 from [\[13\]](#page-14-1), and we shall leave this to the reader. Alternatively, one can use Theorem [3](#page-5-0) and combine it with the following variational formula for the entropy:

$$
H(\mu \mid \delta_x P_T) = \sup_f \left\{ \int_M f \, d\mu - \log P_T(e^f)(x) \right\}
$$

(again details are left to the reader). Besides, as in the Euclidean case, there is actually an optimal drift, whose energy is exactly the relative entropy of  $\mu$ . This is the purpose of the next result.

**Theorem 7.** Let  $\phi$  be a fixed element of  $\mathcal{O}(M)$ . Let  $x = \pi(\phi)$  and let T be a time horizon. Let  $\mu$  have density f with respect to  $\delta_x P_T$ , and assume that f is Lipschitz and bounded away from 0. The stochastic differential equation

<span id="page-10-0"></span>
$$
\begin{cases} \n\Phi_0 = \phi \\ \nd\Phi_t = \mathcal{H}(\Phi_t) \circ (dB_t + \Phi_t^* \nabla \log P_{T-t} f(Y_t)) \\ \nY_t = \pi(\Phi_t) \n\end{cases} \n(14)
$$

has a unique strong solution on  $[0, T]$ . The law of the process  $(Y_t)$  is given by the following formula: For every functional  $H: \mathcal{C}([0,T];M) \to \mathbb{R}$  we have

<span id="page-10-2"></span>
$$
\mathsf{E}[H(Y)] = \mathsf{E}[H(X)f(X_1)],\tag{15}
$$

where  $(X_t)$  is a Brownian motion on M starting from x. In other words  $Y_T$  has law  $\mu$  and the bridges of Y equal those of the Brownian motion on M, in law. Moreover, letting U be the drift given by

<span id="page-10-1"></span>
$$
U_t = \int_0^t \Phi_s^* \nabla \log P_{T-s} f(Y_s) ds, \quad t \le T,
$$
\n(16)

we have

$$
H(\mu \mid \delta_x P_T) = \frac{1}{2} \mathsf{E} \left[ \|U\|_{\mathbb{H}}^2 \right].
$$

*Proof.* Since Ric  $\geq -\lambda g$ , we have the following estimate for the Lipschitz norm of  $f$ :

$$
||P_t f||_{\text{Lip}} \leq e^{\lambda t/2} ||f||_{\text{Lip}}.
$$

One way to see this is to use Kendall's coupling for Brownian motions on a manifold, see for instance [\[10,](#page-14-6) section 6.5]. Alternatively, it is easily derived from the commutation property  $|\nabla P_t f|^2 \leq e^{\lambda t} P_t(|\nabla f|^2)$  which, in turn, follows from Bochner's formula, see [\[2,](#page-14-8) Theorem 3.2.3]. Recall that  $f$  is assumed to be bounded away from 0, and for every  $t \leq T$  let  $F_t = \log P_{T-t}f$ . Then  $(t, x) \mapsto \nabla F_t(x)$  is smooth and bounded on  $[0, T] \times M$ , which is enough to insure the existence of a unique strong solution to [\(14\)](#page-10-0). Besides an easy computation shows that

<span id="page-10-3"></span>
$$
\partial_t F_t = -\frac{1}{2} (\Delta F_t + |\nabla F_t|^2). \tag{17}
$$

Then using  $(14)$  and Itô's formula we get

$$
dF(t, Y_t) = \langle \nabla F_t(Y_t), \Phi_t dB_t \rangle + \frac{1}{2} |\nabla F_t(Y_t)|^2 dt.
$$
  
=  $\langle \dot{U}_t, dB_t \rangle + \frac{1}{2} |\dot{U}_t|^2 dt$ 

(recall the definition  $(16)$  of U). Therefore

<span id="page-11-0"></span>
$$
\frac{1}{f(Y_T)} = e^{-F_T(Y_T)} = \exp\left(-\int_0^T \langle \dot{U}_t, dB_t \rangle - \frac{1}{2} ||U||_{\mathbb{H}}^2\right).
$$
 (18)

Observe that the variable  $||U||_{\mathbb{H}}$  is bounded (just because  $\nabla F$  is bounded). So Girsanov's formula applies:  $1/f(Y_T)$  has expectation 1 and under the measure Q given by  $dQ = (1/f(Y_T)) dP$  the process  $B+U$  is a standard Brownian motion on  $\mathbb{R}^n$ . Since Y is the stochastic development of  $B + U$  starting from x, this shows that under  $Q$  the process Y is a Brownian motion on M starting from x. This is a mere reformulation of  $(15)$ . For the entropy equality observe that since  $Y_T$  has law  $\mu$ , we have

$$
H(\mu \mid \delta_x P_T) = \mathsf{E}[\log f(Y_T)].
$$

Using [\(18\)](#page-11-0) again and the fact that  $\int \langle \dot{U}_t, dB_t \rangle$  is a martingale we get the desired equality.  $\Box$ 

To conclude this article, let us derive from this formula the log–Sobolev inequality for a manifold having a positive lower bound on its Ricci curvature. This is of course well–known, but our point is only to illustrate how the previous theorem can be used to prove inequalities. Recall the definition of the Fisher information: if  $\mu$  is a probability measure on M having Lipschitz and positive density  $f$  with respect to some reference measure  $m$ , the relative Fisher information of  $\mu$  with respect to  $m$  is defined by

$$
I(\mu \mid m) = \int_M \frac{|\nabla f|^2}{f} dm = \int_M |\nabla \log f|^2 d\mu.
$$

By Bishop's Theorem, if Ric  $\geq \kappa g$  pointwise for some positive  $\kappa$  then the volume measure on  $M$  is finite. We let  $m$  be the volume measure normalized to be a probability measure.

**Theorem 8.** If Ric  $\geq \kappa g$  pointwise for some  $\kappa > 0$ , then for any probability measure  $\mu$  on M having a Lipschitz and positive density with respect to m we have

<span id="page-11-3"></span>
$$
H(\mu \mid m) \le \frac{n}{2} \log \left( 1 + \frac{I(\mu \mid m)}{n \kappa} \right). \tag{19}
$$

Remarks. Since  $log(1 + x) \leq x$  this inequality is a dimensional improvement of the more familiar inequality

<span id="page-11-1"></span>
$$
H(\mu \mid m) \le \frac{1}{\kappa} I(\mu \mid m). \tag{20}
$$

It is known (see for instance [\[1,](#page-13-0) section 5.6]) that [\(20\)](#page-11-1) admits yet another sharp form that takes the dimension into account, namely

<span id="page-11-2"></span>
$$
H(\mu \mid m) \le \frac{n-1}{\kappa n} I(\mu \mid m),\tag{21}
$$

but we were not able to recover this one with our method. Note also that depending on the measure  $\mu$ , the right hand side of [\(21\)](#page-11-2) can be smaller than that of [\(19\)](#page-11-3), or the other way around.

*Proof.* By the Bonnet–Myers theorem M is compact. Fix  $x \in M$  and a time horizon T. Let  $p_T(x, \cdot)$  be the density of the measure  $\delta_x P_T$  with respect to m (in other words let  $(p_t)$  be the heat kernel on M). If  $d\mu = \rho dm$  then  $\mu$  has density  $f = \rho / p_T(x, \cdot)$  with respect to  $\delta_x P_T$ . Since  $p_T(x, \cdot)$  is smooth and positive (see for instance  $[8,$  chapter  $6]$ )  $f$  satisfies the technical assumptions of the previous theorem. Let  $F_t = \log P_{T-t} f$  and let  $(Y_t)$  be the process given by [\(14\)](#page-10-0). We know from the previous theorem that

<span id="page-12-2"></span>
$$
H(\mu \mid \delta_x P_T) = \frac{1}{2} \mathsf{E} \left[ \int_0^T |\nabla F_t(Y_t)|^2 dt \right]. \tag{22}
$$

Using [\(17\)](#page-10-3) we easily get

$$
\partial_t(|\nabla F|^2) = -\langle \nabla \Delta F, \nabla F \rangle - \langle \nabla |\nabla F|^2, \nabla F \rangle.
$$

Applying Itô's formula we obtain after some computations (omitting variables in the right hand side)

$$
d|\nabla F(t,Y_t)|^2 = \langle \nabla |\nabla F|^2, \Phi_t dB_t \rangle - \langle \nabla \Delta F, \nabla F \rangle dt + \frac{1}{2} \Delta |\nabla F|^2 dt.
$$

Now recall Bochner's formula

$$
\frac{1}{2}\Delta|\nabla F|^2 = \langle \nabla \Delta F, \nabla F \rangle + ||\nabla^2 F||_{HS}^2 + \text{Ric}(\nabla F, \nabla F).
$$

So that

<span id="page-12-0"></span>
$$
d|\nabla F(t, Y_t)|^2 = \langle \nabla |\nabla F|^2, \Phi_t dB_t \rangle + \|\nabla^2 F\|_{HS}^2 dt + \text{Ric}(\nabla F, \nabla F) dt. \tag{23}
$$

Since  $\nabla F$  is bounded and the Ricci curvature non negative, the local martingale part in the above equation is bounded from above. So by Fatou's lemma it is a sub–martingale, and its expectation is non decreasing. So taking expectation in [\(23\)](#page-12-0) and using the hypothesis Ric  $\geq \kappa g$  we get

<span id="page-12-1"></span>
$$
\frac{d}{dt} \mathsf{E}\left[|\nabla F_t(Y_t)|^2\right] \ge \mathsf{E}\left[\|\nabla^2 F_t(Y_t)\|_{HS}^2\right] + \kappa \mathsf{E}\left[|\nabla F_t(Y_t)|^2\right].\tag{24}
$$

Throwing away the Hessian term would lead us to the inequality [\(20\)](#page-11-1). Let us exploit this term instead. Using Cauchy-Schwartz and Jensen's inequality we get

$$
\mathsf{E}\left[\|\nabla^2 F_t(Y_t)\|_{HS}^2\right] \geq \frac{1}{n} \mathsf{E}\left[\Delta F_t(Y_t)^2\right] \geq \frac{1}{n} \mathsf{E}[\Delta F_t(Y_t)]^2.
$$

Also, by [\(15\)](#page-10-2) and recalling that  $(P_t)$  is the heat semigroup on M we obtain

$$
\mathsf{E}\left[\Delta F_t(Y_t) + |\nabla F_t(Y_t)|^2\right] = \mathsf{E}\left[\frac{\Delta P_{T-t}f(Y_t)}{P_{T-t}f(Y_t)}\right] = \mathsf{E}\left[\frac{\Delta P_{T-t}f(X_t)}{P_{T-t}f(X_t)}f(X_T)\right]
$$

$$
= \mathsf{E}\left[\Delta P_{T-t}f(X_t)\right] = \Delta P_T(f)(x).
$$

Letting  $\alpha(t) = \mathsf{E} \left[ |\nabla F_t(Y_t)|^2 \right]$  and  $C_T = \Delta P_T f(x)$  we thus get from [\(24\)](#page-12-1)

$$
\alpha'(t) \ge \kappa \alpha(t) + \frac{1}{n} (\alpha(t) - C_T)^2
$$

$$
\ge \frac{1}{n} \alpha(t) (\alpha(t) + n\kappa - 2C_T)
$$

Since  $P_T f$  tends to a constant function as T tends to  $+\infty$ ,  $C_T$  tends to 0. So if T is large enough  $n\kappa - 2C_T$  is positive and the differential inequality above yields

$$
\alpha(t) \le \frac{n \kappa(T) \alpha(T)}{e^{\kappa(T)(T-t)} \left(n \kappa(T) + \alpha(T)\right) - \alpha(T)}, \quad t \le T
$$

where  $\kappa(T) = \kappa - 2C_T/n$ . Integrating this between 0 and T we get

<span id="page-13-1"></span>
$$
\int_0^T \alpha(t) dt \le n \log \left( 1 + \frac{\alpha(T)(1 - e^{-\kappa(T)T})}{n \kappa(T)} \right). \tag{25}
$$

Observe that  $\kappa(T) \to \kappa$  as T tends to  $+\infty$ . By [\(22\)](#page-12-2) and since  $(\delta_x P_t)$  converges to  $m$  measure on  $M$  we have

$$
\int_0^T \alpha(t) dt \to 2 \,\mathrm{H}(\mu \mid m),
$$

as T tends to  $+\infty$ . Also, since  $Y_T$  has law  $\mu$ 

$$
\alpha(T) = \mathsf{E}\left[|\nabla \log f(Y_T)|^2\right] = \mathbf{I}(\mu \mid \delta_x P_T) \to \mathbf{I}(\mu \mid m).
$$

Therefore, letting T tend to  $+\infty$  in [\(25\)](#page-13-1) yields

$$
H(\mu \mid m) \leq \frac{n}{2} \log \left( 1 + \frac{I(\mu \mid m)}{n \kappa} \right),
$$

which is the result.

Let us give an open problem to finish this article. We already mentioned that Borell recovered the Prékopa–Leindler inequality from  $(1)$ . It is natural to ask whether there is probabilistic proof of the Riemannian Prékopa–Leindler inequality of Cordero, McCann and Schmuckenschlger [\[9\]](#page-14-12) based on Theorem [3.](#page-5-0) Copying naively Borell's argument, we soon face the following difficulty: If  $X$ and Y are two Brownian motions on a manifold coupled by parallel transport, then unless the manifold is flat, the midpoint of  $X$  and  $Y$  is not a Brownian motion. We believe that there is a way around this but we could not find it so far.

#### <span id="page-13-0"></span>References

[1] C. Ané et al. Sur les inégalités de Sobolev logarithmiques. Panoramas et Synthèses 10. SMF, Paris, 2000.

口

- <span id="page-14-8"></span>[2] D. Bakry, I. Gentil and M. Ledoux. Analysis and geometry of Markov diffusion operators. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 348. Springer, Cham, 2014.
- <span id="page-14-9"></span>[3] F. Barthe, D. Cordero–Erausquin and B. Maurey. Entropy of spherical marginals and related inequalities. J. Math. Pures Appl. (9) 86 (2006), no. 2, 89–99.
- <span id="page-14-10"></span>[4] F. Barthe, D. Cordero–Erausquin, M. Ledoux and B. Maurey. Correlation and Brascamp–Lieb inequalities for Markov semigroups. Int. Math. Res. Not. IMRN 2011, no. 10, 2177–2216.
- <span id="page-14-2"></span>[5] C. Borell. Diffusion equations and geometric inequalities, Potential Anal. 12 (2000), no 1, 49–71.
- <span id="page-14-0"></span>[6] M. Boué and P. Dupuis. A variational representation for certain functionals of Brownian motion. Ann. Probab. 26 (1998), no 4, 1641–1659.
- <span id="page-14-7"></span>[7] E.A. Carlen, E.H. Lieb and M. Loss. A sharp analog of Young's inequality on  $\mathbb{S}^N$  and related entropy inequalities. J. Geom. Anal. 14 (2004), no. 3, 487–520.
- <span id="page-14-11"></span>[8] I. Chavel. Eigenvalues in Riemannian geometry. Including a chapter by Burton Randol. With an appendix by Jozef Dodziuk. Pure and Applied Mathematics, 115. Academic Press, Inc., Orlando, FL, 1984.
- <span id="page-14-12"></span>[9] D. Cordero–Erausquin, R.J McCann and M. Schmuckenschlger. A Riemannian interpolation inequality à la Borell, Brascamp and Lieb. Invent. Math. 146 (2001), no. 2, 219–257.
- <span id="page-14-6"></span>[10] E.P. Hsu. Stochastic analysis on manifolds. Graduate Studies in Mathematics, 38. American Mathematical Society, Providence, RI, 2002.
- <span id="page-14-4"></span>[11] N. Ikeda and S. Watanabe. Stochastic differential equations and diffusion processes. Second edition. North–Holland Mathematical Library, 24. North–Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, 1989.
- <span id="page-14-5"></span>[12] I. Karatzas and S.E. Shreve. Brownian motion and stochastic calculus. Second edition. Graduate Texts in Mathematics, 113. Springer–Verlag, New York, 1991.
- <span id="page-14-1"></span>[13] J.Lehec, Representation formula for the entropy and functional inequalities. Ann. Inst. Henri Poincaré Probab. Stat. 49 (2013), no. 3, 885–899.
- <span id="page-14-3"></span>[14] J. Lehec, Short probabilistic proof of the Brascamp–Lieb and Barthe theorems. Canad. Math. Bull. 57 (2014), no. 3, 585–597.

<span id="page-15-0"></span>[15] L.C.G. Rogers and R. Williams. Diffusions, Markov processes, and martingales. Vol. 2. Itô calculus. Reprint of the second (1994) edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2000.