An integral inequality for the invariant measure of some finite dimensional stochastic differential equation

Giuseppe Da Prato Scuola Normale Superiore di Pisa

Dedicated to Björn Schmalfuss

Abstract

We prove an integral inequality for the invariant measure ν of a stochastic differential equation with additive noise in a finite dimensional space $H = \mathbb{R}^d$. As a consequence, we show that there exists the Fomin derivative of ν in any direction $z \in H$ and that it is given by $v_z = \langle D \log \rho, z \rangle$, where ρ is the density of ν with respect to the Lebesgue measure. Moreover, we prove that $v_z \in L^p(H,\nu)$ for any $p \in [1,\infty)$. Also we study some properties of the gradient operator in $L^p(H,\nu)$ and of his adjoint.

2000 Mathematics Subject Classification AMS: 60H07, 60H30, 37L40.

Key words: stochastic differential equations, invariant measure, Fomin derivative, gradient operator.

1 Introduction and preliminaries

In the recent paper [DaDe14] the following inequality involving the invariant measure ν of the Burgers equation was proved

$$\left| \int_{H} \langle RD\varphi, z \rangle \, d\nu \right| \le C_p \|\varphi\|_{L^p(H,\nu)} \, |z|, \tag{1.1}$$

for all $\varphi \in C_b^1(H)$, all $z \in H$ and all p > 1, R being a suitable negative power of the Laplace operator equipped with Dirichlet boundary conditions.

As noted in [DaDe14], by estimate (1.1) it follows that RD is closable in $L^p(H,\nu)$ for all p > 1. Moreover, for each $z \in H$ there exists $v_z \in L^p(H,\nu)$ such that

$$\int_{H} \langle RD\varphi, z \rangle \, d\nu = \int_{H} v_{z} \, \varphi \, d\nu, \quad \forall \, \varphi \in C_{b}^{1}(H).$$
(1.2)

Identity (1.2) implies that ν is Fomin differentiable in all directions of the range of R(H) of R. We recall that if $\nu = N_Q$ (the Gaussian measure of mean 0 and covariance Q) identity (1.2) is well known in Malliavin Calculus. In this case the adjoint $(Q^{1/2}D)^*$ of $Q^{1/2}D$ is called the Skorhood operator.

The aim of the present paper is to show that the inequality (1.1), with R replaced by the identity operator, can also be proved for the invariant measures of some stochastic differential equations in $H = \mathbb{R}^d$ of the form

$$\begin{cases} dX(t) = b(X(t))dt + dW(t), \\ X(0) = x \in H, \end{cases}$$
(1.3)

where W is an \mathbb{R}^d -valued standard Brownian motion and b fulfills the following assumptions.

Hypothesis 1.1. (i) There exist $\omega > 0$, $a \ge 0$ such that

$$\langle b(x), x \rangle \le -\omega |x|^2 + a, \quad \forall x \in \mathbb{R}^d,$$
(1.4)

(ii) $b: H \to H$ is continuously differentiable and there exists K > 0, $N \in \mathbb{N}$ such that

$$|b(x)| + ||b'(x)|| \le K(1+|x|^{2N}), \quad \forall x \in \mathbb{R}^d.$$
(1.5)

By (ii) it follows that b is Lipschitz continuous on bounded sets of H, whereas (i) allows to estimate $|X(t,x)|^2$ by Itô' formula; therefore existence and uniqueness of a strong solution $X(\cdot, x)$ of (1.3) is classical, see e.g. the monograph [Kr95]. We shall denote by P_t the transition semigroup

$$P_t\varphi(x) = \mathbb{E}[\varphi(X(t,x))], \quad t \ge 0, \ x \in H, \ \varphi \in B_b(H)$$
(1.6)

For proving (1.1) we argue as in [DaDe14] starting from the elementary identity, see (3.2)

$$P_t(\langle D\varphi, h\rangle) = \langle DP_t\varphi, h\rangle - \int_0^t P_{t-s}(\langle Db \cdot h, DP_s\varphi\rangle)ds.$$

Then we prove suitable estimates for $DP_t\varphi$ and their integrals with respect to ν . These estimates require some work because, due to the polynomial growth of the derivative of b, see (1.5), we cannot exploit the classical Bismut–Elworthy–Li formula, see [El92]. To overcome this problem we shall argue as in [DaDe03], [DaDe07] and [DaDe14], introducing a suitable potential (in the present case $V(x) = K(1 + |x|^{2N})$) and the Feynman–Kac semigroup

$$S_t \varphi(x) = \mathbb{E}[\varphi(X(t,x)) e^{-\int_0^t V(X(s,x)) \, ds}].$$
(1.7)

We shall first estimate $\langle DS_t\varphi(x),h\rangle$ then $\langle DP_t\varphi(x),h\rangle$, by taking advantage of the identity

$$P_t \varphi = S_t \varphi + \int_0^t S_{t-s}(V P_s \varphi) \, ds, \qquad (1.8)$$

which follows from the variation of constants formula, see Section 2 below.

In Section 3 we prove that inequality (1.1) and identity (1.2) hold with R = I. Moreover, for any $z \in H$ we show that the Fomin derivative v_z in the direction $z \in H$ is given by $\langle D \log \rho, z \rangle$, where ρ is the density of ν with respect to the Lebesgue measure. Moreover $v_z \in L^p(H, \nu)$ for all $p \in [1, \infty)$. Finally, we prove a formula for the adjoint D^* of D and also for the elliptic operator $-\frac{1}{2}D^*D$ which can be seen as a generalisation of the Ornstein–Uhlenbeck operator.

We end this section with some notations. We set $H = \mathbb{R}^d$, $d \ge 1$ (norm $|\cdot|$, inner product $\langle \cdot, \cdot \rangle$) and denote by L(H) the space of all linear bounded operators from H into H. Moreover, $C_b(H)$ is the space of all real continuous and bounded mappings $\varphi \colon H \to \mathbb{R}$ endowed with the sup norm

$$\|\varphi\|_{\infty} = \sup_{x \in H} |\varphi(x)|$$

whereas $C_b^k(H)$, k > 1, is the space of all real functions which are continuous and bounded together with their derivatives of order lesser than k. Finally, $B_b(H)$ will represent the space of all real, bounded and Borel mappings on H.

2 Estimates of the derivative of the transition semigroup

Let us start by giving an estimate of $\mathbb{E}(|X(t,x)|^{2m})$, $m \in \mathbb{N}$. The following lemma is standard, we shall give some details of the proof for the reader's convenience.

Lemma 2.1. Assume Hypothesis 1.1(i). Then for any $m \in \mathbb{N}$ there exists $a_m > 0$ such that

$$\mathbb{E}[|X(t,x)|^{2m}] \le e^{-2m\omega t} |x|^{2m} + a_m, \quad \forall \ x \in H, \ t \ge 0.$$
(2.1)

Proof. Let first consider the case m = 1. Then by Itô's formula, taking into account (1.4) we find

$$\frac{d}{dt} \mathbb{E}[|X(t,x)|^2] = 2\mathbb{E}[\langle X(t,x), b(X(t,x)) \rangle] + d$$
$$\leq -2\omega\mathbb{E}[|X(t,x)|^2] + 2a + d.$$

We deduce that

$$\frac{d}{dt} \mathbb{E}\left[|X(t,x)|^2\right] \le -2\omega \mathbb{E}\left[|X(t,x)|^2\right] + 2a + d.$$

By a standard comparison result it follows that

$$\mathbb{E}\left[|X(t,x)|^{2}\right] \le e^{-2\omega t}|x|^{2} + a_{2}, \quad \forall \ x \in H, \ t \ge 0,$$
(2.2)

where

$$a_2 = \frac{1}{\omega} \left(2a + d\right).$$

Now let m > 1 and $\varphi_m(x) = |x|^{2m}$. Then we have

$$D\varphi_m(x) = 2m|x|^{2m-2}x$$

and

$$D^{2}\varphi_{m}(x) = 4m(m-1)|x|^{2m-4} x \otimes x + 2m|x|^{2m-2} I,$$

where I represents identity in H. Consequently

$$\frac{1}{2} \operatorname{Tr} \left[D^2 \varphi_m(x) \right] = m(2m - 2 + d) |x|^{2m - 2}$$

Then again by Itô's formula we have

$$\frac{d}{dt} \mathbb{E} \left[|X(t,x)|^{2m} \right] = 2m \mathbb{E} \left[X(t,x)|^{2m-2} \right] \langle X(t,x), b(X(t,x)) \rangle \right] + m(2m-2+d) \mathbb{E} [X(t,x)|^{2m-2}] \leq -2m\omega \mathbb{E} \left[|X(t,x)|^{2m} \right] + m(2a+2m-2+d) \mathbb{E} \left[X(t,x)|^{2m-2} \right].$$

It follows that

$$\mathbb{E}[|X(t,x)|^{2m}] \leq e^{-2m\omega t}|x|^{2m}$$

$$+m(2a+2m-2+d)\int_0^t e^{-2m\omega(t-s)}\mathbb{E}[X(s,x)]^{2m-2}]ds.$$

The conclusion follows easily by recurrence.

Now we are going to prove an estimate for the derivative
$$D_x X(t,x)h$$
,
which we denote by $\eta^h(t,x)$, $h \in H$. As well known $\eta^h(t,x)$ is a solution to
the random equation

$$\begin{cases} \frac{d}{dt} \eta^h(t,x) = b'(X(t,x) \cdot \eta^h(t,x), \\ \eta^h(0,x) = h \end{cases}$$
(2.3)

Lemma 2.2. Assume Hypothesis 1.1. Then the following estimate holds

$$|\eta^{h}(t,x)| \le e^{K \int_{0}^{t} (1+|X(s,x)|^{2N}) ds} |h|, \quad t \ge 0, \ x,h \in H.$$
(2.4)

Proof. By
$$(2.3)$$
 we deduce, taking into account (1.5) , that

$$\frac{1}{2}\frac{d}{dt}|\eta^{h}(t,x)|^{2} = \langle b'(X(t,x)\cdot\eta^{h}(t,x),\eta^{h}(t,x))\rangle \leq K(1+|X(t,x)|^{2N})|\eta^{h}(t,x)|^{2}.$$

So, the conclusion follows from Gronwall's lemma.

Now we are going to estimate of $D_x P_t \varphi$.

2.1 Pointwise estimate

As we said in the introduction, we cannot estimate $D_x P_t \varphi$ for $\varphi \in C_b(H)$ using the Bismut–Elworthy–Li formula see [El92], because we do not know whether the expectation on the right hand side of (2.4) does exist. For this reason, we introduce the potential

$$V(x) = K(1 + |x|^{2N}), \quad x \in H$$

and the Feynman–Kac semigroup

$$S_t\varphi(x) = \mathbb{E}[\varphi(X(t,x)) e^{-\int_0^t V(X(s,x)) ds}].$$

We recall that the Bismut–Elworthy–Li formula generalises to S_t , see [DaZa97]. In fact for all $\varphi \in C_b(H)$, setting

$$\beta(t) = \int_0^t V(X(s,x))ds,$$

the following identity holds

$$\langle DS_t \varphi(x), h \rangle = \frac{1}{t} \mathbb{E} \left[\varphi(X(t,x)) e^{-\beta(t)} \int_0^t \langle \eta^h(s,x), dW(s) \rangle \right]$$
$$-\mathbb{E} \left[\varphi(X(t,x)) e^{-\beta(t)} \int_0^t \left(1 - \frac{s}{t}\right) \left\langle V'(X(s,x), \eta^h(s,x)) \right\rangle \right] ds \tag{2.5}$$

$$=: I_1(\varphi, x, h, t) + I_2(\varphi, x, h, t) = I_1 + I_2.$$

We shall first estimate $\langle DS_t\varphi(x),h\rangle$, then $\langle DP_t\varphi(x),h\rangle$. In the latter case, we take advantage of the identity

$$P_t\varphi = S_t\varphi + \int_0^t S_{t-s}(VP_s\varphi)\,ds,$$

which follows from the variation of constants formula; in fact, denoting by \mathscr{L} and \mathscr{K} the infinitesimal generators of P_t and S_t respectively, it holds

$$\mathscr{L} = \mathscr{K} + V.$$

Lemma 2.3. Let $\varphi \in C_b(H)$, $t \ge 0$, $x \in H$. Then for p > 1, there exists a constant $C_p > 0$ such that

$$|D_x S_t \varphi(x)| \le C_p (1 + t^{-1/2}) (1 + |x|^{2N-1}) \left[\mathbb{E} \left(\varphi^p (X(t, x)) \right)^{1/p} \right].$$
(2.6)

Proof. We start by estimating I_1 . By Hölder's inequality with exponents $p, q = \frac{p}{p-1}$ we have

$$|I_1| \leq \frac{1}{t} \left[\mathbb{E} \left(\varphi^p(X(t,x)) \right]^{1/p} \left[\mathbb{E} \left(e^{-q\beta(t)} \left| \int_0^t \langle \eta^h(s,x), dW(s) \rangle \right|^q \right) \right]^{1/q}$$

=: $\frac{1}{t} \left[\mathbb{E} \left(\varphi^p(X(t,x)) \right]^{1/p} \left[\mathbb{E} \left(|z(t)|^q \right) \right]^{1/q} ,$ (2.7)

where

$$z(t) = e^{-\beta(t)} \int_0^t \langle \eta^h(s, x), dW(s) \rangle, \quad t \ge 0.$$
(2.8)

We now apply Itô's formula to g(z(t)) where $g(r) = |r|^q, r \in \mathbb{R}$. Since

$$g'(r) = q|r|^{q-2}r, \quad g''(r) = q(q-1)|r|^{q-2},$$

and

$$dz(t) = -\beta'(t)e^{-\beta(t)} \int_0^t \langle \eta^h(s,x), dW(s) \rangle \, ds + e^{-\beta(t)} \langle \eta^h(t,x), dW(t) \rangle$$
$$= -\beta'(t)z(t) + e^{-\beta(t)} \langle \eta^h(t,x), dW(t) \rangle,$$

we find

$$\begin{aligned} d|z(t)|^{q} &= q|z(t)|^{q-2}z(t)(-\beta'(t)z(t) + e^{-\beta(t)}\langle \eta^{h}(t,x), dW(t)\rangle) \\ &+ \frac{1}{2}q(q-1)|z(t)|^{q-2}e^{-2\beta(t)}|\eta^{h}(t,x)|^{2}dt. \end{aligned}$$

Integrating from 0 to t, yields

$$|z(t)|^{q} = -q \int_{0}^{t} |z(s)|^{q} \beta'(s) ds$$

+ $q \int_{0}^{t} |z(s)|^{q-2} z(s) e^{-\beta(s)} \langle \eta^{h}(s, x), dW(s) \rangle$ (2.9)
+ $\frac{1}{2} q(q-1) \int_{0}^{t} e^{-2\beta(s)} |z(s)|^{q-2} |\eta^{h}(s, x)|^{2} ds.$

Neglecting the negative first term in the previous identity and taking expectation, we find

$$\mathbb{E}\left(\sup_{r\in[0,t]}|z(r)|^{q}\right)$$

$$\leq q \mathbb{E}\left(\sup_{r\in[0,t]}\left|\int_{0}^{r}e^{-\beta(s)}|z(s)|^{q-2}z(s)\langle\eta^{h}(s,x),dW(s)\rangle\right|\right)$$

$$+\frac{1}{2}q(q-1)\mathbb{E}\left(\int_{0}^{t}e^{-2\beta(s)}|z(s)|^{q-2}|\eta^{h}(s,x)|^{2}ds\right)$$

$$=: A_{1} + A_{2}.$$
(2.10)

By the Burkholder inequality we have, taking into account Lemma 2.1

$$A_{1} \leq 3q\mathbb{E}\left[\left|\int_{0}^{t} e^{-2\beta(s)}|z(s)|^{2(q-1)}|\eta^{h}(s,x)|^{2}ds\right|^{1/2}\right]$$

$$\leq 3q\mathbb{E}\left[\sup_{r\in[0,t]}|z(r)|^{q-1}\left(\int_{0}^{t} e^{-2\beta(s)}|\eta^{h}(s,x)|^{2}ds\right)^{1/2}\right]$$

$$\leq 3qt^{1/2}\mathbb{E}\left[\sup_{r\in[0,t]}|z(r)|^{q-1}\right]|h|.$$
(2.11)

By Hölder's inequality with exponents $q,\,\frac{q}{q-1},$ it follows that

$$A_1 \le 3q t^{1/2} |h| \left[\mathbb{E} \left(\sup_{r \in [0,t]} |z(r)|^q \right) \right]^{\frac{q-1}{q}}.$$
 (2.12)

Now by the Young inequality

$$ab \le \frac{1}{u}a^u + \frac{1}{v}a^v, \quad a > 0, \ b > 0, \ \frac{1}{u} + \frac{1}{v} = 1$$
 (2.13)

with u = q, $v = \frac{q-1}{q}$, there exists $c_1 > 0$ such that

$$A_1 \le \frac{1}{4} \mathbb{E}\left(\sup_{r \in [0,t]} |z(r)|^q\right) + c_1 t^{q/2} |h|^q.$$
(2.14)

Concerning A_2 , using again Lemma 2.1, we find

$$A_{2} = \frac{1}{2}q(q-1)\mathbb{E}\left(\int_{0}^{t} e^{-2\beta(s)}|z(s)|^{q-2}|\eta^{h}(s,x)|^{2}ds\right)$$

$$\leq \frac{1}{2}q(q-1)\mathbb{E}\left[\left(\sup_{r\in[0,t]}|z(r)|^{q-2}\right)\int_{0}^{t} e^{-2\beta(s)}|\eta^{h}(s,x)|^{2}ds\right]$$

$$\leq \frac{1}{2}q(q-1)\mathbb{E}\left[\left(\sup_{r\in[0,t]}|z(r)|^{q-2}\right)\right]|h|^{2}t.$$

By Hölder's inequality with exponents $\frac{q}{2},\,\frac{q}{q-2}$ we have

$$A_2 \le \frac{1}{2} q(q-1) |h|^2 \left[\mathbb{E} \left(\sup_{r \in [0,t]} |z(r)|^q \right) \right]^{\frac{q-2}{q}}.$$

By the Young inequality (2.13) with $u = \frac{q}{2}$ and $v = \frac{q}{q-2}$, it follows that there exists $c_2 > 0$ such that

$$A_2 \le \frac{1}{4} \mathbb{E}\left(\sup_{r \in [0,t]} |z(r)|^q\right) + c_2 |h|^q t^{q/2}.$$
(2.15)

Taking into account (2.10), (2.14) and (2.15) we conclude that

$$\mathbb{E}\left(\sup_{r\in[0,t]}|z(r)|^{q}\right) \leq \frac{1}{2} \mathbb{E}\left(\sup_{r\in[0,t]}|z(r)|^{q}\right) + (c_{1}+c_{2})|h|^{q} t^{q/2}.$$

Therefore

$$\mathbb{E}\left(\sup_{r\in[0,t]}|z(r)|^{q}\right) \le (c_{1}+c_{2}).\,|h|^{q}\,t^{q/2}.$$
(2.16)

Finally, by (2.7) it follows that

$$I_1 \le (c_1 + c_2)t^{-1/2} |h| \left[\mathbb{E} \left(\varphi^p(X(t, x)) \right)^{1/p} \right].$$
(2.17)

Now let us consider I_2 , and write

$$I_2 \le 2KN \left[\mathbb{E}\left[\varphi^p(X(t,x))\right]^{1/p} (\Lambda(t))^{1/q},$$
(2.18)

where $\frac{1}{p} + \frac{1}{q} = 1$ and

$$\Lambda(t) = \mathbb{E}\left[e^{-q\beta(t)}\left(\int_0^t |X(s,x)|^{2N-1} |\eta^h(s,x)| \, ds\right)^q\right]$$

$$\leq \mathbb{E}\left[\left(\int_0^t e^{-\beta(s)} |X(s,x)|^{2N-1} |\eta^h(s,x)| \, ds\right)^q\right]$$

$$\leq \mathbb{E}\left[\sup_{r \in [0,t]} \left(|X(r,x)|^{(2N-1)q}\right) \left(\int_0^t e^{-\beta(s)} |\eta^h(s,x)| \, ds\right)^q\right]$$

$$\leq \mathbb{E}\left[\sup_{r \in [0,t]} \left(|X(r,x)|^{(2N-1)q}\right)\right] |h|^q t^q.$$
(2.19)

 So

$$I_{2} \leq 2KN \left[\mathbb{E}\left[\varphi^{p}(X(t,x))\right]^{1/p} \left(\mathbb{E}\left[\sup_{r \in [0,t]} \left(|X(r,x)|^{(2N-1)q}\right)\right]\right)^{1/q} |h| t.$$
(2.20)

Recalling finally (2.1) we see that there exists $c_3 > 0$ such that

$$\left(\mathbb{E}\left[\sup_{r\in[0,t]}\left(|X(r,x)|^{2N-1}\right)\right]\right)^{q} \le c_{3}(1+|x|^{2N-1}),$$

so that

$$I_2 \le 2KNc_3(1+|x|^{2N-1}) \left[\mathbb{E}\left[\varphi^p(X(t,x))\right]^{1/p} |h| t.$$
(2.21)

Finally, by (2.5), (2.17) and (2.21), the conclusion follows easily.

2.2 The invariant measure ν

We shall denote by $\pi_{t,x}$ the law of X(t,x) so that for each $\varphi \in B_b(H)$ we have

$$P_t\varphi(x) = \int_H \varphi(y)\pi_{t,x}(dy), \quad x \in H, \ t > 0.$$
(2.22)

Lemma 2.4. Assume Hypothesis 1.1(i). Then there is an invariant measure ν of P_t , moreover for all $m \in \mathbb{N}$ we have

$$\int_{H} |x|^{2m} \nu(dx) \le a_m, \tag{2.23}$$

where a_m is the constant in (2.1).

Proof. Let r > 0 and fix $x \in H$. Set $B_r^c = \{y \in H : |y| \ge r\}$. Then, taking into account (2.2) it follows that

$$\pi_{t,x}(B_r^c) = \int_{\{|y| \ge r\}} \pi_{t,x}(dy) \le \frac{1}{r^2} \int_H |y|^2 \pi_{t,x}(dy)$$

$$= \frac{1}{r^2} \mathbb{E}\left[|X(t,x)|^2\right] \le \frac{|x|^2 + a_2}{r^2}.$$
(2.24)

Therefore by the Krylov–Bogoliubov theorem, see e.g [DaZa96], there exists a sequence $T_n \uparrow +\infty$ such that

$$\lim_{n \to +\infty} \frac{1}{T_n} \int_0^{T_n} \pi_{t,x} dt = \nu \quad \text{weakly},$$
(2.25)

where ν is an invariant measure of P_t .

Now we can prove (2.23). By (2.1) we deduce

$$\int_{H} |y|^{2m} \pi_{t,x}(dy) \le e^{-m\omega t} |x|^{2m} + a_m, \quad \forall \ x \in H, \ t \ge 0.$$
(2.26)

It follows that for any $\epsilon > 0$

$$\int_{H} \frac{|y|^{2m}}{1+\epsilon|y|^{2m}} \pi_{t,x}(dy) \le e^{-m\omega t} |x|^{2m} + a_m, \quad \forall x \in H, \ t \ge 0.$$
(2.27)

Consequently integrating both sides with respect to t over $[0, T_n]$ and dividing by T_n , yields

$$\frac{1}{T_n} \int_0^{T^n} dt \int_H \frac{|y|^{2m}}{1+\epsilon |y|^{2m}} \pi_{t,x}(dy) \le \frac{1}{m\omega T_n} \left(1 - e^{-m\omega T_n}\right) |x|^{2m} + a_m, \quad (2.28)$$

for all $x \in H$, $t \ge 0$. Finally, letting $n \to +\infty$ and taking into account (2.25), we find

$$\int_{H} \frac{|y|^{2m}}{1+\epsilon |y|^{2m}} \nu(dy) \le a_m$$

and the conclusion follows letting ϵ tend to 0.

2.3 Integral estimates

Let us start with an estimate of $\int_H \langle D_x S_t \varphi(x), h(x) \rangle \nu(dx)$.

Lemma 2.5. Let p > 1, q > 1, $\frac{1}{p} + \frac{1}{q} < 1$. Then there is $C_1 > 0$ such that

$$\left| \int_{H} \langle D_x S_t \varphi(x), h(x) \rangle \, \nu(dx) \rangle \right| \le C_1 (1 + t^{-1/2}) \, \|\varphi\|_{L^p(H,\nu)} \, \|h\|_{L^q(H,\nu)}.$$
(2.29)

Proof. Taking into account (2.6) we have

$$\left| \int_{H} \langle D_{x} S_{t} \varphi(x), h(x) \rangle \, \nu(dx) \right| \leq C_{p} \left(1 + t^{-1/2} \right)$$

$$\times \int_{H} (1 + |x|^{2N-1}) \left[\left(|P_{t} \varphi^{p}(x)| \right) \right]^{1/p} |h(x)| \, \nu(dx).$$
(2.30)

Let

$$\frac{1}{r} = 1 - \frac{1}{p} - \frac{1}{q},$$

then by the triple Hölder inequality with exponents r, p, q we have, taking into account the invariance of ν ,

$$\begin{split} \left| \int_{H} \langle D_{x} S_{t} \varphi(x), h(x) \rangle \, \nu(dx) \right| &\leq c \left(1 + t^{-1/2} \right) \\ &\times \left[\int_{H} (1 + |x|^{N-1})^{r} \nu(dx) \right]^{1/r} \left(\int_{H} |P_{t} \varphi^{p}(x)| \nu(dx) \right)^{1/p} \|h\|_{L^{q}(H,\nu)} \quad (2.31) \\ &\leq c \left(1 + t^{-1/2} \right) \left[\int_{H} (1 + |x|^{N-1})^{r} \nu(dx) \right]^{1/r} \|\varphi\|_{L^{p}(H,\nu)} \|h\|_{L^{q}(H,\nu)}. \end{split}$$
The conclusion follows from (2.23).

The conclusion follows from (2.23).

Now we are ready to estimate $\int_H \langle D_x P_t \varphi(x), h(x) \rangle \, \nu(dx)$. We start from the identity

$$P_t\varphi(x) = S_t\varphi(x) + K \int_0^t S_{t-s}(1+|x|^{2N}) P_s\varphi(x) ds, \qquad (2.32)$$

from which

$$\langle D_x P_t \varphi(x), h(x) \rangle$$

$$= \langle D_x S_t \varphi(x), h(x) \rangle + K \int_0^t \langle D_x S_{t-s}((1+|x|^{2N}) P_s \varphi)(x), h(x) \rangle ds.$$

$$(2.33)$$

Proposition 2.6. Let p > 1, q > 1, $\frac{1}{p} + \frac{1}{q} < 1$. Then there is C_p^1 such that

$$\left| \int_{H} \langle D_{x} P_{t} \varphi(x), h(x) \rangle \, \nu(dx) \rangle \right| \leq C_{p}^{1} (1 + t^{-1/2}) \, \|\varphi\|_{L^{p}(H,\nu)} \, \|h\|_{L^{q}(H,\nu)}.$$
(2.34)

Proof. The first term of (2.33) is bounded by (2.29). Let us estimate the second one. Again by (2.29) we have

$$\left| \int_{0}^{t} \int_{H} \langle D_{x} S_{t-s}((1+|x|^{2N}) P_{s}\varphi), h(x) \rangle ds \right| \nu(dx)$$

$$\leq C_{1} \int_{0}^{t} (1+(t-s)^{-\frac{1}{2}}) \| (1+|x|^{2N}) P_{s}\varphi \|_{L^{p}(H,\nu)} \| h \|_{L^{q}(H,\nu)} ds$$

$$(2.35)$$

Now let us chose $\epsilon > 0$ such that

$$\frac{1}{p+\epsilon}+\frac{1}{q}<1.$$

Then by Hölder's inequality with exponents $\frac{p+\epsilon}{\epsilon}$ and $\frac{p+\epsilon}{p}$ it follows that

$$\begin{split} \|(1+|x|^{2N}) P_s \varphi)\|_{L^p(H,\nu)}^p &= \int_H (1+|x|^{2N})^p (P_s \varphi)^p \, d\nu \\ &\leq \left(\int_H (1+|x|^{2N})^{\frac{p(p+\epsilon)}{\epsilon}} \, d\nu\right)^{\frac{\epsilon}{p+\epsilon}} \left(\int_H (P_s \varphi)^{p+\epsilon} \, d\nu\right)^{\frac{p}{p+\epsilon}} \\ &\leq \left(\int_H (1+|x|^{2N})^{\frac{p(p+\epsilon)}{\epsilon}} \, d\nu\right)^{\frac{\epsilon}{p+\epsilon}} \|\varphi\|_{L^{p+\epsilon}(H,\nu)}^p, \end{split}$$

by the invariance of ν . Now by (2.2) there exists a constant C' such that

$$\int_{H} (1+|x|^{2N})^{\frac{p(p+\epsilon)}{\epsilon}} d\nu \le C',$$

Therefore

$$\|(1+|x|^{2N})P_{s}\varphi)\|_{L^{p}(H,\nu)}^{p} \leq (C')^{\frac{\epsilon}{p+\epsilon}} \|\varphi\|_{L^{p+\epsilon}(H,\nu)}^{p}.$$
(2.36)

Substituting in (2.35), yields

$$\left| \int_{0}^{t} \langle D_{x} S_{t-s}(|x|_{N}^{N} P_{s} \varphi), h(x) \rangle ds \right|$$

$$\leq C_{1}(C')^{\frac{\epsilon}{p+\epsilon}} \int_{0}^{t} (1 + (t-s)^{-\frac{1}{2}}) \|\varphi\|_{L^{p+\epsilon}(H,\nu)} \|h|_{L^{q}(H,\nu)} ds$$
(2.37)

Non the conclusion follows by the arbitrariness of ϵ, p, q .

3 The main inequality and its consequences

Theorem 3.1. For all p > 1 there exists a constant $C_p > 0$ such that for all $\varphi \in L^p(H, \nu)$ and all $h \in H$ we have

$$\left| \int_{H} \langle D_x \varphi(x), h \rangle \rangle \, \nu(dx) \right| \le c \|\varphi\|_{L^p(H,\nu)} \, |h|. \tag{3.1}$$

Proof. Step 1. For any $\varphi \in C_b^1(H)$ and any $h \in H$ the following identity holds.

$$P_t(\langle D\varphi, h\rangle) = \langle DP_t\varphi, h\rangle - \int_0^t P_{t-s}(\langle Db \cdot Hh, DP_s\varphi\rangle)ds, \quad t > 0.$$
(3.2)

To prove (3.2) we consider a sequence (b_n) of mappings $H \to H$ of class C^∞ such that

- (i) $\lim_{n\to\infty} b_n(x) = b(x)$, uniformly on bounded sets of H.
- (ii) $\langle b_n(x), x \rangle \le -\omega |x|^2 + a, \quad \forall x \in H.$

To construct (b_n) we first set

$$f_n(x) = \frac{b(x) + \omega x}{1 + n^{-1} |x|^{2N+2}} - \omega x,$$

so that

$$\langle f_n(x), x \rangle \le -\omega |x|^2 + a, \quad \forall \ x \in H,$$

and f_n is sub–linear, then we regularise f_n using mollifiers.

Now we prove the identity

$$P_t^n(\langle D\varphi,h\rangle) = \langle DP_t^n\varphi,h\rangle - \int_0^t P_{t-s}(\langle Db\cdot h, DP_s^n\varphi\rangle)ds, \qquad (3.3)$$

where P_t^n is the transition semigroup corresponding to b_n . It is enough to show (3.3) for each $\varphi \in C_b^3(H)$. In such a case set $u_n(t,x) = P_t^n \varphi(x)$ and write

$$\begin{cases} D_t u_n(t,x) = \frac{1}{2} \Delta u_n(t,x) + \langle D u_n(t,x), b_n(x) \rangle, \\ u_n(0,x) = \varphi(x). \end{cases}$$
(3.4)

Now, taking $h \in H$ and setting

$$v_n(t,x) = \langle Du_n(t,x), h \rangle$$

we see, by a simple computation, that

$$\begin{cases} D_t v_n(t,x) = \frac{1}{2} \Delta v_n(t,x) + \langle D v_n(t,x), b_n(x) \rangle \\ + \langle D u_n(t,x), b'_n(x)h \rangle, \\ v_b(0,x) = \langle D \varphi(x), h \rangle. \end{cases}$$
(3.5)

By the variation of constants formula it follows that

$$v_n(t,x) = P_t^n(\langle D\varphi(x),h\rangle) + \int_0^t P_{t-s}^n \langle Du_n(s,x),Ah + b'_n(x)h\rangle ds, \quad (3.6)$$

which coincides with (3.3). Letting $n \to \infty$, yields (3.2).

Step 2. Conclusion.

Integrating (3.2) with respect to ν over H and taking into account the invariance of ν , yields

$$\int_{H} \langle D\varphi(x), h \rangle |\nu(dx) = \int_{H} \langle DP_{t}\varphi(x), h \rangle \nu(dx)$$

$$- \int_{H} \int_{0}^{t} \langle b'(x)h, DP_{s}\varphi(x) \rangle \, ds \, \nu(dx) =: J_{1} + J_{2}$$
(3.7)

Setting and t = 1 we deduce

$$|J_1| \le \left| \int_H \langle D_x P_t \varphi(x), h \rangle \,\nu(dx) \right| \le 2C_p^1 \, \|\varphi\|_{L^p(H,\nu)} \, |h|. \tag{3.8}$$

Concerning J_2 we have by (2.34) and taking into account (1.5)

$$|J_{2}| \leq \int_{0}^{t} \int_{H} C_{p}^{1} (1 + (t - s)^{-1/2}) \|\varphi\|_{L^{p}(H,\nu)} \|b'(\cdot)h\|_{L^{q}(H,\nu)} ds$$

$$\leq K \int_{0}^{t} \int_{H} C_{p}^{1} (1 + (t - s)^{-1/2}) \|\varphi\|_{L^{p}(H,\nu)} \|(1 + |x|^{2N})\|_{L^{q}(H,\nu)} ds |h|.$$
(3.9)

Finally, recalling (2.23) and setting t = 1 the conclusion follows.

3.1 Consequences of the integral inequality (3.1)

The following result can be proved exactly as in [DaDe14], replacing R by I so, we omit the proof.

Proposition 3.2. Assume Hypothesis 1.1 and let ν be the invariant measure of problem (1.3). Then for any p > 1 the gradient

$$D: C^1_b(H) \subset L^p(H,\nu) \to L^p(H,\nu;H), \quad \varphi \to D\varphi,$$

is closable.

For any p > 1 we shall denote by D_p the closure of D and by D_p^* the adjoint operator of D_p . D_p is a mapping

$$D_p: D(D_p) \subset L^p(H,\nu) \to L^p(H,\nu;H)$$

and D_p^* is a mapping

$$D_p^*: D(D_p^*) \subset L^q(H,\nu;H) \to L^q(H,\nu),$$

where $q = \frac{1}{1-p}$. We have obviously

$$\int_{H} \langle D_{p}\varphi, F \rangle \, d\nu = \int_{H} \varphi \, D_{p}^{*}(F) \, d\nu, \qquad (3.10)$$

for any $\varphi \in D(D_p)$ and any $F \in D(D_p^*)$. We recall that $F \in D(D_p^*)$ if and only if there exists a positive constant K_F such that

$$\left| \int_{H} \langle D\varphi, F \rangle \, d\nu \right| \le K_F \|\varphi\|_{L^p(H,\nu)}, \quad \forall \varphi \in C_b^1(H).$$
(3.11)

In this case we have

$$\|D_p^*(F)\|_{L^q(H,\nu)} \le K_F.$$
(3.12)

If no confusion may arise we shall omit sub-indices p in D_p and D_p^* .

Proposition 3.3. For any $z \in H$ there is $v_z \in L^q(H, \nu)$ for all $q \in [1, +\infty)$ such that

$$\int_{H} \langle D\varphi, z \rangle \, d\nu = \int_{H} v_{z} \, \varphi \, d\nu. \tag{3.13}$$

Proof. Let $z \in H$ and set $F_z(x) = z$, $\forall x \in H$. Then by (3.1) it follows that

$$\left| \int_{H} \langle D\varphi, F_{z} \rangle \, d\nu \right| \le C_{1,p} \, \|\varphi\|_{L^{p}(H,\nu)} \, |z| \tag{3.14}$$

This implies $F_z \in D(D_p^*)$ and $||D_p^*(F_z)||_{L^q(H,\nu)} \leq C_{1,p}|z|$. Setting $D_q^*(F_z) = v_z$, identity (3.13) follows.

Remark 3.4. By Proposition 3.3 ν possesses the Fomin derivative of ν at the direction z which is given precisely by v_z and so, it belongs to $L^q(H,\nu)$ for all $q \in [1,\infty)$,

Now we are going to identify v_z .

Proposition 3.5. Assume Hypothesis 1.1. Then for any $z \in H$ we have $v_z = \langle D \log \rho, z \rangle$, where ρ is the density of ν with respect to the Lebesgue measure on \mathbb{R}^d . Therefore $\langle D \log \rho, z \rangle$ belong to $L^p(H, \nu)$ for any $p \in [1, +\infty)$.

Proof. First notice that by (3.13) it follows in particular that

$$\left| \int_{H} \langle D\varphi, z \rangle \, d\nu \right| \le \|v_z\|_{L^1(H,\nu)} \, \|\varphi\|_{\infty}. \tag{3.15}$$

Therefore, by an argument due to Malliavin, ν has a density ρ with respect to the Lebesgue measure on \mathbb{R}^d with $\rho \in L^{\frac{d}{d-1}}(\mathbb{R}^d)$, see [Nu95].

To prove the last statement, we write (3.13) as

$$\int_{H} \langle D\varphi, z \rangle \, \rho \, dx = \int_{H} \varphi \, v_z \, \rho \, dx.$$

This implies in the sense of distributions that

$$v_z = \langle D \log \rho, z \rangle.$$

Now the conclusion follows from Proposition 3.3.

Remark 3.6. The fact that ν has a density ρ with respect to the Lebesgue measure, together with several properties of ρ have already been proved in [MePaRh05], [BoKrRo01] and [BoKrRo05].

Let us finally study some properties of operators D^* and D^*D .

Proposition 3.7. Let

$$F(x) = \sum_{h=1}^{d} f_h(x)e_h, \quad x \in H,$$
(3.16)

where $(e_1, ..., e_d)$ is an orthonormal basis in H and $f_h \in C_b^1(H)$, h = 1, ..., d. Then F belongs to the domain of D^* and it results

$$D^*(F) = -\text{div } F + \sum_{h=1}^d v_{e_h} f_h.$$
 (3.17)

Moreover, if $\varphi \in C_b^2(H)$ we have

$$-\frac{1}{2}D^*D(\varphi) = \frac{1}{2}\Delta\varphi - \frac{1}{2}\sum_{h=1}^d v_{e_h} D_h\varphi.$$
 (3.18)

Proof. Write

$$\int_{H} \langle D\varphi, F \rangle \, d\nu = \sum_{h=1}^{d} \int_{H} D_{h} \varphi \, f_{h} \, d\nu$$

$$= \sum_{h=1}^{d} \int_{H} D_{h} (\varphi \, f_{h}) \, d\nu - \sum_{h=1}^{d} \int_{H} \varphi \, D_{h} f_{h} \, d\nu$$
(3.19)

Since, in view of (3.13)

$$\int_{H} D_{h}(\varphi f_{h}) \, d\nu = \int_{H} \varphi \, v_{e_{h}} f_{h} \, d\nu,$$

(3.17) follows. Now (3.18) follows as well setting $F = D\varphi$ in (3.17).

Acknowledgement

G. Da Prato is partially supported by GNAMPA from INDAM.

References

- [BoKrRo01] V. I. Bogachev, N. V. Krylov and M. Röckner, On regularity of transition probabilities and invariant measures of singular diffusions under minimal conditions, Comm. Partial Differential Equations 26, no. 11-12, 2037–2080, 2001.
- [BoKrRo05] V. I. Bogachev, N. V. Krylov and M. Röckner, Regularity and global bounds for the densities of invariant measures of diffusion processes. Dokl. Akad. Nauk 405, 2005.
- [DaDe03] G. Da Prato and A. Debussche, Ergodicity for the 3D stochastic Navier-Stokes equations, J. Math. Pures Appl., 82, 877–947, 2003.
- [DaDe07] G. Da Prato and A. Debussche, m-dissipativity of Kolmogorov operators corresponding to Burgers equations with space-time white noise, Potential Anal, 26, 31-55, 2007.
- [DaDe14] G. Da Prato and Debussche, Estimate for P_tD for the stochastic Burgers equation, arXiv:1412.7426, 2014, to appear in Ann. Inst. Poincaré.
- [DaZa96] G. Da Prato and J. Zabczyk, Ergodicity for Infinite Dimensional Systems, London Mathematical Society Lecture Notes, no. 229, Cambridge University.
- [DaZa97] G. Da Prato and J. Zabczyk, Differentiability of the Feynman-Kac semigroup and a control application, Rend. Mat. Acc. Lincei. s.9, v. 8, 183–188, 1997.
- [El92] K. D. Elworthy, Stochastic flows on Riemannian manifolds, in "Diffusion processes and related problems in analysis, Vol. II, M. A. Pinsky and V. Wihstutz (eds.), Birkhäuser, 33–72, 1992.
- [Kr95] N. V. Krylov, Introduction to the theory of diffusion processes. Translations of Mathematical Monographs, 142, 1995.
- [Ma97] P. Malliavin (1997) Stochastic analysis, Springer-Verlag, 1997.
- [MePaRh05] G. Metafune, D. Pallara and A. Rhandi, Global properties of invariant measures, J. Funct. Analysis, 223, 396–424, 2005.
- [Nu95] D. Nualart, *The Malliavin calculus and related topics*. Probability and its Applications, Springer-Verlag, Berlin (1995).