The Ponzano-Regge asymptotic of the supersymmetric 6jS symbols

Lionel Bréhamet

July 7, 2019

Abstract

We adapt the Gurau's proof (2008) about the asymptotic limit of Ponzano-Regge formula to supersymmetric 6jS symbols according to their intrinsic parities alpha, beta, gamma. The behaviour at a large scaling shows significant differences depending on these parities. The decay is slowed and the angles of the oscillating parts in cosine are generally shifted or more altered. Our results should be relevant in 3-D Quantum Supergravity and Spin Foam models.

PACS: 03.65.Fd Algebraic methods PACS: 04.60.-m Quantum gravity PACS: 11.30.Pb Supersymmetry MSC: 81Q60; 83D05. Keywords. 6j symbol; Quantum Gravity; supersymmetric 6jS symbol.

1 Introduction

Since the growing expansion of quantum gravity theory [1] the SU(2) 6j symbols acquired a considerable importance by becoming the basic building blocks of all spin networks. They appear to represent a quantum tetrahedron with quantized edges and even can be viewed as eigenfunctions of a discrete Schrödinger equation [2]. As well known the classical Ponzano-Regge partition function Z_{PR} [3] was expressed as a sum of products of 6j symbols. In [4] the occurrence of supersymmetric $6j^S$ symbols [8, 9], related to OSP(1|2), was also studied and led to a similar supersymmetric partition function, called Z_{sugra} . A possible different divergence compared to the classical case was sketched.

Guided by the conceptual approach of Gurau [10], our present task is to give also an elementary proof of the Ponzano-Regge asymptotic of the $6j^S$ supersymmetric symbols.

Let us recall the formulation of Gurau:

Under a rescaling of all its arguments by a large k the 6j symbol associated to an Euclidean tetrahedron behaves like

$$\begin{cases} kj_1 \ kj_2 \ kj_3\\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} = \frac{1}{\sqrt{12\pi k^3 V}} \cos\left\{\frac{\pi}{4} + \sum_{\iota=1}^3 \left[\left(kj_\iota + \frac{1}{2}\right)\theta_{j_\iota} + \left(kJ_\iota + \frac{1}{2}\right)\theta_{J_\iota}\right]\right\},\tag{1}$$

where V is the tetrahedron volume and $\theta_{j_{\iota}}$, $\theta_{J_{\iota}}$ are the exterior dihedral angles of the tetrahedron corresponding to the edges j_{ι} and J_{ι} respectively.



Tetrahedron of volume V

2 Recalls about 6j and $6j^S$ symbols and notations

We rewrite our formulas [9] with notations as close as possible to those used by Gurau [10] and replace $j_i \leftrightarrow J_i, p_1 \rightarrow v_4, p_2 \rightarrow v_2, p_3 \rightarrow v_3, p_4 \rightarrow v_1, q \rightarrow p$. Notations used in [10] were

$$v_1 = j_1 + j_2 + j_3, v_2 = J_1 + j_2 + J_3, v_3 = J_1 + J_2 + j_3, v_4 = j_1 + J_2 + J_3,$$

$$(2)$$

$$p_1 = j_2 + J_2 + j_3 + J_3, \ p_2 = j_3 + J_3 + j_1 + J_1, \ p_3 = j_1 + J_1 + j_2 + J_2.$$
(3)

Diagrammatic representation of a 6j symbol $\begin{cases} j_1 & j_2 & j_3 \\ J_1 & J_2 & J_3 \end{cases}$:

the four triangles of any $6j$			the three columns pairs			
· · ·	· · · ·	••••	0 <u>0</u> 0	\cdot	$\circ \cdot \circ$	
v_4	v_2	v_3	v_1	p_1	p_2	p_3

 v_i is the sum of the values of the three circled spins just above v_i in the diagrams. In the same way, p_j is the sum of the values of the four circled spins above p_j . The variables p,v satisfy the equation

$$\sum_{j=1}^{3} p_j = \sum_{i=1}^{4} v_i.$$
(4)

As a result any spin is determined by two v_i and one p_j according to

$$2j_1 = v_1 + v_4 - p_1, \ 2j_2 = v_1 + v_2 - p_2, \ 2j_3 = v_1 + v_3 - p_3, \tag{5}$$

$$2J_1 = v_2 + v_3 - p_1, \, 2J_2 = v_3 + v_4 - p_2, \, 2J_3 = v_2 + v_4 - p_3, \tag{6}$$

or by two p_i and two v_i 'complementary' according to

$$2j_1 = p_2 + p_3 - v_2 - v_3, \ 2j_2 = p_1 + p_3 - v_3 - v_4, \ 2j_3 = p_1 + p_2 - v_2 - v_4, \tag{7}$$

$$2J_1 = p_2 + p_3 - v_1 - v_4, \ 2J_2 = p_1 + p_3 - v_1 - v_2, \ 2J_3 = p_1 + p_2 - v_1 - v_3.$$
(8)

Formulas for a 6j symbol are well known, however here we shall use an expression [11], used elsewhere long ago. That avoids the usual triangle coefficients \triangle and contains only terms in v_i, p_j .

$$\begin{cases} j_1 \ j_2 \ j_3 \\ J_1 \ J_2 \ J_3 \end{cases} = \sqrt{\mathsf{R}} \sum_t \frac{(-1)^t (t+1)!}{\prod_{i=1}^{i=4} (t-v_i)! \prod_{j=1}^{j=3} (p_j-t)!} ,$$
(9)

where the radical R under square root is the prefactor used by Gurau

$$\mathbf{R} = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} (p_j - v_i)!}{\prod_{i=1}^{i=4} (v_i + 1)!}.$$
(10)

A short historical review about the supersymmetric $6j^S$ symbols:

The conceptualization of supersymmetric 3j or 6j symbols was done in several years spread over time, with sometimes different notations. It started with Pais and Rittenberg [5] in 1975 with a semisimple graded Lie algebra named "graded $\mathfrak{su}(2)$ ". Later in 1977, Scheunert *et al.* [6] have computed the super-rotation Clebsch-Gordan coefficients as a product of usual rotation Clebsch-Gordan coefficients with a scalar factor. Finally, in 1981, Berezin and Tolstoy [7] suggested that the (iso)scalar factors form a pseudo-orthogonal matrix. Subsequently these works were materialized in the paper by Daumens *et al.* [8] dated 1993 with the introduction of properly defined $6j^S$ symbols.

Our definitions of supersymmetric $6j_{\pi}^{S}$ symbols [9] of parity $\pi = \alpha, \beta, \gamma$ were published in 2006, more than a decade after the paper of Daumens *et al.*¹.

These $6j_{\pi}^{S}$ symbols were expressed by a single sum formula over an index t [involving monomials $\Pi_{\pi}(t)$] as shown below:

$$\begin{cases} j_1 \ j_2 \ j_3 \\ J_1 \ J_2 \ J_3 \end{cases} \Big\}_{\pi}^{S} = (-1)^{4\sum j_{\iota}J_{\iota}} \sqrt{\mathsf{R}_{\pi}^{S}} \sum_{t} \frac{(-1)^{t}t!\Pi_{\pi}(t)}{\prod_{i=1}^{i=4} \left(t - [v_i + \frac{1}{2}]\right)! \prod_{j=1}^{j=3} \left([p_j + \frac{1}{2}] - t\right))!} , \tag{11}$$

where \mathtt{R}^S_{π} is a supersymmetric prefactor given by

$$\mathbf{R}_{\pi}^{S} = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} [p_{j} - v_{i}]!}{\prod_{i=1}^{i=4} [v_{i} + \frac{1}{2}]!}.$$
(12)

Detailed expressions for \mathbf{R}^{S}_{α} , \mathbf{R}^{S}_{γ} , \mathbf{R}^{S}_{β} are given in Appendix. Notation [] means 'integer part of'. In contrast to SU(2), OSP(1|2) triangles v_i can be integer or half-integer, however they still satisfy the well known triangular inequalities.

Let us recall the definitions of the parities π and monomials $\Pi_{\pi}(t)$ of degree 0 or 1 in t:

$$\begin{cases} \pi = \alpha \text{ if } \forall i \in [1, 4] v_i \text{ integer}, \\ \pi = \beta \text{ if } \exists \text{ only two distinct } v_i, v_j \text{ integer}, \\ \pi = \gamma \text{ if } \forall i \in [1, 4] v_i \text{ half-integer}. \end{cases}$$
(13)

¹Ref. [8] was explicitly used in 2004 by Livine and Oeckl [4] . On that date our work [9] was not known.

For a parity β , both integer triangles shall be denoted by v, v', both other half-integer by $\overline{v}, \overline{v'}$. The single integer quadrangle is denoted by p, both other half-integer by $\overline{p}, \overline{p'}$. In this case eq. (4) transforms into

$$p + \overline{p} + \overline{p}' = v + v' + \overline{v} + \overline{v}'. \tag{14}$$

Thus a parity π just depends on the quality half-integral or integer of a triangle v_i . Let us note that only the γ parity may contain supersymmetric symbols whose all the spins are half-integers. Our monomials $\Pi_{\pi}(t)$ were defined in [9] as

$$\Pi_{\alpha}(t) = 1, \tag{15}$$

$$\Pi_{\beta}(t) = -t(2j^{\star}_{\beta} + 1) + (\overline{p} + \frac{1}{2})(\overline{p}' + \frac{1}{2}) - vv', \qquad (16)$$

$$\Pi_{\gamma}(t) = -t + 2(J_1j_1 + J_2j_2 + J_3j_3) + (J_1 + j_1 + J_2 + j_2 + J_3 + j_3) + \frac{1}{2}.$$
 (17)

All constants $\Pi_{\alpha}(0)$, $\Pi_{\beta}(0)$, $\Pi_{\gamma}(0)$ are positive integers.

The special spin $\mathfrak{j}_{\beta}^{\star}$ was identified [9] as the vertex common to both half-integer triangles $\overline{v}, \overline{v}'$:

$$2\mathbf{j}_{\beta}^{\star} = \overline{p} + \overline{p}' - v - v' = \overline{v} + \overline{v}' - p.$$
(18)

Other shorter formulas are available:

$$J_1 j_1 + J_2 j_2 + J_3 j_3 = \sum_{i=\iota}^3 j_\iota J_\iota \quad \text{and} \quad J_1 + j_1 + J_2 + j_2 + J_3 + j_3 = \frac{1}{2} \sum_{j=1}^3 p_j.$$
(19)

Oddly enough these 'supersymmetric' quantities are reflected in most parameters of the discriminant necessary for the saddle points computation [10], while the background is that of standard 6*j*.

An appropriate formulation more adapted to the manipulations of \sum_t in [10] is the following

$$\Pi_{\beta}(t) = -(t+1)(2j^{\star}_{\beta}+1) + [(2j^{\star}_{\beta}+1) + (\overline{p}+\frac{1}{2})(\overline{p}'+\frac{1}{2}) - vv'], \qquad (20)$$

$$\Pi_{\gamma}(t) = -(t+1) + \left[2\sum_{\iota=1}^{3} j_{\iota}J_{\iota} + \frac{1}{2}\sum_{j=1}^{3} p_{j} + \frac{3}{2}\right].$$
(21)

The supersymmetric frontal phase $(-1)^{4\sum j_{\iota}J_{\iota}}$ is worthwhile of attention. From eq. (21) in [10]

$$A = 2\sum_{\iota=1}^{3} j_{\iota} J_{\iota} = \sum_{i < j} v_{i} v_{j} - \sum_{k < l} p_{k} p_{l}, \qquad (22)$$

it can be proved hat

$$4\sum j_{\iota}J_{\iota} = \begin{cases} 0 \pmod{2} \text{ if } \pi = \alpha, \\ 1 + \sum_{j} p_{j} \pmod{2} \text{ if } \pi = \gamma, \\ v + v' - p \pmod{2} \text{ if } \pi = \beta. \end{cases}$$
(23)

As $(-1^{k^2N} = +1$ if k even and $(-1)^N$ if k odd, rescaling any spin by k odd leads to

$$(-1)^{4k^2 \sum j_{\iota} J_{\iota}}_{|\mathbf{k} \text{ odd}} = \begin{cases} +1 \text{ if } \pi = \alpha, \\ (-1)^{(1+\sum_j p_j)} \text{ if } \pi = \gamma, \\ (-1)^{(v+v'-p)} \text{ if } \pi = \beta. \end{cases}$$
(24)

In the same way if v_i is integer, kv_i remains integer, whereas if v_i is half-integer kv_i remains half-integer only if k is odd and the formulas for initial parities β, γ can change. Thus care should be taken of k parity itself. Indeed the following transformations can occur:

$$\begin{cases} j_1 \ j_2 \ j_3 \\ J_1 \ J_2 \ J_3 \end{cases}^S_{\boldsymbol{\alpha}} \xrightarrow{\times k} \begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases}^S_{\boldsymbol{\alpha}} \quad \forall k \text{ even or odd,}$$
 (25)

$$\begin{cases} j_1 \ j_2 \ j_3 \\ J_1 \ J_2 \ J_3 \end{cases} \overset{S}{\underset{\beta,\gamma}{\longrightarrow}} \xrightarrow{\times k} \begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \overset{S}{\underset{\alpha}{\longrightarrow}} \quad \text{if } k \text{ even},$$
 (26)

$$\begin{cases} j_1 \ j_2 \ j_3 \\ J_1 \ J_2 \ J_3 \end{cases} \overset{\times k}{\underset{\beta,\gamma}{\longrightarrow}} \overset{\times k}{\underset{kJ_1 \ kJ_2 \ kJ_3}{\longrightarrow}} \begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \overset{S}{\underset{\beta,\gamma}{\longrightarrow}} \quad \text{if } k \text{ odd.}$$
 (27)

The precisions given about 'subleading contributions' in [10] are not altered by the fact that $k \to \infty$ by even or odd integer values.

Due to the presence of integer parts, in \mathbb{R}^{S}_{π} and \sum_{t} , the Stirling's approximation

$$n! = \sqrt{2\pi} e^{(n+\frac{1}{2})\ln(n) - n} = \sqrt{2\pi} e^{n\ln(n) - n + \frac{1}{2}\ln(n)},$$
(28)

could give rise to different formulas when all spins are rescaled by a factor k. Namely

$$[kn]! = (kn)! = \sqrt{2\pi} \ e^{[k\ln(k) - k]n + kn\ln(n) + \frac{1}{2}\ln(kn)} \quad \text{if } n \text{ integer.}$$
(29)

$$[kn]! = \begin{cases} (kn)! & \text{if } k \text{ even and } n \text{ half-integer,} \\ (kn - \frac{1}{2})! & \text{if } k \text{ odd and } n \text{ half-integer.} \end{cases}$$
(30)

$$[kn + \frac{1}{2}]! = \begin{cases} (kn)! & \text{if } k \text{ even and } n \text{ half-integer,} \\ (kn + \frac{1}{2})! & \text{if } k \text{ odd and } n \text{ half-integer.} \end{cases}$$
(31)

It can be checked that only two approximations are modified for k odd and n half-integer:

$$[kn]! = \sqrt{2\pi} \ e^{[k\ln(k) - k]n + kn\ln(n)},\tag{32}$$

$$[kn + \frac{1}{2}]! = \sqrt{2\pi} e^{[k\ln(k) - k]n + kn\ln(n) + \ln(kn)}.$$
(33)

As a direct consequence of eq. (4), we have single out the term $e^{[k \ln(k)-k]}$ because in all the summations to carry out with parities α, β, γ the resulting multiplicative factor will yield a null quantity.

Since we look for a dominant behaviour in first order approximation, the equalities are valid up to a multiplicative factor 1 + 1/k. Throughout this paper we will keep the original functions of Gurau H, h, F, f, denoted here by Gothic letters as $\mathfrak{H}, \mathfrak{H}, \mathfrak{F}, \mathfrak{f}$.

A preliminary analysis shows that, for the three supersymmetric parities α, β, γ , both functions $\mathfrak{h}(j, J)$ and $\mathfrak{f}(x)$ remain unchanged as well as the saddle points values x_{\pm} .

These functions appear under an exponentiated form as $e^{k\mathfrak{h}(j,J)}$ and $e^{k\mathfrak{f}(x)}$. The results were presented as a linear expansion over the six spins as

$$\mathfrak{h}(j,J) = j_1 \mathfrak{h}_{j_1} + j_2 \mathfrak{h}_{j_2} + j_3 \mathfrak{h}_{j_3} + J_1 \mathfrak{h}_{J_1} + J_2 \mathfrak{h}_{J_2} + J_3 \mathfrak{h}_{J_3}, \tag{34}$$

$$\mathfrak{h}_{j_1} = \frac{1}{2} \ln \left\{ \frac{(j_1 + j_2 - j_3)(j_1 - j_2 + j_3)(j_1 + J_2 - J_3)(j_1 - J_2 + J_3)}{(j_1 + j_2 + j_3)(-j_1 + j_2 + j_3)(j_1 + J_2 + J_3)(-j_1 + J_2 + J_3)} \right\} \text{ and so on,}$$
(35)

$$f(x) = i\pi x + x \ln(x) - \sum (x - v_i) \ln(x - v_i) - \sum (p_j - x) \ln(p_j - x).$$
(36)

Equation (35) can be rewritten in terms of p, v:

$$\mathfrak{h}_{j_1} = \frac{1}{2} \ln \left\{ \frac{(p_3 - v_3)(p_2 - v_2)(p_3 - v_2)(p_2 - v_3)}{(p_1 - v_4)(p_1 - v_1)v_1v_4} \right\}.$$
(37)

So we have

$$h_{\alpha}(j,J) = h_{\beta}(j,J) = h_{\gamma}(j,J) \equiv \mathfrak{h}(j,J), \qquad (38)$$

$$h_{j_1}^{\alpha} = h_{j_1}^{\beta} = h_{j_1}^{\gamma} \equiv \mathfrak{h}_{j_1} \text{ and so on.}$$
(39)

$$f_{\alpha}(x) = f_{\beta}(x) = f_{\gamma}(x) \equiv \mathfrak{f}(x). \tag{40}$$

Equation (18) in [10] is the saddle points equation and remains unchanged, then the same holds for the saddle points solutions x_{\pm} lying in the appropriate integration interval. We recall that

$$f'(x) = i\pi \ln(x) - \sum \ln(x - v_i) + \sum \ln(p_j - x) = 0.$$
(41)

Changes with respect to the usual 6j symbols can arise either from \mathbb{R}^S_{π} via $e^{H_{\pi}((j,J)\neq\mathfrak{H}((j,J))}$ or from Σ_x via $e^{F_{\pi}(x)\neq\mathfrak{H}(x)}$. This will lead to distinct variants of Gurau's proof.

3 parity lpha

3.1 Change in the prefactor \mathbb{R}^{S}_{α}

Multiplying the spins of the prefactor \mathtt{R}^S_α by a scaling factor k yields

$$\mathbf{R}_{\alpha}^{S}(k) = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} \left(k(p_{j} - v_{i}) \right)!}{\prod_{i=1}^{i=4} (kv_{i})!},$$
(3.1)

From eq. (29) the numerator and denominator of $\mathtt{R}^S_\alpha(k)$ become

$$\begin{aligned} \text{NumPre}\alpha &= (2\pi)^{6} e^{[k \ln(k) - k](4\sum_{j} p_{j} - 3\sum_{i} v_{i}) + k[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i})]} \\ &\times e^{\frac{1}{2}\sum_{j,i} \ln(p_{j} - v_{i}) + \frac{1}{2}\sum_{j,i} \ln(k)} \\ &= (2\pi)^{6} e^{[k \ln(k) - k](4\sum_{j} p_{j} - 3\sum_{i} v_{i}) + k[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i})]} \\ &\times e^{\frac{1}{2}\sum_{j,i} \ln(p_{j} - v_{i}) + 6 \ln(k))}. \end{aligned}$$
(3.2)

$$DenPre\alpha = (2\pi)^{2} e^{[k \ln(k) - k] \sum_{i} v_{i} + k \sum_{i} v_{i} \ln(v_{i}) + \frac{1}{2} \sum_{i} \ln(k) + \frac{1}{2} \sum_{i} \ln(v_{i})}$$

$$= (2\pi)^{2} e^{[k \ln(k) - k] \sum_{i} v_{i} + k \sum_{i} v_{i} \ln(v_{i}))}$$

$$\times e^{\frac{3}{2} \sum_{i} \ln(v_{i}) - \sum_{i} \ln(v_{i}) + 2 \ln(k)}$$

$$= \frac{(2\pi)^{2}}{\Pi_{i}(v_{i})} e^{[k \ln(k) - k] \sum_{i} v_{i} + k \sum_{i} v_{i} \ln(v_{i}) + \frac{1}{2} \sum_{i} \ln(v_{i}^{3}) + 2 \ln(k)}.$$
(3.3)

That is

$$R^{S}_{\alpha}(k) = (2\pi)^{4} k^{4} \Pi_{i}(v_{i}) e^{k \left[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i}) - \sum_{i} v_{i} \ln(v_{i})\right]} \\ \times e^{\frac{1}{2} \sum_{j,i} \ln \frac{(p_{j} - v_{i})}{v_{i}^{3}}}.$$
(3.4)

The similarities with eqs. (8-9) in [10] are obvious whence

$$\sqrt{\mathbf{R}^{S}_{\alpha}(k)} = (2\pi)^{2} k^{2} \sqrt{\Pi_{i}(v_{i})} e^{\mathfrak{H}(j,J) + k\mathfrak{h}(j,J)}$$
(3.5)

3.2 Changes in the factors Σ_{α} and $F_{\alpha}(x)$

$$\Sigma_{\alpha} = \sum_{\max v_i}^{\min p_j} \frac{(-1)^t t!}{\prod_i (t - v_i)! \prod_j (p_j - t)!}.$$
(3.6)

The only difference with [10] lies in the lack of factor $e^{\ln(kx+1))} \approx e^{\frac{1}{2}\ln(k^2x^2)}$ whence

$$\Sigma_{\alpha} = \frac{1}{(2\pi)^3} \sum_{x=\max v_i}^{x=\min p_j} G_{\alpha}(x), \qquad (3.7)$$

with

$$G_{\alpha}(x) = \frac{1}{k^3} e^{\frac{1}{2} \ln \frac{x}{\Pi(x-v_i)\Pi(p_j-x)}} \times e^{k \{i\pi x + x \ln(x) - [\sum_i (x-v_i) \ln(x-v_i) + \sum_j (p_j-x) \ln(p_j-x)]\}},$$
(3.8)

so that

$$\Sigma_{\alpha} = \frac{1}{(2\pi)^3} \sum_{x=\max v_i}^{x=\min p_j} \frac{1}{k^3} e^{F_{\alpha}(x) + k\mathfrak{f}(x)}.$$
(3.9)

with

$$F_{\alpha}(x) = \frac{1}{2} \ln \frac{x}{\Pi(x - v_i)\Pi(p_j - x)} = \mathfrak{F}(x) - \ln(x).$$
(3.10)

Accordingly

$$\Sigma_{\alpha} = \frac{1}{(2\pi)^3} \frac{1}{k^3} \sum_{x=\max v_i}^{x=\min p_j} \frac{1}{x} e^{\mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(3.11)

3.3 Contribution to the saddle points (parity α)

Identifying Σ_{α} of eq. (3.11) as a Riemann sum with $\frac{1}{k} \to dx$ leads to

$$\Sigma_{\alpha} = \frac{1}{(2\pi)^3 k^2} \int_{\max v_i}^{\min p_j} dx \frac{1}{x} e^{\mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(3.12)

After collecting the three header factors from eqs. (3.5, 3.12) and from the Laplace's approximation [in complex plane] we find

$$\operatorname{Sdl}_{\alpha}(x_s) = \frac{\sqrt{\Pi v_i}}{\sqrt{2\pi k}} \frac{1}{x_s} \frac{1}{\sqrt{-f''(x_s)}} e^{\mathfrak{H}(j,J) + \mathfrak{F}(x_s) + k[\mathfrak{h}(j,J) + \mathfrak{f}(x_s)]},$$
(3.13)

with the same values x_s than those used by Gurau. For reducing the equation lengths let us define

$$\varphi_k^{\theta} = \sum_{\iota=1}^{\iota=3} (kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}} + (kJ_{\iota} + \frac{1}{2})\theta_{J_{\iota}}.$$
(3.14)

Then eq. (45) in [10] changes into

$$\frac{\sqrt{\Pi_{i=1}^{i=4}(v_i)}}{\sqrt{48\pi kV}} \left[\frac{1}{x_+} e^{i\frac{\pi}{4} + i\varphi_k^{\theta}} + \frac{1}{x_-} e^{-i\frac{\pi}{4} - i\varphi_k^{\theta}} \right] \\
= \frac{\sqrt{\Pi v_i}}{\sqrt{48\pi kV}} \left[\left[\frac{1}{x_+} + \frac{1}{x_-} \right] \cos\left(\frac{\pi}{4} + \varphi_k^{\theta}\right) + i \left[\frac{1}{x_+} - \frac{1}{x_-} \right] \sin\left(\frac{\pi}{4} + \varphi_k^{\theta}\right) \right].$$
(3.15)

By using the results and notations of [10] we have

$$\frac{1}{x_{\pm}} = \frac{2A}{(B \pm i\sqrt{\Delta})}$$
$$= \frac{2A(B \mp i\sqrt{\Delta})}{B^2 + \Delta} = \frac{2A(B \mp i\sqrt{\Delta})}{4AC} = \frac{(B \mp i\sqrt{\Delta})}{2C}, \qquad (3.16)$$

whence

$$\frac{1}{x_{+}} + \frac{1}{x_{-}} = \frac{B}{C}$$
 and $\frac{1}{x_{+}} - \frac{1}{x_{-}} = -\frac{i\sqrt{\Delta}}{C}$. (3.17)

As $C = \Pi v_i$ and $\sqrt{\bigtriangleup} = 24V$ we obtain finally

$$\frac{1}{\sqrt{48\pi kV}\sqrt{\Pi v_i}} \Big[B\cos\left(\frac{\pi}{4} + \varphi_k^\theta\right) + 24V\sin\left(\frac{\pi}{4} + \varphi_k^\theta\right) \Big]. \tag{3.18}$$

We recall that coefficient B defined in [10] has the dimension of a volume and writes as

$$B = \left(\sum_{i=1}^{3} j_i J_i \sum_{j=1}^{3} p_j\right) + 2(j_1 j_2 j_3 + j_1 J_2 J_3 + j_2 J_3 J_1 + j_3 J_1 J_2).$$
(3.19)

The square root $\sqrt{\Pi v_i}$ has the dimension of an area.

4 PARITY γ and k odd

4.1 Change in the prefactor R^S_{γ}

Multiplying the spins of the prefactor \mathtt{R}^S_γ by a scaling odd factor k yields

$$\mathbf{R}_{\gamma}^{S}(k) = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} \left(k(p_{j} - v_{i}) - \frac{1}{2} \right)!}{\prod_{i=1}^{i=4} \left(kv_{i} + \frac{1}{2} \right)!},\tag{4.1}$$

$$\operatorname{NumPre}_{\gamma} = (2\pi)^{6} e^{[k \ln(k) - k](4\sum_{j} p_{j} - 3\sum_{i} v_{i}) + k[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i})]}.$$
(4.2)

$$DenPre\gamma = (2\pi)^2 e^{[k\ln(k)-k]\sum_i v_i + \sum_i \ln(k) + \sum_i \ln(v_i)}$$

= $(2\pi)^2 k^4 e^{[k\ln(k)-k]\sum_i v_i + \sum_i \ln(v_i)}.$ (4.3)

Then

$$\sqrt{\mathbb{R}^{S}_{\gamma}(k)} = (2\pi)^{2} \frac{1}{k^{2}} e^{\frac{1}{2}k \left[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i}) - \sum_{i} \ln(v_{i})\right]}.$$
(4.4)

This means that the prefactor $\sqrt{\mathtt{R}_{\gamma}^S(k)}$ is given by

$$\sqrt{\mathbb{R}_{\gamma}^{S}(k)} = (2\pi)^{2} \frac{1}{k^{2}} e^{H_{\gamma}(i,J) + kh_{\gamma}(i,J)}, \qquad (4.5)$$

with

$$H_{\gamma}(i,J) = 0$$
 and $h_{\gamma}(i,J) \equiv \mathfrak{h}(i,J).$ (4.6)

4.2 Changes in the factors Σ_{γ} and $F_{\gamma}(x)$

From eqs. (11) and (21) let be two sums $\Sigma'_{\gamma}, \Sigma''_{\gamma}$ to be gathered to form a Σ_{γ} :

$$\Sigma'_{\gamma} = \sum_{t=\max v_i + \frac{1}{2}}^{t=\min p_j} \frac{(-1)^t (t+1)!}{\prod_i (t-v_i - \frac{1}{2})! \prod_j (p_j - t)!},$$
(4.7)

$$\Sigma_{\gamma}^{\prime\prime} = \sum_{t=\max v_i + \frac{1}{2}}^{t=\min p_j} \frac{(-1)^t t!}{\prod_i (t - v_i - \frac{1}{2})! \prod_j (p_j - t)!},$$
(4.8)

so that

$$\Sigma_{\gamma} = -\Sigma_{\gamma}' + \left[2\sum_{i=1}^{3} j_i J_i + \frac{1}{2}(\sum_{j=1}^{3} p_j) + \frac{3}{2}\right]\Sigma_{\gamma}''.$$
(4.9)

In the first sum Σ'_{γ} we carry out simultaneously the variable changes $t \to kx$ and $v_i \to kv_i, p_j \to kp_j$ and use eqs. (29-30). The result is

$$\Sigma'_{\gamma} = \frac{1}{(2\pi)^3} \sum_{x=\max v_i + \frac{1}{2}}^{x=\min p_j} G'_{\gamma}(x), \qquad (4.10)$$

where

$$G'_{\gamma}(x) = e^{\frac{1}{2}\ln\frac{x^3}{\Pi_j(p_j - x)} + k\{i\pi x + x\ln(x) - [\sum_i (x - v_i)\ln(x - v_i) + \sum_j (p_j - x)\ln(p_j - x)]\}}$$

= $e^{F_{\gamma'}(x) + k\mathfrak{f}(x)},$ (4.11)

$$F_{\gamma'}(x) = e^{\frac{1}{2}\ln\frac{x^3}{\Pi_j(p_j - x)}},$$
(4.12)

so that

$$G'_{\gamma}(x) = \frac{\sqrt{x^3}}{\sqrt{\Pi_j(p_j - x)}} e^{k\mathfrak{f}(x)}.$$
(4.13)

The second sum $\Sigma_{\gamma''}$ is handled in the same way as the first with slight differences

$$\Sigma_{\gamma}^{''} = \frac{1}{(2\pi)^3} \sum_{x=\max v_i + \frac{1}{2}}^{x=\min p_j} G_{\gamma}^{''}(x), \qquad (4.14)$$

with

$$\begin{aligned} G_{\gamma}''(x) &= \\ e^{\frac{1}{2}\ln\frac{1}{k^{2}}\frac{x}{\Pi_{j}(p_{j}-x)} + k\{i\pi x + x\ln(x) - [\sum_{i}(x-v_{i})\ln(x-v_{i}) + \sum_{j}(p_{j}-x)\ln(p_{j}-x)]\}} \\ &= \frac{1}{k}\frac{\sqrt{x}}{\sqrt{\Pi_{j}(p_{j}-x)}}e^{k\mathfrak{f}(x)}. \end{aligned}$$
(4.15)

4.3 Contribution to the saddle points (parity γ)

From eq. (4.13) for \sum_{γ}' and the value of $-f''(x_{\pm})$ found by Gurau, see its equation (33),

$$-f''(x_{\pm}) = \frac{\mp i \sqrt{\Delta}}{x_{\pm} \Pi_j (p_j - x_{\pm})},$$
(4.16)

we derive

$$\sqrt{\frac{2\pi}{k(-f''(x_{\pm}))}} \times \frac{\sqrt{x_{\pm}^3}}{\sqrt{\Pi_j(p_j - x_{\pm})}} \\
= \sqrt{\frac{2\pi}{k}} \sqrt{\frac{x_{\pm}}{\mp i\sqrt{\Delta}}} \sqrt{\Pi_j(p_j - x_{\pm})} \times \frac{\sqrt{x_{\pm}^3}}{\sqrt{\Pi_j(p_j - x_{\pm})}} \\
= \sqrt{\frac{2\pi}{k(\mp i\sqrt{\Delta})}} x_{\pm}^2.$$
(4.17)

The k-dependent header factor comes from eq. (4.5) with $(2\pi)^2 \frac{1}{k^2}$, from eq. (4.10) with $\frac{1}{(2\pi)^3}$ and from the equation just above with $\sqrt{\frac{2\pi}{k}}$ what yields finally $\frac{1}{k} \frac{1}{\sqrt{2\pi k^3}}$. As $\frac{1}{k} \to dx$ the remaining contributions for \sum_{γ}' are

$$\frac{x_{\pm}^2}{\sqrt{2\pi k^3(\mp i\sqrt{\Delta})}} e^{k[\mathfrak{h}(j,J)+\mathfrak{f}(x_{\pm})]}.$$
(4.18)

For $\sum_{\gamma}^{\prime\prime}$ we have the following contribution

$$\frac{1}{k} \frac{x_{\pm}}{\sqrt{2\pi k^3 (\mp i\sqrt{\Delta})}} e^{k[\mathfrak{h}(j,J) + \mathfrak{f}(x_{\pm})]},\tag{4.19}$$

which must be weighted by its multiplicative coefficient $\left[2\sum_{\iota=1}^{3} j_{\iota}J_{\iota} + \frac{1}{2}(\sum_{j=1}^{3} p_{j}) + \frac{3}{2}\right] \approx k^{2}\sum_{\iota=1}^{3} 2j_{\iota}J_{\iota}$. Then \sum_{γ}' is no longer dominant that is the only one contribution comes from \sum_{γ}'' has the form

$$\sum_{\iota=1}^{3} 2j_{\iota}J_{\iota} \Big] \frac{k^{2}}{k} \frac{x_{\pm}}{\sqrt{2\pi k^{3}(\mp i\sqrt{\Delta})}} e^{k[\mathfrak{h}(j,J)+\mathfrak{f}(x_{\pm})]} \\ = \Big[\sum_{\iota=1}^{3} 2j_{\iota}J_{\iota}\Big] \frac{\sqrt{\pm i} x_{\pm}}{\sqrt{48\pi kV}} e^{k[\mathfrak{h}(j,J)+\mathfrak{f}(x_{\pm})]}.$$
(4.20)

As a function of the parameter A defined in eq. (21) in [10] for the saddle points equation we obtain

$$\operatorname{Sdl}_{\gamma}(x_{\pm}) = A \frac{x_{\pm}}{\sqrt{48\pi kV}} e^{\pm \imath \frac{\pi}{4} + k[\mathfrak{h}(j,J) + \mathfrak{f}(x_{\pm})]}.$$
(4.21)

From eq. (44) in [10] the real part of

$$e^{k[j_1(\mathfrak{h}_{j_1}+\mathfrak{f}_{j_1})+j_2(\mathfrak{h}_{j_2}+\mathfrak{f}_{j_2})+j_3(\mathfrak{h}_{j_3}+\mathfrak{f}_{j_3})+J_1(\mathfrak{h}_{J_1}+\mathfrak{f}_{J_1})+J_2(\mathfrak{h}_{J_2}+\mathfrak{f}_{J_2})+J_3(\mathfrak{h}_{J_3}+\mathfrak{f}_{J_3})]}$$

is zero, then its remains only the imaginary part (the h_{j_i,J_i} being real) $e^{\pm i k [j_1 \theta_{j_1} + j_2 \theta_{j_2} + j_3 \theta_{j_3} + J_1 \theta_{J_1} + J_2 \theta_{J_2} + J_3 \theta_{J_3}]}$ whence

$$Sdl_{\gamma}(x_{\pm}) = A \frac{x_{\pm}}{\sqrt{48\pi kV}} e^{\pm i\frac{\pi}{4}} \\ \times e^{\pm ik[j_{1}\theta_{j_{1}} + j_{2}\theta_{j_{2}} + j_{3}\theta_{j_{3}} + J_{1}\theta_{J_{1}} + J_{2}\theta_{J_{2}} + J_{3}\theta_{J_{3}}]}.$$
(4.22)

Let us define a new angle $\varphi^{\theta}_{k_{\gamma}}$ as

$$\varphi_{k\gamma}^{\theta} = \sum_{\iota=1}^{\iota=3} k(j_{\iota}\theta_{j_{\iota}} + J_{i}\theta_{J_{\iota}}).$$
(4.23)

Then the final expression of $\operatorname{Sdl}_{\gamma}(x_{\pm})$ writes

$$\operatorname{Sdl}_{\gamma}(x_{\pm}) = A \frac{x_{\pm}}{\sqrt{48\pi kV}} e^{\pm i \{\frac{\pi}{4} + \varphi_{k\gamma}^{\theta}\}}.$$
(4.24)

Finally

$$\operatorname{Sdl}_{\gamma}(x_{+}) + \operatorname{Sdl}_{\gamma}(x_{-}) = \frac{1}{\sqrt{48\pi kV}} \times \left[(Ax_{+})e^{i\left\{\frac{\pi}{4} + \varphi_{k\gamma}^{\theta}\right\}} + (Ax_{-})e^{-i\left\{\frac{\pi}{4} + \varphi_{k\gamma}^{\theta}\right\}} \right].$$
(4.25)

Since $(A x_{\pm}) = \frac{1}{2} [B \pm i \sqrt{\triangle}]$ the sum in (4.25) writes as

$$\operatorname{Sdl}_{\gamma}(x_{+}) + \operatorname{Sdl}_{\gamma}(x_{-}) = \frac{1}{\sqrt{48\pi kV}} \times \left\{ B\cos\left(\frac{\pi}{4} + \varphi_{k_{\gamma}}^{\theta}\right) - 24V\sin\left(\frac{\pi}{4} + \varphi_{k_{\gamma}}^{\theta}\right) \right\}.$$
(4.26)

5 PARITY β and k odd

A useful correlation table for parity β is shown below.

spin \mathfrak{j}_{β}^* vertex common to both triangles $\overline{v}, \overline{v}'$								
$\overline{v}, \overline{v}' \frac{1}{2}$ integer	\mathfrak{j}_{eta}^*	p integer	v, v' integer	$\overline{p}, \overline{p}' \frac{1}{2}$ integer				
v_4, v_1	j_1	p_1	v_2, v_3	p_2, p_3				
v_2, v_1	j_2	p_2	v_{3}, v_{4}	p_3, p_1				
v_{3}, v_{1}	j_3	p_3	v_4, v_2	p_1, p_2				
v_2, v_3	J_1	p_1	v_4, v_1	p_2, p_3				
v_3, v_4	J_2	p_2	v_2, v_1	p_3, p_1				
v_4, v_2	J_3	p_3	v_3, v_1	p_1, p_2				

Let us define \mathfrak{J}_{β}^* as the column companion of the spin \mathfrak{j}_{β}^* and represent it by an empty oval. An appropriate diagrammatic representation looks as follows

 \mathfrak{j}_β^* represented by $\bullet,\,\mathfrak{J}_\beta^*$ by an oval, only both half-integer $\overline{v},\overline{v}'$ are drawn

•					\mathbf{X}°
$\mathfrak{j}_\beta^*=j_1$	$\mathfrak{j}_\beta^*=j_2$	$\mathfrak{j}_\beta^*=j_3$	$\mathfrak{j}_\beta^*=J_3$	$\mathfrak{j}_\beta^*=J_2$	$\mathfrak{j}_\beta^*=J_3$
$\mathfrak{J}_{\beta}^* = J_1$	$\mathfrak{J}_{eta}^*=J_2$	$\mathfrak{J}_{eta}^* = J_3$	$\mathfrak{J}_{eta}^{*}=j_{3}$	$\mathfrak{J}_{eta}^{*}=j_{2}$	$\mathfrak{J}_{eta}^*=j_3$

5.1 Change in the prefactor \mathbb{R}^S_β

Multiplying the spins of the prefactor \mathbb{R}^S_β by a scaling odd factor k yields

$$\begin{aligned} \mathbf{R}_{\beta}^{S}(k) &= (k(p-v))!(k(p-v'))!k((p-\overline{v}) - \frac{1}{2})!(k(p-\overline{v}') - \frac{1}{2})!\\ &\times (k(\overline{p} - \overline{v}))!(k(\overline{p} - \overline{v}'))!(k(\overline{p} - v) - \frac{1}{2})!(k(\overline{p} - v') - \frac{1}{2})!\\ &\times (k(\overline{p}' - \overline{v}))!(k(\overline{p}' - \overline{v}'))!(k(\overline{p}' - v) - \frac{1}{2})!(k(\overline{p}' - v') - \frac{1}{2})!\\ &\times (kv!kv'!)^{-1}(k\overline{v} + \frac{1}{2})!(k\overline{v}' + \frac{1}{2})!)^{-1}. \end{aligned}$$
(5.1)

From eqs. (32-33) the numerator of $\mathtt{R}^S_{\beta}(k)$ becomes

$$\operatorname{NumPre}_{\beta} = (2\pi)^{6} e^{[k \ln(k) - k](4\sum_{j} p_{j} - 3\sum_{i} v_{i}) + k[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i})]} \times e^{\frac{1}{2}[\ln(p-v) + \ln(p-v') + \ln(\overline{p}-\overline{v}) + \ln(\overline{p}-\overline{v}') + \ln(\overline{p}'-\overline{v}')] + 3\ln(k)}.$$
(5.2)

and the denominator is

$$DenPre\beta = (2\pi)^2 e^{[k\ln(k)-k](v+v'+\overline{v}+\overline{v}')}$$

$$\times e^{k[v\ln(v)+v'\ln(v')+\overline{v}\ln(\overline{v})+\overline{v}'\ln(\overline{v}')]}$$

$$\times e^{\frac{1}{2}[\ln(kv)+\ln(kv')]+\ln(k\overline{v})+\ln(k\overline{v}')}$$

$$= (2\pi)^2 e^{[k\ln(k)-k]\sum_i v_i+k[\sum_i v_i\ln(v_i)]}$$

$$\times e^{3\ln(k)} e^{\frac{1}{2}\{\ln(v)+\ln(v')\}+\ln(\overline{v})+\ln(\overline{v}')}.$$
(5.3)

Note the lack of frontal factor in $\frac{1}{k^m}$ in $\mathbf{R}^S_{\beta}(k)$. Whence

$$R_{\beta}^{S}(k) = (2\pi)^{4} e^{k \left[\sum_{j,i} (p_{j} - v_{i}) \ln(p_{j} - v_{i}) - \sum_{i} v_{i} \ln(v_{i})\right]} \\ \times e^{\frac{1}{2} \ln \left\{ \frac{(p - v)(p - v')(\overline{p} - \overline{v})(\overline{p} - \overline{v})(\overline{p}' - \overline{v})(\overline{p}' - \overline{v}')}{(vv')(\overline{v}^{2} \overline{v}'^{2})} \right\}}.$$
(5.4)

Thanks to the correlation table and the definition of the spin \mathfrak{J}_{β}^{*} it can be proved that

$$e^{\frac{1}{2}\ln\left\{\frac{(p-v)(p-v')(\overline{p}-\overline{v})(\overline{p}-\overline{v}')(\overline{p}'-\overline{v})(\overline{p}'-\overline{v}')}{(vv')(\overline{v}^2\,\overline{v}'^2)}\right\}} = \frac{(p-v)(p-v')}{\overline{v}\,\overline{v}'}e^{\mathfrak{h}\mathfrak{J}^*_{\beta}},\tag{5.5}$$

where the header fraction is dimensionless. Then

$$\mathbf{R}^{S}_{\beta}(k) = (2\pi)^{4} e^{k[\sum_{j,i}(p_{j}-v_{i})\ln(p_{j}-v_{i})-\sum_{i}v_{i}\ln(v_{i})]} \times \frac{(p-v)(p-v')}{\overline{v}\,\overline{v'}} e^{\mathfrak{h}_{\mathfrak{J}^{*}_{\beta}}}.$$
(5.6)

Finally

$$\sqrt{\mathbf{R}_{\beta}^{S}(k)} = (2\pi)^{2} \sqrt{\frac{(p-v)(p-v')}{\overline{v}\,\overline{v'}}} e^{\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}} + k\mathfrak{h}(i,J)}.$$
(5.7)

5.2 Changes in the factors Σ_{β} and $F_{\beta}(x)$

From eqs.(11) and (20) we define two sums $\Sigma'_{\beta}, \Sigma''_{\beta}$ to be gathered to form a a Σ_{β} :

$$\Sigma'_{\boldsymbol{\beta}} = \sum_{t_{\min}}^{t_{\max}} \frac{(-1)^t (t+1)!}{(t-v)! (t-v')! (t-\overline{v}-\frac{1}{2})! (t-\overline{v}'-\frac{1}{2})! (p-t)! (\overline{p}+\frac{1}{2}-t)! (\overline{p}'+\frac{1}{2}-t)!},$$
(5.8)

$$\Sigma_{\beta}^{\prime\prime} = \sum_{t_{\min}}^{t_{\max}} \frac{(-1)^{t} t!}{(t-v)!(t-v^{\prime})!(t-\overline{v}-\frac{1}{2})!(t-\overline{v}^{\prime}-\frac{1}{2})!(p-t)!(\overline{p}+\frac{1}{2}-t)!(\overline{p}^{\prime}+\frac{1}{2}-t)!},$$
(5.9)

so that

$$\Sigma_{\beta} = -(2j_{\beta}^{\star} + 1)\Sigma_{\beta}' + \left[(2j_{\beta}^{\star} + 1) + (\overline{p} + \frac{1}{2})(\overline{p}' + \frac{1}{2}) - vv'\right]\Sigma_{\beta}'',$$
(5.10)

with

$$t_{\min} = \max\left(v, v', \overline{v} + \frac{1}{2}, \overline{v}' + \frac{1}{2}\right) \text{ and } t_{\max} = \min\left(p, \overline{p} + \frac{1}{2}, \overline{p}' + \frac{1}{2}\right).$$
(5.11)

Computation of Σ'_{β} is very similar to that of the standard sum Σ analyzed by Gurau whereas Σ''_{β} looks like our Σ_{α} explicited in sect. **3.2**.

With all spins multiplied by the factor k, and changes $t \to kx, p, v \to kp, kv, \Sigma'_{\beta}(k)$ rewrites as the sum $\sum_{x=t_{\min}}^{x=t_{\max}}$ over the quotient below

$$\frac{(-1)^{kx}(kx+1)!}{(k(x-v'))!(k(x-\overline{v})-\frac{1}{2})!(k(x-\overline{v}')-\frac{1}{2})!(k(p-x))!(k(\overline{p}-x)+\frac{1}{2})!(k(\overline{p}'-x)+\frac{1}{2})!}.$$
(5.12)

By using factorial approximations for large k we write $\Sigma'_{\pmb{\beta}}$ in detail as

$$\Sigma'_{\beta} = \frac{1}{(2\pi)^3} \sum_{x=t\,\text{min}}^{x=t\,\text{max}} G'_{\beta}(x), \qquad (5.13)$$

with

$$\begin{aligned} G'_{\beta}(x) &= e^{[k\ln(k)-k]x+k\{i\pi x+x\ln(x)\}+\frac{1}{2}\ln(k^{3}x^{3})} \\ &\times e^{-\left\{\frac{1}{2}\ln k(x-v)+[k\ln(k)-k](x-v')+k(x-v')\ln(x-v')+\frac{1}{2}\ln k(x-v')\right\}} \\ &\times e^{-\left\{[k\ln(k)-k](x-v)+k(x-v)\ln(x-v)\right\}} \\ &\times e^{-\left\{[k\ln(k)-k](x-\overline{v})+k(x-\overline{v})\ln(x-\overline{v})+[k\ln(k)-k](x-\overline{v}')+k(x-\overline{v}')\ln(x-\overline{v}')\right\}} \\ &\times e^{-\left\{[k\ln(k)-k](p-x)+k(p-x)\ln(p-x)+\frac{1}{2}\ln k(p-x)\right\}} \\ &\times e^{-\left\{[k\ln(k)-k](\overline{p}-x)+k(\overline{p}-x)\ln(\overline{p}-x)+\ln k(\overline{p}-x)\right\}} \\ &\times e^{-\left\{[k\ln(k)-k](\overline{p}'-x)+k(\overline{p}'-x)\ln(\overline{p}'-x)+\ln k(\overline{p}'-x)\right\}}. \end{aligned}$$
(5.14)

All rearrangements done it remains

$$G'_{\beta}(x) = \frac{1}{k^2} e^{\frac{1}{2} \ln \frac{x^3}{\Pi(x-v_i)\Pi(p_j-x)} + \frac{1}{2} \ln \frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)}} \times e^{k\{i\pi x + x\ln(x) - \sum(x-v_i)\ln(x-v_i) - \sum(p_j-x)\ln(p_j-x)\}}.$$
(5.15)

It results in

$$\Sigma'_{\beta}(k) = \frac{1}{(2\pi)^3} \sum_{x=t_{\min}}^{x=t_{\max}} \frac{1}{k^2} e^{\frac{1}{2} \ln \frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + F_{\beta'}(x) + kf_{\beta'}(x)},$$
(5.16)

with

$$F_{\beta'}(x) = \frac{1}{2} \ln \frac{x^3}{\Pi(x - v_i)\Pi(p_j - x)} \equiv \mathfrak{F}(x),$$
(5.17)

$$f_{\beta'}(x) \equiv \mathfrak{f}(x). \tag{5.18}$$

In the same way we derive

$$\Sigma_{\beta}^{\prime\prime}(k) = \frac{1}{(2\pi)^3} \sum_{x=t_{\min}}^{x=t_{\max}} \frac{1}{k^3} e^{\frac{1}{2} \ln \frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + F_{\beta^{\prime\prime}}(x) + kf_{\beta^{\prime\prime}}(x)},$$
(5.19)

with

$$F_{\beta''}(x) = \frac{1}{2} \ln \frac{x}{\Pi(x - v_i)\Pi(p_j - x)} \equiv \mathfrak{F}(x) - \ln(x),$$
(5.20)

so that

$$e^{F_{\beta^{\prime\prime}}(x)} = \frac{1}{x}e^{\mathfrak{F}(x)}$$
 and $f_{\beta^{\prime\prime}}(x) \equiv \mathfrak{f}(x).$ (5.21)

5.3 Contribution to the saddle points (parity β)

Identifying Σ'_{β} of eq. (5.16) as a Riemann sum with $\frac{1}{k} \to dx$ leads to

$$\Sigma'_{\beta}(k) = \frac{1}{(2\pi)^3 k} \int_{t_{\min}}^{t_{\max}} dx \, e^{\frac{1}{2} \ln \frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + \mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(5.22)

Equation (5.10), with $j^{\star}_{\beta} \to k j^{\star}_{\beta}$, gives the following dominant contribution

$$-\left(2k\mathfrak{j}_{\beta}^{\star}+1\right)\Sigma_{\beta}'(k) = -\frac{2\mathfrak{j}_{\beta}^{\star}}{(2\pi)^3}\int_{t_{\min}}^{t_{\max}}dx\,e^{\frac{1}{2}\ln\frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]}+\mathfrak{F}(x)+k\mathfrak{f}(x)}.$$
(5.23)

In the same way

$$\Sigma_{\beta}^{\prime\prime}(k) = \frac{1}{(2\pi)^3 k^2} \int_{t_{\min}}^{t_{\max}} dx \, \frac{1}{x} \, e^{\frac{1}{2} \ln \frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + \mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(5.24)

Its coefficient in (5.10), $[(2j^{\star}_{\beta}+1)+(\overline{p}+\frac{1}{2})(\overline{p}'+\frac{1}{2})-vv']$, is rescaled into

$$[(2kj_{\beta}^{\star}+1) + (k\overline{p} + \frac{1}{2})(k\overline{p}' + \frac{1}{2}) - k^2vv'] = k^2[\overline{p}\,\overline{p}' - vv'] + k[2j_{\beta}^{\star} + \frac{1}{2}(\overline{p} + \overline{p}')] + \frac{3}{4}.$$
 (5.25)

The $\Sigma_{\pmb{\beta}}^{\prime\prime}(k)$ contribution is also dominant and can be written down as

$$\frac{1}{x} \frac{\left[\overline{p}\,\overline{p}' - vv'\right]}{(2\pi)^3} \int_{t_{\min}}^{t_{\max}} dx \, e^{\frac{1}{2}\ln\frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + \mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(5.26)

Finally the integral approximation has the form

$$\Sigma_{\boldsymbol{\beta}} = \left(-2\mathfrak{j}_{\boldsymbol{\beta}}^{\star} + \frac{\left[\overline{p}\,\overline{p}' - vv'\right]}{x}\right) \frac{1}{(2\pi)^3} \int_{x=t_{\min}}^{x=t_{\max}} e^{\frac{1}{2}\ln\frac{(x-\overline{v})(x-\overline{v}')}{(\overline{p}-x)(\overline{p}'-x)]} + \mathfrak{F}(x) + k\mathfrak{f}(x)}.$$
(5.27)

Writing f(x) as the equation (27) used by Gurau for the saddle points

$$\mathfrak{f}(x_{\pm}) = j_1 \mathfrak{f}_{j_1}(x_{\pm}) + j_2 \mathfrak{f}_{j_2}(x_{\pm}) + j_3 \mathfrak{f}_{j_3}(x_{\pm}) + J_1 \mathfrak{f}_{J_1}(x_{\pm}) + J_2 \mathfrak{f}_{J_2}(x_{\pm}) + J_3 \mathfrak{f}_{J_3}(x_{\pm}), \tag{5.28}$$

using its result, see eqs. (26-27-44) in [10],

$$\Re(\mathfrak{f}_{\mathfrak{j}_{\beta}^{*}}(x_{+})) = \Re(\mathfrak{f}_{\mathfrak{j}_{\beta}^{*}}(x_{-})) = -\mathfrak{h}_{\mathfrak{j}_{\beta}^{*}} \quad (\text{not depending on } x_{+} \text{ or } x_{-}),$$
(5.29)

our correlation table for \mathfrak{j}_β^* shows that the following identification holds

$$\Re\left\{\ln\frac{(x_{\pm}-\overline{v})(x_{\pm}-\overline{v}')}{(\overline{p}-x_{\pm})(\overline{p}'-x_{\pm})}\right\} \equiv \Re\{\mathfrak{f}_{\mathfrak{j}_{\beta}^{*}}(x_{\pm})\} = -\mathfrak{h}_{\mathfrak{j}_{\beta}^{*}}.$$
(5.30)

In contrast the imaginary part is

$$\Im\left(\mathfrak{f}_{\mathfrak{j}_{\beta}^{*}}(x_{\pm})\right) = \pm \theta_{\mathfrak{j}_{\beta}^{*}}.$$
(5.31)

It follows that

$$\Sigma_{\boldsymbol{\beta}}(x_{\pm}) = e^{-\frac{1}{2}\mathfrak{h}_{j_{\boldsymbol{\beta}}^{\star}}} \left(-2\mathfrak{j}_{\boldsymbol{\beta}}^{\star} + \frac{\left[\overline{p}\,\overline{p}' - vv'\right]}{x_{\pm}}\right) \frac{1}{(2\pi)^3} e^{\pm\frac{\imath}{2}\theta_{j_{\boldsymbol{\beta}}^{\star}} + \mathfrak{F}(x_{\pm}) + k\mathfrak{f}(x_{\pm})}.$$
(5.32)

After the Laplace's approximation the final header factor is simply $\frac{1}{\sqrt{2\pi k}}$. Ultimately from eq. (5.7) the saddle points contributions for parity β have the form

$$\operatorname{Sdl}_{\beta}(x_{\pm}) = \frac{1}{\sqrt{2\pi k}} \sqrt{\frac{(p-v)(p-v')}{\overline{v}\,\overline{v}'}} e^{\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}} \\ \times e^{-\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}} \frac{\mathfrak{F}(x_{\pm})}{\sqrt{-f''(x_{\pm})}} \left(-2\mathfrak{j}_{\beta}^{\star} + \frac{[\overline{p}\,\overline{p}'-vv']}{x_{\pm}}\right) \\ \times e^{\frac{\pm i}{2}\theta_{\mathfrak{J}_{\beta}^{\star}} + k[\mathfrak{h}(i,J) + \mathfrak{f}(x_{\pm})]}.$$
(5.33)

From eqs. (32-33) in [10] clearly

$$-f''(x_{-}) = \frac{+i\sqrt{\Delta}}{x_{-}\Pi((p_{j} - x_{-}))}.$$
(5.34)

From eq. (35) in [10] we obtain

$$\begin{aligned} \mathsf{Sdl}_{\beta}(x_{\pm}) &= \frac{1}{\sqrt{2\pi k}} \sqrt{\frac{(p-v)(p-v')}{\overline{v} \, \overline{v}'}} e^{\frac{1}{2} \mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}} \\ &\times e^{k[j_{1}(\mathfrak{h}_{j_{1}}+\mathfrak{f}_{j_{1}})+j_{2}(\mathfrak{h}_{j_{2}}+\mathfrak{f}_{j_{2}})+j_{3}(\mathfrak{h}_{j_{3}}+\mathfrak{f}_{j_{3}})]} \\ &\times e^{k[J_{1}(\mathfrak{h}_{J_{1}}+\mathfrak{f}_{J_{1}})+J_{2}(\mathfrak{h}_{J_{2}}+\mathfrak{f}_{J_{2}})+J_{3}(\mathfrak{h}_{J_{3}}+\mathfrak{f}_{J_{3}})]} \\ &\times e^{\frac{1}{2}[\mathfrak{f}_{j_{1}}+\mathfrak{f}_{j_{2}}+\mathfrak{f}_{j_{3}}+\mathfrak{f}_{J_{1}}+\mathfrak{f}_{J_{2}}+\mathfrak{f}_{J_{3}}]} \\ &\times e^{-\frac{1}{2}\mathfrak{h}_{j^{*}}} \frac{1}{\sqrt{\mp i\sqrt{\Delta}}} \left(-2\mathfrak{j}_{\beta}^{\star}+\frac{[\overline{p}\,\overline{p}'-vv']}{x_{\pm}}\right) e^{\pm\frac{i}{2}\theta_{j^{*}_{\beta}}}. \end{aligned}$$
(5.35)

From $\triangle = (24)^2 V^2$ and $\frac{1}{\sqrt{\mp i}} = e^{\pm i \frac{\pi}{4}}$ we derive

$$\frac{1}{\sqrt{\mp i\sqrt{\Delta}}} = \frac{1}{\sqrt{24V}} e^{\pm i\frac{\pi}{4}}.$$
(5.36)

The result for $\operatorname{Sdl}_{\beta}(x_{\pm})$ becomes the complex expression

$$\begin{aligned} \mathsf{Sdl}_{\beta}(x_{\pm}) &= \frac{1}{\sqrt{48\pi kV}} \sqrt{\frac{(p-v)(p-v')}{\overline{v}\,\overline{v}'}} e^{\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}}} \\ &\times e^{k[j_{1}(\mathfrak{h}_{j_{1}}+\mathfrak{f}_{j_{1}})+j_{2}(\mathfrak{h}_{j_{2}}+\mathfrak{f}_{j_{2}})+j_{3}(\mathfrak{h}_{j_{3}}+\mathfrak{f}_{j_{3}})+J_{1}(\mathfrak{h}_{J_{1}}+\mathfrak{f}_{J_{1}})+J_{2}(\mathfrak{h}_{J_{2}}+\mathfrak{f}_{J_{2}})+J_{3}(\mathfrak{h}_{J_{3}}+\mathfrak{f}_{J_{3}})] \\ &\times e^{\frac{1}{2}[\mathfrak{f}_{j_{1}}+\mathfrak{f}_{j_{2}}+\mathfrak{f}_{j_{3}}+\mathfrak{f}_{J_{1}}+\mathfrak{f}_{J_{2}}+\mathfrak{f}_{J_{3}}]} \\ &\times e^{-\frac{1}{2}\mathfrak{h}_{j_{\beta}^{*}}} e^{\pm\imath\frac{\pi}{4}} \left(-2\mathfrak{j}_{\beta}^{*}+\frac{[\overline{p}\,\overline{p}'-vv']}{x_{\pm}}\right) e^{\pm\frac{\imath}{2}\theta_{j_{\beta}^{*}}}. \end{aligned}$$
(5.37)

From eq. (44) in [10] the real part of

 $e^{k[j_1(\mathfrak{h}_{j_1}+\mathfrak{f}_{j_1})+j_2(\mathfrak{h}_{j_2}+\mathfrak{f}_{j_2})+j_3(\mathfrak{h}_{j_3}+\mathfrak{f}_{j_3})+J_1(\mathfrak{h}_{J_1}+\mathfrak{f}_{J_1})+J_2(\mathfrak{h}_{J_2}+\mathfrak{f}_{J_2})+J_3(\mathfrak{h}_{J_3}+\mathfrak{f}_{J_3})]}$

is zero, then its remains only the imaginary part (the h_{j_i,J_i} being real) which is $e^{\pm ik[j_1\theta_{j_1}+j_2\theta_{j_2}+j_3\theta_{j_3}+J_1\theta_{J_1}+J_2\theta_{J_2}+J_3\theta_{J_3}]}$ whence

$$\begin{aligned} \operatorname{Sdl}_{\beta}(x_{\pm}) &= \frac{1}{\sqrt{48\pi kV}} \sqrt{\frac{(p-v)(p-v')}{\overline{v}\,\overline{v}'}} e^{\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}} \\ &\times e^{\pm ik[j_{1}\theta_{j_{1}}+j_{2}\theta_{j_{2}}+j_{3}\theta_{j_{3}}+J_{1}\theta_{J_{1}}+J_{2}\theta_{J_{2}}+J_{3}\theta_{J_{3}}]} \\ &\times e^{\pm i\frac{1}{2}[\theta_{j_{1}}+\theta_{j_{2}}+\theta_{j_{3}}+\theta_{J_{1}}+\theta_{J_{2}}+\theta_{J_{3}}]} \\ &\times e^{\frac{1}{2}\Re[\mathfrak{f}_{j_{1}}+\mathfrak{f}_{j_{2}}+\mathfrak{f}_{j_{3}}+\mathfrak{f}_{J_{1}}+\mathfrak{f}_{J_{2}}+\mathfrak{f}_{J_{3}}]} \\ &\times e^{-\frac{1}{2}\mathfrak{h}_{j_{\beta}^{\star}}} e^{\pm i\frac{\pi}{4}} \left(-2\mathfrak{j}_{\beta}^{\star}+\frac{[\overline{p}\,\overline{p}'-vv']}{x_{\pm}}\right) e^{\pm \frac{s}{2}\theta_{j_{\beta}^{\star}}}. \end{aligned}$$
(5.38)

That is

$$\begin{aligned} \operatorname{Sdl}_{\beta}(x_{\pm}) &= \frac{1}{\sqrt{48\pi kV}} \sqrt{\frac{(p-v)(p-v')}{\overline{v}\,\overline{v}'}} \\ &\times e^{-\frac{1}{2}[\mathfrak{h}_{j_{1}}+\mathfrak{h}_{j_{2}}+\mathfrak{h}_{j_{3}}+\mathfrak{h}_{J_{1}}+\mathfrak{h}_{J_{2}}+\mathfrak{h}_{J_{3}}+\mathfrak{h}_{j_{\beta}^{*}}-\mathfrak{h}_{\mathfrak{I}_{\beta}^{*}}]} \\ &\times \left(\left[v+v'-\overline{p}-\overline{p}'\right] + \frac{\left[\overline{p}\,\overline{p}'-vv'\right]}{x_{\pm}} \right) e^{\pm i(\frac{\pi}{4}+\varphi_{k_{\beta}}^{\theta})}, \end{aligned}$$
(5.39)

where a new angle $\varphi^{\theta}_{k_{\beta}}$ depending on $\theta_{\mathbf{j}^{*}_{\beta}}$ is defined via

$$\varphi_{k_{\beta}}^{\theta} = \sum_{\iota=1}^{\iota=3} \left[(kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}} + (kJ_{\iota} + \frac{1}{2})\theta_{J_{\iota}} \right] + \frac{1}{2}\theta_{j_{\beta}^{*}}.$$
(5.40)

We have

$$\frac{1}{x_{+}} = \frac{(B - i\sqrt{\Delta})}{2C} \quad \text{and} \quad \frac{1}{x_{-}} = \frac{(B + i\sqrt{\Delta})}{2C}, \tag{5.41}$$

whence adding the contribution + and -, taken into account $C = \Pi(v_i)$, $\sqrt{\Delta} = 24V$, we obtain

$$\frac{1}{\Pi(v_i)}\bigg\{\Big\{2\Pi(v_i)[v+v'-\overline{p}-\overline{p}']+B[\overline{p}\,\overline{p}'-vv']\Big\}\cos(\frac{\pi}{4}+\varphi^{\theta}_{k_{\beta}})+24V[\overline{p}\,\overline{p}'-vv']\sin(\frac{\pi}{4}+\varphi^{\theta}_{k_{\beta}})\bigg\}.$$

A preliminary result is then

$$\begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \overset{S}{\underset{\beta}{\overset{\beta}{\Rightarrow}}} \approx \frac{(-1)^{(v+v'-p)}}{\sqrt{48\pi kV} \Pi(v_i)} \sqrt{\frac{(p-v)(p-v')}{\overline{v} \,\overline{v}'}} \\ \times e^{-\frac{1}{2}[\mathfrak{h}_{j_1}+\mathfrak{h}_{j_2}+\mathfrak{h}_{j_3}+\mathfrak{h}_{J_1}+\mathfrak{h}_{J_2}+\mathfrak{h}_{J_3}] - \frac{1}{2}[\mathfrak{h}_{j_\beta}^* - \mathfrak{h}_{\mathfrak{J}_\beta}^*]} \\ \times \left\{ \left\{ 2\Pi(v_i)[v+v'-\overline{p}-\overline{p}'] + B[\overline{p}\,\overline{p}'-vv'] \right\} \cos(\frac{\pi}{4}+\varphi_{k_\beta}^\theta) + 24V[\overline{p}\,\overline{p}'-vv'] \sin(\frac{\pi}{4}+\varphi_{k_\beta}^\theta) \right\}. \tag{5.42}$$

Factor $\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}$ is compensatory, *i.e.* $\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}$ term vanishes whereas $\frac{1}{2}\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}$ becomes $\mathfrak{h}_{\mathfrak{J}_{\beta}^{\star}}$. Unfortunately this evidence will be more or less lost in the general formulas we try to explicit below.

Computation of $e^{\mathfrak{h}_{\mathfrak{I}_{\beta}^{*}}-\mathfrak{h}_{j_{\beta}^{*}}}$ yields

$$e^{\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}}-\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}}} = e^{\frac{1}{2}\ln\left\{\frac{(\overline{p}-\overline{v})(\overline{p}-\overline{v}')(\overline{p}'-\overline{v})(\overline{p}'-\overline{v}')(p-\overline{v}')(\overline{p}-\overline{v}')(\overline{p}-\overline{v}')}{(p-v)(\overline{p}-v)(\overline{p}-v')(\overline{p}'-v')vv'}\right\}},\tag{5.43}$$

whence

$$e^{-\frac{1}{2}[\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}}-\mathfrak{h}_{\mathfrak{J}_{\beta}^{*}}]} = \sqrt[4]{\frac{(p-v)(p-v')(\overline{p}-v)(\overline{p}-v'(\overline{p}'-v)(\overline{p}'-v')vv'}{(\overline{p}-\overline{v})(\overline{p}-\overline{v}')(\overline{p}'-\overline{v})(\overline{p}'-\overline{v})(p-\overline{v}')vv'}}.$$
(5.44)

Besides from eqs. (8-9-11) in [10] we can write

$$e^{-\mathfrak{H}(i,J)} = e^{-\frac{1}{2}\sum_{j=1}^{3}(\mathfrak{h}_{j_{j}} + \mathfrak{h}_{J_{j}})} = e^{-\frac{1}{4}\ln\Pi_{i,j}\frac{(p_{j} - v_{i})}{v_{i}^{3}}} = \sqrt[4]{\Pi_{i,j}[v_{i}^{3}/(p_{j} - v_{i})]}$$
(5.45)

We conclude that

$$\begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \right\}_{\beta}^{S} \approx \frac{(-1)^{(v+v'-p)}}{\sqrt{48\pi kV} \Pi(v_i)} \sqrt{\frac{(p-v)(p-v')}{\overline{v} \overline{v'}}} \\ \times \sqrt[4]{\Pi_{i,j}[v_i^3/(p_j-v_i)]} \sqrt[4]{\frac{(p-v)(p-v')(\overline{p}-v)(\overline{p}-v)(\overline{p}-v'(\overline{p'}-v)(\overline{p'}-v')v')}{(\overline{p}-\overline{v})(\overline{p}-\overline{v'})(\overline{p'}-\overline{v'})(p-\overline{v})(p-\overline{v'})v'}} \\ \times \left\{ \left\{ 2\Pi(v_i)[v+v'-\overline{p}-\overline{p'}] + B[\overline{p}\overline{p'}-vv'] \right\} \cos(\frac{\pi}{4}+\varphi_{k_{\beta}}^{\theta}) + 24V[\overline{p}\overline{p'}-vv'] \sin(\frac{\pi}{4}+\varphi_{k_{\beta}}^{\theta}) \right\} \\ = \frac{(-1)^{(v+v'-p)}}{\sqrt{48\pi kV}} \sqrt[4]{\Pi_{i,j}[(p_j-v_i)v_i]} \\ \times \sqrt{\frac{(p-v)(p-v')}{\overline{v}\overline{v'}}} \sqrt{\frac{(p-v)(p-v')(\overline{p}-v)(\overline{p}-v'(\overline{p'}-v)(\overline{p'}-v')v')v'}{(\overline{p}-\overline{v})(\overline{p'}-\overline{v'})(\overline{p'}-\overline{v'})(\overline{p'}-v')v'}} \\ \times \left\{ \left\{ 2\Pi(v_i)[v+v'-\overline{p}-\overline{p'}] + B[\overline{p}\overline{p'}-vv'] \right\} \cos(\frac{\pi}{4}+\varphi_{k_{\beta}}^{\theta}) + 24V[\overline{p}\overline{p'}-vv'] \sin(\frac{\pi}{4}+\varphi_{k_{\beta}}^{\theta}) \right\}.$$
(5.46)

Note that denominator $\sqrt[4]{\Pi_{i,j}[(p_j - v_i)v_i]}$ has the dimension of an area like $\sqrt{\Pi v_i}$ in the formula (3.18) for approximating the supersymmetric symbol $6j_{\alpha}^S$. The following line with $\sqrt{}$ and $\sqrt[4]{}$ is dimensionless.

6 Results for the supersymmetric limits

Prior to properly present results as similar as possible with standard 6j symbols we will use a formula to shift the angular arguments, namely:

$$a\cos(x) + b\sin(x) = \mathcal{N}\cos(x - \psi), \tag{6.1}$$

where

$$\mathcal{N} = \sqrt{a^2 + b^2}$$
 and $\tan(\psi) = \frac{b}{a}$. (6.2)

Then for each parity $\boldsymbol{\alpha}, \, \boldsymbol{\gamma}, \, \boldsymbol{\beta}$, we get

$$\mathcal{N}_{\alpha} = \sqrt{B^2 + (24V)^2}, \quad \psi_{\alpha} = \arctan\left(\frac{24V}{B}\right),$$
(6.3)

$$\mathcal{N}_{\gamma} = \mathcal{N}_{\alpha}, \quad \psi_{\gamma} = -\psi_{\alpha},$$
(6.4)

$$\mathcal{N}_{\beta} = \sqrt{(2\Pi(v_i)[v+v'-\overline{p}-\overline{p'}])^2 + (24V[\overline{p}\,\overline{p'}-vv'])^2},\tag{6.5}$$

$$\psi_{\beta} = \arctan\left(\frac{24V[\overline{p}\,\overline{p}'-vv']}{2\Pi(v_i)[v+v'-\overline{p}-\overline{p'}]}\right). \tag{6.6}$$

We recall that V is the tetrahedron volume and B which also has the dimension of a volume is given by

$$B = \left(\sum_{\iota=1}^{3} j_{\iota} J_{\iota} \sum_{j=1}^{3} p_{j}\right) + 2(j_{1} j_{2} j_{3} + j_{1} J_{2} J_{3} + j_{2} J_{3} J_{1} + j_{3} J_{1} J_{2}).$$
(6.7)

Parity $\boldsymbol{lpha},:$

Under a rescaling of all its arguments by a large k a supersymmetric $6j_{\alpha}^{S}$ symbol behaves like

$$\begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \right\}_{\alpha}^{S} = \frac{1}{\sqrt{48\pi kV}\sqrt{\Pi v_i}} \Big[\mathcal{N}_{\alpha} \cos\left(\frac{\pi}{4} + \varphi_k^{\theta} - \psi_{\alpha}\right) \Big], \tag{6.8}$$

where the angle φ_k^{θ} is defined by

$$\varphi_k^{\theta} = \sum_{\iota=1}^{\iota=3} (kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}} + (kJ_{\iota} + \frac{1}{2})\theta_{J_{\iota}}.$$
(6.9)

 $\theta_{j_{\iota}}, \theta_{J_{\iota}}$ are the exterior dihedral angles of the tetrahedron corresponding to the edges j_{ι} and J_{ι} respectively.

Parity γ : [If k is even, refer to formula (6.8)]

Under a rescaling of all its arguments by a large $k \pmod{2}$ a supersymmetric $6j_{\gamma}^{S}$ behaves like

$$\begin{cases} kj_1 \ kj_2 \ kj_3 \\ kJ_1 \ kJ_2 \ kJ_3 \end{cases} \Big\}_{\gamma}^{S} = \frac{(-1)^{(1+\sum p_j)}}{\sqrt{48\pi kV}} \Big[\mathcal{N}_{\alpha} \cos\left(\frac{\pi}{4} + \varphi_{k_{\gamma}}^{\theta} + \psi_{\alpha}\right) \Big], \tag{6.10}$$

where

$$\varphi_{k\gamma}^{\theta} = \sum_{\iota=1}^{\iota=3} k(j_{\iota}\theta_{j_{\iota}} + J_{i}\theta_{J_{\iota}}).$$
(6.11)

Parity β : [If k is even, refer to formula (6.8)]

Under a rescaling of all its arguments by a large k (odd) a supersymmetric $6j^S_\beta$ behaves like

$$\begin{cases} kj_{1} \ kj_{2} \ kj_{3} \\ kJ_{1} \ kJ_{2} \ kJ_{3} \end{cases} \stackrel{S}{=} \frac{(-1)^{(v+v'-p)}}{\sqrt{48\pi kV} \sqrt[4]{\Pi_{i,j}[(p_{j}-v_{i})v_{i}]}} \sqrt{\frac{(p-v)(p-v')}{\overline{v} \overline{v}'}} \\ \times \sqrt[4]{\frac{(p-v)(p-v')(\overline{p}-v)(\overline{p}-v'(\overline{p}'-v)(\overline{p}'-v')v'}{(\overline{p}-v)(\overline{p}'-\overline{v}')(\overline{p}'-\overline{v}')(\overline{p}-\overline{v}')v'}} \left[\mathcal{N}_{\beta} \cos\left(\frac{\pi}{4} + \varphi^{\theta}_{k_{\beta}} - \psi_{\beta}\right) \right], \tag{6.12}$$

where a new angle depending on $\theta_{j^*_\beta}$ is defined as

$$\varphi_{k_{\beta}}^{\theta} = \sum_{\iota=1}^{\iota=3} \left[(kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}} + (kJ_{\iota} + \frac{1}{2})\theta_{J_{\iota}} \right] + \frac{1}{2}\theta_{j_{\beta}^{*}}.$$
(6.13)

This means that only one term among the six $(kj_{\iota}+\frac{1}{2})\theta_{j_{\iota}}, (kJ_{\iota}+\frac{1}{2})\theta_{J_{\iota}}$ transforms into $(kj_{j_{\beta}^{*}}+1)\theta_{j_{\beta}^{*}}$. Denominator in 4 has the dimension of an area like $\sqrt{\Pi v_{i}}$.

7 Conclusion

Compared to a standard 6j symbol [SU(2)] a major difference with the supersymmetric $6j_{\pi}^{S}$ [OSP(1|2)] lies in a slower decay as a function of the scaling parameter k. Indeed it becomes $\sqrt{\frac{1}{k}}$ instead of $\sqrt{\frac{1}{k^3}}$. The standard term in *cos* contains always angular arguments depending on the six tetrahedral angles $\theta_{j_{\iota}}, \theta_{J_{\iota}}$. However all the angles are shifted from their standard values by an angle different according to each parity π .

For parity α the usual expressions like $(kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}}$ are unchanged. For parity γ (and k odd) the terms in $\frac{1}{2}$ vanish so that it remains $(kj_{\iota})\theta_{j_{\iota}}$ and so on. For parity β (and k odd) the angular dependence takes a form where a special angle $\theta_{j_{\beta}^{*}}$ modifies the standard formula into $(kj_{j_{\beta}^{*}} + 1)\theta_{j_{\beta}^{*}} + \sum_{\iota} (kj_{\iota} + \frac{1}{2})\theta_{j_{\iota}} + (kj_{\iota} + \frac{1}{2})\theta_{J_{\iota}}|_{j_{\iota},J_{\iota} \neq j_{\sigma}^{*}}$.

If k is even all supersymmetric $6j_{\pi}^{S}$ have the same asymptotic behaviour, *ie* that of parity α . Pertinent interpretations are clearly within the Quantum Supergravity framework.

Appendix: Expression of supersymmetric prefactors

$$\mathbf{R}_{\alpha}^{S} = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} (p_{j} - v_{i})!}{\prod_{i=1}^{i=4} (v_{i})!},$$
(A.1)

$$\mathbf{R}_{\gamma}^{S} = \frac{\prod_{j=1}^{j=3} \prod_{i=1}^{i=4} (p_{j} - v_{i} - \frac{1}{2})!}{\prod_{i=1}^{i=4} (v_{i} + \frac{1}{2})!},$$
(A.2)

$$R_{\beta}^{S} = (p-v)!(p-v')!(p-\overline{v}-\frac{1}{2})!(p-\overline{v}'-\frac{1}{2})! \times (\overline{p}-\overline{v})!(\overline{p}-\overline{v}')!(\overline{p}-v-\frac{1}{2})!(\overline{p}-v'-\frac{1}{2})! \times (\overline{p}'-\overline{v})!(\overline{p}'-\overline{v}')!(\overline{p}'-v-\frac{1}{2})!(\overline{p}'-v'-\frac{1}{2})! \times (v!v'!)^{-1}((\overline{v}+\frac{1}{2})!(\overline{v}'+\frac{1}{2})!)^{-1}$$
(A.3)

References

- [1] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004).
- [2] V. Aquilanti, D. Marinelli and A Marzuoli, "Hamiltonian dynamics of a quantum of space: hidden symmetries and spectrum of the volume operator, and discrete orthogonal polynomials", Journal of Physics A: Mathematical and Theoretical. 46(17), 175001-175303 (2013).
- [3] G. Ponzano and T. Regge, *Semiclassical limit of Racah coefficients*, (Spectroscopic and group theoretical methods in physics, North-Holland, New York, 1968).
- [4] E.R. Livine and R. Oeckl, "Three-dimensional Quantum SuperGravity and Supersymmetric Spin Foam Models", Adv. Theor Math. Phys. 7(6), 951-1001(2004).
- [5] A. Pais and V. Rittenberg, "Semisimple graded Lie algebras", J. Math. Phys. 16(10), 2062-2073 (1975).
- [6] M. Scheunert, W. Nahm and V. Rittenberg, "Irreducible representations of the osp(2,1) and spl(2,1) graded Lie algebras", J. Math. Phys. 18(1), 155-162 (1977).
- [7] F. A. Berezin, V. N. Tolstoy, "The group with Grassmann structure UOSP(1.2)", 78(3), 409-428 (1981).

- [8] M. Daumens, P. Minnaert, M. Mozrzymas and S. Toshev, "The super-rotation Racah-Wigner calculus revisited", J. Math. Phys. 34(6), 2475-2507 (1993).
- [9] L. Bréhamet, "Analytical complements to the parity-independent Racah-Wigner calculus for the superalgebra $\mathfrak{osp}(1|2)$ Part I", Il Nuovo Cimento 121B(3), 241-274 (2006).
- [10] R. Gurau, "The Ponzano-Regge Asymptotic of the 6j Symbol: An Elementary Proof", Ann. Henri Poincaré 9(7), 1413-1420 (2008).
- [11] L. Bréhamet, "Regge Symmetry of 6-j or super 6-jS Symbols: a Re-Analysis with Partition Properties", Pioneer Journal of Mathematical Physics and its Applications **6**(1), 1-20 (2014). arXiv:1505.05408v1 [math-ph]