

RANDOM TESSELLATIONS, RESTRICTED ISOMETRIC EMBEDDINGS, AND ONE BIT SENSING

DMITRIY BILYK AND MICHAEL T. LACEY

ABSTRACT. We obtain improved bounds for one bit sensing. For instance, let K_s denote the set of s -sparse unit vectors in the sphere \mathbb{S}^n in dimension $n+1$ with sparsity parameter $0 < s < n+1$ and assume that $0 < \delta < 1$. We show that for $m \gtrsim \delta^{-2}s \log \frac{n}{s}$, the one-bit map

$$x \mapsto [\operatorname{sgn}\langle x, g_j \rangle]_{j=1}^m,$$

where g_j are iid gaussian vectors on \mathbb{R}^{n+1} , with high probability has δ -RIP from K_s into the m -dimensional Hamming cube. These bounds match the bounds for the *linear* δ -RIP given by $x \mapsto \frac{1}{m} [\langle x, g_j \rangle]_{j=1}^m$, from the sparse vectors in \mathbb{R}^n into ℓ^1 . In other words, the one bit and linear RIPs are equally effective. There are corresponding improvements for other one-bit properties, such as the sign-product RIP property.

CONTENTS

1. Introduction	1
2. Small Cells: Proof of Theorem 1.5	8
3. δ -RIP into the Hamming Cube: Proof of Theorem 1.14	12
4. Sign-Product Embedding Property	14
5. Background on Stochastic Processes	17
References	21

1. INTRODUCTION

We study questions associated with one bit sensing and dimension reduction. We shall work with subsets K of the unit sphere \mathbb{S}^n in dimension $n+1$. A natural class of such subsets, which arises in compressed sensing, are the *s-sparse vectors*

$$(1.1) \quad K_s := \{x = (x_1, \dots, x_{n+1}) \in \mathbb{S}^n : \#\{j : x_j \neq 0\} \leq s\},$$

where $0 < s < n+1$

Research supported in part by.

Research supported in part by grant NSF-DMS 1265570.

We denote by $d(x, y)$ the geodesic distance on \mathbb{S}^n , normalized so that the distance between antipodal points is one, that is

$$d(x, y) = \frac{\cos^{-1}(x \cdot y)}{\pi}.$$

For iid uniform samples θ_j , $1 \leq j \leq m$, we will consider the linear map

$$Ax = \frac{1}{m} \left[\langle \theta_j, x \rangle \right]_{1 \leq j \leq m}$$

from \mathbb{R}^{n+1} to \mathbb{R}^m , where \mathbb{R}^{n+1} is equipped with the usual Euclidean ℓ^2 metric, while we use the ℓ^1 metric on \mathbb{R}^m .

More generally, we will consider the *sign-linear map*

$$(1.2) \quad \text{sgn}(Ax) := \left[\text{sgn}(\langle \theta_j, x \rangle) \right]_{1 \leq j \leq m}$$

which is a map from \mathbb{R}^{n+1} into the Hamming cube $\{-1, 1\}^m$ with the metric

$$d_H(x, y) = \frac{1}{2m} \sum_{i=1}^m |x_i - y_i| = \frac{|\{1 \leq j \leq m : x_j \neq y_j\}|}{m},$$

i.e. the fraction of coordinates in which $x, y \in \{-1, +1\}^m$ differ. In other words, a sign-linear map keeps only one bit of information from each linear measurement. This is an example of one-bit sensing, a subject initiated by Boufounos and Baraniuk [5], and further studied by several authors [9, 10, 13, 15].

The motivation for the one-bit map is that it is a canonical non-linearity in measurement, and so its study opens the door for a broader non-linear theory in compressed sensing. It has proven to be useful in other settings, see the applications of one-bit sensing in [3, 9]. Furthermore, this topic has deep connections to the Dvoretzky Theorem, as explained in [15], as well as important relations to geometric discrepancy theory, explored in a companion paper of the authors [4].

The basic question, with different meanings and interpretations being attached to it, is how well these maps preserve the structure of K . Generally, it is also of interest to study this question when the iid samples θ_j are, for instance, standard gaussians on \mathbb{R}^{n+1} . However, in the context of the one-bit map, the magnitudes of both x and θ_j are lost, so that we concern ourselves mostly with the θ_j being uniform on the sphere \mathbb{S}^n .

The first crude property begins with the observation that the map $\text{sgn}(\langle \theta_j, x \rangle)$ divides \mathbb{S}^n into two hemispheres, bounded by the hyperplane θ_j^\perp . Therefore, a number of one bit observations induce a tessellation of a subset $K \subset \mathbb{S}^n$ into cells bounded by the hyperplanes. Two points are in the same cell if and only if their sequence of one bit measurements are the same. See Figure 1.

Definition 1.3. We say that the tessellation of K induced by $\{\theta_j\}$ is has δ -small cells if for all $x, y \in K$, if $\text{sgn}\langle x, \theta_j \rangle = \text{sgn}\langle y, \theta_j \rangle$, for all j , then $d(x, y) < \delta$. Equivalently, all the cells of the induced tessellation have maximal diameter at most δ .




FIGURE 1. A region K , bounded by the solid line, with an induced tessellation from hyperplanes represented by the dashed lines.

One can then ask how small the cells are in the induced tessellation of K . The relevant characteristic of K that will control this property is the *metric entropy* $N(K, \delta)$, i.e. the least number of d -balls of radius $\delta > 0$ needed to cover K .

We shall repeatedly make use of the following observation. Let $M(K, \delta)$ be the maximal cardinality of a δ -separated subset $K_0 \subset K$ (i.e. any $x, y \in K_0$ with $x \neq y$ satisfy $d(x, y) > \delta$). Then the following obvious inequalities hold.

$$(1.4) \quad M(K, 2\delta) \leq N(K, \delta) \leq M(K, \delta).$$

The next result should be compared to [13, Thm 4.2], where the bound on m is substantially larger, both in terms of the power of δ , as well as in the fact that it involves the gaussian mean width, see (1.9), instead of the smaller metric entropy of K .

Theorem 1.5. [δ -Small Cells] *There are constants $0 < c < 1 < C$ so that for all integers $n \geq 2$, for any $K \subset \mathbb{S}^n$, and any $0 < \delta < 1$, if $m > C\delta^{-1} \log N(K, c\delta)$, then with probability at least $1 - [2N(K, c\delta)]^{-2}$ the following holds:*

- (1) *The random vectors $\{\theta_j\}_{j=1}^m$ induce a tessellation of K with δ -small cells.*
- (2) *For any pair of points $x, y \in K$, with $|x - y| > \delta$, there are at least $c\delta m$ choices of j such that*

$$\langle x, \theta_j \rangle < -c\delta < c\delta < \langle y, \theta_j \rangle.$$

- (3) *And, for iid gaussians g_j , are at least $c\delta m$ choice of j with*

$$\langle x, g_j \rangle < -c\delta \sqrt{n} < c\delta \sqrt{n} < \langle y, g_j \rangle.$$

Our proof is a modification of the standard ‘occupation time problem’, which states that it takes about $n \log n$ independent uniform $(0,1)$ observations to occupy all of the intervals $[j/n, (j+1)/n)$, for $0 \leq j < n$. The heuristic here is the following: *the small cells property is governed by metric entropy at scale δ .*

This heuristic is quite useful in the next corollary, which improves the result of Plan and Vershynin [13, Thm 2.1], which proves the statement below with δ^{-3} replaced by δ^{-5} . This fact was basic to the main results of that paper concerning logistic regression in a compressed one-bit setting.

Corollary 1.6. *Let $0 < s < n + 1$ and $0 < \delta < 1$ be given. Define a convex variant of the sparse vectors to be*

$$(1.7) \quad K_{n,s} := \{x \in \mathbb{R}^{n+1} : \|x\|_2 = 1, \|x\|_1 \leq s\}.$$

If $m \gtrsim \delta^{-3}s \log_+ \frac{n}{s}$, then the random vectors $\{\theta_j\}_{j=1}^m$ induce a tessellation of $K_{n,s}$ with δ -small cells.

We now turn to questions that are *multi-scale in nature*, namely the *restricted isometry property*, which will be abbreviated to RIP [6].

Definition 1.8. Let (X, d_X) and (Y, d_Y) be metric spaces. We say that map $\varphi : X \mapsto Y$ has the δ -RIP if

$$|d_X(x, x') - d_Y(\varphi(x), \varphi(x'))| < \delta, \quad x, x' \in X.$$

In many situations, the dimension of Y is much smaller than that of X , i.e. restricted isometries are a form of dimension reduction. In this direction, considering *linear* isometries, Klartag and Mendelson [11] have proved

Theorem A. [*δ -RIP into ℓ^1*] *There is a constant $C > 0$ so that for all integers n , $K \subset \mathbb{S}^n$, and $0 < \delta < 1$, if $m \geq C\delta^{-2}\omega(K)$, with probability at least $1 - \exp(-C\delta^{-2}\omega(K)^2)$, the linear map $\frac{1}{m}A : \ell_n^2 \mapsto \ell_m^1$, where the rows of A are iid gaussian vectors, has the δ -RIP.*

Above, $\omega(K)$ is the gaussian mean width, defined by

$$(1.9) \quad \omega(K) = \mathbb{E} \sup_{x,y \in K} \langle x - y, \gamma \rangle,$$

where γ is a standard normal on \mathbb{R}^n . The motivation here comes from standard examples. Observe that the gaussian mean width of the sphere is

$$\omega(\mathbb{S}^n) = \mathbb{E}|\gamma| \simeq \sqrt{n}.$$

Moreover, for the s -sparse vectors K_s as in (1.1), one has $\omega(K_s) \simeq \sqrt{s \log \frac{n}{s}}$. That is, in many cases $\omega(K)^2$ represents an intrinsic notion of dimension.

To compare the bounds in the two different theorems, first observe that the metric for the gaussian process $\gamma_x := \langle x, \gamma \rangle$ satisfies

$$(1.10) \quad \|\gamma_x - \gamma_y\|_2 = \left[\mathbb{E} \langle x - y, \gamma \rangle^2 \right]^{1/2} \simeq d(x, y).$$

Second, use Theorem 5.1, Sudakov's lower bound for the supremum of gaussian processes, to see that

$$\delta^{-1} \log N(K, \delta) = \delta^{-3} [\delta^2 \log N(K, \delta)] \lesssim \delta^{-3} \omega(K)^2.$$

See Schechtman [17] for a detailed discussion of the theorem above, its relation to Dvoretzky Theorem, and remarks about optimal bounds.

We turn to the subject of the one bit RIP. Namely, the construction of δ -RIPs from $K \subset \mathbb{S}^n$ into a Hamming cube of dimension m . Plan and Vershynin [11] proved that




FIGURE 2. An illustration of the wedge $W_{x,y} = H_x \Delta H_y$.

if $m \gtrsim \delta^{-6} \omega(K)^2$, then the one bit map (1.2) satisfies δ -RIP from the sphere into the Hamming cube, with high probability. They also suggested that $\omega(K)^2$ would also be the correct measure of the size of K for δ -isometries into the Hamming cube. We cannot verify this, but can prove an estimate of the form $m \gtrsim \delta^{-2} H(K)^2$, where $H(K)$ is a different measure of the intrinsic dimension of K . In the case of sparse vectors, we will see that $H(K_s) \approx \omega(K_s) \approx \sqrt{s \log \frac{n}{s}}$.

To set the stage for this new measure, define $H_x := \{\theta \in \mathbb{S}^n : \langle \theta, x \rangle \geq 0\}$ to be the positive hemisphere relative to x . Observe that a hyperplane θ^\perp point θ separates two points $x, y \in \mathbb{S}^n$ if and only if $\theta \in W_{x,y} := H_x \Delta H_y$ (W stands for 'wedge'). The symmetric difference between two hemispheres, is related to the geodesic distance on the sphere through the essential equality $\mathbb{P}(W_{x,y}) = d(x, y)$, which is a simple instance of the Crofton formula. See Figure 2 and [18, p. 36-40].

Let us use randomly selected points $\{\theta_1, \dots, \theta_m\} \subset \mathbb{S}^n$ to define the one bit map given by (1.2), namely

$$\varphi(x) = \{\text{sgn}(\langle x, \theta_j \rangle) : 1 \leq j \leq m\} \in \{-1, +1\}^m.$$

Restricting this map to $K \subset \mathbb{S}^n$, the δ -RIP property becomes

$$\sup_{x,y \in K} \left| \frac{1}{m} \sum_{j=1}^m \mathbf{1}_{W_{x,y}}(\theta_j) - \mathbb{P}(W_{x,y}) \right| \leq \delta.$$

That is, we need to bound, in non-asymptotic fashion, a standard empirical process over the class of wedges $W_{x,y}$ defined by K . Essential to such an endeavor is to understand the asymptotic behavior of the associated empirical process.

We state the following theorem without proof. It is an asymptotic result, while our emphasis is on non-asymptotic ones. However, it points to a distinguished role played by the gaussian process that appears in the conclusion. We rely upon the well-known fact that a gaussian process is uniquely determined by its mean and covariance structure.

Theorem 1.11. *As $m \rightarrow \infty$, the empirical process indexed by hemispheres converges weakly*

$$\frac{1}{\sqrt{m}} \sum_{j=1}^m \mathbf{1}_{H_x}(\theta_j) - \frac{1}{2} \xrightarrow{d} G_x$$

where x ranges over \mathbb{S}^n , and G_x is the mean-zero gaussian process with

$$(1.12) \quad \mathbb{E}G_x^2 = \frac{1}{4}, \quad (\mathbb{E}(G_x - G_y)^2)^{1/2} = \mathbb{P}(W_{x,y})^{1/2} = \sqrt{d(x,y)}.$$

We call the gaussian process above the *hemisphere gaussian process*. It is a key innovation of this paper. Particularly relevant in comparing our results with those stated in terms of gaussian mean width, note that the metric for γ_x (1.10) and that of G_x are related (essentially) through a square root.

We then define the *hemisphere mean width* of $K \subset \mathbb{S}^n$ by

$$(1.13) \quad H(K) := \mathbb{E} \sup_{x,y \in K} (G_x - G_y)$$

It always holds that $\omega(K) \lesssim H(K) \lesssim \sqrt{n}$, and for the s -sparse vectors K_s , $\omega(K) \simeq H(K) = \sqrt{s \log_+ \frac{n}{s}}$. We comment more on this measure in §3 below.

The next result concerns the bounds for the one-bit δ -RIP. It has two parts, one for general $K \subset \mathbb{S}^n$, and the other for the unit sparse vectors defined in (1.1).

Theorem 1.14. [δ -RIP into Hamming Cube] *There is a constant $C > 0$ so that for all dimensions $n \in \mathbb{N}$, subsets $K \subset \mathbb{S}^n$, and $0 < \delta < 1$, the following two conclusions hold.*

General K : *There exists a δ -RIP map from (K, d) to the Hamming cube $(\{-1, 1\}^m, d_H)$, provided that $m \geq C\delta^{-2}H(K)^2$.*

Sparse vectors: *For $1 \leq s \leq n + 1$, with probability $1 - \exp(-s \log_+ \frac{n}{s})$, the map $x \mapsto \text{sgn}(Ax)$ has δ -RIP from (K_s, d) to the Hamming cube $(\{-1, 1\}^m, d_H)$, provided that $m \geq C\delta^{-2}s \log_+ \frac{n}{s}$.*

Notice that for general K , we do not assert that $x \mapsto \text{sgn}(Ax)$ is the isometry, whereas Plan and Vershynin [15, Thm 1.2] show that it is, but with $m \gtrsim \delta^{-6}\omega(K)^2$. We comment more on this in §3. But, note that in the sparse vector case, we only require m to be as big as in the linear δ -RIP. That is we have this surprising conclusion.

For sparse vectors the one-bit map is just as effective as the linear map.

We note that [9] very nearly proves this result for sparse vectors, missing by a factor of $\log 1/\delta$, and using a proof that is more involved than ours. Also, the Jacques [8] considers quantized maps from sets $K \subset \mathbb{R}^n$ into \mathbb{Z}^m , using gaussian mean width. It seems plausible that the hemisphere process is relevant to that paper as well.

This corollary, in which K is a finite set, is immediate, and is a one bit analog of the Johnson-Lindenstrauss lemma.

Corollary 1.15. *If $K \subset \mathbb{S}^n$ is finite, then there is a δ -isometry from (K, d) into the Hamming cube $(\{-1, 1\}^m, d_H)$, where $m \lesssim \delta^{-2} \log|K|$.*

Recall that the Johnson-Lindenstrauss lemma states that a finite set $K \subset \mathbb{R}^n$ has a linear Lipschitz embedding into \mathbb{R}^m , where $m \gtrsim \delta^{-2} \log|K|$. Namely, for linear $A : \mathbb{R}^n \mapsto \mathbb{R}^m$,

$$\left| \|Ax - Ax'\|_2 - \|x - x'\|_2 \right| \leq \delta \|x - x'\|_2.$$

Here, A is $\frac{1}{m}$ times a standard $m \times n$ gaussian matrix, the inequality holding with high probability. Embeddings of ℓ^2 into ℓ^1 are of significant interest ([7] and references therein), from the perspective of the best possible bounds, as well as implementations with few random inputs.

Another relevant property is *sign-product RIP* property. This we specialize to the case of sparse vectors, and also add modest restrictions on s and δ in order get a value of m that matches those of the other Theorems. Namely, we require a weak lower bound on δ , as a function of s/n .

Theorem 1.16. *There is a constant C so that for all integers $0 < s \leq n$, and $0 < \delta < 1$ such that*

$$(1.17) \quad (s/10n)^4 < \delta$$

with probability at least $1 - (\frac{s}{2n})^{2s}$, for all $m \simeq C\delta^{-2}s \log_+ \frac{n}{s}$, there holds

$$(1.18) \quad \sup_{x, y \in K_s} \frac{1}{m} \left| \sum_{j=1}^m \operatorname{sgn}(\langle x, g_j \rangle) \langle y, g_j \rangle - \lambda \langle x, y \rangle \right| \leq \delta$$

where the g_j are iid standard gaussians on \mathbb{R}^{n+1} , and $\lambda = \sqrt{\frac{2}{\pi}}$.

This RIP property arises from [9], and a quantitative estimate $m \gtrsim \delta^{-6} \omega(K)^2$ is proved in [14, Prop. 4.3] for general K . Furthermore, this latter bound is heavily used in [3, §4] in the sparse vector case. Above we have seemingly optimal lower bounds on m , subject to mild restrictions on δ .

One can ask what is the best value of $m = m(K, \delta)$ that can achieve the bounds in these different estimates, from small cells to the restricted isometry properties. The estimates of Plan and Vershynin [15] were in terms of the gaussian mean width. Using Sudakov's lower bound, for the supremum of gaussian processes, Theorem 5.1, note that

$$(1.19) \quad \sqrt{\log N(K, \delta)} \lesssim \begin{cases} \delta^{-1} \omega(K) \\ \delta^{-1/2} H(K) \end{cases}$$

with the difference in the right hand side being a consequence in the difference of a square root in the metrics of the two gaussian processes. Indeed, $N(K, d_G, \delta) = N(K, \delta^2)$, where

d_G is the metric of the hemisphere process. Therefore,

$$\delta^{-1} \log N(K, \delta) \lesssim \begin{cases} \delta^{-3} \omega(K)^2 \\ \delta^{-2} H(K)^2 \end{cases}$$

The left hand side is the bound for the δ -small cell property. The lower part of the right hand side is the bound required for the δ -RIP into the Hamming cube. It would appear that δ^{-3} is the smallest multiple of $\omega(K)^2$ that could appear in such theorems, and, for instance, the bounds of Plan and Vershynin [15] were of the form $\delta^{-6} \omega(K)^2$.

The small cells Theorem 1.5, proved in § 2, will be seen as a consequence of a standard occupation time calculation. The RIP Theorem 1.14 is a calculation involving empirical processes, after reducing a general K to a finite approximating set. Specializing K to sparse vectors leads naturally to the notion of VC dimension. This is detailed in § 3, with the sign-product RIP Theorem 1.16 proved in § 4. Powerful inequalities for empirical processes for VC sets simplify the analysis of these theorems. The background information on empirical processes and the properties we need are collected in § 5.

Acknowledgment. This work was completed as part of a the program in High Dimensional Approximation, Fall 2014, at ICERM, at Brown University. We thank Simon Foucart for asking us about the sign-product RIP.

2. SMALL CELLS: PROOF OF THEOREM 1.5

We introduce the notion of a hyperplane *transversely separating* vectors $x, y \in \mathbb{S}^n$. Assuming x and y are not antipodal, there is a unique geodesic τ from x to y . Any hyperplane H that separates x and y must intersect τ . We further say that H *transversely separates* x and y if (a) the angle of intersection between τ and H is at least $\pi/4$, and (b) that the distance of the point of intersection $\tau \cap H$ to both x and y be at least $d(x, y)/4$. The heuristic motivation for this definition is that if a hyperplane transversely separates x and y , it also has to separate point lying close to x and y , which allows one to pass to a finite subset. Let

$$\widetilde{W}_{x,y} := \{\theta \in W_{x,y} : \theta^\perp \text{ transversely separates } x \text{ and } y\}.$$

Now, for a randomly selected $\theta \in W_{x,y}$, the angle of intersection between θ^\perp and τ is uniformly distributed on the circle, as is the point of intersection $\theta^\perp \cap \tau$ on τ . Moreover, these two quantities are statistically independent. From this, it follows that

$$\mathbb{P}(\widetilde{W}_{x,y}) = \frac{1}{4} d(x, y).$$

The main line of argument begins with selection of $K_0 \subset K$, a maximal subset such that each pair of distinct points $x, y \in K_0$ satisfies $d(x, y) \geq \delta/64$. Then, as is well known, see (1.4), $|K_0| \leq N(K, \delta/128)$.

Observe the following: if \mathcal{H} is a collection of hyperplanes for which each pair of distinct elements of K_0 is transversely separated by a hyperplane in \mathcal{H} , then \mathcal{H} induces a tessellation




FIGURE 3. The points x and y are transversely separated by θ^\perp , but not $\tilde{\theta}^\perp$.

with δ -small cells on K . Indeed, for two points $x, y \in K$, separated by δ , let τ be the geodesic between x and y , and then, take $x', y' \in K_0$ to be the points closest to x , and y , respectively. So, $d(x, x'), d(y, y') \leq \frac{1}{64}\delta$, and $d(x', y') \geq \frac{31}{32}\delta$. Let τ' be the geodesic between x' and y' .

Now, x' and y' are transversely separated by a hyperplane $H \in \mathcal{H}$, by assumption. Then, (i) the point $z' = H \cap \tau'$ is at distance at most $\delta/8$ from τ , (ii) z' is at least at distance $\frac{1}{5}d(x, y)$ from both x and y , and hence (iii) H separates x and y , and moreover, the point $z = \tau \cap H$ satisfies $d(z, x) \geq \frac{1}{20}d(x, y)$, and similarly for y .

Indeed, to see (i) the lengths of τ and τ' very close.

$$|d(x, y) - d(x', y')| \leq \frac{1}{32}\delta.$$

Let τ'' be the geodesic obtained by a rigid motion of τ' so that τ'' has x as an endpoint. Then, parameterizing τ and τ'' by arc length s , and identifying $\tau(0) = \tau''(0) = x$, the quantity $d(\tau(s), \tau''(s))$ is monotone increasing. But, at $s = d(x, y)$, we will have

$$\begin{aligned} \text{dist}(z', \tau) &\leq \frac{1}{64}\delta + d(\tau(s), \tau''(s)) \\ &= \frac{1}{64}\delta + d(y, \tau''(s)) \leq \frac{1}{64}\delta + d(y, y') + d(y', \tau''(s)) \\ &\leq \frac{1}{32}\delta + d(y', \tau''(d(x', y'))) + |d(x', y') - d(x, y)| \leq \frac{1}{8}\delta. \end{aligned}$$

To see (ii), just use the triangle inequality to get a better result.

$$d(z', x) \geq d(z', x') - d(x', x) \geq \frac{1}{4}d(x', y') - \frac{1}{64}\delta \geq \frac{29}{128}d(x, y).$$

And, (iii) would be obvious on the plane. We are however on the sphere, but with spherical triangles of small diameter. The spherical corrections are small, so the result will follow.

We can now argue for the first conclusion of the Theorem, concerning small cells. Take $\mathcal{H} = \{\theta_\ell^\perp : 1 \leq \ell \leq m\}$ to be iid uniform random hyperplanes, where $m \geq C\delta^{-1} \log N(K, \delta/128)$. For a pair of points $x, y \in K_0$, $x \neq y$, the chance that no θ_ℓ transversely separates x and y is, for some fixed $0 < c < 1$,

$$\begin{aligned} \mathbb{P}(\theta_\ell \notin \widetilde{W}_{x,y}, 1 \leq \ell \leq m) &\leq (1 - c\delta)^m \\ &\leq \exp(-c'\delta m) \leq (2 \cdot N(K, \delta/128))^{-4}, \end{aligned}$$

with appropriate choice of C . Now, there are at most $N(K, \delta/128)^2$ pairs of points $x, y \in K_0$, $x \neq y$, hence, by the union bound, the probability that K_0 is not transversely

separated by \mathcal{H} is at most $(2 \cdot N(K, \delta/128))^{-2}$. So the proof of the first conclusion of Theorem 1.5 is complete.

For the second conclusion of the Theorem, begin by observing that if θ^\perp transversely separates x and y , then, we have necessarily

$$\langle x, \theta \rangle < -c_0 d(x, y) < c_0 d(x, y) < \langle y, \theta \rangle,$$

where above, we can take $c_0 = \frac{1}{10}$. Above, we assume as we may that $\langle x, \theta \rangle < 0$.

We will then prove the conclusion of the Theorem for K_0 . Namely if $m > C\delta^{-1} \log N(K, c\delta)$, with probability at least $1 - [2N(K, c\delta)]^{-2}$, for any pair of points $x, y \in K$, with $|x - y| > \delta$, there are at least $c'\delta m$ choices of j such that

$$(2.1) \quad \langle x, \theta_j \rangle < -c_0 \delta < c_0 \delta < \langle y, \theta_j \rangle.$$

We can then draw the same conclusion for all of K , provided we replace c_0 above by $\frac{1}{20}$, by property (ii) above.

Now, $p_{x,y} = \mathbb{P}(\theta \in \widetilde{W}_{x,y}) > \frac{1}{256}\delta$, so for sufficiently small c ,

$$\begin{aligned} \mathbb{P}\left(\sum_{j=1}^m \mathbf{1}_{\theta_j \in \widetilde{W}_{x,y}} < c\delta m\right) &\leq \mathbb{P}\left(\sum_{j=1}^m \mathbf{1}_{\theta_j \in \widetilde{W}_{x,y}} - p_{x,y} < -\frac{c}{2}\delta m\right) \\ &\lesssim \exp\left(-\frac{c^2\delta^2 m^2}{c'\delta m}\right) \lesssim \exp(-C \log N(K, \delta/64)) \end{aligned}$$

by choice of m . We can make C large, so that a simple union bound shows that with probability at least $1 - (2N(K, \delta/64))^{-4}$, for each pair of $x, y \in K_0$, $x \neq y$, there are at least $c\delta m$ choices of j for which θ_j satisfies (2.1). The proof is complete.

For the third conclusion of the Theorem, we want to make the same conclusion as in the the second part, but for gaussian r.v.s. But, we know the result for θ_j chosen at random on the sphere. And, when making a gaussian observation g on \mathbb{R}^{n+1} , we say that it is *moderate* if $\frac{\sqrt{n+1}}{2} \leq |g| \leq 2\sqrt{n+1}$. Note that

$$\mathbb{P}\left(\frac{\sqrt{n+1}}{2} \leq |g| \leq 2\sqrt{n+1}\right) = \mathbb{P}\left(-\frac{3(n+1)}{4} \leq \sum_{\ell=1}^{n+1} g_\ell^2 - 1 \leq 3(n+1)\right)$$

where the g_ℓ are one dimensional standard gaussian r.v.s. It follows that the probability of g being moderate tends to one, as n tends to ∞ .

Now, for iid gaussians $\{g_j\}$ taking values in \mathbb{R}^{n+1} , set $\theta_j = g_j/|g_j|$. If θ_j satisfies (2.1), and g_j is moderate, then $g_j/\sqrt{n+1}$ satisfies (2.1), with different constants. The event that g_j being moderate, and the distribution of θ_j are statistically independent, hence we can condition on the $\{g_j\}$ being moderate, and apply the second conclusion, to deduce our result.

2.1. **Proof of Corollary 1.6.** The next lemma improves upon [13, Lemma 3.4] and, together with Theorem 1.5, clearly proves the Corollary.

Lemma 2.2. *The set $K_{n,s}$ defined in (1.7) satisfies the metric entropy bound*

$$\delta^2 \log N(K_{n,s}, \delta) \lesssim s \log_+ n/s.$$

Proof. To estimate this quantity, we use Sudakov's bound in the form of the upper estimate in (1.19). Thus, we have

$$\delta^2 \log N(K_{n,s}, \delta) \lesssim \omega(K_{n,s})^2.$$

Therefore, it suffices to show that $\omega(K_{n,s})^2 \lesssim s \log_+ \frac{n}{s}$, which is a well-known fact.

This follows from: (A) the convexity result $K_{n,s} \subset 2 \operatorname{convex}(K_s)$, where the latter is the convex hull of K_s ; (B) the further convexity estimate $\omega(\operatorname{convex}(K)) \lesssim \omega(K)$; and (C) $\omega(K_s)^2 \lesssim s \log_+ \frac{n}{s}$.

These three points are as follows. The point (A) is seen by a common technique in compressed sensing. Given $x \in K_{n,s}$, write it as $x = \sum_j x_j$, where x_1 is x restricted to the s -largest coefficients, x_2 has the next s -largest coefficients, and so on. It suffices to show that

$$\sum_j \|x_j\|_2 \leq 2.$$

Now, $\|x_1\|_2 \leq 1$. And, for $j > 1$, each coordinate of x_j is dominated by the least coordinate of x_j , which is at most $\|x_{j-1}\|_1/s$. Hence $\|x_j\|_2 \leq \|x_{j-1}\|_1/\sqrt{s}$. But, we have a bound on $\|x\|_1$, hence

$$\begin{aligned} \sum_{j=1}^{\infty} \|x_j\|_2 &\leq \|x_1\|_2 + \sum_{j=2}^{\infty} \|x_j\|_2 \\ &\leq 1 + \frac{1}{\sqrt{s}} \sum_{j=1}^{\infty} \|x_j\|_1 \leq 2. \end{aligned}$$

Point (B) is a deep implication of Talagrand's majorizing measure theorem. And (C) is a well-known fact, see for instance [14, Lemma 2.3]. □

Remark 2.3. We have chosen the proof above for convenience. The gaussian mean width is used, namely the upper half of the Sudakov estimate (1.19), so that we could appeal to Talagrand's convexity inequality. (With the hemisphere process, the convexity argument would be more complicated.) Thus, the power on δ we get is δ^{-3} , which as (1.19) suggests, is optimal for this strategy. Potentially, there is an additional improvement in the power of δ , but we do not pursue it here.

3. δ -RIP INTO THE HAMMING CUBE: PROOF OF THEOREM 1.14

3.1. **The Case of General $K \subset \mathbb{S}^n$.** The distinction between the case of general K and sparse K_s is that in the general case, we avoid making an entirely specific choice of RIP. Let K_0 be a maximal cardinality subset of K so that for all $x \neq y \in K_0$, one has $d(x, y) > \delta/4$. Of course $|K_0| \leq N(K, \delta/8)$, see (1.4). Letting $\pi_0 : K \mapsto K_0$ be the map sending $x \in K$ to the element of K_0 closest to x , one then has $d(x, \pi_0 x) \leq \delta/4$. That is $\pi_0 : K \mapsto K_0$ has $\delta/2$ -RIP.

We construct a one-bit map $\varphi_0 : K_0 \mapsto \{-1, 1\}^m$, which is also a $\delta/2$ -RIP map, i.e.

$$\sup_{x, y \in K_0} \left| d_H(\varphi_0(x), \varphi_0(y)) - d(x, y) \right| \leq \delta/2.$$

Observe that if we extend φ_0 to K by the formula $\varphi(x) := \varphi_0(\pi_{s_0} x)$, we will have then proved Theorem 1.14, since the composition of a δ_1 -RIP and a δ_2 -RIP is a $\delta_1 + \delta_2$ -RIP.

The mapping φ_0 is natural map $x \mapsto \text{sgn}(Ax)$, as in (1.2), but restricted to K_0 . Note that for $x, y \in K_0$ the Hamming distance is

$$\begin{aligned} d_H(x, y) &= \frac{1}{m} \sum_{\ell=1}^m \mathbf{1}\{ \theta_\ell^\perp \text{ separates } x \text{ and } y \} \\ &= \frac{1}{m} \sum_{\ell=1}^m \mathbf{1}_{W_{x,y}}(\theta_\ell). \end{aligned}$$

In expectation, this is the geodesic distance: $\mathbb{E} d_H(x, y) = d(x, y)$. We will show that

$$\mathbb{E} \sup_{x, y \in K_0} |d_H(x, y) - d(x, y)| \leq H(K) / \sqrt{m}$$

where $H(K)$ is defined in (1.13). Since $m \geq C\delta^{-2}H(K)^2$, and by the general large deviation result for empirical processes Lemma 5.9, we can then conclude our claim in high probability. In fact, appealing to concentration of measure again, we do not directly estimate the expectation above, but rather show that a sufficiently small quantile of the r.v. is dominated by $CH(K) / \sqrt{m}$, hence the expectation is as well.

To bring the hemisphere gaussian process into play, define

$$Z_{x,y} := \frac{1}{\sqrt{m}} \sum_{k=1}^m \varepsilon_j \mathbf{1}_{W_{x,y}}(\theta_j)$$

where $\{\varepsilon_j\}$ is an iid sequence of rademacher random variables. We will show that if $m \gtrsim \delta^{-2}H(K)$,

$$(3.1) \quad \mathbb{E} \sup_{x, y \in K_0} |Z_{x,y}| \lesssim H(K_0) \leq H(K).$$

This is sufficient, since we can apply the symmetrization estimate (5.7),

$$\mathbb{E} \sup_{x, y \in K} \left| \frac{1}{m} \sum_{k=1}^m \mathbf{1}_{W_{x,y}}(\theta_j) - d(x, y) \right| \lesssim \frac{1}{\sqrt{m}} \mathbb{E} \sup_{x, y \in K} |Z_{x,y}| \lesssim H(K) / \sqrt{m} \lesssim \delta,$$

where the last quantity follows from our assumption on m .

Holding the θ_j fixed, associated to the process $Z_{x,y}$ is the conditional metric

$$D(x, y)^2 := \frac{1}{m} \sum_{j=1}^m \mathbf{1}_{W_{x,y}}(\theta_j).$$

It remains to show that it is very close to the metric for the hemisphere gaussian process.

Lemma 3.2. *For any $\epsilon > 0$, there is a constant C_ϵ sufficiently large, so that for all $m \geq C_\epsilon \delta^{-2} H(K)$, with probability at $1 - \epsilon$, there holds*

$$\sup_{x,y \in K_0} \frac{|D(x, y)^2 - \mathbb{P}(W_{x,y})|}{\mathbb{P}(W_{x,y})} \leq 1$$

Thus, on a set of large measure the conditional sub-gaussian process given by $Z_{x,y}$ has a metric dominated by twice that of the metric for the hemisphere gaussian process. Thus, for any $0 < \epsilon < 1$, there is a $C_\epsilon > 0$ so that

$$\mathbb{P}\left(\sup_{x,y \in K_0} |Z_x - Z_y| \geq C_\epsilon H(K)\right) \leq 2\epsilon.$$

It follows from the deviation inequality (5.10) and standard concentration of measure estimates for gaussian processes that (3.1) holds, completing the proof.

We now turn to the proof of Lemma 3.2.

Proof. Fix $x, y \in K_0$ and let $p = \mathbb{P}(W_{x,y}) = d(x, y)$. It is a consequence of Bernstein's inequality that

$$\mathbb{P}(|D(x, y)^2 - \mathbb{P}(W_{x,y})| > p) \leq C_0 \exp\left(-\frac{p^2}{p/m + 3p/m}\right) \leq C_0 \exp(-mp/4).$$

By construction of K_0 , $p = d(x, y) > \delta/4$. Hence $mp \gtrsim m\delta > C\delta^{-1}H(K) \geq C' \log N(K, \delta/8)$, with the last inequality following from Sudakov's lower bound for the supremum of gaussian processes, see Theorem 5.1 as well as the discussion of (1.19). The constant C' can be made as large as desired, and there are at most $N(K, \delta/8)^2$ pairs of $x, y \in K_0$, so the proof is immediate from the union bound. \square

3.2. Remarks. Plan and Vershynin [15] proved analogs of the results above using the gaussian process $\gamma_x = \langle x, \gamma \rangle$, which has a metric that differs from that of the hemisphere process G_x by essentially a square root, compare (1.10) and (1.12). Recall that the set K_0 is a $c\delta$ packing in K , relative to the geodesic metric $d(x, y)$. This allows a direct comparison of $\omega(K_0)$ and $H(K_0)$, namely $H(K_0) \lesssim \delta^{-1/2} \omega(K_0)$, which follows from Talagrand's majorizing measure theorem. The key inequality (3.1) proved about K_0 shows that if $m \gtrsim \delta^{-3} \omega(K)^2 \gtrsim \delta^{-2} H(K_0)$, then there is a δ -RIP map from K to the Hamming cube of dimension m . And, when restricted to K_0 , this is the map $x \mapsto \text{sgn}(Ax)$, where the rows of A are the iid θ_j .

This has the disadvantage of not making the RIP explicit on all of K . Plan-Vershynin [15] show more, at the cost of additional powers of δ^{-1} . If $m \gtrsim \delta^{-6}\omega(K)^2$, then the map $x \mapsto \text{sgn}(Ax)$, on all of K , is a δ -RIP, with high probability.

It is reasonable to conjecture that the hemisphere constant $H(K)$ is the correct quantity governing the RIP property into the Hamming cube. Namely, that if $m \gtrsim \delta^{-2}H(K)^2$, then with high probability, the map $x \mapsto \text{sgn}(Ax)$ is a δ -RIP map from K into the m -dimensional Hamming cube. Verifying this conjecture seems to bump up against subtle questions about empirical processes indexed by hemispheres, and attendant issues related to concentration of measure in the sphere. That is why we resort to a small ambiguity about exactly what the RIP is. These issues do not arise in the sparse vector case, however, as is argued below.

3.3. The Case of Sparse Vectors in Theorem 1.14. We turn to the case of sparse vectors, where we can give a proof that the natural map $x \mapsto \text{sgn}(Ax)$ is a δ -isometry of the set $K = K_s$, for integer $0 < s \leq n$ into the Hamming cube. The very short proof is based on observations about VC classes and empirical processes that are collected in the concluding section of the paper.

Observe that for $x, y \in K_s$, we have the empirical process identity

$$|d_H(x, y) - d(x, y)| = \left| \frac{1}{m} \sum_{j=1}^m \mathbf{1}_{W_{x,y}}(\theta_j) - d(x, y) \right|.$$

Apply the empirical process inequality (5.16) with $\eta = 1$, and $u = C\sqrt{s \log_+ \frac{n}{s}}$. We conclude that the bound below holds with probability at least $1 - \exp(-Cs \log_+ \frac{n}{s})$.

$$\sup_{x, y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j=1}^m (\mathbf{1}_{W_{x,y}}(\theta_j) - d(x, y)) \right| \lesssim \sqrt{s \log_+ \frac{n}{s}}$$

With the normalization by \sqrt{m} above, our condition $m \gtrsim \delta^{-2}s \log_+ \frac{n}{s}$ clearly gives the desired conclusion.

4. SIGN-PRODUCT EMBEDDING PROPERTY

We turn to the analysis of Theorem 1.16. For a standard gaussian g on \mathbb{R}^n and $\lambda = \sqrt{\frac{2}{\pi}}$, we argue that this inequality holds for $x, y \in \mathbb{S}^n$.

$$\mathbb{E} \text{sgn}(\langle x, g \rangle) \langle y, g \rangle = \lambda \langle x, y \rangle.$$

Now, if $\langle x, y \rangle = 0$, then $\langle x, g \rangle$ and $\langle y, g \rangle$ are independent, verifying the equality above. And, by linearity in expectation and y , we can then reduce to the case of $x = y$. But then, the random variable above is the absolute value of Z , a standard gaussian on \mathbb{R} . And $\mathbb{E}|Z| = \lambda$. Thus, the bound in (1.18) fits within the empirical process framework.

We will provide a proof that with probability at least $1 - (\frac{s}{2n})^{2s}$, we have

$$\sup_{x, y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j=1}^m \operatorname{sgn}(\langle x, g_j \rangle) \langle y, g_j \rangle - \lambda \langle x, y \rangle \right| \leq C \sqrt{s \log_+ \frac{n}{s}},$$

where K_s is the collection of s -sparse vectors in \mathbb{S}^n . For each fixed $x, y \in K_s$, we certainly have

$$\mathbb{P} \left(\left| \frac{1}{\sqrt{m}} \sum_{j=1}^m \operatorname{sgn}(\langle x, g_j \rangle) \langle y, g_j \rangle - \lambda \langle x, y \rangle \right| > 2 \right) \leq \frac{1}{2}.$$

It follows that the symmetrization inequality Lemma 5.5 holds. Namely it suffices to show that with probability at least $1 - \frac{1}{4} (\frac{s}{2n})^{2s}$, we have

$$(4.1) \quad \sup_{x, y \in K_s} \sup_{x', y' \in K_s} \underbrace{\left| \frac{1}{\sqrt{m}} \sum_{j=1}^m \varepsilon_j \operatorname{sgn}(\langle x, g_j \rangle) \langle y, g_j \rangle \right|}_{:=Z(x, y)} \leq C \sqrt{s \log_+ \frac{n}{s}}.$$

Above, we take ε_j to be an independent set of rademacher r.v.s.

To ease notations, call the sum above $Z(x, y)$, set $\bar{x}_j = \operatorname{sgn}(\langle x, g_j \rangle)$, and $\bar{y}_j = \langle y, g_j \rangle$. It is essential to observe that $\varepsilon_j \bar{x}_j \bar{y}_j$ is distributed like a one dimensional mean zero gaussian, of variance $\|y\|^2$. (On the other hand $\varepsilon_j (\bar{x}_j - \bar{x}'_j) \bar{y}_j$ is not gaussian, since it will equal zero if $g_j / \|g_j\| \notin W_{x, x'}$.)

The proof of (4.1) then combines three ingredients. (1) For fixed $x \in K_s$, the r.v. $\sup_{y \in K_s} |Z(x, y)|$ is controlled, and has concentration of measure. We can, essentially for free, form a supremum over x in net $X \subset K_s$ of small diameter. (2) As $x', x'' \in K_s$ vary over all pairs of vectors that are close, the selector random variables $\bar{x}'_j - \bar{x}''_j \in \{-1, 0, 1\}$ pick out a set of small cardinality. And (3) the supremum over the sum of the \bar{y}_j over sets of small cardinality is controlled.

The details are as follows. For fixed x , the process $\{Z(x, y) : y \in K_s\}$ is a gaussian process, with mean given in terms of the gaussian mean width of K_s .

$$\mathbb{E} \sup_{y \in K_s} |Z(x, y)| \lesssim \omega(K_s) \lesssim \sqrt{s \log_+ \frac{n}{s}}.$$

Furthermore, gaussian processes have a concentration of measure around their means. Therefore, for any finite set $X \subset K_s$, provided

$$(4.2) \quad \log \#X \lesssim s \log_+ \frac{n}{s},$$

we have, with probability at least $1 - (\frac{s}{10n})^{2s}$,

$$\sup_{x \in X, y \in K_s} |Z(x, y)| \lesssim \sqrt{s \log_+ \frac{n}{s}}.$$

Take $X \subset K_s$ to be a minimal cardinality collection of vectors so that for all $y \in K_s$, there is an $x \in X$ with $d(x, y) \leq \eta = (\frac{s}{100n})^6$. Observe that by the VC property of wedges and Lemma 5.13,

$$\#X \lesssim \binom{n+1}{s} \eta^{-3(s+1)} \lesssim (100 \frac{n}{s})^{20(s+1)}.$$

This is the estimate (5.14), applied with the uniform probability measure on the sphere. Therefore, (4.2) holds. This completes the first stage of the argument.

In the second stage, we have the differences $Z(x, y) - Z(x', y)$, which involve $\bar{x}_j - \bar{x}'_j \in \{-1, 0, 1\}$. But these last differences will be non-zero on a small set of indices. Indeed, with probability at least $1 - (\frac{s}{10n})^{2s}$, the inequality below holds.

$$(4.3) \quad \sup_{\substack{x, x' \in K_s \\ d(x, x') < \eta}} \sum_{j=1}^m |\bar{x}_j - \bar{x}'_j| \leq C \sqrt{m\eta s \log_+ \frac{n}{s}} = u.$$

The left side of (4.3) is the un-normalized Hamming metric between the natural mapping of x and x' in the Hamming metric. With our choice of $\eta = (\frac{s}{100n})^6$, note that

$$\eta \leq \sqrt{m\eta s \log_+ \frac{n}{s}}.$$

Subtracting the actual Hamming metric, we should bound the supremum

$$(4.4) \quad \sup_{\substack{x, x' \in K_s \\ d(x, x') < \eta}} \left| \sum_{j=1}^m \mathbf{1}_{W_{x, x'}}(\theta_j) - d(x, x') \right|$$

where $\theta_j = g_j / \|g_j\|$. Note that this is an empirical process for which we have the concentration of measure of Lemma 5.9. Thus, the high probability part of our assertion will follow from this fact.

Now, Theorem 5.15 proves that the expectation in (4.4) is bounded by

$$\sqrt{m} \int_0^{\eta^{1/2}} \sqrt{s \log_+ \frac{n}{s} + s \log_+ \frac{1}{t}} dt \lesssim \sqrt{m\eta s \log_+ \frac{n}{s}}$$

So, (4.3) holds, and the second stage of the argument is complete.

The third and final stage is to show that with probability $1 - (\frac{s}{10n})^{2s}$,

$$(4.5) \quad \sup_{\substack{J \subset \{1, \dots, m\} \\ |J| \leq u}} \sup_{y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j \in J} \varepsilon_j \bar{y}_j \right| \lesssim \sqrt{s \log_+ \frac{n}{s}}.$$

Above, u is as in (4.3).

The sum in (4.5) is that of a gaussian process, hence to deduce the claim, it suffices to bound the expectation on the left in (4.5). For fixed $J \subset \{1, \dots, m\}$, of cardinality at

most u , the supremum below

$$\sup_{y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j \in J} \varepsilon_j \bar{y}_j \right|$$

is that of a gaussian process with metric dominated by $\sqrt{u/m} \|y - y'\|_2$. Therefore, the estimate below is immediate.

$$\mathbb{E} \sup_{y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j \in J} \varepsilon_j \bar{y}_j \right| \lesssim \sqrt{u/m} \omega(K_s) \lesssim \sqrt{\frac{u}{m} \cdot s \log_+ \frac{n}{s}}.$$

Since this is universal in J , the next step is to appeal to concentration of measure above the mean. For this, we just need to count the number of sets J of cardinality at most u , of which there are clearly at most $2^{\binom{m}{u}}$. We conclude that

$$(4.6) \quad \mathbb{E} \sup_{\substack{J \subset \{1, \dots, m\} \\ |J| \leq u}} \sup_{y \in K_s} \left| \frac{1}{\sqrt{m}} \sum_{j \in J} \varepsilon_j \bar{y}_j \right| \lesssim \sqrt{\frac{u}{m}} \left(\sqrt{s \log_+ \frac{n}{s}} + \sqrt{u \log_+ \frac{m}{u}} \right).$$

It remains to see that the right side of (4.6) is at most $C \sqrt{s \log_+ \frac{n}{s}}$, and from this fact, the conclusion (4.5) follows from concentration of measure. The right side of (4.6) has two terms, the first is obviously less than $\sqrt{s \log_+ \frac{n}{s}}$, since $u \leq m$. The second is

$$\begin{aligned} \frac{u}{\sqrt{m}} \sqrt{\log_+ \frac{m}{u}} &= \sqrt{\eta s \log_+ \frac{n}{s}} \times \sqrt{\log_+ \frac{m}{\eta s \log_+ \frac{n}{s}}} \\ &\lesssim \sqrt{s \log_+ \frac{n}{s}} \left(\frac{s}{100n} \right)^3 \sqrt{\log_+ \frac{m}{s \log_+ \frac{n}{s}}} \\ &\lesssim \sqrt{s \log_+ \frac{n}{s}} \left(\frac{s}{100n} \right)^3 \sqrt{\log_+ \delta^{-2}} \lesssim \sqrt{s \log_+ \frac{n}{s}}. \end{aligned}$$

Here, we have used the choice of $\eta = \left(\frac{s}{100n} \right)^6$, $m \approx \delta^{-2} s \log_+ \frac{n}{s}$, and the definition of u in (4.3). The last inequality follows from the condition on δ in (1.17). This completes the proof.

5. BACKGROUND ON STOCHASTIC PROCESSES

5.1. Gaussian Processes. A process $\{Z_t : t \in T\}$ is a mean zero gaussian process iff for every finite subset $T' \subset T$, the restriction $\{Z_{t'} : t \in T'\}$ is a finite dimensional mean zero gaussian vector. The process Z_t induces a metric on T by

$$d_Z(s, t) = \|X_s - X_t\|_2 = \left(\mathbb{E} |X_t - X_s|^2 \right)^{1/2}.$$

We shall be interested in the bounded sample path properties, which are measured by

$$\mathbb{E} \sup_t |Z_t|, \quad \mathbb{E} \sup_{s, t} |Z_s - Z_t|.$$

The second expression can be written without absolute values.

Since a gaussian process is uniquely defined by its mean and covariance, any subset K of a Hilbert space H generates a mean zero gaussian process $\{Z_t : t \in K\}$, where the metric d_Z is given by the Hilbert space metric, $d_Z(s, t) = \|s - t\|_H$. This is in fact how the gaussian mean width and hemisphere processes could be defined. If the set K is symmetric, then the suprema of the gaussian processes, with or without the absolute values, are the same.

The quantity $\mathbb{E} \sup_{t \in T} Z_t$ is of basic interest for us. A foundational result concerning this quantity is the Sudakov lower bound.

Theorem 5.1. [*Sudakov Lower Bound*] For a gaussian process $\{Z_t : t \in T\}$, one has

$$\sup_{\delta} \delta \sqrt{\log N(T, d_Z, \delta)} \lesssim \mathbb{E} \sup_{s, t} |Z_s - Z_t|,$$

where $N(T, d_Z, \delta)$ is the covering number of T with respect to the metric of the gaussian process Z , i.e. the least number of d_Z -balls of radius δ needed to cover T .

Characterizing the bounded sample path properties of gaussian processes is an important accomplishment of Talagrand in the majorizing measure theorem [19], to which we referred in the text.

We also make use of the concentration of measure.

Theorem 5.2. [*Gaussian Concentration of Measure*] Let Z_t be a mean zero Gaussian process with $\sigma^2 = \sup_{t \in T} \mathbb{E} Z_t^2$ the maximal variance. There is a constant $c > 0$ so that for all $\lambda > 0$

$$\mathbb{P}(c \sup_t Z_t > \mathbb{E} \sup_{t \in T} Z_t + \lambda) \lesssim \exp(-(\lambda/\sigma)^2).$$

A random variable X is said to be *sub-gaussian with parameter σ* if

$$(5.3) \quad \mathbb{P}(|X| > \lambda) \lesssim \exp(-(\lambda/\sigma)^2), \quad \lambda > 0.$$

The supremum of a finite number of sub-gaussian random variables grows very slowly.

Lemma 5.4. Let X_1, \dots, X_N be sub-gaussian r.v.s with maximal parameter σ . Then,

$$\mathbb{E} \sup_{1 \leq n \leq N} |X_n| \lesssim \sqrt{\log N}.$$

5.2. Empirical Processes. Let (T, μ) be a probability space, and X_1, \dots, X_m are iid random variables, taking values in T with distribution μ . For a class of functions $f \in \mathcal{F}$, mapping T into \mathbb{R} , we define the empirical process

$$S_m f = \frac{1}{m} \sum_{j=1}^m f(X_j) - \int_T f d\mu, \quad f \in \mathcal{F}.$$

Above, we must assume that $f \in L^1(\mathbb{T}, \mu)$, and while the case of general functions is quite interesting, but for our purposes we can always assume that f are bounded functions. Indeed, we mostly work with the case $\mathcal{F} = \{\mathbf{1}_{W_{x,y}}\}$. One is then interested in variants of the law of large numbers and central limit theorem in this context. The quantity important to us is

$$S_m(\mathcal{F}) := \sup_{f \in \mathcal{F}} |S_m f|.$$

Basic to the analysis of these processes is *symmetrization*, which we state in two forms.

Lemma 5.5. [2] *Fix $s > 0$ so that*

$$\sup_{f \in \mathcal{F}} \mathbb{P}(|S_m f| > s) \leq \frac{1}{2}.$$

Then, for $t > 0$, there holds

$$\mathbb{P}\left(\sup_{f \in \mathcal{F}} |S_m f| > t\right) \leq 2\mathbb{P}\left(\sup_{f \in \mathcal{F}} |S_m f - S'_m f| > t - s\right)$$

where $S'_m f$ is an independent copy of $S_m f$.

A second symmetrization lemma for expectation is as follows.

Lemma 5.6. [Symmetrization] *For any class of μ -integrable functions \mathcal{F} one has*

$$(5.7) \quad \mathbb{E} S_m(\mathcal{F}) \leq 2\mathbb{E} \sup_{f \in \mathcal{F}} \left| \frac{1}{m} \sum_{j=1}^m r_j f(X_j) \right|$$

$$(5.8) \quad \leq 2\mathbb{E} \sup_{f \in \mathcal{F}} \left| \frac{1}{m} \sum_{j=1}^m g_j f(X_j) \right|.$$

Above, $\{r_j\}$ denote Rademacher and $\{g_j\}$ standard gaussian random variables, both independent of the $\{X_j\}$.

In the first line (5.7), conditional on the $\{X_j\}$, the process is a *rademacher process*, which is subgaussian, in the sense of (5.3). Hence, it is dominated by the conditional gaussian process in (5.8). The conditional metric on the set \mathcal{F} is of fundamental importance. It is

$$d(f, g) = \frac{1}{m} \sum_{j=1}^m |f(X_j) - g(X_j)|^2.$$

Estimating the term $S_m(\mathcal{F})$ is fundamental in the case when \mathcal{F} consists of functions bounded by one, in view of the following deviation inequality, which is an application of the Hoeffding inequality, and is sometimes called the McDiarmid inequality.

Lemma 5.9. [16] *Suppose that \mathcal{F} consist of functions bounded by one. There holds*

$$(5.10) \quad \mathbb{P}(S_m(\mathcal{F}) > \lambda + \mathbb{E} S_m(\mathcal{F})) \lesssim \exp(-m\lambda^2/2)$$

To estimate $S_m(\mathcal{F})$, where \mathcal{F} consists of indicator sets, one should divide \mathcal{F} into parts which are either governed by the gaussian theory, or have no cancellative part, and so should be controlled by Poisson like behavior. While there are several techniques here, one should not forget that concentration of measure on the sphere is an obstacle to the use of techniques such as ‘bracketing.’ A succinct summation of the main conjectures in the subject are in [19, Chap. 9].

5.3. VC Dimension. Let (M, \mathcal{M}) be a measure space, and $\mathcal{C} \subset \mathcal{M}$ a class of measurable sets. For integers n , let

$$S(n, \mathcal{C}) := \sup_{\substack{A \subset M \\ |A|=n}} |\{B : B = A \cap C, C \in \mathcal{C}\}|$$

That is, $S(n)$ is the largest number of subsets that can be formed by intersecting a set A of cardinality n with sets $C \in \mathcal{C}$. It is clear that $S(n) \leq 2^n$. The Vapnik-Cervonenkis dimension (VC-dimension) $\nu(\mathcal{C})$ of \mathcal{C} is the least integer d such that $S(n) < 2^n$ for all $n > d$. The Sauer Lemma then states that

$$(5.11) \quad S(n) < \left(\frac{ne}{d}\right)^d, \quad n > 1.$$

For us, the class of relevant sets are *hemispheres*. More generally, for spherical caps in \mathbb{S}^n , the VC-dimension is $n + 1$.

Proposition 5.12. *The family \mathcal{C}_n of spherical caps on \mathbb{S}^n has VC-dimension $n + 1$.*

Proof. This is a modification of the proof in [1, Prop. 8]. A collection A of points on the unit sphere that is shattered by spherical caps has the additional property that for any partition of A into disjoint subsets A' and A'' , the convex hulls of A' and A'' can't intersect. This is just because a spherical cap is itself the intersection of the sphere with a half-space. But Radon's Theorem, a basic result in convex geometry, says that any $n + 2$ points in \mathbb{R}^n can be partitioned into two disjoint sets whose convex hulls intersect. Hence $\nu(\mathcal{C}_n) < n + 2$.

We show that spherical caps shatter the collection of $n + 1$ vectors

$$A = \{\mathbf{e}_1, \dots, \mathbf{e}_n, \mathbf{1}/\sqrt{n}\}$$

where the \mathbf{e}_j is the standard basis elements, and $\mathbf{1}$ is the vector of all ones. Consider a partition of A into A' and A'' , supposing that $\mathbf{1}/\sqrt{n+1} \in A''$. If either A' or A'' is a singleton, a small radius spherical cap around the singleton provides this decomposition. Otherwise, observe that there is a $\theta \in \mathbb{S}^n$ with

$$\langle x, \theta \rangle = 0 \quad x \in A',$$

and $\langle x, \theta \rangle > 0$ for $x \in A''$. Indeed, the first condition only requires that the j th coordinates of θ be zero, for $\mathbf{e}_j \in A'$. And otherwise we require that j th coordinate of θ be positive, which we can do since A'' is not a singleton. \square

Thus, the class of hemispheres in \mathbb{S}^n has dimension at most $n + 1$. Now, we are also interested in the class of symmetric differences of hemispheres. But, it follows from estimate (5.11) that forming the symmetric difference of a class \mathcal{C} of finite VC-dimension increases the dimension by at most a factor of 3. In symbols, $\nu(\mathcal{C}\Delta\mathcal{C}) \leq 3\nu(\mathcal{C})$.

The next important property of VC-classes is that they are universal for empirical processes. This is encoded in an inequality of this form.

Lemma 5.13. *Let \mathcal{C} be a collection of sets on a measure space (M, \mathcal{M}) with VC-dimension d . Let P be any probability measure on (M, \mathcal{M}) , and set metric d_P on \mathcal{C} by $d_P(C_1, C_2) = P(C_1\Delta C_2)$. For the metric entropy numbers $N(\mathcal{C}, d_P, \delta)$, we have the estimate*

$$N(\mathcal{C}, d_P, \delta) \lesssim (\delta/2)^{-4d}, \quad 0 < \delta < 1.$$

5.4. Uniform Entropy Classes. It is convenient to recall a consequence of Lemma 5.13 in somewhat greater generality. The class $\mathcal{W}_s := \{W_{x,y} : x, y \in K_s\}$ of symmetric differences of hemispheres associated to sparse vectors is not VC, unless we further restrict the coordinates in which x is not zero. But, this class of sets is still ‘universal’ in this sense. For any probability measure P on \mathbb{S}^n , define the metric d_P as in Lemma 5.13. We have the following estimate on the metric entropy of \mathcal{W}_s .

$$(5.14) \quad N(\mathcal{W}_s, d_P, \delta) \lesssim \binom{n+1}{s}^2 (\delta/2)^{-12s-24}, \quad 0 < \delta < 1.$$

This is a direct consequence of Proposition 5.12 and Lemma 5.13, after holding the coordinates in which x and y are supported fixed.

The class \mathcal{W}_s satisfies the *uniform entropy bounds* of Panchenko. And, as a corollary to [12, Cor. 1], we derive the result below. (See the beginning of [12, §1.1] for relevant definitions. The proof of the Corollary is a standard, but somewhat delicate, chaining argument, using the uniform entropy bounds.)

Theorem 5.15. *For $\{\theta_j\}$ iid and uniform on \mathbb{S}^n , and $0 < \eta < 1$, and $u > 0$, with probability at least $1 - 2e^{-u^2}$,*

$$(5.16) \quad \sup_{\substack{x,y \in K_s \\ d(x,y) < \eta}} \frac{1}{\sqrt{m}} \sum_{j=1}^m (\mathbf{1}_{W_{x,y}}(\theta_j) - d(x,y)) \lesssim \int_0^{\eta^{1/2}} \sqrt{s \log_+ \frac{n}{s} + s \log_+ \frac{1}{t}} dt + u \sqrt{\eta}.$$

Proof. In the language of Panchenko [12], the collection of wedges $\{W_{x,y} : x, y \in K_s\}$ satisfy *uniform entropy bounds*, namely the inequality (5.14) holds. And the statement above is then a particular instance of [12, Corollary 1]. \square

REFERENCES

- [1] C. Aistleitner, J. S. Brauchart, and J. Dick, *Point sets on the sphere \mathbb{S}^2 with small spherical cap discrepancy*, Discrete Comput. Geom. **48** (2012), no. 4, 990–1024. \uparrow 20

- [2] Kenneth S. Alexander, *Probability inequalities for empirical processes and a law of the iterated logarithm*, Ann. Probab. **12** (1984), no. 4, 1041–1067. MR757769 (86b:60048) ↑19
- [3] R. Baraniuk, S. Foucart, D. Needell, Y. Plan, and M. Wootters, *Exponential decay of reconstruction error from binary measurements of sparse signals* (2014), available at <http://arxiv.org/abs/1407.8246>. ↑2, 7
- [4] Dmitriy Bilyk and Michael T. Lacey, *One Bit Sensing, Discrepancy, and Stolarsky Principle* (2015), available at <http://arxiv.org/abs/arXiv:1511.08452>. ↑2
- [5] P. Boufounos and R. Baraniuk, *1-bit compressive sensing*, Proc. 42nd Annu. Conf. Inf. Sci. Syst., Princeton, NJ, (Mar. 2008). ↑2
- [6] Emmanuel J. Candes and Terence Tao, *Near-optimal signal recovery from random projections: universal encoding strategies?*, IEEE Trans. Inform. Theory **52** (2006), no. 12, 5406–5425. ↑4
- [7] Piotr Indyk and Stanislaw Szarek, *Almost-Euclidean subspaces of ℓ_1^N via tensor products: a simple approach to randomness reduction*, Approximation, randomization, and combinatorial optimization, Lecture Notes in Comput. Sci., vol. 6302, Springer, Berlin, 2010, pp. 632–641. ↑7
- [8] Laurent Jacques, *Small width, low distortions: quasi-isometric embeddings with quantized sub-Gaussian random projections* (2015). ↑6
- [9] L. Jacques and K. Degraux, *Quantized Iterative Hard Thresholding: Bridging 1-bit and High-Resolution Quantized Compressed Sensing*, Proceedings of International Conference on Sampling Theory and Applications 2013, Bremen, Germany, 2013, available at <http://hdl.handle.net/2078.1/134466>. ↑2, 6, 7
- [10] Laurent Jacques, Jason N. Laska, Petros T. Boufounos, and Richard G. Baraniuk, *Robust 1-bit compressive sensing via binary stable embeddings of sparse vectors*, IEEE Trans. Inform. Theory **59** (2013), no. 4, 2082–2102. ↑2
- [11] B. Klartag and S. Mendelson, *Empirical processes and random projections*, J. Funct. Anal. **225** (2005), no. 1, 229–245. ↑4
- [12] Dmitriy Panchenko, *Some extensions of an inequality of Vapnik and Chervonenkis*, Electron. Comm. Probab. **7** (2002), 55–65 (electronic). ↑21
- [13] Yaniv Plan and Roman Vershynin, *One-bit compressed sensing by linear programming*, Comm. Pure Appl. Math. **66** (2013), no. 8, 1275–1297. ↑2, 3, 11
- [14] ———, *Robust 1-bit compressed sensing and sparse logistic regression: a convex programming approach*, IEEE Trans. Inform. Theory **59** (2013), no. 1, 482–494. ↑7, 11
- [15] ———, *Dimension reduction by random hyperplane tessellations*, Discrete Comput. Geom. **51** (2014), no. 2, 438–461. ↑2, 6, 7, 8, 13, 14
- [16] WanSoo T. Rhee and Michel Talagrand, *Martingale inequalities and NP-complete problems*, Math. Oper. Res. **12** (1987), no. 1, 177–181. ↑19
- [17] Gideon Schechtman, *Two observations regarding embedding subsets of Euclidean spaces in normed spaces*, Adv. Math. **200** (2006), no. 1, 125–135. ↑4
- [18] Kenneth B. Stolarsky, *Sums of distances between points on a sphere. II*, Proc. Amer. Math. Soc. **41** (1973), 575–582. ↑5
- [19] Michel Talagrand, *Upper and lower bounds for stochastic processes*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 60, Springer, Heidelberg, 2014. Modern methods and classical problems. ↑18, 20

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS MN 55455, USA

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA GA 30332, USA
E-mail address: lacey@math.gatech.edu