

Neurofeedback Tunes Long-Range Temporal Correlations in Spontaneous Brain Activity

T. Ros ^{a,1}, P. Frewen ^b, J. Théberge^c, R. Kluetsch ^d, A. Mueller ^e, G. Candrian ^e, R. Jetly ^f, P. Vuilleumier ^{a*}, R.A. Lanius ^{b*}

^a Laboratory for Neurology and Imaging of Cognition, Department of Neurosciences, University of Geneva, Switzerland

^b Department of Psychiatry, University of Western Ontario, London, Ontario, Canada

^c Department of Medical Imaging, Lawson Health Research Institute, London, Ontario

^d Department of Psychosomatic Medicine and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim Heidelberg University, Mannheim, Germany

^e Brain and Trauma Foundation, Chur, Switzerland

^f Directorate of Mental Health, Canadian Forces Health Services, Ottawa, Canada

¹ To whom correspondence may be addressed. E-mail: tomasino.ros@gmail.com

* These authors contributed equally to this work.

Abstract

The amplitude dynamics of neuronal oscillations, namely their scale-free, long-range temporal correlations (LRTCs), are considered supporting evidence for criticality in the brain, a balanced state between order and disorder theoretically optimal for information processing. This idea is compatible with observations of abnormally random (decreased LRTCs) or self-sustained (increased LRTCs) oscillation amplitudes in a variety of brain disorders. Interestingly, oscillation amplitude may be volitionally controlled with the aid of an electroencephalogram (EEG) brain-computer interface, utilizing a closed neurofeedback loop. However, despite emerging clinical applications, the nature of neurofeedback's impact on brain function remains controversial and poorly understood. Here, we directly examined lasting changes in spontaneous LRTCs after a single NFB training session, consisting of a tonic suppression of alpha oscillation amplitude for 30 minutes. By applying detrended fluctuation analysis (DFA), we detected a post-neurofeedback enhancement of spontaneous alpha-amplitude LRTCs in a group of healthy young adults, between real versus sham conditions. In a second experiment, we examined psychiatric patients with posttraumatic stress disorder (PTSD) for an equivalent mechanism, replicating a comparable boost of alpha-amplitude LRTCs, which correlated with significant reductions in hyperarousal. Interestingly, neurofeedback rescued LRTCs in regions that were abnormally low in value (i.e., increased randomness) compared to a healthy control sample. Lastly, we identified an inverse-U relationship between strength of LRTC and alpha amplitude, suggesting a gradual breakdown of long-range dependence at high/low synchronization extremes, and consistent with theoretical predictions about criticality. By uncovering a substantially independent mechanism of action, our findings offer a novel framework for motivating research and clinical applications of neurofeedback, encompassing brain disorders with perturbed LRTCs.

Introduction

The multiple spatiotemporal scales through which brain activity can be studied presents a veritable challenge in neuroscientists' quest to link brain and behaviour. Although traditional approaches have been mostly restricted to discrete spatial or temporal scales, recent investigations are bearing witness to an emerging interest in scale-free (or 'fractal') measures of brain function (He et al., 2010; Kello et al., 2010). As their name implies, such measures are able to capture relationships *across*

different levels of brain organization, inherent for example, in the temporal structure of electrocortical patterns (Van de Ville et al., 2010) or the topology of functional connectivity networks (Liu et al., 2014). Here, scale-free relationships are frequently represented by the Hurst scaling exponent (H), as a measure of self-similarity within time-series, which indicates the degree of long-range temporal (auto)correlations (LRTCs) present between shorter and longer time scales (e.g. ranging from seconds to minutes). A larger H value generally reflects the presence of a long-range (yet transient) trend in the data, for example when values alternate between high and low values, but do so for a prolonged period at a time in each state. Given that H integrates a signal's temporal evolution, it may be regarded to index its long-term dependence, or *memory*. Thereby, H essentially estimates the extent of temporal memory in a signal, with random white-noise and a smooth line taking $H = 0.5$ and $H = 1$ values, respectively. A fascinating implication of scale-free indices is their link with the nascent science of complex systems, which reputedly operate at a 'critical' balance between spatiotemporal order and disorder (Chialvo, 2010). Intriguingly, spontaneous neural activity has been found to exhibit temporal memory reflected in positive, long-range correlations (i.e. $H > 0.5$) (Linkenkaer-Hansen, 2001; Nikulin and Brismar, 2004), lending support to hypotheses that the brain may display a 'self-organised' form of criticality (SOC) (Bak et al., 1987; Hesse and Gross, 2014): a homeostatic state which maximizes the dynamic range and memory required for information processing (Hsu and Beggs, 2006; Shew and Plenz, 2012). This notion is consistent with studies in patients with brain disorders, which report abnormal resting state H -values relative to the healthy population, thus signalling disrupted LRTCs. For example, schizophrenia (Nikulin et al., 2012) and Alzheimer's disease (Montez et al., 2009) patients exhibit attenuated (i.e. more random) LRTCs in the amplitude fluctuations of alpha-band oscillations, while theta-amplitude LRTCs seem to negatively correlate with major depression severity (Linkenkaer-Hansen et al., 2005). Elsewhere, LRTCs have been directly linked to fluctuations in behavioural performance (Palva et al., 2013; Smit et al., 2013), brain maturation (Smit et al., 2011), and levels of consciousness (Tagliazucchi et al., 2013). Notably, both computational modelling (Poil et al., 2012) and subdural recordings (Monto et al., 2007) suggest that network changes in the excitation/inhibition balance may underpin disordered LRTCs.

In light of this interdependence between temporally-complex neural dynamics, behaviour, and pathology, a therapeutic means by which to normalize LRTCs might unlock a promising treatment avenue for brain disorders, collectively the leading contributor to global disease burden (Collins et al., 2011). In this work, we report on the exciting possibility of tuning LRTCs of spontaneous brain oscillations following non-invasive, closed-loop neurofeedback (NFB) training (Kamiya, 2011). During NFB, a sensory description of real-time brain activity is fed-back to users via a brain-computer interface (BCI), enabling top-down control of network oscillations, including amplitude (Hardt and Kamiya, 1978), frequency (Angelakis et al., 2007) and functional connectivity (Brunner et al., 2006). Although NFB is showing early promise for treating brain disorders (Niv, 2013; Ros et al., 2014), there is still a poor mechanistic understanding of its influence on brain function. Interestingly, as NFB has been shown to concomitantly alter the amplitude of cortical oscillations and the balance of cortical excitation/inhibition (Ros et al., 2010), we recently hypothesized this might be associated with changes in signature(s) of critical brain dynamics (e.g. LRTCs) (Ros et al., 2014). In this work, we begin by demonstrating that one session of NFB training may reliably enhance alpha-band LRTCs in healthy subjects relative to a sham-control group. We then go on to show that NFB is able to rescue abnormally reduced alpha-band LRTCs in patients with post-traumatic stress disorder (PTSD), together with a significant and correlated improvement in symptoms of hyperarousal.

Materials and Methods

Subjects

Experiments were approved by the Research Ethics Board of Western University, Ontario, Canada and all participants were recruited from the university neighbourhood. *Experiment 1* involved a group of 40 healthy adult subjects (mean age 33.6, SD 11.1, 25 female). These participants were carefully screened for the presence of neurological or psychiatric disorders during a structured SCID-I- Interview. *Experiment 2* involved 21 adult PTSD patients (mean age 39.9, SD 13.7, 18 female). All met the DSM-IV (1) criteria for a primary diagnosis of PTSD related to childhood maltreatment. Axis I diagnoses were assessed by a trained psychologist using the Structured Clinical Interview for DSM-IV Axis I Disorders (42) and the Clinician-Administered PTSD Scale (CAPS, total cut-off score >50) (43). The CAPS indicated the PTSD group met the necessary criteria for trait hyperarousal (a score of >2/5). Exclusion criteria comprised a lifetime diagnosis of a psychotic disorder, bipolar disorder, substance use disorders, a history of head trauma, serious medical or neurological illness. Eleven participants were currently taking psychotropic medications. The healthy control group consisted of 30 healthy adult subjects (mean age 39.4, SD 8.7, 26 female), and was matched for age/gender from the Human Brain Institute (HBI) normative database (<http://www.hbimed.com/>). Lastly, an additional set of healthy adults (n=32) were recruited for a resting-state *eyes closed* recording. All participants' EEG recordings were performed using the same amplifier type (Mitsar-201).

Experimental Timeline

For all NFB, SHAM and PTSD participants, the overall protocol consisted of 3 sequential parts that occurred within the same daytime visit: fMRI scan before neurofeedback (~30 min), EEG neurofeedback (30 min), and fMRI scan after neurofeedback (~30 min). fMRI analyses of Experiments 1 and 2 have been previously reported in (Ros et al., 2013) and (Kluetsch et al., 2014), respectively, while we additionally recruited another 6 participants to boost the overall sample of Experiment 1. In this paper, we focus on the EEG resting-state recordings that directly flanked the start and end of one individual neurofeedback session. Upon arrival to the examination facility, healthy participants were randomized to one of two experimental groups: EEG-neurofeedback (NFB, n=20) or sham-neurofeedback (SHAM, n=20). For ethical reasons, PTSD patients received only EEG-neurofeedback. Each session consisted of a 3-min resting-state (no-feedback), followed by 30 min of continuous feedback (real or sham), and lastly another 3-min resting-state (all in eyes open). Immediately before and after the resting-state, participants completed Thayer's Activation–Deactivation Checklist questionnaire. Here the “uncalmness” subscale was used as a self-reported measure of state arousal. No adverse effects were reported after NFB or SHAM feedback.

EEG recording and feedback training

Scalp voltages of all participants were recorded using a 19 channel electrode cap (Electro-cap International, Inc.) according to the 10-20 international system. The ground electrode was placed at site AFz. Electrical signals were amplified with the Mitsar 21-channel EEG system (Mitsar-201, CE0537, Mitsar, Ltd) and all impedances were kept under 5 k Ω . EEG was recorded continuously, digitized at a sampling rate of 250 Hz, and stored on hard disk for off-line analysis. In parallel, a bridged Pz channel was specifically used for neurofeedback and was connected to a ProComp+ amplifier (Thought Technology, Canada) interfacing with EEGer 4.2 neurofeedback software (EEG Spectrum Systems, CA). During (feedback-free) resting-state recordings, participants were asked to

relax with their eyes open and gaze at a blank wall. SHAM group participants did not receive veridical feedback from their brain activity, but were re-played EEG signal from a previously recorded session of a NFB-successful participant (their whole-scalp EEG activity was nevertheless recorded passively). For online training, the EEG signal was IIR (infinite impulse response) band-pass filtered to extract alpha (8–12 Hz) amplitude with an epoch size of 0.5 seconds. Here, subjects were rewarded upon reduction of their absolute alpha amplitude, where threshold for reward was set to occur 60% of the time below the initial 3-min baseline average (i.e., 40% negative-feedback). Visual feedback was clearly displayed on a 17" monitor via 1) a dynamic bar graph at the center of the screen whose height was proportional to real-time alpha fluctuations, and 2) a "Space Race" game, where a spaceship advanced through space when amplitude was below threshold, and became stationary when above threshold. No explicit instructions were given on how to achieve control over the spaceship, and all participants were told to be guided by the visual feedback process. The 30-min session was divided into ten 3-min periods, with a break of 10 seconds between each period.

EEG analysis

For offline analyses, EEG signals were re-referenced to common-average reference. Low- and high-pass filters were set to 0.5 and 40 Hz, respectively, with a 55–65 Hz notch filter. EEG data was analysed with the Neurophysiological Biomarker Toolbox (NBT, <http://www.nbtwiki.net/>) in Matlab (MathWorks Inc., MA, USA). We used ICA decomposition to first remove stereotypical artifacts using the Infomax algorithm (blinking and lateral eye movement). Statistically-defined artifacting was then carried out with the FASTER plugin (Nolan et al., 2010) removing segments based on extremal deviations of amplitude and variance from the mean. Based on the NFB protocol, our hypotheses focused on: i) alpha amplitude long-range temporal correlation (LRTC) and ii) mean alpha amplitude. We restricted LRTC analyses to the resting-state given that training data was more contaminated and of variable length between subjects when artifacted. Each 3-minute EEG recording was firstly decomposed with a discrete wavelet transform (DWT) (Tumari et al., 2013; Kitzbichler and Bullmore, 2015) using the Wavos toolkit (Harang et al., 2012). Here, a Daubechies 4-tap wavelet was used and D5 detail extracted, corresponding to 7.8-15.6 Hz (i.e., alpha-band). Then, the instantaneous alpha amplitude (envelope) across time was determined using the absolute value of the Hilbert transform (see Fig S1). LRTCs were subsequently estimated via detrended fluctuation analysis (DFA), which estimates the scaling of the root-mean-square fluctuation of the integrated and linearly detrended signals, $F(t)$, as a function of time window size, t . As shown in Fig S2, H (Hurst) exponents were calculated using a linear-regression fit on a log-log graph with a range of 10 equally spaced points between 1-30 seconds, in line with previous work (Linkenkaer-Hansen, 2001; Montez et al., 2009). A linear goodness-of fit was additionally calculated for each DFA estimation using the R-squared statistic (Nikulin and Brismar, 2004). Lastly, absolute alpha amplitude (8-16 Hz) was estimated with a standard FFT approach using Welch's method (Matlab *pwelch* function) and a Hanning windowing function (4 second epoch, 50% overlap). Relative alpha amplitude was calculated as the ratio of the mean alpha amplitude and the broadband amplitude (1.5 - 40 Hz)

Statistical analysis

Multivariate analysis of variance (MANOVA, Pillai's trace statistic) F-tests were used to assess global changes of alpha amplitudes and H-exponent. The within-subject factors were *Time* (T1 and T2, i.e. pre and post feedback) and *Channels* (19 channels, see above). The between-subject factor was *Group* (NFB and SHAM). In order to account for multicollinearity among EEG channels, multivariate Hotelling's T^2 tests were conducted to examine directionality of the *Time* (T_1 and T_2) and *Group* (NFB and SHAM, or PTSD and CONTROLS) factors. Secondly, in order to reveal topographical (univariate) effects, two-tailed Student t-tests were performed channel-wise, within- and between- subjects

(dependent and independent tests, respectively); P-values below 0.05 and 0.01 are indicated on topographic plots. Following Montez et al (Montez et al., 2009), a correction for multiple comparisons was not necessary, because the number of channels with P-values below 0.05 ranged from 5 and 10 and the likelihood of having this many channels out of 19 by chance was less than 2% (cf. binomial distribution). Moreover, the channels were anatomically clustered in the topographic plots. Grand-average amplitudes and H-exponents across *all channels* are reported with standard error of mean (SEM) for respective experimental conditions and subject groups. Group differences in these grand-average means were computed using a two-tailed t-test with $P < 0.05$. Lastly, least-squares correlations were estimated using the Pearson coefficient R. All analyses had a statistical significance threshold of $\alpha = 0.05$, and all post-hoc comparisons were Bonferroni corrected.

Results

Experiment 1: Investigating *real vs sham* neurofeedback on LRTCs in healthy subjects

We investigated whether neurofeedback (NFB) regulation of the dominant human EEG rhythm, the alpha oscillation, would reliably alter alpha-band LRTCs in the post-NFB resting state. Here, we provided real-time feedback of alpha fluctuations on a computer visual display, and asked subjects to volitionally attenuate their parietal alpha amplitude for a total of 30 minutes; *before* and *after* which a 3-minute full-scalp EEG was recorded in the eyes open condition. This procedure was randomized to two groups: an experimental group (n=20) receiving veridical (i.e., real-time) feedback (NFB), and a control group (n=20) receiving sham (i.e., pre-recorded EEG) feedback (SHAM).




Figure S1: Summary of signal processing steps. **A.** Wavelet-filtered alpha oscillation (blue trace) recorded from Subject #2 (Pz channel) during resting-state, including its instantaneous amplitude envelope (red trace). **B.** Power spectrum of the same alpha oscillation. **C.** Instantaneous alpha amplitude (red trace) can also be visualised on a wavelet time-frequency plot where bursts of high/low alpha amplitude (red/blue, respectively) are evident during the whole resting-state recording. Long range temporal correlations (LRTCs) of alpha amplitude fluctuations are then estimated for the entire time-range of 3 minutes.

As shown in Figure S1, and in accordance with prior work (Linkenkaer-Hansen, 2001), LRTCs were estimated from the Hilbert envelope (i.e., amplitude) of wavelet-filtered alpha oscillations (8-16 Hz) over a resting-state period of 3 minutes. For each channel time-series, the Hurst (H) scaling exponent was estimated using detrended fluctuation analysis (DFA), and computed as the slope of the linear regression line on a log-log plot, with temporal windows ranging from 1 to 30 seconds (Figure S2). In accordance with previous work (Nikulin and Brismar, 2004), the linear model accounted for >99.5% of the variance explained in the entire dataset, while no significant differences of goodness-of-fit were present between experimental groups or conditions ($p < 0.05$), excluding the possibility that group/condition differences in H-exponent were driven by departures from the linear model.




Figure S2: Graphical representation of detrended fluctuation analysis (DFA). The Hurst exponent was defined by the slope of a linear regression line on a log-log plot, fitted within a min-max range of different window sizes (blue vertical lines). Here, the slope of the DFA log-log plot is numerically equivalent to the Hurst (H) exponent for values < 1 . For illustration, H-exponents (red lines) were calculated here for NFB-group subjects at baseline (Pz channel), with a fitting interval of 1 to 30 seconds; the black line highlights the fit for subject #11 ($H = 0.74$).

One-way MANOVAs (*Group*) of NFB and SHAM groups showed no significant *baseline* (T_1) differences in overall H-exponent ($F_{19,20} = 1.4$, n.s.), nor alpha amplitude ($F_{19,20} = 1.0$, n.s.). Regarding outcomes of feedback training, a two-way MANOVA revealed a significant *Group x Time* interaction in H-exponent ($F_{1,38} = 5.9$, $p < 0.05$), demonstrating a dissociation in pre-to-post feedback resting-state values between NFB and SHAM groups. As seen in Fig 1A/B, testing for directionality using multivariate paired tests ($T_2 - T_1$) indicated a significant increase of H-exponent values in the NFB group (Hotelling's $T^2 = 15.8$, $p < 0.05$), while no reliable change was present in the SHAM group (Hotelling's $T^2 = 0.5$, n.s.). On the other hand, as illustrated in Fig 1 C/D, we observed a main effect of *Time* for alpha amplitude ($F_{1,38} = 18.8$, $P < 0.01$). However, alpha amplitude did not exhibit a *Group x Time* interaction ($F_{1,38} = 0.1$, n.s.). Moreover, Pearson correlation analyses confirmed that pre-to-post shifts in H-exponent values were not correlated to intra-individual changes in alpha amplitude, either in NFB ($R = 0.15$, n.s.) or SHAM groups ($R = 0.09$, n.s.), further demonstrating their independence.




Figure 1. Pre (T_1) and post (T_2) resting-state changes in neurofeedback (NFB) and sham (SHAM) groups. (A) Topography of Hurst-exponents in the alpha-band reflecting LRTCs; **(B)** Topography of alpha amplitudes; **(C)** mean alpha Hurst-exponents; and **(D)** mean alpha amplitudes of individual NFB subjects (blue circles) and SHAM subjects (red diamonds). Grand averages indicate all subjects, error-bars denote \pm SEM. * significant difference at $p < 0.05$, n.s. not significant.

Experiment 2: Neurofeedback normalization of LRTCs in post-traumatic stress disorder

As post-traumatic stress disorder (PTSD) is associated with cognitive-affective dysregulation and anomalous EEG signatures (Jokić-Begić and Begić, 2003), we reasoned that it may also manifest perturbed LRTCs based on recent evidence in psychological distress (Churchill et al., 2015). We therefore sought to test and replicate the neuromodulatory effects of NFB in 21 adult patients with PTSD. Remarkably, and as shown in Fig 3A/B, paired tests confirmed the hypothesized boost of alpha H-exponent values post-neurofeedback (Hotelling's $T^2 = 9.24$, $p < 0.05$), which was maximally expressed along midline regions. Mean alpha amplitude experienced a parallel, albeit topographically non-overlapping enhancement ($T^2 = 24.7$, $p < 0.05$), as evidenced by Fig 3C/D. Here, global H-exponent changes correlated strongly with those of global alpha amplitude ($R = 0.72$, $p < 0.05$). In order to clarify whether these values moved towards or away from those of the normal population, we sampled an additional group of 30 healthy adults, matched for age and gender.

As illustrated in Fig 3, Hurst values for PTSD subjects were found to be significantly lower than CONTROLS at *baseline* (Hotelling's $T^2 = 65.3$, $p < 0.05$), indicating relatively more *random* alpha burst dynamics. This novel signature in PTSD patients was echoed by a deficit in mean alpha amplitude (Hotelling's $T^2 = 168.8$, $p < 0.05$), consistent with previous work (Jokić-Begić and Begić, 2003). Critically, NFB was found to restore both parameters towards control-group values, thereby reversing previous significant differences (H-exponent: Hotelling's $T^2 = 40.5$, n.s.; Alpha amplitude: Hotelling's $T^2 = 45.9$, n.s.). Thus post-neurofeedback, dynamical (i.e., LRTCs) and static (i.e., *mean* amplitude) measures of alpha oscillations exhibited a “rebound” from abnormal lows, so that anomalous baseline differences were restored towards healthy population levels.




Figure 2: Pre (T₁) and post (T₂) resting-state changes in post-traumatic stress disorder (PTSD), in comparison to a healthy control group (CONTROLS) **(A)** Topography of Hurst-exponents in the alpha-band reflecting LRTCs; **(B)** mean alpha Hurst-exponents; **(C)** Topography of alpha amplitudes; and **(D)** mean alpha amplitudes of individual PTSD patients (blue circles) and CONTROL subjects (red diamonds). Grand averages indicate all subjects, error-bars denote \pm SEM. * significant difference at $p < 0.05$; n.s. not significant.

Changes in hyperarousal in PTSD patients

The Clinician Administered PTSD Scale (CAPS) indicated that the PTSD group met the necessary criteria for trait hyperarousal. A comparison of state arousal (Thayer Activation Checklist) pre-to-post neurofeedback revealed a significant decrease in arousal in the PTSD group ($t = -2.72$; $p < 0.05$) after NFB training. Crucially, as depicted in Fig 3, individual decreases in arousal score were significantly correlated with increases in Hurst exponent ($R = -0.45$, $p < 0.05$) at the feedback channel Pz. A similar (marginally-significant) trend was identified for mean alpha amplitude ($R = -0.39$, $p = 0.08$).




Figure 3: Changes in state arousal vs. LRTCs in PTSD patients. State arousal (pre-to-post neurofeedback training) was estimated from the Thayer Activation state questionnaire. LRTCs are denoted by alpha-band Hurst exponents calculated at the parietal feedback site (channel Pz). * significant Pearson correlation at $p < 0.05$.

Relationship between LRTCs and mean alpha amplitude

We reasoned that a large span of individual data could prove crucial for uncovering a potential non-linear relationship between LRTCs and oscillation amplitude. Hence, we collected and analysed an additional sample comprising of eyes-closed recordings (32 healthy subjects, *before* and *after* neurofeedback). Combining data from all experiments ($n=123$ subjects) on a single scatter-plot, we conducted a regression analysis to estimate the best curve-fit. As illustrated in Fig 4, a quadratic fit provided the best model ($R^2 = 0.21$, $p < 0.05$), explaining around 20% of the total variance. This quadratic relationship held for *absolute* alpha amplitude ($R^2 = 0.24$), indicating a superior fit ($Z = 2.03$, $p < 0.05$) versus a linear relationship ($R^2 = 0.11$), which would result from a simple increase in signal-to-noise arising with higher alpha amplitudes (Linkenkaer-Hansen, 2001). The statistical presence of an inverted-U relationship is striking, revealing a middle zone of maximal LRTCs, and attenuated LRTCs at higher and lower alpha extremes. Considering reports that alpha amplitude positively covaries with the synchrony of local neuronal populations (Bollimunta et al., 2011; Haegens et al., 2011; Musall et al., 2014), this seems to indicate that both hypo- and hyper-synchronization could be associated with a breakdown of long-range dependence, consistent with theoretical predictions about criticality (Poil et al., 2012; Botcharova et al., 2014; Tomen et al., 2014; Shanahan, 2015).




Figure 4: Relationship between individual values of global Hurst exponent and global alpha amplitude (i.e., mean of all channels). Colours represent different subject groups: real-neurofeedback (NFB), sham-feedback (SHAM), post-traumatic stress disorder (PTSD), PTSD healthy control subjects (CONTROLS), extra eyes closed subjects (EYES CLOSED). Pre (T_1) and post (T_2) neurofeedback values are plotted for all subjects (except T_1 for CONTROLS only). Dotted line denotes quadratic fit, with upper and lower lines the 95% confidence intervals * significant at $p < 0.05$.

Alpha training dynamics during neurofeedback

To confirm that closed-loop training (neurofeedback) resulted in differential changes of the controlled parameter (absolute alpha amplitude), we report the 30-minute training session dynamics for the 3 experimental groups: NFB, SHAM, and PTSD. As depicted in Fig S3, and consistent with the neurofeedback protocol, NFB and PTSD groups exhibited a more sustained reduction of alpha amplitudes compared to the SHAM group. For the feedback channel Pz (Fig S3A,) the alpha amplitude time-course significantly differed between NFB vs SHAM groups (Group x Time interaction: $F_{10, 380} = 3.4$, $p < 0.05$). Similarly, a significant difference was observed between PTSD vs SHAM groups ($F_{10, 390} = 2.7$, $p < 0.05$). As seen in Fig S3B, analogous training dynamics were found on the global level, averaging over all channels.




Figure S3: Temporal evolution of alpha amplitude *during* feedback training, for real-neurofeedback healthy subjects (NFB), sham-feedback healthy subjects (SHAM), and post-traumatic stress disorder patients (PTSD). *Rest* represents the initial 3-minute resting-state recording (i.e., T_1). The subsequent feedback training was subdivided into 10 periods (3 – 30 minutes). **(A)** absolute alpha amplitude at the feedback site (channel Pz); **(B)** absolute alpha amplitude globally (i.e., mean of all channels).

Discussion

Overall, our findings demonstrate that sustained control of brain activity, via neurofeedback (NFB), may induce changes in the scale-free dynamics of spontaneous brain oscillations, evidenced by increased critical exponent(s) of electroencephalogram (EEG) long-range temporal correlations (LRTCs). Specifically, a significant post-neurofeedback increase in resting-state alpha-band LRTCs was observed in a group of healthy adults, as compared to a sham control group that received false feedback (Fig. 1). NFB thereby appears to have induced a plastic ‘re-tuning’ of spontaneous neuronal dynamics, consisting of a tendency for alpha oscillations to remain alternatively high and low in amplitude for a longer duration of time (Hardstone et al., 2012). Fascinatingly, this result was reproduced in patients with post-traumatic stress disorder (PTSD), where abnormal alpha rhythm dynamics (LRTCs) normalized towards values seen in the healthy population (Fig. 2). Here, the inter-individual degree of LRTC re-organization correlated with reductions in self-reported hyperarousal, which significantly decreased at the group level after NFB.

Potential mechanism of LRTC modulation and its relationship to brain function

A question that naturally arises is how scale-free oscillations are linked to brain function and behaviour. Recent studies report a temporally-direct correlation between LRTC exponents of cortical oscillations and those of individual performance errors (e.g., sensory detection (Palva et al., 2013) and time-estimation (Smit et al., 2013)). Cortical LRTCs are known to increase during development (Smit et al., 2011), and are more attenuated ('random') in a number of brain disorders, including depression (Linkenkaer-Hansen et al., 2005), schizophrenia (Nikulin et al., 2012), Alzheimer's (Montez et al., 2009), and autism (Lai et al., 2010). Patients with depression also express reduced LRTCs in their sleep (Leistedt et al., 2007b), which is interestingly reversed during states of symptom remission (Leistedt et al., 2007a). Hence, self-report data from our PTSD sample (Fig. 3) corroborate observations that normalization of LRTCs appears to track state-related improvements of key psychiatric symptoms. A more provocative implication is that such neuromodulation is able to occur at remarkably rapid timescales (i.e. after only 30 minutes of training), and do so endogenously, without the need of external agents (e.g. pharmaceuticals, electrostimulation). This points to the existence of residual and/or homeostatic plasticity mechanisms in the brain (Marder and Goaillard, 2006) that may be capable of reversing long-term pathological brain activity. Our results thus pave the way for future work investigating whether these effects can maintain themselves following multiple sessions of NFB training, paralleling evidence of long-term changes in EEG amplitude (Becerra et al., 2006; Gevensleben et al., 2009).

LRTCs may also be over-pronounced, as has been reported in epilepsy (Parish et al., 2004; Monto et al., 2007). Here, down-regulating the excitation to inhibition (E/I) balance via GABAergic agonists seems to dampen the highly-correlated oscillations of epileptogenic cortex (Monto et al., 2007). This, together with evidence from computational modelling (Poil et al., 2012), tentatively links LRTC changes to fluctuations in E/I balance. Interestingly, we have previously shown that reducing alpha amplitude via NFB may lead to a plastic increase in cortical excitability, as well as a decrease in intracortical inhibition (Ros et al., 2010). It is plausible that a similar neurophysiological mechanism may be at work here, whereby more sustained (i.e., correlated) oscillations could be subserved by a stronger cortical excitatory drive following NFB (Poil et al., 2012). Yet, the contrary facet remains unresolved: could NFB be equally used to reduce pathologically-elevated LRTCs? Here, it is noteworthy that promising studies of NFB in epilepsy already exist (Sterman and Egner, 2006; Strehl et al., 2014).

Long-range dependence versus oscillation amplitude: a hidden link?

The observation of an apparent inverted-U relationship between LRTC magnitude and mean alpha amplitude (Fig 4) is reassuring as it firstly confirms that LRTC changes cannot be ascribed to a trivial increase in signal-to-noise ratio, as may be expected with ever-greater levels of oscillation amplitude. Secondly, depending on the location of sampled data on the curve, the inverted-U shape provides a parsimonious explanation for the reported differences in association between LRTC and oscillation amplitude: i.e., positive (left-side) (Poil et al., 2011), null (middle) (Linkenkaer-Hansen, 2001), or negative (right-side) (Linkenkaer-Hansen, 2001). Intriguingly, this finding also suggests there may be a physiologically limited range of oscillation amplitudes that are associated with maximal LRTCs. We were initially perplexed by this result, but were intrigued when it was juxtaposed with a theoretical framework suggesting a bell-shaped relationship between complexity and synchronization (Tononi et al., 1998). According to this perspective, "true complexity" (as opposed

to basic entropy) is maximal at intermediate states between high synchronization (order) and low synchronization (disorder). Here, alpha amplitude itself is known to be positively associated with the synchronization of local neuronal populations (Bollimunta et al., 2011; Haegens et al., 2011; Musall et al., 2014). After this transformation, our data appear consistent with the framework of Tononi et al (Tononi et al., 1998) when considering LRTCs as a substitute measure of dynamical complexity: as once oscillations shift to excessively desynchronised (more disorderly) or synchronized (more orderly) levels there is a parallel shift of LRTCs towards randomness, reflecting a decrease in variability (shallower slope of critical exponent, i.e., dynamic range) and long-range dependence (i.e., memory), in line with computational models (Poil et al., 2012). Not incidentally (Tagliazucchi and Chialvo, 2013), critical phenomena such as long-range correlations are found near order-to-disorder transitions (Chialvo, 2010; Botcharova et al., 2014; Hesse and Gross, 2014; Shanahan, 2015), where dynamic range and memory are maximized, both features favourable for information processing (Shew and Plenz, 2012). This could be a reason why, in conjunction with deviant LRTCs (Stam et al., 2005; Poil et al., 2011), signatures of hyper- and hypo-synchronization frequently feature in brain disorders (Coburn et al., 2006; Uhlhaas and Singer, 2006). It may therefore be possible that abnormal oscillation amplitudes and LRTCs share a common pathophysiological mechanism (Poil et al., 2008, 2011; Zhigalov et al., 2015). According to our own data, PTSD exhibits both pathologically reduced alpha amplitude and LRTCs, which are normalized in tandem post-NFB. Hence, by means of a sustained modulation of oscillation amplitude, in line with conventional approaches (Lubar, 1997; Heinrich et al., 2007), NFB might concurrently reinstate a self-tuning of more complex temporal dynamics (i.e. LRTCs, which are scale-free and intractable as a real-time parameter for feedback).

The potential utility of neurofeedback in a spectrum of brain disorders

We have shown here the capability of NFB as a neuromodulatory tool in healthy participants as well those with PTSD, and have previously put forward a conceptual framework how such 'endogenous training' of neuronal oscillations might help normalize brain dynamics in a range of psychiatric and neurological conditions (Ros et al., 2014). Although the current study was limited to exploring the effects induced by a single training session, earlier investigations have demonstrated a robust long-term impact of repeated NFB sessions on cortical oscillations (Ros et al., 2014). Given that numerous NFB studies report normalization of targeted EEG amplitudes following long-term treatment (e.g. attentional-deficit hyperactivity disorder (Arns et al., 2009; Gevensleben et al., 2009), tinnitus (Hartmann et al., 2013), and learning disability (Becerra et al., 2006), we hypothesize whether this may have also restored normal LRTCs in these disorders (see Zhigalov et al., 2015 for their typical distribution across cortical regions). Moreover, LRTC normalization could potentially underpin the clinical improvements observed following other types of therapy, whether they be endogenous (e.g. meditation (Lomas et al., 2015)) or exogenous (e.g. transcranial alternating current stimulation (Vossen et al., 2014)); seeing that many of these treatments are also found to modulate EEG amplitude. For example, transcranial magnetic stimulation (rTMS) in schizophrenia has been reported to decrease negative symptoms in proportion to post-treatment increases in alpha amplitude (Jin et al., 2006), while recovery from severe brain injury was found to be associated with reductions of theta amplitude after pharmacotherapy (Williams et al., 2013). Should a unifying pathophysiological mechanism between LRTCs and oscillation amplitude be confirmed in future work, it would underline the scientific rationale for harnessing the unique safety-profile of NFB in a potentially much wider spectrum of clinical disorders, where it is demonstrating early promise -

including schizophrenia (Surmeli et al., 2012), major depression (Escolano et al., 2014), obsessive-compulsive disorder (Kopřivová et al., 2013), insomnia (Schabus et al., 2014), autism (Friedrich et al., 2015) and stroke (Várkuti et al., 2013; Young et al., 2014).

Collectively, our findings establish novel empirical and mechanistic evidence for an intrinsic 'self-tuning' of scale-free spontaneous oscillations following closed-loop brain training, and directly suggest that NFB could find valuable therapeutic applications in disorders with perturbed LRTCs, such as schizophrenia, major depression, and epilepsy (Linkenkaer-Hansen et al., 2005; Monto et al., 2007; Nikulin et al., 2012).

Acknowledgements

We thank The Foundation for Neurofeedback and Applied Neuroscience for generously providing the neurofeedback hardware and software. We are also grateful to Claire Braboszcz for helpful comments, as well as to Abele Michela, Suzy Southwell, Stephanie Nevill, Melody Chow, and Nancy Mazza for their assistance with data collection. This study was supported by grants from the EU Marie-Curie COFUND program, Lawson Health Research Institute and Canadian Institute for Military & Veteran Health Research.

References

- Angelakis E, Stathopoulou S, Frymiare JL, Green DL, Lubar JF, Kounios J (2007) EEG neurofeedback: a brief overview and an example of peak alpha frequency training for cognitive enhancement in the elderly. *Clin Neuropsychol* 21:110–129.
- Arns M, de Ridder S, Strehl U, Breteler M, Coenen A (2009) Efficacy of neurofeedback treatment in ADHD: the effects on inattention, impulsivity and hyperactivity: a meta-analysis. *Clin EEG Neurosci* 40:180–189.
- Bak P, Tang C, Wiesenfeld K (1987) Self-organized criticality: An explanation of the 1/f noise. *Phys Rev Lett* 59:381–384.
- Becerra J, Fernández T, Harmony T, Caballero MI, García F, Fernández-Bouzas A, Santiago-Rodríguez E, Prado-Alcalá RA (2006) Follow-up study of learning-disabled children treated with neurofeedback or placebo. *Clin EEG Neurosci* 37:198–203.
- Bollimunta A, Mo J, Schroeder CE, Ding M (2011) Neuronal mechanisms and attentional modulation of corticothalamic α oscillations. *J Neurosci* 31:4935–4943.
- Botcharova M, Farmer S, Berthouze L (2014) Markers of criticality in phase synchronisation. *arXiv Prepr arXiv14045774* 8:1–24.
- Brunner C, Scherer R, Graimann B, Supp G, Pfurtscheller G (2006) Online control of a brain-computer interface using phase synchronization. *IEEE Trans Biomed Eng* 53:2501–2506.
- Chialvo DR (2010) Emergent complex neural dynamics. *Nat Phys* 6:744–750.
- Churchill NW, Cimprich B, Askren MK, Reuter-Lorenz PA, Jung MS, Peltier S, Berman MG (2015) Scale-free brain dynamics under physical and psychological distress: pre-treatment effects in women diagnosed with breast cancer. *Hum Brain Mapp* 36:1077–1092.
- Coburn KL, Lauterbach EC, Boutros NN, Black KJ, Arciniegas DB, Coffey CE (2006) The value of quantitative electroencephalography in clinical psychiatry: a report by the Committee on Research of the American Neuropsychiatric Association. *J Neuropsychiatry Clin Neurosci* 18:460–500.

- Collins PY et al. (2011) Grand challenges in global mental health. *Nature* 475:27–30.
- Escolano C, Navarro-Gil M, Garcia-Campayo J, Congedo M, De Ridder D, Minguez J (2014) A controlled study on the cognitive effect of alpha neurofeedback training in patients with major depressive disorder. *Front Behav Neurosci* 8:296.
- Friedrich EVC, Sivanathan A, Lim T, Suttie N, Louchart S, Pillen S, Pineda JA (2015) An Effective Neurofeedback Intervention to Improve Social Interactions in Children with Autism Spectrum Disorder. *J Autism Dev Disord*.
- Gevensleben H, Holl B, Albrecht B, Schlamp D, Kratz O, Studer P, Wangler S, Rothenberger A, Moll GH, Heinrich H (2009) Distinct EEG effects related to neurofeedback training in children with ADHD: A randomized controlled trial. *Int J Psychophysiol* 74:149–157.
- Haegens S, Nacher V, Luna R, Romo R, Jensen O (2011) γ -Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal spiking. *Proc Natl Acad Sci* 108:19377–19382.
- Harang R, Bonnet G, Petzold LR (2012) WAVOS: a MATLAB toolkit for wavelet analysis and visualization of oscillatory systems. *BMC Res Notes* 5:163.
- Hardstone R, Poil S-S, Schiavone G, Jansen R, Nikulin V V, Mansvelder HD, Linkenkaer-Hansen K (2012) Detrended fluctuation analysis: a scale-free view on neuronal oscillations. *Front Physiol* 3:450.
- Hardt J V, Kamiya J (1978) Anxiety change through electroencephalographic alpha feedback seen only in high anxiety subjects. *Science* 201:79–81.
- Hartmann T, Lorenz I, Müller N, Langguth B, Weisz N (2013) The Effects of Neurofeedback on Oscillatory Processes Related to Tinnitus. *Brain Topogr*.
- He BJ, Zempel JM, Snyder AZ, Raichle ME (2010) The temporal structures and functional significance of scale-free brain activity. *Neuron* 66:353–369.
- Heinrich H, Gevensleben H, Strehl U (2007) Annotation: Neurofeedback - Train your brain to train behaviour. *J Child Psychol Psychiatry Allied Discip* 48:3–16.
- Hesse J, Gross T (2014) Self-organized criticality as a fundamental property of neural systems. *Front Syst Neurosci* 8:166.
- Hsu D, Beggs JM (2006) Neuronal avalanches and criticality: A dynamical model for homeostasis. *Neurocomputing* 69:1134–1136.
- Jin Y, Potkin SG, Kemp AS, Huerta ST, Alva G, Thai TM, Carreon D, Bunney WE (2006) Therapeutic effects of individualized alpha frequency transcranial magnetic stimulation (alphaTMS) on the negative symptoms of schizophrenia. *Schizophr Bull* 32:556–561.
- Jokić-Begić N, Begić D (2003) Quantitative electroencephalogram (qEEG) in combat veterans with post-traumatic stress disorder (PTSD). *Nord J Psychiatry* 57:351–355.
- Kamiya J (2011) The First Communications About Operant Conditioning of the EEG. *J Neurother* 15:65–73.
- Kello CT, Brown GD a, Ferrer-i-cancho R, Holden JG, Linkenkaer-Hansen K, Rhodes T, Orden GC Van, Van Orden GC (2010) Scaling laws in cognitive sciences. *Top Cogn Sci* 14:223–232.
- Kitzbichler MG, Bullmore ET (2015) Power Law Scaling in Human and Empty Room MEG Recordings. *PLoS Comput Biol* 11:e1004175.
- Kluetsch RC, Ros T, Théberge J, Frewen P a, Calhoun VD, Schmahl C, Jetly R, Lanius R a (2014) Plastic modulation of PTSD resting-state networks and subjective wellbeing by EEG neurofeedback. *Acta Psychiatr Scand*:1–14.
- Kopřivová J, Congedo M, Raszka M, Praško J, Brunovský M, Horáček J (2013) Prediction of Treatment Response and the Effect of Independent Component Neurofeedback in Obsessive-Compulsive Disorder: A Randomized, Sham-Controlled, Double-Blind Study. *Neuropsychobiology* 67:210–223.

- Lai MC, Lombardo M V., Chakrabarti B, Sadek S a., Pasco G, Wheelwright SJ, Bullmore ET, Baron-Cohen S, Suckling J (2010) A shift to randomness of brain oscillations in people with autism. *Biol Psychiatry* 68:1092–1099.
- Leistedt S, Dumont M, Coumans N, Lanquart JP, Jurysta F, Linkowski P (2007a) The modifications of the long-range temporal correlations of the sleep EEG due to major depressive episode disappear with the status of remission. *Neuroscience* 148:782–793.
- Leistedt S, Dumont M, Lanquart J-P, Jurysta F, Linkowski P (2007b) Characterization of the sleep EEG in acutely depressed men using detrended fluctuation analysis. *Clin Neurophysiol* 118:940–950.
- Linkenkaer-Hansen K (2001) Long-range temporal correlations and scaling behavior in human brain oscillations. *J Neurosci* 21:1370–1377.
- Linkenkaer-Hansen K, Monto S, Rytsälä H, Suominen K, Isometsä E, Kähkönen S (2005) Breakdown of long-range temporal correlations in theta oscillations in patients with major depressive disorder. *J Neurosci* 25:10131–10137.
- Liu X, Ward BD, Binder JR, Li SJ, Hudetz AG (2014) Scale-free functional connectivity of the brain is maintained in anesthetized healthy participants but not in patients with unresponsive wakefulness syndrome Zang Y-F, ed. *PLoS One* 9:e92182.
- Lomas T, Ivtzan I, Fu CHY (2015) A systematic review of the neurophysiology of mindfulness on EEG oscillations. *Neurosci Biobehav Rev*.
- Lubar JF (1997) Neocortical dynamics: implications for understanding the role of neurofeedback and related techniques for the enhancement of attention. *Appl Psychophysiol Biofeedback* 22:111–126.
- Marder E, Goaillard J-M (2006) Variability, compensation and homeostasis in neuron and network function. *Nat Rev Neurosci* 7:563–574.
- Montez T, Poil S-S, Jones BF, Manshanden I, Verbunt JP a, van Dijk BW, Brussaard AB, van Ooyen A, Stam CJ, Scheltens P, Linkenkaer-Hansen K (2009) Altered temporal correlations in parietal alpha and prefrontal theta oscillations in early-stage Alzheimer disease. *Proc Natl Acad Sci U S A* 106:1614–1619.
- Monto S, Vanhatalo S, Holmes MD, Palva JM (2007) Epileptogenic neocortical networks are revealed by abnormal temporal dynamics in seizure-free subdural EEG. *Cereb Cortex* 17:1386–1393.
- Musall S, Von Pfösti V, Rauch A, Logothetis NK, Whittingstall K (2014) Effects of neural synchrony on surface EEG. *Cereb Cortex* 24:1045–1053.
- Nikulin V V., Brismar T (2004) Long-range temporal correlations in alpha and beta oscillations: Effect of arousal level and test-retest reliability. *Clin Neurophysiol* 115:1896–1908.
- Nikulin V V., Jönsson EG, Brismar T (2012) Attenuation of long-range temporal correlations in the amplitude dynamics of alpha and beta neuronal oscillations in patients with schizophrenia. *Neuroimage* 61:162–169.
- Niv S (2013) Clinical efficacy and potential mechanisms of neurofeedback. *Pers Individ Dif* 54:676–686.
- Nolan H, Whelan R, Reilly RB (2010) FASTER: Fully Automated Statistical Thresholding for EEG artifact Rejection. *J Neurosci Methods* 192:152–162.
- Palva JM, Zhigalov A, Hirvonen J, Korhonen O, Linkenkaer-Hansen K, Palva S (2013) Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. *Proc Natl Acad Sci U S A* 110:3585–3590.
- Parish LM, Worrell G a., Cranstoun SD, Stead SM, Pennell P, Litt B (2004) Long-range temporal correlations in epileptogenic and non-epileptogenic human hippocampus. *Neuroscience* 125:1069–1076.
- Poil S-S, Hardstone R, Mansvelder HD, Linkenkaer-Hansen K (2012) Critical-State Dynamics of Avalanches and Oscillations Jointly Emerge from Balanced Excitation/Inhibition in Neuronal Networks. *J Neurosci* 32:9817–9823.
- Poil SS, Jansen R, van Aerde K, Timmerman J, Brussaard AB, Mansvelder HD, Linkenkaer-Hansen K (2011) Fast

- network oscillations invitro exhibit a slow decay of temporal auto-correlations. *Eur J Neurosci* 34:394–403.
- Poil SS, Van Ooyen A, Linkenkaer-Hansen K (2008) Avalanche dynamics of human brain oscillations: Relation to critical branching processes and temporal correlations. *Hum Brain Mapp* 29:770–777.
- Ros T, Baars BJ, Lanius RA, Vuilleumier P (2014) Tuning pathological brain oscillations with neurofeedback: A systems neuroscience framework. *Front Hum Neurosci* 8.
- Ros T, Munneke MAM, Ruge D, Gruzelier J, Rothwell JC (2010) Endogenous control of waking brain rhythms induces neuroplasticity in humans. *Eur J Neurosci* 31:770–778.
- Ros T, Théberge J, Frewen P a., Kluetsch R, Densmore M, Calhoun VD, Lanius R a. (2013) Mind over chatter: Plastic up-regulation of the fMRI salience network directly after EEG neurofeedback. *Neuroimage* 65:324–335.
- Schabus M, Heib DPJ, Lechinger J, Griessenberger H, Klimesch W, Pawlizki A, Kunz AB, Sterman BM, Hoedlmoser K (2014) Enhancing sleep quality and memory in insomnia using instrumental sensorimotor rhythm conditioning. *Biol Psychol* 95:126–134.
- Shanahan M (2015) Local and Global Criticality within Oscillating Networks of Spiking Neurons.
- Shew WL, Plenz D (2012) The Functional Benefits of Criticality in the Cortex. *Neurosci* 19:88–100.
- Smit DJ a, Linkenkaer-Hansen K, de Geus EJC (2013) Long-range temporal correlations in resting-state α oscillations predict human timing-error dynamics. *J Neurosci* 33:11212–11220.
- Smit DJ a., de Geus EJC, van de Nieuwenhuijzen ME, van Beijsterveldt CEM, van Baal GCM, Mansvelter HD, Boomsma DI, Linkenkaer-Hansen K (2011) Scale-Free Modulation of Resting-State Neuronal Oscillations Reflects Prolonged Brain Maturation in Humans. *J Neurosci* 31:13128–13136.
- Stam CJ, Montez T, Jones BF, Rombouts S a RB, Van Der Made Y, Pijnenburg Y a L, Scheltens P (2005) Disturbed fluctuations of resting state EEG synchronization in Alzheimer’s disease. *Clin Neurophysiol* 116:708–715.
- Sterman MB, Egner T (2006) Foundation and practice of neurofeedback for the treatment of epilepsy. *Appl Psychophysiol Biofeedback* 31:21–35.
- Strehl U, Birkle SM, Wörz S, Kotchoubey B (2014) Sustained reduction of seizures in patients with intractable epilepsy after self-regulation training of slow cortical potentials - 10 years after. *Front Hum Neurosci* 8:604.
- Surmeli T, Ertem A, Eralp E, Kos IH (2012) Schizophrenia and the efficacy of qEEG-guided neurofeedback treatment: a clinical case series. *Clin EEG Neurosci* 43:133–144.
- Tagliazucchi E, Chialvo DR (2013) Brain complexity born out of criticality. In: *ArXiv: 1211.0309*, pp 4–13.
- Tagliazucchi E, von Wegner F, Morzelewski A, Brodbeck V, Jahnke K, Laufs H (2013) Breakdown of long-range temporal dependence in default mode and attention networks during deep sleep. *Proc Natl Acad Sci U S A* 110:15419–15424.
- Tomen N, Rotermond D, Ernst U (2014) Marginally subcritical dynamics explain enhanced stimulus discriminability under attention. *Front Syst Neurosci* 8:151.
- Tononi G, Edelman GM, Sporns O (1998) Complexity and coherency: Integrating information in the brain. *Trends Cogn Sci* 2:474–484.
- Tumari SZM, Sudirman R, Ahmad a. H (2013) Selection of a Suitable Wavelet for Cognitive Memory Using Electroencephalograph Signal. *Engineering* 05:15–19.
- Uhlhaas PJ, Singer W (2006) Neural Synchrony in Brain Disorders: Relevance for Cognitive Dysfunctions and Pathophysiology. *Neuron* 52:155–168.
- Van de Ville D, Britz J, Michel CM (2010) EEG microstate sequences in healthy humans at rest reveal scale-free dynamics. *Proc Natl Acad Sci U S A* 107:18179–18184.
- Várkuti B, Guan C, Pan Y, Phua KS, Ang KK, Kuah CWK, Chua K, Ang BT, Birbaumer N, Sitaram R (2013) Resting

state changes in functional connectivity correlate with movement recovery for BCI and robot-assisted upper-extremity training after stroke. *Neurorehabil Neural Repair* 27:53–62.

Vossen A, Gross J, Thut G (2014) Alpha power increase after transcranial alternating current stimulation at alpha-frequency (α -tACS) reflects plastic changes rather than entrainment. *Brain Stimul*.

Williams ST, Conte MM, Goldfine AM, Noirhomme Q, Gosseries O, Thonnard M, Beattie B, Hersh J, Katz DI, Victor JD, Laureys S, Schiff ND (2013) Common resting brain dynamics indicate a possible mechanism underlying zolpidem response in severe brain injury. *Elife* 2013:e01157.

Young BM, Nigogosyan Z, Walton LM, Song J, Nair V a, Grogan SW, Tyler ME, Edwards DF, Caldera K, Sattin J a, Williams JC, Prabhakaran V (2014) Changes in functional brain organization and behavioral correlations after rehabilitative therapy using a brain-computer interface. *Front Neuroeng* 7:26.

Zhigalov A, Arnulfo G, Nobili L, Palva S, Palva JM (2015) Relationship of fast- and slow-timescale neuronal dynamics in human MEG and SEEG. *J Neurosci* 35:5385–5396.

