

# On asymptotic validity of naive inference with an approximate likelihood

Helen Ogden

University of Southampton, UK

## Abstract

Many statistical models have likelihoods which are intractable: it is impossible or too expensive to compute the likelihood exactly. In such settings, a common approach is to replace the likelihood with an approximation, and proceed with inference as if the approximate likelihood were the exact likelihood. In this paper, we describe conditions on the approximate likelihood which guarantee that this naive inference with an approximate likelihood has the same first-order asymptotic properties as inference with the exact likelihood. We investigate the implications of these results for inference using a Laplace approximation to the likelihood in a simple two-level latent variable model, and using reduced dependence approximations to the likelihood in an Ising model on a lattice.

**Keywords:** Intractable likelihood, Ising model, Laplace approximation, Latent variable model

## 1 Introduction

For many models, it is impossible or infeasibly expensive to evaluate the likelihood function, typically because it involves a high-dimensional sum or integral. In such cases, a common approach is to find an approximation  $\tilde{L}(\cdot)$  to the likelihood  $L(\cdot)$ , and to use  $\tilde{L}(\cdot)$  in place of  $L(\cdot)$  to conduct inference about the parameters of the model.

For instance, one could construct a point estimate of the parameters by maximizing the approximate likelihood, and form confidence intervals based on the curvature of the approximated log-likelihood about its maximum. From a Bayesian perspective, an approximate posterior  $\tilde{\pi}(\theta; y) \propto \tilde{L}(\theta; y)\pi(\theta)$  could be formed by substituting the approximate likelihood in place of the exact likelihood.

Such an approach is commonly used in practice. In latent variable models, where the likelihood is an integral over the latent variables, naive inference using a Laplace approximation to the likelihood is used in both maximum likelihood (Pinheiro and Bates, 1995; Bates et al., 2015) and Bayesian (Rue et al., 2009) settings. In Markov random field models, where the likelihood involves an intractable normalizing constant, inference is often conducted by substituting an approximation to this normalizing constant in place of the exact

normalizing constant into the expression for the likelihood (Friel et al., 2009; Tjelmeland and Austad, 2012).

In this paper, we provide conditions under which the naive approach is asymptotically justified. Under these conditions, the approximate maximum likelihood estimator is consistent and has the same asymptotic normal distribution as the exact maximum likelihood estimator, hypothesis tests based on the approximate likelihood remain valid, and in Bayesian analysis the distance between the approximate posterior and the true posterior shrinks to zero.

Douc et al. (2004) show that the approximate maximum likelihood estimator will have the correct asymptotic normal distribution provided that the error in the log-likelihood,

$$\epsilon_n(\theta) = \log \tilde{L}_n(\theta) - \log L_n(\theta),$$

tends in probability to zero as  $n \rightarrow \infty$ , uniformly in  $\theta$ . We argue that this measure is too strict in many practical examples, in which  $\epsilon_n(\theta)$  grows rapidly with  $n$  and yet the inference remains asymptotically valid. Our conditions are based instead on  $\nabla_{\theta} \epsilon_n(\theta)$ , the error in the approximation to the score function.

We provide two examples to demonstrate how the conditions may be used in practice. The first is a simple two-level latent variable model, with  $m_n$  repeated observations for each of  $n$  items. We deduce the rate at which  $m_n$  must grow with  $n$  in order for the Laplace approximation to give asymptotically valid inference. If  $m_n$  grows with  $n$  at a slower rate, the estimator remains consistent, but loses efficiency relative to the exact maximum likelihood estimator, and naively constructed confidence intervals have lower than nominal coverage.

The second example is an Ising model on an  $m \times m$  lattice, with the class of reduced dependence approximations (Friel et al., 2009) used to approximate the likelihood. For parameter values associated with weak dependence, we show that the reduced dependence approximation may be used to obtain asymptotically valid inference at cost polynomial in  $m$ , in contrast with the exponential cost of computing the likelihood exactly.

## 2 Asymptotic validity of approximate likelihood inference

### 2.1 Setup and notation

Consider a sequence of models indexed by  $n$ , with common parameter  $\theta \in \Theta \subseteq \mathbb{R}^p$ . Write  $\ell_n(\theta; y)$  for the log-likelihood given observed data  $y$  under model  $n$ ,  $u_n(\theta; y) = \nabla_{\theta} \ell_n(\theta; y)$  for the corresponding score function and  $J_n(\theta; y) = -\nabla_{\theta}^T \nabla_{\theta} \ell_n(\theta; y)$  for the observed information. We sometimes drop the data  $y$  from the notation for convenience. Suppose that the data were generated from the model for some  $\theta_0 \in \Theta$ , and that as  $n \rightarrow \infty$ , the amount of information provided by the data about the parameter grows at some rate  $r_n$ , such that  $J_n(\theta_0) = O(r_n)$  in probability. This includes the case with  $n$  independent replications as a special case, with  $r_n = n$ , but we also wish to allow for more complex settings. Write  $I(\theta)$  for the Fisher information matrix, chosen such that  $\bar{J}_n(\theta) = r_n^{-1} J_n(\theta) \rightarrow I(\theta)$  in probability.

Let  $\ell_n(\cdot; y)$  be an approximate log-likelihood, which in general may be any function of the parameters  $\theta$ , which will be used in place of  $\ell_n(\cdot; y)$ . Write  $\hat{\theta}_n$

and  $\tilde{\theta}_n$  for the estimators maximizing  $\ell_n(\theta)$  and  $\tilde{\ell}_n(\theta)$  respectively. Suppose that  $\ell_n(\theta)$  and  $\tilde{\ell}_n(\theta)$  are both three times differentiable, and write  $\tilde{u}_n(\theta) = \nabla_{\theta}\tilde{\ell}_n(\theta)$  and  $\tilde{J}_n(\theta) = -\nabla_{\theta}^T\nabla_{\theta}\tilde{\ell}_n(\theta)$  for the approximate score and information.

Write

$$\epsilon_n(\theta) = \tilde{\ell}_n(\theta) - \ell_n(\theta)$$

for the pointwise error in the log-likelihood,

$$\delta_n(\theta) = \|\nabla_{\theta}\epsilon_n(\theta)\| = \|\tilde{u}_n(\theta) - u_n(\theta)\|$$

for the absolute error in the score, and

$$\gamma_n(\theta) = \|\nabla_{\theta}^T\nabla_{\theta}\epsilon_n(\theta)\| = \|J_n(\theta) - \tilde{J}_n(\theta)\|$$

for the absolute error in the observed information matrix. For concreteness, we use the  $L_1$  norms  $\|a\| = \sum_i |a_i|$  for a vector  $a$ , and  $\|A\| = \max_j \{\sum_i |A_{ij}|\}$  for a matrix  $A$ , although the same results hold for any choice of norms.

Write  $\delta_n^{\infty}(S) = \sup_{\theta \in S} \delta_n(\theta)$  for the uniform error in the score over any set  $S \subseteq \Theta$ , and let  $\delta_n^{\infty} = \delta_n^{\infty}(\Theta)$ . Similarly, define  $\gamma_n^{\infty}(S) = \sup_{\theta \in S} \gamma_n(\theta)$  and  $\gamma_n^{\infty} = \gamma_n^{\infty}(\Theta)$ . For any  $\theta_0 \in \Theta$ , write  $B_t(\theta_0) = \{\theta \in \Theta : \|\theta - \theta_0\| \leq t\}$  for the ball of radius  $t$  about  $\theta_0$ .

## 2.2 Approximate maximum likelihood inference

First, we describe sufficient conditions to ensure that  $\tilde{\theta}_n$  is consistent. The proofs of all results are given in the appendix.

We will assume some standard regularity conditions on the model. Writing  $\bar{u}_n(\theta) = r_n^{-1}u_n(\theta)$ , and  $\bar{u}(\theta)$  for the limit as  $n \rightarrow \infty$ , we assume that  $\sup_{\theta \in \Theta} \|\bar{u}_n(\theta) - \bar{u}(\theta)\| \rightarrow 0$  in probability. We assume  $\bar{u}(\cdot)$  is such that, for any  $\epsilon > 0$ ,  $\int_{\theta: d(\theta, \theta_0) \geq \epsilon} \|\bar{u}(\theta)\| > \bar{u}(\theta_0) = 0$ . These conditions are stronger than necessary, and we expect the same result to hold in many other situations where the exact maximum likelihood estimator is consistent.

**Theorem 1.** *Suppose  $\delta_n^{\infty} = o_p(r_n)$  as  $n \rightarrow \infty$ . Then  $\tilde{\theta}_n \rightarrow \theta_0$  in probability, as  $n \rightarrow \infty$ .*

We now give conditions to ensure that  $\tilde{\theta}_n$  retains the same limiting distribution as  $\hat{\theta}_n$ . Since  $\hat{\theta}_n - \theta_0$  is  $O_p(r_n^{-1/2})$ , this is equivalent to finding conditions under which  $\hat{\theta}_n - \tilde{\theta}_n$  is  $o_p(r_n^{-1/2})$ . The following lemma bounds the distance between  $\tilde{\theta}_n$  and  $\hat{\theta}_n$  in terms of the error in the score function near  $\theta_0$ .

**Lemma 1.** *Suppose that  $\delta_n^{\infty} = o_p(r_n)$ , and that there exists  $t > 0$  such that  $\delta_n^{\infty}\{B_t(\theta_0)\} = o_p(a_n)$ . Then  $\tilde{\theta}_n - \hat{\theta}_n = o_p(a_n r_n^{-1})$ .*

Applying Lemma 1 with  $a_n = r_n^{1/2}$  leads directly to the asymptotic normality result.

**Theorem 2.** *Suppose that  $\delta_n^{\infty} = o_p(r_n)$ , and that there exists  $t > 0$  such that  $\delta_n^{\infty}\{B_t(\theta_0)\} = o_p(r_n^{1/2})$ . Then*

$$r_n^{1/2}(\tilde{\theta}_n - \theta_0) \rightarrow N(0, I(\theta_0)^{-1})$$

*in distribution, as  $n \rightarrow \infty$ .*

It is also desirable for hypothesis tests constructed by using the approximate likelihood in place of the exact likelihood to have the correct asymptotic distribution.

Consider testing the hypothesis  $H_0 : \theta \in \Theta_R$ , where  $\Theta_R \subset \Theta$  and  $\dim(\Theta_R) = q$ . Write  $\hat{\theta}_n^R$  for the restricted maximum likelihood estimator, and  $\Lambda_n = 2\{\ell_n(\hat{\theta}_n) - \ell_n(\hat{\theta}_n^R)\}$  for the likelihood ratio statistic. The approximate version of the likelihood ratio test statistic is  $\tilde{\Lambda}_n = 2\{\tilde{\ell}_n(\tilde{\theta}_n) - \tilde{\ell}_n(\tilde{\theta}_n^R)\}$ , where  $\tilde{\theta}_n^R$  is the restricted approximate likelihood estimator.

Under the same conditions that were used to show that  $\tilde{\theta}_n$  has the correct limiting distribution, plus a bound on the error in the information around  $\theta_0$ ,  $\tilde{\Lambda}_n$  is asymptotically equivalent to  $\Lambda_n$  under  $H_0$ .

**Theorem 3.** *Suppose that  $\delta_n^\infty = o_p(r_n)$ , and that there exists  $t > 0$  such that  $\delta_n^\infty\{B_t(\theta_0)\} = o_p(r_n^{1/2})$  and  $\gamma_n^\infty\{B_t(\theta_0)\} = o_p(r_n)$ . Then, under  $H_0$ ,  $\tilde{\Lambda}_n - \Lambda_n = o_p(1)$ .*

The Wald and score test statistics,  $W_n$  and  $S_n$ , are asymptotically equivalent to the likelihood ratio test, so under  $H_0$ , all three statistics have limiting distribution  $\chi_{p-q}^2$ . Under the conditions of Theorem 3,  $I(\theta_0)$  is consistently estimated by  $r_n^{-1}\tilde{J}_n(\tilde{\theta}_n)$ , so the approximate Wald and score test statistics  $\tilde{W}_n$  and  $\tilde{S}_n$  are also asymptotically equivalent to  $\Lambda_n$ .

### 2.3 Approximate Bayesian inference

We now consider the approximate posterior

$$\tilde{\pi}(\theta|y) \propto \tilde{L}_n(\theta; y)\pi(\theta),$$

where we suppose that the prior is such that  $\log \pi(\cdot)$  is three times differentiable. Under the same conditions that were used to show asymptotic correctness of maximum likelihood inference, the total variation distance between the approximate and exact posteriors,

$$d_{TV}\{\tilde{\pi}(\theta|y), \pi(\theta|y)\} = \frac{1}{2} \int_{\Theta} |\tilde{\pi}(\theta|y) - \pi(\theta|y)| d\theta,$$

tends to zero as  $n \rightarrow \infty$ .

**Theorem 4.** *Suppose that  $\delta_n^\infty = o_p(r_n)$ , and that there exists  $t > 0$  such that  $\delta_n^\infty\{B_t(\theta_0)\} = o_p(r_n^{1/2})$  and  $\gamma_n^\infty\{B_t(\theta_0)\} = o_p(r_n)$ . Then*

$$d_{TV}\{\tilde{\pi}(\theta|y), \pi(\theta|y)\} = o_p(1)$$

as  $n \rightarrow \infty$ .

If the Bernstein-von Mises Theorem holds for the exact posterior distribution, under the conditions of Theorem 4, it will also hold for the approximate posterior distribution  $\tilde{\pi}(\theta|y)$ . In that case, credible regions formed from the approximate posterior distribution will also be valid confidence sets.

## 2.4 Adjusted approximate likelihood inference

If Lemma 1 holds for some  $a_n > r_n^{1/2}$ , then  $\tilde{\theta}_n$  will still be a consistent estimator, but might not have the same limiting distribution as  $\hat{\theta}_n$ . The approximate likelihood may still be useful in practice, provided that the inference is adjusted accordingly.

The sandwich information matrix (Godambe, 1960) is  $G_n(\theta) = I_n(\theta)H_n(\theta)^{-1}I_n(\theta)$  where  $H_n(\theta) = \text{var}\{\nabla_{\theta}\tilde{u}_n(\theta)\}$  and  $I_n(\theta) = E\{\tilde{J}_n(\theta)\}$ . Under suitable regularity conditions,

$$s_n^{1/2}(\tilde{\theta}_n - \theta_0) \rightarrow N(0, \tilde{G}(\theta_0)^{-1})$$

in distribution, where  $s_n$  is the rate of convergence of  $\tilde{\theta}_n$ , chosen such that  $G_n(\theta_0) = O(s_n)$ , and  $\tilde{G}(\theta) = \lim_{n \rightarrow \infty} s_n^{-1}G_n(\theta)$ .

Composite likelihood estimators (Lindsay, 1988) also have this type of asymptotic behaviour, and many methods which have been proposed to adjust inference using a composite likelihood could also be used to adjust the inference with an approximate likelihood, provided that  $\tilde{\theta}_n$  is a consistent estimator. For example, Varin et al. (2011) describe various methods to approximate the variance of a composite likelihood estimator, and list some modification to the composite likelihood ratio test statistic designed to ensure that the resulting test statistic has approximately  $\chi_{p-q}^2$  distribution. From a Bayesian perspective, the adjustments proposed by Pauli et al. (2011) and Ribatet et al. (2012) to posterior distributions based on composite likelihoods could also be used in the context of approximate likelihood inference.

## 3 Examples

### 3.1 Laplace approximation to the likelihood in a simple latent variable model

#### 3.1.1 A two-level model

Suppose  $Y_i \sim \text{Binomial}(m, p_i)$ , where  $\text{logit}(p_i) = b_i$  and  $b_i \sim N(0, \theta^2)$ , for  $i = 1, \dots, n$ . The likelihood is

$$\begin{aligned} L_n(\theta) &= \int_{\mathbb{R}^n} \prod_{i=1}^n \{\text{logit}^{-1}(b_i)\}^{y_i} \{1 - \text{logit}^{-1}(b_i)\}^{m-y_i} \phi(b_i; 0, \theta) db \\ &= \prod_{i=1}^n L_{(i)}(\theta), \end{aligned}$$

where

$$L_{(i)}(\theta) = \int_{-\infty}^{\infty} \{\text{logit}^{-1}(b_i)\}^{y_i} \{1 - \text{logit}^{-1}(b_i)\}^{m_n - y_i} \phi(b_i; 0, \theta) db_i$$

and  $\phi(\cdot; \mu, \sigma)$  is the  $N(\mu, \sigma^2)$  density function.

If we take a Laplace approximation  $\tilde{L}_n(\theta)$  to the likelihood, it is intuitively clear that  $m = m_n$  will have to grow with  $n$  to give valid inference as  $n \rightarrow \infty$ . It is less obvious whether any choice of  $m_n$  that grows with  $n$  will give valid inference, or whether  $m_n$  needs to grow with  $n$  at some minimum rate. Rue et al. (2009) suggest that any  $m_n$  which grows with  $n$  will suffice, conjecturing that

“the error rate” is [number of latent variables / number of observations], although they note that this rate is not established rigorously. In this case, the error rate refers to the error in approximating  $\pi(\theta|y)$  with  $\tilde{\pi}(\theta|y)$ , found by using a Laplace approximation to the likelihood. The Integrated Nested Laplace Approximations proposed by Rue et al. (2009) are based on this  $\tilde{\pi}(\theta|y)$ , with further approximations used to approximate the marginal posterior distribution of each component of  $\theta$ , if  $p > 1$ . In this example  $\theta$  is a scalar, so the Integrated Nested Laplace Approximation to the posterior distribution is precisely  $\tilde{\pi}(\theta|y)$ .

### 3.1.2 Theoretical analysis

The factorization of the likelihood allows us to study the error for each item  $\epsilon_{(i)}(\theta) = \ell_{(i)}(\theta) - \ell_{(i)}(\theta)$  separately, and then combine the errors. In the supplementary materials, we show that for each fixed  $\theta \in \Theta$ ,

$$\delta_{(i)}(\theta) = \|\nabla_{\theta} \epsilon_{(i)}(\theta)\| = O_p(m_n^{-2}),$$

so  $\delta_n(\theta) \leq \sum_{i=1}^n \delta_{(i)}(\theta) = O_p(nm_n^{-2})$ .

However, the conditions are in terms of uniform rather than pointwise errors. Since  $\delta_{(i)}(\theta)$  is maximized at a point  $\theta^* = O_p(m_n^{-1/2})$ , and decreasing for all  $\theta > \theta^*$ ,

$$\delta_{(i)}^{\infty} = \delta_{(i)}(\theta^*) = O_p(m_n^{-1/2}),$$

so  $\delta_n^{\infty} = O_p(nm_n^{-1/2})$ .

The amount of information that the data provides about  $\theta$  is bounded for fixed  $n$  as  $m_n \rightarrow \infty$ , by the available information on  $\theta$  given the value of each  $b_i$ . So if  $m_n \rightarrow \infty$  as  $n \rightarrow \infty$ , then  $r_n = O(n)$ , and since  $\delta_n^{\infty} = o_p(r_n)$ ,  $\hat{\theta}_n$  will be consistent.

Provided that  $\theta_0 \neq 0$ , choose any fixed  $t \in (0, \theta_0)$ . Then, for sufficiently large  $n$ ,

$$\delta_n^{\infty} \{B_t(\theta_0)\} = \delta_n(\theta_0 - t) = O_p(nm_n^{-2})$$

and

$$\gamma_n^{\infty} \{B_t(\theta_0)\} = \gamma_n(\theta_0 - t) = O_p(nm_n^{-2}).$$

If  $m_n$  grows at a rate faster than  $n^{1/4}$ , then  $\delta_n^{\infty} \{B_t(\theta_0)\} = o_p(n^{1/2})$  and  $\gamma_n^{\infty} \{B_t(\theta_0)\} = o_p(n)$ , so the Laplace approximation to the likelihood will give first-order correct inference.

### 3.1.3 Numerical demonstration

To verify these results numerically, we simulate 10000 realizations from the model with  $\theta_0 = 0.5$ , various values of  $n$  between 1000 and 10000, and  $m_n = \min\{1, 5 + 4(n^a - 1000^a)\}$ , for  $a = 0.2, 0.25$  or  $0.3$ . The three choices of  $m_n$  are shown in Figure 1a.

A very accurate approximation to the likelihood may be obtained by using adaptive Gaussian quadrature with 20 quadrature points to approximate each of the univariate integrals  $L_{(i)}(\theta)$ . We use this approximation as a proxy for the exact likelihood  $L(\theta)$ .

As  $n \rightarrow \infty$ , we have  $r_n = O(n)$ , but for smaller sample sizes  $E(\|J_n(\hat{\theta}_n)\|)$  still grows with  $m_n$ .  $E(\|J_n(\hat{\theta}_n)\|)$  may be approximated by  $\hat{r}_n = \hat{E}(\|J_n(\hat{\theta}_n)\|)$ , where  $\hat{E}(\cdot)$  the sample mean over the 10000 realizations. The functional form




Figure 1: Comparison of exact and approximate likelihood inference, for a two-level model with  $m_n$  observations on each of  $n$  items, where  $m_n = \min\{1, 5 + 4(n^a - 1000^a)\}$ , for  $a = 0.2$  (dashed lines),  $0.25$  (solid lines) and  $0.3$  (dotted lines).  $\text{RMSE}(\cdot)$  denotes root mean squared error.

of  $m_n$  was chosen to make  $\hat{r}_n^{-1/2}\delta_n(\theta_0)$  approximately constant in the  $a = 0.25$  case, as shown in Figure 1b. The same quantity grows with  $n$  for  $a = 0.2$  and shrinks with  $n$  for  $a = 0.3$ .

Figure 1c shows the root mean squared error in the Laplace estimator, and as expected, the estimator is consistent for all three choices of  $a$ . The root mean squared error of the Laplace estimator divided by that of the maximum likelihood estimator, shown in Figure 1d, grows in the  $a = 0.2$  case, stays approximately constant if  $a = 0.25$ , and shrinks towards 1 if  $a = 0.3$ .

Figure 1e shows the actual coverage of likelihood ratio type confidence intervals for  $\theta$ , of nominal 90% coverage. The upper three lines show the coverage of the intervals constructed using the exact likelihood, which have very close to nominal coverage for each  $a$ . The lower three lines show the coverage of the approximate likelihood intervals, which decreases with  $n$  for  $a = 0.2$ , and increases towards the nominal 90% level for  $a = 0.3$ .

Figure 1f shows the total variation distance between  $\tilde{\pi}(\theta|y)$  and  $\pi(\theta|y)$ , with prior  $\pi(\theta) \propto 1/\theta$ . The distance between the approximate posterior and the exact posterior grows in the  $a = 0.2$  case, stays approximately constant if  $a = 0.25$ , and shrinks towards zero if  $a = 0.3$ . The behaviour for  $a = 0.2$  refutes the conjecture of Rue et al. (2009) that the error in the approximate posterior distribution should shrink to zero provided that  $m_n$  grows with  $n$ .

## 3.2 Reduced dependence approximation to the likelihood in an Ising model

### 3.2.1 An Ising model

We consider a simple Ising model for  $n = rc$  variables  $y_i \in \{-1, 1\}$ , arranged on an  $r \times c$  lattice, with parameters  $\theta = (\alpha, \beta)$ , so that

$$pr(Y = y; \theta) = \{Z_{r,c}(\theta)\}^{-1} \exp\{\alpha V_0(y) + \beta V_1(y)\},$$

where  $V_0(y) = \sum_i y_i$ , and  $V_1(y) = \sum_{i \sim j} y_i y_j$ . Here  $i \sim j$  indicates that  $i$  and  $j$  have an edge between them in the lattice, and  $Z_{r,c}(\theta) = \sum_{y \in \{-1,1\}^n} \exp\{\alpha V_0(y) + \beta V_1(y)\}$  is a normalizing constant. The likelihood function  $L(\theta; y) = pr(Y = y; \theta)$  depends on  $Z_{r,c}(\theta)$ , and it is the computation of this normalizing constant which makes evaluation of the likelihood function difficult. By using variable elimination (e.g. Jordan, 2004),  $Z_{r,c}(\theta)$  may be computed at cost  $O(rc2^{\min\{r,c\}})$ , which remains infeasibly expensive if both  $r$  and  $c$  are large.

### 3.2.2 Reduced dependence approximations

Many methods for approximating  $Z_{r,c}(\theta)$  have been proposed. We will study properties of inference using the reduced dependence approximations introduced by Friel et al. (2009), a family of approximations controlled by a positive integer tuning parameter, which we call  $k$ . The approximation for fixed  $k$  is  $\tilde{Z}_{r,c}^{(k)}(\theta) = \{Z_{k,c}(\theta)\}^{r-k+1} / \{Z_{k-1,c}(\theta)\}^{r-k}$ .

We consider the case  $r = c = m$ , using a reduced dependence approximation at level  $k$  to approximate to the likelihood, giving  $\tilde{L}_m^{(k)}(\beta) = \{\tilde{Z}_{m,m}^{(k)}(\theta)\}^{-1} \exp\{\alpha V_0(y) + \beta V_1(y)\}$ . The exact likelihood may be computed at cost  $O(m^2 2^m)$ , and the reduced dependence approximation at level  $k$  at cost  $O(m^2 + km2^k)$ . The aim



is to understand how  $k = k_m$  should vary with  $m$  to give asymptotically valid inference as  $m \rightarrow \infty$ . The error in the log-likelihood,

$$\epsilon_m^{(k)}(\beta) = \tilde{\ell}_m^{(k)}(\beta) - \ell_m(\beta) = \log Z_{m,m}(\beta) - \log \tilde{Z}_{m,m}^{(k)}(\beta),$$

does not depend on the data  $y$ , so we do not need to consider the statements in probability: all of the errors are deterministic in this case.

### 3.2.3 A special case where the exact likelihood is available

If  $\alpha = 0$ , and the lattice has periodic boundary conditions, so that the top row of variables are joined to the bottom row, and the left row joined to the right, Kaufman (1949) provides a relatively simple expression for  $Z_{r,c}(0, \beta)$ , so that it is possible to compute the likelihood exactly, even for large lattices.

We restrict the parameter space to  $\alpha = 0$ , and assume  $\beta \in [0, 0.43]$ . This guarantees that  $\beta < \beta_c = \log(1 + \sqrt{2})/2 \approx 0.44$ , where  $\beta_c$  is a critical value at which the behaviour of the Ising model suddenly changes, so that for  $\beta > \beta_c$  large areas of all plus ones or all minus ones are observed. If  $\beta_0 = \beta_c$ , the maximum likelihood estimator may not have a normal limiting distribution, so our results do not apply to this case.

The information provided by the data about  $\beta$  grows at rate  $r_m = m^2$ . In the supplementary material, we show that if  $k \rightarrow \infty$  as  $m \rightarrow \infty$ , then  $\delta_m^{(k)}(\beta) = O\{m^2 k \exp(-b_\beta k)\} + o(1)$ , where  $b_\beta = 2 \cosh^{-1}\{-1 + \cosh(2\beta)^2 / \sinh(2\beta)\}$ .

For any choice of  $k$  which grows with  $m$ ,  $\sup_{\beta \in [0, 0.43]} \delta_m^{(k)}(\beta) = o(m^2)$ , so the approximate likelihood estimator will be consistent.


In order to meet the conditions of Theorem 2,  $k = k_m$  should be chosen so that  $\delta_m^{(k)}\{B_t(\beta_0)\} = o(m)$ . For any  $t < \beta_c - \beta_0$ , for sufficiently large  $m$ ,  $\delta_m^{(k)}\{B_t(\beta_0)\} = \delta_m^{(k)}(\beta_0 + t) = O\{m^2 k \exp(-b_{\beta_0+t} k)\} + o(1)$ . Since  $b_\beta$  is a continuous function of  $\beta$ , any  $k$  such that  $m^2 k \exp(-b_{\beta_0} k) = o(m)$  will meet this condition for sufficiently small  $t$ . This may be achieved by taking  $k_m = c_{\beta_0} \log m$ , for any  $c_{\beta_0} > b_{\beta_0}^{-1}$ . Figure 2a shows how  $b_\beta^{-1}$  varies with  $\beta$ .

For any  $k \rightarrow \infty$  as  $m \rightarrow \infty$ ,  $\gamma_m^{(k)}\{B_t(\beta_0)\} = o(m^2)$ , so the reduced dependence approximation to the likelihood with  $k_m = c_{\beta_0} \log m$  will provide asymptotically valid inference for any  $c_{\beta_0} > b_{\beta_0}^{-1}$ . The cost of computing this approximation is  $O(m \log m m^{c_{\beta_0} \log 2}) < O(m^{c_{\beta_0} \log 2 + 2})$ , polynomial in the size of the model, but increasing rapidly as  $\beta_0$  approaches the critical value.

### 3.2.4 Extension to the more general case

If  $\alpha \neq 0$ , no simple expression for  $Z_{r,c}(\alpha, \beta)$  is known. Instead, we investigate the behaviour of  $m^{-1} \delta_m^{(k)}(\alpha, \beta)$  numerically, by using  $\tilde{\ell}_m^{(K)}$  for some large  $K$  as a proxy for the exact log-likelihood. Care is needed to choose  $K$  sufficiently large to ensure that the results are not sensitive to the choice of  $K$ .

Taking  $K = 16$ , a contour plot of this approximation to  $\log\{m^{-1} \delta_m^{(k)}(0.1, 0.3)\}$  against  $\log m$  and  $k$  is shown in Figure 2b, for  $k = 2, \dots, 12$ ,  $m = 50, \dots, 300$ . The same plots with  $K = 15$  (dashed) and  $K = 14$  (dot-dash) are overlaid, and the differences are barely visible at this scale. Given this stability, it seems reasonable to assume a contour plot of the exact  $\log\{m^{-1} \delta_m^{(k)}(0.1, 0.3)\}$  would look very similar to this. To obtain asymptotically valid inference,  $k_m$  should



(a)  $b_\beta^{-1}$ , for a restricted Ising model with  $\alpha = 0$ . (b) Contour plot of  $\log\{m^{-1}\delta_m^{(k)}(0.1, 0.3)\}$  against  $\log m$  and  $k$ .

Figure 2: Inference for an Ising model with a reduced dependence approximation to the likelihood.

be chosen so that this rescaled error shrinks with  $m$ , which seems to occur if  $k_m$  grows at rate  $c(0.1, 0.3) \log m$ , for  $c(0.1, 0.3)$  larger than about 1.5. This pattern of behaviour is very similar to the  $\alpha = 0$  case. In both cases, reduced dependence approximations with an appropriate choice of  $k$  will give asymptotically valid inference at cost polynomial in the size of the model, in contrast with the exponential cost of computing the likelihood exactly.

## 4 Discussion

The results obtained here can also be applied to other approximations to the likelihood and to other types of model. The conditions on the approximate likelihood, such as showing that  $\delta_n^\infty\{B_t(\theta_0)\} = o_p(r_n)$ , may be difficult to verify in practice, as the exact likelihood is assumed unavailable. If the approximation to the likelihood is a truncation of a series expansion for the exact likelihood, as is the case for the Laplace approximation, the conditions can be checked by examining the contribution from higher-order terms in the expansion. In other cases, it may only be possible to investigate the size of the errors numerically, by using a more accurate and expensive approximation to the likelihood as a proxy for the exact likelihood, as described in Section 3.2.4 for the Ising model example.

Many approximation methods have a tuning parameter,  $k$  say, where increasing  $k$  allows computation of a more accurate likelihood approximation at increased cost. The reduced dependence approximation at level  $k$  described in Section 3.2.1 is one example of this. In order for the approximate likelihood inference to be close to the exact likelihood inference,  $k = k_n$  should be allowed to vary with  $n$ . Given an understanding of how the error in the score function varies with  $k$  and  $n$ , the results obtained here could be applied to determine how  $k_n$  should scale with  $n$ . This has the potential to allow the construction of approximate likelihoods which match the inference with the exact likelihood

closely for all  $n$ , but which scale well to large data sizes.

## Acknowledgements

I am grateful to David Firth, Nancy Reid, Cristiano Varin and two referees for helpful comments which have greatly improved this paper. This work was supported by the Engineering and Physical Sciences Research Council through Intractable Likelihood: New Challenges from Modern Applications [grant number EP/K014463/1].

## Appendix: proofs of results

### Proof of Theorem 1

*Proof.* We apply Theorem 5.9 of van der Vaart (2000), with  $\Psi_n(\theta) = r_n^{-1}\tilde{u}_n(\theta)$  and  $\Psi(\theta) = \bar{u}(\theta)$ . Then

$$\begin{aligned} \sup_{\theta \in \Theta} \|\Psi_n(\theta) - \Psi(\theta)\| &= \sup_{\theta \in \Theta} \|\bar{u}_n(\theta) + r_n^{-1}\nabla_{\theta}\epsilon_n(\theta) - \bar{u}(\theta)\| \\ &\leq \sup_{\theta \in \Theta} \|\bar{u}_n(\theta) - \bar{u}(\theta)\| + r_n^{-1}\delta_n \\ &= o_p(1) \end{aligned}$$

The conditions of Theorem 5.9 of van der Vaart (2000) are met, so since  $\Psi_n(\tilde{\theta}_n) = 0$ , we have that  $\tilde{\theta}_n \rightarrow \theta_0$  in probability, as  $n \rightarrow \infty$ .  $\square$

### Proof of Lemma 1

*Proof.* Taking a Taylor expansion of  $\bar{u}_n$  about  $\hat{\theta}_n$ ,

$$\bar{u}_n(\theta) = \bar{u}_n(\hat{\theta}_n) - (\theta - \hat{\theta}_n)^T \bar{J}_n(\theta_n^*) = -(\theta - \hat{\theta}_n)^T \bar{J}_n(\theta_n^*),$$

for some  $\theta_n^*$  between  $\theta$  and  $\hat{\theta}_n$ . Define  $\tilde{\bar{u}}_n(\theta) = r_n^{-1}\tilde{u}_n(\theta)$ . Then

$$\tilde{\bar{u}}_n(\theta) = \bar{u}_n(\theta) + r_n^{-1}\nabla_{\theta}\epsilon_n(\theta) = -(\theta - \hat{\theta}_n)^T \bar{J}_n(\theta_n^*) + r_n^{-1}\nabla_{\theta}\epsilon_n(\theta),$$

so any  $\tilde{\theta}_n$  solving  $\tilde{\bar{u}}_n(\tilde{\theta}_n) = 0$  solves

$$\tilde{\theta}_n - \hat{\theta}_n = \bar{J}_n^{-1}(\theta_n^*)r_n^{-1}\nabla_{\theta}\epsilon_n(\tilde{\theta}_n),$$

for some  $\theta_n^*$  between  $\tilde{\theta}_n$  and  $\hat{\theta}_n$ .

But  $\theta_n^*$  is a consistent estimator of  $\theta$ , because  $\tilde{\theta}_n$  is by Theorem 1, so  $\bar{J}_n(\theta_n^*) \rightarrow I(\theta_0)$  in probability. So

$$\tilde{\theta}_n - \hat{\theta}_n = O_p(r_n^{-1}\delta_n(\tilde{\theta}_n)).$$

Write  $A_n$  for the event  $\{\tilde{\theta}_n \in B_t(\theta_0)\}$ . Then  $\text{pr}(A_n) \rightarrow 1$  as  $n \rightarrow \infty$  since  $\tilde{\theta}_n$  is consistent, and conditional on  $A_n$ ,

$$\tilde{\theta}_n - \hat{\theta}_n = O_p[r_n^{-1}\delta_n^\infty\{B_t(\theta_0)\}] = o_p(r_n^{-1}a_n).$$

For any  $\epsilon > 0$ ,

$$\begin{aligned}
& \text{pr}(\|\tilde{\theta}_n - \hat{\theta}_n\| \geq \epsilon a_n r_n^{-1}) \\
&= \text{pr}(\|\tilde{\theta}_n - \hat{\theta}_n\| \geq \epsilon a_n r_n^{-1} | A_n) \text{pr}(A_n) + \text{pr}(\|\tilde{\theta}_n - \hat{\theta}_n\| \geq \epsilon a_n r_n^{-1} | A_n^C) \text{pr}(A_n^C) \\
&\leq \text{pr}(\|\tilde{\theta}_n - \hat{\theta}_n\| \geq \epsilon a_n r_n^{-1} | A_n) + \text{pr}(A_n^C) \\
&\rightarrow 0
\end{aligned}$$

as  $n \rightarrow \infty$ . □

## Proof of Theorem 2

*Proof.* Applying Lemma 1 with  $a_n = r_n^{1/2}$ ,

$$\tilde{\theta}_n - \hat{\theta}_n = o_p(r_n^{1/2} r_n^{-1}) = o_p(r_n^{-1/2}).$$

So

$$r_n^{1/2}(\tilde{\theta}_n - \theta_0) = r_n^{1/2}(\hat{\theta}_n - \theta_0) + o_p(1) \rightarrow N(0, I(\theta_0)^{-1})$$

in distribution, as required. □

## Proof of Theorem 3

*Proof.* We have

$$\begin{aligned}
(\tilde{\Lambda}_n - \Lambda_n)/2 &= \{\tilde{\ell}_n(\tilde{\theta}_n) - \tilde{\ell}_n(\tilde{\theta}_n^R)\} - \{\ell_n(\hat{\theta}_n) - \ell_n(\hat{\theta}_n^R)\} \\
&= \{\tilde{\ell}_n(\tilde{\theta}_n) - \tilde{\ell}_n(\hat{\theta}_n)\} + \{\tilde{\ell}_n(\hat{\theta}_n) - \ell_n(\hat{\theta}_n)\} + \{\tilde{\ell}_n(\hat{\theta}_n^R) - \tilde{\ell}_n(\tilde{\theta}_n^R)\} + \{\ell_n(\hat{\theta}_n^R) - \tilde{\ell}_n(\hat{\theta}_n^R)\}.
\end{aligned}$$

Taking a Taylor expansion of the first term,

$$\begin{aligned}
\tilde{\ell}_n(\hat{\theta}_n) - \tilde{\ell}_n(\tilde{\theta}_n) &= (\hat{\theta}_n - \tilde{\theta}_n)^T \tilde{u}_n(\tilde{\theta}_n) + (\hat{\theta}_n - \tilde{\theta}_n)^T \tilde{J}_n(\tilde{\theta}_n)(\hat{\theta}_n - \tilde{\theta}_n) \\
&= (\hat{\theta}_n - \tilde{\theta}_n)^T \tilde{J}_n(\tilde{\theta}_n)(\hat{\theta}_n - \tilde{\theta}_n)
\end{aligned}$$

for some  $\bar{\theta}_n$  between  $\hat{\theta}_n$  and  $\tilde{\theta}_n$ , so  $\tilde{\ell}_n(\hat{\theta}_n) - \tilde{\ell}_n(\tilde{\theta}_n) = o_p(1)$ , since  $\|\hat{\theta}_n - \tilde{\theta}_n\| = o_p(r_n^{-1/2})$  and  $\tilde{J}_n(\tilde{\theta}_n) = J_n(\tilde{\theta}_n) + o_p(r_n) = O_p(r_n)$ . Similarly,  $\tilde{\ell}_n(\hat{\theta}_n^R) - \tilde{\ell}_n(\tilde{\theta}_n^R) = o_p(1)$ , so

$$\begin{aligned}
(\tilde{\Lambda}_n - \Lambda_n)/2 &= \epsilon_n(\hat{\theta}_n) - \epsilon_n(\hat{\theta}_n^R) + o_p(1) \\
&= (\hat{\theta}_n - \hat{\theta}_n^R)^T \nabla_{\theta} \epsilon_n(\theta_n^*) + o_p(1)
\end{aligned}$$

for some  $\theta_n^*$  between  $\hat{\theta}_n$  and  $\hat{\theta}_n^R$ . But  $\hat{\theta}_n - \hat{\theta}_n^R = O_p(r_n^{-1/2})$ , and, for sufficiently large  $n$ ,  $\|\nabla_{\theta} \epsilon_n(\theta_n^*)\| \leq \delta_n^{\infty} \{B_t(\theta_0)\}$ , so  $\tilde{\Lambda}_n - \Lambda_n = o_p(1)$ , as required. □

## Results needed to prove Theorem 4

In order to prove Theorem 4, it is helpful to first consider properties of inference with the penalized loglikelihood  $\ell_n^{\pi}(\theta) = \ell(\theta) + \log \pi(\theta)$  with log-prior penalty. We write  $\hat{\theta}_n^{\pi}$  for the corresponding penalized likelihood estimator, which is the posterior mode. Similarly, write  $A^{\pi}$  for the version of the quantity  $A$  computed

with  $\ell_n^\pi(\cdot)$  in place of  $\ell_n(\cdot)$ , and  $\tilde{A}^\pi$  for the approximate version. Since  $\epsilon_n^\pi(\theta) = \epsilon_n(\theta)$ , all the error terms remain unchanged.

Under the regularity conditions assumed on the model, the penalized likelihood estimator  $\hat{\theta}_n^\pi$  will be consistent, and for any  $b_n = o_p(r_n)$ , the posterior probability that  $\theta \in B_{b_n^{-1/2}}(\theta_0)$  will tend to one as  $n \rightarrow \infty$ .

To prove Theorem 4, we will use the following lemma, which says that the error in the penalized log-likelihood ratio may be approximated in terms of the error in the score function.

**Lemma 2.** *Suppose that  $\delta_n^\infty = o_p(r_n)$ , and that there exists  $t > 0$  such that  $\delta_n^\infty \{B_t(\theta_0)\} = o_p(r_n^{1/2})$ . Then  $\hat{\theta}_n^\pi$  and  $\tilde{\theta}_n^\pi$  are consistent estimators of  $\theta_0$ , and  $\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi = o_p(r_n^{-1/2})$ .*

*If  $\gamma_n^\infty \{B_t(\theta_0)\} = o_p(b_n)$ , for some  $b_n = o(r_n)$ , then*

$$\sup_{\theta \in B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)} \left| \left[ \{ \tilde{\ell}_n^\pi(\tilde{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\theta) \} - \{ \ell_n^\pi(\hat{\theta}_n^\pi) - \ell_n^\pi(\theta) \} \right] - [(\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta)] \right| = o_p(1).$$

*Proof.* That  $\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi = o_p(r_n^{-1/2})$  follows by a similar argument to the proof of Theorem 2. Write

$$\begin{aligned} C_n(\theta) &= \{ \tilde{\ell}_n^\pi(\tilde{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\theta) \} - \{ \ell_n^\pi(\hat{\theta}_n^\pi) - \ell_n^\pi(\theta) \} \\ &= \{ \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) - \ell_n^\pi(\hat{\theta}_n^\pi) \} - \{ \tilde{\ell}_n^\pi(\theta) - \ell_n^\pi(\theta) \} - \{ \ell_n^\pi(\hat{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) \} \\ &= \epsilon_n(\hat{\theta}_n^\pi) - \epsilon_n(\theta) - \{ \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\tilde{\theta}_n^\pi) \}. \end{aligned}$$

Then

$$\tilde{\ell}_n(\hat{\theta}_n^\pi) - \tilde{\ell}_n(\tilde{\theta}_n^\pi) = (\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)^T \tilde{J}_n^\pi(\bar{\theta}_n)(\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)$$

for some  $\bar{\theta}_n$  between  $\hat{\theta}_n^\pi$  and  $\tilde{\theta}_n^\pi$ , and

$$\epsilon_n(\hat{\theta}_n^\pi) - \epsilon_n(\theta) = (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta) + (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta^T \nabla_\theta \epsilon_n(\theta_n^*(\theta))(\hat{\theta}_n^\pi - \theta)$$

for some  $\theta_n^*(\theta)$  between  $\hat{\theta}_n^\pi$  and  $\theta$ . Write

$$\begin{aligned} D_n(\theta) &= C_n(\theta) - (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta) \\ &= (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta^T \nabla_\theta \epsilon_n(\theta_n^*(\theta))(\hat{\theta}_n^\pi - \theta) - (\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)^T \tilde{J}_n^\pi(\bar{\theta}_n)(\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi). \end{aligned}$$

Then  $\sup_{\theta \in B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)} |D_n(\theta)|$  may be expressed as

$$\begin{aligned} &\sup_{\theta \in B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)} \left\{ |(\hat{\theta}_n^\pi - \theta)^T \nabla_\theta^T \nabla_\theta \epsilon_n(\theta_n^*(\theta))(\hat{\theta}_n^\pi - \theta) - (\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)^T \tilde{J}_n^\pi(\bar{\theta}_n)(\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)| \right\} \\ &\leq \sup_{\theta \in B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)} \left\{ |(\hat{\theta}_n^\pi - \theta)^T \nabla_\theta^T \nabla_\theta \epsilon_n(\theta_n^*(\theta))(\hat{\theta}_n^\pi - \theta)| \right\} + |(\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)^T \tilde{J}_n^\pi(\bar{\theta}_n)(\hat{\theta}_n^\pi - \tilde{\theta}_n^\pi)| \\ &\leq b_n^{-1} \delta_n^\infty \{B_t(\theta_0)\} + o_p(1) \end{aligned}$$

for  $n$  sufficiently large. But  $\delta_n^\infty \{B_t(\theta_0)\} = o_p(b_n)$ , so  $\sup_{\theta \in B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)} |D_n(\theta)| = o_p(1)$ , as required.  $\square$

## Proof of Theorem 4

*Proof.* The normalized exact and approximate posterior distributions are  $\pi(\theta|y) = L_n(\theta)\pi(\theta)/Z_n$  and  $\tilde{\pi}(\theta|y) = \tilde{L}_n(\theta)\pi(\theta)/\tilde{Z}_n$ , where  $Z_n = \int L_n(\theta)\pi(\theta)d\theta$  and  $\tilde{Z}_n = \int \tilde{L}_n(\theta)\pi(\theta)d\theta$ .

First, we find a Laplace approximation

$$\hat{Z}_n = (2\pi)^{-p/2} |J_n^\pi(\hat{\theta}_n^\pi)|^{-1/2} L_n^\pi(\hat{\theta}_n^\pi)$$

to  $Z_n$ . Then  $\log Z_n - \log \hat{Z}_n = o_p(1)$ , because  $Z_n$  is a  $p$ -dimensional integral, where  $p$  remains fixed as  $n \rightarrow \infty$ . Similarly,  $\tilde{Z}_n$  may be approximated by using a Laplace approximation. Then

$$\begin{aligned} \log Z_n - \log \tilde{Z}_n &= \log \hat{Z}_n - \log \hat{\tilde{Z}}_n + o_p(1) \\ &= (\log |J_n^\pi(\hat{\theta}_n^\pi)| - \log |\tilde{J}_n^\pi(\hat{\theta}_n^\pi)|)/2 + \ell_n^\pi(\hat{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) + o_p(1). \end{aligned}$$

Since  $\gamma_n^\infty\{B_t(\theta_0)\} = o_p(r_n)$ , both  $r_n^{-1}J_n^\pi(\hat{\theta}_n^\pi)$  and  $r_n^{-1}\tilde{J}_n^\pi(\hat{\theta}_n^\pi)$  converge towards  $I(\theta_0)$ , so

$$\begin{aligned} \log |J_n^\pi(\hat{\theta}_n^\pi)| - \log |\tilde{J}_n^\pi(\hat{\theta}_n^\pi)| &= \log |r_n^{-1}J_n^\pi(\hat{\theta}_n^\pi)| - \log |r_n^{-1}\tilde{J}_n^\pi(\hat{\theta}_n^\pi)| \\ &= \log |I(\theta_0) + o_p(1)| - \log |I(\theta_0) + o_p(1)| \\ &= o_p(1). \end{aligned}$$

So

$$\log Z_n - \log \tilde{Z}_n = \ell_n^\pi(\hat{\theta}_n^\pi) - \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) + o_p(1).$$

Since  $\delta_n^\infty\{B_t(\theta_0)\} = o_p(r_n^{-1/2})$  and  $\gamma_n^\infty\{B_t(\theta_0)\} = o_p(r_n)$ , we may choose  $b_n = o(r_n)$  such that  $\delta_n^\infty\{B_t(\theta_0)\} = o_p(b_n^{-1/2})$  and  $\gamma_n^\infty\{B_t(\theta_0)\} = o_p(b_n)$ . Writing  $S_n = B_{b_n^{-1/2}}(\hat{\theta}_n^\pi)$ ,

$$\begin{aligned} &\sup_{\theta \in S_n} \left\{ \left| \log \tilde{\pi}(\theta|y) - \log \pi(\theta|y) - (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta) \right| \right\} \\ &= \sup_{\theta \in S_n} \left\{ \left| \tilde{\ell}_n^\pi(\theta) - \log \tilde{Z}_n - \ell_n^\pi(\theta) + \log Z_n - (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta) \right| \right\} \\ &\leq \sup_{\theta \in S_n} \left\{ \left| \tilde{\ell}_n^\pi(\theta) - \tilde{\ell}_n^\pi(\hat{\theta}_n^\pi) - \ell_n^\pi(\theta) + \hat{\ell}_n^\pi(\hat{\theta}_n^\pi) - (\hat{\theta}_n^\pi - \theta)^T \nabla_\theta \epsilon_n(\theta) \right| \right\} + o_p(1) \\ &= o_p(1), \end{aligned}$$

by Lemma 2. Then

$$\begin{aligned}
2d_{TV}\{\tilde{\pi}(\theta|y), \pi(\theta|y)\} &= \int_{\Theta} |\tilde{\pi}(\theta|y) - \pi(\theta|y)| d\theta \\
&= \int_{S_n} |\tilde{\pi}(\theta|y) - \pi(\theta|y)| d\theta + \int_{S_n^c} |\tilde{\pi}(\theta|y) - \pi(\theta|y)| d\theta \\
&\leq \int_{S_n} |\tilde{\pi}(\theta|y) - \pi(\theta|y)| d\theta + \int_{S_n^c} 2|\pi(\theta|y)| d\theta + \int_{S_n^c} 2|\tilde{\pi}(\theta|y)| d\theta \\
&= \int_{S_n} \left| \frac{\tilde{\pi}(\theta|y)}{\pi(\theta|y)} - 1 \right| \pi(\theta|y) d\theta + o_p(1) \\
&\leq \sup_{\theta \in S_n} \left| \frac{\tilde{\pi}(\theta|y)}{\pi(\theta|y)} - 1 \right| + o_p(1) \\
&= \sup_{\theta \in S_n} \left| \exp\{\log \tilde{\pi}(\theta|y) - \log \pi(\theta|y)\} - 1 \right| + o_p(1) \\
&\leq \left| \exp \left\{ \sup_{\theta \in S_n} |\log \tilde{\pi}(\theta|y) - \log \pi(\theta|y)| \right\} - 1 \right| + o_p(1).
\end{aligned}$$

But

$$\begin{aligned}
\sup_{\theta \in S_n} |\log \tilde{\pi}(\theta|y) - \log \pi(\theta|y)| &\leq \sup_{\theta \in S_n} |(\hat{\theta}_n^\pi - \theta)^T \nabla_{\theta} \epsilon_n(\theta)| + o_p(1) \\
&= b_n^{-1/2} \delta_n^\infty \{B_t(\theta_0)\} + o_p(1) \\
&= o_p(1).
\end{aligned}$$

So  $d_{TV}\{\tilde{\pi}(\theta|y), \pi(\theta|y)\} = o_p(1)$ , as claimed.  $\square$

## References

- Douglas Bates, Martin Mächler, Ben Bolker, and Steve Walker. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1):1–48, 2015. doi: 10.18637/jss.v067.i01.
- Randal Douc, Éric Moulines, and Tobias Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. *The Annals of Statistics*, 32(5):2254–2304, 2004.
- Nial Friel, AN Pettitt, Robert Reeves, and Ernst Wit. Bayesian inference in hidden markov random fields for binary data defined on large lattices. *Journal of Computational and Graphical Statistics*, 18(2):243–261, 2009.
- V. P. Godambe. An optimum property of regular maximum likelihood estimation. *Ann. Math. Statist.*, 31(4):1208–1211, 12 1960. doi: 10.1214/aoms/1177705693. URL <http://dx.doi.org/10.1214/aoms/1177705693>.
- Michael I. Jordan. Graphical models. *Statist. Sci.*, 19(1):140–155, 02 2004. doi: 10.1214/088342304000000026. URL <http://dx.doi.org/10.1214/088342304000000026>.
- Bruria Kaufman. Crystal statistics. II. Partition function evaluated by spinor analysis. *Physical Review*, 76(8):1232, 1949.

- Bruce G. Lindsay. Composite likelihood methods. *Contemporary Mathematics*, 80:221–239, 1988.
- Francesco Pauli, Walter Racugno, and Laura Ventura. Bayesian composite marginal likelihoods. *Statistica Sinica*, pages 149–164, 2011.
- José C. Pinheiro and Douglas M. Bates. Approximations to the log-likelihood function in the nonlinear mixed-effects model. *Journal of Computational and Graphical Statistics*, 4(1):12–35, 1995. ISSN 10618600. URL <http://www.jstor.org/stable/1390625>.
- Mathieu Ribatet, Daniel Cooley, and Anthony C Davison. Bayesian inference from composite likelihoods, with an application to spatial extremes. *Statistica Sinica*, pages 813–845, 2012.
- Håvard Rue, Sara Martino, and Nicolas Chopin. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(2):319–392, 2009.
- Håkon Tjelmeland and Haakon Michael Austad. Exact and approximate recursive calculations for binary markov random fields defined on graphs. *Journal of Computational and Graphical Statistics*, 21(3):758–780, 2012. doi: 10.1080/10618600.2012.632236. URL <http://dx.doi.org/10.1080/10618600.2012.632236>.
- Aad W van der Vaart. *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2000.
- Cristiano Varin, Nancy Reid, and David Firth. An overview of composite likelihood methods. *Statistica Sinica*, 21(1):5–42, 2011.



# Supplementary material for On asymptotic validity of naive inference with an approximate likelihood

Helen Ogden

University of Southampton, UK

## 1 Finding $\delta_{(i)}(\theta)$ in Example 3.1

Recall  $Y_i$  is the number of successes out of  $m = m_n$  trials on item  $i$ . We study how  $\delta_{(i)}(\theta) = \|(d/d\theta)\epsilon_{(i)}(\theta)\|$  varies with  $m$ .

Write

$$f(b; y_i) = -y_i \log \{\text{logit}^{-1}(b)\} + (m - y_i) \log \{1 - \text{logit}^{-1}(b)\}$$

and

$$g(b; \theta, y_i) = f(b; y_i) - \log \phi(b; 0, \theta),$$

so that

$$L_{(i)}(\theta) = \int_{-\infty}^{\infty} \exp\{-g(b; \theta, y_i)\} db.$$

In the following, we drop the data  $y_i$  from the notation for convenience. Write  $\hat{b}(\theta)$  for the maximizer of  $g(\cdot, \theta)$ , and

$$\hat{g}_r(\theta) = \frac{\partial^r g}{\partial b^k}(\hat{b}(\theta); \theta).$$

By equation (4) of Shun and McCullagh (1995), the error in the Laplace approximation to the log-likelihood  $\ell_{(i)}(\theta)$  is

$$\epsilon_{(i)}(\theta) = \sum_{l=1}^{\infty} \frac{1}{2l!} \sum_{P \in \mathcal{P}_{2l}} n_2(P) (-1)^v \hat{g}_{|p_1|}(\theta) \dots \hat{g}_{|p_v|}(\theta) \{\hat{g}_2(\theta)\}^{-l}, \quad (\text{S1})$$

where  $P = p_1 | \dots | p_v$  is a partition of  $2l$  indices into  $v$  blocks of size 3 or more, and  $n_2(P)$  is the number of partitions  $Q$  of  $2l$  indices into  $l$  blocks of size 2, such that  $Q$  is complementary to  $P$ .

Write  $h_P(\theta) = \hat{g}_{|p_1|}(\theta) \dots \hat{g}_{|p_v|}(\theta) \{\hat{g}_2(\theta)\}^{-l}$ . Then  $h_P(\theta) = O_p(m^{v-l})$ , since  $\hat{g}_r(\theta) = O_p(m)$  for each  $r$ .

Differentiating (S1) gives

$$\frac{d}{d\theta} \epsilon_{(i)}(\theta) = \sum_{l=1}^{\infty} \frac{1}{2l!} \sum_{P \in \mathcal{P}_{2l}} n_2(P) (-1)^v \frac{d}{d\theta} h_P(\theta), \quad (\text{S2})$$

and

$$\frac{d}{d\theta} h_P(\theta) = \sum_{i=1}^v \left[ \hat{g}'_{|p_i|}(\theta) \prod_{j \neq i} \hat{g}_{|p_j|}(\theta) \{\hat{g}_2(\theta)\}^{-l} - \prod_{j=1}^v l \hat{g}_{|p_j|}(\theta) \hat{g}'_2(\theta) \{\hat{g}_2(\theta)\}^{-(l+1)} \right], \quad (\text{S3})$$

where  $\hat{g}'_r(\theta) = (d/d\theta)\hat{g}_r(\theta)$ .

For each  $r$ , we have

$$\begin{aligned} \hat{g}'_r(\theta) &= \frac{d}{d\theta} \left\{ g^{(r)}(\hat{b}(\theta); \theta) \right\} \\ &= \frac{\partial g^{(r)}}{\partial \theta}(\hat{b}(\theta); \theta) + \frac{d\hat{b}(\theta)}{d\theta} \hat{g}_{r+1}(\theta). \end{aligned} \quad (\text{S4})$$

We now study the size of each of the terms in (S4). We have

$$\frac{\partial g^{(r)}}{\partial \theta}(b; \theta) = \frac{\partial}{\partial \theta} \left\{ -\frac{\partial^r}{\partial b^r} \log \phi(b; 0, \theta) \right\} = O_p(1), \quad (\text{S5})$$

and

$$\hat{g}_{r+1}(\theta) = O_p(m). \quad (\text{S6})$$

For each  $\theta$ ,  $\hat{b}(\theta)$  satisfies  $g_1(\hat{b}(\theta); \theta) = 0$ . Differentiating this with respect to  $\theta$ ,

$$\frac{d\hat{b}(\theta)}{d\theta} g_2(\hat{b}(\theta); \theta) + \frac{\partial g_1}{\partial \theta}(\hat{b}(\theta); \theta) = 0.$$

But

$$\frac{\partial g_1}{\partial \theta}(b; \theta) = -2b\theta^{-3},$$

so

$$\frac{d\hat{b}(\theta)}{d\theta} = 2\hat{b}(\theta)\theta^{-3} \{\hat{g}_2(\theta)\}^{-1} = O_p(m^{-1}). \quad (\text{S7})$$

Substituting (S5), (S6) and (S7) into (S4) gives that  $\hat{g}'_r(\theta) = O_p(1)$  for each  $r$ . From (S3) we then have

$$\frac{d}{d\theta} h_P(\theta) = O_p(m^{v-l-1}).$$

The highest order terms in (S2) come from partitions with  $(l, v) = (2, 1)$  or  $(3, 2)$ , and so  $\delta_{(i)}(\theta) = O_p(m^{-2})$ .

## 2 Finding $\delta_m^{(k)}(\beta)$ in Example 3.2

Kaufman (1949) provides an exact expression for the normalizing constant for an Ising model on an  $n \times m$  lattice, with  $\alpha = 0$  and periodic boundary, as

$$Z_{n \times m}(0, \beta) = \{2 \sinh(2\beta)\}^{nm/2} \bar{A}_{n,m}(\beta)/2,$$

where

$$\bar{A}_{n,m}(\beta) = A_{n,m}^{(1)}(\beta) + A_{n,m}^{(2)}(\beta) + A_{n,m}^{(3)}(\beta) + A_{n,m}^{(4)}(\beta),$$

and

$$A_{n,m}^{(1)}(\beta) = \prod_{q=0}^n 2 \cosh(m a_{2q+1,n}(\beta)/2), \quad A_{n,m}^{(2)}(\beta) = \prod_{q=0}^n 2 \sinh(m a_{2q+1,n}(\beta)/2),$$

$$A_{n,m}^{(3)}(\beta) = \prod_{q=0}^n 2 \cosh(m a_{2q,n}(\beta)/2), \quad A_{n,m}^{(4)}(\beta) = \prod_{q=0}^n 2 \sinh(m a_{2q,n}(\beta)/2)$$

where

$$a_{l,n}(\beta) = \cosh^{-1} \{ \cosh(2\beta)^2 / \sinh(2\beta) - \cos(\pi l/n) \}$$

for  $l \geq 1$ , and  $a_{0,n}(\beta) = a_0(\beta) = 2\beta + \log \{ \tanh(\beta) \}$ .

Using the approximation  $Z_{m \times m}^{(k)}(\beta)$  to  $Z_{m \times m}(\beta)$ , the error in the log-likelihood is

$$\epsilon_m^{(k)}(\beta) = (m-k+1) \log \bar{A}_{k,m}(\beta) - (m-k) \log \bar{A}_{k-1,m}(\beta) - \log \bar{A}_{m,m}(\beta).$$

Differentiating this with respect to  $\beta$ ,

$$\frac{d}{d\beta} \epsilon_m^{(k)}(\beta) = (m-k+1) \frac{d}{d\beta} \{ \log \bar{A}_{k,m}(\beta) \} - (m-k) \frac{d}{d\beta} \{ \log \bar{A}_{k-1,m}(\beta) \} - \frac{d}{d\beta} \{ \log \bar{A}_{m,m}(\beta) \}, \quad (\text{S8})$$

and

$$\frac{d}{d\beta} \{ \log \bar{A}_{n,m}(\beta) \} = \sum_{i=1}^4 \frac{d}{d\beta} \{ \log A_{n,m}^{(i)}(\beta) \} r_{n,m}^{(i)}(\beta),$$

where  $r_{n,m}^{(i)}(\beta) = A_{n,m}^{(i)}(\beta) / \bar{A}_{n,m}(\beta)$ .

We have

$$\begin{aligned} \frac{d}{d\beta} \{ \log A_{n,m}^{(1)}(\beta) \} &= m/2 \sum_{q=0}^n a'_{2q+1,n}(\beta) \tanh(m a_{2q+1,n}(\beta)/2) \\ &= m/2 \sum_{q=0}^n a'_{2q+1,n}(\beta) + O(m \exp\{-a_0(\beta)m\}) \end{aligned}$$

as  $m \rightarrow \infty$ , since  $\tanh(x) = 1 + O(\exp\{-2x\})$  as  $x \rightarrow \infty$ , and  $a_{2q+1,n}(\beta) \geq a_0(\beta) > 0$ .

Similar expressions may be obtained for the other  $\frac{d}{d\beta} \{ \log A_{n,m}^{(i)}(\beta) \}$ , and combining these gives

$$\frac{d}{d\beta} \{ \log \bar{A}_{n,m}(\beta) \} = m S_n^{(o)}(\beta) r_{n,m}^{(o)}(\beta) + m S_n^{(e)}(\beta) r_{n,m}^{(e)}(\beta) + O(m \exp\{-a_0(\beta)m\})$$

where  $S_n^{(o)} = \sum_{q=0}^n a'_{2q+1,n}(\beta)$ ,  $S_n^{(e)} = \sum_{q=0}^n a'_{2q,n}(\beta)$ ,  $r_{n,m}^{(o)}(\beta) = r_{n,m}^{(1)}(\beta) + r_{n,m}^{(2)}(\beta)$  and  $r_{n,m}^{(e)}(\beta) = r_{n,m}^{(3)}(\beta) + r_{n,m}^{(4)}(\beta)$ . Define

$$f(x; \beta) = d_\beta \{-1 + c_\beta - \cos(x)\}^{-1/2} \{1 + c_\beta - \cos(x)\}^{-1/2}$$

where  $d_\beta = 4 \cosh(2\beta) - 2 \cosh(2\beta) \coth(2\beta)^2$  and  $c_\beta = \cosh(2\beta)^2 / \sinh(2\beta)$ . Then  $a'_{j,n}(\beta) = f(j\pi/n; \beta)$ , and  $n^{-1} S_n^{(o)}(\beta)$  and  $n^{-1} S_n^{(e)}(\beta)$  are both trapezium rule approximations to  $I(\beta) = \frac{1}{2\pi} \int_0^{2\pi} f(x; \beta) dx$ . Write  $R_n^{(o)}(\beta) = n^{-1} S_n^{(o)}(\beta) - I(\beta)$  and  $R_n^{(e)}(\beta) = n^{-1} S_n^{(e)}(\beta) - I(\beta)$  for the error in each of these approximations to the integral.

**Lemma 3.** For each  $\beta < \beta_c$ ,  $R_n(\beta) = \max\{|R_n^{(e)}(\beta)|, |R_n^{(o)}(\beta)|\} = O(\exp\{-b_\beta n\})$ , where  $b_\beta = 2 \cosh^{-1}\{-1 + \cosh(2\beta)^2 / \sinh(2\beta)\}$ .

*Proof.* We apply the results of Trefethen and Weideman (2014) to show exponentially fast convergence of these trapezium rule approximations to  $I(\beta)$ . These results depend on properties of the integrand  $f(z, \beta)$ , considered as a function of complex-valued  $z$ . There are a branch points of  $f(z, \beta)$  at a distance  $a_\beta = \cosh^{-1}\{-1 + \cosh(2\beta)^2 / \sinh(2\beta)\}$  from the real axis, and the function is analytic for  $-a_\beta < \text{Im } z < a_\beta$ , so by Theorem 3.2 of Trefethen and Weideman (2014),  $|R_n^{(o)}(\beta)| = O(\exp\{-2a_\beta n\}) = O(\exp\{-b_\beta n\})$ .

The same argument holds with  $R_n^{(e)}(\beta)$  in place of  $R_n^{(o)}(\beta)$ , so  $R_n(\beta) = O(\exp\{-b_\beta n\})$ , as required.  $\square$

We now prove the main result.

**Lemma 4.** If  $k \rightarrow \infty$  as  $m \rightarrow \infty$ ,  $\delta_m^{(k)}(\beta) = O(m^2 k \exp\{-b_\beta k\}) + o(1)$ .

*Proof.* We have

$$\frac{d}{d\beta} \{\log \bar{A}_{n,m}(\beta)\} = mnI(\beta) + mnt_{n,m}(\beta) + O(m \exp\{-a_0(\beta)m\})$$

where  $t_{n,m}(\beta) = R_n^{(o)}(\beta)r_{n,m}^{(o)}(\beta) + R_n^{(e)}(\beta)r_{n,m}^{(e)}(\beta)$ .

Substituting this into (S8), the contributions from the  $mnI(\beta)$  terms cancel, and the combined remainder terms are always  $o(1)$ , since  $m^2 \exp\{-a_0(\beta)m\} = o(1)$ . We are left with

$$\frac{d}{d\beta} \epsilon_m^{(k)}(\beta) = (m - k + 1)mt_{k,m}(\beta) - (m - k)mt_{k-1,m}(\beta) - mt_{m,m}(\beta) + o(1).$$

Then

$$\begin{aligned} \delta_m^{(k)}(\beta) &= \left| \frac{d}{d\beta} \epsilon_m^{(k)}(\beta) \right| \\ &\leq (m - k + 1)mk|t_{k,m}(\beta)| + (m - k)m(k - 1)|t_{k-1,m}(\beta)| + m^2|t_{m,m}(\beta)| + o(1) \\ &\leq (m - k + 1)mkR_k(\beta) + (m - k)m(k - 1)R_{k-1}(\beta) + m^2R_m(\beta) + o(1) \end{aligned}$$

since  $|t_{n,m}(\beta)| \leq |R_n^{(o)}(\beta)|r_{n,m}^{(o)}(\beta) + |R_n^{(e)}(\beta)|r_{n,m}^{(e)}(\beta) \leq R_n(\beta)$

$$= O(m^2 k \exp\{-b_\beta k\}) + o(1)$$

by Lemma 3, as required.  $\square$

## References

- Bruria Kaufman. Crystal statistics. II. Partition function evaluated by spinor analysis. *Physical Review*, 76(8):1232, 1949.
- Zhenming Shun and Peter McCullagh. Laplace approximation of high dimensional integrals. *Journal of the Royal Statistical Society. Series B (Methodological)*, 57(4):749–760, 1995.
- Lloyd N Trefethen and JAC Weideman. The exponentially convergent trapezoidal rule. *SIAM Review*, 56(3):385–458, 2014.