

An apparent paradox concerning the field of an ideal dipole

Edward Parker*

University of California, Santa Barbara

(Dated: November 4, 2018)

The electric or magnetic field of an ideal dipole is known to have a Dirac delta function at the origin. The usual textbook derivation of this delta function is rather ad hoc and cannot be used to calculate the delta-function structure for higher multipole moments. Moreover, a naive application of Gauss's law to the ideal dipole field appears to give an incorrect expression for the dipole's effective charge density. We derive a general procedure for calculating the delta-function structure at the origin of an arbitrary ideal multipole field; this procedure leads to the nontrivial result that the divergence of a singular vector field can contain a *derivative* of a Dirac delta function even if the field itself does not contain a delta function. We also argue that a physical interpretation of the delta function in the dipole field previously given in the literature is incorrect.

I. INTRODUCTION

If we expand the electric potential due to a localized source of electric charge in terms of multipole moments, we find that (in CGS units) the dipole term is [1]

$$\phi(\mathbf{x}) = \frac{\mathbf{p} \cdot \mathbf{x}}{r^3} = \frac{p_i x_i}{r^3}, \quad (1)$$

where \mathbf{p} is the charge distribution's dipole moment (chosen to lie at the coordinate origin parallel to \hat{z}), $r := |\mathbf{x}|$, and repeated indices are summed. Far away from the charge distribution, the far field due to the dipole moment is

$$E_{\text{ff},i}(\mathbf{x}) = \frac{3n_i p_j n_j - p_i}{r^3}, \quad (2)$$

where the radial unit vector $\mathbf{n} := \mathbf{x}/r$. For example, if the charge distribution consists of two particles with charge q and $-q$ separated by a displacement vector \mathbf{d} with $\mathbf{p} = q\mathbf{d}$, then \mathbf{E}_{ff} describes the field at distances $r \gg d$. In the idealized limit where $q \rightarrow \infty$ and $\mathbf{d} \rightarrow \mathbf{0}$ with their product \mathbf{p} held constant, the potential is given exactly by (1) for all $\mathbf{x} \neq \mathbf{0}$.

Our experience with the identity $\nabla^2(1/r) = -4\pi\delta^3(\mathbf{x})$ tells us that potentials with singularities can have derivatives with Dirac delta functions that cannot be captured by a naive differentiation, suggesting that the far-field expression (2) may need to be modified at the origin for an ideal point dipole. Indeed, the correct expression is

$$E_i(\mathbf{x}) = \frac{3n_i p_j n_j - p_i}{r^3} - \frac{4\pi}{3} p_i \delta^3(\mathbf{x}). \quad (3)$$

One standard argument [1] justifying the delta-function term is that if V is the interior of a sphere containing all of the electric charge in a system with net electric dipole moment \mathbf{p} , then

$$\int_V d^3x \mathbf{E}(\mathbf{x}) = -\frac{4\pi}{3} \mathbf{p}. \quad (4)$$

Evaluating the volume integral over (2) is tricky, because the integrand diverges at the origin. The natural regularization procedure is to adopt the convention that all integrals are evaluated in spherical coordinates, with the angular integrals performed first. (This is a higher-dimensional analog of the convention that all integrals over singular functions are given by their Cauchy principal values. In the more familiar case of a monopole, the electric field only diverges as $1/r^2$ so the integral converges in spherical coordinates, and we do not need to specify a regularization scheme.) By symmetry, only the z -component could be nonzero, and

$$\begin{aligned} & \int_V d^3x \left(\frac{3n_z p_j n_j - p_z}{r^3} \right) \\ &= p \int_V d^3x \left(\frac{3 \cos^2 \theta - 1}{r^3} \right) \\ &= 4\pi p \int_0^R \frac{dr}{r} \int_0^\pi d\theta \sin \theta (3 \cos^2 \theta - 1) \\ &= 0, \end{aligned}$$

so the far-field term does contribute to the integral in identity (4), and with this choice of regularization the expression (3) satisfies the identity.

A second argument [2] uses the identity $\partial_j \left(\frac{1}{r} \right) = -\frac{x_j}{r^3}$ to get

$$E_i(\mathbf{x}) = -\partial_i \phi(\mathbf{x}) = -p_j \partial_i \left(\frac{x_j}{r^3} \right) = p_j \partial_i \partial_j \left(\frac{1}{r} \right). \quad (5)$$

The mixed partial derivative is given by the identity [2, 3]

$$\partial_i \partial_j \left(\frac{1}{r} \right) = \frac{3n_i n_j - \delta_{ij}}{r^3} - \frac{4\pi}{3} \delta_{ij} \delta^3(\mathbf{x}), \quad (6)$$

and (5) and (6) together give (3). This identity was also explicitly derived using the convention that angular integrals are always evaluated first. (This convention is known as spherical regularization; different regularizations of the improper integral give expressions different from (6) [4]. Roughly speaking, identity (4) requires that the two terms in (3) must have a "total" delta function of

* tparker@physics.ucsb.edu

$-\frac{4\pi}{3}\mathbf{p}\delta^3(\mathbf{x})$ between them, and the spherical regularization scheme puts it entirely into the second term. Under a regularization scheme in which $\int_V d^3x \mathbf{E}_{\text{ff}}(\mathbf{x}) = c\mathbf{p}$, the coefficient of the delta function in (6) becomes $-4\pi/3-c$.)

A third argument [5, 6] modifies the potential (1) by multiplying it by the unit step function $\theta(r)$ and using the fact that $\frac{d\theta}{dr} = \delta(r)$.

The first argument is quite ad hoc - we simply noticed that the naive expression (2) fails to satisfy one particular identity, and manually added a term to satisfy it. The second argument is much more satisfying, as [2] derives the delta-function term together with the far-field term. The third argument is quite tricky to formulate consistently, as it requires the use of expressions like $\frac{\mathbf{p}}{r}\delta^3(\mathbf{x})$ and integrals in which a delta function lies exactly at one limit of integration, which raise extremely subtle mathematical questions. We would like to find a simple, intuitive algorithm for finding the delta-function structure at the origin of an arbitrary multipole field.

The delta-function term in (3) has physically measurable effects. The simplest one actually occurs in the corresponding expression for an ideal magnetic dipole. The magnetic field $\mathbf{B}(\mathbf{x})$ is the curl of a vector potential, so if a sphere V contains all the current in a system then $\int_V d^3x \mathbf{B}(\mathbf{x}) = \frac{8\pi}{3}\mathbf{m}$, where \mathbf{m} is the current distribution's magnetic dipole moment. We must therefore add to the magnetic dipole far-field term $\mathbf{B}_{\text{ff}}(\mathbf{x}) \equiv \mathbf{E}_{\text{ff}}(\mathbf{x})$ (with \mathbf{p} replaced by \mathbf{m}) a term $\frac{8\pi}{3}\mathbf{m}\delta^3(\mathbf{x})$ under a spherical regularization scheme [1]. In the nonrelativistic limit, particles with quantum-mechanical spin correspond to (so far as we know) ideal magnetic dipoles. Moreover, a particle's wavefunction can probe the magnetic field precisely at another particle's location, so the delta-function term can affect the particles' interaction. This can be seen most simply in the hyperfine splitting of the ground-state energy levels of the hydrogen atom due to the coupling between the proton's and electron's spins. If we treat the spins' dipole-dipole interaction as a perturbation to the usual classical Coulomb potential, then it is straightforward to calculate that the first-order contribution to the hyperfine splitting is

$$\Delta E_{\text{hf}} = \frac{8\pi}{3} \frac{\gamma_e \gamma_p \hbar^2}{\pi a^3} = 5.884 \times 10^{-6} \text{ eV},$$

where γ_e and γ_p represent the electron's and proton's gyromagnetic ratios, respectively, and a is the Bohr radius [7]. The prefactor $8\pi/3$ comes from the prefactor of the delta-function term in the ideal magnetic dipole field. This energy level splitting is responsible for the famous 21-cm hydrogen line measured by radio astronomy, which is one of the most common forms of radiation in the universe and has been measured extremely accurately. The prediction above agrees with experiment to 99.8% accuracy, and quantum electrodynamics corrections further improve the accuracy [8].

For simplicity, we will now only consider electric multipole fields. Similar considerations apply to magnetic multipoles, but the vector nature of the potential intro-

duces mathematical complications which do not significantly affect our conclusions.

II. AN APPARENT PARADOX

Another motivation for considering the delta-function structure at the origin more carefully and generally is given by an apparent paradox that arises in computing the effective charge density of an ideal dipole, which is

$$\rho(\mathbf{x}) = -\mathbf{p} \cdot \nabla \delta^3(\mathbf{x}) = -p\delta(x)\delta(y)\delta'(z). \quad (7)$$

The form of the expression is intuitively clear when we consider the ideal dipole as the limit of a physical dipole as $\mathbf{d} \rightarrow 0$ and $q \rightarrow \infty$, and that the distribution $\delta'(x)$ corresponds to a function that is strongly peaked at $(-\epsilon, p/\epsilon)$ and $(\epsilon, -p/\epsilon)$ for infinitesimal ϵ . We can derive it more rigorously in two different ways [1]. One way is to note that

$$\begin{aligned} \phi(\mathbf{x}) &= \int d^3x' \frac{\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} = -\mathbf{p} \cdot \int d^3x' \frac{\nabla' \delta^3(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} \\ &= \mathbf{p} \cdot \int d^3x' \delta^3(\mathbf{x}') \nabla' \left(\frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) = \frac{\mathbf{p} \cdot \mathbf{x}}{r^3} \end{aligned}$$

in accordance with (1) (where ∇' denotes the gradient with respect to \mathbf{x}'). A second, similar method is to verify that the potential energy of the dipole in an external potential $\phi_{\text{ext}}(\mathbf{x})$ (which does not include the potential from the dipole itself) gives the correct expression

$$\begin{aligned} U &= \int d^3x \phi_{\text{ext}}(\mathbf{x}) \rho(\mathbf{x}) = -\mathbf{p} \cdot \int d^3x \phi_{\text{ext}}(\mathbf{x}) \nabla \delta^3(\mathbf{x}) \\ &= \mathbf{p} \cdot \int d^3x \delta^3(\mathbf{x}) \nabla \phi_{\text{ext}}(\mathbf{x}) = -\mathbf{p} \cdot \mathbf{E}_{\text{ext}}(\mathbf{0}). \quad (8) \end{aligned}$$

But applying Gauss's law to (3) gives

$$\nabla \cdot \mathbf{E}(\mathbf{x}) = \nabla \cdot \mathbf{E}_{\text{ff}}(\mathbf{x}) - \frac{4\pi}{3}\mathbf{p} \cdot \nabla \delta^3(\mathbf{x}). \quad (9)$$

The divergence $\nabla \cdot \mathbf{E}_{\text{ff}}$ is clearly zero away from the origin because there is no charge away from the dipole. $\mathbf{E}_{\text{ff}}(\mathbf{x})$ diverges at the origin, so we must specify a regularization scheme in order to evaluate its divergence there. In order to be consistent with the derivations in Section I, we must again adopt the spherical regularization scheme in which all integrals are performed in spherical coordinates and the angular integrals are performed first. Under this scheme, \mathbf{E}_{ff} does not "contain" a delta function at the origin (as discussed above), so taking the divergence of \mathbf{E}_{ff} should not produce a derivative of a delta function. We therefore seem to have that $\nabla \cdot \mathbf{E}(\mathbf{x}) \stackrel{?}{=} -\frac{4\pi}{3}\mathbf{p} \cdot \nabla \delta^3(\mathbf{x})$, from which Gauss's law $\nabla \cdot \mathbf{E}(\mathbf{x}) = 4\pi\rho(\mathbf{x})$ implies that $\rho(\mathbf{x}) \stackrel{?}{=} -\frac{1}{3}\mathbf{p} \cdot \nabla \delta^3(\mathbf{x})$, which disagrees with (7).

This apparent paradox occurs under any choice of regularization, because as discussed in Section I, identity (4) requires that the "total" delta function across both terms

in (3) must be $-\frac{4\pi}{3}\mathbf{p}\delta^3(\mathbf{x})$, so the ‘‘total’’ derivative of a delta function in $\nabla \cdot \mathbf{E}(\mathbf{x})$ should be $-\frac{4\pi}{3}\mathbf{p} \cdot \nabla\delta^3(\mathbf{x})$, implying that $\rho(\mathbf{x}) \stackrel{?}{=} -\frac{1}{3}\mathbf{p} \cdot \nabla\delta^3(\mathbf{x})$ under any choice of regularization. We also cannot resolve the paradox by changing the prefactor of the delta-function term in (3) to -4π , because doing so would contradict both (4) and experimental results, as discussed above.

III. DISCUSSION

Let us generalize to an arbitrary ideal multipole potential $\phi^{(lm)}(\mathbf{x})$ corresponding to a multipole moment q_{lm} , which we will define by

$$\phi^{(lm)}(\mathbf{x}) = q_{lm} \frac{Y_{lm}(\Omega)}{r^{l+1}}$$

where Ω denotes the angular coordinates, $Y_{lm}(\Omega)$ the usual spherical harmonics, and there is no sum on repeated indices. (Note that we use a different normalization convention from [1] for q_{lm} .) Away from the origin, the electric field is given by the far-field expression $\mathbf{E}_{\text{ff}}^{(lm)}(\mathbf{x}) = -\nabla\phi^{(lm)}(\mathbf{x})$.

Delta functions and their derivatives are defined by their integrals against arbitrary smooth test functions $f(\mathbf{x})$, so in order to calculate $\nabla \cdot \mathbf{E}_{\text{ff}}^{(lm)}(\mathbf{x})$ at the origin we must evaluate

$$\int_V d^d x [f(\mathbf{x}) \nabla \cdot \mathbf{E}_{\text{ff}}^{(lm)}(\mathbf{x})]$$

over a volume V that includes the origin. We integrate by parts and suppress the superscripts, subscripts, and arguments \mathbf{x} for clarity:

$$\int_V d^3 x [f \nabla \cdot \mathbf{E}] = \oint_{\partial V} d\mathbf{S} \cdot (f \mathbf{E}) - \int_V d^3 x [\mathbf{E} \cdot \nabla f]. \quad (10)$$

Only the neighborhood of the origin contributes to the integral on the LHS, so its value does not depend on the volume V (as long as it contains the origin). WLOG, we take it to be a ball of radius R centered at the origin. We are interested in fields \mathbf{E} that diverge at the origin, so we also need to specify a regularization scheme for the volume integral on the RHS of (10). We use spherical regularization again:

$$\begin{aligned} \int_V d^3 x [f \nabla \cdot \mathbf{E}] &= R^2 \oint d\Omega \cdot (f \mathbf{E}) \Big|_{r=R} \\ &\quad - \lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^R dr r^2 \oint d\Omega (\mathbf{E} \cdot \nabla f). \end{aligned} \quad (11)$$

We now expand $f(\mathbf{x})$ in complex conjugate spherical harmonics:

$$f(\mathbf{x}) = \sum_{l'=0}^{\infty} \sum_{m'=-l'}^{l'} c_{l'm'}(r) Y_{l'm'}^*(\Omega). \quad (12)$$

The polar angle dependence of $Y_{l'm'}^*(\Omega)$ is given by an order- l' associated Legendre polynomial of $\sin\theta$ and $\cos\theta$, so in order for $f(\mathbf{x})$ to be smooth at the origin, $c_{l'm'}(r)$ must have a zero of order at least l' at $r=0$.

The radial component of the electric field

$$E_r = (l+1) \frac{q_{lm}}{r^{l+2}} Y_{lm}(\Omega), \quad (13)$$

and because of the orthonormality of the spherical harmonics, the angular integration over the surface term in (11) gives

$$R^2 \oint d\Omega \cdot (f \mathbf{E}) \Big|_{r=R} = (l+1) q_{lm} \frac{c_{lm}(R)}{R^l}. \quad (14)$$

The volume term can be expressed as

$$\int_{\epsilon}^R dr r^2 \oint d\Omega [(E_r \partial_r + \mathbf{E}_{\perp} \cdot \nabla_{\perp}) f]$$

where the subscript \perp denotes the angular coordinates. (12) and (13) together give

$$\oint d\Omega E_r \partial_r f = (l+1) q_{lm} \frac{c'_{lm}(r)}{r^{l+2}}. \quad (15)$$

We can also integrate

$$\begin{aligned} \oint d\Omega [(\mathbf{E}_{\perp} \cdot \nabla_{\perp}) f] &= - \oint d\Omega [\nabla_{\perp} \phi^{(lm)} \cdot \nabla_{\perp} f] \\ &= \oint d\Omega [\phi^{(lm)} \nabla_{\perp}^2 f] \end{aligned}$$

by parts with no surface term because the surface of integration is closed. Using the eigenvalue identity $\nabla_{\perp}^2 Y_{lm}(\Omega) = -\frac{l(l+1)}{r^2} Y_{lm}(\Omega)$, the angular integral gives

$$\oint d\Omega [(\mathbf{E}_{\perp} \cdot \nabla_{\perp}) f] = -l(l+1) q_{lm} \frac{c_{lm}(r)}{r^{l+3}}. \quad (16)$$

Combining (15) and (16), the volume term in (11) becomes

$$\begin{aligned} &\int_{\epsilon}^R dr r^2 \oint d\Omega (\mathbf{E} \cdot \nabla f) \\ &= (l+1) q_{lm} \int_{\epsilon}^R dr \left[\frac{c'_{lm}(r)}{r^l} - l \frac{c_{lm}(r)}{r^{l+1}} \right] \\ &= (l+1) q_{lm} \int_{\epsilon}^R dr \frac{d}{dr} \left(\frac{c_{lm}(r)}{r^l} \right) \\ &= (l+1) q_{lm} \left(\frac{c_{lm}(R)}{R^l} - \frac{c_{lm}(\epsilon)}{\epsilon^l} \right). \end{aligned} \quad (17)$$

The first term is cancelled by the surface term (14), so

$$\int_V d^3 x [f \nabla \cdot \mathbf{E}_{\text{ff}}^{(lm)}] = (l+1) q_{lm} \lim_{\epsilon \rightarrow 0^+} \frac{c_{lm}(\epsilon)}{\epsilon^l}.$$

Since $c_{lm}(r)$ has a zero of order at least l at $r=0$, this limit converges and we finally arrive at

$$\int_V d^3 x [f \nabla \cdot \mathbf{E}_{\text{ff}}^{(lm)}] = \frac{l+1}{l!} q_{lm} \frac{d^l c_{lm}}{dr^l} \Big|_{r=0}. \quad (18)$$

This result allows us to easily extract the delta-function structure of the far field of an ideal multipole. We see that an order- l multipole has an order- l *derivative* of a delta function at the origin.

For example, for an order $l = 0$ multipole (a monopole) with charge q , we only need to know the value of $f(\mathbf{x})$ at the origin: $f(\mathbf{x}) = c_{00}(r)Y_{00}^*(\Omega) + \dots = (4\pi)^{-1/2}c_{00}(0) + o(r)$. With our choice of normalization conventions, $q_{00} = \sqrt{4\pi}q$ [1] so

$$\begin{aligned} \int_V d^3x \left[f \nabla \cdot \mathbf{E}_{\text{ff}}^{(00)} \right] &= q_{00} c_{00}(0) = 4\pi q f(\mathbf{0}) \\ \nabla \cdot \mathbf{E}_{\text{ff}}^{(00)}(\mathbf{x}) &= 4\pi q \delta^3(\mathbf{x}) \\ \rho(\mathbf{x}) &= q \delta^3(\mathbf{x}). \end{aligned}$$

In the case of a dipole $\mathbf{p} \parallel \hat{z}$, we have $q_{10} = \sqrt{4\pi/3}p$ [1] and we need to keep the term

$$\begin{aligned} f(\mathbf{x}) &= c_{10}(r) Y_{10}^*(\Omega) + \dots \\ &= \sqrt{\frac{3}{4\pi}} \cos\theta c_{10}(r) + \dots \end{aligned}$$

in expansion (12). The easiest way to proceed is to Taylor expand $c_{10}(r)$ (recalling that $c_{10}(0) = 0$) and then convert to Cartesian coordinates:

$$\begin{aligned} f(\mathbf{x}) &= \sqrt{\frac{3}{4\pi}} \cos\theta c'_{10}(0)r + \dots \\ &= \sqrt{\frac{3}{4\pi}} c'_{10}(0)z + \dots \\ \left. \frac{\partial f}{\partial z} \right|_{\mathbf{x}=\mathbf{0}} &= \sqrt{\frac{3}{4\pi}} c'_{10}(0). \end{aligned}$$

Combining this with (18),

$$\begin{aligned} \int_V d^3x \left[f \nabla \cdot \mathbf{E}_{\text{ff}}^{(10)} \right] &= \frac{8\pi}{3} p \left. \frac{\partial f}{\partial z} \right|_{\mathbf{x}=\mathbf{0}} \\ \nabla \cdot \mathbf{E}_{\text{ff}}^{(10)}(\mathbf{x}) &= -\frac{8\pi}{3} \mathbf{p} \cdot \nabla \delta^3(\mathbf{x}). \end{aligned} \quad (19)$$

Even though $\mathbf{E}_{\text{ff}}^{(10)}(\mathbf{x})$ does not have delta function at the origin (under spherical regularization), its divergence nevertheless has the derivative of a delta function at the origin! We come to the surprising conclusion that by taking the divergence of a singular vector field, it is possible to directly produce the derivative of a delta function without ever “passing through” a delta function (either implicit or explicit) in the vector field.

This resolves the apparent paradox discussed in Section II: we see that the mistaken step in our reasoning was in assuming that because \mathbf{E}_{ff} does not contain a delta function at the origin, $\nabla \cdot \mathbf{E}_{\text{ff}}$ does not contain the derivative of a delta function. But in fact the “far field” and “near field” terms on the RHS of (9) *both* have nonzero divergence at the origin and contribute to the charge density. The two terms on the RHS of (9) add up to $\nabla \cdot \mathbf{E}(\mathbf{x}) = -4\pi \mathbf{p} \cdot \nabla \delta^3(\mathbf{x})$. Gauss’s law then

gives the correct answer $\rho(\mathbf{x}) = -\mathbf{p} \cdot \nabla \delta^3(\mathbf{x})$, in agreement with (7). ([5] derives a result equivalent to (19) using a very different method that, as mentioned above, raises subtle mathematical issues. Using the method of [5], one can separately calculate the far-field contribution to the charge density from any multipole moment, but the calculation becomes increasingly complicated for higher moments, whereas the general result (18) captures the contributions from all multipole moments at once.)

The fact that the far-field term \mathbf{E}_{ff} gives a contribution to $\nabla \cdot \mathbf{E}$ that is necessary for the consistence of the theory means that [6] and [7] are incorrect in interpreting the field of an ideal dipole “on the understanding that the [far-field] term applies only to the region *outside* an infinitesimal sphere about the point $r = 0$.” The far-field term actually corresponds to two-thirds of the charge density at the origin, and is therefore important for capturing the physics arbitrarily close to the dipole.

If the arbitrary function $f(\mathbf{x})$ itself happens to obey the Laplace equation, then $c_{l'm'}(r) \propto r^{l'}$ for all $l' \geq 0$. In this case, the surface term (14) is actually independent of the radius R of the region of integration, and the volume term (17) becomes zero so only the surface term contributes to (11). Since the surface term does not go to zero at long distances, we must be careful to always retain it when integrating by parts. This can pose some subtleties; for example, when treating $f(\mathbf{x})$ as an external potential and trying to calculate the potential energy of a dipole in a manner similar to (8), one must be careful to retain the surface term in (10). Of course, a potential of this form which grows unboundedly with distance is not very physical, as it would need to be generated by an infinitely extended charge distribution.

Finally, there is also a less-rigorous shortcut for calculating (19). Applying Gauss’s law to (5) gives

$$\rho(\mathbf{x}) = \frac{1}{4\pi} p_j \partial_i \partial_i \partial_j \left(\frac{1}{r} \right) = p_j C_j(\mathbf{x}),$$

where $C_j(\mathbf{x}) := \frac{1}{4\pi} \partial_i \partial_i \partial_j \left(\frac{1}{r} \right)$. (6) then gives

$$C_j(\mathbf{x}) = \frac{1}{4\pi} \partial_i \left(\frac{3n_i n_j - \delta_{ij}}{r^3} \right) - \frac{1}{3} \partial_j \delta^3(\mathbf{x}). \quad (20)$$

$C_j(\mathbf{x})$ is clearly an extremely pathological distribution, and it is not at all clear whether its mixed partial derivatives commute. If we nevertheless assume that they do, then we get the alternative expression

$$C_j(\mathbf{x}) = \frac{1}{4\pi} \partial_j \left(\partial_i \partial_i \left(\frac{1}{r} \right) \right) = -\partial_j \delta^3(\mathbf{x}). \quad (21)$$

(Evaluating the partial derivatives in this order roughly physically corresponds to first finding the charge distribution corresponding to an ideal monopole, then spatially differentiating that distribution to “split” the monopole into an ideal dipole.) Equating (20) and (21) gives

$$\partial_i \left(\frac{3n_i n_j - \delta_{ij}}{r^3} \right) = -\frac{8\pi}{3} \partial_j \delta^3(\mathbf{x}),$$

and contracting both sides with p_j gives (19).

ACKNOWLEDGMENTS

The author would like to thank Mark Srednicki, Brayden Ware, Alex Rasmussen, and Dominic Else for helpful

discussions, and Bruno Klajn for bringing [5] to the author's attention.

-
- [1] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, 1999) pp. 145 - 150, 170, 187-188.
 - [2] C. P. Frahm, *American Journal of Physics* **51**, 826 (1983).
 - [3] J. M. Bowen, *American Journal of Physics* **62**, 511 (1994).
 - [4] V. Hnizdo, *European Journal of Physics* **32**, 287 (2011).
 - [5] V. Namias, *International Journal of Mathematical Education in Science and Technology* **18**, 767 (1987).
 - [6] S. M. Blinder, *American Journal of Physics* **71**, 816 (2003).
 - [7] D. J. Griffiths, *American Journal of Physics* **50**, 698 (1982).
 - [8] S. J. Brodsky and S. D. Drell, *Annual Reviews of Nuclear Science* **20**, 183 (1970).