

An ecological resilience perspective on cancer: insights from a toy model

Artur C. Fassoni¹ and Hyun M. Yang²

¹IMC, UNIFEI, Itajubá, MG, Brazil

²IMECC, UNICAMP, Campinas, SP, Brazil

Abstract

In this paper we propose an ecological resilience point of view on cancer. This view is based on the analysis of a simple ODE model for the interactions between cancer and normal cells. The model presents two regimes for tumor growth. In the first, cancer arises due to three reasons: a partial corruption of the functions that avoid the growth of mutated cells, an aggressive phenotype of tumor cells and exposure to external carcinogenic factors. In this case, treatments may be effective if they drive the system to the basin of attraction of the cancer cure state. In the second regime, cancer arises because the repair system is intrinsically corrupted. In this case, the complete cure is not possible since the cancer cure state is no more stable, but tumor recurrence may be delayed if treatment is prolonged. We review three indicators of the resilience of a stable equilibrium, related with size and shape of its basin of attraction: latitude, precariousness and resistance. A novel method to calculate these indicators is proposed. This method is simpler and more efficient than those currently used, and may be easily applied to other population dynamics models. We apply this method to the model and investigate how these indicators behave with parameters changes.

Keywords: Tumor growth; Chemotherapy; Basins of Attraction; Regime shifts; Critical transitions.

1 Introduction

The ecological resilience perspective is an emerging approach for understanding the dynamics of social-ecological systems [1, 2, 3, 4, 5, 6]. While the stability point of view emphasizes the equilibrium and the maintenance of present state, the resilience point of view focus on shifts between alternative basins of attraction, thresholds, uncertainty and unexpected disturbances. External forces or random events may cause state variable perturbations that drive a nonlinear system, which is initially near a stable state, to enter a undesirable basin of attraction. In this case, the resilience of the original steady state is related with the size and shape of its basin of attraction, and the capacity of the system to persist in this basin of attraction when subject to state variable perturbations. Three different indicators are established in the literature as measures of the resilience of a stable state with respect to state variable perturbations [7, 8]: *latitude*, which is a measure of the volume of its basin of attraction; *precariousness*, which is related with the minimal state space disturbance needed to drive the system outside its basin of attraction; and *resistance*, which is a measure of the deepness of its basin of attraction.

On the other hand, changes in system parameters occur in a slow time scale, due to evolutionary forces or by modifying the intensities of interactions and forces governing such system.

In this case, parameters modify the resilience of the system with respect to state variable perturbations. Further, when parameters do change enough, the system may undergo several bifurcations and the phase portrait may change substantially. In this case, one can measure the resilience of the system with respect to parameters changes as the distance to the threshold values for which bifurcations occur. As a consequence of such bifurcations, an undesirable alternative stable state may be created, and its basin of attraction can be achieved by state variable perturbations, as commented above. A more dramatic outcome happens when parameters changes lead to loss of stability of the original steady state or even its disappearance. In this case, a regime shift occurs and the system moves to another state. Now, the question of reversibility takes place. Of first importance is the question whether it is possible or not to return the parameters to their original values. When parameters change due to evolutionary factors, it is more likely that this change can not be undone. Changes due to external forces can be undone more easily through the correct manipulation of those forces (if possible). However, even if the original values can be restored, the reversal to the original stable state may not be completely achieved if the system exhibits hysteresis.

In this paper we illustrate how these concepts of ecological resilience can be applied to cancer, a complex disease whose causes are far from being well understood and whose cure is far from being achieved. Indeed, despite the intense efforts that led the elucidation of many biochemical mechanisms developed by cancer cells to survive [9], there is a current debate on which are the major factors that allow the onset of cancer cells. While some argue that alterations in intrinsic cellular processes are the main reasons that some tissues become cancerous [10], others defend the view that most cases of cancer result from extrinsic factors such environmental exposure to toxic chemicals and radiation [11]. With respect to cancer treatment, although the development of new drugs and strategies to treat cancer in the last fifty years achieved good results in many cases, another large portion of cancer patients did not respond well to treatments, or presented tumor recurrence, indicating that there is still a long road in the fight against cancer [12, 13].

We propose a toy model for tumor growth and apply the above concepts to suggest a framework for viewing the arising of cancer and its effective treatment as critical transitions between two alternative stable states. In this framework, tumor growth and tumor treatment depend ultimately on ecological resilience questions. Further, we briefly review the three resilience indicators commented above, propose a method to calculate these indicators and apply this method to the model. As far as we know, this novel method we propose is simpler and more efficient than those currently used, and can be applied to other population dynamics models to improve their analysis through this resilience perspective.

The paper is organized as follows. In Section 2 the model is presented. In Section 3 the analysis of the model is performed. In Section 4, the results are discussed in the ecological resilience perspective. In Section 5, the method to calculate resilience indicators is presented and applied to the model. Finally, conclusions are presented in Section 6.

2 A toy model for tumor growth

We present a toy model consisting of a system of ODEs describing tumor growth and its effect on normal tissue, together with the tissue response to tumor. Our goal is not to consider the several aspects of tumor growth and to reproduce quantitative behavior with very accuracy, but to use the model to give some insights through a resilience point of view. The model equations

are given by

$$\frac{dN}{dt} = r_N - \mu_N N - \beta_1 N A, \quad (1a)$$

$$\frac{dA}{dt} = r_A A \left(1 - \frac{A}{K_A}\right) - \beta_3 N A - (\mu_A + \epsilon_A) A, \quad (1b)$$

where N and A stand for normal and tumor cells, respectively. This system is a limit case of a three-dimensional model for oncogenesis encompassing mutations and genetic instability [14].

Parameter r_N represents the total constant reproduction of normal cells, and μ_N is their natural mortality. A constant flux for normal cells is considered in the vital dynamics, and not a density-dependent one, like the logistic growth generally assumed [15, 16, 17, 18]. The reason for this choice is that at a normal and already formed tissue the imperative dynamics is not the cells intraspecific competition by nutrients, but the maintenance of a homeostatic state, through the natural replenishment of old and dead cells [19].

On the contrary, cancer cells have a certain independence on growth signals released by the tissue and keep their own growth program, like an embryonic tissue in growth phase [20]. Thus a density dependent growth is considered. Several growth laws could be used, such as the Gompertz, generalized logistic, Von Bertalanffy and others [21]. We choose the logistic growth due its simplicity, and a natural mortality μ_A . An extra mortality rate ϵ_A due to apoptosis [22] is also included.

Several models for tumor growth consider the phenomena of tumor angiogenesis, i.e., the formation of new blood vessels to feed the tumor, in response to signals released by tumor cells [23, 24]. In order to keep the model as simple as possible, we do not consider angiogenesis here.

Parameter β_3 encompasses, in the simplest way, all negative responses of the many cell types in normal tissue due to the presence of tumor cells. These interactions include the release of anti-growth and death signals by host cells [9], the immune-system response [25], competition by nutrients with tumor cells and so on. Similarly, parameter β_1 covers all mechanisms developed by tumor cells which damage the normal tissue, like increasing local acidity [26], suppression of immune cells [27], release of death signals [9], and competition with normal cells.

System (1) is similar to the classical Lotka-Volterra model of competition [28], commonly used in models for tumor growth [15, 16, 17, 18] and biological invasions [29], but has a fundamental difference. The use of a constant flux instead a logistic growth to normal cells breaks the symmetry present in the classical Lotka-Volterra model, so that no equilibrium with $N = 0$ will exist. Thus, normal cells will never be extinct, on the contrary to those models. We believe that this is not a problem, but, on the contrary, is a realistic outcome. Indeed, roughly speaking, cancer ‘wins’ not by the fact that it kills all cells in the tissue, but by the fact that it reaches a dangerous size that disrupt the well functioning of the tissue and threaten the health of the individual. A constant flux term was already taken in other well-know models for cancer, specifically, to describe the growth of immune cells [30, 31, 32].

3 Analysis of the model

We now present the analysis of system (1). Biological implications are discussed in Section 4.

3.1 Equilibrium points

System (1) has a trivial equilibrium

$$P_0 = \left(\frac{r_N}{\mu_N}, 0 \right),$$

and up to two nontrivial equilibria

$$P_i = (N_i, A_i) = \left(\frac{r_N}{\mu_N + \beta_1 A_i}, A_i \right), \quad i = 1, 2.$$

Here, A_1 and A_2 are the roots of the second degree equation

$$aA^2 + bA + c = 0, \quad (2)$$

with coefficients

$$a = \frac{\beta_1 r_A}{K_A} > 0, \quad b = l_A (\beta_1^{th} - \beta_1), \quad c = r_N (\beta_3 - \beta_3^{th}).$$

where

$$\beta_1^{th} = \frac{\mu_N r_A}{l_A K_A}, \quad \text{and} \quad \beta_3^{th} = \frac{\mu_N}{r_N} l_A. \quad (3)$$

When A_1 and A_2 are real, we label them in the order $A_1 < A_2$. Conditions for having positive equilibria P_1 and P_2 are obtained by Descartes' Rule of Signs. Together with the trivial equilibrium P_0 , the results are summarized as follows:

- I) If $\beta_3 > \beta_3^{th}$ and $\beta_1 < \beta_{1,\Delta}^{th}$, the unique nonnegative equilibrium is the trivial equilibrium P_0 .
- II) If $\beta_3 > \beta_3^{th}$ and $\beta_1 > \beta_{1,\Delta}^{th}$, three nonnegative equilibria are P_0 , P_1 and P_2 .
- III) If $\beta_3 < \beta_3^{th}$, the nonnegative equilibria are P_0 and P_2 .

The threshold $\beta_{1,\Delta}^{th}$, defined for $\beta_3 > \beta_3^{th}$, is the value of $\beta_1 > \beta_1^{th}$ for which the discriminant $\Delta = b^2 - 4ac$ is zero, and is given by

$$\beta_{1,\Delta}^{th} = \beta_1^{th} + 2\eta + 2\sqrt{\eta(\beta_1^{th} + \eta)}, \quad (4)$$

where $\eta = r_A r_N (\beta_3 - \beta_3^{th}) / (K_A l_A^2)$.

The threshold β_3^{th} can be written as $\beta_3^{th} = l_A / N_0$, where $N_0 = r_N / \mu_N$ is the number of normal cells in the absence of tumor cells. As we will see, condition $\beta_3 > \beta_3^{th}$ implicates on more difficult regimes for cancer to arise. Thus, favorable regimes for tumor growth occur for larger values of net reproduction rate l_A of tumor cells and for smaller values of carrying capacity of normal cells N_0 . The threshold β_1^{th} for tumor aggressiveness increases as the mortality μ_N of normal cells increases, and decreases as the effective carrying capacity of tumor cells $l_A K_A / r_A$ increases.

3.2 Local Stability

Stability of P_0 is easily determined. The eigenvalues of the Jacobian matrix of system (1) evaluated at P_0 are given by

$$\lambda_1 = -\mu_N, \quad \text{and} \quad \lambda_2 = \frac{r_N}{\mu_N} (\beta_3^{th} - \beta_3).$$

Thus, P_0 is locally asymptotically stable if $\beta_3 > \beta_3^{th}$, and is a saddle otherwise.

We now study the local stability of P_i , $i = 1, 2$. Using the fact that N_i and A_i satisfy

$$l_A - \frac{r_A}{K_A} A_i - \beta_3 N_i = 0, \quad (5)$$

we find that the Jacobian matrix of system (1) evaluated at P_i is given by

$$j(P_i) = \begin{bmatrix} -\beta_1 A_i - \mu_N & -\frac{\beta_1}{\beta_3} (l_A - \frac{r_A}{K_A} A_i) \\ -\beta_3 A_i & -\frac{r_A}{K_A} A_i \end{bmatrix}.$$

Whenever P_i is a positive equilibrium, the trace of $j(P_i)$ is negative. Thus, when P_i is positive, both eigenvalues of $j(P_i)$ will have negative real part if $\det(j(P_i)) > 0$, and we have opposite signs in the other case. Solving equation (2) for A_i , and calculating the determinant we obtain

$$\det(j(P_i)) = A_i(2aA_i + b).$$

As $A_1 < A_2$, we have, whenever P_i is a positive equilibrium, $i = 1, 2$,

$$\det(j(P_1)) = -A_1\sqrt{\Delta} < 0, \quad \text{and} \quad \det(j(P_2)) = A_2\sqrt{\Delta} > 0,$$

where Δ is the discriminant of (2). Thus, P_1 will be a saddle point whenever it is positive (case II above), and P_2 will be stable whenever it is positive (cases II and III above).

3.3 Asymptotic behavior and global stability

Let us show the boundedness of trajectories of (1). By noting that

$$\frac{dN}{dt} \leq r_N - \mu_N N,$$

and

$$\frac{dA}{dt} \leq l_A A \left(1 - \frac{r_A}{l_A K_A} A \right),$$

we may apply classical comparison principles [33] and conclude that all solutions $(N(t), A(t))$ with non-negative initial values remain restricted in the box

$$B = \left[0, \frac{r_N}{\mu_N} \right] \times \left[0, \frac{l_A}{r_A} K_A \right] \quad (6)$$

when $t \rightarrow \infty$.

In order to rule out periodic orbits for system (1) we apply the Dulac Criterion [34] with $u(N, A) = 1/NA$, obtaining

$$\nabla \cdot \left(\frac{1}{NA} \left(\frac{dN}{dt}, \frac{dA}{dt} \right) \right) = -\frac{r_A NA + K_A r_N}{K_A AN^2} < 0$$

for $(N, A) \in B$. Thus, system (1) has no periodic orbits.

By the Poincaré-Bendixson Theorem we conclude that all trajectories converge to an equilibrium point [34]. It implies that equilibria P_0 and P_2 are globally stable in cases I and III, respectively (in the latter, P_2 is globally stable for initial conditions $A(0) > 0$, since the N axis is the stable manifold of P_0). In case II, the plane $N \times A$ is divided in the basins of attraction of P_0 and P_2 . The stable manifold of P_1 is the separatrix between these basins. All these results are summarized in Theorem 1.

Theorem 1. *System (1) has the following behavior:*

- I) *If $\beta_3 > \beta_3^{th}$ and $\beta_1 < \beta_{1,\Delta}^{th}$, then P_0 is globally stable.*
- II) *If $\beta_3 > \beta_3^{th}$ and $\beta_1 > \beta_{1,\Delta}^{th}$, then P_0 and P_2 are locally stable. Equilibrium P_1 is a saddle point whose stable manifold is the separatrix between the basins of attraction of P_0 and P_2 .*
- III) *If $\beta_3 < \beta_3^{th}$, then P_2 is globally stable for initial conditions $A(0) > 0$.*

The division of the $\beta_1 \times \beta_3$ plane into regions I, II and III is showed in Figure 1.




Figure 1: Left: Regions I, II and III in the $\beta_3 \times \beta_1$ plane of parameters space, together with equilibria behavior. Right: bidimensional diagram of the A coordinates of equilibria P_2 (red, stable), P_1 (orange, unstable) and P_0 (blue, stable for $\beta_3 > \beta_3^{th}$), when β_3 and β_1 vary. When $\beta_3 = \beta_3^{th}$ a transcritical bifurcation occurs between P_0 and P_2 if $\beta_1 < \beta_1^{th}$, and between P_0 and P_1 if $\beta_1 > \beta_1^{th}$. When $\beta_1 = \beta_{1,\Delta}^{th}$ we have a saddle-node bifurcation between P_1 and P_2 .

3.4 Numerical simulations

We now present numerical simulations in order to stimulate discussions in next sections. Figure 2 shows simulations of system (1) in cases II and III. Parameters values were based on data from the literature, specially for breast cancer, according the procedure below. A summary of the parameter values is presented in Table 1.




Figure 2: Solutions of system (1) when parameters correspond to regimes II (left) and III (right), using values in Table 1. Initial conditions are $N(0) = r_N/\mu_N - A_0$ and $A(0) = A_0$, representing that the tissue was initially at its homeostatic state when A_0 cells have become cancerous. The values of A_0 are: $A_0 = 0.06 \times 10^8$ and $A_0 = 0.07 \times 10^8$ (left); $A_0 = 1$ and $A_0 = 0.07 \times 10^8$ (right). In the left panel, the gray dotted curve represents the separatrix between the basins of attraction of P_0 and P_2 . The green and blue numbers indicate the time (in years) corresponding to the trajectory.

We assume that the lifetime of a normal cell is 100 days, thus $\mu_N = 1/100 \text{ days}^{-1}$. The number of normal cells in the breast cannot pass $N_0 = 10^8$ cells [35]. Thus, in order to adjust the equilibrium r_N/μ_N of normal cells in the absence of cancer to be 10^8 cells, we consider $r_N = 10^6$

Table 1: Parameters description and values adopted in simulations.

Parameter	Description	Value
μ_N	$1/\mu_N$ is the lifetime of a normal cell	0.01 day^{-1}
r_N	total constant reproduction of normal cells	$10^6 \text{ cell day}^{-1}$
r_A	tumor cells growth rate	0.05 day^{-1}
K_A	tumor carrying capacity	$0.75 \times 10^8 \text{ cells}$
μ_A	natural mortality rate of cancer cells	0.01 day^{-1}
ϵ_A	extra mortality rate of cancer cells	0.01 day^{-1}
β_1	cancer cells aggressiveness	$0.40 \times 10^{-9} \text{ cell}^{-1}\text{day}^{-1}$
β_3^{II}	tissue response to cancer cells - case II	$0.28 \times 10^{-9} \text{ cell}^{-1}\text{day}^{-1}$
β_3^{III}	tissue response to cancer cells - case III	$0.32 \times 10^{-9} \text{ cell}^{-1}\text{day}^{-1}$

cells/day. For cancer cells, we assume the same natural mortality, $\mu_A = 1/100 \text{ days}^{-1}$. For the apoptotic rate of A cells, we use $\epsilon_A = 1/100 \text{ days}^{-1}$. The ratio *birth rate/death rate* is 1 for a normal tissue, in order to maintain a homeostatic state. Following [35], we assume that cancer cells have increased this ratio by a factor of five. Thus, we have $r_A = 5/100 \text{ days}^{-1}$. In order to have the maximum number of cancer cells being 75% of the normal cells, we consider $K_A = 7.5 \times 10^7$. The values of interacting parameters β_1 and β_3 , in units of $\text{cell}^{-1}\text{day}^{-1}$, are unknown a priori but, by substituting the values of other parameters, we obtain the thresholds for β_1 and β_3 : the threshold β_3^{th} which separates cases II and III has the value $\beta_3^{th} = 0.30 \times 10^{-9}$. So, we assume two possible values for β_3 : $\beta_3^{III} = 0.28 \times 10^{-9}$, and $\beta_3^{II} = 0.32 \times 10^{-9}$. Each of these values will originate a different behavior of subsystem NA (see Figure 1). If $\beta_3 = \beta_3^{III}$ we are in case III, for every value of β_1 . If $\beta_3 = \beta_3^{II}$, we have $\beta_{1,\Delta}^{th} = 0.37 \times 10^{-9}$, so we assume $\beta_1 = 0.40 \times 10^{-9} > \beta_3^{II}$, which is reasonable since cancer cells are supposed to cause more damage to normal cells than the contrary. With these values we are in case II. In all numerical simulations in this paper, we use these parameter values, and $\beta_3 = \beta_3^{III}$ or $\beta_3 = \beta_3^{II}$, depending on the interest to simulate cases III or II.

4 An ecological resilience perspective on cancer

We now discuss the biological implications of the previous analysis. Our look to system (1) as being a simple cartoon, a toy model, of the underlying system governing tumor growth in a cancer patient and thus we apply the perspective of ecological resilience to discuss the results above. Although this is a rough approximation, it may be instructive illustration on our understanding of cancer onset and cancer treatment. By cartoon or an approximation, we mean that the underlying system of cancer in real life, despite being very complex, may presents three qualitative distinct regimes, corresponding to regimes I, II and III of system (1). In this analogy, an equilibrium state corresponding to the presence of a tumor is not necessarily a static equilibrium, but a state of the system where a tumor is growing and developing. Let us discuss the differences between these regimes.

4.1 Cancer onset as a critical transition

Initially, we look to cancer onset as a critical transition. Let us first comment at the ‘natural repair system of the patient’, a mechanism which is operated at a variety of levels and by many agents. In the tissue level, it is operated by the immune system, trough lymphocytes and natural killers cells for example [25]. The presence of cancer cells at a given site stimulate the locomotion of immune system cells to that site in order to eliminate the cancer cells. In the

cellular level, many cell components watch some parameters of the own cell and its neighbors, as DNA integrity, the products of cellular metabolism, the concentration of growth factors, etc. When abnormal conditions are detected, the cell may kill itself through apoptosis [22], or the normal cells may release death or inhibitor factors to control the undesired growth [9]. In system (1), these mechanisms are roughly described by the parameters β_3 and ϵ_A . These are the parameters most subject to changes in a slow-time scale, through the multistep process of genetic alterations which transform the descendants of a normal cell in a malign tumor [9]. Parameter β_1 also is thought to be a varying parameter in this slow-time scale, since it encompasses the many types of negative interactions which cancer cells impose to the host tissue, specially due to changes in their metabolism which increase local acidity or lead to starvation of oxygen and nutrients for normal cells.

In the first regime (region I), we have a healthy person, since P_0 is globally stable. In this case, we have an efficient tissue repair system, since condition $\beta_3 > \beta_3^{th}$ can be written as

$$\beta_3 + \epsilon_A \frac{\mu_N}{r_N} > \frac{\mu_N}{r_N} (r_A - \mu_A).$$

Further, we have a limited aggressiveness of cancer cells, $\beta_1 < \beta_{1,\Delta}^{th}$. This condition depends also on β_3 , because $\beta_{1,\Delta}^{th}$ depends on β_3 (see Figure 1, left). Therefore, for a fixed β_1 , condition $\beta_1 < \beta_{1,\Delta}^{th}$ is equivalent to $\beta_3 > \beta_{3,\Delta}^{th}$, where $\beta_{3,\Delta}^{th}$ is the inverse function of $\beta_{1,\Delta}^{th}$. Thus, for each level of aggressiveness of cancer cells, we have a second threshold that the tissue response must be above in order to completely eliminate the chance of cancer. More aggressive are the cancer cells, higher is this threshold. Thus, region I corresponds to parameters such that, although new mutant cells may arise all the time, they are not so much aggressive and the intrinsic repair system is capable to eliminate them.

In the second regime (region II), we have the possibility of having cancer, since P_0 and P_2 are both stable. Condition $\beta_3 > \beta_3^{th}$ implies that the tissue response is efficient, but condition $\beta_1 > \beta_{1,\Delta}^{th}$, which is equivalent to $\beta_3 < \beta_{3,\Delta}^{th}$, implies that the tissue response is not completely capable to face the aggressiveness of cancer cells. Thus, region II corresponds to a partially corrupted repair system due to the aggressiveness of cancer cells. In this region, the resilience of the cancer cure equilibrium P_0 plays an important role. The survival and installation of a tumor mass depends on an increased genetic instability and/or exposure to external carcinogenic factors, in such way the mutations from normal to cancer cells increase the initial conditions of cancer cells allow them to surpass the threshold separating the basins of attraction of P_0 and P_2 ; see Figure 2 left. Therefore, it is important to analyze how the resilience of P_0 behave as key parameters are changed. In the next Section we develop this ‘resilience analysis’. We refer to [14] for a model that comprises an intermediary pre-cancer population that continuously ‘feed’ the population A with new individuals.

Finally, case III represents a dramatic corruption of the repair system, $\beta_3 < \beta_3^{th}$. Now, the cancer equilibrium P_2 is globally stable, and the onset and development of cancer is possible for any initial number of cancer cells. However, distinct quantitative behaviors may be observed, depending on the initial number of cancer cells. In Figure 2 we see that the time that the tumor reaches a detectable size (approximately 10^6 cells [36]) is more than ten years if a single mutant cell arises. As the number of initial cancer cells increases, this time decreases.

We present in Figure 3 bifurcation diagrams when β_3 and β_1 vary. When β_3 varies with β_1 fixed, two different cases occur (Figure 3, top panels). In the case i), the tumor is not much aggressive to normal cells, $\beta_1 < \beta_1^{th}$, and we have a transition between regimes I and III through a transcritical bifurcation. In case ii), the tumor is very aggressive, $\beta_1 > \beta_1^{th}$, and we have transitions from regimes I to II (saddle-node bifurcation) and from II to III (transcritical bifurcation). Before entering regime III, there is a previous and additional interval, $\beta_3^{th} < \beta_3 < \beta_{3,\Delta}^{th}$ that allows cancer onset depending on initial conditions (regime II). The

comparison of cases (i) and (ii) contributes to the debate of whether intrinsic or extrinsic factors are the major responsible for cancer onset [11, 10]. Due to the possibility of a direct transition from regime I to II when $\beta_1 > \beta_1^{th}$, we conclude that extrinsic factors appear to be the major cause of aggressive tumors, but combined with small contribution of intrinsic factors ($\beta_3 < \beta_{3,\Delta}^{th}$). On the other hand, the possibility of direct transition from I to III in case (i) implicates that non-aggressive tumors may arise due to intrinsic factors only.




Figure 3: Top: bifurcation diagrams of coordinate A of equilibrium points when β_3 varies, with $\beta_1 < \beta_1^{th}$ (left), and $\beta_1 > \beta_1^{th}$ (right). Down: bifurcation diagrams when β_1 varies, with $\beta_3 < \beta_3^{th}$ (left), and $\beta_3 > \beta_3^{th}$ (right). Continuous plot corresponds to stable equilibrium, while dashed plot corresponds to unstable equilibrium. These diagrams are particular cases (horizontal sections i and ii and vertical sections a and b) of the bidimensional one, shown in Figure 1.

When β_1 varies while β_3 is kept constant, we have a different effect (Figure 3, bottom panels). In the case when the tissue response to tumor is low, $\beta_3 < \beta_3^{th}$, no bifurcation occurs and we remain in regime III for all β_1 . In the second case, the tissue response is high, $\beta_3 > \beta_3^{th}$, and a transition between regimes I and II occurs through a saddle-node bifurcation. We remain in regime II for all values of $\beta_1 > \beta_{1,\Delta}^{th}$, and no transition to regime III occurs, on the contrary to case (b) when β_3 varies with a high aggressiveness of tumor. Comparison of diagrams (ii) and (b) lead to an interesting conclusion. While normal cells are able with their own characteristics (strong repair system) to guarantee tissue integrity against aggressive tumors, aggressive cancer cells, on the other hand, depend on genetic instability (initial conditions) when fighting aggressive normal cells.

4.2 Cancer treatment as an attempt to allow a critical transition

Now, let us consider cancer treatment. We focus on application of chemotherapy due to its widely use, but some of the general results may be extend to other types of treatment, like

radiotherapy or surgery. A simple way to include chemotherapy in system (1) is to consider the following equations:

$$\frac{dN}{dt} = r_N - \mu_N N - \beta_1 N A - \alpha_N \gamma_N N D, \quad (7a)$$

$$\frac{dA}{dt} = r_A A \left(1 - \frac{A}{K_A}\right) - (\mu_A + \epsilon_A) A - \beta_3 N A - \alpha_A \gamma_A A D, \quad (7b)$$

$$\frac{dD}{dt} = v(t) - \gamma_N N D - \gamma_A A D - \tau D. \quad (7c)$$

Here, D is the chemotherapeutic drug. It is administered according to a treatment function $v(t)$, and has a clearance rate τ . The terms $\gamma_N N D$ and $\gamma_A A D$ describe drug absorption by normal and cancer cells, while the killing terms $\alpha_A \gamma_A A D$ and $\alpha_N \gamma_N N D$ follow the log-kill hypotheses [13]. The treatment function can be described as a finite sum of Dirac Deltas,

$$v(t) = \sum_{i=1}^n \rho \delta(t - iT),$$

representing n doses of ρ mg of drug each, every T days. Figures 4 and 5 show simulations of system (7) in cases II and III respectively. In these simulations, we assumed the values $\tau = 2.5 \text{ day}^{-1}$, $\gamma_A = 0.3 \times 10^{-8} \text{ cell}^{-1} \text{ day}^{-1}$, and $\alpha_A = 0.5 \times 10^8 \text{ cell mg}^{-1}$ and $\gamma_N = 0.6\gamma_A$ and $\alpha_N = 0.6\alpha_A$ since the chemotherapeutic agent is supposed to be more specific to cancer cells than normal cells. The treatment parameters were $\rho = 10$ mg, $T = 7$ days, and $n = 4, 5, 6, 7$ doses.

More important than the form used to modeling the treatment $v(t)$, the general property is that all chemotherapeutic treatments cease after some time $t_f > 0$, i.e., $v(t) = 0$ for all $t > t_f$. Thus, $D(t) \leq -\tau D$ and so $D \rightarrow 0$ when $t \rightarrow \infty$. Therefore *all solutions of system (7) approach solutions of system (1) when $t \rightarrow \infty$* . This fact has an important consequence in our ecological perspective. In the real system underlying tumor growth, the treatment would have only the effect of state space disturbance, without altering the intrinsic dynamics. This important feature reveals that the possibility of cure, above all, concerns questions of stability and resilience. Figures 4 and 5 provide an illustration of this fact.

If tumor growth in a patient is described by some underlying dynamical system which does not have a cancer cure stable equilibrium, then a complete cure is not possible. This is what happens in case III, due to the weakness of the repair system, and it is illustrated in Figure 5. After getting near this equilibrium through the treatment, the system moves back to the cancer equilibrium, even if it takes a long time. Thus, it is necessary a systemic change that alters the dynamics permanently. However, as also shown in Figure 5, even in this regime of instability of the cure equilibrium, treatment may lead to large time survival. Indeed, closer the system gets to the cure equilibrium, longer it takes to tumor recurrence be observable. This fact is related with the large time necessary to pass through a saddle-point. In this case, an important feature will be the slope of the basin of attraction of the cancer equilibrium. A flat basin implicates in a large time to tumor recurrence.

On the other hand, even if the system has a cancer cure stable equilibrium, as in case II here, the treatment may be ineffective if the solution does not achieve the basin of attraction of the cure equilibrium. It is the case of simulation with $n = 4$ in case II, shown in Figure 4. In other words: a necessary condition to a treatment be effective is that the underlying system must have a stable cancer cure equilibrium; and the sufficient condition for the treatment be effective is that the treatment must move the trajectory to the basin of attraction of this equilibrium




Figure 4: Solutions of system (7) when parameters correspond to regime II, with the number of doses given by $n = 4$ (continuous), $n = 5$ (dashed), $n = 6$ (dot-dashed) and $n = 7$ (dotted). Initial conditions were $(A(0), N(0)) = P_2$, representing that the treatment was initiated only after the tumor reached the steady state P_2 . In panel (A), the small black (large blue) numbers on phase portrait indicate the time in months (weeks) in which the solution was at each point. In panels (A) and (B) the gray dotted curve represents the separatrix between the basins of attraction of P_0 and P_2 .

(simulations with $n = 5, 6, 7$ in case II). Once it has been reached, the treatment can stop, since the own patient repair system will eliminate the reminiscent cancer cells, and move the trajectory in direction to the cure equilibrium. Thus, the resilience of cancer equilibrium plays an important role: if this equilibrium has a large and deep basin, and is located at a large distance from the basin boundary, then more doses, or more intense doses, will be necessary in order to make the treatment be effective. Further, it also suggests a mechanism through which two individuals with similar diagnosis and treatments may have different fates: some of those treatments which end very near the separatrix (simulation with $n = 5$ in Figure 4) may become ineffective due to stochastic fluctuations which can drive the system to the cancer basin again, while other may continue in the cure basin. This indicates that truly effective treatments should drive the system to a safe distance from the boundary of the basin. This rationale agrees with the fact that treatments which consist of single surgery or radiotherapy (which would correspond to large state space disturbances) may be reinforced if followed by adjuvant treatment in order to preclude tumor relapse [37].




Figure 5: Solutions of system (7) when parameters correspond to regime III, with the number of doses given by $n = 4$ (continuous), $n = 5$ (dashed), $n = 6$ (dot-dashed) and $n = 7$ (dotted). Initial conditions were $(A(0), N(0)) = P_2$, representing that the treatment was initiated only after the tumor reached the steady state P_2 . In panel (A), the small black (large blue) numbers on phase portrait indicate the time in months (weeks) in which the solution was at each point.

5 Resilience Analysis

The ecological point of view on cancer discussed above is illustrated through the stability landscape in Figure 6. Of course, this view oversimplifies the reality of cancer, since there are many intermediary stages in the disease progression, but it provides at least a theoretical framework which sheds some light on the understanding of begin and the end of the disease. The better view of this landscape is that where it is not static, but a shaking landscape, due to the stochastic disturbances occurring all the time both on parameters and on state variables [38]. Also, some features discussed above may not be correctly represented in Figure 6, where one can easily think that the escape from the cancer cure equilibrium in case III, a saddle point, will be as fast as a ball falls from the top of the hill, which is indeed false.

In the stability landscape in Figure 6, the creation of the cancer equilibrium and the loss of stability by the cancer cure equilibrium (large red and orange arrows) is achieved by the sequential acquiring of genetic alterations that improve the fitness of cancer cells (the hallmarks of cancer [9]), both by deregulating mechanisms of control or by creating mechanisms which favor cancer cell functions. The transition from I to III (large orange arrow) occurs only if tumor aggressiveness is low. Destruction of cancer equilibrium and creation of stability of the cancer cure equilibrium (opposite directions of large red and orange arrows) will be achieved only by a permanent change in the intrinsic system, through, for instance, restoring the control




Figure 6: An ecological view on cancer onset and treatment as the switching between two states, a cancer cure state (positions of blue balls) and a cancer state (positions of red balls). Perturbations on state variables are due to exposure to external carcinogenic factors and genetic instability (small red arrows), which favor the moving in the direction to the cancer state, or to chemotherapy, radiotherapy and surgery (small blue arrows), which move the system towards the cancer cure equilibrium. Perturbations in parameters (large arrows) lead to transitions between the three different regimes (I, II and III), and may create or destroy equilibria, or lead to changes of stability in these equilibria, what can makes impossible either the cure or the onset of cancer.

systems (immunostimulation for example) or by limiting cancer cells functions (continuous anti-angiogenic treatments for example, which would decrease the value of K_A in theory). Finally, combination of small changes in parameters with perturbations on state variables also have a fundamental role, since the former can diminish the basin of attraction while the latter push the system to cross the basin boundary.

Let us analyze the effect of parameter changes on the resilience of stable equilibria of system (1), i.e., on size and shape of their basins of attraction. To perform this analysis we briefly review the measurements which may be of importance when analyzing the resilience of an equilibrium [8, 6, 7, 5]. As far as we know, very few papers have dealt with this kind of analysis for population dynamics models [8, 28, 29], although it is not an uncommon approach for systems modeling power grids [5]. Recently, Mitra *et al.* applied these measures to the nonlinear pendulum, the daisy-world model, and to an one-dimensional model of desertification in Amazon forest [8]. Also, as far as we know, the methods provided here to calculate these measures are novel and more efficient than the currently used ones, which consist in integrating the system in many points of the phase space, assigning each point to some basin of attraction. Throughout this section, we use the notation $\mathbb{R}^2 = \{X = (N, A); N, A \in \mathbb{R}\}$.

5.1 The latitude $L(P)$ of an equilibrium point

The *latitude* of an equilibrium point corresponds to the volume of the basin of attraction. It measures the resilience of that equilibrium with respect to state-space perturbations. Larger

the latitude of an equilibrium, smaller is the chance that external or probabilistic events will drive the system outside the basin of attraction. For two dimensional systems, the latitude corresponds to the area of the basin of attraction. However, as it may happens that the basin of attraction has infinite area, it may be the case to consider its area inside a relevant bounded region. In the case of system (1), all trajectories remain in the box B given in (6). This box could be this bounded region of interest. However, biological relevant perturbations of $P_2 = (N_2, A_2)$ will diminish the value of both coordinates, and relevant perturbations of $P_0 = (r_N/\mu_N, 0)$ will diminish the first coordinate and increase the second. Thus, we consider the smallest box C which contains P_0 and P_2 as our bounded region of interest. It is clear that

$$C = [0, r_N/\mu_N] \times [0, A_2].$$

Thus, if $\mathcal{A}(P)$ denotes the basin of attraction of P , then the latitude of P in our case is defined by

$$L(P) = \frac{\text{Area}(\mathcal{A}(P) \cap C)}{\text{Area}(C)}. \quad (8)$$

We divide by the total area of C to obtain a non-dimensional quantity normalized between 0 and 1. In the case when system presents bistability (region II of parameters space) it is clear that

$$L(P_0) + L(P_2) = 1.$$

Extreme cases are regions I and III. In region I, when P_0 is globally stable, we have

$$L(P_0) = 1, \quad L(P_2) = 0,$$

In region III, when P_2 is globally stable,

$$L(P_0) = 0, \quad L(P_2) = 1.$$

We present a simple and efficient method to calculate $L(P)$ for two dimensional systems. It consists of two steps. The first step is to obtain the terms of a series expansion for a parametric representation of the stable manifold of saddle point in a two-dimensional system. The second step involves the use of Green's Theorem to transform the area of the basin of attraction in a line integral calculated along the stable manifold approximated in the first step.

Summary of first step Let X^* be a saddle point of the two-dimensional system

$$X' = F(X), \quad X \in \mathbb{R}^2,$$

where F is a C^1 vector field. In the vicinity of X^* , this system can be re-written as

$$X' = J(X - X^*) + G(X),$$

where $G(X) = \mathcal{O}(\|X - X^*\|^2)$ and $J = F'(X^*)$ is the Jacobian matrix evaluated at X^* . Let $J = MKM^{-1}$ its Jordan decomposition, with $k_{11} < 0 < k_{22}$ (which are real since X^* is a saddle-point). With the change of coordinates

$$U = M^{-1}(X - X^*),$$

the previous system becomes

$$U' = KU + R(U),$$

where $R(U) = M^{-1}G(X^* + MU) = \mathcal{O}(\|U\|^2)$. This system can be written in coordinates $U = (u, v)$ as

$$u' = f(u, v), \quad v' = g(u, v).$$

The origin is a saddle point for this system. Its stable manifold is tangent to $v = 0$, and is locally the graph of a function $v = p(u)$. Substituting it into $v' = g(u, v)$ we obtain a nonlinear ODE for $p(u)$:

$$p'(u)f(u, p(u)) = g(u, p(u)).$$

Although this ODE cannot be solved in general, we can write a series expansion

$$p(u) = c_2u^2 + c_3u^3 + \dots$$

and solve term by term, obtaining the coefficients c_j recursively. For large k , the truncated polynomial $p_k(u) = c_2u^2 + \dots + c_ku^k$ provides a good approximation of the stable manifold near the origin. Backing to the original coordinates,

$$s_k(u) = X^* + M(u, p_k(u))^T$$

is a parametric approximation of the stable manifold of X^* , for small $\|u\|$.

Summary of the second step Now we explain how to calculate

$$L(P) = \frac{\text{Area}(\mathcal{A}(P) \cap C)}{\text{Area}(C)} = \frac{\iint_{\mathcal{A}(P) \cap C} dAdN}{\iint_C dAdN}. \quad (9)$$

The infinitesimal area element is denoted by $dAdN$. The integral in the denominator is easy to calculate in general; in our case C is a rectangle. The difficult lies in calculating the integral in the numerator. However, it can be easily calculated by using the approximation of the stable manifold obtained in the previous step and the Green's Theorem. Indeed, the boundary of the region $\mathcal{R} = \mathcal{A}(P) \cap C$ is formed by four curves, $\partial\mathcal{R} = S \cup L_1 \cup L_2 \cup L_3$, where S is the part of the stable manifold of P_1 which is contained in C , and L_i , $i = 1, 2, 3$, are line segments contained in the boundary of C . By Greens' Theorem, the integral can be written as

$$\iint_{\mathcal{A}(P) \cap C} dAdN = \oint_{\partial\mathcal{R}} NdA = \int_S NdA + \sum_{i=1}^3 \int_{L_i} NdA,$$

with the correct orientation defined for these curves. The summation terms are easily computed, and a good approximation for the integral in S is given by using the parametric approximation obtained in the first step. With this approach, we have a very good approximation of $L(P)$.

Let us comment on the applicability of this method. First of all, the method works when the boundary of the basin is formed by invariant manifolds of saddle points [39]. If many saddle points lie in the boundary, then the first step needs to be applied to each point. Second, the bounded region C must be such that this area is easily calculated. These first two requirements are very general [39]. The last and most restrictive condition is that, for each saddle point on the boundary, the local approximation obtained in the first step must be a good approximation in all points inside C . The method fails when this requirement is not satisfied. In our case, for parameters values in Table 1, the approximation for the stable manifold of P_1 with 25 terms was obtained in a few seconds by *Mathematica*[®], and provided a very good approximation for points inside C . The boundary of the basin in Figures 2, 4 and 7 was plotted with this approximation.

5.2 The precariousness $Pr(P)$ of an equilibrium point

Another important measure is the *precariousness* of an equilibrium point. Roughly speaking, it is defined as the minimum perturbation required to drive the system to another basin of

attraction. As a basin of attraction of interest may be large but the equilibrium may be located near the boundary, the precariousness is an important measure. It can be defined as

$$Pr(P) = \inf\{\text{dist}(P, X), X \in \partial\mathcal{A}(P)\},$$

where $\partial\Omega$ stands for the boundary of the set Ω , and $\text{dist}(X, Y)$ is the Euclidean distance. However, it may happens that perturbations of relevance may not occur in the direction where this minimum distance is achieved. Thus, one can consider

$$Pr(P) = \inf\{\text{dist}(P, X), X \in \partial\mathcal{A}(P) \cap Z(P)\},$$

where $Z(P)$ is a specified set containing the relevant directions for perturbations from P [6]. For system (1), all relevant perturbations of P_2 will diminish both coordinates. Thus we define

$$Pr(P_2) = \inf\{\text{dist}(P_2, X), X \in \partial\mathcal{A}(P_2) \cap Z(P_2)\}, \quad Z(P_2) = [0, N_2] \times [0, A_2]. \quad (10)$$

Let us now consider which are the relevant perturbations of P_0 . As new cancer cells may arise in the tissue, we may consider perturbations which increase the value of A . Thus, it would be the case to consider a single direction, $\vec{v}_1 = (0, 1)$. However, as cancer cells arise during mitosis, roughly speaking, the number of normal cells diminish by one when a cancer cell arise. Thus, it would be the case to consider the direction given by $\vec{v}_2 = (-1, 1)$. Therefore, we consider all directions between \vec{v}_1 and \vec{v}_2 , what give us the set

$$Z(P_0) = \{(N, A) \in \mathbb{R}_+^2, N + A \geq r_N/\mu_N\}.$$

Thus, the precariousness of P_0 is given by

$$Pr(P_0) = \inf\{\text{dist}(P_0, X), X \in \partial\mathcal{A}(P_0) \cap Z(P_0)\}. \quad (11)$$

An illustration of regions $Z(P_1)$ and $Z(P_2)$ can be seen in Figure 7.

In extreme cases I and III we do not have bistability, and $Pr(P)$ must be defined appropriately. In case III, when P_2 is globally stable, we define $Pr(P_2)$ as the tumor volume, $Pr(P_2) = A_2$, meaning that the removal of the entire tumor leads the system outside the basin of attraction of P_2 . Indeed, this removal leads the system to the N -axis, which is the invariant manifold of P_0 . Analogously, in case I, when P_0 is globally stable, we define $Pr(P_0) = N_0$. However, in this case the A -axis is not invariant, and another choice would be $Pr(P_0) = \infty$.

The result obtained in the first step of the previous section can be used to calculate the precariousness of an equilibrium straightforwardly. In our case, let be $s_k(u) = (s_k^1(u), s_k^2(u))$ the approximation for the stable manifold of P_1 obtained in that step. To calculate $Pr(P_0)$, for instance, we first must find the interval I_0 of values of u such that $s_k(u) \in Z(P_0)$. By the definition of $Z(P_0)$, this can be done solving equations

$$s_k^{(1)}(u) = r_N/\mu_N$$

and

$$s_k^{(1)}(u) + s_k^{(2)}(u) = r_N/\mu_N$$

for u . The solutions are the extrema of interval I_0 . With this, $Pr(P_0)$ is obtained by solving the minimization problem

$$\min_{u \in I_0} \|s_n(u) - P_0\|$$

which can be easily solved. The precariousness of P_2 is calculated in an analogous way.




Figure 7: Phase portrait of system (1). $Z(P_2)$ is the box with P_2 in one of the vertices. $Z(P_0)$ is the triangular region with P_0 in its vertices. The blue points are the nearest points of P_0 and P_2 inside regions $Z(P_0)$ and $Z(P_2)$ respectively. The intensity of color on background correspond to the value of the local resistance $R(N, A)$ defined in sub-section 5.3.

5.3 The resistance $R(P)$ of an equilibrium point

Finally, the third important measure concerning resilience of an equilibrium is termed as the *resistance* of this equilibrium. It refers to “the ease or difficulty of changing the system, related to the topology of the basin - deep basins of attraction indicate that greater forces or perturbations are required to change the current state of the system away from the attractor” [7]. Thus, a resistant system will overcome perturbations rapidly, while a system with small resistance can be driven to a basin transition through a series of small perturbations that are not absorbed enough. This description concerns exactly the illustration provided by Figure 4. There, if P_2 was more resistant, more chemotherapeutic doses, or more intense doses, would be necessary to drive the system to the basin of attraction of P_0 .

The task of characterizing the sizes and frequencies of perturbations that a basin of attraction can absorb is currently a research area [6]. Recently, Mitra and others [8] proposed an approach to measure the resistance of a point in state-space to local perturbations using local Lyapunov exponents [40]. Again, consider the two-dimensional system

$$X' = F(X),$$

where F is a C^1 vector field. The *instantaneous Jacobian matrix* at X is defined as

$$\mathcal{J}_{dt}(X) = I_{2 \times 2} + J(X)dt,$$

where $J(X)$ is the Jacobian at X and dt is a infinitesimal time. Let $\sigma_i(X)$, $i = 1, 2$, the square roots of the eigenvalues of the right Cauchy-Green tensor $\mathcal{J}_{dt}(X)^T \mathcal{J}_{dt}(X)$. The $\sigma_i(X)$'s are also the singular values of $\mathcal{J}_{dt}(X)$. They measure the instantaneous stretching of the neighborhood of the trajectory at X . The local Lyapunov exponents evaluated at the state X are defined as

$$\lambda_i(X) = \frac{1}{dt} \ln(\sigma_i(X)), \quad i = 1, 2.$$

and measure the rate of stretching at X . The local resistance $R(N, A)$ at the state $X = (N, A)$ may be defined as

$$R(X) = -\max\{\lambda_1(X), \lambda_2(X)\}. \quad (12)$$

Thus, the resistance of an equilibrium point can be measured as

$$Res(P) = \frac{\iint_{\mathcal{A}(P) \cap C} R(N, A) dNdA}{\iint_C R(N, A) dNdA}. \quad (13)$$

The division by the total resistance on C is made to obtain a non-dimensional quantity normalized between 0 and 1.

For system (1), analytical expressions for $\lambda_i(X)$, $i = 1, 2$, at state $X = (N, A)$ can be calculated in *Mathematica*[©]. The density plot of $R(N, A)$ is showed in Figure 7. With these expressions, we can calculate integrals in (13). The unique concern is with respect to the region of integration in the first integral, $\mathcal{A}(P) \cap C$. However, this integral can be calculated using the approximations for the stable manifold of P_1 obtained in the first step of 5.1.

5.4 Application of resilience analysis to system (1)

We analyze the behavior of the above measures when parameters of system (1) vary. Figure 8 shows the results when β_3 varies while other parameters are kept constant, with $\beta_1 > \beta_1^{th}$, which corresponds to bifurcation diagram (ii) in Figure 3 (center), where a transition I-II-III is observed. Results when other parameters vary are similar (not shown here).

To discuss these results, we first consider the point of view of cancer onset and analyze the resilience measures of P_0 . In region I, P_0 is globally stable and all these measures are equal to the unity. When β_3 becomes lesser than $\beta_{3,\Delta}^{th}$ and enters region II, equilibrium P_0 is no longer globally stable, and its resilience measures undergo an abrupt jump and decay rapid in a small strip near $\beta_{3,\Delta}^{th}$. The most notorious jump occurs with $Pr(P_0)$. For values in the midpoint between the two thresholds separating region II from regions I and III, the values of $Pr(P_0)$ and $L(P_0)$ are very small. These features are due to the up-concave shape of the graphs of $L(P_0)$, $Pr(P_0)$ and $Res(P_0)$, and indicate that in the bistable regime the healthy state P_0 is threaten by small disturbances which may easily drive the system to the basin of attraction of P_2 .

On the other hand, let us consider the point of view of treatment, and discuss the results concerning P_2 . In regime III, this equilibrium is globally stable and $L(P_2)$, $Pr(P_2)$ and $Res(P_2)$ are equal to the unity. As β_3 becomes larger than β_3^{th} , these measures decay very slowly and are greater than the respective measures of P_0 , until β_3 reaches the small strip near the next threshold, $\beta_{3,\Delta}^{th}$. Now, these features are due to the down-concave shape of the graphs, and implicate that P_2 is relatively protected against small perturbations.

By comparing these differences on the resilience measures of P_0 and P_2 we conclude that it is much more easy to drive the system outside the basin of attraction of the cure equilibrium P_0 when it loses its global stability, than driving the system out the basin of attraction of the cancer equilibrium P_2 when it reaches the bistable regime, unless the parameters get very near the next bifurcation threshold at which P_2 loses its stability. In other words, our analysis reveal that, in the bistable regime, although these are different phenomena, it is more likely to mutations or exposure to carcinogenic factors drive cancer onset than chemotherapy, surgery or radiotherapy lead to tumor regression.

6 Conclusion

In this paper, an ecological resilience framework to think of cancer as the alternance between two states is presented. This framework is based on the analysis of a simple ODE model for tumor growth considering the interaction with the host tissue. Despite the simplicity of the




Figure 8: Values of $L(P_i)$ (top), $Pr(P_i)$ (center) and $Res(P_i)$ (bottom), $i = 0, 2$, as β_3 varies, with $\beta_1 > \beta_1^{\text{th}}$, which corresponds to the bifurcation diagram (ii) in Figure 3 (top, right). As β_3 varies, we observe transitions between regimes III, II and I.

model, the approach adopted here gives interesting theoretical insights that shed some light on several relevant issues concerning cancer onset and treatment, and may help to improve the way we view cancer.

The model exhibits three regimes. These regimes are used to illustrate three different possibilities which may occur in clinical cases in general. The first regime corresponds to a healthy person where cancer onset is not possible since the cancer cure state is globally stable. The second regime corresponds to a person which can develop cancer if exposed to external carcinogenic factors, due to a partially corrupted repair system and/or a high aggressive phenotype of tumor cells. This regime presents bistability between the cancer and the cancer cure states. The third regime corresponds to a person in which cancer will arise due to intrinsic factors, i.e., the total corruption of repair systems. In this regime, the cancer state is globally

stable.

Based on the general property that treatments are finite and, therefore, do not change the global dynamics in the phase space, we discuss the possibility of cure, which concerns stability and resilience questions above all. In the bistable regime the cure is possible if the treatment is able to drive the system to the basin of attraction of the cure equilibrium. Tumor recurrence in this case is associated with treatments which are unable to cross the separatrix between the basins, or which do not end at a safe distance from the separatrix. In the third regime, the complete cure is not possible at all, since the repair system is intrinsically weak, but tumor recurrence may be delayed if the treatment is prolonged, because the system takes a long time to pass around the cure equilibrium, which is a saddle point. However, toxicity is not assessed in this model.

Besides perturbations on state variables, a view in the switching between these three regimes due to parameters changes in a slow time scale is discussed and the roles of the most important parameters in these transitions are assessed. Results indicate that only aggressive tumors may arise if intrinsic repair systems are not totally corrupted. Further, these aggressive tumors depends on exposure to external carcinogenic factors for arising.

Finally, we review the use of three different measures to assess the resilience of a stable equilibrium. We propose simple and efficient methods to calculate these measures. After applying this analysis to the model we conclude that in the bistable regime the cancer equilibrium has much more resilience than the cure equilibrium, with respect to state variable perturbations as parameters changes.

This paper contributes to the current understanding on cancer by raising some issues in an ecological view, and also demonstrates how a ‘resilience analysis’ may be applied to population dynamics models in order to improve the understanding of nonlinear phenomena. Further, the application of the ‘resilience analysis’ presented here to detailed and accurate models for specific tumor types has the potential to generate measures which can be accounted for in the design of treatment plans for cancer patients and also in the development of adaptive treatments [13]. The size and shape of the basin of attraction of the cancer equilibria in those models may be used as indicators in the design and planning of personalized treatments.

References

- [1] Crawford S Holling. “Resilience and stability of ecological systems”. In: *Annual Review of Ecology and Systematics* (1973), pp. 1–23.
- [2] Robert M May. “Thresholds and breakpoints in ecosystems with a multiplicity of stable states”. In: *Nature* 269.5628 (1977), pp. 471–477.
- [3] Marten Scheffer et al. “Catastrophic shifts in ecosystems”. In: *Nature* 413.6856 (2001), pp. 591–596.
- [4] Carl Folke. “Resilience: The emergence of a perspective for social–ecological systems analyses”. In: *Global Environmental Change* 16.3 (2006), pp. 253–267.
- [5] Peter J Menck et al. “How basin stability complements the linear-stability paradigm”. In: *Nature Physics* 9.2 (2013), pp. 89–92.
- [6] Katherine Meyer. “A Dynamical Systems Framework for Resilience in Ecology”. In: *arXiv preprint arXiv:1509.08175* (2015).
- [7] Brian Walker et al. “Resilience, adaptability and transformability in social–ecological systems”. In: *Ecology and Society* 9.2 (2004), p. 5.

- [8] Chiranjit Mitra, Jürgen Kurths, and Reik V Donner. “An integrative quantifier of multistability in complex systems based on ecological resilience”. In: *Scientific Reports* 5 (2015).
- [9] Douglas Hanahan and Robert A Weinberg. “Hallmarks of cancer: the next generation”. In: *Cell* 144.5 (2011), pp. 646–674.
- [10] Cristian Tomasetti and Bert Vogelstein. “Variation in cancer risk among tissues can be explained by the number of stem cell divisions”. In: *Science* 347.6217 (2015), pp. 78–81.
- [11] Song Wu et al. “Substantial contribution of extrinsic risk factors to cancer development”. In: *Nature* 529.7584 (2016), pp. 43–47.
- [12] Robert S Kerbel and Barton A Kamen. “The anti-angiogenic basis of metronomic chemotherapy”. In: *Nature Reviews Cancer* 4.6 (2004), pp. 423–436.
- [13] Sébastien Benzekry et al. “Metronomic reloaded: theoretical models bringing chemotherapy into the era of precision medicine”. In: *Seminars in Cancer Biology*. Vol. 35. Elsevier. 2015, pp. 53–61.
- [14] Artur C Fassoni. “Mathematical modeling in cancer addressing the early stage and treatment of avascular tumors”. PhD thesis. Unicamp, 2016.
- [15] Robert A Gatenby. “Models of tumor-host interaction as competing populations: implications for tumor biology and treatment”. In: *Journal of Theoretical Biology* 176.4 (1995), pp. 447–455.
- [16] Robert A Gatenby and Edward T Gawlinski. “A reaction-diffusion model of cancer invasion”. In: *Cancer Research* 56.24 (1996), pp. 5745–5753.
- [17] Lisette G de Pillis and Ami Radunskaya. “A mathematical tumor model with immune resistance and drug therapy: an optimal control approach”. In: *Computational and Mathematical Methods in Medicine* 3.2 (2001), pp. 79–100.
- [18] Jessica B McGillen et al. “A general reaction–diffusion model of acidity in cancer invasion”. In: *Journal of Mathematical Biology* 68.5 (2014), pp. 1199–1224.
- [19] Benjamin D Simons and Hans Clevers. “Strategies for homeostatic stem cell self-renewal in adult tissues”. In: *Cell* 145.6 (2011), pp. 851–862.
- [20] P Fedi, SR Tronick, and SA Aaronson. “Growth factors”. In: *Cancer Medicine* (1997), pp. 41–64.
- [21] Elizabeth A Sarapata and LG de Pillis. “A comparison and catalog of intrinsic tumor growth models”. In: *Bulletin of Mathematical Biology* 76.8 (2014), pp. 2010–2024.
- [22] Nika N Danial and Stanley J Korsmeyer. “Cell death: critical control points”. In: *Cell* 116.2 (2004), pp. 205–219.
- [23] Robert S Kerbel. “Tumor angiogenesis”. In: *New England Journal of Medicine* 358.19 (2008), pp. 2039–2049.
- [24] Hyun M Yang. “Mathematical modeling of solid cancer growth with angiogenesis”. In: *Theoretical Biology and Medical Modelling* 9.2 (2012).
- [25] Eric Vivier et al. “Targeting natural killer cells and natural killer T cells in cancer”. In: *Nature Reviews Immunology* 12.4 (2012), pp. 239–252.
- [26] Robert A Gatenby et al. “Acid-mediated tumor invasion: a multidisciplinary study”. In: *Cancer Research* 66.10 (2006), pp. 5216–5223.
- [27] Andrea Facciabene, Gregory T Motz, and George Coukos. “T-regulatory cells: key players in tumor immune escape and angiogenesis”. In: *Cancer Research* 72.9 (2012), pp. 2162–2171.

- [28] Artur C Fassoni, Lucy T Takahashi, and Laércio J dos Santos. “Basins of attraction of the classic model of competition between two populations”. In: *Ecological Complexity* 18 (2014), pp. 39–48.
- [29] Artur C Fassoni and Marcelo L Martins. “Mathematical analysis of a model for plant invasion mediated by allelopathy”. In: *Ecological Complexity* 18 (2014), pp. 49–58.
- [30] Vladimir A Kuznetsov et al. “Nonlinear dynamics of immunogenic tumors: parameter estimation and global bifurcation analysis”. In: *Bulletin of Mathematical Biology* 56.2 (1994), pp. 295–321.
- [31] Lisette G de Pillis, Ami E Radunskaya, and Charles L Wiseman. “A validated mathematical model of cell-mediated immune response to tumor growth”. In: *Cancer Research* 65.17 (2005), pp. 7950–7958.
- [32] Raluca Eftimie, Jonathan L Bramson, and David JD Earn. “Interactions between the immune system and cancer: a brief review of non-spatial mathematical models”. In: *Bulletin of Mathematical Biology* 73.1 (2011), pp. 2–32.
- [33] Robert Stephen Cantrell and Chris Cosner. “Practical persistence in ecological models via comparison methods”. In: *Proceedings of the Royal Society of Edinburgh: Section A Mathematics* 126.02 (1996), pp. 247–272.
- [34] Steven H Strogatz. *Nonlinear dynamics and chaos: with applications to physics, biology and chemistry*. Perseus publishing, 2001.
- [35] Sabrina L Spencer et al. “An ordinary differential equation model for the multistep transformation to cancer”. In: *Journal of Theoretical Biology* 231.4 (2004), pp. 515–524.
- [36] Frank M Schabel. “Concepts for systemic treatment of micrometastases”. In: *Cancer* 35.1 (1975), pp. 15–24.
- [37] Roger Stupp et al. “Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma”. In: *New England Journal of Medicine* 352.10 (2005), pp. 987–996.
- [38] Marten Scheffer et al. “Generic indicators of ecological resilience: inferring the chance of a critical transition”. In: *Annual Review of Ecology, Evolution, and Systematics* 46 (2015), pp. 145–167.
- [39] Hsiao-Dong Chiang, Morris W Hirsch, and Felix F Wu. “Stability regions of nonlinear autonomous dynamical systems”. In: *Automatic Control, IEEE Transactions on* 33.1 (1988), pp. 16–27.
- [40] Henry DI Abarbanel, Reggie Brown, and Matthew B Kennel. “Variation of Lyapunov exponents on a strange attractor”. In: *Journal of Nonlinear Science* 1.2 (1991), pp. 175–199.