

A DRAFT GENOME ASSEMBLY OF SOUTHERN BLUEFIN TUNA *THUNNUS MACCOYII*

S. MCWILLIAM

CSIRO Agriculture, 306 Carmody Road, St. Lucia 4067, Australia

P. M. GREWE

CSIRO Oceans and Atmosphere, Castray Esplanade, Hobart, Tasmania, Australia

AND

R. J. BUNCH AND W. BARENDSE¹

CSIRO Agriculture, 306 Carmody Road, St. Lucia 4067, Australia

arxiv preprint—not yet submitted

ABSTRACT

Tuna are large pelagic fish whose populations are close to panmixia. In addition, they are threatened species, so it is important for the maintenance and monitoring of genetic diversity that genetic information at a genome level be obtained. Here we report the draft assembly of the southern bluefin tuna genome and the collection of genome-wide sequence data for five other tuna species. We sampled five tuna species of the genus *Thunnus*, the northern and southern bluefin, yellowfin, albacore, and bigeye, as well as the skipjack (*Katsuwonis pelamis*), a tuna-like species. Genome assembly was facilitated at k-mer=25 while k-mer=51 generated assembly artefacts. The estimated size of the southern bluefin tuna genome was 795 Mb. We assembled two southern bluefin tuna individuals independently using both paired end and mate pair sequence. This resulted in scaffolds with N50>174,000 bp and maximum scaffold lengths>1.4 Mb. Our estimate of the size of the assembled genome was the scaffolded sequences in common to both assemblies, which amounted to 721 Mb of the 795 Mb of the southern bluefin tuna genome sequence. Using BLAST, there were matches between 13,039 of 14,341 (91%) refseq mRNA of the zebrafish *Danio rerio* to the tuna assembly indicating that most of a generic fish transcriptome was covered by the assembly.

Subject headings: genome; DNA; tuna

1. INTRODUCTION

The annual wild tuna harvest is of the order of 4 million tonnes per year and, with the decline or stringent management of fisheries, monitoring of stocks and movement of animals is important to the long term sustainability of tuna fisheries (Campbell *et al.* 2000; MacKenzie *et al.* 2009; Group 2014). True tuna are the species in the genus *Thunnus*, such as bluefin (Atlantic (*T. thynnus*), northern Pacific (*T. orientalis*), and southern (*T. maccoyii*)), yellowfin (*T. albacares*), albacore (*T. alalunga*), and bigeye (*T. obesus*). Many of these species are threatened or critically endangered and are being replaced as a food source by related ‘tuna’ fish such as the skipjack (*Katsuwonis pelamis*).

Genetic studies of these species have a long history and have shown that, as expected, the populations of these species are close to panmixia. In general, pelagic marine fish have low F_{ST} values across ocean-wide samples and tuna species are no exception (Elliott & Ward 1992; Ward *et al.* 1995; Takagi *et al.* 1999; Ward 2000; Appleyard *et al.* 2001; Terol *et al.* 2002; Martinez *et al.* 2006; Rooker *et al.* 2007; Lowenstein *et al.* 2009; Riccioni *et al.* 2010; Montes *et al.* 2012). Such panmixia makes monitoring difficult if there are only a small number of DNA markers and so it is important that a large number of polymorphisms are available for these species.

One option for generating large numbers of polymorphisms is to use short read genome sequencing (Bentley *et al.* 2008). Initial genome sequencing of a tuna species has occurred, for the northern Pacific bluefin tuna and the existence of a preliminary assembly has been reported (Nakamura *et al.* 2013). Single nucleotide polymorphisms (SNP) for monitoring populations have been identified in yellowfin (Grewe *et al.* 2015; Pecoraro *et al.* 2016) and DNA based methods for identification of tuna species have recently been published (Takashima *et al.* 2014; Santaclara *et al.* 2015). Next generation sequencing has also recently been used to obtain the complete mitochondrial sequence of two tuna species (Chen *et al.* 2016; Li *et al.* 2016), to start the identification of microsatellites across the Scombridae (Nikolic *et al.* 2015), and to develop functional genomics tools for the study of tuna in aquaculture (Yasuike *et al.* 2016). Here we report additional genome sequence data for a set of tuna species particularly the southern bluefin tuna (SBT). Our main objective is to provide genome sequence information that might contribute to the identification of SNPs, to make tools to monitor tuna fisheries, and to generate public data to improve tuna genome assemblies. Other objectives are to quantify, where possible, the genetic differences between species revealed through genome sequencing and to provide a resource for the further genomic study of pelagic ocean fish. In this study we report the collection of tissue samples, of genome sequence for the six species, and the draft assembly of the southern bluefin tuna genome.

bill.barendse@gmail.com

¹ School of Veterinary Science, The University of Queensland, Gatton 4343, Australia.

2. MATERIALS AND METHODS

2.1. DNA samples

Tissue samples of tuna fish were obtained from wild caught tuna as part of commercial, authorised fishing quotas. The various tuna fish are easy to discriminate from each other using colour, length, and shape of dorsal and pectoral fins, colour of the back and sides, size of eyes, presence or absence of body stripes (Collette *et al.* 2001; Itano 2005; Joshi *et al.* 2012) and we visually checked the species identity of each individual (see www.iotc.org/science/species-identification-cards checked 7 July 2016). Samples were taken post mortem, chilled, and then frozen once they reached the laboratory. DNA for genome sequencing was purified using a modified CTAB (cetyl trimethylammonium bromide) method (Doyle & Doyle 1987).

DNA for genome sequencing was checked for quality control using agarose gel electrophoresis, followed by quantification, DNA fragmentation, and size selection using a Nanodrop, a Covaris sonicator, and Qubit-HS DNA kit, as described (Barendse *et al.* 2015). DNA quality was determined by running a subsample on a 1% TAE agarose gel to check for DNA smears due to DNA degradation and the 260:280 nm absorbance ratio on the Nanodrop was required to be between 1.8 and 2.0. Each sample was normalized to 20 ng/ μ l and 55 μ l was fragmented to an average insert length of 300 to 400 bp using a Covaris S220 ultra-sonicator (Woburn, MA, USA).

2.2. DNA libraries and short sequence reads

DNA libraries for sequencing were prepared from at least 1.0 μ g of genomic DNA of each individual using the Illumina TruSeq DNA Sample Prep Kit v2-Set A kit following the manufacturers instructions (Illumina Australia P/L, Scoresby, Vic). A TruSeq PE Cluster Kit v3 was used to generate sequencing clusters using the Illumina cBot on Flow Cell v3. 100 bp paired end (PE) reads were obtained using the Illumina TruSeq SBS Kit v3 - HS kit following the manufacturers instructions on the SQ module of an Illumina HiScan instrument or a HiSeq instrument. Individuals were multiplexed at 8 samples per lane. Two of the southern bluefin samples were sequenced using the Illumina Nextera Mate Pair Library Kit (MP) following the manufacturer's instructions at insert sizes of 2 kb, 5 kb, and 8 kb.

2.3. Sequence analysis and genome assembly

Initial quality control of DNA sequence reads followed a standard genome analysis pipeline (Barendse *et al.* 2015), except that there was no reference genome sequence for any tuna species. Initial quality control consisted of removing sequences that failed chastity filtering using Illumina Casava 1.8. Sequences passing chastity filtering were quality trimmed to remove bases below phred Q20. Using the fastq trimmer module of the Fastx toolkit (http://hannonlab.cshl.edu/fastx_toolkit/command-line.html) sequences less than 50 bp following trimming were removed from further analysis. The k-mer distribution of the reads was evaluated using Jellyfish (Marcais & Kingsford 2011) and used to estimate the genome size of tuna. All short sequence reads of tuna fish that survived trimming were assembled using SOAPdenovo2

v2.04 (Li *et al.* 2010; Luo *et al.* 2012) at the CSIRO High Performance Cluster at two different k-mer sizes, 25 and 51. A small k-mer size was examined to cope with the potentially large number of DNA polymorphisms that could be encountered in pelagic marine species with large population sizes. Contigs reported as part of the assembly had lengths greater than 100 bp. The assembly settings for the PE reads were average insert size 300 bp, maximum read length 100 bp, asm flags 3, pair number cutoff 3, and map length 32. Assembly settings for the MP reads were average insert size 3000 or 5000, asm flags 2, pair number cutoff 5, and map length 35. To form scaffolds, contigs greater than 200 bp were used. Using the draft SBT genome assembly as a reference, we compared the RNA or contigs of the other species to the SBT using BLAST (Altschul *et al.* 1990). Fastq format reads of the Pedrabranca3 (PDB3) and Pedrabranca5 (PDB5) individuals were lodged in the Short Read Archive associated with BioProject ID PRJNA327759 which will continue to be updated as the project progresses.

3. RESULTS

3.1. Sequence collection and genome size estimate

In total, there were 15 southern bluefin, 3 northern bluefin, 3 albacore, 3 bigeye, 4 skipjack, and 2 yellowfin individual tuna samples that passed DNA quality control and were sequenced using PE methods. The number of reads from each individual after merging and trimming is shown in Table 1, indicating for each species that there was more than 27X genome coverage. GC content of the genomes was similar, from 39.8% to 40.8%. Two of the SBT animals, PDB3 and PDB5, were chosen for additional PE and for MP sequencing based on the amount and quality of DNA. We assembled sequence reads of PDB3 by itself, of reads of PDB5 by itself, of reads of PDB3 and PDB5 combined, and of reads of all available SBT animals combined. When assembling reads of PDB3 or PDB5 by itself, only the MP sequence reads of a single individual was used, but for SBT assemblies using reads of multiple individuals, combined MP sequence of PDB3 and PDB5 was used. The 8 kb MP reads proved to have low complexity and were excluded from assemblies.

We estimated the genome size of tuna to be 795 Mb based on the k-mer distribution. A plot of the k-mer distribution is shown in Figure 1. This shows the count of 17-mers on the horizontal axis where 40 represents a 17-mer that occurs 40 times. The vertical axis shows the sum of the count of 17-mers so that the curve of the graph shows the total number of 17-mers that occurred a particular number of times. For example, the total number of all 17-mers that occurred 40 times is 13,211,979. For a k-mer length of 17, there was read coverage of 44 for the genome and a peak k-mer depth of 37.

3.2. Assembly information for species lacking MP reads

Although we did not have MP sequence for five of the species, we attempted to assemble the reads because the contigs would be useful for discovering DNA variants. In addition, once the SBT was assembled using MP reads, the contigs of the other five species obtained in this process could be aligned to that assembly better than merely matching heterologous fastq reads to the SBT assembly.

The information on these five species is shown in Table 2 which includes part of the SBT dataset as a comparison. The SBT sequences in this comparison are a mixture of all the PE sequences of PDB3 and PDB5, giving a similar dataset to the other species. In these comparisons, the SBT PE only assembly showed no special characteristics compared to the others and had similar statistics to the northern bluefin tuna. Between 67% and 74% of a tuna genome was covered by contigs of longer than 100 bp (Table 2). Estimates of coverage, N50, and maximum length of contigs or scaffolds did not appear to be strongly related to amount or type of data. The scaffolds generated in this process showed a several fold improvement in N50 and maximum length by a factor of two to five compared to contigs before scaffolding occurred. These multi-contig scaffolds covered less of the genome than the full list of contigs, and in the case of the skipjack, the coverage by scaffolds was markedly less than the true tuna species.

3.3. Draft assembly of the southern bluefin tuna

We constructed draft SBT assemblies using both PE and MP sequence using several datasets to determine the best approach for that species (Table 3). In particular we wanted to know whether increasing the number of reads and number of individuals would lead to superior results for this species. Although PDB3 did but PDB5 did not quite reach 30X coverage of the tuna genome, both showed massive reductions in scaffold number and several order of magnitude increases in scaffold length compared to lengths of contigs prior to scaffolding. This occurred whether the k-mer length was 25 or 51. As more reads from different individuals were added to the assembly, contig number increased, N50 and maximum lengths decreased, and scaffolds became shorter. This feature became worse at longer k-mer lengths. The likely cause is increased genetic variability in the dataset as more individuals were added.

Longer k-mers did not improve the sequence and may have decreased the quality of the assemblies. Although the number of contigs and the apparent coverage of the tuna genome is greater with k=51 (Table 3), this coverage exceeds the known length of the tuna genome by 18% or more than 140 Mb. Two factors may contribute to this result. Firstly, with shorter k-mers and a threshold of 100 bp for inclusion in the list of counted contigs, more short contigs are excluded with k=25. These short contigs have reduced sequence variability compared to longer contigs and may represent segments of the same repeat structure (Figure 2). Contigs such as those containing mononucleotide runs are excluded at shorter k-mer lengths. As contig length increases, the major change in the graph in Figure 2 is that the minimum diversity values increase, and at short contig lengths the low diversity values are driven by poly(A), poly(C), poly(G), and poly(T) segments. Secondly, the distribution of contig length by contig frequency showed a clear artefact in the distribution of the k=51 contigs for both PDB3 and PDB5 (Figure 3). This artefact would inflate the coverage by contigs of size 100 to 200 bp. Further analyses of the draft assemblies were restricted to k=25.

To estimate the coverage of the SBT tuna genome, first we counted the length of each assembly (Table 3). This showed values for scaffolds that exceeded the length of

the SBT genome. A BLAST analysis of the PDB3 scaffolds against the PDB5 scaffolds showed 725 Mb in common, i.e., the intersection of the two draft assemblies was 725 Mb. This is likely to be an overestimate of the coverage of the genome because there would be some short contigs that only match for 33 bp and at that level of similarity they may be referencing different parts of the genome. Excluding BLAST matches of ≤ 50 bp resulted in joint coverage of 721 Mb of the 795 Mb SBT genome. These BLAST analyses showed that alignment of the two assemblies filled sequence missing from one or the other assembly (Figure 4), sequence represented as runs of Ns in one or the other assembly. There were 143,134 segments where the BLAST alignment filled in missing sequence in this way. An additional 1,402 matching segments had gaps in the same locations in both assemblies.

3.4. Gene content of the draft SBT assembly

A BLAST analysis was used to quantify the RNA coverage of the assembly using the RNA complement of the zebrafish *Danio rerio*. The RNA sequences contained in GCF_000002035.5.GRCz10.rna.fna were matched to either assembly using BLAST. Most scaffolds showed a single match but some scaffolds in both assemblies showed more than 50 matches. The distribution of BLAST matches is the same for PDB3 compared to PDB5 (Figure 5). As increased stringency is used, fewer RNA sequences matched and the overlap between PDB3 and PDB5 decreased, as expected. Nevertheless, at a BLAST probability threshold of 1-e15 (Table 4), 45,135 of the RNAs were located to either of the two assemblies, of which 30,868 were found on both assemblies. At the more stringent threshold of 1-e40 28,461 of the RNA were located to either assembly of which 11,160 were found on both assemblies. The total number of RNA items tested from *D. rerio* was 54,437. For reference sequences beginning with 'NM_', taken to be the refseq mRNA complement, there were a total of 14,341 in the *D. rerio* set of which 13,039 were located to either of the assemblies at a threshold of 1-e15 and 8,222 were located to either of the assemblies at a threshold of 1-e40 (Table 4). In conclusion, a very large part of the transcriptome of a generic fish genome was located either totally or in part to the SBT genome.

An examination of one of the large scaffolds, scaffold269 of the PDB3 assembly, showed greater synteny between tuna and fugu (*Takifugu rubripes*) than tuna and zebrafish in that scaffold. Scaffold269 was chosen because it matched to 18 RNAs. Genome locations were sought in the zebrafish genome for these genes. The locations were Dre15: sgcg, trappc4, sacs; Dre11: foxglb, foxglc; Dre10: rps25, XM.005172584.2, XM.009305610.1, XM.699844.6; Dre18: cfap45, pgllyrp5, snrpd2, polr2i, c5ar1, vaspb, XM.001336435.5, XM.001340960.3; and Dre13: foxglc. Not all of the *D. rerio* RNA were found in fugu using a search on names. Of those that were, the locations were Tru11: sgcg, trappc4, sacs, foxglb, rps25, cfap45, snrpd2, polr2i, foxglc; Tru4: foxglc; not found in Tru: pgllyrp5, c5ar1, vaspb, XM.005172584.2, XM.009305610.1, XM.699844.6, XM.001336435.5, XM.001340960.3.

We report the genome sequencing of six tuna species and the draft genome assembly of the southern bluefin tuna. Our estimate of the tuna genome was 795 Mb, which is consistent with the 800 Mb genome size estimated using other methods (Hardie & Hebert 2004; Gregory *et al.* 2007). Although it is often recommended that assemblies be performed with larger kmers, we found that at $k=51$ artefacts were generated in smaller contigs which inflated statistics of genome coverage. Although five of the six genomes were analysed primarily to obtain DNA variants for polymorphism discovery, MP sequence was obtained to generate a *de novo* assembly of the southern bluefin tuna. The *de novo* assemblies of the SBT resulted in scaffolds with $N50 > 174,000$ bp and maximum scaffold lengths > 1.4 Mb. Comparison of the PDB3 and PDB5 assemblies showed that 721 Mb of the 795 Mb SBT genome was aligned to scaffolds and could find a matching sequence in the other alignment. Examination of the matching scaffolds showed 143,134 segments where PDB3 sequence filled in a poly(N) sequence of PDB5 and *vice versa*. Of the refseq mRNA of *Danio rerio*, 13,039 of 14,341 sequences were located to either PDB3 or PDB5, with 10,464 sequences located to both PDB3 and PDB5. This confirmed that a large proportion of the coding sequence of the SBT was contained in the scaffolds of the SBT assembly. Further annotation of the gene complement of the SBT genome will require a bioinformatics community effort. Preliminary examination of synteny in a scaffold with 18 different coding sequences showed that synteny of those genes was highly conserved to fugu but there was less conservation of synteny observed to the zebrafish. Further analyses of synteny will require detailed mapping of elements and confirmation of homology, which is beyond the scope of this communication. Such analyses will be required to demonstrate whether the conservation of synteny continued to be greater for the pair of tuna and fugu compared to the pair of tuna and zebrafish. Further analyses of the DNA variants uncovered in this study will be performed and reported in a subsequent manuscript.

We found that with increasing stringency that an RNA sequence would match to one rather than both assemblies. This is unlikely to be due to genetic variability but rather due to places in the assembly where one assembly has poly(N) where the other assembly has reportable sequence. Since the sequence that is sampled is partly dependent upon chance, the presence of reportable sequence in one animal but poly(N) in another indicates that even with around 30X genome coverage there are many areas of the genome that are counted as covered but which have short sections of unknown bases. Our results suggest that it is preferable to keep the genome data for each animal separate but that it may be advantageous to fully sequence more than one individual and compare the assemblies to fill in missing sequence. This may be because of the random nature of sequence capture, genome segments that may be well captured in one animal during one sequence run may not be well captured at some other time in a different animal. Our data do not say whether it is preferable to sequence one individual to 60X coverage compared to two individuals each to 30X coverage with separate assemblies for the two individuals. However, where the description of DNA variability is important then the sequencing of two individuals each

to 30X coverage will yield more DNA variants than sequencing one individual to 60X coverage. Our results suggest that future analyses of the other five tuna species should try to assemble each individual separately, compare the resulting assemblies using BLAST or some other tool, and then compare the resulting consensus assembly to the SBT draft assembly to construct larger scale scaffolds of those tuna genomes. This method should be used even if the total number of reads per individual does not reach the 30X level of sequence coverage.

To quantify the sequence diversity of short contigs we used the Shannon Information index because it is a standard method. The plot in Figure 3 clearly shows that as contig length increases so the occurrence of contigs with low information declines, low information in this case meaning mononucleotide runs of DNA bases or similar low diversity patterns. There are contigs where the maximum index value can be found for both short and longer contigs, but it is only among the short contigs that very low index values are found. These correspond to contigs that have for example 26 out of 26 sequential adenine nucleotides. Consequently, as contig length increased the mean index value tracked upwards towards the maximum value and the standard deviation reduced as the extremely small index values disappeared. The data show that much of this low diversity occurs for contigs below 50 bp in size, and so setting $k\text{-mer}=25$ acts as a way of flushing out low diversity contigs.

The BLAST alignment of the PDB3 and PDB5 assemblies totals 721 Mb and we take that as our estimate of genome coverage of the SBT genome. This estimate is very conservative but we see no point in including small contigs that are not strongly placed or which might represent repeat sequences that are located in multiple regions of the tuna genome. The accurate localization of such repeats, whether they are telomeres, centromeres, short interspersed nuclear elements, or other low complexity regions, requires special techniques beyond the scope of this study. To include them in the count of the percent of the genome covered would give a false impression. Much further research is needed to complete the genome assembly of a tuna species and to make it as useful as possible. The following still needs to be done: 1) a survey of the DNA variants and their frequency distribution; 2) a cataloging of the coding sequences and the many decisions on homology that need to be made; 3) the identification of the full transcriptome of tuna; 4) the assignment of scaffolds to chromosomes; 5) the identification of promoter, enhancer, and other elements relevant to gene expression; 6) and the identification of the genome sections association with sex determination. While those are a wish-list for the future, the draft assembly presented here is a step forwards towards that goal.

Acknowledgements—We thank CSIRO for support and access to the High Performance Computing Cluster and to CSIRO and the University of Queensland for access to their library services. We thank Ross Tellam and Brian Dalrymple who read the manuscript and made comments which improved the text. Both noted the absence of Biology. We thank Blair Harrison and Janette Edson for technical assistance. Fish samples were obtained or coord-

inated by PMG. DNA was extracted by PMG and RJB. Sequencing libraries were prepared by RJB. Quality control analysis of fastq files for sequence analysis was performed by SM and RJB. Assembly of genomes and anal-

ysis of data was performed by WB and SM. Manuscript drafted by WB with comments from the other authors. The manuscript passed CSIRO internal review 6 July 2016.

REFERENCES

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. 1990. Basic local alignment search tool. *Journal of Molecular Biology*, **215**(3), 403–410.
- Appleyard, S. A., Grewe, P. M., Innes, B. H., & Ward, R. D. 2001. Population structure of yellowfin tuna (*Thunnus albacares*) in the western Pacific Ocean, inferred from microsatellite loci. *Marine Biology*, **139**(2), 383–393.
- Barendse, William, McWilliam, Sean, Bunch, Rowan J., & Harrison, Blair E. 2015. Adaptive divergence in the bovine genome. *bioRxiv*, DOI, 10.1101/022764.
- Bentley, D. R., Balasubramanian, S., Swerdlow, H. P., Smith, G. P., Milton, J., Brown, C. G., & et al. 2008. Accurate whole human genome sequencing using reversible terminator chemistry. *Nature*, **456**(7218), 53–59.
- Campbell, D., Brown, D., & Battaglione, T. 2000. Individual transferable catch quotas: Australian experience in the southern bluefin tuna fishery. *Marine Policy*, **24**(2), 109–117.
- Chen, C., Li, Y. L., Yu, H., Peng, S. M., Sun, S. G., Wang, L., Meng, X. J., Huang, Y., & Kong, X. D. 2016. The complete mitochondrial genome of the juvenile Pacific bluefin tuna *Thunnus orientalis* (Perciformes, Scombridae). *Mitochondrial DNA*, **27**(1), 71–72.
- Collette, B. B., Reeb, C., & Block, B. A. 2001. Systematics of the tunas and mackerels (Scombridae). *Fish Physiology*, **19**, 1–33.
- Doyle, J.J., & Doyle, J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin*, **19**, 11–15.
- Elliott, N. G., & Ward, R. D. 1992. Enzyme variation in orange roughly, *Hoplostethus atlanticus* (Teleostei, Trachichthyida), from Southern Australian and New Zealand waters. *Australian Journal of Marine and Freshwater Research*, **43**(6), 1561–1571.
- Gregory, T. R., Nicol, J. A., Tamm, H., Kullman, B., Kullman, K., Leitch, I. J., Murray, B. G., Kapraun, D. F., Greilhuber, J., & Bennett, M. D. 2007. Eukaryotic genome size databases. *Nucleic Acids Research*, **35**, D332–D338.
- Grewe, P. M., Feutry, P., Hill, P. L., Gunasekera, R. M., Schaefer, K. M., Itano, D. G., Fuller, D. W., Foster, S. D., & Davies, C. R. 2015. Evidence of discrete yellowfin tuna (*Thunnus albacares*) populations demands rethink of management for this globally important resource. *Scientific Reports*, **5**.
- Group, Pacific Bluefin Tuna Working. 2014. Stock assessment of Pacific Bluefin Tuna. [url](http://isc.fra.go.jp/pdf/ISC14/Annex%2016-%20PBFWG%20Report%2010Sep14.pdf), <http://isc.fra.go.jp/pdf/ISC14/Annex%2016-%20PBFWG%20Report%2010Sep14.pdf>.
- Hardie, D. C., & Hebert, P. D. N. 2004. Genome-size evolution in fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, **61**(9), 1636–1646.
- Itano, D. 2005. A handbook for the identification of Yellowfin and Bigeye Tunas in Fresh Condition. *Pelagic Fisheries Research Program, Honolulu, Hawaii, USA.. Ver*, **2**, 1–27.
- Joshi, K. K., Abdussamad, E. M., Koya, K. P. S., Rohit, P., Ghosh, S., Sreenath, K. R., Beni, M., Bineesh, K. K., & V., Akhilesh K. 2012. Taxonomy and key for the identification of tuna species exploited from the Indian EEZ. *Indian Journal of Fisheries*, **59**(3), 53–60.
- Li, R. Q., Zhu, H. M., Ruan, J., Qian, W. B., Fang, X. D., Shi, Z. B., Li, Y. R., Li, S. T., Shan, G., Kristiansen, K., Li, S. G., Yang, H. M., & Wang, J. 2010. De novo assembly of human genomes with massively parallel short read sequencing. *Genome Research*, **20**(2), 265–272.
- Li, Y. L., Chen, C., Yu, H., Peng, S. M., Sun, S. G., Wang, L., Meng, X. J., Huang, Y., & Kong, X. D. 2016. The complete mitochondrial genome of the juvenile Atlantic bluefin tuna *Thunnus thynnus* (Perciformes, Scombridae). *Mitochondrial DNA*, **27**(1), 96–97.
- Lowenstein, J. H., Amato, G., & Kolokotronis, S. O. 2009. The Real maccoyii: Identifying Tuna Sushi with DNA Barcodes - Contrasting Characteristic Attributes and Genetic Distances. *Plos One*, **4**(11), e7866.
- Luo, R. B., Liu, B. H., Xie, Y. L., Li, Z. Y., Huang, W. H., Yuan, J. Y., He, G. Z., Chen, Y. X., Pan, Q., Liu, Y. J., Tang, J. B., Wu, G. X., Zhang, H., Shi, Y. J., Liu, Y., Yu, C., Wang, B., Lu, Y., Han, C. L., Cheung, D. W., Yiu, S. M., Peng, S. L., Zhu, X. Q., Liu, G. M., Liao, X. K., Li, Y. R., Yang, H. M., Wang, J., & Lam, T. W. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience*, **1**, 18.
- MacKenzie, B. R., Mosegaard, H., & Rosenberg, A. A. 2009. Impending collapse of bluefin tuna in the northeast Atlantic and Mediterranean. *Conservation Letters*, **2**(1), 25–34.
- Marcais, G., & Kingsford, C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics*, **27**(6), 764–770.
- Martinez, P., Gonzalez, E. G., Castilho, R., & Zardoya, R. 2006. Genetic diversity and historical demography of Atlantic bigeye tuna (*Thunnus obesus*). *Molecular Phylogenetics and Evolution*, **39**(2), 404–416.
- Montes, I., Iriondo, M., Manzano, C., Arrizabalaga, H., Jimenez, E., Pardo, M. A., Goni, N., Davies, C. A., & Estonba, A. 2012. Worldwide genetic structure of albacore *Thunnus alalunga* revealed by microsatellite DNA markers. *Marine Ecology Progress Series*, **471**, 183–U202.
- Nakamura, Y., Mori, K., Saitoh, K., Oshima, K., Mekuchi, M., Sugaya, T., Shigenobu, Y., Ojima, N., Muta, S., Fujiwara, A., Yasuike, M., Oohara, I., Hirakawa, H., Chowdhury, V. S., Kobayashi, T., Nakajima, K., Sano, M., Wada, T., Tashiro, K., Ikeo, K., Hattori, M., Kuhara, S., Gojobori, T., & Inouye, K. 2013. Evolutionary changes of multiple visual pigment genes in the complete genome of Pacific bluefin tuna. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(27), 11061–11066.
- Nikolic, N., Duthoy, S., Destombes, A., Bodin, N., West, W., Puech, A., & Bourjea, J. 2015. Discovery of Genome-Wide Microsatellite Markers in Scombridae: A Pilot Study on Albacore Tuna. *Plos One*, **10**(11), e0141830.
- Pecoraro, C., Babbucci, M., Villamor, A., Franch, R., Papetti, C., Leroy, B., Ortega-Garcia, S., Muir, J., Rooker, J., Arocha, F., Murua, H., Zudaire, I., Chassot, E., Bodin, N., Tinti, F., Bargelloni, L., & Cariani, A. 2016. Methodological assessment of 2b-RAD genotyping technique for population structure inferences in yellowfin tuna (*Thunnus albacares*). *Marine Genomics*, **25**, 43–48.
- Riccioni, G., Landi, M., Ferrara, G., Milano, I., Cariani, A., Zane, L., Sella, M., Barbujani, G., & Tinti, F. 2010. Spatio-temporal population structuring and genetic diversity retention in depleted Atlantic Bluefin tuna of the Mediterranean Sea. *Proceedings of the National Academy of Sciences of the United States of America*, **107**(5), 2102–2107.
- Rooker, J. R., Bremer, J. R. A., Block, B. A., Dewar, H., De Metrio, G., Corriero, A., Kraus, R. T., Prince, E. D., Rodriguez-Marin, E., & Secor, D. H. 2007. Life history and stock structure of Atlantic bluefin tuna (*Thunnus thynnus*). *Reviews in Fisheries Science*, **15**(4), 265–310.
- Santaclara, F. J., Velasco, A., Perez-Martin, R. I., Quinteiro, J., Rey-Mendez, M., Pardo, M. A., Jimenez, E., & Sotelo, C. G. 2015. Development of a multiplex PCR-ELISA method for the genetic authentication of *Thunnus* species and *Katsuwonus pelamis* in food products. *Food Chemistry*, **180**, 9–16.
- Takagi, M., Okamura, T., Chow, S., & Taniguchi, N. 1999. PCR primers for microsatellite loci in tuna species of the genus *Thunnus* and its application for population genetic study. *Fisheries Science*, **65**(4), 571–576.
- Takahshima, Y., Iguchi, J., Namikoshi, A., Yamashita, Y., & Yamashita, M. 2014. DNA Analysis for Identification of Species and Geographical Origins of Fishery Products. *Bunseki Kagaku*, **63**(10), 797–807.

- Terol, J., Mascarell, R., Fernandez-Pedrosa, V., & Perez-Alonso, M. 2002. Statistical validation of the identification of tuna species: Bootstrap analysis of mitochondrial DNA sequences. *Journal of Agricultural and Food Chemistry*, **50**(5), 963–969.
- Ward, R. D. 2000. Genetics in fisheries management. *Hydrobiologia*, **420**, 191–201.
- Ward, R. D., Elliott, N. G., & Grewe, P. M. 1995. Allozyme and mitochondrial DNA separation of Pacific northern bluefin tuna, *Thunnus thynnus orientalis* (Temminck and Schlegel), from southern bluefin tuna, *Thunnus maccoyii* (Castelnau). *Marine and Freshwater Research*, **46**(6), 921–930.
- Yasuike, M., Fujiwara, A., Nakamura, Y., Iwasaki, Y., Nishiki, I., Sugaya, T., Shimizu, A., Sano, M., Kobayashi, T., & Ototake, M. 2016. A functional genomics tool for the Pacific bluefin tuna: Development of a 44K oligonucleotide microarray from whole-genome sequencing data for global transcriptome analysis. *Gene*, **576**(2), 603–609.




FIG. 1.— The k-mer distribution of 17-mers of southern bluefin tuna yielding a genome size estimate of 795 Mb




FIG. 2.— The increase in contig sequence diversity with increased contig length. Diversity is expressed as the Shannon Information Index. Minimum diversity is shown with squares, maximum diversity with diamonds, the mean with circles, and the standard deviation with triangles




FIG. 3.— The size distribution of contigs in base pairs for PDB3 $k=25$ in black $k=51$ in brown and PDB5 $k=25$ in red $k=51$ in blue




FIG. 4.— The distribution of number of matches per scaffold for PDB3 (open circles) and for PDB5 (closed circles) for *Danio rerio* RNA

```

Query 27940  GTCACCGGCTGATGAACACTGAGACGCTGTTTCCATCTCAAATCACAGCCTCTTTTCTTG 27999
             |||
Sbjct 159872  GTCACCGGCTGATGAACACTGAGACGCTGTTTCCATCTCAAATCACAGCCTCTTTTCTTG 159931

Query 28000  TGTGGTTCAGTGGAGAATAGGTTGTCATTTGAAGCCCCCCCACCCCTCTCTCCCTCTCCC 28059
             |||
Sbjct 159932  TGTGGTTCAGTGGAGAATAGGTTGTCATTTGAAGCCCCCCNNNNNNNNNNNNNNNNNNNN 159991

Query 28060  CTCCCTGTCGACTCTCTGTTGCCACTTTATGGCTCTGTTAAGTATTCCTTATAGCACCTC 28119
             |
Sbjct 159992  NNNNNNNNNNNNNNNNNNNNNNA-----TGGCTCTGTTAAGTATTCCTTATAGCACCTC 160046

Query 28120  TATGAAACTGTGTGGACCTCATGGCTCACCTGACAATCTGTACCATACAGTGTAAACACAC 28179
             |||
Sbjct 160047  TATGAAACTGTGTGGACCTCATGGCTCACCTGACAATCTGTACCATACAGTGTAAACACAC 160106

Query 28180  CAGAAACCTGCACCTTAGCTAATCAAGACATTTTAAATTCGAGCTCCTTCTTGCTACTT 28239
             |||
Sbjct 160107  CAGAAACCTGCACCTTAGCTAATCAAGACATTTTAAATTCGAGCTCCTTCTTGCTACTT 160166

```

FIG. 5.— BLAST alignment between part of a PDB3 and PDB5 scaffold where sequence complementarity increases coverage

TABLE 1
 GENOME SEQUENCE USED FOR TUNA GENOME ASSEMBLY.

Species	N	GC percent	PE reads	MP reads	PE Coverage
albacore (ALB)	3	40.26	533,707,152	-	66X
bigeye (BET)	3	39.84	220,316,910	-	28X
northern bluefin (NBT)	3	39.99	352,965,874	-	44X
skipjack (SJK)	4	40.82	734,059,014	-	91X
southern bluefin (SBT) ^a	13	40.72	1,694,446,949	-	213X
SBT PDB3&PDB5	2	39.94	479,524,357	1,266,757,744	60X
yellowfin (YFT)	2	39.81	240,409,742	-	30X

NOTE. — The number of reads reported is for merged, trimmed, and paired reads, representing the reads used in the assembly, which are a subset of the reads collected for each species

^a Total SBT is 2,173,971,306 reads, which is the sum of these reads and the SBT Pedrabranca reads

TABLE 2
TUNA ASSEMBLY STATISTICS.

Species	Contig Number	Coverage (Gb)	N50 (bp)	Max (bp)	Scaffold ^a Number	Coverage (Gb)	N50 (bp)	Max (bp)
albacore	1,826,784	545	398	7,187	176,177	343	1,406	43,393
bigeye	1,533,456	606	640	15,453	147,654	454	3,207	65,966
northern bluefin	1,741,562	575	479	11,837	174,297	391	1,837	45,910
skipjack	1,812,717	422	271	5,119	134,886	194	608	45,518
southern bluefin ^b	1,756,506	590	489	8,422	174,892	407	1,973	44,602
yellowfin	1,584,634	607	616	15,538	157,596	446	2,760	63,041

NOTE. — These assemblies were constructed without MP sequences and with k-mer=25

^a Number of scaffolds consisting of more than one contig

^b SBT PDB3 and PDB5 PE sequences only, k=25

TABLE 3
SOUTHERN BLUEFIN TUNA ALTERNATIVE ASSEMBLY STATISTICS.

Assembly	Contig Number	Coverage (Gb)	N50 (bp)	Max (bp)	Scaffold ^a Number	Coverage (Gb)	N50 (bp)	Max (bp)
PDB3-pe ^b	1,523,729	629	720	16,076	162,973	459	2,708	59,033
PDB3-mp ^c	1,523,729	629	720	16,076	12,828	890	174,777	1,412,496
PDB3x5-mp ^d	1,523,729	629	720	16,076	12,827	945	222,019	1,707,042
PDB3-mp51 ^e	2,630,814	938	709	37,065	14,814	930	147,567	1,412,291
PDB5-pe ^f	1,625,821	627	626	13,874	169,576	455	2,538	50,265
PDB5-mp ^g	1,625,821	627	626	13,874	11,740	963	245,122	2,315,601
PDB5x3-mp ^h	1,625,821	627	626	13,874	13,215	879	181,542	1,323,743
PDB5-mp51 ⁱ	2,472,157	920	768	30,018	11,139	998	243,461	2,056,612
PDB3&PDB5-pe ^j	1,756,506	590	489	8,422	174,892	407	1,973	44,602
PDB3&PDB5-mp ^k	1,756,506	590	489	8,422	244,152	298	922	9,801
PDB3&PDB5-mp51 ^l	4,629,720	1,089	280	20,743	223,853	340	754	22,277
All-pe ^m	1,813,209	439	285	4,957	144,536	228	801	27,046
All-mp ⁿ	1,813,209	439	285	4,957	193,004	178	522	6,402
All-mp51 ^o	10,794,475	1,516	112	14,818	18,716	21	86	14,767

NOTE. — There are significant differences in the quality of these assemblies by changing the k-mer size and the data set

^a Number of scaffolds consisting of more than one contig

^b PDB3 PE sequences only, k=25

^c PDB3 including MP sequences, k=25

^d PDB3 using PDB5 MP sequences, k=25

^e PDB3 including MP sequences, k=51

^f PDB5 PE sequences only, k=25

^g PDB5 including MP sequences, k=25

^h PDB5 using PDB3 MP sequences, k=25

ⁱ PDB5 including MP sequences, k=51

^j PDB3 and PDB5 PE sequences only

^k PDB3 and PDB5 including MP sequences

^l PDB3 and PDB5 including MP sequences, k=51

^m All available SBT PE sequences, k=25

ⁿ All available SBT PE and MP sequences, k=25

^o All available SBT PE and MP sequences, k=51

TABLE 4
 COVERAGE OF DANIO RERIO RNA ON SOUTHERN BLUEFIN TUNA ASSEMBLIES.

Assembly	NM ^a 1-e05	NM 1-e10	NM 1-e15	NM 1-e20	NM 1-e25	NM 1-e30	NM 1-e35	NM 1-e40	NM 1-e45	NM 1-e50
PDB3	44801	41539	38158	34657	30903	27127	23464	20027	16742	13916
PDB5	44665	41289	37845	34291	30742	26954	23069	19594	16249	13426
PDB3 and PDB5 ^b	-	-	10464	-	-	-	-	3598	-	-
PDB3 or PDB5 ^c	-	-	13039	-	-	-	-	8222	-	-
PDB3 and PDB5 ^d	-	-	30868	-	-	-	-	11160	-	-
PDB3 or PDB5 ^e	-	-	45135	-	-	-	-	28461	-	-

^a number of RNA matching to assembly at BLAST probability 1-e05

^b The D. rerio mRNA matching to both assemblies at that BLAST probability

^c The total D. rerio mRNA matching to either assembly at that BLAST probability

^d The D. rerio RNA matching to both assemblies at that BLAST probability

^e The total D. rerio RNA matching to either assembly at that BLAST probability