

Risk minimization by median-of-means tournaments ^{*}

Gábor Lugosi^{†‡} Shahar Mendelson [§]

August 3, 2016

Abstract

We consider the classical statistical learning/regression problem, when the value of a real random variable Y is to be predicted based on the observation of another random variable X . Given a class of functions \mathcal{F} and a sample of independent copies of (X, Y) , one needs to choose a function \hat{f} from \mathcal{F} such that $\hat{f}(X)$ approximates Y as well as possible, in the mean-squared sense. We introduce a new procedure, the so-called median-of-means tournament, that achieves the optimal tradeoff between accuracy and confidence under minimal assumptions, and in particular outperforms classical methods based on empirical risk minimization.

1 Introduction

Estimation and prediction problems are of central importance in statistics and learning theory. In the standard regression setup, (X, Y) is a pair of random variables: X takes its values in some (measurable) set \mathcal{X} and is distributed according to an unknown probability measure μ , while Y is real valued that is also unknown. Given a class \mathcal{F} of real-valued functions defined on \mathcal{X} , one wishes to find $f \in \mathcal{F}$ for which $f(X)$ is a good prediction of Y . Although one may consider various notions of ‘a good prediction’, we restrict our attention to the—perhaps most commonly used—*squared error*: the learner is penalized by $(f(X) - Y)^2$ for predicting $f(X)$ instead of Y . Thus, one would like to find a function $f \in \mathcal{F}$ for which the expected loss $\mathbb{E}(f(X) - Y)^2$, known as the *risk*, is as small as possible. Naturally, the best performance one may hope for is of the risk minimizer in the class, that is, that of

$$f^* = \operatorname{argmin}_{f \in \mathcal{F}} \mathbb{E}(f(X) - Y)^2 .$$

We assume in what follows that the minimum is attained and $f^* \in \mathcal{F}$ exists and is unique, as is the case when $\mathcal{F} \subset L_2(\mu)$ is a closed, convex set.

One may formulate two natural goals in estimation and prediction problems. One of them is to find a function $f \in \mathcal{F}$ whose $L_2(\mu)$ distance to f^*

$$\left(\mathbb{E} (f(X) - f^*(X))^2 \right)^{1/2} \tag{1.1}$$

^{*}Gábor Lugosi was supported by the Spanish Ministry of Economy and Competitiveness, Grant MTM2015-67304-P and FEDER, EU. Shahar Mendelson was supported in part by the Israel Science Foundation.

[†]Department of Economics and Business, Pompeu Fabra University, Barcelona, Spain, gabor.lugosi@upf.edu

[‡]ICREA, Pg. Llus Companys 23, 08010 Barcelona, Spain

[§]Department of Mathematics, Technion, I.I.T, and Mathematical Sciences Institute, The Australian National University, shahar@tx.technion.ac.il

is as small as possible. The other is to ensure that the *excess risk* of the selected function

$$R(f) = \mathbb{E}(f(X) - Y)^2 - \mathbb{E}(f^*(X) - Y)^2, \quad (1.2)$$

is a small.

The crucial difference between this type of problems and standard questions in approximation theory is that the available information is limited to a random sample. One observes $\mathcal{D}_N = ((X_1, Y_1), \dots, (X_N, Y_N))$, that is, N independent pairs, where each (X_i, Y_i) has the same distribution as (X, Y) and \mathcal{D}_N is independent of (X, Y) . The fact that the distribution of the pair (X, Y) is not known makes it impossible to invoke approximation-theoretical methods and identify directly the true minimizer of the risk.

Given a sample size N , a *learning procedure* is a map $\Phi : (\mathcal{X} \times \mathbb{R})^N \rightarrow \mathcal{F}$ that assigns to each sample $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$ a (random) function in \mathcal{F} , which we denote by \hat{f}_N . The *mean squared error* (also called the *estimation error*) of Φ is the squared $L_2(\mu)$ distance between the true minimizer f^* and the function selected by Φ given the data \mathcal{D}_N , that is, the conditional expectation

$$\|\hat{f}_N - f^*\|_{L_2}^2 = \mathbb{E}((\hat{f}_N(X) - f^*(X))^2 | \mathcal{D}_N) \equiv \mathcal{E}_e^2,$$

where throughout the article, for $q \geq 1$, we use the notation

$$\|f - g\|_{L_q} = (\mathbb{E}|f(X) - g(X)|^q)^{1/q} \quad \text{and also} \quad \|f - Y\|_{L_q} = (\mathbb{E}|f(X) - Y|^q)^{1/q}.$$

The *excess risk*, also known as the *prediction error*, compares the ‘predictive capabilities’ of \hat{f}_N to that of the best in the class, and is defined by the conditional expectation

$$R(\hat{f}_N) = \mathbb{E}((\hat{f}_N(X) - Y)^2 | \mathcal{D}_N) - \mathbb{E}(f^*(X) - Y)^2 \equiv \mathcal{E}_p^2.$$

Both the mean squared error and the excess risk are functions of the given data \mathcal{D}_N , and as such are random quantities. It is worth noting here that in the special situation when $f^*(X) = \mathbb{E}(Y|X)$, we have $R(\hat{f}_N) = \|\hat{f}_N - f^*\|_{L_2}^2$. This is the case, for example, when $Y = f(X) + W$ for some $f \in \mathcal{F}$ and a zero-mean random variable W that is independent of X . However, in general, a small mean squared error does not automatically imply a small excess risk, or vice-versa.

In what follows we refer to both \mathcal{E}_e and \mathcal{E}_p as the *accuracy*¹ of learning procedure Φ . The *confidence* of Φ for an error rate of \mathcal{E} is the probability (with respect to the product measure on $(\mathcal{X} \times \mathbb{R})^N$ endowed by the pair (X, Y)) with which Φ performs with accuracy smaller than \mathcal{E} .

Note that, up to this point, Y was an arbitrary square-integrable real-valued random variable, and obviously one would like to be able to treat as wide a variety of targets as possible. Clearly, the accuracy and confidence one may establish may depend on some features of the target—for example, some a-priori estimate on its L_q norm, or on its ‘distance’ to \mathcal{F} , etc. We consider a broad set of *admissible targets* \mathcal{Y} , and the accuracy and confidence of Φ relates to its performance for *any* admissible target $Y \in \mathcal{Y}$. Thus, a *learning problem* is the triplet (\mathcal{F}, Y, X) , when X and Y are not known, though the learner does know that $Y \in \mathcal{Y}$.

It is clear that there is a tradeoff between the accuracy and confidence in a given learning problem: the smaller the error is, the harder it is to attain it. The question of this accuracy/confidence tradeoff is of utmost importance in statistical learning theory, and has been

¹Sometimes the accuracy is defined by \mathcal{E}_e^2 and \mathcal{E}_p^2 .

investigated extensively in numerous manuscripts since the early days of the area in the late 1960’s (see, for example, the books [42, 13, 41, 2, 39, 28, 22, 38, 10] for a sample of the work devoted to this question). To find the right accuracy/confidence tradeoff one must first identify a lower bound on the tradeoff in terms of the sample size, the structure of \mathcal{F} and possibly some additional information on X and Y , and then come up with a learning procedure that attains the tradeoff.

Roughly put, one should explore the tradeoff for the set of “achievable accuracies” of each learning problem. An accuracy \mathcal{E} is achievable if there is a learning procedure in \mathcal{F} that achieves the accuracy \mathcal{E} for the problem (\mathcal{F}, Y, X) with constant probability—say at least $3/4$ —, and because Y is not known, this has to hold for any $Y \in \mathcal{Y}$. We define the *accuracy edge* as the smallest achievable accuracy of a problem.

The primary question is to find the correct accuracy/confidence tradeoff for any (reasonable) learning problem and identify a learning procedure that attains that tradeoff all the way down to the accuracy edge. It should be noted that up to this point in time, and other than in a few isolated examples, no learning procedure even came close to the optimal tradeoff at *any* nontrivial accuracy level.

In this article we solve this problem by presenting an *optimal learning procedure*: it yields the best possible accuracy/confidence tradeoff (almost) up to the achievable edge, and under minimal assumptions on the learning problem.

The minor reservation “almost” is due to fact that more often than not, the identity of the accuracy edge of a learning problem is not known. As it is explained in what follows, while one may provide lower estimates on the accuracy edge, there is a very real possibility that such estimates are too optimistic and the real accuracy edge is larger. Regardless, the procedure we introduce “gets as close” to the accuracy edge as any other known procedure—or better, and with a dramatically better confidence. It also exhibits the optimal accuracy/confidence tradeoff for larger errors, something that no other procedure is known to do.

It should be clarified at this point that by “optimal accuracy” we mean an accuracy that is optimal up to a constant factor and optimality in the confidence means that the probability with which the claimed accuracy does not hold is optimal up to a constant factor in the exponent.

Before we present a more accurate formulation of our main results and describe the optimal procedure, let us explain what our procedure is not: it is not *empirical risk minimization* (ERM), nor any of its “family members”.

1.1 ERM—the wrong choice

Perhaps the most natural way of choosing \hat{f}_N is by empirical risk minimization, that is, by *least squares regression*,

$$\hat{f}_N = \operatorname{argmin}_{f \in \mathcal{F}} \sum_{i=1}^N (f(X_i) - Y_i)^2 .$$

Again, we assume that the minimum is attained, while if there are several minimizers, \hat{f}_N may be chosen among them in an arbitrary way.

The performance of least squares regression has been thoroughly studied in many different scenarios. A sample of the rich literature includes Györfi, Kohler, Krzyzak, Walk [17], van de Geer [39], Bartlett, Bousquet, and Mendelson [6], Koltchinskii [23], Massart [28].

The simple idea behind empirical risk minimization is that, for each $f \in \mathcal{F}$, the empirical risk $(1/N) \sum_{i=1}^N (f(X_i) - Y_i)^2$ is a good estimate of the risk $\mathbb{E}(f(X) - Y)^2$ and the minimizer of the empirical risk should nearly match that of the “true” risk.

It turns out that the performance of ERM changes dramatically according to the tail behaviour of the functions involved in the given learning problem. One may show (see, e.g., [24]) that if \mathcal{F} is convex and the functions in \mathcal{F} (more precisely, the random variables $\{f(X) : f \in \mathcal{F}\}$) and the target Y have well-behaved tails (and by “well-behaved” we mean sub-Gaussian), ERM performed in \mathcal{F} yields good results: for an accuracy that is not far from the accuracy edge, it attains the optimal confidence, though it does not maintain the optimal accuracy/confidence tradeoff for larger errors. Unfortunately, the situation deteriorates considerably when either members of \mathcal{F} or one of the admissible targets is heavy-tailed in some sense. In such cases, the performance of ERM is significantly weaker than the known theoretical limitations of the accuracy/confidence tradeoff. Moreover, replacing ERM with a different procedure is of little use: other than in few and rather special learning problems, there have been no known alternatives to ERM whose performance comes close to the known theoretical limitations of the accuracy/confidence tradeoff, and certainly not when the problem is heavy-tailed.

The reason for ERM’s diminished capacity is that it is sensitive to even a small number of atypical points in the sample $(X_i, Y_i)_{i=1}^N$: since ERM selects a minimizer of the empirical mean of the squared loss, atypical values may distort the selection and send ERM to the wrong part of \mathcal{F} . This sensitivity is clearly reflected in the confidence with which ERM operates in heavy-tailed situations: roughly put, one can guarantee that ERM performs with the right accuracy only on samples that are not contaminated by a significant number of atypical values. However, in heavy-tailed situations, the latter does not occur frequently, and having atypical values is simply a fact of life one has to deal with.

In contrast, the procedure we suggest as an alternative to ERM leads to the optimal accuracy/confidence tradeoff even in heavy-tailed situations. Unlike ERM, it is not sensitive even to a large number of atypical sample points.

Before we dive into a more technical description of our results, let us present the following classical example of linear regression in \mathbb{R}^n , exhibiting the limitations of empirical risk minimization in heavy-tailed problems, and comparing its performance to that of the procedure we introduce.

Let $\mathcal{F} = \{\langle t, \cdot \rangle : t \in \mathbb{R}^n\}$ be the class of linear functionals on \mathbb{R}^n . Let X be an isotropic random vector in \mathbb{R}^n (i.e., $\mathbb{E} \langle t, X \rangle^2 = 1$ for every t in the Euclidean unit sphere) and assume that X exhibits some (very weak) norm equivalence in the following sense: there are $q > 2$ and $L > 1$ for which, for every $t \in \mathbb{R}^n$, $\|\langle X, t \rangle\|_{L_q} \leq L \|\langle X, t \rangle\|_{L_2}$.

Assume that one is given N noisy measurements of $\langle t_0, \cdot \rangle$ for a fixed but unknown $t_0 \in \mathbb{R}^n$. Specifically, assume that $Y = \langle t_0, X \rangle + W$ for some symmetric random variable W that is independent of X and has variance σ^2 . One observes the “noisy” data $(X_i, Y_i)_{i=1}^N$ and the aim is to approximate t_0 with a small error (accuracy) and with high probability (confidence) using this random data only.

One may show that a nontrivial estimate is possible only when $N \geq cn$ for a suitable absolute constant c , and we consider only such values of N . Also, there are known estimates on the theoretical limitations of this problem: a lower bound on the accuracy edge is of the order of $\sigma \sqrt{n/N}$, and for an accuracy level that is proportional to the accuracy edge, say, $c_0 \sigma \sqrt{n/N}$ for a suitable absolute constant c_0 , the conjectured confidence is $1 - 2 \exp(-c_1 n)$.

If there is no information on higher than q -th moments for linear functionals, and no

information beyond the second moment for W is available, this clearly is a (potentially) heavy-tailed scenario. It turns out (the claims made here follow from results of [24], see the next section for the general statements) that the best that one can guarantee using ERM is a choice of $\hat{t} \in \mathbb{R}^n$, for which the Euclidean norm $\|\hat{t} - t_0\|_2 = \|\langle \hat{t}, X \rangle - \langle t_0, X \rangle\|_{L_2} \leq r$ with probability at least $1 - \delta - 2 \exp(-cN)$; the error r is defined as the smallest number for which

$$(*) = \left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N W_i X_i \right\|_2 \leq c\sqrt{N}r \quad \text{with probability at least } 1 - \delta .$$

Since X is isotropic, one has $\mathbb{E}\|X\|_2^2 = n$. Therefore, the mean of $(*)$ is bounded as

$$\mathbb{E} \left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N W_i X_i \right\|_2 \leq \left(\mathbb{E} \left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N W_i X_i \right\|_2^2 \right)^{1/2} = \sigma(\mathbb{E}\|X\|_2^2)^{1/2} = \sigma\sqrt{n} .$$

Moreover, because of the minimal assumptions on W and X , the best estimate one can hope for on $(*)$ and that holds with probability $1 - \delta$ follows from Chebyshev's inequality. In particular, it leads to the following, rather unsatisfactory, estimate on the performance of ERM:

$$\|\hat{t} - t_0\|_2 \leq \frac{c_0(q, L)}{\delta} \sigma \sqrt{\frac{n}{N}} \quad \text{with probability } 1 - \delta - 2 \exp(-c_1 N) .$$

Also, if one wishes for an error that is proportional to the (conjectured) accuracy edge, that is, of the order of $\sigma\sqrt{n/N}$, the best that one can hope for is a constant confidence—a very different estimate from the conjectured confidence of $1 - 2 \exp(-c_1 n)$.

Although what we have described is an upper bound, one may show (see [24]) that this estimate captures the performance of empirical risk minimization, and in particular exhibits ERM's inability to deal with atypical sample points. The reality is that ERM performs with an accuracy of the order of $\sigma\sqrt{n/N}$ only on the relatively few samples that contain almost no misleading data.

The main result of this article, when applied to this example, shows that under the same assumptions, the procedure we suggest selects \hat{t} for which

$$\|\hat{t} - t_0\|_2 \leq C\sigma\sqrt{\frac{n}{N}} \quad \text{with probability } 1 - 2 \exp(-cn) \tag{1.3}$$

for some numerical constants $c, C > 0$; that is, it performs with optimal confidence at a level that is proportional to the accuracy edge. In fact, our procedure gives the optimal confidence for any accuracy $r \geq c'\sigma\sqrt{n/N}$.

Note that for the special case of linear regression described above, Hsu and Sabato [19] achieve slightly (by a factor logarithmic in n) weaker bounds than (1.3) under slightly stronger $((4 + \epsilon)$ -th moment) assumptions. We also refer to Minsker [34] for related bounds for sparse regression under possibly heavy-tailed variables. Of course, these results hold in a rather special example, while our main result yields optimal estimates for almost any convex class \mathcal{F} and target Y , and not just for linear regression in \mathbb{R}^n .

In the next section we present the required definitions, outline the current state of the art, and formulate our main results. In Section 3, we describe the new procedure in detail. In Section 4 we illustrate the power of the main results on some canonical examples, before turning to the proofs of our results.

Let us point out the well-understood fact that the behaviour of the accuracy and confidence in learning problems in which \mathcal{F} is not convex is trivial in some sense, and totally different from the convex case, which is why focus on the latter. As it happens, the dominating factor in non-convex problems is the ‘location’ of the targets Y relative to \mathcal{F} rather than the structure of \mathcal{F} , and an ‘unfavourable location’ of a target Y completely distorts the accuracy and confidence of the learning problem. However, even if \mathcal{F} is not convex, all the targets of the form $Y = f_0(X) + W$ for $f_0 \in \mathcal{F}$ and W that is symmetric and independent of X happen to be in a ‘favourable’ location and thus our results apply to such problems as well.

2 The accuracy edge and the accuracy/confidence tradeoff

We begin by describing the known theoretical limitations on the accuracy edge and on the accuracy/confidence tradeoff for a given learning problem. To this end, let us introduce some notation, following the path of [29, 30].

Let $D = \{f : \|f\|_{L_2} \leq 1\}$ be the unit ball in $L_2(\mu)$ and set $S = \{f : \|f\|_{L_2} = 1\}$ to be the unit sphere. For $h \in L_2(\mu)$ and $r > 0$, put $rD_h = \{f : \|f - h\|_{L_2} \leq r\}$. Let

$$\text{star}(\mathcal{F}, h) = \{\lambda f + (1 - \lambda)h : 0 \leq \lambda \leq 1, f \in \mathcal{F}\}.$$

Thus, $\text{star}(\mathcal{F}, h)$ is the star-shaped hull of \mathcal{F} around h , that is, the union of all segments for which one end-point is h and the other is in \mathcal{F} .

The star-shaped hull $\text{star}(\mathcal{F}, h)$ adds regularity to the class around the fixed centre h : on the one hand, it does not increase the size of the class by much, while on the other hand, it implies that every function of the form $f - h$ has a ‘scaled-down’ version when one moves towards 0. In particular, the level sets $\text{star}(\mathcal{F} - h, 0) \cap rS$ become ‘richer’ as r gets smaller: each one of them contains scaled-down copies of all ‘higher’ levels.

Consider the localization of \mathcal{F}

$$\mathcal{F}_{h,r} = \text{star}(\mathcal{F} - h, 0) \cap rD,$$

which is given by the shift that maps the designated point h to 0. Then the resulting class is made ‘more regular’ by taking its star-shaped hull around 0, and finally it is localized, by considering its intersection with rD , the $L_2(\mu)$ ball of radius r , centred in 0.

Observe that if \mathcal{F} is convex then for any $h \in \mathcal{F}$, $\text{star}(\mathcal{F}, h) = \mathcal{F}$. Also, in that case,

$$\mathcal{F}_{h,r} = \{f - h : f \in \mathcal{F}, \|f - h\|_{L_2} \leq r\},$$

and if, in addition, \mathcal{F} is centrally symmetric (that is, if $f \in \mathcal{F}$ then $-f \in \mathcal{F}$), then $\mathcal{F}_{h,r} \subset 2\mathcal{F} \cap rD$.

One way of deriving lower estimates on the accuracy edge and on the accuracy/confidence tradeoff is based on the packing numbers of the localizations $\mathcal{F}_{h,r}$.

Definition 2.1. *Given a set $H \subset L_2(\mu)$ and $\varepsilon > 0$, denote by $\mathcal{M}(H, \varepsilon D)$ the cardinality of a maximal ε -separated subset of H . That is, $\mathcal{M}(H, \varepsilon D)$ is the maximal cardinality of a subset $\{h_1, \dots, h_m\} \subset H$, for which $\|h_i - h_j\|_{L_2} \geq \varepsilon$ for every $i \neq j$.*

Note that if H' is a maximal ε -separated subset of H , then it is also an ε -cover of H in the sense that for every $h \in H$ there is some $h' \in H'$ that satisfies $\|h' - h\|_{L_2} \leq \varepsilon$.

Definition 2.2. For $\kappa, \eta > 0$ and $h \in \mathcal{F}$, set

$$\lambda_{\mathbb{Q}}(\kappa, \eta, h) = \inf\{r : \log \mathcal{M}(\mathcal{F}_{h,r}, \eta r D) \leq \kappa^2 N\} , \quad (2.1)$$

and let

$$\lambda_{\mathbb{Q}}(\kappa, \eta) = \sup_{h \in \mathcal{F}} \lambda_{\mathbb{Q}}(\kappa, \eta, h) .$$

For every fixed $h \in \mathcal{F}$, the parameter $\lambda_{\mathbb{Q}}$ pin-points the level r at which the localization $\mathcal{F}_{h,r}$ becomes “too rich” in the following sense: given the sample size N , $\mathcal{F}_{h,r}$ contains a subset of cardinality $\exp(\kappa^2 N)$ that is ηr -separated with respect to the $L_2(\mu)$ norm. Note that $\lambda_{\mathbb{Q}}$ is not affected by the fine structure of $\mathcal{F}_{h,r}$. Indeed, the set $\mathcal{F}_{h,r} \cap (\eta r/2)D$ cannot contain more than two points that are ηr -separated, and thus it does not contribute to the existence of a large ηr -separated set in $\mathcal{F}_{h,r}$.

One may show that $\lambda_{\mathbb{Q}}$ serves as a lower bound on the accuracy edge of a learning problem, when \mathcal{F} is convex and centrally symmetric and the admissible targets are noise-free: that is, $\mathcal{Y} = \{f(X) : f \in \mathcal{F}\}$.

Proposition 2.3. [30] *There exist absolute constants κ and η for which the following holds. Let $\mathcal{F} \subset L_2(\mu)$ be convex and centrally symmetric. For any learning procedure Φ there exists an $f_0 \in \mathcal{F}$ and target $Y = f_0(X)$ for which, with probability at least $1/4$,*

$$\|\Phi(\mathcal{D}_N) - f_0\|_{L_2}^2 \geq \lambda_{\mathbb{Q}}(\kappa, \eta) .$$

The following variant of $\lambda_{\mathbb{Q}}$ also serves as a lower bound on the accuracy edge, this time, because of ‘noisy’ targets.

Definition 2.4. For $\kappa > 0$, $0 < \eta < 1$ and $h \in \mathcal{F}$, set

$$\lambda_{\mathbb{M}}(\kappa, \eta, h) = \inf\{r : \log \mathcal{M}(\mathcal{F}_{h,r}, \eta r D) \leq \kappa^2 N r^2\} \quad (2.2)$$

and let

$$\lambda_{\mathbb{M}}(\kappa, \eta) = \sup_{h \in \mathcal{F}} \lambda_{\mathbb{M}}(\kappa, \eta, h) .$$

Proposition 2.5. (See, e.g., [30].) *There exist absolute constants κ and η for which the following holds. Let $\mathcal{F} \subset L_2(\mu)$ and set W to be a centred Gaussian variable with variance $\sigma > 0$ that is independent of X . Then, for any learning procedure Φ there exists $f_0 \in \mathcal{F}$ and a target $Y = f_0(X) + W$ for which, with probability at least $1/4$,*

$$\|\Phi(\mathcal{D}_N) - f_0\|_{L_2}^2 \geq \lambda_{\mathbb{M}}(\kappa/\sigma, \eta) .$$

In particular, if \mathcal{Y} contains all the targets of the form $f_0(X) + W$, for $f_0 \in \mathcal{F}$ and W that is a centred Gaussian variable with variance σ that is independent of X , then the accuracy edge is at least $\lambda_{\mathbb{M}}(\kappa/\sigma, \eta)$.

Combining these two facts, we have a lower bound on the accuracy edge:

$$\lambda^* \equiv \max\{\lambda_{\mathbb{Q}}(\kappa_1, \eta_1), \lambda_{\mathbb{M}}(\kappa_2/\sigma, \eta_2)\} ,$$

for some constants κ_i, η_i , $i = 1, 2$. However, there is no guarantee that this lower estimate is sharp. As we explain in what follows, it is very possible that the true accuracy edge is larger.

Let us turn to the theoretical limitations of the accuracy/confidence tradeoff, assuming that the set of admissible targets \mathcal{Y} is not too trivial. By that we mean that it at least contains all the targets of the form $Y = f_0(X)$ (the noise-free problems) and $Y = f_0(X) + W$, for $f_0 \in \mathcal{F}$ and W that is a centred Gaussian variable with variance σ , and is independent of X . We call this set of targets *minimal*, and of course, \mathcal{Y} could be much larger.

Applying the results in [24] and [30] one has the following:

Proposition 2.6. *There exists an absolute constant c for which the following holds. Let $\mathcal{F} \subset L_2(\mu)$ be a class that is star-shaped around one of its points (i.e., for some $f_0 \in \mathcal{F}$ and every $f \in \mathcal{F}$, $[f_0, f] \subset \mathcal{F}$). Consider \mathcal{Y} that contains the minimal set of targets. If Φ is a learning procedure that performs with accuracy r and confidence $1 - \delta$ for every such target, then*

$$\delta \geq \exp(-cN \min\{1, r^2/\sigma^2\}).$$

These facts set our first benchmark (which may be too optimistic, of course): the lower bound on the accuracy edge

$$\lambda^* = \max\{\lambda_{\mathbb{Q}}(\kappa_1, \eta_1), \lambda_{\mathbb{M}}(\kappa_2/\sigma, \eta_2)\}, \quad (2.3)$$

and the bound on the accuracy/confidence tradeoff for $r \geq c_0\lambda^*$,

$$1 - 2 \exp(-c_1N \min\{1, \sigma^{-2}r^2\}). \quad (2.4)$$

As we noted earlier, λ^* is an optimistic, and perhaps not very realistic, lower bound on the accuracy edge. A more reasonable conjecture relies on more “global” parameters that take into account the fine structure of \mathcal{F} at an arbitrarily small level, defined next.

From here on, let $(\varepsilon_i)_{i=1}^N$ be independent, symmetric $\{-1, 1\}$ -valued random variables that are independent of $(X_i, Y_i)_{i=1}^N$.

Definition 2.7. *For $\kappa > 0$ and $h \in \mathcal{F}$ let*

$$r_E(\kappa, h) = \inf \left\{ r : \mathbb{E} \sup_{u \in \mathcal{F}_{h,r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i u(X_i) \right| \leq \kappa \sqrt{Nr} \right\}, \quad (2.5)$$

and set $r_E(\kappa) = \sup_{h \in \mathcal{F}} r_E(\kappa, h)$.

The parameter $r_E(\kappa, h)$ measures the empirical oscillation around h . It does not depend on the identity of the target Y and is a purely intrinsic parameter of the class \mathcal{F} . However, it may be highly affected by functions in \mathcal{F} that are close to h , and as such it is more “global” than $\lambda_{\mathbb{Q}}$.

The other “global” parameter we require does depend on Y . It is used to calibrate the interaction between \mathcal{F} and the target.

Definition 2.8. *For $\kappa > 0$ and $h \in \mathcal{F}$ set $\bar{r}_{\mathbb{M}}(\kappa, h)$ to be*

$$\bar{r}_{\mathbb{M}}(\kappa, h) = \inf \left\{ r : \mathbb{E} \sup_{u \in \mathcal{F}_{h,r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i u(X_i) \cdot (h(X_i) - Y_i) \right| \leq \kappa \sqrt{Nr^2} \right\}. \quad (2.6)$$

For $\sigma > 0$ put $\mathcal{F}_Y^{(\sigma)} = \{f \in \mathcal{F} : \|f(X) - Y\|_{L_2} \leq \sigma\}$ and let $\tilde{r}_{\mathbb{M}}(\kappa, \sigma) = \sup_{h \in \mathcal{F}_Y^{(\sigma)}} \bar{r}_{\mathbb{M}}(\kappa, h)$.

Remark. The role of σ and of $\mathcal{F}_Y^{(\sigma)}$ in Definition 2.8 deserves some explanation. The mean oscillation in Definition 2.8 involves the “multipliers” $(h(X_i) - Y_i)_{i=1}^N$, and the right choice for the centre h is the unknown f^* . While one may take into account the “worst” $h \in \mathcal{F}$, doing so makes little sense, as f^* is the minimizer of the L_2 distance between \mathcal{F} and Y , and for the worst h , $\|h - Y\|_{L_2}$ could be significantly larger than $\|f^* - Y\|_{L_2}$. To overcome this obstacle we assume that an a-priori estimate on the L_2 distance between Y and \mathcal{F} (i.e., a value σ such that $\|f^* - Y\|_{L_2} \leq \sigma$) is available. With this information one still needs to consider the worst centre h , but only among all functions $h \in \mathcal{F}$ that satisfy $\|h - Y\|_{L_2} \leq \sigma$. As we explain in Section 4, thanks to known estimates for the expectation of the supremum of a multiplier process, one only needs to keep in mind that the multipliers $\xi_i = h(X_i) - Y_i$ are independent copies of some random variable ξ that satisfies some moment condition, such as $\|\xi\|_{L_q} \leq L\sigma$ for some $q > 2$ and a suitable constant L .

In light of the results from [29], a realistic alternative to λ^* is

$$r^* = \max\{\lambda_{\mathbb{Q}}(c_1, c_2), \lambda_{\mathbb{M}}(c_1/\sigma, c_2), r_E(c_1), \tilde{r}_{\mathbb{M}}(c_1, \sigma)\}, \quad (2.7)$$

for some constants c_1, c_2 , and when for the given (and unknown) target $Y \in \mathcal{Y}$ one has $\|Y - f^*(X)\|_{L_2} \leq \sigma$. Indeed, in [29] it was shown that under some mild conditions on the learning problem, specified below, ERM performs in \mathcal{F} with accuracy cr^* and constant confidence. Thus, r^* is a potential (and to-date, the best) candidate for the accuracy edge.

Therefore, up to the issue of the true identity of the accuracy edge, the (somewhat vaguely formulated) question of the accuracy/confidence tradeoff is as follows:

Question 2.9. *Is there a learning procedure which, for any reasonable learning problem and any $r \geq c_1 r^*$, performs with accuracy r and confidence $1 - 2 \exp(-c_2 N \min\{1, \sigma^{-2} r^2\})$, thus achieving the optimal accuracy/confidence tradeoff?*

Our main result answers Question 2.9 in the affirmative where our notion of a “reasonable learning problem” is formulated rigorously below.

The first theorem we present requires the following conditions:

Assumption 2.1. *Let L be a constant and let X be distributed according to the measure μ on \mathcal{X} . Given a locally compact, convex class of functions $\mathcal{F} \subset L_2(\mu)$ and $Y \in L_2$, assume that*

- for every $f, h \in \mathcal{F}$, $\|f - h\|_{L_4} \leq L\|f - h\|_{L_2}$;
- for every $f \in \mathcal{F}$, $\|f - Y\|_{L_4} \leq L\|f - Y\|_{L_2}$;
- $\|f^* - Y\|_{L_2} \leq \sigma$ for some known constant $\sigma > 0$.

Theorem 2.10. *Let $L \geq 1$, $\sigma > 0$, and suppose Assumption 2.1. There exist constants c, c_0, c_1 and c_2 that depend only on L for which the following holds. Let*

$$r^*(f^*) = \max\{\lambda_{\mathbb{Q}}(c_1, c_2, f^*), \lambda_{\mathbb{M}}(c_1/\sigma, c_2, f^*), r_E(c_1, f^*), \bar{r}_{\mathbb{M}}(c_1, f^*)\},$$

and fix $r \geq 2r^*(f^*)$.

There exists a procedure that, based on the data $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$ and the values of L , σ and r , selects a function $\hat{f} \in \mathcal{F}$ such that, with probability at least

$$1 - \exp(-c_0 N \min\{1, \sigma^{-2} r^2\}),$$

$$\|\hat{f} - f^*\|_{L_2} \leq cr \quad \text{and} \quad \mathbb{E}((\hat{f}(X) - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}(f^*(X) - Y)^2 + (cr)^2.$$

Of course, the identity of f^* is not known, and therefore, it is not reasonable to expect that $r^*(f^*)$ is known beforehand. A “legal” data-independent choice is any $r \geq 2r^*$ that is larger than $2r^*(f^*)$ regardless of the identity of f^* . In particular, Theorem 2.10 gives the optimal accuracy/confidence tradeoff for any accuracy $r \geq 2r^*$. Alternatively, one may consider the “right” choice of the parameter r as a model selection problem that may be selected using cross validation if independent data are available. We do not discuss the rather straightforward details further here.

Remark. In our main assumption of Theorem 2.10 we use the equivalence between the L_4 and L_2 norms for functions of the form $Y - f(X)$. This allows us to derive bounds in terms of the variance of $Y - f^*(X)$. In fact, if instead of norm equivalence we just have a bound on $\sigma_4 = \|Y - f^*(X)\|_{L_4}$, the arguments work equally well, with σ_4 replacing σ . The sets \mathcal{F}_Y^σ need to be adjusted as well: it should be replaced by all functions in \mathcal{F} whose L_4 distance to Y is at most σ_4 .

It turns out that, when dealing with independent noise, the assumptions required in Theorem 2.10 may be relaxed even further. In particular, we do not require convexity of the class \mathcal{F} and the assumption of norm equivalence may be relaxed:

Theorem 2.11. *Let $q > 2$, $L > 1$, and $\sigma > 0$. There exist constants c, c_0, c_1 and c_2 that depend only on q and L for which the following holds. Let \mathcal{F} be a locally compact class of functions and assume that for every $f \in \text{span}(\mathcal{F})$, $\|f\|_{L_q} \leq L\|f\|_{L_2}$. Assume further that $Y = f_0(X) + W$ for some $f_0 \in \mathcal{F}$ and W that is mean-zero, independent of X , and satisfies $\|W\|_{L_2} \leq \sigma$.*

Let $r^(f^*)$ be as above and fix $r \geq 2r^*(f^*)$. There exists a procedure that, based on the data $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$ and the values of L, q, σ and r , selects a function $\hat{f} \in \mathcal{F}$ such that, with probability at least*

$$1 - \exp(-c_0 N \min\{1, \sigma^{-2} r^2\}) ,$$

$$\|\hat{f} - f_0\|_{L_2} \leq cr \quad \text{and} \quad \mathbb{E}((\hat{f}(X) - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}(f_0(X) - Y)^2 + (cr)^2 .$$

Note that in the case of independent additive noise, the assumptions of Theorem 2.11 are almost the minimal needed for the learning problem to be well defined: norm equivalence for q that may be arbitrarily close to 2 and $W \in L_2$ that perhaps does not have any higher moments. Even under these minimal assumptions, we still obtain the optimal accuracy/confidence tradeoff.

2.1 State of the art

To put Theorem 2.10 in perspective, we describe the current sharpest estimates on the accuracy and confidence of a learning problem, focusing on possibly heavy-tailed distributions.

Firstly, there were no known results that are based on the “averaged” parameter \bar{r}_M . Instead, the interaction between class members and the target were measured using the following “in-probability” version of \bar{r}_M :

Definition 2.12. *For every $\kappa > 0$, $0 \leq \delta \leq 1$ and $h \in \mathcal{F}$, set $r_M(\kappa, \delta, h)$ to be the infimum of the set of all values of r for which*

$$\mathbb{P} \left(\sup_{u \in \mathcal{F}_{h,r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i u(X_i) \cdot (h(X_i) - Y_i) \right| \leq \kappa \sqrt{N} r^2 \right) \geq 1 - \delta .$$

The best known estimate on prediction and estimation in a general convex class are based on a quite weak condition, rather than the norm equivalence we use. Recall that a class \mathcal{F} satisfies a small-ball condition with constants κ_0 and ρ_0 if for every $f, h \in \mathcal{F} \cup \{0\}$,

$$\mathbb{P}(|f - h| \geq \kappa_0 \|f - h\|_{L_2}) \geq \rho_0 .$$

Theorem 2.13. (Mendelson [29].) *Let $\mathcal{F} \subset L_2(\mu)$ be a convex class that satisfies the small-ball condition with constants κ_0 and ρ_0 , and let $Y \in L_2$. If $r = \max\{r_E(c_1), r_M(c_2, \delta, f^*)\}$ and \hat{f} is selected in \mathcal{F} using empirical risk minimization, then, with probability at least*

$$1 - \delta - 2 \exp(-c_3 N) , \tag{2.8}$$

$$\|\hat{f} - f^*\|_{L_2} \leq r \quad \text{and} \quad \mathbb{E}((\hat{f}(X) - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}(f^*(X) - Y)^2 + r^2, \tag{2.9}$$

for constants c_1, c_2 and c_3 that depend only on κ_0 and ρ_0 .

The obvious weakness of Theorem 2.13 is the poor tradeoff between the accuracy term $r_M(c_2, \delta, f^*)$ and the confidence δ . There is no hope of obtaining a high confidence result—say as in (2.4)—, unless both \mathcal{F} and $Y - f^*(X)$ exhibit a sub-Gaussian tail behaviour. If not, then for δ as in (2.4), the value $r_M(c_2, \delta, f^*)$ is very large, and the resulting accuracy estimate is rather useless—far worse than r^* . Moreover, replacing the small-ball assumption with some norm equivalence as in Theorem 2.10 does not improve the outcome. Thus, Theorem 2.13 is significantly weaker than Theorem 2.10 in every aspect. This phenomenon exhibits the nature of ERM: it does not perform with both high accuracy *and* high confidence in heavy-tailed situations, and falls well short of our benchmarks, but still, it was the best available alternative prior to this work.

Regression in \mathbb{R}^n revisited

We now show how our general results imply the optimal accuracy/confidence tradeoff for linear regression in \mathbb{R}^n all the way to a number proportional to the conjectured accuracy edge.

Recall that the class of functions in question is $\mathcal{F} = \{\langle t, \cdot \rangle : t \in \mathbb{R}^n\}$, X is an isotropic random vector on \mathbb{R}^n (i.e., $\mathbb{E} \langle t, X \rangle^2 = 1$ for every t in the Euclidean unit sphere), and $Y = \langle t_0, \cdot \rangle + W$, for $t_0 \in \mathbb{R}^n$ and a symmetric random variable $W \in L_2$ that is independent of X and has variance σ^2 . Thus, $f^* = f_0$ and $\mathcal{F}_{f^*, r} = \{\langle t - t_0, \cdot \rangle : \|t - t_0\|_2 \leq r\}$. If W_1, \dots, W_N are independent copies of W , then by a standard symmetrization argument and since X is isotropic,

$$\begin{aligned} \mathbb{E} \sup_{u \in \mathcal{F}_{h, r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i u(X_i) \cdot (f^*(X_i) - Y_i) \right| &\leq 2 \mathbb{E} \sup_{t \in r B_2^n} \left| \frac{1}{\sqrt{N}} \left\langle \sum_{i=1}^N \varepsilon_i W_i X_i, t \right\rangle \right| \\ &= \frac{2r}{\sqrt{N}} \mathbb{E} \left\| \sum_{i=1}^N \varepsilon_i W_i X_i \right\|_2 \leq 2r \|W\|_{L_2} (\mathbb{E} \|X\|_2^2)^{1/2} = 2r \sigma \sqrt{n} , \end{aligned}$$

where B_2^n is the Euclidean unit ball in \mathbb{R}^n . Now (1.3) follows from Theorem 2.11, assuming that the $L_q(\mu)$ and $L_2(\mu)$ norms are equivalent on $\text{span}(\mathcal{F})$; that is, for every $t \in \mathbb{R}^n$,

$\| \langle t, X \rangle \|_{L_q} \leq L \| \langle X, t \rangle \|_{L_2}$. Indeed, the above shows that $\bar{r}_{\mathbb{M}}(c_1, f^*) \leq 2c_1^{-1} \sigma \sqrt{n/N}$. A similar argument leads to $r_E(c_1) = 0$ when $N \geq cn$ for a constant c that depends only on c_1 . Also, by a volumetric estimate,

$$\log \mathcal{M}(\mathcal{F}_{f^*, r}, \eta r D) = \log \mathcal{M}(r B_2^n, \eta r B_2^n) \sim n \log(2/\eta) ;$$

hence, for $N \geq cn$, $\lambda_{\mathbb{Q}}(c_1) = 0$ and $\lambda_{\mathbb{M}}(c_1/\sigma, c_2) \sim \sigma \sqrt{n/N}$, implying in particular that the lower bound on the accuracy edge is $c\sigma \sqrt{n/N}$.

In other words, when $N \geq cn$, the procedure exhibits the optimal accuracy/confidence tradeoff for any $r \geq c_3 \sigma \sqrt{n/N}$. \blacksquare

In Section 4 we present two more examples, in which we obtain the (previously unknown) optimal accuracy/confidence tradeoff. The first example studies regression in an arbitrary convex, centrally symmetric subset of \mathbb{R}^n when the underlying measure is sub-Gaussian, but the target may be heavy-tailed. In the other example we focus on regression in $\rho B_1^n = \{t \in \mathbb{R}^n : \|t\|_1 \leq \rho\}$, but under significantly weaker assumptions on the underlying measure. Regression in ρB_1^n is of central importance in *sparse recovery*, specifically, in the study of the *basis pursuit* procedure and the *LASSO*. We refer the reader to the books [22, 16, 40] for more information on sparse recovery and on these procedures.

3 The median-of-means tournament

The key to obtaining sharp estimates for both the accuracy and the confidence is identifying a procedure that is not sensitive to atypical values that may occur on a small part of the given sample. Thus, a natural starting point is the conceptually simple and attractive mean estimator, the so-called *median-of-means* estimator. It was proposed, independently, by Nemirovsky and Yudin [35], Jerrum, Valiant, and Vazirani [21], Alon, Matias, and Szegedy [1], and is defined as follows.

Let Z_1, \dots, Z_N be independent, identically distributed real random variables with a finite second moment. The median-of-means estimator of $\mu = \mathbb{E}Z_1$ has parameter $\delta \in [e^{-N/2}, 1)$. Setting $n = \lceil \ln(1/\delta) \rceil$, one may partition $\{1, \dots, N\}$ into n blocks I_1, \dots, I_n , each of cardinality $|I_j| \geq \lfloor N/n \rfloor \geq 2$. Compute the sample mean in each block

$$W_j = \frac{1}{|I_j|} \sum_{i \in I_j} Z_i$$

and define $\hat{\mu}_N^{(\delta)}$ as the median of W_1, \dots, W_n . (If the median is not uniquely defined, here, and in the rest of the paper, we choose the smallest one. Any other choice would work equally well.) It is straightforward to verify that for any $N \geq 4$,

$$\mathbb{P} \left\{ |\hat{\mu}_N^{(\delta)} - \mu| > 2e \sqrt{2 \text{Var}(Z)} \sqrt{\frac{(1 + \ln(1/\delta))}{N}} \right\} \leq \delta , \quad (3.1)$$

where $\text{Var}(Z)$ denotes the variance of Z . In other words, the median-of-means estimator achieves a high (sub-Gaussian) confidence under the minimal assumption that the variance $\text{Var}(Z)$ is finite. Note that the high confidence is valid even though Z can be heavy-tailed, and thus a nontrivial part of the sample $(Z_i)_{i=1}^N$ may be atypical in the sense that Z_i is ‘far away’ from μ .

For properties, applications, and extensions of the median-of-means estimator, we refer to Bubeck, Cesa-Bianchi, and Lugosi [9]. Devroye, Lerasle, Lugosi, and Oliveira [14] Hsu and Sabato [18], Lerasle and Oliveira [27], Minsker [34], Audibert and Catoni [3].

It is tempting to try to replace the empirical means $(1/N) \sum_{i=1}^N (f(X_i) - Y_i)^2$ by some median-of-means estimate of the risk for each $f \in \mathcal{F}$, and select a function in \mathcal{F} that minimizes the estimate. However, due to the nonlinear nature of the median-of-means, it is difficult to control the process of the estimated losses. Instead, the alternative we propose is to estimate the *difference* of the risk for all pairs f, h and organize a two-stage ‘tournament’.

We mention here that Brownlees, Joly, and Lugosi [8] propose empirical minimization based on a different robust mean estimator, a carefully designed M-estimator proposed by Catoni [11]. Under general loss functions they derive analogs of Dudley’s chaining bound for the excess risk. However, the derived bounds are far from giving the optimal rate of convergence under the squared loss.

Without loss of generality and for convenience in the notation, we use $3N$ instead of N for the sample size and assume that $\mathcal{D}_{3N} = (X_i, Y_i)_{i=1}^{3N}$ is the given sample. The sample is split into three equal parts $(X_i, Y_i)_{i=1}^N$, $(X_i, Y_i)_{i=N+1}^{2N}$ and $(X_i, Y_i)_{i=2N+1}^{3N}$. Given the wanted degree of accuracy $r \geq r^*$, the first part of the sample—in fact, just $(X_i)_{i=1}^N$ —is used to estimate pairwise distances within \mathcal{F} , in a sense that will be clarified below. The second part is used in the preliminary round of the tournament. We show that the outcome of the preliminary round is a set $H \subset \mathcal{F}$ that contains f^* and possibly some other functions whose $L_2(\mu)$ distance to f^* is at most cr . The final part of the tournament is a ‘champions league’ round. Participants in that final round are the elements in H (the ‘qualifiers’ of the preliminary round), and the goal of that round is to identify a function $\hat{f} \in H$ whose predictive capabilities are almost optimal, in the sense that

$$\mathbb{E}((\hat{f}(X) - Y)^2 | \mathcal{D}_{3N}) \leq \mathbb{E}(f^*(X) - Y)^2 + cr^2,$$

as required. This last round is only needed to guarantee the desired excess risk. The first two rounds suffice to output a function whose $L_2(\mu)$ distance to f^* is at most cr : one may simply select an arbitrary element of H . Also note that in the setup of Theorem 2.11, the third round is not required as the conditions of the theorem imply that $f^*(X) = \mathbb{E}(Y|X)$, and the excess risk equals the mean squared error; thus, any $\hat{f} \in H$ has the desired performance.

Let us now describe the three stages of the median-of-means tournament in detail.

3.1 The ‘referee’: the distance oracle

Like all good tournaments, ours too requires a ‘referee’, whose role is to decide whether a *match* (described below) is allowed to take place. The referee’s decision is based on a *distance oracle*—a data dependent functional that allows one to crudely identify distances between functions in \mathcal{F} . The functional is constructed via the median-of-means philosophy. Without loss of generality and for ease of exposition, we may assume that N is an integer multiple of ℓ in the next definition.

Definition 3.1. *Let $1 \leq \ell \leq N$ and set I'_j to be the partition of $\{1, \dots, N\}$ to disjoint intervals of cardinality ℓ . Set $k = N/\ell$ and for $v \in \mathbb{R}^N$ let $\text{Med}_\ell(v)$ be the median of the means $(\ell^{-1} \sum_{i \in I'_j} v_i)_{j=1}^k$.*

Recall that one of our assumptions is an L_q - L_2 norm equivalence, that is, that there are $q > 2$ and $L \geq 1$, such that, for every $f \in \text{span}(\mathcal{F})$, $\|f\|_{L_q} \leq L\|f\|_{L_2}$. (In Theorem 2.10

we only consider $q = 4$.) Let $\ell = \ell(q, L)$ to be specified later. For $\mathcal{C}_N = (X_i)_{i=1}^N$ and every $f, h \in \mathcal{F}$, set $v = (|f(X_i) - h(X_i)|)_{i=1}^N$ and put

$$\Phi_{\mathcal{C}_N}(f, h) = \text{Med}_\ell(v) .$$

The functional Φ allows one to identify distances in \mathcal{F} in a crude (isomorphic) way, as the next theorem shows:

Proposition 3.2. *There exist constants $\kappa, \eta, \ell, c > 0$ and $0 < \alpha < 1 < \beta$, all of them depending only on q and L for which the following holds. For a fixed $f^* \in \mathcal{F}$, let $d^* = \max\{\lambda_{\mathbb{Q}}(\kappa, \eta, f^*), r_E(\kappa, f^*)\}$. For any $r \geq d^*$, with probability at least $1 - 2\exp(-cN)$, for every $f \in \mathcal{F}$, one has*

- If $\Phi_{\mathcal{C}_N}(f, f^*) \geq \beta r$ then $\beta^{-1}\Phi_{\mathcal{C}_N}(f, f^*) \leq \|f - f^*\|_{L_2} \leq \alpha^{-1}\Phi_{\mathcal{C}_N}(f, f^*)$.
- If $\Phi_{\mathcal{C}_N}(f, f^*) < \beta r$ then $\|f - f^*\|_{L_2} \leq (\beta/\alpha)r$.

Remark. Replacing d^* by the larger $\max\{\lambda_{\mathbb{Q}}(\kappa, \eta), r_E(\kappa)\}$, which is independent of f^* , a similar assertion to Proposition 3.2 holds for all the pairs $f, h \in \mathcal{F}$. The probability bound in that case is essentially unchanged: $1 - 2\exp(-c'N)$. However, Proposition 3.2 is sufficient for our purposes.

Proposition 3.2 is an immediate modification of Theorem 3.3 from [31]. For the sake of completeness we outline the main components of its proof in the appendix.

Next we introduce the ‘‘distance oracle’’, denoted by \mathcal{DO} . Recall the definition of r^* from (2.7) and note that for the right choice of constants, $r^* \geq d^*$. The distance oracle is adapted to the wanted degree of accuracy, that is, to any fixed $r \geq 2r^*$.

Definition 3.3. *Fix $r \geq 2r^*$. Using the notation of Proposition 3.2, if $\Phi_{\mathcal{C}_N}(f, h) \geq \beta r$ set $\mathcal{DO}(f, h) = 1$, otherwise set $\mathcal{DO}(f, h) = 0$.*

The distance oracle determines if a match between f and h takes place: it does if $\mathcal{DO}(f, h) = 1$ and it is abandoned if $\mathcal{DO}(f, h) = 0$. Note that Proposition 3.2 only shows that \mathcal{DO} is a realistic indication of the distance between pairs when one of the functions is the designated function f^* . This serves our purposes since the designated function we are interested in is the minimizer of the true risk in \mathcal{F} , and the success of the procedure only requires having accurate information on matches that involve f^* , even if we do not know which matches those are.

It follows from Proposition 3.2 that with probability at least $1 - 2\exp(-cN)$ relative to $(X_i)_{i=1}^N$, if a match between f^* and f is allowed to proceed then $\|f - f^*\|_{L_2} \geq r$, while if it is abandoned then $\|f - f^*\|_{L_2} \leq (\beta/\alpha)r$.

3.2 The preliminary round

The goal of the preliminary round is to produce a subset $H \subset \mathcal{F}$ that, with overwhelming probability over the samples $(X_i, Y_i)_{i=N+1}^{2N}$, contains f^* and $\|h - f^*\|_{L_2} \leq (\beta/\alpha)r$ for any $h \in H$.

The round consists of ‘matches’ between every pair $f, h \in \mathcal{F}$, and a match can have three possible outcomes: a win by either side, or a draw (the latter includes abandoned matches because of the ruling of the distance oracle).

Each match is ‘played’ using the second part of the sample, $(X_i, Y_i)_{i=N+1}^{2N}$. The sub-sample is partitioned to n blocks $(I_j)_{j=1}^n$ of cardinality $m = N/n$ each, for a choice of n specified later. Let us note that n depends on the desired degree of accuracy r .

- A match between f and h takes place if the distance oracle, using the first part of the sample $(X_i)_{i=1}^N$, declares that $\mathcal{DO}(f, h) = 1$; otherwise, the match is abandoned and results in a draw.
- Each match is decided according to the n blocks generated by the partition of $(X_i, Y_i)_{i=N+1}^{2N}$, with the j -th block played on the coordinate block I_j . Put

$$B_{f,h}(j) = \frac{1}{m} \sum_{i \in I_j} ((f(X_i) - Y_i)^2 - (h(X_i) - Y_i)^2), \quad 1 \leq j \leq n.$$

The function h defeats f on the j -th block if $B_{f,h}(j) < 0$, and f defeats h if $B_{f,h}(j) > 0$.

- A winner of more than $n/2$ blocks is the winner of the match. If neither function wins more than half of the blocks, the match is drawn.

Definition 3.4. *A function $f \in \mathcal{F}$ qualifies from the preliminary round if it has not lost a single match; that is, it has won or drawn all its matches. The set of “champions” H consists of all functions qualified from the preliminary round.*

The key fact regarding the outcome of the preliminary round is as follows:

Proposition 3.5. *Under the assumptions of Theorem 2.10 or of Theorem 2.11, and using their notation, with probability at least*

$$1 - 2 \exp(-c_0 N \min\{1, \sigma^{-2} r^2\}),$$

with respect to $(X_i, Y_i)_{i=1}^{2N}$, for all $h \in \mathcal{F}$, if $\mathcal{DO}(f^, h) = 1$ then f^* defeats h . In particular, $f^* \in H$ and for any $h \in H$, $\mathcal{DO}(f^*, h) = 0$, and therefore $\|h - f^*\|_{L_2} \leq (\beta/\alpha)r$.*

The proof of Proposition 3.5 is presented in Section 5.1. Note that Propositions 3.2 and 3.5 imply Theorem 2.11. In order to prove the general result of Theorem 2.10, another round of matches is necessary to choose a function from H with small excess risk.

3.3 Champions league

The goal of the second round of the tournament is to choose, among the “champions” selected in the preliminary round, a function with a small excess risk. This round consists of different kind of matches, played between functions in H . Since this round consists of matches between functions in H , conditioned on the ‘good event’ from the preliminary round, every qualifier satisfies that $\|h - f^*\|_{L_2} \leq (\beta/\alpha)r$.

The modified matches are decided using the third part of the sample $(X_i, Y_i)_{i=2N+1}^{3N}$. The aim is to produce a function $\hat{f} \in H$ that has a good excess risk, namely,

$$\mathbb{E}((\hat{f}(X) - Y)^2 | (X_i, Y_i)_{i=2N+1}^{3N}) \leq \mathbb{E}(f^*(X) - Y)^2 + r_1^2,$$

for some r_1 that is bounded by a constant multiple of r .

Setting $\Psi_{h,f} = (h(X) - f(X))(f(X) - Y)$, the significant observation here is that if $\mathbb{E}\Psi_{f^*,f}$ is not very negative, then the prediction error associated with f is small:

Lemma 3.6. *For $\gamma > 0$, if $f \in \mathcal{F}$ satisfies that $\mathbb{E}\Psi_{f^*,f} \geq -\gamma t^2$, then*

$$\mathbb{E}(f(X) - Y)^2 - \mathbb{E}(f^*(X) - Y)^2 \leq 2\gamma t^2.$$

Proof. Observe that for every $f \in \mathcal{F}$

$$(f(X) - Y)^2 - (f^*(X) - Y)^2 = (f(X) - f^*(X))^2 + 2(f(X) - f^*(X))(f^*(X) - Y) ,$$

and

$$\begin{aligned} (f(X) - f^*(X))(f^*(X) - Y) &= (f(X) - f^*(X))((f^*(X) - f(X)) + (f(X) - Y)) \\ &= - (f(X) - f^*(X))^2 - \Psi_{f^*,f} . \end{aligned}$$

Therefore,

$$\mathbb{E}(f(X) - Y)^2 - \mathbb{E}(f^*(X) - Y)^2 \leq -2\mathbb{E}\Psi_{f^*,f} \leq 2\gamma t^2 .$$

■

The role of the “champions league” round is to use the third part of the sample to select $\hat{f} \in H$ for which $\mathbb{E}(\Psi_{f^*,\hat{f}} | (X_i, Y_i)_{i=1}^{2N}) \geq -\gamma r_1^2$, for a suitable constant $\gamma > 0$ and r_1 that is proportional to r .

The matches in the champions league consist of “home-and-away” legs:

Definition 3.7. Given a sample $(X_i, Y_i)_{i=2N+1}^{3N}$, let $(I_j)_{j=1}^n$ be the partition of $\{2N+1, 3N\}$ to n blocks, for the same value of n as in the preliminary round. Let β and α be as in Proposition 3.5 and set $r_1 = 2(\beta/\alpha)r$. The function f wins its home match against h if

$$\frac{2}{m} \sum_{i \in I_j} \Psi_{h,f}(X_i, Y_i) \geq -r_1^2/10$$

on more than $n/2$ of the blocks I_j .

We select as \hat{f} any “champion” in H that wins all of its home matches.

The main result regarding the champions league is as follows:

Proposition 3.8. Let $H \subset \mathcal{F}$ as above. Under the assumptions of Theorem 2.10 and using its notation, with probability at least

$$1 - 2 \exp(-c_0 N \min\{1, \sigma^{-2} r^2\})$$

with respect to $(X_i, Y_i)_{i=2N+1}^{3N}$, one has:

- f^* wins all of its home matches, and
- if $\mathbb{E}\Psi_{f^*,f} \leq -2r_1^2$, then f loses its home match against f^* .

Thus, on this event, the set of possible champions is nonempty (since it contains f^*), and any other champion satisfies that $\mathbb{E}\Psi_{f^*,f} \geq -2r_1^2$ and therefore, by Lemma 3.6,

$$\mathbb{E}((\hat{f}(X) - Y)^2 | (X_i, Y_i)_{i=2N+1}^{3N}) - \mathbb{E}(f^*(X) - Y)^2 \leq 4r_1^2 .$$

The proof of Proposition 3.8 is presented in Section 5.2.

The combination of Propositions 3.2, 3.5, and 3.8 yields the proof of Theorem 2.10.

4 Examples

Before turning to the proofs of our main results, let us present some explicit examples of applications of Theorem 2.10.

It is an unrealistic hope to obtain a simple characterization of all involved complexity parameters in every example. Indeed, to get good bounds for $\lambda_{\mathbb{Q}}$ and $\lambda_{\mathbb{M}}$, one has to obtain sharp estimates on covering numbers of what is almost an arbitrary set and with respect to the $L_2(\mu)$ norm for an arbitrary probability measure μ . On the other hand, $\bar{r}_{\mathbb{M}}$ and r_E depend on the oscillation of general multiplier and empirical processes, respectively. Both types of estimates are of central importance in modern mathematics and have been the subject of thorough research, but they are by no means completely understood.

Having said this, there are many interesting cases in which sharp estimates may be derived. In what follows we focus on two such examples. The first is rather general: linear regression performed in a convex, centrally symmetric set $T \subset \mathbb{R}^n$, under the assumption that the underlying random vector X is L -sub-Gaussian (see the definition below). However, and unlike the results in [24], the ‘noise’ $Y - f^*(X)$ may be heavy-tailed.

The second example is similar to the first one and is motivated by questions in *sparse recovery*: linear regression in the set $T = \rho B_1^n = \{t : \|t\|_1 \leq \rho\}$. The difference between this example and the first one lies in the assumption on X . In the second example X is not assumed to be L -sub-Gaussian, but rather satisfies a much weaker moment condition, the same condition that is needed to ensure that the *basis pursuit* algorithm has a unique solution with the optimal number of measurements (see [25]).

Both examples lead to explicit estimates on the accuracy and confidence of the median-of-means tournament. The estimates are better than the known bounds and hold with optimal confidence for accuracy larger than cr^* (though in general, the true identity of the accuracy edge is an open question). Moreover, in the second example, of ρB_1^n , one may show that r^* is proportional to the accuracy edge, and the optimal tradeoff holds all the way down to that value.

4.1 Coverings and Gaussian processes

Let X be an isotropic random vector in \mathbb{R}^n , that is, for every $t \in \mathbb{R}^n$, $\mathbb{E} \langle X, t \rangle^2 = \|t\|_2^2$. The assumption that X is isotropic only serves clarity of the illustration. Indeed, the $L_2(\mu)$ metric endowed on \mathbb{R}^n via the identification of $t \in \mathbb{R}^n$ with the linear functional $\langle \cdot, t \rangle$ is the standard Euclidean metric. Thus, D (the unit ball in $L_2(\mu)$) can be identified with the Euclidean unit ball in \mathbb{R}^n . While it is possible to extend the results presented below to X with a general covariance structure (in which case, D is identified with an ellipsoid in \mathbb{R}^n), the isotropic example is interesting enough to serve as a proof of concept.

Thanks to the isotropicity assumption, if T is a convex and centrally-symmetric set and $F = \{\langle \cdot, t \rangle : t \in T\}$, then for every $h \in F$,

$$\lambda_{\mathbb{Q}}(\kappa, \eta, h) \leq \inf\{r : \log \mathcal{M}(2T \cap rB_2^n, \eta rB_2^n) \leq \kappa^2 N\} ,$$

and

$$\lambda_{\mathbb{M}}(\kappa, \eta, h) \leq \inf\{r : \log \mathcal{M}(2T \cap rB_2^n, \eta rB_2^n) \leq \kappa^2 Nr^2\} .$$

A standard, though sometimes suboptimal, method to estimate covering/packing numbers relies on the theory of Gaussian processes, specifically, on Sudakov’s inequality. We formulate

it only in the case we need here. Denote by $G = (g_i)_{i=1}^n$ a standard Gaussian vector in \mathbb{R}^n . For $T \subset \mathbb{R}^n$ let

$$\ell_*(T) = \mathbb{E} \sup_{t \in T} \langle G, t \rangle$$

the *mean-width* of T with respect to the Gaussian measure.

Proposition 4.1. (Sudakov [36].) *There exists an absolute constant c , such that, for any $T \subset \mathbb{R}^n$ and every $\varepsilon > 0$,*

$$\varepsilon \sqrt{\log \mathcal{M}(T, \varepsilon B_2^n)} \leq c \ell_*(T) . \quad (4.1)$$

Applying Proposition 4.1 it follows that

$$\log \mathcal{M}(2T \cap rB_2^n, \eta r B_2^n) \leq \left(c \frac{\ell_*(2T \cap rB_2^n)}{\eta r} \right)^2 .$$

Hence,

$$\lambda_{\mathbb{Q}}(\kappa, \eta, h) \leq \inf \{ r : \ell_*(2T \cap rB_2^n) \leq (\kappa \eta / c) r \sqrt{N} \} , \quad (4.2)$$

and

$$\lambda_{\mathbb{M}}(\kappa, \eta, h) \leq \inf \{ r : \ell_*(2T \cap rB_2^n) \leq (\kappa \eta / c) r^2 \sqrt{N} \} . \quad (4.3)$$

We emphasize again that replacing $\lambda_{\mathbb{Q}}$ and $\lambda_{\mathbb{M}}$ with these upper estimates, is, at times, suboptimal. We refer the reader to [24] for more details on this issue.

The other two parameters involved in Theorem 2.10, namely, r_E and $\bar{r}_{\mathbb{M}}$, measure the oscillation of multiplier and empirical processes. The analysis of such processes is highly nontrivial—even when just considering their limits as the sample size N tends to infinity, and one expects convergence to the limiting Gaussian process (see, for example, the book [15] for a detailed exposition of such limit theorems). Because the estimates we require are non-asymptotic, in general they are much harder to obtain.

The following notion makes the task of obtaining such bounds more manageable, though still nontrivial.

Definition 4.2. *A random vector X in \mathbb{R}^n is L -sub-Gaussian if for every $t \in \mathbb{R}^n$ and any $p \geq 2$,*

$$\| \langle X, t \rangle \|_{L_p} \leq L \sqrt{p} \| \langle X, t \rangle \|_{L_2} .$$

Note that if X is L -sub-Gaussian and isotropic, then for every $t \in \mathbb{R}^n$, $\| \langle X, t \rangle \|_{L_p} \leq L \sqrt{p} \| t \|_2$, because $\| \langle X, t \rangle \|_{L_2} = \| t \|_2$.

The simplest examples of isotropic, L -sub-Gaussian random vectors are vectors with independent, mean-zero, variance 1 components that are sub-Gaussian. For instance, the standard Gaussian vector $(g_i)_{i=1}^n$, and $(\varepsilon_i)_{i=1}^n$, whose components are independent, symmetric random signs are both L -sub-Gaussian for a constant that is independent of the dimension n . Another family of examples consists of the random vectors whose density is uniform on sets of the form $\{ t : \| t \|_p \leq cn^{1/p} \}$ for some $p \geq 2$, normalized to have volume 1. Again, L is an absolute constant, independent of n and p (see, e.g., [5, 4]).

The reason for considering a sub-Gaussian random vector X is that, by Talagrand's theory of *generic chaining* (see the book [37] for an extensive exposition on the subject), the oscillations in question may be controlled using the oscillation of the corresponding Gaussian process. For example, the next result describes how the expected supremum of a multiplier process is upper bounded in terms of the Gaussian mean-width ℓ_* . Although it is formulated in \mathbb{R}^n , it holds in a far more general context (see [33]).

Proposition 4.3. *Let $q > 2$ and let X be an isotropic, L -sub-Gaussian random vector in \mathbb{R}^n . There exists a constant $c = c(q)$ such that the following holds. Let $\xi \in L_q$ be a random variable (not necessarily independent of X) and let $(X_i, \xi_i)_{i=1}^N$ be independent copies of (X, ξ) . Then, for any $T \subset \mathbb{R}^n$,*

$$\mathbb{E} \sup_{t \in T} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \xi_i \langle X_i, t \rangle \right| \leq cL \|\xi\|_{L_q} \ell_*(T) , \quad (4.4)$$

where $(\varepsilon_i)_{i=1}^N$ are independent, symmetric signs that are independent of $(X_i, \xi_i)_{i=1}^N$.

Note that if ξ is heavy-tailed there is no hope of obtaining a high-probability version of Proposition 4.3. In fact, if all one knows is that ξ belongs to L_q , one cannot hope that

$$\sup_{t \in T} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \xi_i \langle X_i, t \rangle \right| \leq cL \|\xi\|_{L_q} \ell_*(T)$$

with a probability higher than $1 - c_1/N^{(q/2)-1}$.

This exhibits, yet again, the weakness of Theorem 2.13, which is based on the ‘‘in-probability’’ parameter $r_{\mathbb{M}}$. Setting $f^*(X) = \langle X, t^* \rangle$ and if $\xi = f^*(X) - Y$ is heavy-tailed, the oscillation is simply too big on an event with the required confidence (2.4), but the mean oscillation is well behaved.

Applying Proposition 4.3, it is evident that

$$r_E(\kappa) \leq \inf \left\{ r : \ell_*(2T \cap rB_2^n) \leq (\kappa/cL)\sqrt{Nr} \right\} \quad (4.5)$$

and

$$\bar{r}_{\mathbb{M}}(\kappa, t) = \inf \left\{ r : \|Y - \langle X, t \rangle\|_{L_q} \ell_*(2T \cap rB_2^n) \leq (\kappa/cL)\sqrt{Nr^2} \right\} . \quad (4.6)$$

Thus, assuming that $\|Y - \langle X, t \rangle\|_{L_q} \leq L\|Y - \langle X, t \rangle\|_{L_2}$, as we do in Assumption 2.1, it follows that

$$\tilde{r}_{\mathbb{M}}(\kappa, \sigma) \leq \inf \left\{ r : \ell_*(2T \cap rB_2^n) \leq (\kappa/cL^2\sigma)\sqrt{Nr} \right\} .$$

Definition 4.4. *For constants c_1, c_2 and σ , let*

$$s_{\mathbb{M}}(c_1, \sigma) = \inf \left\{ r : \ell_*(2T \cap rB_2^n) \leq (c_1/\sigma)\sqrt{Nr} \right\} ,$$

and

$$s_{\mathbb{Q}}(c_2) = \inf \left\{ r : \ell_*(2T \cap rB_2^n) \leq c_2\sqrt{Nr} \right\} .$$

Hence, for every $t^* \in T$,

$$\max\{\lambda_{\mathbb{Q}}(c_1, c_2, t^*), \lambda_{\mathbb{M}}(c_1/\sigma, c_2, t^*), r_E(c_1, t^*), \bar{r}_{\mathbb{M}}(c_1, t^*)\} \leq \max\{s_{\mathbb{Q}}(c_3, \sigma), s_{\mathbb{M}}(c_4/\sigma)\} ,$$

for constants c_1, c_2, c_3, c_4 that depend only on q and L .

We obtain the following consequence of Theorem 2.10:

Theorem 4.5. *Let X be an isotropic, L -sub-Gaussian random vector in \mathbb{R}^n , and let $T \subset \mathbb{R}^n$ be a convex, centrally-symmetric set. Let $q > 2$ and assume that $\|Y - \langle X, t \rangle\|_{L_q} \leq L\|Y - \langle X, t \rangle\|_{L_2}$ for every $t \in T$. Then, with probability at least $1 - 2\exp(-c_1 N \min\{1, \sigma^{-2} s_M^2(c_2, \sigma)\})$, the median-of-means tournament produces \hat{t} such that*

$$\|\hat{t} - t^*\|_{L_2} \leq c_3 s^* \quad \text{and} \quad \mathbb{E}((\langle X, \hat{t} \rangle - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}(\langle X, t^* \rangle - Y)^2 + (c_4 s^*)^2,$$

where

$$s^* = \max\{s_{\mathbb{M}}(c_2, \sigma), s_{\mathbb{Q}}(c_5)\},$$

and the constants c_1, \dots, c_5 depend only on L and q .

Moreover, for any $s \geq s^*$,

$$\|\hat{t} - t^*\|_{L_2} \leq c_3 s \quad \text{and} \quad \mathbb{E}((\langle X, \hat{t} \rangle - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}(\langle X, t^* \rangle - Y)^2 + (c_4 s)^2,$$

with probability at least $1 - 2\exp(-c_1 N \min\{1, \sigma^{-2} s^2\})$, exhibiting the optimal accuracy-confidence tradeoff.

Theorem 4.5 improves Theorem A from [24], where it was shown that ERM produces \hat{t} with the same accuracy and confidence as in the first part of Theorem 4.5, but only when $\|Y - \langle X, t \rangle\|_{L_q} \leq L\sqrt{q}\|Y - \langle X, t \rangle\|_{L_2}$ for every $q > 2$ and every $t \in T$. In other words, Theorem A from [24] is based on the assumption that each $Y - \langle X, t \rangle$ is an L -sub-Gaussian random variable, and holds only for the accuracy level s^* . In contrast, Theorem 4.5 shows that the median-of-means tournament performs in an optimal way in heavy-tailed situations that are totally out of reach for ERM and for the entire range $s \geq cs^*$.

Observe that the only range of accuracies in which Theorem 4.5 is (perhaps) suboptimal, is when

$$\lambda^* = \max\{\lambda_{\mathbb{Q}}(\kappa_1, \eta_1), \lambda_{\mathbb{M}}(\kappa_2/\sigma, \eta_2)\} \leq s \leq s^* \quad (4.7)$$

for well chosen values of κ_i, η_i , $i = 1, 2$; that is, for values that are larger than the known lower estimate on the accuracy edge for such problems. As noted in [24], there are many examples in which λ^* and s^* are equivalent (roughly speaking, this happens when Sudakov's inequality is sharp). In such cases the median-of-means tournament is optimal in the entire range of accessible accuracies.

One important class of sets in which this equivalence is true is $\rho B_1^n = \{t : \|t\|_1 \leq \rho\}$ (see [24] for the proof). In light of Theorem 4.5, the median-of-means tournament performs in an optimal way in ρB_1^n . Moreover, it turns out that one may relax the sub-Gaussian assumption on X and still obtain the optimal behaviour ρB_1^n , as we show next.

4.2 ρB_1^n – Sparse recovery sets

It is well understood that classical sparse recovery procedures, such as *basis pursuit* or *LASSO* rely heavily on the geometry of B_1^n . Indeed, LASSO selects \hat{t} , the minimizer in \mathbb{R}^n of the functional

$$t \rightarrow \frac{1}{N} \sum_{i=1}^N (Y_i - \langle X_i, t \rangle)^2 + \lambda \|t\|_1.$$

Being a penalized version of ERM, the analysis of LASSO is equivalent to the study of ERM in the sets ρB_1^n for an arbitrary choice of ρ .

While we defer the question of a “LASSO-tournament” procedure to future work, it is clear that the first step in that direction is to explore the median-of-means tournament in ρB_1^n . Instead of the sub-Gaussian assumption used in Theorem 4.5, the assumption we use follows the path of [25]:

Assumption 4.1. *Let X be an isotropic random vector and $\kappa \geq 1$. Assume that for every $t \in \mathbb{R}^n$ and any $2 \leq p \leq \kappa \log n$, $\|\langle X, t \rangle\|_{L_p} \leq L\sqrt{p}\|\langle X, t \rangle\|_{L_2}$.*

Note that the coordinates of X need not be independent and X may be far from being an L -sub-Gaussian random vector. Indeed, it is required that linear functionals $\langle \cdot, t \rangle$ satisfy a sub-Gaussian moment growth only up to the logarithm of the dimension, and it is possible that some do not have any higher moment beyond $p = \kappa \log n$.

It turns out (see [25]) that if X satisfies Assumption 4.1, then $N^{-1/2} \sum_{i=1}^N \langle X_i, \cdot \rangle e_i$, the random matrix whose rows are independent copies of X , exhibits the best possible sparse recovery features. For example, one requires $N \sim s \log(en/s)$ random measurements $\langle X_i, t \rangle$ to recover any s -sparse vector t using basis pursuit, and a similar type of estimate holds in the “noisy” setup for the LASSO. Moreover, one cannot relax the moment condition in Assumption 4.1 and still get the same recovery properties.

Of course, since both basis pursuit and LASSO are variations of ERM, they suffer from the same weaknesses as ERM. As such, when the given measurements are $(\langle X_i, t \rangle)_{i=1}^N$ for a heavy-tailed X , the confidence with which the recovery properties hold is suboptimal, and very different from the confidence one has when X is the standard Gaussian vector in \mathbb{R}^n .

We show that as far as regression in ρB_1^n goes, the median-of-means tournament yields the optimal, “Gaussian” behaviour even when X only satisfies Assumption 4.1 and $Y - \langle t^*, X \rangle$ is heavy-tailed. To this end, we need sharp bounds on the parameters that are used to define r^* in Theorem 2.10.

As was noted earlier, $\lambda_{\mathbb{Q}}$ and $\lambda_{\mathbb{M}}$ depend only on the covariance structure endowed on \mathbb{R}^n by $L_2(\mu)$, and since X is isotropic, the $L_2(\mu)$ metric corresponds to the standard Euclidean norm. Therefore, the difficulty lies in bounding r_E and $\tilde{r}_{\mathbb{M}}$, and specifically, in extending Proposition 4.3 beyond the L -sub-Gaussian case. While Theorem 4.3 is general and holds for any subset of \mathbb{R}^n , here it is needed for very specific sets, namely $T = \rho B_1^n \cap r B_2^n$. Such indexing sets fall within the scope of Theorem 1.6 from [32].

Proposition 4.6. *Let $q > 2$ and let X satisfy Assumption 4.1 for $\kappa = c_1(q)$. Let $\xi \in L_q$ be a random variable (not necessarily independent of X) and let $(X_i, \xi_i)_{i=1}^N$ be independent copies of (X, ξ) . Then*

$$\mathbb{E} \sup_{t \in \rho B_1^n \cap r B_2^n} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \xi_i \langle X_i, t \rangle \right| \leq c_2(q) L \|\xi\|_{L_q} \ell_*(\rho B_1^n \cap r B_2^n). \quad (4.8)$$

Thus, Assumption 4.1 suffices to ensure that r_E and $\tilde{r}_{\mathbb{M}}(\kappa, \sigma)$ may be controlled as if X were L -sub-Gaussian. All that remains is to estimate

$$\log \mathcal{M}(\rho B_1^n \cap r B_2^n, \eta r B_2^n) \quad \text{and} \quad \ell_*(\rho B_1^n \cap r B_2^n)$$

which are well-understood quantities.

To put them in a more familiar form, set $\sqrt{s} = \rho/r$ and observe that

$$\ell_*(\rho B_1^n \cap r B_2^n) = r \ell_*((\rho/r) B_1^n \cap B_2) = r \ell_*(\sqrt{s} B_1^n \cap B_2^n)$$

and

$$\mathcal{M}(\rho B_1^n \cap r B_2^n, \eta r) = \mathcal{M}((\rho/r)B_1^n \cap B_2^n, \eta B_2^n) = \mathcal{M}(\sqrt{s}B_1^n \cap B_2^n, \eta B_2^n) .$$

Recall that if $1 \leq s \leq n$ and V_s is the set of s -sparse vectors in the Euclidean unit sphere (i.e., those with at most s nonzero components), then $\text{conv}(V_s) \subset \sqrt{s}B_1^n \cap B_2^n \subset C \cdot \text{conv}(V_s)$ for a suitable absolute constant C . Using what are by now standard estimates (see, e.g., [24]),

$$\ell_*(\sqrt{s}B_1^n \cap B_2^n) \sim \sqrt{s \log(en/s)}, \quad \text{and} \quad \log \mathcal{M}(\sqrt{s}B_1^n \cap B_2^n, \eta B_2^n) \sim s \log(en/\eta s) .$$

The estimates are simpler outside the range $1 \leq s \leq n$: when $\rho/r \leq 1$ then $\rho B_1^n \cap r B_2^n = \rho B_1^n$ and when $\rho/r \geq \sqrt{n}$ then $\rho B_1^n \cap r B_2^n = r B_2^n$. Again, the required estimates on \mathcal{M} and ℓ_* are standard and may be found, for example, in [24].

Using these observations, and with the same (tedious) computation as in [24], one obtains the following: let c_1 and c_2 be well-chosen absolute constants and set

$$v_{\mathbb{M}}^2 = \begin{cases} \frac{\rho\sigma}{\sqrt{N}} \sqrt{\log\left(\frac{2c_1 n \sigma}{\sqrt{N}\rho}\right)} & \text{if } N \leq c_1 n^2 \sigma^2 / \rho^2 \\ \frac{\sigma^2 n}{N} & \text{if } N > c_1 n^2 \sigma^2 / \rho^2 , \end{cases}$$

and

$$v_{\mathbb{Q}}^2 = \begin{cases} \frac{\rho^2}{N} \log\left(\frac{2c_2 n}{N}\right) & \text{if } N \leq c_2 n , \\ 0 & \text{if } N > c_2 n . \end{cases}$$

Then

- λ^* , the lower estimate on the accuracy edge, satisfies $\lambda^* \geq c_3 \max\{v_{\mathbb{Q}}, v_{\mathbb{M}}\}$; thus, there is no learning procedure in ρB_1^n that can perform with a better accuracy than $c_3 \max\{v_{\mathbb{Q}}, v_{\mathbb{M}}\}$ with a higher confidence than 3/4;
- For any $v \geq c_4 \max\{v_{\mathbb{Q}}, v_{\mathbb{M}}\}$, the median-of-means tournament achieves the accuracy v with the optimal confidence $1 - 2 \exp(-c_5 N \min\{1, \sigma^{-2} v^2\})$, thus exhibiting the optimal accuracy/confidence tradeoff up to a level that is proportional to λ^* .

Formally:

Corollary 4.7. *Let $q > 2$ and assume that X satisfies Assumption 4.1 with a constant $\kappa = c(q)$. Assume further that for every $t \in \rho B_1^n$, $\|Y - \langle X, t \rangle\|_{L_q} \leq L \|Y - \langle X, t \rangle\|_{L_2}$. Then for every $v \geq c_4 \max\{v_{\mathbb{Q}}, v_{\mathbb{M}}\}$, with probability at least $1 - 2 \exp(-c_5 N \min\{1, \sigma^{-2} v^2\})$, the median-of-means tournament produces $\hat{t} \in \rho B_1^n$ that satisfies*

$$\|\hat{t} - t^*\|_{\ell_2^n} \leq c_6 v \quad \text{and} \quad \mathbb{E}((\langle X, \hat{t} \rangle - Y)^2 | \mathcal{D}_N) \leq \mathbb{E}((\langle X, t^* \rangle - Y)^2) + (c_6 v)^2 ,$$

for constants c_4, c_5, c_6 that depend only on q and L .

The advantage of the median-of-means tournament over ERM is clear: it performs in ρB_1^n with the optimal accuracy and confidence, starting a constant factor away from the level of accuracy that can be attained only with constant confidence, and it does so under a heavy-tailed assumption both on X and on Y . In contrast, ERM (which was the ‘‘record holder’’ prior to this work) achieves the optimal performance only in a purely sub-Gaussian setup and does so only for one level of accuracy, of the order of $\max\{v_{\mathbb{Q}}, v_{\mathbb{M}}\}$.

5 Proofs

Let us begin by considering the structure of the various indexing sets involved in the proofs, paying particular attention to the way their structure affects the regularity of the parameters we defined earlier.

For any class \mathcal{F} and every $h \in \mathcal{F}$, $\mathcal{F} - h \subset \mathcal{F} - \mathcal{F}$, and in particular,

$$\mathcal{F}_{r,h} = \text{star}(\mathcal{F} - h, 0) \cap rD \subset \text{star}(\mathcal{F} - \mathcal{F}, 0) \cap rD.$$

Thus, for all the complexity parameters defined above, one may avoid the need to take the supremum over all possible choices of centres by considering a slightly larger indexing set, namely, $\text{star}(\mathcal{F} - \mathcal{F}, 0) \cap rD$. Moreover, if \mathcal{F} is convex, then $\mathcal{F} - \mathcal{F}$ is both convex and centrally symmetric, and if \mathcal{F} happens to be convex and centrally symmetric then $\mathcal{F} - \mathcal{F} = 2\mathcal{F}$ and $\text{star}(\mathcal{F} - \mathcal{F}, 0) \cap rD = 2\mathcal{F} \cap rD$.

The fact that $\text{star}(\mathcal{F} - h, 0)$ is star-shaped around 0 leads to important regularity properties of the parameters we use. Recall that S is the unit sphere in $L_2(\mu)$ and observe that if V is star-shaped around 0 and $v \in V \cap rS$, then for every $r' \leq r$, V contains a ‘scaled-down’ version of v , of norm r' : for $\alpha = r'/r < 1$, $\alpha v \in V$. Hence, if $\phi(r) = \mathbb{E} \sup_{v \in V \cap rD} \left| \sum_{i=1}^N \varepsilon_i v(X_i) \right|$ and $r' \leq r$, then $\phi(r') \geq (r'/r)\phi(r)$.

This argument shows that when $r > r_E(\kappa, h)$,

$$\mathbb{E} \sup_{v \in \mathcal{F}_{h,r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i v(X_i) \right| \leq \kappa \sqrt{N} r ,$$

and when $r < r_E(\kappa, h)$, the reverse inequality holds. In a similar fashion, if $r \geq \bar{r}_M(\kappa, h)$, then

$$\mathbb{E} \sup_{v \in \mathcal{F}_{h,r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i v(X_i) (h(X_i) - Y_i) \right| \leq \kappa \sqrt{N} r^2 ,$$

whereas if $r < \bar{r}_M(\kappa, h)$, the reverse inequality holds.

Similar arguments apply for $\lambda_{\mathbb{Q}}$ and $\lambda_{\mathbb{M}}$: if V is star-shaped around 0 and if $\{v_1, \dots, v_M\}$ is an ηr -separated subset of $V \cap rD$ and $0 < \alpha < 1$, then $\{\alpha v_1, \dots, \alpha v_M\}$ is an $\eta(\alpha r)$ -separated subset of $V \cap (\alpha r)D$. Therefore,

$$\log \mathcal{M}(V \cap \alpha rD, \eta \alpha rD) \geq \log \mathcal{M}(V \cap rD, \eta rD) , \quad (5.1)$$

implying that the function $r \rightarrow \log \mathcal{M}(V \cap rD, \eta rD)$ is monotone decreasing. Moreover, if $\log \mathcal{M}(V \cap r'D, \eta r'D) \leq \kappa^2 N$, or if $\log \mathcal{M}(V \cap r'D, \eta r'D) \leq \kappa^2 N (r')^2$, the same is true for every $r \geq r'$. Therefore, if $r > \lambda_{\mathbb{Q}}(\kappa, \eta, h)$ then

$$\log \mathcal{M}(\mathcal{F}_{h,r}, \eta rD) \leq \kappa^2 N ,$$

and if $r < \lambda_{\mathbb{Q}}(\kappa, \eta, h)$, the reverse inequality holds. In a similar fashion, if $r > \lambda_{\mathbb{M}}(\kappa, \eta, h)$ then

$$\log \mathcal{M}(\mathcal{F}_{h,r}, \eta rD) \leq \kappa^2 N r^2 ,$$

while if $r < \lambda_{\mathbb{M}}(\kappa, \eta, h)$, the reverse inequality holds.

These observations allow one to choose a level of accuracy such as r^* by ‘intersecting’ multiple conditions like the ones appearing in the definitions of $\lambda_{\mathbb{Q}}$, $\lambda_{\mathbb{M}}$, r_E and \bar{r}_M , simply because

each one of the required inequalities holds for any level larger than $\lambda_{\mathbb{Q}}, \lambda_{\mathbb{M}}, r_E$ and $\bar{r}_{\mathbb{M}}$, respectively. An additional feature is that one may “combine” conditions by decreasing the constants involved in the definitions. For example, $\max\{r_E(\kappa_1, h), r_E(\kappa_2, h)\} = r_E(\min\{\kappa_1, \kappa_2\}, h)$ and similar observations hold for $\lambda_{\mathbb{Q}}, \lambda_{\mathbb{M}}$ and $\bar{r}_{\mathbb{M}}$.

For the sake of transparency, we do not specify all the constants involved in the definitions of the fixed points right from the start. Instead, we collect conditions on these constants and use the fact that one may “combine” and “intersect” them. It turns out that all the constants depend on only two parameters: $q > 2$, for which the L_q norm is equivalent to the L_2 norm, and the constant L . In what follows, we denote by $c(q, L)$ a constant that depends only on q and L , for the values of q and L in Theorem 2.10 (where only $q = 4$ is considered) or in Theorem 2.11, when q can be arbitrarily close to 2—at the price of worse constants, of course.

With this in mind, let $\kappa_1, \kappa_2, \kappa_3$ and η be constants that will be specified later and that depend only on q and L . Fix $f^* \in \mathcal{F}$ and consider r^* for which, for every $r > r^*$,

$$\mathbb{E} \sup_{f \in \mathcal{F}_{f^*, r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i (f - f^*)(X_i) \right| \leq \kappa_1 \sqrt{N} r, \quad (5.2)$$

$$\mathbb{E} \sup_{f \in \mathcal{F}_{f^*, r}} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i (f^*(X_i) - Y_i) (f - f^*)(X_i) \right| \leq \kappa_2 \sqrt{N} r^2 \quad (5.3)$$

and

$$\log \mathcal{M}(\mathcal{F}_{f^*, r}, \eta r D) \leq \kappa_3^2 N \min\{1, r^2/\sigma^2\}. \quad (5.4)$$

The value of r^* from (2.7) will be given by selecting right constants $\kappa_1, \kappa_2, \kappa_3$ and η . We consider an accuracy $r \geq 2r^*$, and for the rest of this section we fix its value.

Let us introduce the main parameter n used in the tournament, that is, the number of blocks into which the second part of the sample $(X_i, Y_i)_{i=N+1}^{2N}$, and the third part of the sample $(X_i, Y_i)_{i=2N+1}^{3N}$ are partitioned. To this end, let $0 < \tau < 1/4$, set $0 < \theta \leq \tau$, to be specified later, and put $\lambda_{\mathbb{M}} = \lambda_{\mathbb{M}}(\kappa_3/\sigma, \eta, f^*)$. Set

$$n = \theta N \min \left\{ 1, \left(\frac{r}{\sigma} \right)^2 \right\}. \quad (5.5)$$

Hence, n depends on the wanted accuracy r . Also, $n \leq \tau N$ and without loss of generality we may assume that both n and $m = N/n$ are integers. Also, note that $m \geq 1/\tau$.

5.1 The preliminary round—proof

In this section we prove Proposition 3.5.

Recall that by Proposition 3.2 and the resulting condition on r , one has that, with probability at least $1 - 2 \exp(-c_0 N)$ with respect to $(X_i)_{i=1}^N$, if a match involving f^* is allowed to take place (i.e., if $\mathcal{DO}(f^*, f) = 1$), then $\|f^* - f\|_{L_2} \geq r$; and if the match is abandoned (i.e., $\mathcal{DO}(f^*, h) = 0$), then $\|f^* - f\|_{L_2} \leq (\beta/\alpha)r$. The constants $\alpha < 1 < \beta$ and c_0 depend only on L and q .

Therefore, to establish Proposition 3.5 it is enough to show that with probability $1 - 2 \exp(-cn)$, if $\|f - f^*\|_{L_2} \geq r$, then f^* wins its match against f . In other words, that $B_{f, f^*}(j) > 0$ for the majority of the blocks in any such match.

Let us begin by exploring the situation in a match between f^* and f , knowing that $\|f^* - f\|_{L_2} \geq r \geq 2r^*$, as above.

Clearly, for every $f, h \in \mathcal{F}$,

$$(f(X) - Y)^2 - (h(X) - Y)^2 = (f(X) - h(X))^2 + 2(f(X) - h(X)) \cdot (h(X) - Y) .$$

Set

$$\mathbb{Q}_{f,h} = \frac{1}{m} \sum_{i=1}^m (f(X_i) - h(X_i))^2 \quad \text{and} \quad \mathbb{M}_{f,h} = \frac{2}{m} \sum_{i=1}^m (f(X_i) - h(X_i)) \cdot (h(X_i) - Y_i) .$$

We introduce the notation $\xi = f^*(X) - Y$ and $\xi_i = f^*(X_i) - Y_i$. Partitioning the sample $(X_i, Y_i)_{i=N+1}^{2N}$ into the n blocks $(I_j)_{j=1}^n$, each one of cardinality m , one has that for $1 \leq j \leq n$, $\mathbb{M}_{h,f^*}(j) = \frac{2}{m} \sum_{i \in I_j} \xi_i (f(X_i) - f^*(X_i))$ and

$$B_{f,f^*}(j) = \frac{1}{m} \sum_{i \in I_j} (f(X_i) - f^*(X_i))^2 + \frac{2}{m} \sum_{i \in I_j} \xi_i (f(X_i) - f^*(X_i)) = \mathbb{Q}_{f,f^*}(j) + \mathbb{M}_{f,f^*}(j) .$$

It follows that f^* defeats f in the j -th block if $\mathbb{Q}_{f,f^*}(j) + \mathbb{M}_{f,f^*}(j) > 0$, and, because $\tau < 1/4$, f^* wins the match if $\mathbb{Q}_{f,f^*}(j) \geq C \|f - f^*\|_{L_2}^2$ on more than $(1 - \tau)n$ of the blocks, while $\mathbb{M}_{f,f^*}(j) \leq -C \|f - f^*\|_{L_2}^2$ on at most τn of the blocks, .

This is summarized in the following lemma, established in the next two sections, and for the right choice of the constants $\kappa_1, \kappa_2, \kappa_3, \eta, \tau$, and θ that depend only on L and q .

Lemma 5.1. *There exists an absolute constant c and a constant $C_1 = C_1(L, q)$ for which the following holds. With probability at least $1 - 2 \exp(-c\tau^2 n)$, for every $f \in \mathcal{F}$ with $\|f - f^*\|_{L_2} \geq r$,*

$$\left| \left\{ j : \mathbb{Q}_{f,f^*}(j) \geq C_1 \|f - f^*\|_{L_2}^2 \right\} \right| \geq (1 - \tau) n$$

and

$$\left| \left\{ j : \mathbb{M}_{f,f^*}(j) \leq -\frac{3C_1}{4} \|f - f^*\|_{L_2}^2 \right\} \right| \leq \tau n .$$

The analysis takes place in the set

$$\mathcal{F}_1 = \text{star}(F, f^*) \cap (f^* + rS) ,$$

consisting of all the functions in the star-shaped hull of \mathcal{F} and f^* whose distance to f^* is precisely r . Once the estimates in Lemma 5.1 are verified for functions in \mathcal{F}_1 , extending them to the set $\{f \in \text{star}(\mathcal{F}, f^*) : \|f - f^*\|_{L_2} \geq r\}$, is straightforward, by invoking homogeneity properties of \mathbb{Q}_{f,f^*} and \mathbb{M}_{f,f^*} in $f - f^*$, and because $\text{star}(\mathcal{F}, f^*)$ is star-shaped around f^* .

The quadratic component

Here we establish the first part of Lemma 5.1, that states that, on an event of high probability, whenever $\|f - f^*\|_{L_2}$ is sufficiently large, a significant majority of the $(\mathbb{Q}_{f,f^*}(j))_{j=1}^n$ are at least a large fixed proportion of $\|f - f^*\|_{L_2}^2$. The size of the fixed proportion depends only on the small-ball property satisfied by the class \mathcal{F} , which, in turn, follows from the L_q - L_2 norm equivalence in $\text{span}(F)$. Indeed, if $\|f - f^*\|_{L_q} \leq L \|f - f^*\|_{L_2}$, then by the Paley-Zygmund inequality (see, e.g., [12]),

$$\mathbb{P}(|f - f^*|(X) \geq \kappa_0 \|f - f^*\|_{L_2}) \geq \rho_0 \tag{5.6}$$

for constants κ_0 and ρ_0 that depend only on L and q .

Recall that $(I_j)_{j=1}^n$ are the n blocks, each one of cardinality $m = N/n$, that $\mathcal{F}_1 = \{f \in \text{star}(\mathcal{F}, f^*) : \|f - f^*\|_{L_2} = r\}$, and that $m \geq 1/\tau$. For $t > 0$ and a function u , set

$$R_j(u, t) = |\{i \in I_j : |u(X_i)| \geq t\}| = \sum_{i \in I_j} \mathbb{1}_{\{|u(X_i)| \geq t\}} .$$

Thanks to (5.6), it is evident that if $f \in \mathcal{F}_1$ and $1 \leq j \leq n$ then, with probability at least $1 - 2\exp(-c_0\rho_0 m)$,

$$R_j(f - f^*, \kappa_0 r) \geq \frac{m\rho_0}{2} .$$

Recalling that $m \geq 1/\tau$, if $\tau \leq c_1(\rho_0)$ then

$$1 - 2\exp(-c_0\rho_0 m) \geq 1 - 2\exp(-c_0\rho_0/\tau) \geq 1 - \tau/12 ,$$

and $(\mathbb{1}_{\{R_j(f-f^*, \kappa_0 r) \geq m\rho_0/2\}})_{j=1}^n$ are i.i.d. Bernoulli random variables with mean at least $1 - \tau/12$. By standard concentration properties of Binomial distributions, with probability at least $1 - 2\exp(-c_2\tau^2 n)$,

$$\frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{R_j(f-f^*, \kappa_0 r) \geq m\rho_0/2\}} \geq 1 - \tau/6 . \quad (5.7)$$

In other words, at least $(1 - \tau/6)n$ of the blocks I_j have at least $m\rho_0/2$ coordinates of size at least $\kappa_0 r$. Moreover, in light of the high probability estimate with which (5.7) holds, by the union bound, the same property is satisfied uniformly by any subset of \mathcal{F}_1 of cardinality at most $\exp(c_2\tau^2 n/2)$. The set we consider is a maximal ηr -separated subset of \mathcal{F}_1 with respect to the $L_2(\mu)$ norm and for a well-chosen η specified later.

Let H_1 be such a maximal separated set. Since

$$\mathcal{F}_1 \subset f^* + (\text{star}(\mathcal{F} - f^*, 0) \cap rD) = f^* + \mathcal{F}_{f^*, r} ,$$

it follows from the translation invariance of packing numbers that

$$|H_1| \leq \mathcal{M}(\mathcal{F}_{f^*, r}, \eta r) .$$

Thus, to obtain a uniform control over points in H_1 , it suffices to show that

$$\log \mathcal{M}(\mathcal{F}_{f^*, r}, \eta r) \leq (c_2/2)\tau^2 n .$$

Recalling (5.4) and the choice of n , it suffices to verify that

$$\kappa_3^2 N \min\{1, \sigma^{-2} r^2\} = \kappa_3^2 n / \theta \leq (c_2/2)\tau^2 n ,$$

which is the case as long as

$$\kappa_3^2 \leq (c_2/2)\theta\tau^2 .$$

Observe that since H_1 is maximal, it is also an ηr -net in \mathcal{F}_1 , that is, every $f \in \mathcal{F}_1$ has some $\pi f \in H_1$ for which $\|f - \pi f\|_{L_2} \leq \eta r$. Consider the following event:

(A) (5.7) holds for every $f \in H_1$, and

$$(B) \sup_{f \in \mathcal{F}_1} (1/n) \sum_{i=1}^n \mathbb{1}_{\{R_j(f - \pi f, \kappa_0 r/2) \geq m\rho_0/4\}} \leq \frac{\tau}{12}.$$

On this event, for every $f \in \mathcal{F}_1$ there are at least $(1 - \tau/4)n$ blocks I_j with the following properties:

- $|\pi f(X_i) - f^*(X_i)| \geq \kappa_0 r$ on at least $m\rho_0/2$ coordinates in I_j , and
- $|f(X_i) - \pi f(X_i)| \geq \kappa_0 r/2$ on at most $m\rho_0/4$ coordinates in I_j .

Hence, in each one of the $(1 - \tau/4)n$ well-behaved blocks there are at least $m\rho_0/4$ coordinates X_i that satisfy

$$|f(X_i) - f^*(X_i)| \geq |\pi f(X_i) - f^*(X_i)| - |f(X_i) - \pi f(X_i)| \geq \kappa_0 r/2.$$

In particular, since $\|f - f^*\|_{L_2} = r$, one has

$$\mathbb{Q}_{f, f^*}(j) \geq (\rho_0 \kappa_0^2 / 16) \|f - f^*\|_{L_2}^2. \quad (5.8)$$

Moreover, (5.8) is positive homogeneous in $f - f^*$ and \mathcal{F}_1 is star-shaped around f^* , implying that (5.8) holds for every $f \in \mathcal{F}_1$ as long as $\|f - f^*\|_{L_2} \geq r$.

All that remains is to establish that (B) holds with a sufficiently high probability. To that end, let

$$\Psi(X_1, \dots, X_N) = \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{R_j(f - \pi f, \kappa_0 r/2) \geq m\rho_0/4\}}.$$

First note that, by the bounded differences inequality (see, e.g., [7]), with probability at least $1 - \exp(-c_3 u^2)$,

$$\Psi(X_1, \dots, X_N) \leq \mathbb{E}\Psi + \frac{u}{\sqrt{n}}.$$

Therefore, if $\mathbb{E}\Psi \leq \tau/24$ and $u \leq \sqrt{n}\tau/24$, then with probability at least $1 - \exp(-c_4 \tau^2 n)$, $\Psi(X_1, \dots, X_N) \leq \tau/12$ as required.

Thus, the final step in the proof is to show that

$$(*) = \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{R_j(f - \pi f, \kappa_0 r/2) \geq m\rho_0/4\}} \leq \frac{\tau}{24}.$$

Using that, for $\alpha > 0$, $\mathbb{1}_{\{|x| \geq \alpha\}} \leq \alpha^{-1}|x|$,

$$\mathbb{1}_{\{R_j(f - \pi f, \kappa_0 r/2) \geq m\rho_0/4\}} \leq \frac{4}{\rho_0 m} R_j(f - \pi f, \kappa_0 r/2) = \frac{4}{\rho_0 m} \sum_{i \in I_j} \mathbb{1}_{\{|f(X_i) - \pi f(X_i)| \geq \kappa_0 r/2\}}.$$

Thus, recalling that $nm = N$,

$$\begin{aligned}
(*) &\leq \frac{4}{\rho_0} \cdot \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{N} \sum_{i=1}^N \mathbb{1}_{\{|f - \pi f|(X_i) \geq \kappa_0 r/2\}} \\
&\leq \frac{8}{\rho_0 \kappa_0 r} \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{N} \sum_{i=1}^N |f(X_i) - \pi f(X_i)| \quad (\text{using } \mathbb{1}_{\{|x| \geq \alpha\}} \leq \alpha^{-1}|x| \text{ again}) \\
&\leq \frac{8}{\rho_0 \kappa_0 r} \left(\mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{N} \sum_{i=1}^N (|f(X_i) - \pi f(X_i)| - \mathbb{E}|f(X) - \pi f(X)|) + \sup_{f \in \mathcal{F}_1} \mathbb{E}|f(X) - \pi f(X)| \right) \\
&\leq \frac{16}{\rho_0 \kappa_0 r} \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{N} \sum_{i=1}^N \varepsilon_i (f(X_i) - \pi f(X_i)) \right| + \frac{8}{\rho_0 \kappa_0 r} \cdot \eta r \\
&\quad (\text{where } (\varepsilon_i)_{i=1}^N \text{ are independent symmetric random signs}).
\end{aligned}$$

In the last step we used a standard symmetrization argument, see, for example, [12]. To conclude the proof, observe that $(8/\rho_0 \kappa_0 r) \cdot \eta r \leq \tau/48$ when

$$\eta \leq c_5 \rho_0 \kappa_0 \tau \tag{5.9}$$

and for a suitable absolute constant c_5 . The fact that

$$\frac{16}{\rho_0 \kappa_0 r} \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{N} \sum_{i=1}^N \varepsilon_i (f - f^*)(X_i) \right| \leq \frac{\tau}{48} \tag{5.10}$$

follows because $r \geq 2r^*$ and invoking (5.2), as long as

$$\kappa_1 \leq c_6 \rho_0 \kappa_0 \tau$$

for an absolute constant c_6 . ■

The multiplier component

Next we complete the proof of Lemma 5.1 by showing that, with high probability, if $\|f - f^*\|_{L_2} \geq r$, then

$$\mathbb{M}_{f, f^*}(j) \leq -(3C_1/4) \|f - f^*\|_{L_2}^2$$

on at most τn blocks. In the first step towards this goal, we consider a single function:

Lemma 5.2. *There exists an absolute constant C_2 for which the following holds. Assume that for every $f \in \mathcal{F}$, $\mathbb{E}(f^*(X) - Y)(f(X) - f^*(X)) \geq 0$. If $f \in \mathcal{F}_1$, then with probability at least $1 - 2 \exp(-C_2 \tau^2 n)$,*

$$\left| \left\{ j : \mathbb{M}_{f, f^*}(j) \geq -\frac{C_1 r^2}{2} \right\} \right| \geq (1 - \tau/8)n .$$

In particular, the same assertion holds uniformly for any fixed subset $H_2 \subset \mathcal{F}_1$ of cardinality at most $\exp(C_2 \tau^2 n/2)$.

Before proving Lemma 5.2, consider the assumption that

$$\mathbb{E}(f^*(X) - Y)(f(X) - f^*(X)) \geq 0 \quad \text{for every } f \in \mathcal{F}. \quad (5.11)$$

It is straightforward to verify that (5.11) is satisfied under the assumptions of Theorems 2.10 and 2.11. Indeed, if \mathcal{F} is closed and convex then (5.11) is just the characterization of f^* as the nearest point to Y in \mathcal{F} in the L_2 sense. On the other hand, if $Y = f_0(X) + W$ for $f_0 \in \mathcal{F}$ and W that is mean-zero and independent of X , then $\mathbb{E}(f^*(X) - Y)(f(X) - f^*(X)) = -\mathbb{E}W(f(X) - f_0(X)) = 0$ for every $f \in \mathcal{F}$. In fact, (5.11) is the only structural assumption on the ‘location’ of Y relative to \mathcal{F} that is required for our analysis.

Another observation is that (5.11) passes to $\text{star}(\mathcal{F}, f^*)$, simply because any function in that set is of the form $h = \lambda f + (1 - \lambda)f^*$ for some $f \in \mathcal{F}$ and $0 \leq \lambda \leq 1$ and therefore, $(f^*(X) - Y)(h(X) - f^*(X)) = \lambda(f^*(X) - Y)(f(X) - f^*(X))$.

Proof of Lemma 5.2. Set $\xi = f^*(X) - Y$, put $U = \xi(f(X) - f^*(X))$ and observe that by (5.11), $\mathbb{E}U \geq 0$. Also note that for every j ,

$$\mathcal{M}_{f,f^*}(j) = \frac{1}{m} \sum_{i \in I_j} \xi_i(f(X_i) - f^*(X_i)) = \frac{1}{m} \sum_{i \in I_j} U_i,$$

for $(U_i)_{i=1}^M$ that are independent copies U . It is straightforward to verify that, with $(\varepsilon_i)_{i=1}^N$ defined as independent symmetric random signs,

$$\begin{aligned} \mathbb{P} \left(\left| \frac{1}{m} \sum_{i=1}^m U_i - \mathbb{E}U \right| \geq t \right) &\leq \frac{\mathbb{E} |\sum_{i=1}^m (U_i - \mathbb{E}U)|}{mt} \leq 2 \frac{\mathbb{E} |\sum_{i=1}^m \varepsilon_i U_i|}{mt} \\ &\leq 2 \frac{\|U\|_{L_2}}{\sqrt{mt}} = 2\sqrt{n} \frac{\|U\|_{L_2}}{\sqrt{Nt}} = (*). \end{aligned}$$

By the norm equivalence assumption of Theorem 2.10,

$$\|U\|_{L_2} \leq \|\xi\|_{L_4} \|f - f^*\|_{L_4} \leq L^2 \sigma r,$$

and by the independence assumption of Theorem 2.11,

$$\|U\|_{L_2} = \|\xi\|_{L_2} \|f - f^*\|_{L_2} = \sigma r.$$

Hence, setting $t = (C_1/2)r^2$ and noting that

$$\sqrt{\frac{n}{N}} \leq \sqrt{\theta} \frac{r}{\sigma},$$

it follows that

$$(*) \leq 2L^2 \frac{r\sigma\sqrt{n}}{t\sqrt{N}} = \frac{4L^2}{C_1} \frac{\sqrt{n}\sigma}{\sqrt{N}r} \leq \frac{\tau}{16}$$

whenever

$$\theta \leq (C_1\tau/64L^2)^2.$$

Therefore, with probability at least $1 - \tau/16$,

$$\frac{1}{m} \sum_{i=1}^m U_i \geq \mathbb{E}U - C_1 r^2/2 \geq -C_1 r^2/2.$$

Finally, consider the independent Bernoulli random variables $(\mathbb{1}_{\{\mathcal{M}_{f,f^*}(j) \leq -C_1 r^2/2\}})_{j=1}^n$ that have mean at most $\tau/16$. By concentration of Binomial random variables, there is an absolute constant C_2 such that, with probability at least $1 - 2\exp(-C_2\tau^2 n)$, there are at least $(1 - \tau/8)$ blocks that satisfy $\mathcal{M}_{f,f^*}(j) \geq -C_1 r^2/2$, as claimed. \blacksquare

Just as before, one may select any fixed subset $H_2 \subset \mathcal{F}_1$ of cardinality $\exp(C_2\tau^2n/2)$, and the assertion of Lemma 5.2 holds with high probability and uniformly for every $h \in H_2$. The choice of H_2 requires some care. It cannot be just an arbitrary maximal separated set.

Lemma 5.3. *There exists a subset $H_2 \subset \mathcal{F}_1$ of cardinality at most $\exp(C_2n\tau^2/2)$ such that for every $f \in \mathcal{F}_1$ there is some $h \in H_2$ that satisfies*

$$\|f - h\|_{L_2} \leq 2\eta r, \quad \text{and} \quad \mathbb{E}\xi(f(X) - h(X)) \geq 0 .$$

Proof. Let H' be a maximal ηr -separated subset of \mathcal{F}_1 . Recall that by the choice of r and (5.4), $\log |H'| \leq \kappa_3^2 N \min\{1, \sigma^{-2}r^2\}$, which is smaller than $(C_2/2)\tau^2n$ when

$$\kappa_3^2 \leq (C_2/2)\theta\tau^2 .$$

The class \mathcal{F} is a locally compact subset of $L_2(\mu)$ and therefore, so is \mathcal{F}_1 . Thus, for every $h' \in H'$, the intersections of \mathcal{F}_1 with the L_2 balls $B(h', \eta r)$ are compact and the continuous linear functional on L_2

$$f \rightarrow \mathbb{E}\xi f(X)$$

attains its minimum in each one of the sets $\mathcal{F}_1 \cap B(h', \eta r)$. Let h be such a minimizer and set H_2 to be the collection of these minimizers. Hence, for every $f \in \mathcal{F}_1$ there is some $h \in H_2$ for which $\|f - h\|_{L_2} \leq 2\eta r$ and $\mathbb{E}f(X)\xi \geq \mathbb{E}h(X)\xi$, as claimed. \blacksquare

With Lemma 5.3 at our disposal, for every $f \in \mathcal{F}_1$ define $\pi f \in H_2$ for which $\mathbb{E}\xi(f(X) - \pi f(X)) \geq 0$ and $\|f - \pi f\|_{L_2} \leq 2\eta r$, as above. By Lemma 5.2, with probability at least $1 - 2\exp(-C_2\tau^2n/2)$, for every $h \in H_2$ there are at least $(1 - \tau/8)n$ blocks I_j with

$$\mathbb{M}_{h, f^*}(j) = \frac{2}{m} \sum_{i \in I_j} \xi_i(h(X_i) - f^*(X_i)) \geq -C_1r^2/2 .$$

Hence, to complete the proof it suffices to show that for every $f \in \mathcal{F}_1$, there are at most $(7/8)n\tau$ blocks I_j with

$$\mathbb{M}_{f, f^*}(j) \leq \mathbb{M}_{\pi f, f^*}(j) - C_1r^2/4 .$$

Indeed, on that event, for every $f \in \mathcal{F}_1$ there are at most $n\tau$ blocks I_j that satisfy $\mathbb{M}_{f, f^*}(j) \leq -(3/4)C_1r^2$, as required.

Again, establishing this estimate for \mathcal{F}_1 yields that on the same event, if $\|f - f^*\|_{L_2}^2 \geq r^2$, then $\mathbb{M}_{f, f^*}(j) \leq -(3/4)C_1\|f - f^*\|_{L_2}^2$ on at most $n\tau$ coordinate blocks, since \mathbb{M} is homogeneous in $f - f^*$ and \mathcal{F}_1 is star-shaped around f^* .

Lemma 5.4. *There exists an absolute constant c for which, with probability at least $1 - 2\exp(-c\tau^2n)$,*

$$\sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{\mathbb{M}_{f, f^*}(j) - \mathbb{M}_{\pi f, f^*}(j) \leq -C_1r^2/4\}} \leq \frac{7\tau}{8} .$$

Proof. Recall that by the definition of πf , $\mathbb{E}\xi f(X) \geq \mathbb{E}\xi \pi f(X)$. Therefore,

$$\mathbb{E}(\mathbb{M}_{f, f^*} - \mathbb{M}_{\pi f, f^*}) = 2\mathbb{E}\xi(f(X) - \pi f(X)) \geq 0 .$$

To simplify notation, set

$$W_{f, \pi f}(j) = \mathbb{M}_{f, f^*}(j) - \mathbb{M}_{\pi f, f^*}(j) = \frac{1}{m} \sum_{i \in I_j} \xi_i(f(X_i) - \pi f(X_i)) ,$$

and note that $\mathbb{E}W_{f,\pi f}(j) \geq 0$.

Consider the supremum of Binomial random variables

$$\Psi = \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{W_{f,\pi f}(j) \leq -C_1 r^2/4\}}.$$

By the bounded differences inequality, with probability at least $1 - 2 \exp(-cu^2)$, $\Psi \leq \mathbb{E}\Psi + u/\sqrt{n}$. Setting $u = (6/8)\sqrt{n}\tau$, all that remains is to show that

$$\mathbb{E}\Psi = \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{W_{f,\pi f}(j) \leq -C_1 r^2/4\}} \leq \frac{\tau}{8}.$$

Since $\mathbb{E}W_{f,\pi f} \geq 0$, using $\mathbb{1}_{\{|x| \geq \alpha\}} \leq \alpha^{-1}|x|$,

$$\begin{aligned} \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{W_{f,\pi f}(j) \leq -C_1 r^2/4\}} &\leq \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{W_{f,\pi f}(j) - \mathbb{E}W_{f,\pi f}(j) \leq -C_1 r^2/4\}} \\ &\leq \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{|W_{f,\pi f}(j) - \mathbb{E}W_{f,\pi f}(j)| \geq C_1 r^2/4\}} \\ &\leq \frac{4}{C_1 r^2} \mathbb{E} \sup_{f \in \mathcal{F}_1} \frac{1}{n} \sum_{j=1}^n |W_{f,\pi f}(j) - \mathbb{E}W_{f,\pi f}(j)|. \end{aligned}$$

The next step is to centre the process. We show that for every $f \in \mathcal{F}_1$, the centring term satisfies

$$\frac{4}{C_1 r^2} \mathbb{E}|W_{f,\pi f}(j) - \mathbb{E}W_{f,\pi f}(j)| \leq \frac{\tau}{16}.$$

Indeed, symmetrizing, applying either the assumption of norm equivalence of Theorem 2.10 or the independence assumption of Theorem 2.11, and recalling that $\|f - \pi f\|_{L_2} \leq 2\eta r$, it is evident that

$$\begin{aligned} \frac{4}{C_1 r^2} \mathbb{E}|W_{f,\pi f} - \mathbb{E}W_{f,\pi f}| &\leq \frac{8}{C_1 r^2} \mathbb{E} \left| \frac{1}{m} \sum_{i=1}^m \varepsilon_i \xi_i (f(X_i) - \pi f(X_i)) \right| \\ &\leq \frac{16L^2}{C_1} \frac{\sigma \eta r}{r^2 \sqrt{m}} \\ &= \frac{16L^2}{C_1} \frac{\eta \sigma \sqrt{n}}{r \sqrt{N}} \leq \frac{16L^2}{C_1} \eta \sqrt{\theta}, \end{aligned}$$

again using the fact that $\sqrt{n/N} \leq \sqrt{\theta}(r/\sigma)$. Clearly,

$$\frac{16L^2}{C_1} \eta \sqrt{\theta} \leq \frac{\tau}{16}$$

when

$$\theta \leq (C_3 \tau / L^2 \eta)^2$$

for a suitable absolute constant C_3 .

Finally, we need to bound the centred empirical process. This is done by standard techniques of symmetrization, the contraction theorem for Bernoulli processes (for the Lipschitz function $\phi(t) = |t|$) and then de-symmetrization (see, e.g., [26]):

$$\begin{aligned}
& \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{n} \sum_{j=1}^n |W_{f, \pi f}(j) - \mathbb{E} W_{f, \pi f}| - \mathbb{E} |W_{f, \pi f}(j) - \mathbb{E} W_{f, \pi f}| \right| \\
& \leq 2 \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{n} \sum_{j=1}^n \varepsilon_j |W_{f, \pi f}(j) - \mathbb{E} W_{f, \pi f}| \right| \\
& \leq 2 \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{n} \sum_{j=1}^n \varepsilon_j (W_{f, \pi f}(j) - \mathbb{E} W_{f, \pi f}) \right| \\
& \leq 4 \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{n} \sum_{j=1}^n (W_{f, \pi f}(j) - \mathbb{E} W_{f, \pi f}) \right|.
\end{aligned}$$

Since

$$W_{f, \pi f}(j) = \frac{1}{m} \sum_{i \in I_j} \xi_i (f(X_i) - \pi f(X_i))$$

one has that

$$\begin{aligned}
& \frac{16}{C_1 r^2} \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{N} \sum_{i=1}^N \xi_i (f(X_i) - \pi f(X_i)) - \mathbb{E} \xi (f(X) - \pi f(X)) \right| \\
& \leq \frac{32}{C_1 r^2} \mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \sum_{i=1}^N \varepsilon_i \xi_i (f(X_i) - f^*(X_i)) \right| \leq \frac{\tau}{16}
\end{aligned}$$

provided that

$$\mathbb{E} \sup_{f \in \mathcal{F}_1} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \xi_i (f(X_i) - f^*(X_i)) \right| \leq C_4 \tau \sqrt{N} r^2, \quad (5.12)$$

that is, when

$$\kappa_2 \leq C_4 \tau,$$

concluding the proof. ■

5.2 Champions league–proof

Finally, it remains to prove Proposition 3.8. Recall that, with probability at least $1 - 2 \exp(-c_0 N \min\{1, \sigma^{-2} r^2\})$ with respect to the sample $(X_i, Y_i)_{i=1}^{2N}$, we have been able to identify a set of “qualifiers” that have not lost a single match in the preliminary round subset; that is, $H \subset \mathcal{F}$, consisting of f^* and possibly other functions that satisfy $\|f - f^*\|_{L_2} \leq (\beta/\alpha)r$. In the rest of this section we work conditionally on this “good” event.

While producing a function that is close to f^* solves the estimation problem, the question of prediction requires an additional step: we would like to choose one of the qualifiers that has an almost optimal statistical performance: a function \hat{f} that satisfies

$$\mathbb{E}((\hat{f}(X) - Y)^2 | (X_i, Y_i)_{i=2N+1}^{3N}) \leq \mathbb{E}(f^*(X) - Y)^2 + C r^2 \quad (5.13)$$

for an appropriate C . We show that this is possible with the required high probability for $C = 16(\beta/\alpha)^2$. To this end, set $r_1 = (\beta/\alpha)r$.

Recall that for $f, h \in \text{star}(\mathcal{F}, f^*)$, $\Psi_{h,f} = (h(X) - f(X))(f(X) - Y)$ and note that

$$\mathbb{M}_{f,f^*} = \frac{2}{m} \sum_{i=1}^m \Psi_{f,f^*}(X_i, Y_i) .$$

Also, as noted previously, since $\mathbb{E}\Psi_{f,f^*} \geq 0$ for every $f \in \mathcal{F}$, the same holds for any $f \in \text{star}(\mathcal{F}, f^*)$.

As was stated in Section 3.3, the sub-sample $(X_i, Y_i)_{i=2N+1}^{3N}$ is used to identify a function $\hat{f} \in \mathcal{F}$ for which $\mathbb{E}\Psi_{f^*,\hat{f}} \geq -2r_1^2$. By Lemma 3.6, this indeed suffices to establish (5.13) for $C = 16(\beta/\alpha)^2$.

The ‘‘champions league’’ round is designed to have ‘‘home-and-away’’ legs. For the partition $(I_j)_{j=1}^n$ of $(X_i, Y_i)_{i=2N+1}^{3N}$, f wins its home match against h if

$$\frac{2}{m} \sum_{i \in I_j} \Psi_{h,f}(X_i, Y_i) \geq -r_1^2/10$$

for more than half of the blocks I_j . We show that f^* wins all of its home matches, implying that the set of possible choices of \hat{f} is nonempty, and that if $\mathbb{E}\Psi_{f^*,f} \leq -2r_1^2$, then f loses its home match against f^* . On that event, a function that wins all of its home matches must satisfy that $\mathbb{E}\Psi_{f^*,\hat{f}} \geq -2r_1^2$, and this observation concludes the proof of Theorem 2.10.

The main ingredient in the proof is the following lemma.

Lemma 5.5. *Let $\mathcal{F}_2 = \text{star}(\mathcal{F}, f^*) \cap (f^* + r_1 D)$. Under the conditions of Theorem 2.10, there is an absolute constant c , such that, with probability at least $1 - 2 \exp(-cn)$,*

$$\sup_{f \in \mathcal{F}_2} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{|\mathbb{M}_{f,f^*}(j) - \mathbb{E}\mathbb{M}_{f,f^*}| \geq r_1^2/10\}} < \frac{1}{2} .$$

Proof. Using the assumption of norm equivalence of Theorem 2.10, one may verify that $\|\Psi_{f,f^*}\|_{L_2} \leq L^2 r_1 \sigma$. Therefore,

$$\mathbb{E} |\mathbb{M}_{f,f^*} - \mathbb{E}\mathbb{M}_{f,f^*}| \leq \frac{2}{\sqrt{m}} \|\Psi_{f,f^*}\|_{L_2} \leq \frac{2}{\sqrt{m}} \cdot L^2 r_1 \sigma . \quad (5.14)$$

Setting

$$Z = \sup_{f \in \mathcal{F}_2} \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{\{|\mathbb{M}_{f,f^*}(j) - \mathbb{E}\mathbb{M}_{f,f^*}| \geq r_1^2/10\}},$$

it follows that $\mathbb{E}Z$ is at most

$$\begin{aligned} \mathbb{E} \sup_{f \in \mathcal{F}_2} \frac{10}{nr_1^2} \sum_{j=1}^n (|\mathbb{M}_{f,f^*}(j) - \mathbb{E}\mathbb{M}_{f,f^*}| - \mathbb{E}|\mathbb{M}_{f,f^*}(j) - \mathbb{E}\mathbb{M}_{f,f^*}|) \\ + \frac{10}{nr_1^2} \cdot \sup_{f \in \mathcal{F}_2} \mathbb{E}|\mathbb{M}_{f,f^*}(j) - \mathbb{E}\mathbb{M}_{f,f^*}| = (I) + (II) . \end{aligned}$$

Applying the same argument used in the previous section—namely, symmetrization, followed by contraction for a Bernoulli process and the Lipschitz function $\phi(t) = |t|$ and de-symmetrization—one has that for absolute constants c_1, c_2 , and c_3 ,

$$(I) \leq \frac{c_1}{r_1^2} \cdot \mathbb{E} \sup_{f \in \mathcal{F}_2} \left| \frac{1}{n} \sum_{j=1}^n \mathbb{M}_{f, f^*}(j) - \mathbb{E} \mathbb{M}_{f, f^*}(j) \right| \leq \frac{c_2}{r_1^2} \cdot \mathbb{E} \sup_{f \in \mathcal{F}_2} \left| \frac{1}{N} \sum_{i=1}^N \varepsilon_i \xi_i (f - f^*)(X_i) \right| \leq \frac{1}{8}$$

when

$$\kappa_2 \leq c_3 .$$

Also, by (5.14),

$$(II) \leq \frac{10L^2\sigma}{r_1} \cdot \sqrt{\frac{n}{N}} \leq \frac{1}{8} ,$$

provided that $n \leq c_4 L^4 \cdot (r_1^2/\sigma^2)N$ for an absolute constant c_4 . Since $r_1 = 2(\beta/\alpha)r$, it suffices that

$$\theta \leq c_4 L^4 .$$

Thus, $\mathbb{E}Z \leq 1/4$, and by the bounded differences inequality applied to Z , one has that with probability at least $1 - 2\exp(-cu^2)$, $Z \leq \mathbb{E}Z + u/\sqrt{n}$. The claim follows by selecting $u = \sqrt{n}/8$. \blacksquare

Recall that for every $f \in H$, $\|f - f^*\|_{L_2} \leq (\beta/\alpha)r$ and consider the “good” event from Lemma 5.5. For any $f \in \mathcal{F}_2$, and, in particular, for any qualifier $f \in H$,

$$\mathbb{E} \mathbb{M}_{f, f^*} - r_1^2/10 \leq \mathbb{M}_{f, f^*}(j) \leq \mathbb{E} \mathbb{M}_{f, f^*} + r_1^2/10 \quad (5.15)$$

on more than $n/2$ blocks. Moreover, if $\|f - f^*\|_{L_2} \leq r_1$ and $\mathbb{E} \Psi_{f^*, f} \leq -2r_1^2$, then $\mathbb{E} \Psi_{f, f^*} \geq r_1^2$. Indeed,

$$\begin{aligned} -2r_1^2 &\geq \mathbb{E}(f^*(X) - f(X)) \cdot (f(X) - Y) \\ &= -\|f - f^*\|_{L_2}^2 + \mathbb{E}(f^*(X) - f(X)) \cdot (f^*(X) - Y) \\ &\geq -r^2 + \mathbb{E}(f^*(X) - f(X)) \cdot (f^*(X) - Y) = -r_1^2 - \mathbb{E} \Psi_{f, f^*} . \end{aligned}$$

Therefore, $\mathbb{E} \mathbb{M}_{f, f^*} \geq 2r_1^2$, and on the event (5.15),

$$\mathbb{M}_{f, f^*}(j) \geq \mathbb{E} \mathbb{M}_{f, f^*} - r_1^2/10 \geq r_1^2 .$$

Finally, since $\Psi_{f^*, f} = -(f^*(X) - f(X))^2 - \Psi_{f, f^*}$, on that event and the same coordinate blocks,

$$\frac{1}{m} \sum_{i \in I_j} \Psi_{f^*, f}(j) = -\frac{2}{m} \sum_{i=1}^m (f^*(X_i) - f(X_i))^2 - \mathbb{M}_{f, f^*} \leq -r_1^2 .$$

Thus, f is defeated by f^* in the majority of the blocks, and in particular, loses its home match against f^* . It follows that, with probability at least $1 - 2\exp(-cn)$, any function \widehat{f} selected in the champions league round must satisfy that $\Psi_{f^*, \widehat{f}} \geq -2r_1^2$, and by Lemma 3.6,

$$\mathbb{E}((\widehat{f}(X) - Y)^2 | (X_i, Y_i)_{2N+1}^{3N}) \leq \mathbb{E}(f^*(X) - Y) + 4r_1^2 .$$

\blacksquare

Additional remarks

It should be noted that the difference between $r^*(f^*)$ and r^* is not a major issue in most interesting cases. $\lambda_{\mathbb{Q}}(\kappa, \eta, h)$, $\lambda_{\mathbb{M}}(\kappa, \eta, h)$, and $r_E(\kappa, h)$ typically do not depend heavily on h and little is lost by taking the supremum over $h \in \mathcal{F}$. $\tilde{r}_{\mathbb{M}}(c_1, \sigma)$ becomes a better estimate of $\bar{r}_{\mathbb{M}}(c_1, f^*)$ as σ gets closer to $\|f^* - Y\|_{L_2}$. Even if this value is not known in advance, it is easy to design a two-stage procedure that constructs a data-dependent estimate of $\|f^* - Y\|_{L_2}$ and then uses the procedure of Theorem 2.10 with a tight upper bound for the value of σ obtained from the first stage. In particular, it is easy to find a value of σ that is within a constant factor of the optimum. This is an issue of secondary importance and we omit the straightforward details.

The major problem that remains open is the identity of the accuracy edge. To date, there is a single generic example in which one may attain an accuracy smaller than Cr^* and in that case the accuracy attained is proportional to λ^* and with the optimal confidence at that level, namely, $1 - 2 \exp(-cN \min\{1, \sigma^{-2} \lambda_{\mathbb{M}}^2(\kappa_2/\sigma, \eta_2)\})$.

This fact has recently been established in [30] in a very special situation: when \mathcal{F} is a convex, L -sub-Gaussian class of functions; *all* the admissible targets are of the form $Y = f_0(X) + W$ for $f_0 \in \mathcal{F}$ and W that is sub-Gaussian, zero-mean, independent of X . The procedure used is a modification of ERM: one replaces \mathcal{F} with an appropriate net, thus ‘erasing’ all the fine structure of \mathcal{F} at the right level, and then runs ERM on the net. The idea behind this procedure is straightforward: if one is interested in accuracy r , one is insensitive to perturbations of that order. From that perspective, a net with a mesh that is proportional to r is just as good as the entire class. It turns out that r^* of the net is proportional to λ^* the original class \mathcal{F} . Unfortunately, all the highly restrictive assumptions are essential to the proof and cannot be relaxed at all.

It should be noted that the median-of-means tournament may be modified in exactly the same way as ERM is modified in [30], leading to an accuracy that is proportional to λ^* when \mathcal{F} is a convex, L -sub-Gaussian class and for an independent noise W that may be heavy-tailed. We decided not pursue this point further because it is a very special case, and shifts the emphasis of the article from the question of the optimal accuracy/confidence tradeoff to the nature of the accuracy edge, a problem of a different nature.

As far as the latter is concerned, it is not clear whether the gap between λ^* and r^* can be closed in other cases. It is possible that both independent noise and a sub-Gaussian class are essential in attaining an accuracy proportional to λ^* , and under weaker assumptions, the true accuracy edge lies somewhere between λ^* and r^* . We leave that question for future study.

Finally, it should be noted that in this work we completely ignore the algorithmic aspects of the new procedure. While computing the empirical risk minimizer often leads to thoroughly studied and well understood convex optimization problems, finding the winner of the median-of-means tournament in a computationally efficient manner is a highly nontrivial—and perhaps not hopeless—problem that goes beyond the scope of this paper. Techniques of “derivative-free optimization” with “function comparison oracle” may be useful, see, for example, [20].

References

- [1] N. Alon, Y. Matias, and M. Szegedy. The space complexity of approximating the frequency moments. *Journal of Computer and System Sciences*, 58:137–147, 2002.

- [2] M. Anthony and P. L. Bartlett. *Neural network learning: theoretical foundations*. Cambridge University Press, Cambridge, 1999.
- [3] J.-Y. Audibert and O. Catoni. Robust linear least squares regression. *The Annals of Statistics*, 39:2766–2794, 2011.
- [4] F. Barthe, O. Guédon, S. Mendelson, and A. Naor. A probabilistic approach to the geometry of the l_p^n -ball. *Ann. Probab.*, 33(2):480–513, 2005.
- [5] F. Barthe and A. Koldobsky. Extremal slabs in the cube and the Laplace transform. *Adv. Math.*, 174(1):89–114, 2003.
- [6] P.L. Bartlett, O. Bousquet, and S. Mendelson. Localized Rademacher complexities. *Annals of Statistics*, 33:1497–1537, 2005.
- [7] S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.
- [8] C. Brownlees, E. Joly, and G. Lugosi. Empirical risk minimization for heavy-tailed losses. *Annals of Statistics*, 43:2507–2536, 2015.
- [9] S. Bubeck, N. Cesa-Bianchi, and G. Lugosi. Badits with heavy tail. *IEEE Transactions on Information Theory*, 59:7711–7717, 2013.
- [10] P. Bühlmann and S. van de Geer. *Statistics for high-dimensional data*. Springer Series in Statistics. Springer, Heidelberg, 2011. Methods, theory and applications.
- [11] O. Catoni. Challenging the empirical mean and empirical variance: a deviation study. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 48(4):1148–1185, 2012.
- [12] V.H. de la Peña and E. Giné. *Decoupling: from Dependence to Independence*. Springer, New York, 1999.
- [13] L. Devroye, L. Györfi, and G. Lugosi. *A probabilistic theory of pattern recognition*, volume 31 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1996.
- [14] L. Devroye, M. Lerasle, G. Lugosi, and R.I. Oliveira. Sub-gaussian mean estimators. *Annals of Statistics*, 2016.
- [15] R. M. Dudley. *Uniform central limit theorems*, volume 142 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, New York, second edition, 2014.
- [16] S. Foucart and H. Rauhut. *A mathematical introduction to compressive sensing*. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, New York, 2013.
- [17] L. Györfi, M. Kohler, A. Krzyżak, and H Walk. *A Distribution-Free Theory of Nonparametric Regression*. Springer, New York, 2002.
- [18] D. Hsu and S. Sabato. Approximate loss minimization with heavy tails. *Computing Research Repository*, abs/1307.1827, 2013.

- [19] D. Hsu and S. Sabato. Loss minimization and parameter estimation with heavy tails. *Journal of Machine Learning Research*, 17:1–40, 2016.
- [20] K. Jamieson, R. Nowak, and B. Recht. Query complexity of derivative-free optimization. In *Advances in Neural Information Processing Systems*, pages 2672–2680, 2012.
- [21] M. Jerrum, L. Valiant, and V. Vazirani. Random generation of combinatorial structures from a uniform distribution. *Theoretical Computer Science*, 43:186–188, 1986.
- [22] V. Koltchinskii. *Oracle inequalities in empirical risk minimization and sparse recovery problems*, volume 2033 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2011. Lectures from the 38th Probability Summer School held in Saint-Flour, 2008, École d’Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School].
- [23] V. Koltchinskii. *Oracle inequalities in empirical risk minimization and sparse recovery problems: Lecture notes*. Springer, 2011.
- [24] G. Lecué and S. Mendelson. Learning subgaussian classes: Upper and minimax bounds. In S. Boucheron and N. Vayatis, editors, *Topics in Learning Theory*. Societe Mathematique de France, 2016.
- [25] G. Lecué and S. Mendelson. Sparse recovery under weak moment assumptions. *Journal of the European Mathematical Society*, to appear.
- [26] M. Ledoux and M. Talagrand. *Probability in Banach Space*. Springer-Verlag, New York, 1991.
- [27] M. Lerasle and R.I. Oliveira. Robust empirical mean estimators. *manuscript*, 2012.
- [28] P. Massart. *Concentration inequalities and model selection*. Ecole d’été de Probabilités de Saint-Flour 2003. Lecture Notes in Mathematics. Springer, 2006.
- [29] S. Mendelson. Learning without concentration. *Journal of the ACM*, 62:21, 2015.
- [30] S. Mendelson. Local vs. global parameters breaking the Gaussian complexity barrier. *manuscript*, 2015.
- [31] S. Mendelson. On aggregation for heavy-tailed classes. *Probability Theory and Related Fields*, to appear.
- [32] S. Mendelson. On multiplier processes under weak moment assumptions. *Geometric Aspects of Functional Analysis - GAFA Seminar notes*, to appear.
- [33] S. Mendelson. Upper bounds on product and multiplier empirical processes. *Stochastic Processes and their Applications*, to appear.
- [34] S. Minsker. Geometric median and robust estimation in Banach spaces. *Bernoulli*, 21:23082335, 2015.
- [35] A.S. Nemirovsky and D.B. Yudin. Problem complexity and method efficiency in optimization. 1983.

- [36] V.N. Sudakov. Gaussian measures, Cauchy measures and ϵ -entropy. In *Soviet Math. Dokl*, volume 10, pages 310–313, 1969.
- [37] M. Talagrand. *Upper and lower bounds for stochastic processes*, volume 60 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Heidelberg, 2014. Modern methods and classical problems.
- [38] A. B. Tsybakov. *Introduction to nonparametric estimation*. Springer Series in Statistics. Springer, New York, 2009. Revised and extended from the 2004 French original, Translated by Vladimir Zaiats.
- [39] S. van de Geer. *Applications of empirical process theory*, volume 6 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2000.
- [40] S. van de Geer. *Estimation and Testing Under Sparsity*. Springer International Publishing, 2016.
- [41] A.W. van der Vaart and J.A. Wellner. *Weak convergence and empirical processes*. Springer-Verlag, New York, 1996.
- [42] V.N. Vapnik and A.Ya. Chervonenkis. *Theory of Pattern Recognition*. Nauka, Moscow, 1974. (in Russian); German translation: *Theorie der Zeichenerkennung*, Akademie Verlag, Berlin, 1979.

A The distance oracle – outline of the proof

Here, we sketch the proof of Proposition 3.2. The proof is almost identical to that of Theorem 3.3 from [31], and follows the same path as the study of the quadratic component in the proof of Theorem 2.10.

The first step in the proof involves generating a small-ball estimate that holds with sufficiently high probability, say 9/10. If a random variable Z satisfies $\|Z\|_{L_q} \leq L\|Z\|_{L_2}$ for some $q > 2$, then Z automatically satisfies a small-ball estimate, but with constants ρ_0 and κ_0 that depend on L and q . In particular, it need not be true that ρ_0 is close to 1, for example, that $\mathbb{P}(|Z| \geq \kappa_0\|Z\|_{L_2}) \geq 0.9$. However, a combination of the norm equivalence and a Berry-Esseen type argument suffices to ensure that an average of a small number of independent copies of Z satisfies such a bound. More accurately, if ℓ is an integer that depends only on q and L and Z_1, \dots, Z_ℓ are ℓ independent copies of Z , then

$$\mathbb{P}\left(\frac{1}{\ell} \sum_{i=1}^{\ell} |Z_i| \geq \kappa_0\|Z\|_{L_2}\right) \geq 0.9 . \quad (\text{A.1})$$

Moreover, combining (A.1) with a straightforward application of Chebyshev's inequality, one has that with probability at least 0.8,

$$\kappa_0\|Z\|_{L_2} \leq \frac{1}{\ell} \sum_{i=1}^{\ell} |Z_i| \leq \kappa_1\|Z\|_{L_2}$$

for constants κ_0 and κ_1 that depend only on q and L .

Now consider a partition of $\{1, \dots, N\}$ to k blocks I'_1, \dots, I'_k , where each block is of cardinality ℓ . Let $\mathcal{M}_{Z,j} = \frac{1}{\ell} \sum_{i \in I'_j} |Z_i|$. It follows from a standard binomial estimate that, with probability at least $1 - 2\exp(-c_1k)$, at least $0.7k$ of the random variables $\mathcal{M}_{Z,j}$ satisfy

$$\kappa_0\|Z\|_{L_2} \leq \mathcal{M}_{Z,j} \leq \kappa_1\|Z\|_{L_2} .$$

Let us apply this observation to our setup: fix $f^* \in \mathcal{F}$ and consider the set

$$\text{star}(\mathcal{F} - f^*, 0) \cap rS \subset \text{star}(\mathcal{F} - f^*, 0) \cap rD = \mathcal{F}_{f^*, r}$$

for any fixed $r > d^*$. By the choice of d^* , $\text{star}(\mathcal{F} - f^*, 0) \cap rS$ contains an ηr -net \mathcal{V}_r of cardinality at most $\exp(c_1k/2)$. Therefore, with probability at least $1 - 2\exp(-c_1k/2)$, for every $v \in \mathcal{V}_r$ there is $J_v \subset \{1, \dots, k\}$ of cardinality at least $0.7k$, such that for every $j \in J_v$,

$$\kappa_0 r \leq \mathcal{M}_{v,j} \leq \kappa_1 r . \quad (\text{A.2})$$

Next, for each $v \in \text{star}(\mathcal{F} - f^*, 0) \cap rS$ let $\pi v \in \mathcal{V}_r$ satisfy $\|v - \pi v\|_{L_2} \leq \eta r$. Suppose that one can show that on a high probability event, for every such v there are at most $k/10$ blocks I'_j on which

$$\mathcal{M}_{v-\pi v, j} \geq \kappa_0 r / 2 .$$

Then, on that event, and on at least $0.6k$ blocks,

$$(\kappa_0/2)r \leq \mathcal{M}_{v,j} \leq 2\kappa_1 r ,$$

and in particular, the same holds for the median of means $\text{Med}_\ell(v)$.

This is precisely the isomorphic estimate we require, so far, for elements of $\text{star}(\mathcal{F} - f^*, 0)$ whose $L_2(\mu)$ norm is r . Obtaining the isomorphic estimate for elements with a larger $L_2(\mu)$ norm follows because $\text{star}(\mathcal{F} - f^*, 0)$ is star-shaped around 0 and the required isomorphic estimate is positive homogeneous.

Therefore, to complete the proof of the first part of Proposition 3.2 it suffices to show that

$$\sup_v |\{j : \mathcal{M}_{v-\pi v, j} \geq \kappa_0 r/2\}| \leq k/10, \quad (\text{A.3})$$

where the supremum is taken in $\text{star}(\mathcal{F} - f^*, 0) \cap rS$. The proof of (A.3) is based on an identical argument to the one we used earlier, in the study of the quadratic component, and the fact that $r > d^*$.

The second claim in Proposition 3.2 is that on a high probability event, a one-sided (upper) estimate on $\text{Med}_\ell(v)$ should hold for any $v \in \text{star}(\mathcal{F} - f^*, 0) \cap rD$. Its proof is almost the same as the one we have just described.

For each $v \in \mathcal{V}'_r$, with probability at least 0.8,

$$\mathcal{M}_{v, j} \leq \kappa_1 \|v\|_{L_2} \leq \kappa_1 r, \quad (\text{A.4})$$

and thus, with probability at least $1 - 2 \exp(-c_1 k)$, at least $0.7k$ of the random variables $\mathcal{M}_{v, j}$ are smaller than $\kappa_1 r$. Next, we consider an ηr -net $\mathcal{V}'_r \subset \mathcal{F}_{f^*, r}$, which, by the choice of r is of cardinality at most $\exp(c_1 k/2)$. It follows that, with probability at least $1 - 2 \exp(-c_1 k/2)$, (A.4) holds for every $v \in \mathcal{V}'_r$. The oscillation term is then controlled as we outlined above.

Finally, because $k = c(q, L)N$, the estimates hold with the claimed probability of $1 - 2 \exp(-cN)$ for a constant c that depends only on q and L . ■