The continuum Pólya-like random walk Daniel Krenn^{*}, Hosam Mahmoud, and Mark Daniel Ward[†]

Abstract

The Pólya urn scheme is a discrete-time process concerning the addition and removal of colored balls. There is a known embedding of it in continuous-time called the Pólya process. We deal with a generalization of this stochastic model, where the initial values and the entries of the transition matrix (corresponding to additions or removals) are not necessarily fixed integer values as in the standard Pólya process. In one of the scenarios, we even allow the entries of the matrix to be random variables. As a result, we no longer have a combinatorial model of "balls in an urn," but a broader interpretation as a random walk in a possibly high number of dimensions. In this paper, we study several parametric classes of these generalized continuum Pólya-like random walks.

AMS classification: 60F05, 60G99.

Keywords and phrases: Urn model, Pólya process, random walk, stochastic process, partial differential equation.

1 Introduction

We deal with a generalization of the Pólya process on c colors by embedding each element of the replacement matrix into \mathbb{R} , or by even treating the elements of the replacement matrix as random variables. We are no longer restricted to thinking about integer-valued quantities (such as *counts* of balls in an urn), we can now handle real-valued quantities, such as a random walk with possibly fractional step sizes, in c dimensions, rather than restricting to integer step sizes in each of the c dimensions. We can view the replacement matrix itself as random. The generalized model is mentioned in [9, 13].

^{*}Daniel Krenn is supported by the Austrian Science Fund (FWF): P 24644-N26.

[†]Mark Daniel Ward is supported by NSF Grant DMS-1246818, and by the NSF Science & Technology Center for Science of Information Grant CCF-0939370.

1.1 Background

We first review the standard Pólya process. The Pólya urn scheme is a process underlying an urn that evolves in discrete time. The urn contains balls of up to c colors. The colors are numbered, say the set of colors is $C = \{1, \ldots, c\}$. At each discrete epoch in time, a ball is sampled from the urn. It is then put back in the urn, together with a number of other balls in various colors. If the sampled ball has color $i \in C$, then we add to the urn $A_{i,j}$ balls of color j, for $j \in C$. If $A_{i,j}$ is negative, we remove $A_{i,j}$ balls of color j. It is customary to represent these dynamics by a replacement matrix

$$\mathbf{A} = \begin{pmatrix} A_{1,1} & A_{1,2} & \dots & A_{1,c} \\ A_{2,1} & A_{2,2} & \dots & A_{2,c} \\ \vdots & \vdots & \ddots & \vdots \\ A_{c,1} & A_{c,2} & \dots & A_{c,c} \end{pmatrix}.$$

In the standard Pólya process each element of \mathbf{A} is an integer. (We will deal with an extended view that considers \mathbf{A} as having real-valued entries in Section 1.2.) It is usually assumed that the urn is "tenable" in the sense that the selection of balls can be continued ad infinitum, no matter which stochastic path is followed, i.e., the process will never get stuck.

The Pólya process is an embedding of the Pólya urn scheme in real time. Embedding in real time (poissonization) was suggested by Kac [11] as a general methodology for understanding discrete probability problems. In the context of urns, poissonization was introduced by Athreya and Karlin [1] to understand discrete-time Pólya urn schemes. Poissonization was thus meant as a transform. The inverse transform (depoissonization)—to translate results in the continuous domain back to results in the discrete domain—is fraught with difficulty [1]. For general background on depoissonization, we refer the reader to [8] (Chapter 8), or [16] (Chapter 10).

Some authors developed interest in the continuous-time Pólya process for its own sake (see [2, 3, 4, 15]). In the Pólya process, each ball carries an internal clock that rings in Exp(1) time (a random amount of time, according to an exponential random variable with mean 1), independently of the behavior of all other clocks. Whenever a clock rings, it is instantaneously reset to ring again in Exp(1) time (independently of the clocks on all the other balls). In other words, each ball has the ability to generate a new Poisson process with intensity 1. When the clock associated with a ball of color *i* rings, the addition and removal of balls corresponds to picking a colored ball from a Pólya urn and using the *i*th row of the replacement matrix **A** to determine which balls to add or remove. All replacements are assumed to occur instantaneously, and each new ball is given an independent clock that rings in Exp(1) time.

1.2 Generalized Pólya process

In this paper, we study a generalized Pólya process that can be viewed as a second layer of embedding of the Pólya urn scheme. The starting numbers and the (possibly random) numbers added are in \mathbb{R} (no longer restricted to \mathbb{Z}), so the numbers are no longer counts of balls. A natural (more general) interpretation is a random walk in c dimensions. At time t, the current position of the walk in \mathbb{R}^c is a c-dimensional column vector $\mathbf{X}(t) := (X_1(t), \ldots, X_c(t))^{\top}$.

At any point t in time, the next renewal occurs after a random amount of time, according to a master clock. The waiting time of this master clock follows an exponential random variable with mean $1/\sum_{i\in C} X_i(t)$. Given that a renewal occurs, the probability that the renewal corresponds to color i is $X_i(t)/\sum_{j\in C} X_j(t)$. In other words, the probability of a renewal of type i is proportional to the amount of quantity i present when the transition occurs. In such a case, the *i*th row of the matrix **A** dictates the *c*-dimensional direction in which to move, i.e., we add $A_{i,j}$ units to $X_j(t)$, for each $j \in C$. Thus, it is appropriate then to call **A** the *navigation matrix*, instead of the replacement matrix.

To avoid trivialities, we only consider starting values $X_j(0)$ and matrices $A_{i,j}$ in which the walk is tenable, i.e., the walk always avoids the origin, and each of the coordinates is always nonnegative. We call the row vector $\mathbf{X}^{\mathsf{T}}(t) = (X_1(t), \ldots, X_c(t))$ a continuum Pólya-like process or random walk.

This stochastic process is not a Poisson process, because the rate of the process itself is random, i.e., the rate of replacement is not simply a function of time. The current rate of replacement depends on the number of replacements that have taken place beforehand.

1.3 Organization of the paper

The paper is organized as follows. Section 2 specifies the probability model for the stochastic process. In Section 3, we derive a fundamental partial differential equation that governs the behavior of the generalized Pólya process. In Section 4, we derive a functional equation for the moment generating function of the position of the random walk at time t. Section 5 is perhaps the most novel part of the paper, because we solve the partial differential equations for a balanced upper-triangular case, a case that proved difficult in the usual urn setting. Additionally, in Appendix A, we show how several classical probability models can be generalized with this approach. In all of these classical cases, the partial differential equations from Section 4 can be solved.

2 The probabilistic model

In order to be able to establish the partial differential equations of Section 3, we need to precisely describe the number of renewals, say $N(t, \Delta t)$, that occur in the processes during the interval $(t, t + \Delta t]$. We use the notation C_i to indicate the event that exactly one renewal occurs in the interval $(t, t + \Delta t]$ and that renewal is induced by the *i*th coordinate. To study the behavior of $\mathbf{X}(t+s)$, for s > 0, we condition on the vector $\mathbf{X}(t)$.

Lemma 2.1. The conditional probabilities of either zero renewals, one renewal of type *i*, or two or more renewals, in the interval $(t, t + \Delta t]$, given the value of $\mathbf{X}(t)$, are (respectively) the following:

$$\mathbb{P}\big(N(t,\Delta t) = 0 \mid \mathbf{X}(t)\big) = \exp\Big(-\Delta t \sum_{j \in C} X_j(t)\Big),$$
$$\mathbb{P}\big((N(t,\Delta t) = 1) \cap \mathcal{C}_i \mid \mathbf{X}(t)\big) = \Delta t X_i(t) \exp\Big(-\Delta t \sum_{j \in C} X_j(t)\Big) + O\Big((\Delta t)^2\Big),$$
$$\mathbb{P}\big(N(t,\Delta t) \ge 2 \mid \mathbf{X}(t)\Big) = O\Big((\Delta t)^2\Big),$$

as $\Delta t \to 0$.

Proof. Given the values $\mathbf{X}(t)$, the master clock does not ring during the interval $(t, t + \Delta t]$ with probability

$$\mathbb{P}\Big(N(t,\Delta t) = 0 \mid \mathbf{X}(t)\Big) = \prod_{j \in C} \left(\frac{(\Delta t)^0 e^{-\Delta t}}{0!}\right)^{X_j(t)} = \exp\left(-\Delta t \sum_{j \in C} X_j(t)\right).$$

For fixed t, once we are given the values $\mathbf{X}(t)$, the next ring of the master clock, after time t, occurs at a random time x (with t < x) with probability density function

$$\left(\sum_{j\in C} X_j(t)\right) \exp\left(-(x-t)\sum_{j\in C} X_j(t)\right).$$

When such a clock ring occurs at time x, it is a ring of type i with probability $X_i(t) / \sum_{j \in C} X_j(t)$. Then, for $t < x \leq t + \Delta t$, there are no additional subsequent

rings before time $t + \Delta t$ with probability

$$\prod_{j \in C} \left(\frac{(t + \Delta t - x)^0 e^{-(t + \Delta t - x)}}{0!} \right)^{X_j(t) + A_{i,j}} = \exp\left(-(t + \Delta t - x) \sum_{j \in C} \left(X_j(t) + A_{i,j}\right)\right).$$

Putting all of this together (and again given $\mathbf{X}(t)$), the conditional probability $\mathbb{P}((N(t, \Delta t) = 1) \cap \mathcal{C}_i \mid \mathbf{X}(t))$ that the master clock rings exactly once during the interval $(t, t + \Delta t]$ and yields a renewal of type *i*, is equal to

$$\int_{t}^{t+\Delta t} \left(\sum_{j\in C} X_{j}(t)\right) \exp\left(-(x-t)\sum_{j\in C} X_{j}(t)\right) \left(\frac{X_{i}(t)}{\sum_{j\in C} X_{j}(t)}\right) \\ \times \exp\left(-(t+\Delta t-x)\sum_{j\in C} \left(X_{j}(t)+A_{i,j}\right)\right) \mathrm{d}x.$$

This simplifies to

$$\int_{t}^{t+\Delta t} X_{i}(t) \exp\left(-\Delta t \sum_{j \in C} X_{j}(t) - (t+\Delta t-x) \sum_{j \in C} A_{i,j}\right) dx$$
$$= \frac{X_{i}(t) \exp\left(-\Delta t \sum_{j \in C} X_{j}(t)\right) \left(\Delta t \sum_{j \in C} A_{i,j} + O((\Delta t)^{2})\right)}{\sum_{j \in C} A_{i,j}}$$
$$= \Delta t X_{i}(t) \exp\left(-\Delta t \sum_{j \in C} X_{j}(t)\right) + O\left((\Delta t)^{2}\right).$$

Finally, the conditional probability (given $\mathbf{X}(t)$) that the master clock rings two or more times during the interval $(t, t + \Delta t]$ is

$$\mathbb{P}(N(t,\Delta t) \ge 2 \mid \mathbf{X}(t)) = 1 - \mathbb{P}(N(t,\Delta t) = 0 \mid \mathbf{X}(t)) - \sum_{i \in C} \mathbb{P}((N(t,\Delta t) = 1) \cap \mathcal{C}_i \mid \mathbf{X}(t)) = 1 - \exp\left(-\Delta t \sum_{j \in C} X_j(t)\right) - \sum_{i \in C} \Delta t X_i(t) \exp\left(-\Delta t \sum_{j \in C} X_j(t)\right) + O\left((\Delta t)^2\right) = O\left((\Delta t)^2\right);$$

we arrive at the latter conclusion after a local expansion of the two exponential functions. This completes the proof of Lemma 2.1. $\hfill \Box$

3 The fundamental partial differential equation

We formulate here a partial differential equation for the continuum Pólya-like random walk. We use the vector $\mathbf{u} = (u_j)_{j \in C}$ to mark the colors $C = \{1, \ldots, c\}$. Let

$$\phi(t, \mathbf{u}) = \mathbb{E}\left[\exp\left(\sum_{j \in C} u_j X_j(t)\right)\right]$$

be the joint moment generating function of the coordinates of the random walk $\mathbf{X}(t)$. For $i \in C$, let

$$\psi_i(\mathbf{u}) = \mathbb{E}\left[\exp\left(\sum_{j \in C} u_j A_{i,j}\right)\right]$$

be the joint moment generating function of the random variables on row i of the navigation matrix **A**.

Theorem 3.1. The joint moment generating function $\phi(t, \mathbf{u})$ satisfies

$$\frac{\partial \phi}{\partial t} + \sum_{i \in C} (1 - \psi_i) \frac{\partial \phi}{\partial u_i} = 0.$$

Proof. We use conditional expectation to calculate

$$E := \mathbb{E}\left[\exp\left(\sum_{j \in C} u_j X_j(t + \Delta t)\right) \, \middle| \, \mathbf{X}(t)\right],\$$

i.e., the expectation conditioned on the status $\mathbf{X}(t)$ at time t. We do this by first conditioning on whether there are zero, one or at least two renewals in the interval $(t, t + \Delta t]$. If there is exactly one renewal, we also condition on the color of the chosen direction. The probabilities of these events are calculated in Lemma 2.1. As in Section 2, let $N(t, \Delta t)$ denote the number of renewals that occur in the processes in the interval $(t, t + \Delta t]$.

We obtain

$$E = \mathbb{E}\left[\exp\left(\sum_{j\in C} u_j X_j(t+\Delta t)\right) \middle| \mathbf{X}(t) \text{ and } N(t,\Delta t) = 0\right] \times \mathbb{P}\left(N(t,\Delta t) = 0\right)$$

+ $\sum_{i\in C} \mathbb{E}\left[\exp\left(\sum_{j\in C} u_j X_j(t+\Delta t)\right) \middle| \mathbf{X}(t) \text{ and } (N(t,\Delta t) = 1) \cap \mathcal{C}_i\right]$
 $\times \mathbb{P}\left(N(t,\Delta t) = 1, \mathcal{C}_i\right)$
+ $\mathbb{E}\left[\exp\left(\sum_{j\in C} u_j X_j(t+\Delta t)\right) \middle| \mathbf{X}(t) \text{ and } N(t,\Delta t) \ge 2\right] \times \mathbb{P}\left(N(t,\Delta t) \ge 2\right)$

In the last equality, we have allowed for the possibility that the $A_{i,j}$ themselves (i.e., the entries of the navigation matrix) may be random variables, allowing for more generality. We only assume that these matrix entries are independent of the current state of the process $\mathbf{X}(t)$.

Collecting all these facts and utilizing Lemma 2.1, we obtain

$$E = \exp\left(\sum_{j \in C} u_j X_j(t)\right) \exp\left(-\Delta t \sum_{i \in C} X_i(t)\right)$$

+
$$\sum_{i \in C} \mathbb{E}\left[\exp\left(\sum_{j \in C} u_j A_{i,j}\right)\right] \exp\left(\sum_{j \in C} u_j X_j(t)\right) \Delta t X_i(t)$$

×
$$\exp\left(-\Delta t \sum_{j \in C} X_j(t)\right) + O\left((\Delta t)^2\right).$$

A local expansion of the exponentials gives

$$E = \exp\left(\sum_{j \in C} u_j X_j(t)\right) \left(1 - \Delta t \sum_{i \in C} X_i(t) + O\left((\Delta t)^2\right)\right) + \Delta t \left(1 + O(\Delta t)\right) \exp\left(\sum_{j \in C} u_j X_j(t)\right) \sum_{i \in C} X_i(t) \mathbb{E}\left[\exp\left(\sum_{j \in C} u_j A_{i,j}\right)\right] + O\left((\Delta t)^2\right).$$

Taking expectations over $\mathbf{X}(t)$ yields

$$\phi(t + \Delta t, \mathbf{u}) = \phi(t, \mathbf{u}) - \Delta t \sum_{i \in C} \mathbb{E} \Big[X_i(t) \exp \Big(\sum_{j \in C} u_j X_j(t) \Big) \Big] \\ + \Delta t \Big(1 + O(\Delta t) \Big) \mathbb{E} \Big[\exp \Big(\sum_{j \in C} u_j X_j(t) \Big) \sum_{i \in C} X_i(t) \Big] \psi_i(\mathbf{u}) \\ + O\Big((\Delta t)^2 \Big) \,.$$

We can now write the limiting form

$$\frac{\partial \phi(t, \mathbf{u})}{\partial t} = \lim_{\Delta t \to 0} \frac{\phi(t + \Delta t, \mathbf{u}) - \phi(t, \mathbf{u})}{\Delta t}$$
$$= -\sum_{i \in C} \mathbb{E} \Big[X_i(t) \exp \Big(\sum_{j \in C} u_j X_j(t) \Big) \Big] \Big(1 - \psi_i(\mathbf{u}) \Big)$$
$$= -\sum_{i \in C} \frac{\partial \phi(t, \mathbf{u})}{\partial u_i} \Big(1 - \psi_i(\mathbf{u}) \Big).$$

Remark 3.2. The proof of Theorem 3.1 is a generalization of the proof in Balaji and Mahmoud [2]. However, it needed some new techniques. In [2], there is a conditional argument that uses a sum on the number of balls of a color, given that number. Of course, the number of balls is an integer and such a sum can be carried out. Here, the counterpart of a number of balls of a certain color is a continuous coordinate, and such a conditional sum cannot be written.

4 Moments

We derive functional equations for the moments. For example, we derive a functional equation for the mean position (see Theorem 4.1 at the end of this section) by differentiating (with respect to u_j) on both sides of the partial differential equation in Theorem 3.1. This yields

$$\frac{\partial}{\partial u_j} \left(\frac{\partial \phi}{\partial t} \right) + \sum_{i \in C} \left(1 - \psi_i \right) \frac{\partial^2 \phi}{\partial u_j \, \partial u_i} - \sum_{i \in C} \frac{\partial \psi_i}{\partial u_j} \times \frac{\partial \phi}{\partial u_i} = 0.$$

Evaluation of the summands at $u_i = 0$, for all $i \in C$, yields

$$\frac{\partial}{\partial u_j} \left(\frac{\partial \phi}{\partial t} \right) \Big|_{\mathbf{u}=0} = \frac{\partial}{\partial t} \left(\frac{\partial \phi}{\partial u_j} \right) \Big|_{\mathbf{u}=0}$$
$$= \frac{\partial}{\partial t} \mathbb{E} \Big[X_j(t) \exp \left(\sum_{i \in C} u_i X_i(t) \right) \Big] \Big|_{\mathbf{u}=0}$$
$$= \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E} [X_j(t)] \,.$$

We also have

$$\sum_{i \in C} \left(1 - \psi_i \right) \frac{\partial^2 \phi}{\partial u_j \, \partial u_i} \Big|_{\mathbf{u} = 0} = 0,$$

because $(1 - \psi_i)\Big|_{\mathbf{u}=0} = 0$, and

$$\sum_{i \in C} \frac{\partial \psi_i}{\partial u_j} \times \frac{\partial \phi}{\partial u_i} \Big|_{\mathbf{u}=0} = \sum_{i \in C} \mathbb{E}[A_{i,j}] \mathbb{E}[X_i(t)].$$

Hence, for the jth coordinate, we get the ordinary differential equation

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}[X_j(t)] = \sum_{i \in C} \mathbb{E}[A_{i,j}] \mathbb{E}[X_i(t)].$$

Putting the differential equations (for $j \in C$) together in matrix form, we get the functional equation

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E} \Big[\mathbf{X}(t) \Big] = \mathbb{E} \Big[\mathbf{A}^{\top} \Big] \mathbb{E} \Big[\mathbf{X}(t) \Big],$$

where \mathbf{A}^{\top} is the transpose of the navigation matrix \mathbf{A} . This first-order functional equation has a standard solution; this yields the following theorem.

Theorem 4.1. Let \mathbf{A} be the navigation matrix of a continuum Pólya-like random walk. At time t, the expected value of the coordinates of the walk are

$$\mathbb{E}[\mathbf{X}(t)] = e^{\mathbb{E}[\mathbf{A}^+]t} \mathbf{X}(0).$$

Note that, for a matrix \mathbf{M} , we have used the notation $e^{\mathbf{M}} = \sum_{n=0}^{\infty} \mathbf{M}^n / n!$, and we can compute this by using the Jordan form of \mathbf{M} .

Remark 4.2. The result of Theorem 4.1 can be obtained via an alternative martingale approach. It is discussed in [9] that $e^{-t\mathbb{E}[A^{\top}]}\mathbf{X}(t)$ is martingale. It follows that $\mathbb{E}[e^{-t\mathbb{E}[A^{\top}]}\mathbf{X}(t)] = \mathbf{X}(0)$, or in other words, $\mathbb{E}[\mathbf{X}(t)] = e^{t\mathbb{E}[A^{\top}]}\mathbf{X}(0)$. However, it is harder to obtain higher moments by this technique. On the other hand, martingale convergence theorems can give us (nonconstructively) almost sure limits, without specifying what the limits are. The PDE method being discussed here works hand in hand with that alternative line to give us the distributions.

5 Walks according to a balanced triangular scheme

In this section, we discuss a walk according to a balanced triangular scheme. The case has some historical significance. Pólya urn models came about in the first decades of the 20th century. Perhaps the first urn schemes are Ehrenfest model, constructed to understand the diffusion of gas [6], and Pólya-Eggenberger urn scheme, a model for contagion [5].

Soon thereafter, a theory was developed for many types of urns. However, the triangular flavor remained defiant until very recently. The triangular case has been handled in [10] and limit distributions have been characterized. Alternative characterizations are given in [7, 12, 18].

So, let us consider a walk $\mathbf{X}(t) = (X(t), Y(t))^{\top}$ following a balanced triangular scheme with the navigation matrix

$$\mathbf{A} = \begin{pmatrix} \alpha & \delta - \alpha \\ 0 & \delta \end{pmatrix},$$

where $\alpha < \delta$ are numbers in \mathbb{R}^+ . We are excluding the case $\alpha = \delta$; see Appendix A.1, where this Pólya–Eggenberger like random walk has already been handled.

Theorem 5.1. Suppose we have a balanced triangular scheme as described above. Then, the moment generating function is

$$\phi(t; u, v) = \mathbb{E}\left[e^{uX(t) + vY(t)}\right]$$

= $e^{-X(0)t}(e^{-\alpha u} - e^{-\alpha v} + (e^{-\delta v} - 1 + e^{-\delta t})^{\alpha/\delta})^{-X(0)/\alpha}$
 $\times e^{-Y(0)t}(e^{-\delta v} - 1 + e^{-\delta t})^{-Y(0)/\delta}.$

From this result we can compute the moments in a straightforward way. See Section 5.5 for details.

To prove Theorem 5.1, we solve the corresponding partial differential equation of Theorem 3.1 by using the method of characteristics. To improve readability, the proof is split up into several sections.

5.1 Characteristic curves

We derive the characteristic curves belonging to the partial differential equation by establishing the following lemma.

Lemma 5.2. The functions

$$x_c = e^{\alpha t}(e^{-\alpha u} - e^{-\alpha v})$$
 and $y_c = e^{\delta t}(e^{-\delta v} - 1)$

are characteristic curves for the partial differential equation that corresponds to the balanced triangular scheme specified above.

To prove this lemma, consider the more general matrix

$$\mathbf{A} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}.$$

We want to find a solution $\phi(t; u, v)$ of the partial differential equation

$$\frac{\partial \phi}{\partial t} + (1 - e^{\alpha u} e^{\beta v}) \frac{\partial \phi}{\partial u} + (1 - e^{\gamma u} e^{\delta v}) \frac{\partial \phi}{\partial v} = 0.$$

Parameterizing

As a first step in the method of characteristics, we introduce a new parameter s. We set

$$\frac{\mathrm{d}t}{\mathrm{d}s} = 1,\tag{5.1t}$$

$$\frac{\mathrm{d}u}{\mathrm{d}s} = 1 - e^{\alpha u} e^{\beta v},\tag{5.1u}$$

and

$$\frac{\mathrm{d}v}{\mathrm{d}s} = 1 - e^{\gamma u} e^{\delta v}.\tag{5.1v}$$

By using the chain rule and inserting (5.1), we obtain

$$\frac{\mathrm{d}}{\mathrm{d}s}\phi(t(s);u(s),v(s)) = 0,$$

so our function is constant along the characteristics.

From (5.1t) it follows that $s = t + t_0$. We choose $t_0 = 0$, thus s = t, which we use from now on.

Specialization to the upper triangular case

Since we are interested in the upper triangular case, we now specialize to $\beta = \delta - \alpha$ and $\gamma = 0$. Thus, (5.1v) becomes

$$\frac{\mathrm{d}v}{\mathrm{d}t} = 1 - e^{\delta v},$$

and we can easily solve it by the separation of variables. This gives

$$v - \frac{1}{\delta}\log(1 - e^{\delta v}) = t - y_0,$$

for some (constant) initial condition y_0 . This is equivalent to

$$e^{\delta(t-y_0)} = \frac{e^{\delta v}}{1-e^{\delta v}} = \frac{1}{e^{-\delta v}-1},$$
(5.2a)

$$e^{\delta y_0} = e^{\delta(t-v)}(1-e^{\delta v}) = e^{\delta t}(e^{-\delta v}-1),$$
 (5.2b)

and to

$$e^{\delta v} = \frac{1}{e^{-\delta(t-y_0)} + 1}$$
(5.2c)

as well. In particular, $y_c = e^{\delta y_0}$ (together with Equation (5.2b)) is the second of our two characteristic curves of Lemma 5.2.

It remains to derive the first characteristic curve. Inserting (5.2c) into (5.1u) yields the differential equation

$$\frac{\mathrm{d}u}{\mathrm{d}t} = 1 - \frac{e^{\alpha u}}{\left(e^{-\delta(t-y_0)} + 1\right)^{\beta/\delta}}$$

With the help of a computational symbolic algebra system, we can solve this. We obtain

$$u(t) = t - \frac{1}{\alpha} \log \left(\alpha x_0 + \frac{\alpha}{\alpha + \beta} e^{\alpha t} (e^{\delta(t - y_0)})^{\beta/\delta} H \right),$$

where H denotes the hypergeometric function

$$H = {}_{2}F_{1}\left(\frac{\beta}{\delta}, \frac{\alpha+\beta}{\delta}; \frac{\alpha+\beta}{\delta}+1; -e^{\delta(t-y_{0})}\right),$$

and for some (constant) initial condition x_0 .

Solving for this x_0 gives

$$x_{0} = \frac{1}{\alpha} e^{\alpha(t-u)} - \frac{1}{\alpha+\beta} e^{\alpha t} (e^{\delta(t-y_{0})})^{\beta/\delta} H.$$
 (5.3)

Balancing the triangular scheme

At this point we consider the balanced triangular case and specialize to $\beta = \delta - \alpha$. The characteristic curve (5.3) is now

$$x_{0} = \frac{1}{\alpha} e^{\alpha(t-u)} - \frac{1}{\delta} e^{\alpha t} \left(-Z\right)^{1-\alpha/\delta} {}_{2}F_{1}\left(1 - \frac{\alpha}{\delta}, 1; 2; Z\right)$$

with $Z = -e^{\delta(t-y_0)} = 1/(1-e^{-\delta v})$, cf. (5.2a).

We can simplify the hypergeometric function, namely,

$$_{2}F_{1}(\mu, 1; 2; Z) = \frac{1 - (1 - Z)^{1 - \mu}}{(1 - \mu)Z}.$$

By using $1 - \mu = \alpha/\delta$, since $(Z - 1)/Z = e^{-\delta v}$, we obtain

$$x_0 = \frac{1}{\alpha} e^{\alpha(t-u)} + \frac{1}{\alpha} e^{\alpha t} \frac{1 - (1-Z)^{\alpha/\delta}}{(-Z)^{\alpha/\delta}} = \frac{e^{\alpha t}}{\alpha} (e^{-\alpha u} - e^{-\alpha v} + (e^{-\delta v} - 1)^{\alpha/\delta}).$$

Setting

$$x_c = \alpha x_0 - e^{\alpha y_0} = e^{\alpha t} (e^{-\alpha u} - e^{-\alpha v})$$

(we also used (5.2b)) completes the proof of Lemma 5.2.

5.2 The general solution

As we have both characteristic curves x_c and y_c now (cf. Lemma 5.2), we can write down the general solution to our partial differential equation. For some function $\tilde{\phi}$, we have

$$\phi(t; u, v) = \widetilde{\phi}(e^{\alpha t}(e^{-\alpha u} - e^{-\alpha v}), e^{\delta t}(e^{-\delta v} - 1)).$$
(5.4)

To determine $\tilde{\phi}$, we need to take the initial conditions into account.

5.3 Initial conditions

We have two possible cases to consider for the initial conditions of the solution (5.4). The case covered in this section is u = v. Regardless of the type of transition we have, we are adding δ balls to the urn, each time a transition takes place. In the other case (see Section B.2), we set u = 0, so that we are only considering transitions with the second direction of navigation.

Let $\tau(0) := X(0) + Y(0)$. The time between the ℓ th and $(\ell+1)$ st transition is exponential, with parameter $\tau(0) + \ell \delta$, i.e., with expected time $1/(\tau(0) + \ell \delta)$. When the $(\ell+1)$ st transition occurs (for any $\tau(0) \ge 0$), we walk a total of δ units in the sense of Manhattan taxicab geometry. So, the walk is $\tau(0) + \ell \delta$ Manhattan blocks away from the origin, after ℓ transitions have taken place.

Lemma 5.3. If u = v, then the probability generating function of Y(t) is

$$\mathbb{E}\left[e^{vY(t)}\right] = (1 - e^{\delta t} + e^{\delta(t-v)})^{-(X(0)+Y(0))/\delta}$$

Note that

$$\mathbb{E}\left[e^{vY(t)}\right] = (1+y_c)^{-(X(0)+Y(0))/\delta},$$

where y_c is the characteristic curve of Lemma 5.2.

Lemma 5.3 can be proved by setting up and solving Kolmogorov's Forward Equations; see Appendix B for details.

5.4 Solution to the partial differential equation

We are now ready to determine the function $\tilde{\phi}$ of the general solution (5.4), and thus, we will prove Theorem 5.1.

Proof of Theorem 5.1. We use the initial conditions provided by Lemma 5.3.We compute

$$\begin{split} \phi(t; u, v) &= \mathbb{E}\Big[e^{uX(t)+vY(t)}\Big] \\ &= (x_c + (y_c + 1)^{\alpha/\delta})^{-X(0)/\alpha}(y_c + 1)^{-Y(0)/\delta} \\ &= (e^{\alpha t}(e^{-\alpha u} - e^{-\alpha v}) + (e^{\delta t}(e^{-\delta v} - 1) + 1)^{\alpha/\delta})^{-X(0)/\alpha} \\ &\times (e^{\delta t}(e^{-\delta v} - 1) + 1)^{-Y(0)/\delta} \\ &= e^{-X(0)t}(e^{-\alpha u} - e^{-\alpha v} + (e^{-\delta v} - 1 + e^{-\delta t})^{\alpha/\delta})^{-X(0)/\alpha} \\ &\times e^{-Y(0)t}(e^{-\delta v} - 1 + e^{-\delta t})^{-Y(0)/\delta}. \end{split}$$

This solution satisfies our initial conditions, and our proof is complete. $\hfill \Box$

5.5 Moments

Theorem 5.1 provides us with the moment generating function. The moments now follow by differentiation and evaluating at u = 0 and v = 0. The first moments are

$$\mathbb{E}[X(t)] = X(0)e^{\alpha t},$$

$$\mathbb{E}[Y(t)] = (X(0) + Y(0))e^{\delta t} - X(0)e^{\alpha t}.$$

Note that the two coordinates of the walk grow at different rates, and the drift is much stronger in the vertical direction.

We can also compute the second moments,

$$\begin{split} \mathbb{E}[X^{2}(t)] &= X(0)(\alpha + X(0))e^{2\alpha t} - \alpha X(0)e^{\alpha t},\\ \mathbb{E}[X(t)Y(t)] &= X(0)(\alpha + X(0) + Y(0))e^{(\alpha + \delta)t} - X(0)(\alpha + X(0))e^{2\alpha t},\\ \mathbb{E}[Y^{2}(t)] &= (X(0) + Y(0))(\delta + X(0) + Y(0))e^{2\delta t} \\ &\quad - 2X(0)(\alpha + X(0) + Y(0))e^{(\alpha + \delta)t} \\ &\quad + \delta(X(0) + Y(0))e^{\delta t} + X(0)(\alpha + X(0))e^{2\alpha t} + \alpha X(0)e^{\alpha t}, \end{split}$$

and consequently

$$\begin{aligned} \mathbb{V}\mathrm{ar}(X(t)) &= \alpha X(0)e^{2\alpha t} - \alpha X(0)e^{\alpha t},\\ \mathbb{C}\mathrm{ov}(X(t),Y(t)) &= \alpha X(0)e^{(\alpha+\delta)t} - \alpha X(0)e^{2\alpha t},\\ \mathbb{V}\mathrm{ar}(Y(t)) &= \delta(X(0) + Y(0))e^{2\delta t} - 2\alpha X(0)e^{(\alpha+\delta)t} \\ &- \delta(X(0) + Y(0))e^{\delta t} + \alpha X(0)e^{2\alpha t} + \alpha X(0)e^{\alpha t}. \end{aligned}$$

Remark 5.4. Of course, the first moments follow by Theorem 4.1 as well: We have (

$$\mathbb{E}[\mathbf{X}(t)] = e^{\begin{pmatrix} \alpha & 0\\ \delta - \alpha & \delta \end{pmatrix}^t} \mathbf{X}(0).$$

The matrix in the exponent is diagonal, which makes it an easy computation and we obtain

$$\begin{pmatrix} \mathbb{E}[X(t)] \\ \mathbb{E}[Y(t)] \end{pmatrix} = \begin{pmatrix} e^{\alpha t} & 0 \\ e^{\delta t} - e^{\alpha t} & e^{\delta t} \end{pmatrix} \begin{pmatrix} X(0) \\ Y(0) \end{pmatrix},$$

as derived by differentiation.

5.6 Asymptotic distributions

With the exact moment generating function at hand, we can determine marginal as well as joint distributions of the number of (suitably scaled) white and blue balls.

Let us start with the displacement in the first coordinate. Set v = 0, and evaluate X(t) with the scale $e^{\alpha t}$. We see that

$$\mathbb{E}\left[\exp\left(\frac{X(t)}{e^{\alpha t}}u\right)\right] = \phi\left(t;\frac{u}{e^{\alpha t}},0\right) = \frac{e^{-\tau t}e^{bt}}{(e^{-\alpha u e^{-\alpha t}} - 1 + e^{-\alpha t})^{X(0)/\alpha}}$$

As $t \to \infty$, we find

$$\mathbb{E}\left[\exp\left(\frac{X(t)}{e^{\alpha t}}u\right)\right] = \frac{e^{-\tau t}e^{bt}}{\left(\left(1 - \alpha u e^{-\alpha t} + O(e^{-2\alpha t})\right) - 1 + e^{-\alpha t}\right)^{X(0)/\alpha}}$$
$$\sim \frac{e^{-\tau t}e^{Y(0)t}}{\left(e^{-\alpha t} - \alpha u e^{-\alpha t}\right)^{X(0)/\alpha}}$$
$$\sim \frac{e^{-\tau t}e^{Y(0)t}}{e^{-X(0)t}(1 - \alpha u)^{X(0)/\alpha}},$$

where "~" stands for asymptotic equivalence. Recalling that $\tau(0) = X(0) + Y(0)$, we see a cancellation leading to the convergence

$$\mathbb{E}\left[\exp\left(\frac{X(t)}{e^{\alpha t}}u\right)\right] \to \frac{1}{(1-\alpha u)^{X(0)/\alpha}}.$$

The right-hand side is the moment generating function of a $\text{Gamma}(X(0)/\alpha, \alpha)$ random variable. By Lévy's Continuity Theorem [17] (Theorem 18.1), we have

$$\frac{X(t)}{e^{\alpha t}} \to \operatorname{Gamma}\left(\frac{X(0)}{\alpha}, \alpha\right).$$

By a similar analysis, putting u = 0 and using the scale $e^{\delta t}$ for the displacement along the second axis, we find

$$\frac{Y(t)}{e^{\alpha t}} \to \operatorname{Gamma}\left(\frac{\tau(0)}{\delta}, \delta\right).$$

Joint distributions with the two marginals being gamma distributions are called bivariate gamma distributions. From the variances and covariance computed in Subsection 5.5, we find the asymptotic correlation between $X(t)e^{-\alpha t}$ and $Y(t)e^{-\delta t}$ to be $\sqrt{(\alpha X(0))/(\delta \tau(0))}$, for $\alpha < \delta$. This extends a calculation in [4], under the scenario of a triangular Pólya urn scheme. For the case $\alpha = \delta$, we have 0 correlation, as naturally arises from the independence of the movement along the two axes.

References

- Krishna B. Athreya and Samuel Karlin, Embedding of urn schemes into continuous time Markov branching processes and related limit theorems, Annals of Mathematical Statistics 39 (1968), 1801–1817.
- [2] Srinivasan Balaji and Hosam M. Mahmoud, Exact and limiting distributions in diagonal Pólya processes, Annals of the Institute of Statistical Mathematics 58 (2006), no. 1, 171–185.
- [3] Srinivasan Balaji, Hosam M. Mahmoud, and Osamu Watanabe, Distributions in the Ehrenfest process, Statistics & Probability Letters 76 (2006), no. 7, 666–674.
- [4] Chen Chen and Hosam Mahmoud, The continuous-time triangular Pólya process, 2018, pp. 303–321.
- [5] Florian Eggenberger and George Pólya, über die statistik verketteter vorgänge, Z. Angew. Math. Mech. 3 (1923), 279–290 (German).
- [6] Paul Ehrenfest and Tatjana Ehrenfest, über zwei bekannte einwände gegen das boltzmannsche h-theorem, Phys. Z. 8 (1907), 311–314 (German).
- [7] Philippe Flajolet, Philippe Dumas, and Vincent Puyhaubert, Some exactly solvable models of urn process theory, Fourth Colloquium on Mathematics and Computer Science (Philippe Chassaing, ed.), DMTCS Proceedings, vol. AG, 2006, pp. 59–118.
- [8] Micha Hofri and Hosam Mahmoud, Algorithmics of nonuniformity, Discrete Mathematics and its Applications (Boca Raton), CRC Press, Boca Raton, FL, 2019, Tools and paradigms.
- [9] Svante Janson, Functional limit theorems for multitype branching processes and generalized Pólya urns, Stochastic Process. Appl. 110 (2004), no. 2, 177–245.
- [10] _____, Limit theorems for triangular urn schemes, Probability Theory and Related Fields 134 (2006), 417–452.

- [11] Mark Kac, On deviations between theoretical and empirical distributions, Proceedings of the National Academy of Sciences of the United States of America 35 (1949), 252–257.
- [12] Markus Kuba and Hosam M. Mahmoud, Two-color balanced affine urn models with multiple drawings, Adv. in Appl. Math. 90 (2017), 1–26.
- [13] Nicolas Pouyanne, An algebraic approach to Pólya processes, Ann. Inst. Henri Poincaré Probab. Stat. 44 (2008), no. 2, 293–323.
- [14] Sheldon M. Ross, *Stochastic Processes*, Wiley series in probability and statistics, Wiley, 1996.
- [15] Joshua Sparks and Hosam M. Mahmoud, Phases in the two-color tenable zero-balanced Pólya process, Statistics & Probability Letters 83 (2013), no. 1, 265–271.
- [16] Wojciech Szpankowski, Average case analysis of algorithms on sequences, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2001, With a foreword by Philippe Flajolet.
- [17] David Williams, Probability with martingales, Cambridge Mathematical Textbooks, Cambridge University Press, Cambridge, 1991.
- [18] Panpan Zhang, Chen Chen, and Hosam Mahmoud, Explicit characterization of moments of balanced triangular Pólya urns by an elementary approach, Statistics & Probability Letters 96 (2015), 149–153.

Daniel Krenn Department of Mathematics Alpen-Adria-Universität Klagenfurt Universitätsstraße 65–67 9020 Klagenfurt am Wörthersee, Austria math@danielkrenn.at or daniel.krenn@aau.at

Hosam M. Mahmoud Department of Statistics The George Washington University Washington, D.C. 20052, USA hosam@gwu.edu

Mark Daniel Ward Department of Statistics Purdue University West Lafayette, IN 47907, USA mdw@purdue.edu

A Illustrative examples

We give here some examples. Some of them resemble and extend standard Pólya processes to the random walk counterpart. Some have no solved equivalent in the Pólya world (neither the discrete- or continuous-time versions).

A.1 Pólya–Eggenberger-like random walk

Suppose the navigation matrix \mathbf{A} is in the diagonal matrix form

$$\begin{pmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_c \end{pmatrix},$$

where A_i is a nonnegative random variable. Let us focus on the *i*th component; we set $u = u_i$ and $u_j = 0$ for $j \neq i$. Let $\psi_i(u) = \mathbb{E}[e^{uA_i}]$ be the moment generating function of A_i , and let $\phi_i(t, u) = \mathbb{E}[e^{uX_i(t)}]$ be the moment generating function of $X_i(t)$. The partial differential equation simplifies to

$$\frac{\partial \phi_i(t,u)}{\partial t} + (1 - \psi_i(u))\frac{\partial \phi_i(t,u)}{\partial u} = 0.$$
(A.1)

This equation can be solved for several standard distributions of the A_i . Two examples are discussed below.

Component-wise almost surely constant

Suppose $A_i = \alpha_i \in \mathbb{R}^+$ almost surely. This gives rise to

$$\phi_i(t, u) = (1 - e^{\alpha_i t} (1 - e^{-\alpha_i u}))^{-X_i(0)/\alpha_i}$$

We set $u = se^{-\alpha_i t}$ and take the limit, to obtain

$$\lim_{t \to \infty} \mathbb{E}\left[\exp\left(\frac{sX_i(t)}{e^{\alpha_i t}}\right)\right] = \lim_{t \to \infty} \phi_i\left(t, \frac{s}{e^{\alpha_i t}}\right) = (1 - \alpha_i s)^{-X_i(0)/\alpha_i}.$$

The latter moment generating function is that of a $\text{Gamma}(X_i(0)/\alpha_i, \alpha_i)$ random variable. That is, we have

$$\frac{X_i(t)}{e^{\alpha_i t}} \xrightarrow{\mathcal{D}} \text{Gamma}\left(\frac{X_i(0)}{\alpha_i}, \alpha_i\right).$$

Note that the displacements along the *i*th coordinate affect only changes in that direction. In other words, the limit multivariate distribution has independent marginals, with the *i*th marginal having the latter gamma limit distribution. Also observe that this random walk has a very long memory. It never forgets where it starts. Even the limit is influenced by the initial position vector, which comes in as a parameter in the joint limit distribution.

Component-wise exponential distribution

We illustrate with another instance, in which the navigation matrix itself has random elements. Suppose the Pólya–Eggenberger-like random walk operates under exponentially distributed displacements. That is, the A_i (for $i \in C$) are independent Exp(1) random variables. Thus, for u < 1, we have $\psi_i(u) = 1/(1-u)$ and we can solve (A.1). In this case, we have

$$\phi_i(t,u) = e^{-W(-ue^{t-u})},$$

where $W(\cdot)$ is Lambert's W function (defined implicitly as any complex solution of $z = W(z) e^{W(z)}$). Note that T(z) := -W(-z) is called the *tree function* and appears in the enumeration of trees.

Set $u = se^{-t}$, and take the limit

$$\lim_{t \to \infty} \mathbb{E}\left[\exp\left(\frac{sX_i(t)}{e^t}\right)\right] = \lim_{t \to \infty} \phi_i\left(t, \frac{s}{e^t}\right) = \frac{T(s)}{s}$$

The right-hand side in the latter equation is the moment generating function of a Lambert random variable W^* . Thus, in the limit we have

$$e^{-t} \begin{pmatrix} X_1(t) \\ X_2(t) \\ \vdots \\ X_c(t) \end{pmatrix} \xrightarrow{\mathcal{D}} \begin{pmatrix} W_1^* \\ W_2^* \\ \vdots \\ W_c^* \end{pmatrix},$$

and the components W_i^* of the limiting vector are independent Lambert random variables, each of which is distributed like W^* .

A.2 Ehrenfest-like random walk

In this example we take the Ehrenfest navigation matrix

$$\mathbf{A} = \begin{pmatrix} -\gamma & \gamma \\ \gamma & -\gamma \end{pmatrix}$$

for some positive γ . This Ehrenfest-like random walk is in two dimensions. Let us call the two coordinates of the walk X(t) and Y(t), i.e., in the previously used notation, $\mathbf{X}(t) = (X(t), Y(t))^{\top}$. Note that the two vectors for the choice of movement are in opposite directions and aligned along the 45-degree line

$$X(t) + Y(t) = \lambda. \tag{A.2}$$

for some intercept λ . Thus, the movement is constrained to a linear subspace. For this walk to be tenable, both $X(0)/\gamma$ and $Y(0)/\gamma$ have to be nonnegative integers (or alternatively, λ/γ and $X(0)/\lambda$ must be a positive integer). With u and v being variables of the moment generating function, the equation to solve is

$$\frac{\partial \phi(t; u, v)}{\partial t} + (1 - e^{-\gamma u + \gamma v}) \frac{\partial \phi(t; u, v)}{\partial u} + (1 - e^{\gamma u - \gamma v}) \frac{\partial \phi(t; u, v)}{\partial v} = 0.$$

Set v = 0 and $\eta(t, u) := \phi(t; u, 0)$, and note that $\eta(t, u) = \mathbb{E}[e^{uX(t)}]$ is the moment generating function of X(t). We can rewrite the partial differential equation as

$$\frac{\partial \eta(t,u)}{\partial t} + (1 - e^{-\gamma u})\frac{\partial \eta(t,u)}{\partial u} + (1 - e^{\gamma u}) \mathbb{E}\Big[Y(t)e^{uX(t)}\Big] = 0.$$

Using the invariant in equation (A.2), we write the latter equation as

$$\frac{\partial \eta(t,u)}{\partial t} + (1 - e^{-\gamma u})\frac{\partial \eta(t,u)}{\partial u} + (1 - e^{\gamma u}) \mathbb{E}\Big[(\lambda - X(t))e^{uX(t)}\Big] = 0.$$

We thus have the simplified equation

$$\frac{\partial \eta(t,u)}{\partial t} + (e^{\gamma u} - e^{-\gamma u})\frac{\partial \eta(t,u)}{\partial u} + \lambda(1 - e^{\gamma u})\,\eta(t,u) = 0.$$

This equation has the solution

$$\eta(t,u) = \left(\frac{1 + e^{-2\gamma t + \gamma u} + e^{\gamma u} - e^{-2\gamma t}}{1 - e^{-2\gamma t + \gamma u} + e^{\gamma u} + e^{-2\gamma t}}\right)^{X(0)/\gamma} \left(\frac{1 - e^{-2\gamma t + \gamma u} + e^{\gamma u} + e^{-2\gamma t}}{2}\right)^{\lambda/\gamma}$$

As $t \to \infty$ we have the limit

$$\lim_{t \to \infty} \eta(t, u) = \left(\frac{1 + e^{\gamma u}}{2}\right)^{\lambda/\gamma}.$$

Recall that the tenability requires that λ/γ is a positive integer. Therefore, the limit of the moment generating function is that of $Bin(\lambda/\gamma, 1/2)$, namely,

a binomial random variable that counts the number of successes in λ/γ independent, identically distributed trials, with rate of success 1/2 per trial.

Unlike the Pólya–Eggenberger-like random walk, the Ehrenfest-like random walk is not much affected by where it starts in the first quadrant of the XY-plane. At any t, the exact distribution does have $\mathbf{X}(0)$ in it; however, its influence is attenuated exponentially fast in time, and in the limit it is completely obliterated.

A.3 Walking along a 45-degree hill

For $\gamma \in \mathbb{R}^+$, the navigation matrix

$$\mathbf{A} = \begin{pmatrix} -\gamma & -\gamma \\ \gamma & \gamma \end{pmatrix}$$

takes a walk along an oblique line, like climbing a 45-degree hill. That walk will remain tenable so long as Y(0) > X(0). Even if the navigator walks all the way down to the bottom of the hill (hitting the Y-axis at a positive point), the navigator will come back up along the 45-degree line with probability 1, staying in the first quadrant. In this walk, we always add or subtract increments in the two dimensions that are in the same amount; the difference Y(t) - X(t) = $Y(0) - X(0) = \lambda > 0$ remains the same at all times.

As with the Ehrenfest-like random walk of the previous section, specializing the partial differential equation of Theorem 3.1 to this walk along a 45-degree hill, we again use v = 0 and $\eta(t, u) := \phi(t; u, 0)$. We note that

$$\frac{\partial \eta(t,u)}{\partial t} + (1 - e^{-\gamma u})\frac{\partial \eta(t,u)}{\partial u} + (1 - e^{\gamma u})\left(\frac{\partial \eta(t,u)}{\partial u} + \lambda \eta(t,u)\right) = 0.$$

Rearranging, we get

$$\frac{\partial \eta(t,u)}{\partial t} + (2 - e^{\gamma u} - e^{-\gamma u}) \frac{\partial \eta(t,u)}{\partial u} + \lambda (1 - e^{\gamma u}) \eta(t,u) = 0.$$

This differential equation has the solution

$$\eta(t,u) = \frac{1}{(1 - \gamma t(e^{\gamma u} - 1))^{\lambda/\gamma}} \left(\frac{e^{\gamma u} - \gamma t(e^{\gamma u} - 1)}{1 - \gamma t(e^{\gamma u} - 1)}\right)^{X(0)/\gamma}$$
(A.3)
$$= \frac{(e^{\gamma u} - \gamma t(e^{\gamma u} - 1))^{X(0)/\gamma}}{(1 - \gamma t(e^{\gamma u} - 1))^{Y(0)/\gamma}}$$

From the exact moment generating function we get the exact mean and variance for the coordinates by taking derivatives at u = 0; these quantities are

$$\mathbb{E}[X(t)] = \lambda \gamma t + X(0),$$

$$\mathbb{E}[Y(t)] = \lambda \gamma t + Y(0),$$

$$\mathbb{V}ar(X(t)) = \mathbb{V}ar(Y(t)) = \lambda \gamma^3 t^2 + (2X(0) + \lambda)\gamma^2 t$$

Next, we put u = s/t in (A.3) and use the local expansion

$$e^{\gamma u} = e^{\gamma s/t} = 1 + \frac{\gamma s}{t} + O\left(\frac{1}{t^2}\right).$$

We obtain

$$\mathbb{E}\left[e^{sX(t)/t}\right] = \frac{1}{(1 - \gamma^2 s + O(1/t))^{\lambda/\gamma}} \left(\frac{1 - \gamma^2 s + O(1/t)}{1 - \gamma^2 s + O(1/t)}\right)^{X(0)/\gamma}$$

which implies

$$\lim_{t \to \infty} \mathbb{E} \Big[e^{sX(t)/t} \Big] = \frac{1}{(1 - \gamma^2 s)^{\lambda/\gamma}}.$$

The limiting moment generating function is that of a $\text{Gamma}(\lambda/\gamma, \gamma^2)$ random variable, i.e., X(t)/t converges in distribution to a $\text{Gamma}(\lambda/\gamma, \gamma^2)$ random variable. We note that Y(t) has a similar behavior.

B Left-out details of Section 5

B.1 First initial condition

As mentioned in Section 5, we will setup and solve Kolmogorov's Forward Equations to prove Lemma 5.3.

Kolmogorov's forward equations

Let $P_{i,j}(t)$ denote the probability that, starting with *i* balls in the urn at a certain time, then *t* time units later, we have *j* balls in the urn. In particular, $P_{i,i+\ell\delta}(t)$ is the probability that, starting with *i* balls, we have exactly ℓ transitions during the next *t* time units. We follow some of the notation of Ross [14]. We let v_i denote the rate for the exponential distribution of time until the next transition occurs, when there are currently *i* balls in the urn. In our case, since the balls act independently, and each ball has exponential rate

1 of being chosen, we have $v_i = i$. We define $q_{i,j} = v_i P_{i,j}$, so $q_{i,i+\delta} = v_i = i$, and $q_{i,j} = 0$ otherwise.

We have, as in Ross's Lemma 5.4.1,

$$\lim_{t \to 0} \frac{1 - P_{i,i}(t)}{t} = v_i \quad \text{and} \quad \lim_{t \to 0} \frac{P_{i,j}(t)}{t} = q_{i,j}, \quad \text{for } i \neq j$$

So, now we set up Kolmogorov's Forward Equations, following Theorem 5.4.4 of Ross. In our case, these equations are

$$P'_{i,i}(t) = -v_i P_{i,i}(t) = -i P_{i,i}(t),$$
(B.1a)

and for $\ell \geq 1$,

$$P'_{i,i+\ell\delta}(t) = (i + (\ell - 1)\delta)P_{i,i+(\ell-1)\delta}(t) - (i + \ell\delta)P_{i,i+\ell\delta}(t).$$
 (B.1b)

We use the rising factorial notation $(i/\delta)^{\overline{\ell}} := \prod_{k=0}^{\ell-1} (i/\delta + k)$ in the statement of the lemma. The initial conditions on the $P_{i,i+\ell\delta}(t)$ are $P_{i,i}(0) = 1$ and $P_{i,i+\ell\delta}(0) = 0$ for $\ell \ge 1$.

Lemma B.1. The functions

$$P_{i,i+\ell\delta}(t) = \frac{(i/\delta)^{\overline{\ell}}}{\ell!} e^{-it} (1 - e^{-\delta t})^{\ell}$$

are the solutions to the Kolmogorov system of differential equations (B.1).

Before proving Lemma B.1, we first note this indeed is a probability distribution, since $\sum_{\ell \geq 0} P_{i,i+\ell\delta}(t) = 1$ (follows from Lemma 5.3) and since all these summands are nonnegative.

In particular, we have the solutions

$$P_{i,i}(t) = e^{-it}$$
 and $P_{i,i+\delta}(t) = \frac{i}{\delta}e^{-it}(1-e^{-\delta t}).$

Proof of Lemma B.1. Inserting t = 0 shows that the initial conditions are satisfied. We used the conventions that $0^0 = 1$, and that the empty product equals 1. The case $\ell = 0$ follows by a direct calculation.

When $\ell \geq 1$ we have

$$\begin{aligned} P_{i,i+\ell\delta}'(t) &= \frac{\prod_{k=0}^{\ell-1}(i+k\delta)}{\ell!\,\delta^{\ell}} (-ie^{-it}(1-e^{-\delta t})^{\ell} + \ell\delta e^{-it}e^{-\delta t}(1-e^{-\delta t})^{\ell-1}) \\ &= -i\frac{\prod_{k=0}^{\ell-1}(i+k\delta)}{\ell!\,\delta^{\ell}}e^{-it}(1-e^{-\delta t})^{\ell} \\ &-\ell\delta\frac{\prod_{k=0}^{\ell-1}(i+k\delta)}{\ell!\,\delta^{\ell}}e^{-it}(1-e^{-\delta t})(1-e^{-\delta t})^{\ell-1} \\ &+(i+(\ell-1)\delta)\frac{\prod_{k=0}^{\ell-1}(i+k\delta)}{(\ell-1)!\,\delta^{\ell-1}}e^{-it}e^{-\delta t}(1-e^{-\delta t})^{\ell-1} \\ &= -(i+\ell\delta)P_{i,i+\ell\delta}(t) + (i+(\ell-1)\delta)P_{i,i+(\ell-1)\delta}(t), \end{aligned}$$

which proves the lemma.

Probability generating function

By using the solutions to Kolmogorov's Forward Equations (Lemma B.1), it is not hard anymore to derive the moment generating function $\mathbb{E}[e^{vY(t)}]$.

Proof of Lemma 5.3. To calculate $\mathbb{E}[e^{vY(t)}]$, we insert i = X(0) + Y(0) at the end of this proof; but for the moment we still write the *i*. Taking Lemma B.1 and summing yields

$$\mathbb{E}\left[e^{vY(t)}\right] = \sum_{\ell \ge 0} P_{i,i+\ell\delta}(t)e^{u(i+\delta\ell)} = e^{i(v-t)}\sum_{\ell \ge 0} \frac{(i/\delta)^{\overline{\ell}}}{\ell!} (e^{\delta v}(1-e^{-\delta t}))^{\ell}.$$

Since $(1-Z)^{-\mu} = \sum_{\ell \ge 0} \mu^{\overline{\ell}} Z^{\ell} / \ell!$ (again using the rising factorial notation $\mu^{\overline{\ell}} := \mu(\mu+1) \dots (\mu+\ell-1)$), we obtain

$$\mathbb{E}\left[e^{vY(t)}\right] = e^{-i(t-v)}(1-e^{\delta v}(1-e^{-\delta t}))^{-i/\delta}$$

and the result follows by rearranging the terms.

B.2 Second initial condition

For this initial condition, we consider the situation at time t = 0. Again, we have X(0) units of the first type and Y(0) units of the second type. This translates to the probability generating function

$$\mathbb{E}\left[e^{uX(0)+vY(0)}\right] = e^{uX(0)}e^{vY(0)}$$
$$= \left(x_c|_{t=0} + \left(y_c|_{t=0} + 1\right)^{\alpha/\delta}\right)^{-X(0)/\alpha} \left(y_c|_{t=0} + 1\right)^{-Y(0)/\delta}$$

which we have by rewriting in terms of the characteristic curves x_c and y_c of Lemma 5.2.

C Alternative approach for the second moment of balanced triangular schemes

In Remark 5.4 we state an alternative approach for obtaining the first moment of the balanced triangular scheme of Section 5.

Pursuing a similar approach to the one we used to derive the mean in Theorem 4.1, we can try to go forward with the second moment. We only highlight the salient steps. Take the partial derivatives $\partial^2/\partial u^2$, $\partial^2/\partial u \partial v$, and $\partial^2/\partial v^2$ of the moment generating function of Theorem 5.1, and evaluate each equation at u = 0 and v = 0. We obtain the system of ordinary differential equations

$$\begin{aligned} \frac{d}{dt} \mathbb{E}[X^2(t)] &= 2\alpha \mathbb{E}[X^2(t)] + \alpha^2 \mathbb{E}[X(t)] \,, \\ \frac{d}{dt} \mathbb{E}[X(t)Y(t)] &= (\alpha + \delta) \mathbb{E}[X(t)Y(t)] + \beta \mathbb{E}[X^2(t)] + \alpha\beta \mathbb{E}[X(t)] \,, \\ \frac{d}{dt} \mathbb{E}[Y^2(t)] &= 2\delta \mathbb{E}[Y^2(t)] + 2\beta \mathbb{E}[X(t)Y(t)] + \beta^2 \mathbb{E}[X(t)] + \delta^2 \mathbb{E}[Y(t)] \,, \end{aligned}$$

with $\beta = \delta - \alpha$. This system is to be solved under the initial conditions $\mathbb{E}[X^2(0)] = X(0)^2$, $\mathbb{E}[X(0)Y(0)] = X(0)Y(0)$, and $\mathbb{E}[Y^2(0)] = Y(0)^2$.

We can solve it sequentially, starting with $\mathbb{E}[X^2(t)]$, as its differential equation is self contained. We can then plug in the solution of $\mathbb{E}[X^2(t)]$ into the differential equation for $\mathbb{E}[X(t)Y(t)]$, which at this point would have only known components on the right-hand side. Finally, we plug in all the known functions of averages and mixed moments in the differential equation for $\mathbb{E}[Y^2(t)]$.