

Modelling censored losses using splicing: a global fit strategy with mixed Erlang and extreme value distributions

Tom Reynkens¹, Roel Verbelen², Jan Beirlant*^{1,3}, and Katrien Antonio^{2,4}

¹LStat and LRisk, Department of Mathematics, KU Leuven.
Celestijnenlaan 200B, 3001 Leuven, Belgium.

²LStat and LRisk, Faculty of Economics and Business, KU Leuven.
Naamsestraat 69, 3000 Leuven, Belgium.

³Department of Mathematical Statistics and Actuarial Science, University
of the Free State. *P.O. Box 339, Bloemfontein 9300, South Africa.*

⁴Faculty of Economics and Business, University of Amsterdam.
Roetersstraat 11, 1018 WB Amsterdam, The Netherlands.

April 19, 2019

Abstract

In risk analysis, a global fit that appropriately captures the body and the tail of the distribution of losses is essential. Modelling the whole range of the losses using a standard distribution is usually very hard and often impossible due to the specific characteristics of the body and the tail of the loss distribution. A possible solution is to combine two distributions in a splicing model: a light-tailed distribution for the body which covers light and moderate losses, and a heavy-tailed distribution for the tail to capture large losses. We propose a splicing model with a mixed Erlang (ME) distribution for the body and a Pareto distribution for the tail. This combines the flexibility of the ME distribution with the ability of the Pareto distribution to model extreme values. We extend our splicing approach for censored and/or truncated data. Relevant examples of such data can be found in financial risk analysis. We illustrate the flexibility of this splicing model using practical examples from risk measurement.

Keywords: censoring, composite model, expectation-maximisation algorithm, risk measurement, tail modelling

*Corresponding author.

Email addresses: tom.reynkens@kuleuven.be (T. Reynkens), roel.verbelen@kuleuven.be (R. Verbelen), jan.beirlant@kuleuven.be (J. Beirlant) and katrien.antonio@kuleuven.be (K. Antonio).

1 Introduction

In several domains such as insurance, finance and operational risk, modelling financial losses is essential. For example, actuaries use models for claim sizes to set premiums, calculate risk measures and determine capital requirements for solvency regulations. This type of data is typically heavy-tailed and high losses can occur. A standard parametric distribution for the tail is a Pareto-type distribution, which is of key importance in extreme value theory (see e.g. McNeil, 1997). The Pareto distribution or the generalised Pareto distribution (GPD) are used to model exceedances over intermediate thresholds. However, they are not able to capture the characteristics over the whole range of the loss distribution which makes them not suitable as a global fit distribution. It is often imperative to obtain a global fit for the distribution of losses, for example in a risk analysis where focus is not only on extreme events, or when setting up a reinsurance program. Instead of trying many different standard distributions, splicing two distributions (Klugman et al., 2012) is more suitable to model the complete loss distribution. In literature, a splicing model is also called a composite model. We hereby combine a light-tailed distribution for the body which covers light and moderate losses (the so-called attritional losses), and a heavy-tailed distribution for the tail to capture large losses. In the actuarial literature simple splicing models have been proposed. Beirlant et al. (2004); Klugman et al. (2012) consider the splicing of the exponential distribution with the Pareto distribution. Other distributions for the body such as the Weibull distribution (Ciumara, 2006; Scollnik and Sun, 2012) or the log-normal distribution (Cooray and Ananda, 2005; Scollnik, 2007; Pigeon and Denuit, 2011) have also been used. Nadarajah and Bakar (2014); Bakar et al. (2015); Calderín-Ojeda and Kwok (2016) investigate the splicing of the log-normal or Weibull distribution with various tail distributions. Lee et al. (2012) consider the splicing of a mixture of two exponentials and the GPD. The use of a mixture model in the first splicing component gives more flexibility in modelling the light and moderate losses. Fackler (2013) provides an overview of spliced distributions for loss modelling. Note that splicing has not only been considered in an actuarial context. Panjer (2006); Peters and Shevchenko (2015); Aue and Kalkbrener (2006) use this technique to model operational risk data.

The mixed Erlang (ME) distribution became popular in loss modelling because of several reasons (see e.g. Willmot and Woo, 2007; Lee and Lin, 2010; Willmot and Lin, 2011; Klugman et al., 2013). The class of ME distributions with common scale parameter is dense in the space of positive continuous distributions (Tijms, 1994). Any positive continuous distribution can thus be approximated up to any given accuracy by a ME distribution. This class is also closed under mixture, convolution and compounding. Therefore, we can readily obtain aggregate loss distributions removing the need for simulations. Moreover, we can easily compute risk measures such as the Value-at-Risk (VaR), the Tail VaR (TVaR) and premiums of excess-loss insurances.

Fitting the ME distribution using direct likelihood maximisation is difficult. The ME parameters can also be estimated based on the denseness proof of Tijms (1994) but this method converges slowly and leads to overfitting (Lee and Lin, 2010). The preferred strat-

egy is to use the expectation-maximisation (EM) algorithm (Dempster et al., 1977) to fit the ME distribution as proposed by Lee and Lin (2010). An advantage is that the E- and M-steps can be solved analytically. Lee and Lin (2010) use information criteria (IC) like the Akaike information criterion (AIC, Akaike (1974)) or the Bayesian information criterion (BIC, Schwarz (1978)) to select the number of components in the mixture and as such avoid overfitting.

Our work is further inspired by the omnipresence of censoring and truncation in risk analysis and risk modelling, see e.g. Cao et al. (2009); Klugman et al. (2012); Antonio and Plat (2014); Verbelen et al. (2015).

Lower truncation occurs when payments that are below certain thresholds are not observed. In insurance, lower truncation occurs, for example, due to the presence of a deductible in the insurance contract. In some practical applications, there might be a natural bound that upper truncates the tail distribution. For example, earthquake magnitudes, forest fire areas, daily price changes of stocks have distributions that are naturally *upper truncated* (Aban et al., 2006). In an insurance context, where premiums have to be set using the fitted model, introducing an upper truncation point can prevent probability mass being assigned to unreasonably large claim amounts.

Right censoring is highly relevant in the context of loss models and risk measurement for unsettled claims in non-life insurance and reinsurance. The (re)insurer only knows the true cost of a policy when all claims on this policy are settled or closed. However, in the development or lifetime of a non-life insurance claim, a significant time may elapse between the claim occurrence and its final settlement or closure. For such unsettled claims only the payment to date is known and the quantity of interest, i.e. the final cumulative payment on a claim, is right censored. This complicates the calculation of reinsurance premiums for large claims and forces the insurer to predict, with maximum accuracy, the capital buffer that is required to indemnify the insured in the future regarding claims that happened in the past. To support this complex task, actuaries will use additional, expert information called *incurred data*. This is the sum of the actual payment (so far) on a claim and its case estimate. These case estimates are set by an experienced case handler and express the expert's estimate of the outstanding loss on a claim. For large claims, facing very long settlement (e.g. due to legal procedures or severe bodily injury), actuaries consider incurred data as a highly important source of information. We propose to construct upper bounds for the final cumulative payment on a claim using incurreds. When the true final claim amount lies between the cumulative payment up to date and the incurred value, *interval censoring* techniques can be applied as we will demonstrate in this paper.

The previous example covers random censoring. Policy limits introduce another type of censoring, namely type I (right) censoring (see e.g. Klein and Moeschberger, 2003). When the loss corresponding to a claim exceeds the policy limit, no further payments need to be made by the insurer. The loss is thus censored and only known to be larger than the amount that needs to be paid by the insurer, i.e. the policy limit. In the remainder, we only consider random censoring and not type I censoring.

In the splicing context, some work has already been done for censored and/or truncated

data. Teodorescu and Panaitescu (2009) take lower truncation into account for Weibull-Pareto splicing, and Cooray and Ananda (2005) extend their approach to type I right censored data. Verbelen et al. (2015) extend the mixed Erlang approach of Lee and Lin (2010) to censored and/or truncated data. Beirlant et al. (2007); Einmahl et al. (2008) discuss extensions of classical extreme value estimators to right censored data. Extensions to upper truncated data have been investigated by Aban et al. (2006) and Beirlant et al. (2016).

Although the ME distribution has several advantages, as discussed above, one major disadvantage is that it has an asymptotically exponential, and hence: light, tail (Neuts, 1981). Therefore, overfitting can still occur on heavy-tailed data as one needs many components to model the heavy-tailedness appropriately. The simulated sample of the GPD in Verbelen et al. (2015) illustrates this behaviour. As a first contribution, we overcome this drawback by proposing a splicing model with the ME distribution for the body and the Pareto distribution for the tail (Section 2). A global fit for financial loss data then results, which combines the flexibility of the ME distribution to model light and moderate losses with the ability of the Pareto distribution to model heavy-tailed data. Fire and motor third party liability (MTPL) insurance losses, and financial returns are examples of heavy-tailed data which are of Pareto type. This strategy avoids ad hoc combinations of a standard light-tailed distribution, such as the log-normal or the Weibull distribution, for the body with a heavy-tailed distribution for the tail, as explored in many papers on loss modelling. Moreover, a mixture of Erlangs yields more flexibility than a mixture of two exponentials as in Lee et al. (2012) while keeping analytic tractability.

As a second contribution, we extend the global fit strategy based on splicing to take both (random) censoring and truncation into account. Up to our knowledge, this full framework has not yet been considered in the literature. We provide a general fitting procedure for the model using the EM algorithm where the incompleteness is caused by censoring, see Section 3. Instead of using a splicing model, a common technique in extreme value analysis is to combine a non-parametric fit for the body and a parametric model (e.g. Pareto distribution) for the tail. However, when censoring is present, this approach can no longer be applied as we might have interval censored data points where the lower bound of the interval is in the body of the distribution, whereas the upper bound is in the tail. Our general splicing framework can handle observations of this type and can hence be used to provide a global fit. As we provide a general procedure to fit a splicing model to censored and/or truncated data, we could possibly use another extreme value distribution, such as the GPD, instead of the Pareto distribution. For the GPD, however, in case there is censoring, the expectations in the E-step can no longer be computed analytically, in contrast to the Pareto distribution.

In Section 4, we apply the general fitting procedure for censored and/or truncated data to the specific case of our ME-Pareto splicing model. The incompleteness now stems on the one hand from censoring and on the other hand from the mixing of Erlang components. The general fitting procedure is therefore extended using ideas from the procedure of Verbelen et al. (2015) for fitting the ME distribution to censored and/or truncated data.

Finally, we discuss the computation of risk measures using our splicing model in Section 5 and we apply the method to two real life data examples in Section 6.

2 Splicing of ME and Pareto distributions

2.1 General splicing model

Consider two densities f_1^* and f_2^* , and denote the corresponding cumulative distribution functions (CDFs) by F_1^* and F_2^* . Their parameters are contained in the vectors Θ_1 and Θ_2 , respectively. We assume that there are no shared parameters in Θ_1 and Θ_2 . Define now

$$f_1(x; t^l, t, \Theta_1) = \begin{cases} \frac{f_1^*(x; \Theta_1)}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)} & \text{if } t^l \leq x \leq t \\ 0 & \text{otherwise,} \end{cases}$$

$$f_2(x; t, T, \Theta_2) = \begin{cases} \frac{f_2^*(x; \Theta_2)}{F_2^*(T; \Theta_2) - F_2^*(t; \Theta_2)} & \text{if } t \leq x \leq T \\ 0 & \text{otherwise,} \end{cases}$$

where $0 \leq t^l < t < T$ are fixed points. The first density is lower truncated at t^l and upper truncated at t , and the second density is lower truncated at t and upper truncated at T . The density for the body, f_1 , and density for the tail, f_2 , are then valid densities on the intervals $[t^l, t]$ and $[t, T]$, respectively. In case of no upper truncation for the tail distribution, we set $T = +\infty$. The corresponding CDFs are

$$F_1(x; t^l, t, \Theta_1) = \begin{cases} 0 & \text{if } x \leq t^l \\ \frac{F_1^*(x; \Theta_1) - F_1^*(t^l; \Theta_1)}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)} & \text{if } t^l < x < t \\ 1 & \text{if } x \geq t, \end{cases}$$

$$F_2(x; t, T, \Theta_2) = \begin{cases} 0 & \text{if } x \leq t \\ \frac{F_2^*(x; \Theta_2) - F_2^*(t; \Theta_2)}{F_2^*(T; \Theta_2) - F_2^*(t; \Theta_2)} & \text{if } t < x < T \\ 1 & \text{if } x \geq T. \end{cases}$$

Consider the splicing weight $\pi \in (0, 1)$. The spliced density is then defined as

$$f(x; t^l, t, T, \Theta) = \begin{cases} 0 & \text{if } x \leq t^l \\ \pi f_1(x; t^l, t, \Theta_1) & \text{if } t^l < x \leq t \\ (1 - \pi) f_2(x; t, T, \Theta_2) & \text{if } t < x < T \\ 0 & \text{if } x \geq T, \end{cases}$$

where $\Theta = (\pi, \Theta_1, \Theta_2)$ is the parameter vector. We call the point t the splicing point, and the points t^l and T the lower, respectively, upper truncation points. The corresponding,

continuous, CDF is given by

$$F(x; t^l, t, T, \Theta) = \begin{cases} 0 & \text{if } x \leq t^l \\ \pi F_1(x; t^l, t, \Theta_1) & \text{if } t^l < x \leq t \\ \pi + (1 - \pi) F_2(x; t, T, \Theta_2) & \text{if } t < x < T \\ 1 & \text{if } x \geq T. \end{cases} \quad (1)$$

Most authors impose differentiability of the probability density function (PDF) at the splicing point to get a smooth density function and to reduce the number of parameters. The splicing point is then estimated together with the other model parameters using maximum likelihood estimation (MLE). This restriction results in less flexibility. Therefore, we choose to not follow this approach, but determine the splicing point directly using an extreme value analysis, see Section 4.3.

2.2 Mixed Erlang distribution

In our specific case, f_1 is the density of a mixed Erlang (ME) distribution which is lower truncated at $t^l \geq 0$ and upper truncated at $t > t_l$. More specifically, we consider a mixture of M Erlang distributions with common scale parameter $\theta > 0$.

The Erlang distribution is a Gamma distribution with an integer shape parameter. It has density function

$$f_E(x; r, \theta) = \frac{x^{r-1} \exp(-x/\theta)}{\theta^r (r-1)!} \quad \text{for } x > 0, \quad (2)$$

where r , a positive integer, is the shape parameter, and $\theta > 0$ is the scale parameter. Its inverse $\lambda = 1/\theta$ is called the rate parameter. Integrating (2) r times by parts gives the cumulative distribution function

$$F_E(x; r, \theta) = \int_0^x \frac{z^{r-1} \exp(-z/\theta)}{\theta^r (r-1)!} dz = 1 - \sum_{z=0}^{r-1} \exp(-x/\theta) \frac{(x/\theta)^z}{z!}.$$

The density of the ME distribution is then given by

$$f_1^*(x; \boldsymbol{\alpha}, \mathbf{r}, \theta) = \sum_{j=1}^M \alpha_j \frac{x^{r_j-1} \exp(-x/\theta)}{\theta^{r_j} (r_j-1)!} = \sum_{j=1}^M \alpha_j f_E(x; r_j, \theta) \quad \text{for } x > 0,$$

where the positive integers $\mathbf{r} = (r_1, \dots, r_M)$ with $r_1 < \dots < r_M$ are the shape parameters of the Erlang distributions, and $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_M)$, with $\alpha_j > 0$ and $\sum_{j=1}^M \alpha_j = 1$, are the mixing weights. Similarly, the cumulative distribution function can be written, for $x > 0$, as

$$F_1^*(x; \boldsymbol{\alpha}, \mathbf{r}, \theta) = \sum_{j=1}^M \alpha_j \left(1 - \sum_{z=0}^{r_j-1} \exp(-x/\theta) \frac{(x/\theta)^z}{z!} \right) = \sum_{j=1}^M \alpha_j F_E(x; r_j, \theta).$$

After truncation, with limits t^l and t , the probability density function becomes

$$\frac{f_1^*(x; \mathbf{r}, \Theta_1^*)}{F_1^*(t; \mathbf{r}, \Theta_1^*) - F_1^*(t^l; \mathbf{r}, \Theta_1^*)} = \sum_{j=1}^M \alpha_j \frac{f_E(x; r_j, \theta)}{F_1^*(t; \mathbf{r}, \Theta_1^*) - F_1^*(t^l; \mathbf{r}, \Theta_1^*)}$$

for $t^l \leq x \leq t$ and zero otherwise, where $\Theta_1^* = (\boldsymbol{\alpha}, \theta)$. This can be rewritten as

$$f_1(x; t^l, t, \mathbf{r}, \Theta_1) = \begin{cases} \sum_{j=1}^M \beta_j f_E^t(x; t^l, t, r_j, \theta) & \text{for } t^l \leq x \leq t \\ 0 & \text{otherwise,} \end{cases}$$

with $\Theta_1 = (\boldsymbol{\beta}, \theta)$, which is again a mixture with mixing weights

$$\beta_j = \alpha_j \frac{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)}{F_1^*(t; \mathbf{r}, \Theta_1^*) - F_1^*(t^l; \mathbf{r}, \Theta_1^*)} \quad (3)$$

and component density functions

$$f_E^t(x; t^l, t, r_j, \theta) = \frac{f_E(x; r_j, \theta)}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)}.$$

The component density functions $f_E^t(x; t^l, t, r_j, \theta)$ are truncated versions of the original component density functions $f_E(x; r_j, \theta)$. We obtain the weights β_j by reweighting the original weights α_j using the probability of the corresponding mixing component to lie in the truncation interval. Denote by F_E^t the CDF corresponding to f_E^t . The CDF corresponding to f_1 is then given by

$$F_1(x; t^l, t, \mathbf{r}, \Theta_1) = \begin{cases} 0 & \text{if } x \leq t^l \\ \sum_{j=1}^M \beta_j F_E^t(x; t^l, t, r_j, \theta) = \sum_{j=1}^M \beta_j \frac{F_E(x; r_j, \theta) - F_E(t^l; r_j, \theta)}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)} & \text{if } t^l < x < t \\ 1 & \text{if } x \geq t. \end{cases} \quad (4)$$

The number of Erlang mixtures M and the positive integer shapes \mathbf{r} are fixed when estimating $\Theta_1 = (\boldsymbol{\beta}, \theta)$. They are chosen using the approach described in Section 4 of Verbelen et al. (2016). A short overview of this approach is included in Appendix A.4.

2.3 Pareto distribution

The second density f_2 is the density of the truncated Pareto distribution with scale parameter $t > 0$, shape parameter $\gamma > 0$ and upper truncation point T that can be $+\infty$. Note that the scale parameter t coincides with the fixed lower truncation point of the tail distribution. As mentioned before, we determine it in advance using an extreme value analysis, see Section 4.3. Hence, $\Theta_2 = \gamma$. More precisely, we have

$$f_2(x; t, T, \gamma) = \frac{f_2^*(x; t, \gamma)}{F_2^*(T; t, \gamma)} = \begin{cases} \frac{\frac{1}{\gamma t} \left(\frac{x}{t}\right)^{-\frac{1}{\gamma}-1}}{1 - \left(\frac{T}{t}\right)^{-\frac{1}{\gamma}}} & \text{if } t < x < T \\ 0 & \text{otherwise,} \end{cases}$$

and

$$F_2(x; t, T, \gamma) = \begin{cases} 0 & \text{if } x \leq t \\ \frac{1 - (\frac{x}{t})^{-\frac{1}{\gamma}}}{1 - (\frac{T}{t})^{-\frac{1}{\gamma}}} & \text{if } t < x < T \\ 1 & \text{if } x \geq T. \end{cases} \quad (5)$$

3 Fitting a general splicing model to censored data using the EM algorithm

In this section, we discuss maximum likelihood estimation for fitting a general splicing model, as proposed in Section 2.1, to censored data. The special case of a splicing model that combines a mixed Erlang distribution (as introduced in Section 2.2) and a Pareto distribution (Section 2.3) is treated in the subsequent section. The parameters to be estimated are contained in the vector $\Theta = (\pi, \Theta_1, \Theta_2)$.

3.1 Randomly censored data

We represent the censored sample by $\mathcal{X} = \{(l_i, u_i) \mid i = 1, \dots, n\}$, where l_i and u_i denote the lower and upper censoring points of each data point from the sample of size n . These censoring points must be interpreted as the lower and upper endpoints of the interval that contains the data point x_i , which is not always observed. The censoring status of each data point is determined as follows:

$$\begin{aligned} \text{Uncensored:} & \quad t^l \leq l_i = x_i = u_i \leq T \\ \text{Left censored:} & \quad t^l = l_i < u_i \leq T \\ \text{Right censored:} & \quad t^l \leq l_i < u_i = T \\ \text{Interval censored:} & \quad t^l \leq l_i < u_i \leq T. \end{aligned}$$

The left censored and right censored data points can be treated as a special case of interval censored data points with $l_i = t^l$ and $u_i = T$, respectively. In the splicing context, we make a distinction between five cases of data points:

- i.* Uncensored with $t^l \leq l_i = x_i = u_i \leq t < T$
- ii.* Uncensored with $t^l < t < l_i = x_i = u_i \leq T$
- iii.* Interval censored with $t^l \leq l_i < u_i \leq t < T$
- iv.* Interval censored with $t^l < t \leq l_i < u_i \leq T$
- v.* Interval censored with $t^l \leq l_i < t < u_i \leq T$.

These cases are visualised in Figure 1. In case *v*, we make a further subdivision based on whether the unobserved data point lies below or above the splicing point t .

$$+ \sum_{i \in S_v} \ln \left(\pi + (1 - \pi) F_2(u_i; t, T, \Theta_2) - \pi F_1(l_i; t^l, t, \Theta_1) \right). \quad (6)$$

Direct numerical optimisation of the log-likelihood expression (6) is not straightforward due to the censoring. Data points corresponding to case v , where the censoring interval contains the splicing point t , lead to logarithmic terms of a sum involving the splicing weight π , the parameters Θ_1 of the body distribution as well as the parameters Θ_2 of the tail distribution of the splicing model. This prevents separate optimisation with respect to each of these parameter blocks.

We use the EM algorithm to overcome this hurdle in fitting a splicing model to censored data. This iterative method, first introduced by Dempster et al. (1977), finds the maximum likelihood estimates when the data are incomplete and direct likelihood maximisation is not easy to perform numerically. Consider the complete data \mathcal{Y} containing the uncensored sample $\mathbf{X} = (X_1, \dots, X_n)$. Given the complete version of the data, we can construct a complete likelihood function as

$$\mathcal{L}_{\text{complete}}(\Theta; \mathcal{Y}) = \prod_{i=1}^n (\pi f_1(X_i; t^l, t, \Theta_1))^{I(\{X_i \leq t\})} \prod_{i=1}^n \left((1 - \pi) f_2(X_i; t, T, \Theta_2) \right)^{I(\{X_i > t\})},$$

where $I(\{X_i \leq t\})$ is the indicator function for the event $\{X_i \leq t\}$. The corresponding complete data log-likelihood function is

$$\begin{aligned} \ell_{\text{complete}}(\Theta; \mathcal{Y}) &= \sum_{i=1}^n I(\{X_i \leq t\}) (\ln \pi + \ln f_1(X_i; t^l, t, \Theta_1)) \\ &\quad + \sum_{i=1}^n I(\{X_i > t\}) (\ln(1 - \pi) + \ln f_2(X_i; t, T, \Theta_2)). \end{aligned} \quad (7)$$

The complete version of the log-likelihood (7), as opposed to the incomplete version (6), is easy to optimise as it does no longer contain any CDF terms due to censored data points and allows for a separate optimisation with respect to π , Θ_1 and Θ_2 .

However, as we do not fully observe the complete version \mathcal{Y} of the data sample, the complete log-likelihood is a random variable. Therefore, it is not possible to directly optimise the complete data log-likelihood. The intuitive idea of the EM algorithm for obtaining parameter estimates in case of incomplete data is to take the conditional expectation of the complete data log-likelihood given the incomplete data and then use this expected log-likelihood function to estimate the parameters. However, taking the expectation of the complete data log-likelihood requires the knowledge of the parameter vector, so an iterative approach is needed.

More specifically, starting from an initial guess for the parameter vector, $\Theta^{(0)}$, the EM algorithm iterates between two steps. In the h th iteration of the E-step, we compute the conditional expectation of the complete data log-likelihood with respect to the complete data \mathcal{Y} given the observed data \mathcal{X} and using the current estimate of the parameter vector $\Theta^{(h-1)}$ as true values:

$$E \left(\ell_{\text{complete}}(\Theta; \mathcal{Y}) \mid \mathcal{X}; \Theta^{(h-1)} \right).$$

In the M-step, we maximise the conditional expectation of the complete data log-likelihood obtained in the E-step with respect to the parameter vector:

$$\Theta^{(h)} = \arg \max_{\Theta} E \left(\ell_{\text{complete}}(\Theta; \mathcal{Y}) \mid \mathcal{X}; \Theta^{(h-1)} \right).$$

Both steps are iterated until convergence.

We discuss these steps in detail for a general splicing model in the presence of random censoring in the following subsections.

3.3 Initial step

Before iterating the EM-steps, we need starting values for the splicing weight π and for the parameters of the distributions for the body and the tail: $\Theta^{(0)} = (\pi^{(0)}, \Theta_1^{(0)}, \Theta_2^{(0)})$. Suitable starting values depend on the distributions used for the body and the tail. We discuss starting values for the splicing of the ME and Pareto distributions in Appendix A.1.

3.4 E-step

In the h th iteration of the E-step, we take the conditional expectation of the complete log-likelihood (7) given the incomplete data \mathcal{X} , the points t^l , t and T , and the current estimate $\Theta^{(h-1)}$ for Θ . We distinguish the five cases of data points again to determine the contribution of a data point to the conditional expectation $E \left(\ell_{\text{complete}}(\Theta; \mathcal{Y}) \mid \mathcal{X}, t^l, t, T; \Theta^{(h-1)} \right)$:

- i.* $\ln \pi + E \left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i = u_i \leq t < T; \Theta_1^{(h-1)} \right)$
- ii.* $\ln(1 - \pi) + E \left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t < l_i = u_i \leq T; \Theta_2^{(h-1)} \right)$
- iii.* $\ln \pi + E \left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)} \right)$
- iv.* $\ln(1 - \pi) + E \left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t \leq l_i < u_i \leq T; \Theta_2^{(h-1)} \right)$
- v.* $E \left([\ln \pi + \ln f_1(X_i; t^l, t, \Theta_1)] I(\{X_i \leq t\}) \right. \\ \left. + [\ln(1 - \pi) + \ln f_2(X_i; t, T, \Theta_2)] I(\{X_i > t\}) \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right)$

Note that the event $\{t^l \leq l_i = u_i \leq t < T\}$ indicates that we know t^l , $l_i = u_i$, t and T , and that the ordering $t^l \leq l_i = u_i \leq t < T$ holds. Similar reasonings hold for the other conditional arguments in the expectations. Using the law of total expectation we can rewrite the expectation in *v.* as

$$E \left(\ln \pi + \ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)} \right)$$

$$\begin{aligned}
& \times P\left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right) \\
& + E\left(\ln(1 - \pi) + \ln f_2(X_i; t, T, \Theta_2) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \Theta_2^{(h-1)}\right) \\
& \times P\left(X_i > t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right),
\end{aligned}$$

where $\{t^l \leq l_i < X_i \leq t < u_i \leq T\}$ denotes that t^l, l_i, t, u_i and T are known, that the ordering $t^l \leq l_i < t < u_i \leq T$ holds, and that $\{X_i \leq t\}$. The considered conditional expectation of the complete log-likelihood is then given by

$$\begin{aligned}
& E\left(\ell_{\text{complete}}(\Theta; \mathcal{Y}) \mid \mathcal{X}, t^l, t, T; \Theta^{(h-1)}\right) \\
& = \sum_{i \in S_{ii.}} \left[\ln \pi + E\left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i = u_i \leq t < T; \Theta_1^{(h-1)}\right) \right] \\
& + \sum_{i \in S_{ii.}} \left[\ln(1 - \pi) + E\left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t < l_i = u_i \leq T; \Theta_2^{(h-1)}\right) \right] \\
& + \sum_{i \in S_{iii.}} \left[\ln \pi + E\left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)}\right) \right] \\
& + \sum_{i \in S_{iv.}} \left[\ln(1 - \pi) + E\left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t \leq l_i < u_i \leq T; \Theta_2^{(h-1)}\right) \right] \\
& + \sum_{i \in S_{v.}} \left[\ln \pi + E\left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)}\right) \right] \\
& \quad \times P\left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right) \\
& + \sum_{i \in S_{v.}} \left[\ln(1 - \pi) + E\left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \Theta_2^{(h-1)}\right) \right] \\
& \quad \times P\left(X_i > t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right). \tag{8}
\end{aligned}$$

Using (1), the probability in the second to last term in (8) can be written as

$$\begin{aligned}
& P\left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right) \\
& = \frac{F\left(t; t^l, t, T, \Theta^{(h-1)}\right) - F\left(l_i; t^l, t, T, \Theta^{(h-1)}\right)}{F\left(u_i; t^l, t, T, \Theta^{(h-1)}\right) - F\left(l_i; t^l, t, T, \Theta^{(h-1)}\right)} \\
& = \frac{\pi^{(h-1)} - \pi^{(h-1)} F_1\left(l_i; t^l, t, \Theta_1^{(h-1)}\right)}{\pi^{(h-1)} + (1 - \pi^{(h-1)}) F_2\left(u_i; t, T, \Theta_2^{(h-1)}\right) - \pi^{(h-1)} F_1\left(l_i; t^l, t, \Theta_1^{(h-1)}\right)}, \tag{9}
\end{aligned}$$

and the probability in the last term of (8) is given by 1 minus this expression.

3.5 M-step

We maximise (8) with respect to π , Θ_1 and Θ_2 by computing the partial derivatives and equating them to zero. In case it is not possible to find analytical solutions for one of these parameters, we need to rely on numerical procedures.

3.5.1 Maximisation w.r.t. π

We denote the number of observations in a set S as $\#S$ and use the notations n_1 and n_2 for the number of data points X_i smaller than or equal to t , and above t , respectively. The partial derivative of (8) w.r.t. π is given by

$$\frac{\partial E \left(\ell_{\text{complete}}(\Theta; \mathcal{Y}) \mid \mathcal{X}, t^l, t, T, \Theta^{(h-1)} \right)}{\partial \pi} = \frac{n_1^{(h)}}{\pi} - \frac{n_2^{(h)}}{1 - \pi}$$

with

$$n_1^{(h)} = \#S_{i.} + \#S_{iii.} + \sum_{i \in S_v} P \left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right),$$

and

$$n_2^{(h)} = \#S_{ii.} + \#S_{iv.} + \sum_{i \in S_v} P \left(X_i > t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right).$$

Data points belonging to case v are weighted using probabilities (9) and $1 - (9)$, leading to the estimates $n_1^{(h)}$ and $n_2^{(h)}$ in the h th iteration. Note that $n_1^{(h)} + n_2^{(h)} = n$. Setting the derivative equal to 0 and then solving for π yields

$$\pi^{(h)} = \frac{n_1^{(h)}}{n_1^{(h)} + n_2^{(h)}} = \frac{n_1^{(h)}}{n}. \quad (10)$$

This updated splicing weight can be interpreted as the proportion of data points smaller than or equal to t as estimated in the h th iteration.

3.5.2 Maximisation w.r.t. Θ_1

In order to maximise (8) w.r.t. Θ_1 , we have to maximise

$$\begin{aligned} & \sum_{i \in S_i} E \left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i = u_i \leq t < T; \Theta_1^{(h-1)} \right) \\ & + \sum_{i \in S_{iii.}} E \left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)} \right) \\ & + \sum_{i \in S_v} E \left(\ln f_1(X_i; t^l, t, \Theta_1) \mid t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)} \right) \\ & \quad \times P \left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right). \end{aligned}$$

3.5.3 Maximisation w.r.t. Θ_2

Similarly, to maximise (8) w.r.t. Θ_2 , we have to maximise

$$\begin{aligned} & \sum_{i \in S_{ii}} E \left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t < l_i = u_i \leq T; \Theta_2^{(h-1)} \right) \\ & + \sum_{i \in S_{iv}} E \left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l < t \leq l_i < u_i \leq T; \Theta_2^{(h-1)} \right) \\ & + \sum_{i \in S_{v}} E \left(\ln f_2(X_i; t, T, \Theta_2) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \Theta_2^{(h-1)} \right) \\ & \quad \times P \left(X_i > t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right). \end{aligned}$$

4 Fitting the ME-Pareto model

We focus on some aspects of the specific case of the mixed Erlang – Pareto (ME-Pa) splicing model. In particular, we zoom in on how the complete data log-likelihood is constructed for the EM algorithm when a mixed Erlang distribution is used for the body, discuss how the estimation algorithm simplifies in case of no censoring and comment on the selection of splicing and truncation points.

4.1 Complete data log-likelihood for mixed Erlang distribution

Besides overcoming the estimation problem related to the censored data, as explained in the previous section, the EM algorithm also offers the right estimation framework when one of the splicing components is a mixture distribution. The clue is to view the data points coming from the mixture as being incomplete since the associated component-indicator vectors are not available (McLachlan and Peel, 2001). The complete data \mathcal{Y} introduced above contains the uncensored sample $\mathbf{X} = (X_1, \dots, X_{n_1}, X_{n_1+1}, \dots, X_n)$ where we, without loss of generality, assume that the first n_1 data points are smaller than or equal to t . We further extend \mathcal{Y} with component-indicator vectors for the first n_1 data points, denoted by $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_{n_1})$ where

$$Z_{ij} = \begin{cases} 1 & \text{if } X_i \text{ comes from the } j\text{th component density } f_E^t(\cdot; t^l, t, r_j, \theta) \\ 0 & \text{otherwise} \end{cases} \quad (11)$$

for $i = 1, \dots, n_1$ and $j = 1, \dots, M$. They are distributed according to a multinomial distribution with

$$P(\mathbf{Z}_i = \mathbf{z}_i; \boldsymbol{\beta}) = \beta_1^{z_{i1}} \dots \beta_M^{z_{iM}}$$

for $i = 1, \dots, n_1$, where z_{ij} is equal to 0 or 1 and $\sum_{j=1}^M z_{ij} = 1$. The joint density of (X_i, \mathbf{Z}_i) given $\{X_i \leq t\}$ equals

$$f_{X_i, \mathbf{Z}_i}(x_i, \mathbf{z}_i; t^l, t, \mathbf{r}, \Theta_1) = f_{X_i | \mathbf{Z}_i}(x_i | \mathbf{z}_i; t^l, t, \mathbf{r}, \Theta_1) P(\mathbf{Z}_i = \mathbf{z}_i; \boldsymbol{\beta})$$

$$\begin{aligned}
&= \prod_{j=1}^M (f_E^t(x_i; t^l, t, r_j, \theta))^{z_{ij}} \prod_{j=1}^M \beta_j^{z_{ij}} \\
&= \prod_{j=1}^M (\beta_j f_E^t(x_i; t^l, t, r_j, \theta))^{z_{ij}},
\end{aligned}$$

for $i = 1, \dots, n_1$. Hence the part of the complete data log-likelihood (7) depending on Θ_1 becomes

$$\sum_{i=1}^n I(\{X_i \leq t\}) \ln f_1(X_i; t^l, t, \Theta_1) = \sum_{i=1}^{n_1} \sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)). \quad (12)$$

Full technical details on the EM algorithm for fitting the ME-Pareto model are treated in Appendix A.

4.2 Uncensored data

When no censoring is present, we only have data points from cases i and ii . Hence, the EM steps for π , the ME part and the Pareto part can be performed separately since the parts of the log-likelihood (6) containing π , Θ_1 and Θ_2 , respectively, can then be split. We discuss this simplified setting in Appendix B. The splicing weight π simply gets estimated as the number of data points smaller than or equal to the splicing point t , see (30). The algorithm of Verbelen et al. (2015) is applied to fit a ME distribution to all data points smaller than or equal to t . The γ parameter of the Pareto distribution is determined by (32). In case there is no upper truncation, i.e. $T = +\infty$, the solution for γ is the Hill estimator (Hill, 1975) with threshold t . This estimator is commonly used to estimate the shape parameter γ when modelling the tail with the Pareto distribution.

4.3 Selection of splicing and truncation points

Up to now, we assumed that the lower truncation point t_l , the splicing point t and the upper truncation point T are known. In many applications, there is no lower or upper truncation and we set $t_l = 0$ and $T = +\infty$.

If lower truncation is present, this boundary can often be deduced from the context. For example, in insurance, in case there is a common deductible, the lower truncation point is set to the value of this deductible.

The splicing point t might not always be as straightforward to determine. We do not propose to estimate it using a likelihood approach (see e.g. Cooray and Ananda, 2005; Lee et al., 2012). Rather, we use extreme value analysis to give an expert opinion about the choice of the splicing point. More specifically, we use the mean excess plot (Beirlant et al., 2004) to visualise where a transition from the body to the tail of the distribution is suitable. We demonstrate this type of modelling in the data examples in Section 6.

In situations where the upper truncation point T cannot be set based on the characteristics of the problem, as is for example the case for earthquake magnitudes, we need a strategy to decide whether upper truncation is applicable to the considered problem, and if so, an estimator for T is required. Aban et al. (2006) propose a conditional maximum likelihood estimator (MLE) for the γ parameter and the upper truncation point T of a truncated Pareto distribution when T is unknown. Beirlant et al. (2016) further extend this methodology and provide an improved estimator for T . Both papers also suggest a formal test to decide between a truncated and a non-truncated tail distribution. We explain these methods in Appendix B.5 and illustrate them on the first data example in Section 6. This approach can only be applied in case there is no censoring. For censored data, there is no method available to estimate the parameters of a truncated Pareto distribution when T is unknown.

5 Risk measures

In order to quantify the risk exposure of a company, several risk measures, such as the Value-at-Risk (VaR) and the Tail Value-at-Risk (TVaR), have been developed. Moreover, these risk measures can be used to determine the amount of capital to hold as a buffer against unexpected losses.

When estimating the risk measures using statistical methods, it is essential that the fitted model captures the data well. Especially a good fit of the tail part is crucial since this corresponds to the largest losses. A global fit, hence not only a tail fit, is needed as one might be interested in computing reinsurance premiums or performing a risk analysis where focus is not only on extreme events. Further details on the estimation of risk measures can be found in McNeil et al. (2005); Klugman et al. (2012, 2013); Albrecher et al. (2017).

5.1 Excess-loss insurance premiums

Using a fitted splicing model such as the ME-Pa model presented in this paper, we calculate premiums for an excess-loss insurance. For this type of insurance, the (re)insurer covers all losses above a certain retention level R . This means that she pays $(X - R)_+ = \max\{X - R, 0\}$, where X is the total claim amount. The loss for the insured (also called the cedent) is thus limited to R . This type of contract is typical in reinsurance where the reinsurer acts as the insurer's insurer and covers the losses of an insurance company above the retention level. Insurance premiums can be seen as a compensation for the (re)insurance company for bearing the risk of the insured claims, and the size of the premium is thus a measure for the risk of the claim. The net premium of such an insurance contract is given by

$$\Pi(R; t^l, t, T, \Theta) = E((X - R)_+) = \int_R^{+\infty} (1 - F(z; t^l, t, T, \Theta)) dz. \quad (13)$$

For $t \leq R < T$ we get

$$\begin{aligned}\Pi(R; t^l, t, T, \Theta) &= \int_R^{+\infty} (1 - (\pi + (1 - \pi)F_2(z; t, T, \Theta_2))) dz \\ &= (1 - \pi) \int_R^{+\infty} (1 - F_2(z; t, T, \Theta_2)) dz \\ &= (1 - \pi)\Pi_2(R; t, T, \Theta_2),\end{aligned}$$

whereas for $t^l \leq R < t$ we have

$$\begin{aligned}\Pi(R; t^l, t, T, \Theta) &= \int_R^t (1 - \pi F_1(z; t^l, t, \Theta_1)) dz + \int_t^{+\infty} (1 - (\pi + (1 - \pi)F_2(z; t, T, \Theta_2))) dz \\ &= (t - R) - \pi \int_R^t F_1(z; t^l, t, \Theta_1) dz + (1 - \pi) \int_t^{+\infty} (1 - F_2(z; t, T, \Theta_2)) dz \\ &= (t - R) - (t - R)\pi + \pi \int_R^t (1 - F_1(z; t^l, t, \Theta_1)) dz + (1 - \pi)\Pi_2(t; t, T, \Theta_2) \\ &= (1 - \pi)(t - R) + \pi\Pi_1(R; t^l, t, \Theta_1) + (1 - \pi)\Pi_2(t; t, T, \Theta_2).\end{aligned}$$

Note that $\Pi(R; t^l, t, T, \Theta) = \Pi(t^l; t^l, t, T, \Theta) + (t^l - R)$ for $R < t^l$ and $\Pi(R; t^l, t, T, \Theta) = 0$ for $R \geq T$.

We can rewrite

$$\begin{aligned}\Pi_1(R; t^l, t, \Theta_1) &= \int_R^t \left(1 - \frac{F_1^*(z; \Theta_1) - F_1^*(t^l; \Theta_1)}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)} \right) dz \\ &= \frac{F_1^*(t; \Theta_1)(t - R) - \int_R^t F_1^*(z; \Theta_1) dz}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)} \\ &= \frac{F_1^*(t; \Theta_1)(t - R) - (t - R) + \int_R^t (1 - F_1^*(z; \Theta_1)) dz}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)} \\ &= \frac{(F_1^*(t; \Theta_1) - 1)(t - R) + (\Pi_1^*(R; \Theta_1) - \Pi_1^*(t; \Theta_1))}{F_1^*(t; \Theta_1) - F_1^*(t^l; \Theta_1)}\end{aligned}$$

for $t^l \leq R < t$. For the ME distribution, the premium is given by

$$\Pi_1^*(R; \alpha, \theta) = \theta^2 \sum_{m=1}^M \left(\sum_{l=m}^{M-1} \left(\sum_{j=l+1}^M \alpha_j \right) \right) f_E(R; m, \theta)$$

for $R \geq 0$, see Verbelen et al. (2015). They assume, without loss of generality, that $r_m = m$ for $m = 1, \dots, M$. Note that $\Pi_1(R; t^l, t, \Theta_1) = \Pi_1(t^l; t^l, t, \Theta_1) + (t^l - R)$ for $R < t^l$ and $\Pi_1(R; t^l, t, \Theta_1) = 0$ for $R \geq t$.

Similarly, we get

$$\Pi_2(R; t, T, \Theta_2) = \int_R^T \left(1 - \frac{F_2^*(z; \Theta_2) - F_2^*(t; \Theta_2)}{F_2^*(T; \Theta_2) - F_2^*(t; \Theta_2)} \right) dz$$

$$\begin{aligned}
&= \frac{(F_2^*(T; \Theta_2) - 1)(T - R) + \int_R^T (1 - F_2^*(z; \Theta_2)) dz}{F_2^*(T; \Theta_2) - F_2^*(t; \Theta_2)} \\
&= \frac{(F_2^*(T; \Theta_2) - 1)(T - R) + (\Pi_2^*(R; \Theta_2) - \Pi_2^*(T; \Theta_2))}{F_2^*(T; \Theta_2) - F_2^*(t; \Theta_2)}
\end{aligned}$$

for $t \leq R < T$. For the Pareto distribution we have the following premium when $R \geq t$:

$$\Pi_2^*(R; t, \gamma) = \int_R^\infty \left(\frac{z}{t}\right)^{-\frac{1}{\gamma}} dz = R^{-\frac{1}{\gamma}+1} \frac{t^{\frac{1}{\gamma}}}{\frac{1}{\gamma} - 1}.$$

Note that $\Pi_2(R; t, T, \Theta_2) = \Pi_2(t; t, T, \Theta_2) + (t - R)$ for $R < t$ and $\Pi_2(R; t, T, \Theta_2) = 0$ for $R \geq T$.

In a pure excess-loss insurance, the potential loss for the (re)insurer is unlimited. However, the maximal amount that the (re)insurer has to pay can be limited to L . The excess-loss insurance with retention R and limit L , which is denoted as L xs R , has premium

$$E(\min\{(X - R)_+, L\}) = E((X - R)_+ - (X - (R + L))_+) = \Pi(R) - \Pi(R + L).$$

In practice, the limit L is typically taken as a multiple of R . More details on excess-loss insurance and limits can be found in Albrecher et al. (2017).

5.2 VaR, simulations and TVaR

The Value-at-Risk (VaR) is a popular risk measure and is defined as a quantile of the distribution, $\text{VaR}_{1-p} = F^{-1}(1 - p)$. For the spliced distribution, the quantile function is

$$F^{-1}(p; t^l, t, T, \Theta) = \begin{cases} F_1^{-1}(p/\pi; t^l, t, \Theta_1) & \text{if } 0 \leq p \leq \pi \\ F_2^{-1}((p - \pi)/(1 - \pi); t, T, \Theta_2) & \text{if } \pi < p \leq 1. \end{cases}$$

The quantile function of the ME distribution F_1^{-1} cannot be computed analytically, but can be obtained by numerically inverting the CDF. For the (truncated) Pareto distribution we have

$$\begin{aligned}
F_2^{-1}(p; t, T, \gamma) &= F_2^{*-1}(pF_2^*(T; t, \gamma); t, \gamma) = t(1 - pF_2^*(T; t, \gamma))^{-\gamma} \\
&= t \left(1 - p + p \left(\frac{T}{t}\right)^{-\frac{1}{\gamma}} \right)^{-\gamma}.
\end{aligned}$$

We can simulate losses by applying the expression for the VaR to random numbers generated from a uniform distribution on $[0, 1]$. This technique is called inverse transform sampling. Simulations are useful for aggregate loss calculations, e.g. when losses are not independent, and to determine risk measures, see Chapter 20 in Klugman et al. (2012).

Closely related is the Tail Value-at-Risk (TVaR) which is defined as the expected loss given that the loss is larger than VaR_{1-p} . It can be rewritten as (see e.g. Klugman et al., 2012)

$$\begin{aligned} \text{TVaR}_{1-p} &:= E(X | X > \text{VaR}_{1-p}) = \text{VaR}_{1-p} + E(X - \text{VaR}_{1-p} | X > \text{VaR}_{1-p}) \\ &= \text{VaR}_{1-p} + \frac{E((X - \text{VaR}_{1-p})_+)}{1 - F(\text{VaR}_{1-p})} \\ &= \text{VaR}_{1-p} + \frac{\Pi(\text{VaR}_{1-p})}{p}. \end{aligned}$$

This can thus easily be computed using the expressions for VaR_{1-p} and $\Pi(R)$. Note that the last equality only holds when the CDF is continuous in VaR_{1-p} which is the case for our spliced CDF since it is continuous everywhere.

6 Data examples

6.1 Secura Re

Our first data example concerns the Secura Re dataset from Beirlant et al. (2004) which is available at <http://lstat.kuleuven.be/Wiley/Data/secura1.txt>. It consists of $n = 381$ automobile claims from Europe filed between 1988 and 2001 that are larger than 1 200 000 euro. This means that left truncation occurs at 1 200 000 euro. The claim sizes are, amongst others, corrected for inflation. Our goal is to propose a good global fit and to provide an estimate for the premium of an excess-loss insurance with a certain retention R .

The splicing point t is chosen based on the mean excess plot (Beirlant et al., 2004). This plot consists of estimates for the mean excess values

$$e(v) = E(X - v | X > v) = \frac{\int_v^{+\infty} (1 - F(x)) dx}{1 - F(v)}, \quad (14)$$

in the order statistics $v = X_{n-k,n} = \hat{Q}\left(1 - \frac{k+1}{n+1}\right) = \hat{Q}\left(\frac{n-k}{n+1}\right)$ with $k = 1, \dots, n-1$, where the CDF F is estimated by the empirical CDF \hat{F} , and \hat{Q} is the corresponding empirical quantile function. The horizontal part on the left in the mean excess plot in Figure 2 indicates that a distribution with an exponential-like tail is suitable there, whereas the linear increasing part suggests a Pareto tail. On the right, there is a decreasing trend which suggests that there might be an upper truncation point. The splicing point is chosen at the transition of the horizontal part to the linear increasing part as indicated by the vertical dashed line. This point $t = 2\,600\,000$ lies very close to the value 2 580 026 that is determined in Beirlant et al. (2004) using adaptive threshold selection methods.

We fit the ME-Pareto splicing model starting from $M = 10$, and consider spread factors $s \in \{1, \dots, 10\}$ (see Appendix A.1). The fitted model was obtained using $s = 1$ and is summarised in Table 1. It consists of a single Erlang distribution for the body and the




Figure 2: Secura Re: Mean excess plot.

Pareto distribution for the tail. Based on the mean excess plot, a splicing model with an upper truncated Pareto distribution provides an alternative possible tail model. Fitting a ME and truncated Pareto splicing model, as discussed in Section 4.3, with the same splicing point then gives $\hat{\gamma} = 0.298$ and $\hat{T} = 9\,387\,484$ whereas the other parameters remain the same.

Splicing	ME	Pareto
$\hat{\pi} = 0.744$	$\hat{\alpha} = 1$	$\hat{\gamma} = 0.263$
$t^l = 1\,200\,000$	$\hat{r} = 8$	
$t = 2\,600\,000$	$\hat{\theta} = 217\,084$	
$T = +\infty$		

Table 1: Secura Re: summary of the fitted ME-Pa splicing model.

In order to evaluate the splicing fit with the ME and Pareto distributions, graphical tools, information criteria and goodness-of-fit (GoF) tests are considered. A first graphical tool is the survival plot in Figure 3a where the fitted survival function (black) is plotted together with the empirical survival function (orange). 95% confidence bands for the empirical estimator (dashed blue) and a vertical line indicating the splicing point are also added. These confidence bands are determined using the Dvoretzky-Kiefer-Wolfowitz inequality (Massart, 1990). The fitted spliced survival function follows the empirical survival function closely and lies well within the confidence bands. Next, to inspect this fit in more detail, a QQ-plot is constructed (Figure 3b) comparing the empirical quantiles to the fitted quantiles. The points on the QQ-plot are close to the 45 degree line suggesting a good fit. Closely related is the PP-plot in Figure 4a where the fitted survival function is plotted vs. the empirical survival function. This plot confirms that the model gives a good global

fit. However, it is difficult to assess the quality of the tail fit from the PP-plot. Therefore, a PP-plot with a minus-log scale is also constructed (Figure 4b). The upper right corner then corresponds to the tail of the distribution. As expected, there are some deviations from the 45 degree line for the largest points, but the plot still indicates a good global fit.




Figure 3: Secura Re: (a) Survival plot and (b) QQ-plot of the fitted ME-Pa splicing model.

Additional to the graphical tools, we look at the negative log-likelihood (NLL), AIC and BIC values for each model where lower values are better, see Table 2. The AIC and BIC are defined as

$$\text{AIC} = 2 \times \text{NLL} + 2 \times df \quad \text{and} \quad \text{BIC} = 2 \times \text{NLL} + \ln n \times df$$

where df denotes the degrees of freedom, i.e. the number of estimated parameters in the model. Moreover, we consider the Kolmogorov-Smirnov (KS) and Anderson-Darling (AD) GoF statistics as they are a measure for the distance between the empirical CDF and the fitted CDF of a model. The KS statistic is defined as

$$D_n = \sup_{x \geq t^l} |\hat{F}_n(x) - F(x)|$$

where \hat{F}_n is the empirical CDF based on n observations and F the fitted CDF. The AD statistic is given by

$$A_n = n \int_{t^l}^{+\infty} \frac{(\hat{F}_n(x) - F(x))^2}{F(x)(1 - F(x))} dx.$$

Note that both test statistics take lower truncation at t^l into account. These statistics are commonly used to test if the data sample is drawn from a specified (continuous) distribution. The standard P-values of the test are not valid when the model parameters are estimated from the data (Babu and Rao, 2004). Therefore, we use a bootstrap approach




Figure 4: Secura Re: PP-plots of the fitted ME-Pa splicing model with (a) ordinary and (b) minus-log scale.

that is detailed in Babu and Rao (2004) and Klugman et al. (2012). First, we compute the KS and AD test statistics using the fitted model for the Secura data. Then, we generate 1000 samples with replacement from the Secura data. For each sample, the model is fitted and then the KS and AD statistics are computed. The P-values are then obtained as the proportion of these 1000 test statistics that exceed the test statistic computed in the first step. The R (R Core Team, 2016) packages `stats` (KS) and `ADGofTest` (Gil Bellosta, 2011) (AD) are used to compute the test statistics. The results are also displayed in Table 2 where values closer to 0 indicate a better fit. The corresponding P-values are added between brackets. Apart from the fitted splicing model, we also consider the following models:

- The ME and truncated Pareto splicing model (ME-TPa) as discussed above.
- The ME and GPD splicing model (ME-GPD) with the same splicing point as before: $t = 2\,600\,000$. Hence, it has the same ME distribution for the body of the distribution as the ME-Pa and ME-TPa splicing models. The tail of the distribution is modelled by a GPD with parameters $\hat{\gamma} = 0.3512$ (shape) and $\hat{\sigma} = 626\,554.8$ (scale).
- The splicing model of Beirlant et al. (2004) combining the exponential distribution and the Pareto distribution (Exp-Pa) with the same splicing point as before: $t = 2\,600\,000$. The ML estimates for the parameters are $\hat{\lambda} = 1/1\,397\,147$ (rate of the exponential distribution) and $\hat{\gamma} = 0.263$ (shape of the Pareto distribution).
- The mixed Erlang fit of Verbelen et al. (2015).

Note that the first three models are fitted using our general fitting procedure, and the ME fit is obtained using the approach in Verbelen et al. (2015).

The simpler exponential-Pareto model (Exp-Pa) provides a worse fit than the ME-Pa, ME-GPD and ME models since the NLL, AIC and BIC values are higher. The splicing model with the truncated Pareto distribution (ME-TPa) is performing slightly worse than the untruncated model (ME-Pa). The test for truncation of a Pareto tail (Beirlant et al., 2016) gives a P-value of 0.3889, at the splicing point t , which means the null hypothesis of a non-truncated Pareto tail is not rejected on the 5% significance level. This confirms that the untruncated Pareto distribution might be more suitable. However, for every case one should decide what type of tail behaviour is appropriate and whether a bounded model has any economic or physical meaning. Based on the NLL, AIC and BIC values, we can conclude that the ME-Pa fit slightly improves the ME fit although differences are small. The values of the NLL suggest that the ME-GPD fit is slightly better than the ME-Pa fit. However, when taking both the quality of the fit and the number of parameters into account, as is done in the AIC and BIC, the ME-Pa model is preferred over the ME-GPD model. The P-values of the GoF tests are large for all models suggesting that all models provide an appropriate fit for the data. Based on the graphical tools, the IC's, and the P-values of the KS and AD tests, we propose to use the ME-Pa model when modelling the Secura Re data.

Model	NLL	AIC	BIC	KS	AD
Exp-Pa	5502.27	11 010.53	11 022.28	0.0364 (0.875)	0.6853 (0.740)
ME-Pa	5499.13	11 006.26	11 021.93	0.0221 (0.997)	0.2173 (0.987)
ME-TPa	5498.54	11 007.07	11 026.65	0.0280 (0.967)	0.2764 (0.974)
ME-GPD	5498.96	11 007.91	11 027.49	0.0221 (0.995)	0.1919 (0.995)
ME	5499.99	11 007.99	11 023.65	0.0237 (0.995)	0.1889 (0.999)

Table 2: Secura Re: NLL, AIC and BIC values, and GoF test statistics and P-values.

As an illustration, premiums for excess-loss insurances can be computed using the fitted models. Table 3 shows the computed premiums for different models and different retentions. Additional to the five previously mentioned models, premiums are also computed non-parametrically using (13) with the empirical survival function (Non-par.), and using the combination of a non-parametric fit for the body (below $t = 2\,600\,000$ as before) and the Pareto distribution for the tail (Non-par.-Pa). All five parametric models result in premiums that are close to the ones obtained using the non-parametric model when the retention levels are small. For higher levels the estimates are substantially different. The non-parametric model results in zero premiums when the retention levels are larger than the maximal data value, 7 898 639. Similarly, premium estimates are 0 for insurances with retention levels that are larger than \hat{T} when the ME-TPa splicing model is used. The ME distribution, which has an exponential tail, results unsurprisingly in lower premium estimates for high retentions than the heavy-tailed ME-Pa and ME-GPD models. The Exp-Pa and ME-Pa models have the same fit for the tail, but a different fit for the body. Therefore, the premium estimates for high retentions are the same, but the premiums for retentions below the splicing point $t = 2\,600\,000$ differ. Although the ME-Pa, ME-TPa and

ME-GPD models have the same model for the body of the distribution, the estimates for the premiums also differ for low retentions since the survival function is integrated starting from the retention level when estimating the premiums, see (13).

R	Non-par.	Non-par.-Pa	Exp-Pa	ME-Pa	ME-TPa	ME-GPD	ME
1 200 000	1 030 667	1 031 738	1 031 738	1 031 738	1 042 430	1 040 995	1 030 667
2 000 000	445 330	446 401	456 172	447 054	457 746	456 311	444 751
3 000 000	161 728	159 527	159 527	159 527	168 159	170 187	164 585
4 000 000	74 696	71 350	71 350	71 350	71 036	84 821	78 228
5 000 000	35 888	38 225	38 225	38 225	32 106	51 215	39 696
7 500 000	1075	12 298	12 298	12 298	3103	21 558	4025
10 000 000	0	5501	5501	5501	0	11 987	159

Table 3: Secura Re: estimates for premiums of excess-loss insurance with different retentions R .

6.2 Motor third party liability insurance

The second data example consists of motor third party liability (MTPL) insurance claims in Europe between 1995 and 2010 (Albrecher et al., 2017). They are evaluated at the end of 2010, i.e. right before the beginning of 2011, and 59% of the 837 claims are not closed at that time. All amounts are indexed in order to reflect costs in calendar year 2011, with inflation taken into account. Our goal is again to provide a good overall fit and to estimate excess-loss insurance premiums.

As discussed in Section 1, a significant time may elapse between the claim occurrence and its final settlement (due to e.g. legal procedures or severe bodily injury). In order to illustrate the development of a claim, in Figure 5, we show for four claims the cumulative indexed payment (full line) and the indexed incurred (dashed line) at the end of each year. The incurred at the end of a given year is equal to the sum of the cumulative payment up to that moment and an expert's estimate for the outstanding loss. The claims occurred, respectively, in 1995, 1996, 1997 and 1998. The first and third claim are closed before the end of the observation period (indicated by the vertical dashed line in Figure 5), and hence the cumulative indexed payment and the indexed incurred value at the end of 2010 are equal. The second and fourth claim are still in development at the end of 2010 and the indexed incurred is larger than the cumulative indexed payment at that moment.

We apply the splicing approach for censored data using an interval censoring framework with the cumulative indexed payments at the end of 2010 as lower bound for the final cumulative indexed payment. It makes sense to construct an upper bound based on the incurreds since they are determined conservatively using information on the specific claim: e.g. the severity of the accident, the number of people involved. As an illustration of the method, and by lack of further claim information, we use here the indexed incurreds at the end of 2010 as upper bound. However, when a claim is early in development, i.e. there is a small period between the claim occurrence and the moment of evaluation, the incurreds




Figure 5: MTPL: cumulative indexed payments (full line) and indexed incurred values (dashed line) at the end of each year for four claims. The moment of evaluation, i.e. the end of 2010, is indicated by the vertical dashed line.

might still be too uncertain to be used as an upper bound since the information available to the expert might be limited. After several years of development, the quality of the incurreds has improved a lot, as more information becomes available, making them more suitable as an upper bound. Since claims with accident years between 2006 and 2010 are still early in development, and we do not have more information to improve their incurreds, we omit them for the analysis (as is done in Albrecher et al., 2017). We then have 596 claims left and 45% of them are not closed at the end of 2010. A more prudent approach is to only use the cumulative indexed payments at the end of 2010 as lower bound in a right censoring framework. However, this does not take the valuable information of incurreds into account. Another possibility is to ignore any censoring information and to only consider the indexed incurreds when estimating the final claim amount. In this example we compare these three possible strategies.

As before, we rely on the mean excess plot to choose the splicing point t . We now use the Turnbull estimator (Turnbull, 1976) to estimate the distribution function in (14). This is a non-parametric estimator for the CDF in the case of interval censored data points. It extends the Kaplan-Meier estimator (Kaplan and Meier, 1958), which can only be used for right censored data, to interval censored data. There is no analytical solution for the Turnbull estimator and its computation relies on the EM algorithm. We use the implementation in the R package `interval` (Fay and Shaw, 2010). The resulting mean

excess estimates are

$$\hat{e}(v) = \frac{\int_v^{+\infty} (1 - \hat{F}^{TB}(x)) dx}{1 - \hat{F}^{TB}(v)},$$

where \hat{F}^{TB} is the Turnbull estimator for the CDF. We evaluate this function in $v = \hat{Q}^{TB}(1 - (k + 1)/(n + 1)) = \hat{Q}^{TB}((n - k)/(n + 1))$, for $k = 1, \dots, n - 1$, where \hat{Q}^{TB} is the estimator for the quantile function based on the Turnbull estimator, since in the uncensored case we also used the empirical quantiles corresponding to $1/(n + 1), \dots, (n - 1)/(n + 1)$. The estimates are plotted in Figure 6a. The mean excess plot now has a convex shape indicating that a Pareto tail is suitable. A different slope is visible after 500 000 and we therefore choose the splicing point at $t = 500\,000$ as shown by the vertical line. As discussed in Section 3.1, there are five classes of data points when fitting a splicing model to censored data. Using the splicing point $t = 500\,000$, the number of data points per class is $\#S_{i.} = 296$, $\#S_{ii.} = 34$, $\#S_{iii.} = 175$, $\#S_{iv.} = 25$ and $\#S_{v.} = 66$.

The model is fitted starting from $M = 10$ and with $s \in \{1, \dots, 10\}$ (see Appendix A.1). The fitted model consists of $M = 2$ Erlangs and was obtained using $s = 2$. It is summarised in Table 4.

Splicing	ME	Pareto
$\hat{\pi} = 0.873$	$\hat{\alpha} = (0.171, 0.829)$	$\hat{\gamma} = 0.438$
$t^l = 0$	$\hat{\mathbf{r}} = (1, 4)$	
$t = 500\,000$	$\hat{\theta} = 55\,227$	
$T = +\infty$		

Table 4: MTPL: summary of the fitted ME-Pa splicing model.

Some of the graphical tools used in Section 6.1 can be extended to the censoring case. The fitted survival function can be compared to the non-parametric Turnbull estimate (Figure 6b). Pointwise confidence intervals are obtained using 200 bootstrap samples generated by the R package `interval` (Fay and Shaw, 2010). They are added as dashed blue lines in Figure 6b. The fitted survival function follows the Turnbull estimate closely and stays within the confidence intervals suggesting a good fit. PP-plots are made using the fitted survival function and the Turnbull survival function, see Figures 7a and 7b, where a minus-log scale is used in the second plot. Both lines are close to the 45 degree line indicating that the fitted model is suitable for the data.

Figure 8 shows estimates for premiums of excess-loss insurance for different retentions. The premiums are estimated using the considered splicing model in the interval censoring framework (full line), and compared to estimates obtained using a splicing model based on the right censoring framework (dashed line) and a splicing model without censoring using the indexed incurreds (dash-dot line). The second approach gives higher premium estimates than the first approach since the (censored) total amount paid for each claim is not bounded from above. The incurreds are conservative expert estimates of the final cumulative claim amount. Only using the indexed incurreds in an uncensored framework does not take into




Figure 6: MTPL: (a) Mean excess plot based on the Turnbull estimator and (b) survival plot of the fitted ME-Pa splicing model.

account that the actual total amount that needs to be paid can be lower than the indexed incurreds. Therefore, it leads to higher premium estimates than for the splicing model using interval censored data. Using all information available, the cumulative indexed payments and the indexed incurreds, leads to significantly lower premium estimates.

As mentioned in the introduction, a non-parametric fit for the body and a parametric model (e.g. Pareto distribution) for large losses can be used instead of a splicing model. When censoring is present, this approach can no longer be used as we might have data points of class v (see Figure 1) where the lower bound of the interval is in the body of the distribution, whereas the upper bound is in the tail. As is shown in this example, our general splicing framework can handle observations of this type and can hence be used to provide a global fit. This global fit is e.g. needed to compute premiums for excess-loss insurances.

We discussed the possibility to use the GPD instead of the Pareto distribution in the splicing model. Unlike for the Pareto distribution, the fourth and sixth expectation in the E-step (8) cannot be computed analytically when using the GPD (with $\gamma \neq 0$). This makes the whole procedure numerically more intensive as it requires numerical integration. Without censoring, this drawback is not present as only the first two expectations in the E-step (8) need to be computed.

7 Conclusions

In order to get a suitable global fit for financial loss data we propose a new splicing model. It combines the flexibility of the mixed Erlang distribution to model the body of the




Figure 7: MTPL: PP-plots of the fitted ME-Pa splicing model with (a) ordinary and (b) minus-log scale.

distribution with the Pareto distribution to provide a suitable fit for the tail. Hence, our proposal avoids ad hoc combinations of a standard light-tailed distribution for the body with a heavy-tailed distribution for the tail.

Motivated by real life insurance data sets where censoring and truncation are omnipresent, we provide a general framework for fitting a spliced distribution to censored and/or truncated data. This fitting procedure uses the EM algorithm to handle data incompleteness due to censoring. Moreover, we give details on the application of this procedure to fit the ME-Pa model.

Estimates for excess-loss premiums and risk measures such as the VaR can be easily extended to the splicing context. We illustrate the flexibility of the proposed ME-Pa splicing approach using the lower truncated Secura Re dataset and using the MTPL dataset where censoring is present.

As we provide a general procedure to fit a splicing model to censored and/or truncated data, other distributions for the body and/or tail can be considered. We illustrated the use of the GPD instead of the Pareto distribution for the tail.

We implemented all models and tools in the R package `ReIns` which complements Albrecher et al. (2017). The package is available on CRAN: <https://CRAN.R-project.org/package=ReIns>

Acknowledgements

The authors thank Hansjörg Albrecher for the interesting discussions, and the companies for providing the data for the examples.

Roel Verbelen acknowledges support from the agency for Innovation by Science and




Figure 8: MTPL: Estimates for premiums of excess-loss insurance with different retentions using ME-Pa splicing model with interval censoring (full line), right censoring (dashed line) and no censoring (dash-dot line).

Technology (IWT). Jan Beirlant acknowledges support through a restart project from KU Leuven’s research council. Katrien Antonio acknowledges financial support from the Ageas Continental Europe Research Chair at KU Leuven and from KU Leuven’s research council [project COMPACT C24/15/001].

References

- Aban, I. B., M. M. Meerschaert, and A. K. Panorska (2006). Parameter estimation for the truncated Pareto distribution. *J. Amer. Statist. Assoc.* 101(473), 270–277.
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Trans. Automat. Control* 19(6), 716–723.
- Albrecher, H., J. Beirlant, and J. Teugels (2017). *Reinsurance: Actuarial and Statistical Aspects*. Chichester, UK: John Wiley & Sons, Ltd. To appear.
- Antonio, K. and R. Plat (2014). Micro-level stochastic loss reserving for general insurance. *Scand. Actuar. J.* 2014(7), 649–669.
- Aue, F. and M. Kalkbrener (2006). LDA at work: Deutsche Bank’s approach to quantifying operational risk. *J. Oper. Risk* 1(4), 49–93.
- Babu, G. J. and C. R. Rao (2004). Goodness-of-fit tests when parameters are estimated. *Sankhyā: The Indian Journal of Statistics* 66(1), 63–74.

- Bakar, S. A. A., N. A. Hamzah, M. Maghsoudi, and S. Nadarajah (2015). Modeling loss data using composite models. *Insurance Math. Econom.* 61, 1146–1154.
- Beirlant, J., M. I. Fraga Alves, and M. I. Gomes (2016). Tail fitting for truncated and non-truncated Pareto-type distributions. *Extremes* 19(3), 429–462.
- Beirlant, J., Y. Goegebeur, J. Teugels, and J. Segers (2004). *Statistics of Extremes: Theory and Applications*. Wiley Series in Probability and Statistics. Chichester, UK: John Wiley & Sons, Ltd.
- Beirlant, J., A. Guillou, G. Dierckx, and A. Fils-Villetard (2007). Estimation of the extreme value index and extreme quantiles under random censoring. *Extremes* 10(3), 151–174.
- Calderín-Ojeda, E. and C. F. Kwok (2016). Modeling claims data with composite Stoppa models. *Scand. Actuar. J.* 2016(9), 817–836.
- Cao, R., J. M. Vilar, and A. Devía (2009). Modelling consumer credit risk via survival analysis. *SORT* 33(1), 3–30.
- Ciumara, R. (2006). An actuarial model based on the composite Weibull-Pareto distribution. *Math. Rep. (Bucur.)*, 8(4), 401–414.
- Cooray, K. and M. M. Ananda (2005). Modeling actuarial data with a composite lognormal-Pareto model. *Scand. Actuar. J.* 2005(5), 321–334.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the EM algorithm. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* 39(1), 1–38.
- Einmahl, J. H., A. Fils-Villetard, and A. Guillou (2008). Statistics of extremes under random censoring. *Bernoulli* 14(1), 207–227.
- Fackler, M. (2013). Reinventing Pareto: Fits for both small and large losses. ASTIN Colloquium, Den Haag.
- Fay, M. P. and P. A. Shaw (2010). Exact and asymptotic weighted logrank tests for interval censored data: The interval R package. *J. Stat. Softw.* 36(2), 1–34.
- Gil Bellosta, C. J. (2011). *ADGofTest: Anderson-Darling GoF test*. R package version 0.3.
- Hill, B. M. (1975). A simple general approach to inference about the tail of a distribution. *Ann. Statist.* 3(5), 1163–1174.
- Kaplan, E. L. and P. Meier (1958). Nonparametric estimation from incomplete observations. *J. Amer. Statist. Assoc.* 53(282), 457–481.
- Klein, J. P. and M. L. Moeschberger (2003). *Survival Analysis: Techniques for Censored and Truncated Data*. New York, NY: Springer-Verlag. 2nd edition.

- Klugman, S. A., H. H. Panjer, and G. E. Willmot (2012). *Loss Models: From Data to Decisions*. Wiley Series in Probability and Statistics. Hoboken, NJ: John Wiley & Sons, Inc. 4th edition.
- Klugman, S. A., H. H. Panjer, and G. E. Willmot (2013). *Loss Models: Further Topics*. Wiley Series in Probability and Statistics. Hoboken, NJ: John Wiley & Sons, Inc.
- Lee, D., W. K. Li, and T. S. T. Wong (2012). Modeling insurance claims via a mixture exponential model combined with peaks-over-threshold approach. *Insurance Math. Econom.* 51(3), 538–550.
- Lee, S. C. K. and X. S. Lin (2010). Modeling and evaluating insurance losses via mixtures of Erlang distributions. *N. Am. Actuar. J.* 14(1), 107–130.
- Massart, P. (1990). The tight constant in the Dvoretzky-Kiefer-Wolfowitz inequality. *Ann. Probab.* 18(3), 1269–1283.
- McLachlan, G. and D. Peel (2001). *Finite Mixture Models*. Wiley Series in Probability and Statistics. Hoboken, NJ: John Wiley & Sons, Inc.
- McNeil, A. J. (1997). Estimating the tails of loss severity distributions using extreme value theory. *Astin Bull.* 27(1), 117–137.
- McNeil, A. J., R. Frey, and P. Embrechts (2005). *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton Series in Finance. Princeton, NJ: Princeton University Press.
- Nadarajah, S. and S. A. A. Bakar (2014). New composite models for the Danish fire insurance data. *Scand. Actuar. J.* 2014(2), 180–187.
- Neuts, M. F. (1981). *Matrix-geometric Solutions in Stochastic Models: An Algorithmic Approach*. Baltimore, MD: John Hopkins University Press.
- Panjer, H. H. (2006). *Operational Risk: Modeling Analytics*. Wiley Series in Probability and Statistics. Hoboken, NJ: John Wiley & Sons, Inc.
- Peters, G. W. and P. V. Shevchenko (2015). *Advances in Heavy Tailed Risk Modeling: A Handbook of Operational Risk*. Hoboken, NJ: John Wiley & Sons, Inc.
- Pigeon, M. and M. Denuit (2011). Composite lognormal-Pareto model with random threshold. *Scand. Actuar. J.* 2011(3), 177–192.
- R Core Team (2016). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Schwarz, G. E. (1978). Estimating the dimension of a model. *J. Amer. Statist. Assoc.* 6(2), 461–464.

- Scollnik, D. P. M. (2007). On composite lognormal-Pareto models. *Scand. Actuar. J.* 2007(1), 20–33.
- Scollnik, D. P. M. and C. Sun (2012). Modeling with Weibull-Pareto models. *N. Am. Actuar. J.* 16(2), 260–272.
- Teodorescu, S. and E. Panaitescu (2009). On the truncated composite Weibull-Pareto model. *Math. Rep. (Bucur.)* 11(61), 259–273.
- Tijms, H. C. (1994). *Stochastic Models: an Algorithmic Approach*. Wiley Series in Probability and Statistics. Chichester, UK: John Wiley & Sons, Ltd.
- Turnbull, B. W. (1976). The empirical distribution function with arbitrarily grouped, censored and truncated data. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* 38(3), 290–295.
- Verbelen, R., K. Antonio, and G. Claeskens (2016). Multivariate mixtures of Erlangs for density estimation under censoring. *Lifetime Data Anal.* 22(3), 429–455.
- Verbelen, R., L. Gong, K. Antonio, A. Badescu, and S. Lin (2015). Fitting mixtures of Erlangs to censored and truncated data using the EM algorithm. *Astin Bull.* 45(3), 729–758.
- Willmot, G. E. and X. S. Lin (2011). Risk modelling with the mixed Erlang distribution. *Appl. Stoch. Models Bus. Ind.* 27(1), 2–16.
- Willmot, G. E. and J. Woo (2007). On the class of Erlang mixtures with risk theoretic applications. *N. Am. Actuar. J.* 11(2), 99–115.

A Fitting the ME-Pareto model to censored data using the EM algorithm

This appendix contains all the details of the framework described in Section 3 applied to the splicing of the ME and Pareto distributions.

A.1 Initial step

We estimate the splicing constant π initially by the Turnbull estimator (Turnbull, 1976) in t , i.e. $\pi^{(0)} = \hat{F}_{TB}(t)$ since the estimate for π in the h th iteration ($\pi^{(h)}$) can be seen as the proportion of data points smaller than or equal to t , see (10).

Initial values for the ME parameters follow from Verbelen et al. (2016) which are improvements of the starting values from Verbelen et al. (2015). Consider the data \mathbf{d} consisting of x_i for all uncensored data points, of l_i for all right censored data points, of u_i for all left censored data points and of the interval midpoints, $\frac{l_i+u_i}{2}$, for all interval censored data points. Restricting the data \mathbf{d} to all data points that are smaller than or equal to t gives $\tilde{\mathbf{d}}$

which has length $n_{\tilde{\mathbf{d}}}$. Verbelen et al. (2016) use the following starting values: $\theta^{(0)} = \frac{\max(\tilde{\mathbf{d}})}{s}$, for a given spread factor s ,

$$\mathbf{r} = \left(\left[\frac{\hat{Q}(0; \tilde{\mathbf{d}})}{\theta^{(0)}} \right], \left[\frac{\hat{Q}\left(\frac{1}{M-1}; \tilde{\mathbf{d}}\right)}{\theta^{(0)}} \right], \dots, \left[\frac{\hat{Q}(1; \tilde{\mathbf{d}})}{\theta^{(0)}} \right] \right) \quad (15)$$

and

$$\alpha_j^{(0)} = \frac{\sum_{i=1}^{n_{\tilde{\mathbf{d}}}} I\left(\left\{r_{j-1}\theta^{(0)} < \tilde{d}_i \leq r_j\theta^{(0)}\right\}\right)}{n_{\tilde{\mathbf{d}}}}$$

for $j = 1, \dots, M$, where $r_0 = 0$ for notational convenience and $\hat{Q}(\cdot; \tilde{\mathbf{d}})$ is the empirical quantile function of the data $\tilde{\mathbf{d}}$.

As initial value for γ , we use the Hill estimator (Hill, 1975) with threshold t applied to \mathbf{d} .

This gives $\Theta^{(0)} = (\pi^{(0)}, \beta^{(0)}, \theta^{(0)}, \gamma^{(0)})$ where $\alpha^{(0)}$ is transformed to $\beta^{(0)}$ using (3).

A.2 E-step

In the h th iteration of the E-step, we take the conditional expectation of the complete log-likelihood given the incomplete data \mathcal{X} and using the current estimate $\Theta^{(h-1)}$ for Θ . As said before, we assume that there are no shared parameters in π , Θ_1 and Θ_2 . We can hence split the maximisation in three parts. Therefore, we also consider three parts in the conditional expectation of the complete log-likelihood.

A.2.1 π

We easily obtain the probability (9) using (4) and (5) with $\Theta_1 = \Theta_1^{(h-1)}$ and $\gamma = \gamma^{(h-1)}$. This gives

$$\begin{aligned} & P\left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right) \\ &= \frac{\pi^{(h-1)} - \pi^{(h-1)} F_1\left(l_i; t^l, t, \mathbf{r}, \Theta_1^{(h-1)}\right)}{\pi^{(h-1)} + (1 - \pi^{(h-1)}) \frac{1 - \left(\frac{u_i}{t}\right)^{-\frac{1}{\gamma^{(h-1)}}}}{1 - \left(\frac{T}{t}\right)^{-\frac{1}{\gamma^{(h-1)}}}} - \pi^{(h-1)} F_1\left(l_i; t^l, t, \mathbf{r}, \Theta_1^{(h-1)}\right)}, \end{aligned} \quad (16)$$

where

$$\begin{aligned} F_1\left(l_i; t^l, t, \mathbf{r}, \Theta_1^{(h-1)}\right) &= \sum_{j=1}^M \beta_j^{(h-1)} F_E^t(l_i; t^l, t, r_j, \theta^{(h-1)}) \\ &= \sum_{j=1}^M \beta_j^{(h-1)} \frac{F_E(l_i; r_j, \theta^{(h-1)})}{F_E(t; r_j, \theta^{(h-1)}) - F_E(t^l; r_j, \theta^{(h-1)})}. \end{aligned}$$

A.2.2 ME distribution

After using arguments similar to those used to obtain (8), we get that the conditional expectation of (12), the part of the complete data log-likelihood depending on Θ_1 , is given by

$$\begin{aligned}
& \sum_{i \in S_i} E \left(\sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \middle| t^l \leq l_i = u_i \leq t < T; \Theta_1^{(h-1)} \right) \\
& + \sum_{i \in S_{iii}} E \left(\sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \middle| t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)} \right) \\
& + \sum_{i \in S_v} E \left(\sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \middle| t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)} \right) \\
& \quad \times P \left(X_i \leq t \middle| t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right). \tag{17}
\end{aligned}$$

Uncensored data. Following Verbelen et al. (2015) we obtain that the first part of the sum is equal to

$$\begin{aligned}
& \sum_{i \in S_i} \sum_{j=1}^M {}^i z_{ij}^{(h)} \left[\ln \beta_j + (r_j - 1) \ln x_i - \frac{x_i}{\theta} - r_j \ln \theta \right. \\
& \quad \left. - \ln((r_j - 1)!) - \ln (F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)) \right].
\end{aligned}$$

Here is

$${}^i z_{ij}^{(h)} = P \left(Z_{ij} = 1 \middle| t^l \leq l_i = u_i \leq t < T; \Theta_1^{(h-1)} \right) = \frac{\alpha_j^{(h-1)} f_E(x_i; r_j, \theta^{(h-1)})}{\sum_{m=1}^M \alpha_m^{(h-1)} f_E(x_i; r_m, \theta^{(h-1)})} \tag{18}$$

the posterior probability that an uncensored data point x_i with $x_i \leq t$ (hence $i \in S_i$) belongs to the j th component in the mixture. In the E-step for the uncensored data, only these probabilities need to be computed for all $i \in S_i$ and $j = 1, \dots, M$. They remain the same in the truncated and the untruncated case.

Censored data. We now have to distinguish between the cases *iii* and *v*. The derivations for case *iii* follow from Verbelen et al. (2015).

Denote by ${}^{iii} z_{ij}^{(h)}$ the posterior probability that the data point x_i belongs to the j th component in the mixture for a censored data point with $u_i \leq t$. Then,

$$\sum_{i \in S_{iii}} E \left(\sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \middle| t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)} \right)$$

$$\begin{aligned}
&= \sum_{i \in S_{iii.}} \sum_{j=1}^M iii. z_{ij}^{(h)} E \left(\ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \mid Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)} \right) \\
&= \sum_{i \in S_{iii.}} \sum_{j=1}^M iii. z_{ij}^{(h)} \left[\ln \beta_j + (r_j - 1) E \left(\ln X_i \mid Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)} \right) \right. \\
&\quad \left. - \frac{1}{\theta} E \left(X_i \mid Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)} \right) - r_j \ln \theta - \ln((r_j - 1)!) \right. \\
&\quad \left. - \ln (F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)) \right] \tag{19}
\end{aligned}$$

where we use the law of total expectation in the first equality. Using Bayes' rule, we can compute the posterior probabilities $iii. z_{ij}^{(h)}$, for $i \in S_{iii.}$ and $j = 1, \dots, M$, as

$$\begin{aligned}
iii. z_{ij}^{(h)} &:= P \left(Z_{ij} = 1 \mid t^l \leq l_i < u_i \leq t < T; \Theta_1^{(h-1)} \right) \\
&= \frac{\beta_j^{(h-1)} (F_E^t(u_i; t^l, t, r_j, \theta^{(h-1)}) - F_E^t(l_i; t^l, t, r_j, \theta^{(h-1)}))}{\sum_{m=1}^M \beta_m^{(h-1)} (F_E^t(u_i; t^l, t, r_m, \theta^{(h-1)}) - F_E^t(l_i; t^l, t, r_m, \theta^{(h-1)}))} \\
&= \frac{\beta_j^{(h-1)} \frac{F_E(u_i; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})}{F_E(t; r_j, \theta^{(h-1)}) - F_E(t^l; r_j, \theta^{(h-1)})}}{\sum_{m=1}^M \beta_m^{(h-1)} \frac{F_E(u_i; r_m, \theta^{(h-1)}) - F_E(l_i; r_m, \theta^{(h-1)})}{F_E(t; r_m, \theta^{(h-1)}) - F_E(t^l; r_m, \theta^{(h-1)})}} \\
&= \frac{\alpha_j^{(h-1)} (F_E(u_i; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)}))}{\sum_{m=1}^M \alpha_m^{(h-1)} (F_E(u_i; r_m, \theta^{(h-1)}) - F_E(l_i; r_m, \theta^{(h-1)}))}. \tag{20}
\end{aligned}$$

The expression for the posterior probability in the censored case has the same form as in the uncensored case (18), but with the densities replaced by the probabilities to lay between the lower and upper censoring points. The terms in (19) containing $(r_j - 1) E (\ln X_i \mid Z_{ij} = 1, t^l \leq l_i < t < u_i \leq T; \theta^{(h-1)})$ do not play a role in the EM algorithm as they do not depend on β or θ . We also need to compute the following conditional expected value in the E-step:

$$\begin{aligned}
&E \left(X_i \mid Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)} \right) \\
&= \int_{l_i}^{u_i} x \frac{f_E(x; r_j, \theta^{(h-1)})}{F_E(u_i; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})} dx \\
&= \frac{r_j \theta^{(h-1)}}{F_E(u_i; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})} \int_{l_i}^{u_i} \frac{x^{r_j} \exp(-x/\theta^{(h-1)})}{(\theta^{(h-1)})^{r_j+1} r_j!} dx \\
&= \frac{r_j \theta^{(h-1)} (F_E(u_i; r_j + 1, \theta^{(h-1)}) - F_E(l_i; r_j + 1, \theta^{(h-1)}))}{F_E(u_i; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})}, \tag{21}
\end{aligned}$$

for $i \in S_{iii.}$ and $j = 1, \dots, M$, which has a closed-form expression.

We perform similar calculations for the case v . Denote by $v. z_{ij}^{(h)}$ the posterior probability that the data point x_i belongs to the j th component in the mixture for a censored data

point with $\{X_i \leq t\}$ and $u_i > t$. The third part of the sum in (17), without the probability, is then given by

$$\begin{aligned}
& \sum_{i \in S_v} E \left(\sum_{j=1}^M Z_{ij} \ln (\beta_j f_E^t(X_i; t^l, t, r_j, \theta)) \middle| t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)} \right) \\
&= \sum_{i \in S_v} \sum_{j=1}^M v \cdot z_{ij}^{(h)} \left[\ln \beta_j + (r_j - 1) E (\ln X_i \mid Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)}) \right. \\
&\quad \left. - \frac{1}{\theta} E (X_i \mid Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T, \theta^{(h-1)}) - r_j \ln \theta - \ln((r_j - 1)!) \right. \\
&\quad \left. - \ln (F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)) \right]. \tag{22}
\end{aligned}$$

The posterior probabilities $v \cdot z_{ij}^{(h)}$, for $i \in S_v$ and $j = 1, \dots, M$, are given by

$$\begin{aligned}
v \cdot z_{ij}^{(h)} &= P \left(Z_{ij} = 1 \middle| t^l \leq l_i < X_i \leq t < u_i \leq T; \Theta_1^{(h-1)} \right) \\
&= \frac{\beta_j^{(h-1)} (F_E^t(t; t^l, t, r_j, \theta^{(h-1)}) - F_E^t(l_i; t^l, t, r_j, \theta^{(h-1)}))}{\sum_{m=1}^M \beta_m^{(h-1)} (F_E^t(t; t^l, t, r_m, \theta^{(h-1)}) - F_E^t(l_i; t^l, t, r_m, \theta^{(h-1)}))} \\
&= \frac{\alpha_j^{(h-1)} (F_E(t; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)}))}{\sum_{m=1}^M \alpha_m^{(h-1)} (F_E(t; r_m, \theta^{(h-1)}) - F_E(l_i; r_m, \theta^{(h-1)}))}. \tag{23}
\end{aligned}$$

This expression corresponds to (20) with t instead of u_i because of the conditioning on the event $\{X_i \leq t < u_i\}$. Again, the terms in (22) containing $(r_j - 1) E (X_i \mid Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)})$ do not play a role in the EM algorithm as they do not depend on β or θ . Also,

$$\begin{aligned}
& E (X_i \mid Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)}) \\
&= \int_{l_i}^t x \frac{f_E(x; r_j, \theta^{(h-1)})}{F_E(t; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})} dx \\
&= \frac{r_j \theta^{(h-1)} (F_E(t; r_j + 1, \theta^{(h-1)}) - F_E(l_i; r_j + 1, \theta^{(h-1)}))}{F_E(t; r_j, \theta^{(h-1)}) - F_E(l_i; r_j, \theta^{(h-1)})}, \tag{24}
\end{aligned}$$

for $i \in S_v$ and $j = 1, \dots, M$, which has a closed-form expression. Likewise, (24) corresponds to (21) with t instead of u_i since we condition on the event $\{X_i \leq t < u_i\}$.

A.2.3 Pareto distribution

The relevant part for the Pareto parameter γ is

$$\sum_{i \in S_{ii}} E (\ln f_2(X_i; t, T, \gamma) \mid t^l < t < l_i = u_i \leq T; \gamma^{(h-1)})$$

$$\begin{aligned}
& + \sum_{i \in S_{iv}} E \left(\ln f_2(X_i; t, T, \gamma) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \\
& + \sum_{i \in S_v} E \left(\ln f_2(X_i; t, T, \gamma) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)} \right) \\
& \quad \times P \left(X_i > t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right).
\end{aligned}$$

The first conditional expectation is simply equal to

$$\begin{aligned}
& E \left(\ln f_2(X_i; t, T, \gamma) \mid t^l < t < l_i = u_i \leq T; \gamma^{(h-1)} \right) \\
& = \ln f_2(x_i; t, T, \gamma) = -\ln(\gamma t) - \left(\frac{1}{\gamma} + 1 \right) \ln \left(\frac{x_i}{t} \right) - \ln \left(1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}} \right).
\end{aligned}$$

Starting from the definition we obtain

$$\begin{aligned}
& E \left(\ln f_2(X_i; t, T, \gamma) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \\
& = -\ln(\gamma t) - \left(\frac{1}{\gamma} + 1 \right) E \left(\ln \left(\frac{X_i}{t} \right) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) - \ln \left(1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}} \right).
\end{aligned}$$

Using integration by parts, we get

$$\begin{aligned}
& E \left(\ln \left(\frac{X_i}{t} \right) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \\
& = \int_{l_i}^{u_i} \ln \left(\frac{x}{t} \right) \frac{f_2^*(x; t, \gamma^{(h-1)})}{F_2^*(u_i; t, \gamma^{(h-1)}) - F_2^*(l_i; t, \gamma^{(h-1)})} dx \\
& = \frac{1}{\left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}} \int_{l_i}^{u_i} \frac{\ln \left(\frac{x}{t} \right)}{\gamma^{(h-1)} t} \left(\frac{x}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}-1} dx \\
& = \frac{1}{\left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}} \int_{\frac{l_i}{t}}^{\frac{u_i}{t}} \frac{\ln v}{\gamma^{(h-1)}} v^{-\frac{1}{\gamma^{(h-1)}}-1} dv \\
& = \frac{1}{\left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}} \left(- \left[\ln v v^{-\frac{1}{\gamma^{(h-1)}}} \right]_{\frac{l_i}{t}}^{\frac{u_i}{t}} + \int_{\frac{l_i}{t}}^{\frac{u_i}{t}} v^{-\frac{1}{\gamma^{(h-1)}}-1} dv \right) \\
& = \frac{1}{\left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}} \left(- \left[\ln v v^{-\frac{1}{\gamma^{(h-1)}}} \right]_{\frac{l_i}{t}}^{\frac{u_i}{t}} + \left[-\gamma^{(h-1)} v^{-\frac{1}{\gamma^{(h-1)}}} \right]_{\frac{l_i}{t}}^{\frac{u_i}{t}} \right) \\
& = \frac{\left(\ln \left(\frac{l_i}{t} \right) + \gamma^{(h-1)} \right) \left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\ln \left(\frac{u_i}{t} \right) + \gamma^{(h-1)} \right) \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}}{\left(\frac{l_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}} - \left(\frac{u_i}{t} \right)^{-\frac{1}{\gamma^{(h-1)}}}}. \tag{25}
\end{aligned}$$

We compute the third conditional expectation similarly which gives

$$E \left(\ln f_2(X_i; t, T, \gamma) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)} \right)$$

$$\begin{aligned}
&= -\ln(\gamma t) - \left(\frac{1}{\gamma} + 1\right) E\left(\ln\left(\frac{X_i}{t}\right) \middle| t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)}\right) \\
&\quad - \ln\left(1 - \left(\frac{T}{t}\right)^{-\frac{1}{\gamma}}\right),
\end{aligned}$$

with

$$E\left(\ln\left(\frac{X_i}{t}\right) \middle| t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)}\right) = \frac{\gamma^{(h-1)} - \left(\ln\left(\frac{u_i}{t}\right) + \gamma^{(h-1)}\right) \left(\frac{u_i}{t}\right)^{-\frac{1}{\gamma^{(h-1)}}}}{1 - \left(\frac{u_i}{t}\right)^{-\frac{1}{\gamma^{(h-1)}}}}. \quad (26)$$

Note that the last expression corresponds to (25) with $l_i = t$ since we condition on the event $\{l_i \leq t < X_i\}$.

A.3 M-step

In the M-step, the expected value of the complete data log-likelihood obtained in the E-step is maximised w.r.t. the parameter vector Θ . As said before, we assume that there are no shared parameters in π , Θ_1 and Θ_2 . This assures that we can split the maximisation in three parts.

A.3.1 π

Maximisation with respect to π gives

$$\pi^{(h)} = \frac{n_1^{(h)}}{n},$$

with

$$n_1^{(h)} = \#S_i + \#S_{iii} + \sum_{i \in S_v} P\left(X_i \leq t \middle| t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right),$$

see (10). An expression for the probability can be found in (16).

A.3.2 ME distribution

The expected value of the complete data log-likelihood obtained in the E-step is now maximised with respect to the parameter vector $\Theta_1 = (\beta, \theta)$ over all (β, θ) with $\beta_j > 0$, $\sum_{j=1}^M \beta_j = 1$ and $\theta > 0$.

To maximise over the mixing weights β ,

$$\sum_{j=1}^M \left(\sum_{i \in S_i} i. z_{ij}^{(h)} + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)} \right) \ln \beta_j$$

needs to be maximised where $P_{1,i}^{(h)} = P\left(X_i \leq t \mid t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)}\right)$, see (16). Setting $\beta_M = 1 - \sum_{j=1}^{M-1} \beta_j$ makes sure that the restriction $\sum_{j=1}^M \beta_j = 1$ holds. Equating the partial derivatives at $\beta^{(h)}$ to zero gives

$$\beta_j^{(h)} = \frac{\sum_{i \in S_i} i. z_{ij}^{(h)} + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)}}{\sum_{i \in S_i} i. z_{iM}^{(h)} + \sum_{i \in S_{iii}} iii. z_{iM}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{iM}^{(h)}} \beta_M^{(h)}$$

for $j = 1, \dots, M-1$. Applying the sum constraint gives

$$\beta_M^{(h)} = \frac{\sum_{i \in S_i} i. z_{iM}^{(h)} + \sum_{i \in S_{iii}} iii. z_{iM}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{iM}^{(h)}}{n_1^{(h)}}.$$

The same form also follows for $j = 1, \dots, M-1$:

$$\beta_j^{(h)} = \frac{\sum_{i \in S_i} i. z_{ij}^{(h)} + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)}}{n_1^{(h)}}. \quad (27)$$

The estimate for the prior probability β_j in the truncated mixture is thus the average of the posterior probabilities of belonging to the j th component in the mixture.

In order to maximise with respect to θ , we set the first order partial derivative at $\theta^{(h)}$ equal to zero

$$\begin{aligned} 0 &= \sum_{j=1}^M \left(\sum_{i \in S_i} i. z_{ij}^{(h)} \left(-\frac{r_j}{\theta} - \frac{\frac{\partial}{\partial \theta} [F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)]}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)} + \frac{x_i}{\theta^2} \right) \right. \\ &\quad + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} \left(-\frac{r_j}{\theta} - \frac{\frac{\partial}{\partial \theta} [F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)]}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)} \right. \\ &\quad \left. \left. + \frac{E(X_i \mid Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)})}{\theta^2} \right) \right. \\ &\quad \left. + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)} \left(-\frac{r_j}{\theta} - \frac{\frac{\partial}{\partial \theta} [F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)]}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)} \right. \right. \\ &\quad \left. \left. + \frac{E(X_i \mid Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)})}{\theta^2} \right) \right) \Bigg|_{\theta=\theta^{(h)}} \\ &= -\frac{1}{\theta^{(h)}} \sum_{j=1}^M \left(\sum_{i \in S_i} i. z_{ij}^{(h)} + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)} \right) r_j \\ &\quad - \sum_{j=1}^M \left(\sum_{i \in S_i} i. z_{ij}^{(h)} + \sum_{i \in S_{iii}} iii. z_{ij}^{(h)} + \sum_{i \in S_v} P_{1,i}^{(h)} v. z_{ij}^{(h)} \right) \frac{\frac{\partial}{\partial \theta} [F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)]}{F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta)} \Bigg|_{\theta=\theta^{(h)}} \\ &\quad + \frac{1}{\theta^{(h)2}} \sum_{j=1}^M \left(\sum_{i \in S_i} i. z_{ij}^{(h)} x_i \right. \end{aligned}$$

$$\begin{aligned}
& + \sum_{i \in S_{iii.}} iii. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)}) \\
& + \sum_{i \in S_{v.}} P_{1,i}^{(h)} v. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)}) \Big).
\end{aligned}$$

Using the lower incomplete gamma function

$$\gamma(s, x) = \int_0^x z^{s-1} \exp(-z) dz,$$

we write the cumulative distribution function of an Erlang distribution as

$$F_E(x; r_j, \theta) = \int_0^x \frac{z^{r_j-1} \exp(-z/\theta)}{\theta^{r_j} (r_j - 1)!} dz = \frac{1}{(r_j - 1)!} \int_0^{x/\theta} u^{r_j-1} \exp(-u) du = \frac{\gamma(r_j, x/\theta)}{(r_j - 1)!}.$$

Applying the Leibniz rule gives that the partial derivative of F_E with respect to θ is equal to

$$\frac{\partial F_E(x; r_j, \theta)}{\partial \theta} = \frac{\partial \gamma(r_j, x/\theta)}{\partial \theta} = \frac{\left(\frac{x}{\theta}\right)^{r_j-1} \exp\left(-\frac{x}{\theta}\right) \left(-\frac{x}{\theta^2}\right)}{(r_j - 1)!}.$$

Using (27) and this derivative gives

$$\begin{aligned}
0 = & -\frac{n_1^{(h)}}{\theta^{(h)}} \sum_{j=1}^M \beta_j^{(h)} r_j - n_1^{(h)} \sum_{j=1}^M \beta_j^{(h)} \frac{\left(\frac{t^l}{\theta}\right)^{r_j-1} \exp\left(-\frac{t^l}{\theta}\right) \frac{t^l}{\theta^2} - \left(\frac{t^l}{\theta}\right)^{r_j-1} \exp\left(-\frac{t^l}{\theta}\right) \frac{t^l}{\theta^2}}{(r_j - 1)! (F_E(t; r_j, \theta) - F_E(t^l; r_j, \theta))} \Bigg|_{\theta=\theta^{(h)}} \\
& + \frac{1}{\theta^{(h)2}} \sum_{j=1}^M \left(\sum_{i \in S_{i.}} i. z_{ij}^{(h)} x_i \right. \\
& \quad + \sum_{i \in S_{iii.}} iii. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)}) \\
& \quad \left. + \sum_{i \in S_{v.}} P_{1,i}^{(h)} v. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)}) \right).
\end{aligned}$$

This leads to the following M-step equation for θ :

$$\begin{aligned}
\theta^{(h)} = & \frac{1}{n_1^{(h)} \sum_{j=1}^M \beta_j^{(h)} r_j} \left(\sum_{i \in S_{i.}} x_i \right. \\
& + \sum_{i \in S_{iii.}} \sum_{j=1}^M iii. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < u_i \leq t < T; \theta^{(h-1)}) \\
& \left. + \sum_{i \in S_{v.}} \sum_{j=1}^M P_{1,i}^{(h)} v. z_{ij}^{(h)} E(X_i | Z_{ij} = 1, t^l \leq l_i < X_i \leq t < u_i \leq T; \theta^{(h-1)}) \right)
\end{aligned}$$

$$\frac{\sum_{j=1}^M \beta_j^{(h)} \frac{(t^l)^{r_j} \exp\left(-\frac{t^l}{\theta^{(h)}}\right) - t^{r_j} \exp\left(-\frac{t}{\theta^{(h)}}\right)}{(\theta^{(h)})^{r_j-1} (r_j-1)! (F_E(t; r_j, \theta^{(h)}) - F_E(t^l; r_j, \theta^{(h)}))}}{\sum_{j=1}^M \beta_j^{(h)} r_j}, \quad (28)$$

where expressions for the conditional expectations can be found in (21) and (24). This equation can be seen as the sample mean of all data points that are smaller than or equal to t , divided by the average shape parameter. For the censored data points, the sample mean is replaced by the expected value given the lower and upper bounds (second and third terms). Moreover, there is a correction for truncation (fourth term) which depends on $\theta^{(h)}$ itself in a complicated way. Therefore, it is not possible to find an analytical solution, and we solve (28) numerically using a Newton-type algorithm with the previous estimate $\theta^{(h-1)}$ as the starting value.

The expression for β_j and the M-step equation for θ are similar to the ones in Verbelen et al. (2015) where an extra term (third term) is included for data points of type v .

Then, we transform the estimates for β_j to estimates for α_j using $\tilde{\alpha}_j / \sum_{j=1}^M \tilde{\alpha}_j$ with

$$\tilde{\alpha}_j = \frac{\hat{\beta}_j}{F_E(t; r_j, \hat{\theta}) - F_E(t^l; r_j, \hat{\theta})},$$

and $\hat{\beta}_j$ and $\hat{\theta}$ the estimates obtained from the final EM-step.

A.3.3 Pareto distribution

To maximise the expected log-likelihood with respect to $\Theta_2 = \gamma$, we have to maximise

$$\begin{aligned} & \sum_{i \in S_{ii.}} E(\ln f_2(X_i; t, T, \gamma) \mid t^l < t < l_i = u_i \leq T; \gamma^{(h-1)}) \\ & + \sum_{i \in S_{iv.}} E(\ln f_2(X_i; t, T, \gamma) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)}) \\ & + \sum_{i \in S_v.} P_{2,i}^{(h)} E(\ln f_2(X_i; t, T, \gamma) \mid t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)}) \\ = & \sum_{i \in S_{ii.}} \left[-\ln(\gamma t) - \ln \left(1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}} \right) - \left(\frac{1}{\gamma} + 1 \right) \ln \left(\frac{x_i}{t} \right) \right] \\ & + \sum_{i \in S_{iv.}} \left[-\ln(\gamma t) - \ln \left(1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}} \right) \right. \\ & \quad \left. - \left(\frac{1}{\gamma} + 1 \right) E \left(\ln \left(\frac{X_i}{t} \right) \mid t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \right] \\ & + \sum_{i \in S_v.} P_{2,i}^{(h)} \left[-\ln(\gamma t) - \ln \left(1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}} \right) \right] \end{aligned}$$

$$- \left(\frac{1}{\gamma} + 1 \right) E \left(\ln \left(\frac{X_i}{t} \right) \middle| t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)} \right) \Bigg], \quad (29)$$

where $P_{2,i}^{(h)} = P \left(X_i > t \middle| t^l \leq l_i < t < u_i \leq T; \Theta^{(h-1)} \right)$. Taking the derivative of (29) with respect to γ gives

$$\begin{aligned} & - \frac{1}{\gamma} \left(\#S_{ii.} + \#S_{iv.} + \sum_{i \in S_v.} P_{2,i}^{(h)} \right) \\ & + \frac{1}{\gamma^2} \left(\#S_{ii.} + \#S_{iv.} + \sum_{i \in S_v.} P_{2,i}^{(h)} \right) \frac{\ln \left(\frac{T}{t} \right) \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}}}{1 - \left(\frac{T}{t} \right)^{-\frac{1}{\gamma}}} \\ & + \frac{1}{\gamma^2} \left(\sum_{i \in S_{ii.}} \ln \left(\frac{x_i}{t} \right) + \sum_{i \in S_{iv.}} E \left(\ln \left(\frac{X_i}{t} \right) \middle| t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \right. \\ & \quad \left. + \sum_{i \in S_v.} P_{2,i}^{(h)} E \left(\ln \left(\frac{X_i}{t} \right) \middle| t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)} \right) \right). \end{aligned}$$

Setting this derivative at $\gamma^{(h)}$ equal to 0 and then solving for $\gamma^{(h)}$ results in

$$\begin{aligned} \gamma^{(h)} &= \frac{1}{n_2^{(h)}} \left(\sum_{i \in S_{ii.}} \ln \left(\frac{x_i}{t} \right) \right. \\ & \quad + \sum_{i \in S_{iv.}} E \left(\ln \left(\frac{X_i}{t} \right) \middle| t^l < t \leq l_i < u_i \leq T; \gamma^{(h-1)} \right) \\ & \quad \left. + \sum_{i \in S_v.} P_{2,i}^{(h)} E \left(\ln \left(\frac{X_i}{t} \right) \middle| t^l \leq l_i \leq t < X_i < u_i \leq T; \gamma^{(h-1)} \right) \right) \\ & \quad + \frac{\ln \left(\frac{T}{t} \right)}{\left(\frac{T}{t} \right)^{\frac{1}{\gamma^{(h)}}} - 1}. \end{aligned}$$

where expressions for the conditional expectations can be found in (25) and (26). Note that $P_{2,i}^{(h)} = 1 - P_{1,i}^{(h)}$ and we compute the latter using (16). The first term in the equation can be seen as the sample mean of all $\ln(X_i/t)$ with $\{X_i > t\}$ which are observed for data points belonging to case *ii* and have to be replaced by their expected values for cases *iv* and *v*. Similarly to (28), the last term, which depends on $\gamma^{(h)}$, is a correction term for the upper truncation at point T and requires us to solve this equation numerically. In case there is no upper truncation, i.e. $T = +\infty$, the last term is equal to zero and we obtain an analytical solution for $\gamma^{(h)}$.

A.4 Choice of shape parameters and number of mixtures for ME distribution

To choose the shape parameters \mathbf{r} and the number of mixtures M , we follow the approach from Verbelen et al. (2016) which is a slightly modified version of the approach from Verbelen et al. (2015). Starting from a certain value for M and shapes as given in (15), they try to reduce M using a backward stepwise search where the mixture component with the smallest shape is deleted if this decreases an IC (AIC or BIC). Moreover, after the first reduction of M , M is further reduced using the IC and the shapes \mathbf{r} are adjusted based on maximising the likelihood. For each value for M and \mathbf{r} , the EM algorithm described above is executed. We refer to Section 4 in Verbelen et al. (2016) for more details. Important is that we now consider the (log-)likelihood including the Pareto part, as given in (6).

B Fitting the ME-Pareto model to uncensored data using the EM algorithm

In case there are no censored data points, calculations are much simpler. Now, we only have data points from cases i and ii .

B.1 Starting values

We use the starting values for θ and β from Verbelen et al. (2016), see Section A.1. Since the obtained estimates are the same in each step, we do not need starting values for π and γ .

B.2 π

Maximisation w.r.t. π gives

$$\pi^{(h)} = \frac{\#S_i}{n} = \frac{n_1}{n}. \quad (30)$$

The estimate for π is thus equal to the proportion of points that is smaller than or equal to the splicing point t . It is clear that $\pi^{(h)}$ is constant since it is independent of h .

Taking the splicing point equal to the $(k+1)$ th largest data point, i.e. $t = x_{n-k,n}$, gives

$$\pi^{(h)} = 1 - \frac{k}{n}$$

when there are no ties.

B.3 ME distribution

Similar to before, the following expression is obtained, for $j = 1, \dots, M$,

$$\beta_j^{(h)} = \frac{\sum_{i \in S_{i.}} z_{ij}^{(h)}}{n_1}.$$

The M-step equation for θ now simplifies to

$$\theta^{(h)} = \frac{\sum_{i \in S_{i.}} x_i}{n_1 \sum_{j=1}^M \beta_j^{(h)} r_j} - \frac{\sum_{j=1}^M \beta_j^{(h)} \frac{(t^l)^{r_j} \exp\left(-\frac{t^l}{\theta^{(h)}}\right) - t^{r_j} \exp\left(-\frac{t}{\theta^{(h)}}\right)}{(\theta^{(h)})^{r_j-1} (r_j-1)! (F_E(t; r_j, \theta^{(h)}) - F_E(t^l; r_j, \theta^{(h)}))}}{\sum_{j=1}^M \beta_j^{(h)} r_j}. \quad (31)$$

As before, a Newton-type algorithm is employed to solve (31) numerically using the previous estimate $\theta^{(h-1)}$ as starting value.

B.4 Pareto distribution

For the Pareto parameter γ we get following implicit equation

$$\gamma^{(h)} = \frac{\sum_{i \in S_{ii.}} \ln\left(\frac{x_i}{t}\right)}{\#S_{ii.}} + \frac{\ln\left(\frac{T}{t}\right)}{\left(\frac{T}{t}\right)^{\frac{1}{\gamma^{(h)}}} - 1} = \frac{\sum_{i \in S_{ii.}} \ln\left(\frac{x_i}{t}\right)}{n_2} + \frac{\ln\left(\frac{T}{t}\right)}{\left(\frac{T}{t}\right)^{\frac{1}{\gamma^{(h)}}} - 1}. \quad (32)$$

It is immediately clear that this estimate does not depend on h , so we denote $\hat{\gamma} = \gamma^{(h)}$. The first term of (32) is the Hill estimator (Hill, 1975) with threshold t , and the second term is a correction for the upper truncation at T which vanishes for $T = +\infty$. Setting t equal to the $(k+1)$ th largest data point $x_{n-k,n}$ gives

$$\hat{\gamma} = \frac{1}{k} \sum_{i=1}^k \ln\left(\frac{x_{n-i+1,n}}{x_{n-k,n}}\right) + \frac{\ln\left(\frac{T}{x_{n-k,n}}\right)}{\left(\frac{T}{x_{n-k,n}}\right)^{\frac{1}{\hat{\gamma}}} - 1} = H_{k,n} + \frac{\ln\left(\frac{T}{x_{n-k,n}}\right)}{\left(\frac{T}{x_{n-k,n}}\right)^{\frac{1}{\hat{\gamma}}} - 1}. \quad (33)$$

B.5 Upper endpoint estimation

Aban et al. (2006) show that the conditional MLE for the endpoint T , if it is unknown, is given by the maximum $x_{n,n}$. The corresponding conditional MLE for γ is given by

$$\hat{\gamma} = H_{k,n} + \frac{\ln\left(\frac{x_{n,n}}{x_{n-k,n}}\right)}{\left(\frac{x_{n,n}}{x_{n-k,n}}\right)^{\frac{1}{\hat{\gamma}}} - 1}.$$

This estimator corresponds to (33) where T is estimated by its conditional MLE $x_{n,n}$. Beirlant et al. (2016) use the same estimator $\hat{\gamma}$, but re-estimate the endpoint T as

$$\hat{T} = \max\left\{x_{n-k,n} \left(1 + \frac{k}{n\hat{D}_T}\right)^{\hat{\gamma}}, x_{n,n}\right\},$$

where

$$\hat{D}_T = \max \left\{ \frac{k \left(\frac{x_{n-k,n}}{x_{n,n}} \right)^{\frac{1}{\hat{\gamma}}} - \frac{1}{k+1}}{n \left(1 - \left(\frac{x_{n-k,n}}{x_{n,n}} \right)^{\frac{1}{\hat{\gamma}}} \right)}, 0 \right\}$$

is an estimator for the odds ratio $D_T = \frac{1-F_2^*(T)}{F_2^*(T)}$.