

# Left-ordered inp-minimal groups

Jan Dobrowolski and John Goodrick

## Abstract

We prove that any left-ordered inp-minimal group is abelian, and we provide an example of a non-abelian left-ordered group of dp-rank 2.

## 0 Introduction and preliminaries

One of the model-theoretic properties that gained a lot of interest recently is dp-minimality, which, on one hand, significantly strengthens NIP, and on the other hand, is satisfied by all strongly minimal theories, all (weakly) o-minimal theories, algebraically closed valued fields (more generally, by all C-minimal structures), and the valued field of p-adics. Several interesting results were obtained for dp-minimal structures in the algebraic contexts of groups and fields (sometimes with additional structure), see for example [9, 4, 3, 7].

Throughout this note, we work in the context of a complete first-order theory  $T$ , and “formula” means a first-order formula in the language of  $T$ .

**Definition 0.1.** 1. An inp-pattern of depth  $\kappa$  (in the partial type  $\pi(\bar{x})$ ) is a sequence  $\langle \varphi_i(\bar{x}; \bar{y}_i) : i < \kappa \rangle$  of formulas and an array  $\{\bar{a}_{ij} : i < \kappa, j < \omega\}$  of parameters (from some model of  $T$ ) such that:

- (a) For each  $i < \kappa$ , there is some  $k_i < \omega$  such that  $\{\varphi_{i,j}(\bar{x}; \bar{a}_{i,j}) : j < \omega\}$  is  $k_i$ -inconsistent; and
- (b) For each  $\eta : \kappa \rightarrow \omega$ , the partial type

$$\pi(\bar{x}) \cup \{\varphi_i(\bar{x}; \bar{a}_{i,\eta(i)}) : i < \kappa\}$$

is consistent.

- 2. The inp-rank (or burden) of a partial type  $\pi(\bar{x})$  is the maximal  $\kappa$  such that there is an inp-pattern of depth  $\kappa$  in  $\pi(\bar{x})$ . The inp-rank of  $T$  is the inp-rank of  $x = x$ , and  $T$  is inp-minimal if its inp-rank is 1.
- 3. An ict-pattern of depth  $\kappa$  (in the partial type  $\pi(\bar{x})$ ) is a sequence  $\langle \varphi_i(\bar{x}; \bar{y}_i) : i < \kappa \rangle$  of formulas and an array  $\{\bar{a}_{ij} : i < \kappa, j < \omega\}$  of parameters (from some model of  $T$ ) such that for each  $\eta : \kappa \rightarrow \omega$ , the partial type

$$\pi(\bar{x}) \cup \{\varphi_i(\bar{x}; \bar{a}_{i,\eta(i)}) : i < \kappa\} \cup \{\neg\varphi_i(\bar{x}; \bar{a}_{i,j}) : i < \kappa, j \neq \eta(i)\}$$

is consistent.

- 4. The dp-rank of a partial type  $\pi(\bar{x})$  is the maximal  $\kappa$  such that there is an ict-pattern of depth  $\kappa$  in  $\pi(\bar{x})$ . The dp-rank of  $T$  is the dp-rank of  $x = x$ , and  $T$  is dp-minimal if its dp-rank is 1.

In spite of its name, dp-rank is really more like a cardinal-valued *dimension* than an ordinal-valued rank such as  $SU(p)$ , and in the context of stable theories, dp-minimality is equivalent to every nonalgebraic 1-type having weight 1, as observed in [8]. It turns out that a theory is dp-minimal just in case it is both inp-minimal and NIP (see [1]).

One of the context investigated in [9] is that of (bi)-ordered groups.

**Definition 0.2.** A left-ordering on a group  $(G, \cdot)$  is a total ordering  $<$  on  $G$  such that for any  $f, g, h \in G$ , whenever  $g < h$ , we have that  $f \cdot g < f \cdot h$ . A right-ordering is defined similarly, and a bi-ordering on  $G$  is an ordering which is simultaneously a left-ordering and a right-ordering.

While Pierre Simon claimed that all inp-minimal “ordered groups” are abelian [9], his proof only applies to groups with a definable bi-ordering: his argument uses the fact that for any  $x, y$  in a bi-orderable group and any positive  $n \in \mathbb{N}$ , if  $x^n = y^n$  then  $x = y$ . But in left-orderable groups (such as in the example of the Klein bottle group below), one may have that  $x^2 = y^2$  but  $x \neq y$ .

In this note, we prove that every inp-minimal left-ordered group is abelian, strengthening the result from [9], and we also show that the result fails already in the dp-rank 2 case, by providing a suitable example.

## 1 A non-abelian left-ordered group of dp-rank 2

In this section, we define the “Klein bottle group” (the fundamental group of a Klein bottle) which is presented as  $G = \langle x, y : x^{-1}yx = y^{-1} \rangle$ . In other words,  $y^{-1}x = xy$ , and routine algebraic manipulation shows:

1. Every  $g \in G$  can be uniquely written as  $g = x^n y^m$  for some  $n, m \in \mathbb{Z}$ ,
2.  $(x^n y^m) \cdot (x^{n'} y^{m'}) = x^{n+n'} y^{m'+(-1)^{n'} m}$ , and
3.  $(x^n y^m)^{-1} = x^{-n} y^{(-1)^{n+1} m}$ .

We can define a left ordering  $\leq$  on  $G$  lexicographically on the exponents:  $x^n y^m \leq x^{n'} y^{m'}$  iff either  $n < n'$  or else  $n = n'$  and  $m \leq m'$ . The subgroup generated by  $y$  is the minimal nontrivial convex subgroup of  $G$ , and the order type of  $G$  is  $\mathbb{Z} \times \mathbb{Z}$ .

We note in passing that while  $G$  is non-abelian, it is abelian-by-*finite*: a simple calculation shows that the centralizer  $C(y)$  of  $y$  is  $\{x^{2n} y^m : n, m \in \mathbb{Z}\}$ , which is abelian, and for any  $g \in G$ , we have  $g^2 \in C(y)$ .

**Proposition 1.1.** *The structure  $(G, \cdot, \leq)$  is dp-rank 2.*

*Proof.* To show that it is NIP and dp-rank *at most* 2, it suffices (thanks to the additivity of the dp-rank proved in [5]) to note that an isomorphic copy of  $G$  is definable in the dp-minimal structure  $(\mathbb{Z}, <, +)$  with  $\mathbb{Z} \times \mathbb{Z}$  as the universe of the group; the definition of the group operation depends on the parity of one of the coordinates, but of course  $2\mathbb{Z}$  is a definable subgroup of  $\mathbb{Z}$ .

Now we compute the centralizer  $C(x)$  of the generator  $x$ . For any  $a, b \in \mathbb{Z}$ , we have

$$x \cdot (x^a y^b) = x^{a+1} y^b$$

and

$$(x^a y^b) \cdot x = x^{a+1} y^{-b},$$

so we conclude that  $C(x) = \langle x \rangle$ .

For any  $n \in \mathbb{N}$ , there are pairwise disjoint intervals  $\langle I_{n,k} : k < \omega \rangle$  such that for any  $k$ ,  $x^k \in I_{n,k}$  and  $I_{n,k}$  intersects every right coset  $C(x), C(x)y, \dots, C(x)y^n$ . So by compactness, in an  $\omega$ -saturated extension of  $G$ , we can find pairwise disjoint intervals  $\langle I_k : k < \omega \rangle$  such that  $x^k \in I_k$  and each  $I_k$  intersects every right coset  $C(x)y^n$ . Therefore the formulas expressing  $z \in I_k$  and  $z \in C(x)y^j$  (in the free variable  $z$ ) give an inp-pattern of depth 2, so  $G$  is not inp-minimal, hence by NIP it is not dp-minimal. □

**Question 1.1.** *Is the group  $G$  above dp-minimal in the pure language of groups?*

## 2 Inp-minimal left-ordered groups

In this section, we prove that every inp-minimal left-ordered group is abelian. For a left-ordered group  $G$  and a subset  $A \subseteq G$ , by  $h(A)$  we will denote the convex hull of  $A$  in  $G$ .

**Fact 2.1** ([9]). *Let  $G$  be an inp-minimal group. Then there is a definable normal abelian subgroup  $H$  of  $G$  such that  $G/H$  has finite exponent.*

Note that in a left-ordered group  $G$  a convex hull of a subgroup  $H$  need not be a subgroup of  $G$ , but still it is a union of right  $H$ -cosets. Thus, by repeating the proof of Lemma 3.2 from [9], we obtain:

**Fact 2.2.** *Let  $G$  be an inp-minimal left-ordered group. Let  $H$  be a definable subgroup of  $G$  and let  $C$  be the convex hull of  $H$ . Then  $C$  is a union of finitely many right  $H$ -cosets.*

We will also use a group-theoretic fact about FC-groups.

**Definition 2.3.** *An FC-group is a group in which the centralizer of every element has finite index.*

Note that if  $[G : Z(G)]$  is finite, then  $G$  is an FC-group. The following is Theorem 6.24 from [6]:

**Fact 2.4.** *Every torsion-free FC-group is abelian.*

**Theorem 2.5.** *Every left-ordered inp-minimal group is abelian.*

*Proof.* By Fact 2.4, it is enough to show that  $[G : Z(G)]$  is finite. Let  $H$  be given by Fact 2.1. Notice that  $G = h(H)$ : if  $a \in G$ , say  $a > e$ , then for  $l$  equal to the exponent of  $G/H$ , we have  $e < a < a^2 < \dots < a^l \in H$ , so  $a \in h(H)$ . Hence, by Fact 2.2, we get that  $n := [G : H]$  is finite.

**Claim 1.** *For any positive  $x \in G$ , the interval  $[e, x]$  is covered by finitely many right cosets of a central subgroup of  $G$ .*

*Proof of Claim 1.* We can assume that  $G$  is non-trivial (hence it is infinite so also  $H$  is non-trivial). Notice that any coset  $Hg$  in  $G$  has a positive representative (if  $g$  is negative then one can take  $g^{-(l-1)}$  as such a representative). It follows (thanks to the normality of  $H$ ) that for any  $y, g \in G$  we can find an element  $z \in G$  such that  $Hg = Hz$  and  $z > y$  (by choosing  $z := yw$ , where  $w$  is a positive element of  $Hy^{-1}g$ ).

Now, fix any positive  $x = x_0 \in G$ . Using the above observation, we can choose  $x_0 < x_1 < \dots < x_{n-1} \in G$  such that  $G = \bigcup_{i < n} Hx_i$ . Since  $[G : H]$  is finite, and, by Fact 2.2,  $h(C(x_i))$  is covered by finitely many right  $C(x_i)$ -cosets for each  $i < n$ , we get that

$$C := \bigcap_{i < n} h(C(x_i))$$

is covered by finitely many right cosets of

$$A := H \cap \bigcap_{i < n} C(x_i)$$

(we use here that an intersection of right cosets of subgroups  $G_0, \dots, G_{k-1} < G$  is a right coset of  $\bigcap_{i < k} G_i$ ). But, since  $H$  is abelian, and  $G$  is generated by  $H, x_0, \dots, x_{n-1}$ , we have that  $A \subseteq Z(G)$ . Also, since  $x_0 < x_1 < \dots, x_{n-1}$  and  $\forall_{i < n} x_i \in C(x_i)$ , we get that  $x = x_0 \in C$ , so, by convexity of  $C$ ,  $[e, x_0] \subseteq C$ , which proves the claim.  $\square$

Now, suppose for a contradiction that  $[G : Z(G)]$  is infinite. Note that if some coset  $Z(G)g$  contains only negative elements, then the coset  $Z(G)g^{-1}$  contains only positive elements, so in any case we may choose infinitely many positive representatives  $y_0, y_1, y_2, \dots$  of pairwise distinct right cosets of  $Z(G)$  in  $G$ . Without loss of generality,  $G$  is  $\omega$ -saturated, and there is an element  $x \in G$  greater than all the  $y_i$ 's. Then  $[e, x]$  cannot be covered by finitely many right cosets of  $Z(G)$ , so it cannot be covered by finitely many right cosets of any central subgroup of  $G$ , contradicting the claim.  $\square$

**Corollary 2.6.** *If  $(G, \cdot, <)$  is a left-ordered group which is inp-minimal (in the pure language of ordered groups), then it is dp-minimal.*

*Proof.* By Theorem 2.5,  $G$  is abelian, and any ordered abelian group is NIP, as shown in [2]; since NIP and inp-minimality imply dp-minimality, we are done.  $\square$

## References

- [1] H. Adler, *Strong theories, burden, and weight*, preprint, 2007. arXiv:1507.0391
- [2] Y. Gurevich and P. Schmitt, *The theory of ordered abelian groups does not have the independence property*, Trans. Amer. Math. Soc., 284:171-182, 1984. arXiv:1507.0391
- [3] F. Jahnke, P. Simon, and E. Walsberg, *Dp-minimal valued fields*, 2015. arXiv:1507.0391
- [4] W. Johnson, *On dp-minimal fields*, 2015. arXiv:1507.0274
- [5] I. Kaplan, A. Onshuus, and A. Usvyatsov, *Additivity of the dp-rank*, Trans. Amer. Math. Soc., 365(11):5783-5804, 2013
- [6] T.Y. Lam, *A First Course in Noncommutative Rings*, Springer-Verlag, 1991.
- [7] E. Levi, I. Kaplan, P. Simon *Some remarks on dp-minimal groups*, 2016. arXiv:1605.07867
- [8] A. Onshuus and A. Usvyatsov, *On dp-minimality, strong dependence, and weight*, J. Symb. Log. 76-3, 2011.
- [9] P. Simon, *On dp-minimal ordered structures*, J. Symb. Log., Volume 76-2, 2011.

Jan Dobrowolski

Department of Mathematics, Yonsei University South Korea

e-mail: dobrowol@math.uni.wroc.pl

John Goodrick

Departamento de Matemáticas, Universidad de los Andes, Bogotá, Colombia

e-mail: jr.goodrick427@uniandes.edu.co