

## SPM BULLETIN

## ISSUE 39 (July '16)

## 1. EDITOR'S NOTE

With the approaching TOPOSYM'16 (<http://www.toposym.cz/programme.php>), it is a pleasure to see how the topic of selection principles gains increasing attention and becomes a standard part of topology and set theory. At least eight of the 28 speakers, and a good number of the contributed lecture speakers, made substantial contributions to this topic in their career. For some of these, SPs constitute the main topic of research in the last few years. This is in accordance with the continuous progress on the topic, some of which mentioned below.

With best regards,

*Boaz Tsaban*, [tsaban@math.biu.ac.il](mailto:tsaban@math.biu.ac.il)  
<http://www.cs.biu.ac.il/~tsaban>

## 2. LONG ANNOUNCEMENTS

**2.1. A characterization of barrelledness of  $C_p(X)$ .** We prove that, for a Tychonoff space  $X$ , the space  $C_p(X)$  is barrelled if and only if it is a Mackey group.

<http://arxiv.org/abs/1512.00788>  
*S. Gabrielyan*

**2.2. Countable tightness and  $\mathfrak{G}$ -bases on Free topological groups.** Given a Tychonoff space  $X$ , let  $F(X)$  and  $A(X)$  be respectively the free topological group and the free Abelian topological group over  $X$  in the sense of Markov. In this paper, we discuss two topological properties in  $F(X)$  or  $A(X)$ , namely the countable tightness and  $\mathfrak{G}$ -base. We provide some characterizations of the countable tightness and  $\mathfrak{G}$ -base of  $F(X)$  and  $A(X)$  for various special classes of spaces  $X$ . Furthermore, we also study the countable tightness and  $\mathfrak{G}$ -base of some level of  $F(X)$ . Some open problems in [?] are partially answered.

<http://arxiv.org/abs/1512.07515>  
*Fucai Lin, Alex Ravsky, Jing Zhang*

**2.3. Ideal equal Baire classes.** For any Borel ideal we characterize ideal equal Baire system generated by the families of continuous and quasi-continuous functions, i.e., the families of ideal equal limits of sequences of continuous and quasi-continuous functions.

<http://arxiv.org/abs/1601.01010>  
*Adam Kwela and Marcin Staniszewski*

**2.4. On questions which are connected with Talagrand problem.** We prove the following results.

- (1) If  $X$  is a  $\alpha$ -favourable space,  $Y$  is a regular space, in which every separable closed set is compact, and  $f : X \times Y \rightarrow \mathbb{R}$  is a separately continuous everywhere jointly discontinuous function, then there exists a subspace  $Y_0 \subseteq Y$  which is homeomorphic to  $\beta\mathbb{N}$ .
- (2) There exist a  $\alpha$ -favourable space  $X$ , a dense in  $\beta\mathbb{N} \setminus \mathbb{N}$  countably compact space  $Y$  and a separately continuous everywhere jointly discontinuous function  $f : X \times Y \rightarrow \mathbb{R}$ .

Besides, it was obtained some conditions equivalent to the fact that the space  $C_p(\beta\mathbb{N} \setminus \mathbb{N}, \{0, 1\})$  of all continuous functions  $x : \beta\mathbb{N} \setminus \mathbb{N} \rightarrow \{0, 1\}$  with the topology of point-wise convergence is a Baire space.

<http://arxiv.org/abs/1601.03163>

*Volodymyr V. Mykhaylyuk*

**2.5. Namioka spaces and strongly Baire spaces.** A notion of strongly Baire space is introduced. Its definition is a transfinite development of some equivalent reformulation of the Baire space definition. It is shown that every strongly Baire space is a Namioka space and every  $\beta - \sigma$ -unfavorable space is a strongly Baire space.

<http://arxiv.org/abs/1601.05884>

*Volodymyr V. Mykhaylyuk*

**2.6. Comparing Fréchet-Urysohn filters with two pre-orders.** A filter  $\mathcal{F}$  on  $\omega$  is called Fréchet-Urysohn if the space with only one non-isolated point  $\omega \cup \{\mathcal{F}\}$  is a Fréchet-Urysohn space, where the neighborhoods of the non-isolated point are determined by the elements of  $\mathcal{F}$ . In this paper, we distinguish some Fréchet-Urysohn filters by using two pre-orderings of filters: One is the Rudin-Keisler pre-order and the other one was introduced by Todorčević-Uzcátegui in [?]. In this paper, we construct an RK-chain of size  $\mathfrak{c}^+$  which is RK-above of every FU-filter. Also, we show that there is an infinite RK-antichain of FU-filters.

<http://arxiv.org/abs/1602.06227>

*S. Garcia-Ferreira, J. E. Rivera-Gómez*

**2.7. Selective game versions of countable tightness with bounded finite selections.** For a topological space  $X$  and a point  $x \in X$ , consider the following game – related to the property of  $X$  being countably tight at  $x$ . In each inning  $n \in \omega$ , the first player chooses a set  $A_n$  that clusters at  $x$ , and then the second player picks a point  $a_n \in A_n$ ; the second player is the winner if and only if  $x \in \overline{\{a_n : n \in \omega\}}$ . In this work, we study variations of this game in which the second player is allowed to choose finitely many points per inning rather than one, but in which the number of points they are allowed to choose in each inning has been fixed in advance. Surprisingly, if the number of points allowed per inning is the same throughout the play, then all of the games obtained in this fashion are distinct. We also show that a new game is obtained if the number of points the second player is allowed to pick increases at each inning.

<http://arxiv.org/abs/1603.09715>

*Leandro F. Aurichi, Angelo Bella, Rodrigo R. Dias*

**2.8. Gruff Ultrafilters.** We investigate the question of whether  $\mathbb{Q}$  carries an ultrafilter generated by perfect sets (such ultrafilters were called gruff ultrafilters by van Douwen). We prove that one can (consistently) obtain an affirmative answer to this question in three different ways: by assuming a certain parametrized diamond principle, from the cardinal invariant equality  $\mathfrak{d} = \mathfrak{c}$ , and in the Random real model.

<http://arxiv.org/abs/1604.02473>

*David Fernández-Bretón and Michael Hrušák*

**2.9. Free locally convex spaces with a small base.** The paper studies the free locally convex space  $L(X)$  over a Tychonoff space  $X$ . Since for infinite  $X$  the space  $L(X)$  is never metrizable (even not Fréchet-Urysohn), a possible applicable generalized metric property for  $L(X)$  is welcome. We propose a concept (essentially weaker than first-countability) which is known under the name a  $\mathfrak{G}$ -base. A space  $X$  has a  $\mathfrak{G}$ -base if for every  $x \in X$  there is a base  $\{U_\alpha : \alpha \in \mathbb{N}^{\mathbb{N}}\}$  of neighborhoods at  $x$  such that  $U_\beta \subseteq U_\alpha$  whenever  $\alpha \leq \beta$  for all  $\alpha, \beta \in \mathbb{N}^{\mathbb{N}}$ , where  $\alpha = (\alpha(n))_{n \in \mathbb{N}} \leq \beta = (\beta(n))_{n \in \mathbb{N}}$  if  $\alpha(n) \leq \beta(n)$  for all  $n \in \mathbb{N}$ . We show that if  $X$  is an Ascoli  $\sigma$ -compact space, then  $L(X)$  has a

$\mathfrak{G}$ -base if and only if  $X$  admits an Ascoli uniformity  $\mathcal{U}$  with a  $\mathfrak{G}$ -base. We prove that if  $X$  is a  $\sigma$ -compact Ascoli space of  $\mathbb{N}^{\mathbb{N}}$ -uniformly compact type, then  $L(X)$  has a  $\mathfrak{G}$ -base. As an application we show: (1) if  $X$  is a metrizable space, then  $L(X)$  has a  $\mathfrak{G}$ -base if and only if  $X$  is  $\sigma$ -compact, and (2) if  $X$  is a countable Ascoli space, then  $L(X)$  has a  $\mathfrak{G}$ -base if and only if  $X$  has a  $\mathfrak{G}$ -base.

<http://arxiv.org/abs/1604.02555>

*Saak Gabrielyan and Jerzy Kakol*

**2.10. On separability of the functional space with the open-point and bi-point-open topologies, II.** In this paper we continue to study the property of separability of functional space  $C(X)$  with the open-point and bi-point-open topologies.

<http://arxiv.org/abs/1604.04609>

*Alexander V. Osipov*

**2.11. On sequential separability of functional spaces.** In this paper, we give necessary and sufficient conditions for the space  $B_1(X)$  of first Baire class functions on a Tychonoff space  $X$ , with pointwise topology, to be (strongly) sequentially separable.

<http://arxiv.org/abs/1604.05116>

*Alexander V. Osipov, Evgenii G. Pytkeev*

**2.12. No interesting sequential groups.** We prove that it is consistent with ZFC that no sequential topological groups of intermediate sequential orders exist. This shows that the answer to a 1981 question of P. Nyikos is independent of the standard axioms of set theory. The model constructed also provides consistent answers to several questions of D. Shakhmatov, S. Todorčević and Uzcátegui. In particular, we show that it is consistent with ZFC that every countably compact sequential group is Fréchet-Urysohn.

<http://arxiv.org/abs/1604.08868>

*Alexander Shibakov*

**2.13. On sequential analytic groups.** Alexander Shibakov, <http://arxiv.org/abs/1604.08872>

We answer a question of S. Todorčević and C. Uzcátegui from [?] by showing that the only possible sequential orders of sequential analytic groups are 1 and  $\omega_1$ . Other results on the structure of sequential analytic spaces and their relation to other classes of spaces are given as well. In particular, we provide a full topological classification of sequential analytic groups by showing that all such groups are either metrizable or  $k_\omega$ -spaces, which, together with a result by Zelenyuk, implies that there are exactly  $\omega_1$  non homeomorphic analytic sequential group topologies.

**2.14. On large sequential groups.** We construct, using  $\diamond$ , an example of a sequential group  $G$  such that the only countable sequential subgroups of  $G$  are closed and discrete, and the only quotients of  $G$  that have a countable pseudocharacter are countable and Fréchet. We also show how to construct such a  $G$  with several additional properties (such as make  $G^2$  sequential, and arrange for every sequential subgroup of  $G$  to be closed and contain a nonmetrizable compact subspace, etc.). Several results about  $k_\omega$  sequential groups are proved. In particular, we show that each such group is either locally compact and metrizable or contains a closed copy of the sequential fan. It is also proved that a dense proper subgroup of a non Fréchet  $k_\omega$  sequential group is not sequential extending a similar observation of T. Banach about countable  $k_\omega$  groups.

<http://arxiv.org/abs/1604.08874>

*Alexander Shibakov*

**2.15. The Ascoli property for function spaces.** The paper deals with Ascoli spaces  $C_p(X)$  and  $C_k(X)$  over Tychonoff spaces  $X$ . The class of Ascoli spaces  $X$ , i.e. spaces  $X$  for which any compact subset  $K$  of  $C_k(X)$  is evenly continuous, essentially includes the class of  $k_{\mathbb{R}}$ -spaces. First we prove that if  $C_p(X)$  is Ascoli, then it is  $\kappa$ -Fréchet-Urysohn. If  $X$  is cosmic, then  $C_p(X)$  is Ascoli iff it is  $\kappa$ -Fréchet-Urysohn. This leads to the following extension of a result of Morishita: If for a Čech-complete space  $X$  the space  $C_p(X)$  is Ascoli, then  $X$  is scattered. If  $X$  is scattered and stratifiable, then  $C_p(X)$  is an Ascoli space. Consequently: (a) If  $X$  is a complete metrizable space, then  $C_p(X)$  is Ascoli iff  $X$  is scattered. (b) If  $X$  is a Čech-complete Lindelöf space, then  $C_p(X)$  is Ascoli iff  $X$  is scattered iff  $C_p(X)$  is Fréchet-Urysohn. Moreover, we prove that for a paracompact space  $X$  of point-countable type the following conditions are equivalent: (i)  $X$  is locally compact. (ii)  $C_k(X)$  is a  $k_{\mathbb{R}}$ -space. (iii)  $C_k(X)$  is an Ascoli space. The Ascoli spaces  $C_k(X, [0, 1])$  are also studied.

<http://arxiv.org/abs/1606.01013>

*Saak Gabrielyan, Jan Grebík, Jerzy Kakol, Lyubomyr Zdomskyy*

**2.16. Products of Menger spaces: a combinatorial approach.** We construct Menger subsets of the real line whose product is not Menger in the plane. In contrast to earlier constructions, our approach is purely combinatorial, and the set theoretic hypotheses used are either milder than or incompatible with earlier ones. On the other hand, we establish productive properties for versions of Menger's property parameterized by filters and semifilters. In particular, the hypothesis  $\mathfrak{b} = \mathfrak{d}$  implies that every productively Menger set of real numbers is productively Hurewicz, and each ultrafilter version of Menger's property is strictly between Menger's and Hurewicz's classic properties. We include a number of open problems emerging from this study.

<http://arxiv.org/abs/1603.03361>

*Piotr Szewczak, Boaz Tsaban*

**2.17. Products of Menger spaces, II: general spaces.** We study products of general topological spaces with Menger's covering property, and its refinements based on filters and semifilters. Among other results, we prove that, assuming the Continuum Hypothesis, every productively Lindelöf space is productively Menger, and every productively Menger space is productively Hurewicz. None of these implications is reversible.

<http://arxiv.org/abs/1607.01687>

*Piotr Szewczak, Boaz Tsaban*

**2.18. Topological spaces with a local  $\omega^\omega$ -base have the strong Pytkeev\* property.** Modifying the known definition of a Pytkeev network, we introduce a notion of Pytkeev\* network and prove that a topological space has a countable Pytkeev network at a point  $x \in X$  if and only if  $X$  is countably tight at  $x$  and has a countable Pytkeev\* network at  $x$ . We define a topological space  $X$  to have the strong Pytkeev\* property if  $X$  has a Pytkeev\* network at each point. Our main theorem says that a topological space  $X$  has a countable Pytkeev\* network at point  $x \in X$  if  $X$  has a local  $\omega^\omega$ -base at  $x$  (i.e., a neighborhood base  $(U_\alpha)_{\alpha \in \omega^\omega}$  at  $x$  such that  $U_\beta \subset U_\alpha$  for all  $\alpha \leq \beta$  in  $\omega^\omega$ ). Consequently, each countably tight space with a local  $\omega^\omega$ -base has the strong Pytkeev property.

<http://arxiv.org/abs/1607.03599>

*Taras Banach*

### 3. SHORT ANNOUNCEMENTS

**3.1. Luzin  $\pi$ -bases and operation foliage hybrid.** Mikhail Patrakeev, <http://arxiv.org/abs/1512.02458>

- 3.2. **Separately continuous functions on products and its dependence on  $\aleph$  coordinates.** Volodymyr V. Mykhaylyuk, <http://arxiv.org/abs/1601.02343>
- 3.3. **Reversible filters.** Alan Dow, Rodrigo Hernández-Gutiérrez, <http://arxiv.org/abs/1601.04081>
- 3.4. **The Bi-Compact-Open Topology on  $C(X)$ .** A. Jindal, R. A. McCoy, S. Kundu, <http://arxiv.org/abs/1603.05366>
- 3.5. **Completeness and compactness properties in metric spaces, topological groups and function spaces.** Alejandro Dorantes-Aldama, Dmitri Shakhmatov, <http://arxiv.org/abs/1603.07459>
- 3.6. **Free topological vector spaces.** Saak S. Gabrielyan and Sidney A. Morris, <http://arxiv.org/abs/1604.04005>
- 3.7. **Continuous extension of functions from countable sets.** V. Mykhaylyuk, <http://arxiv.org/abs/1604.06178>
- 3.8. **The left side of Cichoń's diagram.** Martin Goldstern; Diego Alejandro Mejia; Saharon Shelah, <http://www.ams.org/journal-getitem?pii=S0002-9939-2016-13161-4>
- 3.9. **A primitive associated to the Cantor-Bendixson derivative on the real line.** Borys Álvarez-Samaniego and Andrés Merino, <http://arxiv.org/abs/1605.00853>
- 3.10. **On Borel semifilters.** Andrea Medini, <http://arxiv.org/abs/1605.01024>
- 3.11. **Every filter is homeomorphic to its square.** Andrea Medini, Lyubomyr Zdomskyy, <http://arxiv.org/abs/1605.04087>
- 3.12.  **$\mathfrak{G}$ -bases in free (locally convex) topological vector spaces.** Taras Banakh and Arkady Leiderman, <http://arxiv.org/abs/1606.01967>
- 3.13. **Completeness Properties of the open-point and bi-point-open topologies on  $C(X)$ .** Anubha Jindal, R. A. McCoy, S. Kundu and Varun Jindal, <http://arxiv.org/abs/1607.01491>
- 3.14. **On  $\sigma$ -countably tight spaces.** István Juhász and Jan van Mill, <http://arxiv.org/abs/1607.00517>

---

Previous issues. <http://front.math.ucdavis.edu/search?&t=%22SPM+Bulletin%22>

Contributions and free subscription. Email [tsaban@math.biu.ac.il](mailto:tsaban@math.biu.ac.il).