

Bayesian mode and maximum estimation and accelerated rates of contraction

William Weimin Yoo and Subhashis Ghosal
Leiden University and North Carolina State University

Abstract

We study the problem of estimating the mode and maximum of an unknown regression function in the presence of noise. We adopt the Bayesian approach by using tensor product B-splines and endowing the coefficients with Gaussian priors. In the usual fixed-in-advanced sampling plan, we establish posterior contraction rates for mode and maximum and show that they coincide with the minimax rates for this problem. To quantify estimation uncertainty, we construct credible sets for these two quantities that have high coverage probabilities with optimal sizes. If one is allowed to collect data sequentially, we further propose a Bayesian two-stage estimation procedure, where a second stage posterior is built based on samples collected within a credible set arising from a first stage posterior. Under appropriate conditions on the radius of this credible set, we can accelerate optimal contraction rates in the fixed-in-advanced setting to the minimax sequential rates. A simulation experiment shows that our Bayesian two-stage procedure outperforms single-stage procedure and also slightly improves upon a non-Bayesian two-stage procedure, due to shrinkage effect of our Bayesian estimators.

Keywords: Mode, Maximum value, Posterior contraction, Credible set, Nonparametric regression, Tensor product B-splines, Anisotropic Hölder space, Sequential, Two-stage.

MSC2010 Classification: Primary 62G05, 62L12; secondary 62G08, 62G15, 62L05

1 Introduction

Consider noisy measurements Y_1, \dots, Y_n of an unknown smooth function f at locations $\mathbf{X}_1, \dots, \mathbf{X}_n \in [0, 1]^d$ given by the nonparametric regression model

$$Y_i = f(\mathbf{X}_i) + \varepsilon_i, \quad i = 1, \dots, n, \quad (1.1)$$

where the regression errors $\varepsilon_1, \dots, \varepsilon_n$ are modeled as independent and identically distributed (i.i.d.) $N(0, \sigma^2)$ with unknown standard deviation $0 < \sigma < \infty$. The covariates can be deterministic or be drawn as i.i.d samples from some fixed distribution independently of the regression errors.

In this paper, we consider the problem of estimating the mode $\boldsymbol{\mu}$ which marks the location of the maximum of f , and the value of this maximum $M = f(\boldsymbol{\mu}) = \sup\{f(\mathbf{x}) : \mathbf{x} \in [0, 1]^d\}$, assuming that $\boldsymbol{\mu}$ is unique. The problem can be thought of as optimization in the presence of noise and has wide range of applications. For instance, searching for the optimal factor configurations in response surface methodology, locating peaks in bacteria (Silverman [27])

and human (Müller [21]) growth curves, or to classify and compare curves arising from longitudinal endocrinological data (Jørgensen *et al.* [15]).

The problem of estimating the mode and the maximum of an isotropic regression function with smoothness α is well studied in the frequentist literature. Müller [21, 22] and Shoung and Zhang [26] provided convergence rates for univariate regression, with the multivariate case obtained by Facer and Müller [11]. Furthermore, Hasminskii [14], Donoho and Liu [9] and Tsybakov [30] showed that these rates are in fact best possible, and it is now known that the minimax rates for estimating $\boldsymbol{\mu}$ is $n^{-(\alpha-1)/(2\alpha+d)}$ and for M is $n^{-\alpha/(2\alpha+d)}$, under the usual sampling plan of choosing samples that are fixed in advance.

However if one is allowed to choose samples based on information gathered from past samples, the structure of the problem changes and we are in the sequential design setting. In this case, the minimax sequential rates of estimating $\boldsymbol{\mu}$ and M are respectively $n^{-(\alpha-1)/(2\alpha)}$ and $n^{-1/2}$ (Chen [5], Tsybakov [23] and Mokkadem and Pelletier [20]). When compared with the fixed design case, it is clear that sequential rates are uniformly better and in fact M has successfully achieved the parametric rate. Moreover, it also shows that judicious use of past information to guide future actions removes the effect of dimension d on the rates. On the more practical side, Kiefer and Wolfowitz [16] and Blum [2] used Robbins-Monro type procedures that is consistent; while Fabian [10], Dippon [8] and Mokkadem and Pelletier [20] each constructed sequential procedures that actually attain the minimax rates.

In actual practice, fully sequential design is costly to implement, because samples collection time is longer and the required logistics in collecting data in many stages is much more complicated than single-stage procedures. This then gave rise to the idea of a two-stage procedure, which offers a compromise between the added cost of doing a follow-up experiment and the added accuracy gained from it. At the first stage, limited samples are taken to give a pilot estimate of some quantity (e.g., mode), and the second stage samples are collected in the vicinity of this preliminary estimate. It was then shown in Lan *et al.* [19], Tang *et al.* [29] and Belitser *et al.* [1] that an extra second stage is enough to accelerate the convergence rates and in some cases propel them to attain the minimax sequential rates.

To the best of our knowledge however, there are hardly any such results and procedures in the Bayesian literature, whether it is in the fixed design, sequential or two-stage cases. Therefore, it is hoped that this paper will fill in this gap by giving a Bayesian solution to this problem. As we shall see, there are advantages in using the Bayesian approach, as it provides a natural framework to do two-stage estimation, and it can outperform frequentist procedures by exploiting the shrinkage property of Bayesian estimators.

In the first part of this article, we consider the fixed-in-advance sampling plan and establish single stage posterior contraction rates for $\boldsymbol{\mu}$ and M . Our prior consists of tensor product B-splines with Gaussian distributed coefficients, and we endow the error variance with some positive and continuous prior density. The main challenge here is the non-linear and non-smooth nature of the argmax and max functionals of f , and we avoid dealing with them directly by relating the estimation errors of $\boldsymbol{\mu}$ and M with the sup-norm errors of f and its first order partial derivatives. To quantify uncertainty in the estimation procedure, we construct credible sets for $\boldsymbol{\mu}$ and M , and show that they have high asymptotic coverage

with optimal sizes.

Sequential sampling or more specifically a two-stage procedure can naturally be embedded inside a Bayesian framework, as information gained from an earlier stage can be used to adjust or update one’s prior opinion. In the second part of this paper, we propose a Bayesian two-stage procedure for estimating the mode and maximum of f . We split the sample into two parts, and use the first part to compute the first stage posterior distribution of $\boldsymbol{\mu}$ and M . Using this posterior, we construct a credible set based on the techniques discussed in the first part of the paper. Second stage samples are then sampled uniformly over this set, and they are used to compute the second stage posterior of these two quantities.

We show that this second stage posterior is more concentrated around the truth than its single stage counterpart, and it can accelerate single stage minimax rates to the optimal sequential rates, under appropriate conditions on the radius of the credible set used. We test our procedure in a numerical experiment and the results seem to support our theoretical conclusions. Moreover when compared with a non-Bayesian method proposed in the literature, our Bayesian two-stage procedure seems to outperform slightly in terms of root mean square error, and this is due to the shrinkage induced by our choice of prior distributions (see Figure 2 below).

Throughout this paper, we will work with a general class of anisotropic Hölder space, such that we allow f to have different order of smoothness in each dimension. In some of our results above, it will be seen that additional smoothness in other dimensions can help alleviate the loss in accuracy due to less smoothness in some dimensions, and this borrowing of smoothness across dimensions, which is a unique feature of anisotropic spaces, can result in the improvement of the overall rate.

The paper is organized as follows. The next section introduces notations and assumptions. Section 3 describes the prior and the resulting posterior distributions of $\boldsymbol{\mu}$ and M . Section 4 contains main results on posterior contraction rates for these two quantities, and their credible sets are discussed in Section 5, accompanied by results on their coverage probabilities. We introduce the Bayesian two-stage procedure of estimating $\boldsymbol{\mu}$ and M in Section 6. Section 7 contains simulation studies for our proposed Bayesian two-stage method. Proofs of all our main results are given in Section 8 while other auxiliary results and technical lemmas are collected in Section 9 Appendix.

2 Notations and assumptions

Given two numerical sequences a_n and b_n , $a_n = O(b_n)$ or $a_n \lesssim b_n$ means a_n/b_n is bounded, while $a_n = o(b_n)$ or $a_n \ll b_n$ means $a_n/b_n \rightarrow 0$. Also, $a_n \asymp b_n$ means $a_n = O(b_n)$ and $b_n = O(a_n)$. For stochastic sequence Z_n , $Z_n = O_P(a_n)$ means $P(|Z_n| \leq C a_n) \rightarrow 1$ for some constant $C > 0$; while $Z_n = o_P(a_n)$ means $Z_n/a_n \rightarrow 0$ in P-probability.

Inequality for a vector will stand for co-ordinatewise inequality. Let $\|\boldsymbol{x}\|_p = (\sum_{k=1}^d |x_k|^p)^{1/p}$, $\|\boldsymbol{x}\|_\infty = \max_{1 \leq k \leq d} |x_k|$ and $\|\boldsymbol{x}\| = \|\boldsymbol{x}\|_2$. For a symmetric matrix \mathbf{A} , let $\lambda_{\max}(\mathbf{A})$ and $\lambda_{\min}(\mathbf{A})$ stand for its largest and smallest eigenvalues, and $\|\mathbf{A}\|_{(2,2)} = |\lambda_{\max}(\mathbf{A})|$. Given another matrix \mathbf{B} of the same size, $\mathbf{A} \leq \mathbf{B}$ means $\mathbf{B} - \mathbf{A}$ is nonnegative definite. The L_p -norm of a

function f is denoted by $\|f\|_p$.

We say $\mathbf{Z} \sim \mathbb{N}_J(\boldsymbol{\xi}, \boldsymbol{\Omega})$ if \mathbf{Z} has a J -dimensional normal distribution with mean $\boldsymbol{\xi}$ and covariance matrix $\boldsymbol{\Omega}$. By saying that $Z \sim \text{GP}(\xi, \Omega)$, we mean that $\{Z(t), t \in U\}$ is a Gaussian process with $\mathbb{E}Z(t) = \xi(t)$ and $\text{Cov}(Z(s), Z(t)) = \Omega(s, t)$ for any $s, t \in U$.

Multi-indexes will be frequently used. Let $\mathbb{N} = \{1, 2, \dots\}$ be the natural numbers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For $\mathbf{i} = (i_1, \dots, i_d)^T \in \mathbb{N}_0^d$ and $\mathbf{x} \in \mathbb{R}^d$, define $|\mathbf{i}| = \sum_{k=1}^d i_k$, $\mathbf{i}! = \prod_{k=1}^d i_k!$ and $\mathbf{x}^{\mathbf{i}} = \prod_{k=1}^d x_k^{i_k}$. For $\mathbf{r} = (r_1, \dots, r_d)^T \in \mathbb{N}_0^d$, let $D^{\mathbf{r}} = \partial^{|\mathbf{r}|} / \partial x_1^{r_1} \dots \partial x_d^{r_d}$ be a mixed partial differentiation operator. If $\mathbf{r} = \mathbf{0}$, we interpret $D^{\mathbf{0}}f \equiv f$. If $\mathbf{r} = \mathbf{e}_k$, where $\mathbf{e}_k = (0, \dots, 0, 1, 0, \dots, 0)^T$ with 1 in the k th position and zero elsewhere, we write $D^{\mathbf{e}_k}$ as D_k . We denote $\nabla f(\mathbf{x}) = (D_1 f(\mathbf{x}), \dots, D_d f(\mathbf{x}))^T$ to be the gradient of f at \mathbf{x} . If f is twice differentiable, $\mathbf{H}f(\mathbf{x}_0)$ stands for the Hessian matrix of f at \mathbf{x}_0 .

For $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_d)^T \in \mathbb{N}^d$, let us denote α^* to be the harmonic mean, i.e., $(\alpha^*)^{-1} = d^{-1} \sum_{k=1}^d \alpha_k^{-1}$. We define the anisotropic Hölder's norm $\|f\|_{\boldsymbol{\alpha}, \infty}$ as

$$\max \left\{ \|D^{\mathbf{r}} f\|_{\infty} + \sum_{k=1}^d \|D^{(\alpha_k - r_k)\mathbf{e}_k} D^{\mathbf{r}} f\|_{\infty} : \mathbf{r} \in \mathbb{N}_0^d, \sum_{k=1}^d (r_k / \alpha_k) < 1 \right\}. \quad (2.1)$$

The constraint $\sum_{k=1}^d (r_k / \alpha_k) < 1$ is a technical condition and is imposed so that contraction rates for f and its derivatives will decrease to 0 for large n .

Definition 2.1. The anisotropic Hölder space of order $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_d)^T \in \mathbb{N}^d$, denoted as $\mathcal{H}^{\boldsymbol{\alpha}}([0, 1]^d)$, consists of functions $f : [0, 1]^d \rightarrow \mathbb{R}$ such that $\|f\|_{\boldsymbol{\alpha}, \infty} < \infty$, and for some constant $C > 0$ with any $\mathbf{x}, \mathbf{x}_0 \in (0, 1)^d$,

$$|D^{\mathbf{r}} f(\mathbf{x}) - D^{\mathbf{r}} T_{\mathbf{x}_0} f(\mathbf{x})| \leq C \sum_{k=1}^d |x_k - x_{0k}|^{\alpha_k - r_k}, \quad (2.2)$$

where $\mathbf{r} \in \mathbb{N}_0^d$ and $\sum_{k=1}^d (r_k / \alpha_k) < 1$. Here $T_{\mathbf{x}_0} f(\mathbf{x}) = \sum_{\mathbf{i} \leq \mathbf{m}_{\boldsymbol{\alpha}}} D^{\mathbf{i}} f(\mathbf{x}_0) (\mathbf{x} - \mathbf{x}_0)^{\mathbf{i}} / \mathbf{i}!$ is the tensor Taylor polynomial of order $\mathbf{m}_{\boldsymbol{\alpha}} := (\alpha_1 - 1, \alpha_2 - 1, \dots, \alpha_d - 1)^T$ by expanding f around \mathbf{x}_0 .

To study frequentist properties of our posterior distribution, we assume the existence of a true regression function f_0 such that it satisfies the following three assumptions. In what follows, let $\mathcal{B}(\mathbf{x}, r) = \{\mathbf{y} : \|\mathbf{y} - \mathbf{x}\| \leq r\}$ be a ℓ_2 -ball centered at \mathbf{x} .

1. Under the true distribution P_0 , $Y_i = f_0(\mathbf{X}_i) + \varepsilon_i$, such that ε_i are i.i.d. Gaussian with mean 0 and variance σ_0^2 for $i = 1, \dots, n$.
2. $f_0 \in \mathcal{H}^{\boldsymbol{\alpha}}([0, 1]^d)$ for $\alpha_k > 2, k = 1, \dots, d$, and attains its maximum M_0 at a unique point $\boldsymbol{\mu}_0$ in $(0, 1)^d$ which is well-separated: for any constant $\tau_1 > 0$, there exists $\delta > 0$ such that $f_0(\boldsymbol{\mu}_0) \geq f_0(\mathbf{x}) + \delta$ for all $\mathbf{x} \notin \mathcal{B}(\boldsymbol{\mu}_0, \tau_1)$.
3. For any $0 < \tau \leq \tau_1$, there exists $\lambda_0 > 0$ such that $\lambda_{\max}\{\mathbf{H}f_0(\mathbf{x})\} \leq -\lambda_0$ for all $\mathbf{x} \in \mathcal{B}(\boldsymbol{\mu}_0, \tau)$.

Assumption 1 is on the true regression model for (1.1). The well-separation property of Assumption 2 ensures that only points \mathbf{x} that are near $\boldsymbol{\mu}_0$ will give values $f(\mathbf{x})$ that are close to the true maximum M . This property is needed to establish posterior consistency for $\boldsymbol{\mu}$ as we shall see in Theorem 4.1 below. Assumption 3 says that the Hessian of f_0 is locally negative definite around $\boldsymbol{\mu}_0$. Observe that Assumptions 2 and 3 imply $\nabla f_0(\boldsymbol{\mu}_0) = \mathbf{0}$ and the Hessian $\mathbf{H}f_0(\boldsymbol{\mu}_0)$ is symmetric and negative definite. Moreover, $\mathbf{H}f_0(\mathbf{x})$ is continuous in \mathbf{x} . If $\alpha_k = 2$, then we need to make an extra assumption that the second partial derivatives of f_0 is continuous; if not, the Hessian may not be symmetric and its eigenvalues may not be real.

For $x_k \in [0, 1]$, let $B_{j_k, q_k}(x_k)$ be the k th component B-spline of fixed order $q_k \geq \alpha_k$, with knots $0 = t_{k,0} < t_{k,1} < \dots < t_{k,N_k} < t_{k,N_k+1} = 1$, such that $J_k = q_k + N_k$. Assume that the set of knots in each direction is quasi-uniform, i.e., $\max_{1 \leq l \leq N_k} (t_{k,l} - t_{k,l-1}) \asymp \min_{1 \leq l \leq N_k} (t_{k,l} - t_{k,l-1})$. Examples include uniform and nested uniform partitions (cf. Examples 6.6 and 6.7 of Schumaker [24]), and we can always choose a subset of knots from any given knot sequence to form a quasi-uniform sequence (cf. Lemma 6.17 of Schumaker [24]).

For fixed design points $\mathbf{X}_i = (X_{i1}, \dots, X_{id})^T$ with $i = 1, \dots, n$, assume that there is a cumulative distribution function G , with positive and continuous density g on $[0, 1]^d$ such that

$$\sup_{\mathbf{x} \in [0,1]^d} |G_n(\mathbf{x}) - G(\mathbf{x})| = o\left(\prod_{k=1}^d N_k^{-1}\right), \quad (2.3)$$

where $G_n(\mathbf{x}) = n^{-1} \sum_{i=1}^n \mathbb{1}_{\{\mathbf{X}_i \in [0, \mathbf{x}]\}}$ is the empirical distribution of $\{\mathbf{X}_i, i = 1, \dots, n\}$, with $\mathbb{1}_U(\cdot)$ the indicator function on U . The condition holds for the discrete uniform design with G the uniform distribution when $N_k \lesssim n^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$ for $k = 1, \dots, d$. If $\mathbf{X}_i \stackrel{\text{i.i.d.}}{\sim} G$ with a continuous density on $[0, 1]^d$, then (2.3) holds with probability tending to one if $N_k \lesssim n^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$ for $k = 1, \dots, d$, and $\alpha^* > d/2$ by Donsker's theorem. In this paper, we shall prove results on posterior contraction rates and credible sets based on fixed design points. These results will translate to the random case by conditioning on the predictor variables.

3 B-splines tensor product, Gaussian prior and posterior

In the model $Y_i = f(\mathbf{x}_i) + \varepsilon_i$, $i = 1, \dots, n$, we put a finite random series prior on f based on tensor-product B-splines, i.e., $f(\mathbf{x}) = \sum_{j_1=1}^{J_1} \dots \sum_{j_d=1}^{J_d} \theta_{j_1, \dots, j_d} \prod_{k=1}^d B_{j_k, q_k}(x_k) := \mathbf{b}_{\mathbf{J}, \mathbf{q}}(\mathbf{x})^T \boldsymbol{\theta}$, where $\mathbf{b}_{\mathbf{J}, \mathbf{q}}(\mathbf{x}) = \{\prod_{k=1}^d B_{j_k, q_k}(x_k) : 1 \leq j_k \leq J_k, k = 1, \dots, d\}$ is a collection of $J = \prod_{k=1}^d J_k$ tensor-product B-splines, and $\boldsymbol{\theta} = \{\theta_{j_1, \dots, j_d} : 1 \leq j_k \leq J_k, k = 1, \dots, d\}$ are the basis coefficients. Note that $\mathbf{b}_{\mathbf{J}, \mathbf{q}}(\mathbf{x})$ and $\boldsymbol{\theta}$ are vectors indexed by d -dimensional indices and the entries are ordered lexicographically. Then by repeatedly applying equations (15) and (16) of Chapter X from de Boor [7] to each direction $k = 1, \dots, d$, the $\mathbf{r} = (r_1, \dots, r_d)^T$ mixed

partial derivative of f is given by

$$D^r f(\mathbf{x}) = \sum_{j_1=1}^{J_1} \cdots \sum_{j_d=1}^{J_d} \theta_{j_1, \dots, j_d} \prod_{k=1}^d \frac{\partial^{r_k}}{\partial x_k^{r_k}} B_{j_k, q_k}(x_k) = \mathbf{b}_{\mathbf{J}, \mathbf{q}-\mathbf{r}}(\mathbf{x})^T \mathbf{W}_r \boldsymbol{\theta}, \quad (3.1)$$

where \mathbf{W}_r is a $\prod_{k=1}^d (J_k - r_k) \times \prod_{k=1}^d J_k$ matrix whose entries consist of coefficients associated with applying the finite difference operator iteratively on $\boldsymbol{\theta}$ (for exact expressions, see (8.1)–(8.4) of Yoo and Ghosal [34]). We represent the model in (1.1) by an n -variate normal distribution $\mathbf{Y} | \mathbf{X}, \boldsymbol{\theta}, \sigma^2 \sim N_n(\mathbf{B}\boldsymbol{\theta}, \sigma^2 \mathbf{I}_n)$, where $\mathbf{B} = (\mathbf{b}_{\mathbf{J}, \mathbf{q}}(\mathbf{X}_1)^T, \dots, \mathbf{b}_{\mathbf{J}, \mathbf{q}}(\mathbf{X}_n)^T)^T$ is the B-splines basis matrix. Note that we can index the rows and columns of $\mathbf{B}^T \mathbf{B}$ by multi-dimensional indices, such that for $\mathbf{u} = (u_1, \dots, u_d)^T$ and $\mathbf{v} = (v_1, \dots, v_d)^T$, we write $(\mathbf{B}^T \mathbf{B})_{\mathbf{u}, \mathbf{v}} = \sum_{i=1}^n \prod_{k=1}^d B_{u_k, q_k}(X_{ik}) B_{v_k, q_k}(X_{ik})$.

We consider deterministic $\mathbf{J} = (J_1, \dots, J_d)^T$ number of basis functions depending on n, d and $\boldsymbol{\alpha}$. On the basis coefficients, we endow the prior $\boldsymbol{\theta} | \sigma^2 \sim N_J(\boldsymbol{\eta}, \sigma^2 \boldsymbol{\Omega})$. We choose the prior mean such that $\|\boldsymbol{\eta}\|_\infty < \infty$ and the $J \times J$ prior covariance matrix $c_1 \mathbf{I}_J \leq \boldsymbol{\Omega} \leq c_2 \mathbf{I}_J$ for some constants $0 < c_1 \leq c_2 < \infty$. We will use the same multi-dimensional indexing convention of $\mathbf{B}^T \mathbf{B}$ on $\boldsymbol{\Omega}$, and further assume that $\boldsymbol{\Omega}^{-1}$ is $\mathbf{h} = (h_1, \dots, h_d)^T$ -banded, in the sense that $(\boldsymbol{\Omega}^{-1})_{\mathbf{u}, \mathbf{v}} = 0$ if $|u_k - v_k| > h_k$ for some $k = 1, \dots, d$.

By direct calculations, $D^r f | (\mathbf{Y}, \sigma) \sim \text{GP}(\mathbf{A}_r \mathbf{Y} + \mathbf{c}_r \boldsymbol{\eta}, \sigma^2 \Sigma_r)$, where $\mathbf{A}_r, \mathbf{c}_r$ and the covariance kernel are defined for $\mathbf{x}, \mathbf{y} \in (0, 1)^d$ by

$$\mathbf{A}_r(\mathbf{x}) = \mathbf{b}_{\mathbf{J}, \mathbf{q}-\mathbf{r}}(\mathbf{x})^T \mathbf{W}_r (\mathbf{B}^T \mathbf{B} + \boldsymbol{\Omega}^{-1})^{-1} \mathbf{B}^T, \quad (3.2)$$

$$\mathbf{c}_r(\mathbf{x}) = \mathbf{b}_{\mathbf{J}, \mathbf{q}-\mathbf{r}}(\mathbf{x})^T \mathbf{W}_r (\mathbf{B}^T \mathbf{B} + \boldsymbol{\Omega}^{-1})^{-1} \boldsymbol{\Omega}^{-1}, \quad (3.3)$$

$$\Sigma_r(\mathbf{x}, \mathbf{y}) = \mathbf{b}_{\mathbf{J}, \mathbf{q}-\mathbf{r}}(\mathbf{x})^T \mathbf{W}_r (\mathbf{B}^T \mathbf{B} + \boldsymbol{\Omega}^{-1})^{-1} \mathbf{W}_r^T \mathbf{b}_{\mathbf{J}, \mathbf{q}-\mathbf{r}}(\mathbf{y}). \quad (3.4)$$

For σ , we either take an empirical Bayes approach by maximizing the marginal likelihood obtained from $\mathbf{Y} | \sigma \sim N_n[\mathbf{B}\boldsymbol{\eta}, \sigma^2(\mathbf{B}\boldsymbol{\Omega}\mathbf{B}^T + \mathbf{I}_n)]$, or use a hierarchical Bayes approach. In the former approach, the empirical Bayes estimate given by $\tilde{\sigma}_n^2 = (\mathbf{Y} - \mathbf{B}\boldsymbol{\eta})^T (\mathbf{B}\boldsymbol{\Omega}\mathbf{B}^T + \mathbf{I}_n)^{-1} (\mathbf{Y} - \mathbf{B}\boldsymbol{\eta}) / n$ is plugged into the expression of the conditional posterior process $D^r f | (\mathbf{Y}, \sigma)$; while in the latter approach, we further endow σ with a continuous and positive prior density. For example, using a conjugate inverse-gamma (IG) prior $\sigma^2 \sim \text{IG}(\beta_1/2, \beta_2/2)$, where $\beta_1 > 4$ and $\beta_2 > 0$ are hyper-parameters, and we will obtain $\sigma^2 | \mathbf{Y} \sim \text{IG}[(\beta_1 + n)/2, (\beta_2 + n\tilde{\sigma}_n^2)/2]$. The posterior for the function f itself can be recovered as a special case $\mathbf{r} = \mathbf{0}$ by setting $\mathbf{W}_0 = \mathbf{I}$.

We need to ensure that the priors discussed above for f will yield a well-defined maximum point at every realization from its posterior. The following lemma assures this property.

Lemma 3.1. *If f is given the tensor-product B-splines with normal coefficient prior, then $\boldsymbol{\mu}$ is unique for almost all sample paths of f under the posterior distribution given \mathbf{Y} (empirical or hierarchical Bayes).*

In this paper, we simply use $\Pi(\cdot | \mathbf{Y})$ to denote either the empirical or hierarchical posteriors, we do not distinguish between these two cases since both approaches yield the same rates.

4 Posterior contraction rates

The posteriors of $\boldsymbol{\mu}$ and M can be induced from the posterior of f through the argmax and maximum operators. However since these operators are nonlinear and non-differentiable, we take an indirect approach by relating the estimation errors of $\boldsymbol{\mu}$ and M to the sup-norm errors in estimating f and its mixed partial derivatives. By this strategy, results for posterior contraction rates in the supremum norm can be used to induce the corresponding rates for $\boldsymbol{\mu}$ and M as the following theorem shows.

Theorem 4.1. *Let $J_k \asymp (n/\log n)^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$, $k = 1, \dots, d$ and assume that Assumptions 1, 2 and 3 hold. Consider the empirical Bayes approach by plugging-in $\tilde{\sigma}_n$ for σ , or the hierarchical Bayes approach by equipping σ^2 with some continuous and positive prior density. Then*

$$\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| \leq \frac{\sqrt{d}}{\lambda_0} \max_{1 \leq k \leq d} \|D_k f - D_k f_0\|_\infty, \quad (4.1)$$

$$|M - M_0| \leq \|f - f_0\|_\infty. \quad (4.2)$$

Therefore for any $m_n \rightarrow \infty$ and uniformly in $\|f_0\|_{\alpha, \infty} \leq R, R > 0$,

$$E_0 \Pi(\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| > m_n (\log n/n)^{\alpha^* \{1 - (\min_{1 \leq k \leq d} \alpha_k)^{-1}\} / (2\alpha^* + d)} | \mathbf{Y}) \rightarrow 0, \quad (4.3)$$

$$E_0 \Pi(|M - M_0| > m_n (\log n/n)^{\alpha^* / (2\alpha^* + d)} | \mathbf{Y}) \rightarrow 0. \quad (4.4)$$

Clearly, a consequence of the above result is that the posterior mean $E(\boldsymbol{\mu} | \mathbf{Y})$ converges to $\boldsymbol{\mu}_0$ in ℓ_2 -norm at the same rate given in (4.3), and the same can be said for $E(M | \mathbf{Y})$. For the isotropic smooth case $\alpha_1 = \dots = \alpha_d = \alpha$, the norm in (2.1) can be generalized (see Section 2.7.1 of van der Vaart and Wellner [33]) and the B-splines approximation rate is obtained for all smoothness levels (Theorem 22 of Chapter XII in de Boor [7]). This allows generalization of these and subsequent results in this paper for arbitrary smoothness levels. The contraction rates thus obtained, when reduced to the isotropic case, are the minimax rates for this problem up to some logarithmic factor (Hasminskii [14], Donoho and Liu [9] and Tsybakov [30] as discussed in the introduction). In Section 6 below, we will describe another Bayesian procedure that is able to remove this logarithmic factor and accelerate these rates to the optimal sequential rates.

Remark 4.2. In the above rate, clearly the direction which has the least smoothness is the most influential factor in determining the contraction rate for $\boldsymbol{\mu}$ because of the presence of the second factor in the numerator of the exponent. This is unlike the contraction rate for f , which is known to be $(n/\log n)^{-\alpha^*/(2\alpha^*+d)}$ (Theorem 4.4 of Yoo and Ghosal [34]), as it depends only on the harmonic mean α^* of smoothness. The reason is evident from (4.1) in that the largest of the deviations of the function's derivative for different directions bound the accuracy of estimating $\boldsymbol{\mu}$. Nevertheless, it is easily checked that the rate obtained above is better than that obtained by applying the above result on a function of isotropic smoothness $\min_{1 \leq k \leq d} \alpha_k$. In other words, additional smoothness in other directions can help

to alleviate the comparative loss of accuracy for dimensions which are less smooth, and this borrowing of smoothness across directions, which is a unique feature to anisotropic spaces, results in the improvement of the overall rate.

5 Credible regions for mode and maximum

Let us now quantify uncertainty by constructing credible regions for $\boldsymbol{\mu}$ and M . In what follows, we require that these sets have credibility at least some given level $1 - \gamma$, and they have optimal size with asymptotic coverage probability also at least $1 - \gamma$.

We first construct credible sets in the form of supremum-norm balls on the regression function space, and then we map these regions back using the argmax and max functionals into Euclidean spaces, so that they are credible sets for $\boldsymbol{\mu}$ and M . Now, the natural Bayesian approach to this problem is to directly construct these sets from the posterior distributions of $\boldsymbol{\mu}$ and M . The main reason for favoring the proposed method is that it allows tighter control over the size of the induced credible regions in view of (4.1) and (4.2). Such a control is essential for frequentist coverage, and enables us to use them later in the Bayesian two-stage procedure in Section 6.

One last remark before we present the main result of this section. It is well known that in nonparametric models, a credible region may have frequentist coverage asymptotically less than the corresponding credibility level. The asymptotic coverage may even be zero, because the order of bias of the center of a credible set may be comparable with its variation around the truth under optimal smoothing; see Cox [6], Freedman [12] and Knapik *et al.* [18]. This is in sharp contrast with finite dimensional models where Bayesian and frequentist quantification of uncertainty agree because of the Bernstein-von Mises theorem. Knapik *et al.* [18] showed that this low coverage problem can be addressed by undersmoothing. Castillo and Nickl [3, 4] circumvented around this problem by using weaker norms to construct credible sets. Szabo *et al.* [28] and Yoo and Ghosal [34] addressed this same problem by appropriately inflating the size of credible regions to ensure coverage. In our own construction, we will use the latter approach by introducing a constant $\rho > 0$ in the radius and choose it large enough so that we also have asymptotic coverage.

Below by posterior distribution we refer to either the empirical Bayes posterior distribution by substituting $\tilde{\sigma}_n$ for σ , or the hierarchical Bayes posterior distribution obtained by putting a further prior on σ . Denote the posterior mean of f by \tilde{f} , and let $\tilde{\boldsymbol{\mu}}$ be the mode of \tilde{f} and \tilde{M} its maximum value. For $\mathbf{e}_k = (0, \dots, 0, 1, 0, \dots, 0)$ with 1 at entry k and the rest zero, we abbreviate $\mathbf{A}_{\mathbf{e}_k}$ by \mathbf{A}_k , $\mathbf{c}_{\mathbf{e}_k}$ by \mathbf{c}_k and $\Sigma_{\mathbf{e}_k}$ as Σ_k respectively for any $k = 1, \dots, d$, where these quantities were defined in (3.2)–(3.4).

Remark 5.1. Note that $\tilde{\boldsymbol{\mu}}$ is well-defined under P_0 . Indeed since the posterior mean is an affine transformation of \mathbf{Y} , it follows from Assumption 1 that \tilde{f} is a Gaussian process under P_0 . Therefore using the same argument as in the proof of Lemma 3.1, we see that \tilde{f} has unique maximum $\tilde{\boldsymbol{\mu}}$ for every realization.

Let us write $\arg \max\{A\}$ for some set A when we apply $\arg \max$ to every element of A and

likewise for $\max A$. For some $0 < \gamma < 1/2$, consider $\{f : \|D_k f - \mathbf{A}_k \mathbf{Y} - \mathbf{c}_k \boldsymbol{\eta}\|_\infty \leq \rho R_{n,k,\gamma}\}$ as a credible band for $D_k f$, where $\rho > 0$ is a sufficiently large constant and $R_{n,k,\gamma}$ is the $(1 - \gamma)$ -quantile of the posterior distribution of $\|D_k f - \mathbf{A}_k \mathbf{Y} - \mathbf{c}_k \boldsymbol{\eta}\|_\infty$. Similarly, let $R_{n,0,\gamma}$ be the $(1 - \gamma)$ -quantile of the posterior distribution of $\|f - \mathbf{A}_0 \mathbf{Y} - \mathbf{c}_0 \boldsymbol{\eta}\|_\infty$. We proceed by using these sets to induce credible regions for $\boldsymbol{\mu}$ and M through the argmax and maximum functionals, and they are given by

$$\mathcal{C}_\boldsymbol{\mu} = \arg \max \left[\bigcap_{k=1}^d \{\boldsymbol{\mu} : \|D_k f - \mathbf{A}_k \mathbf{Y} - \mathbf{c}_k \boldsymbol{\eta}\|_\infty \leq \rho R_{n,k,\gamma}\} \right], \quad (5.1)$$

$$\mathcal{C}_M = \max \{M : \|f - \mathbf{A}_0 \mathbf{Y} - \mathbf{c}_0 \boldsymbol{\eta}\|_\infty \leq \rho R_{n,0,\gamma}\}. \quad (5.2)$$

The following result establishes properties of these regions.

Theorem 5.2. *If $J_k \asymp (n/\log n)^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$, $k = 1, \dots, d$, then we have uniformly in $\|f_0\|_{\alpha,\infty} \leq R$ for any $R > 0$:*

- (i) *the credibility and coverage probability of $\mathcal{C}_\boldsymbol{\mu}$ for $\boldsymbol{\mu}$ tend to 1,*
- (ii) $\mathcal{C}_\boldsymbol{\mu} \subset \bar{\mathcal{C}}_\boldsymbol{\mu} := \{\boldsymbol{\mu} : \|\boldsymbol{\mu} - \tilde{\boldsymbol{\mu}}\|_\infty \leq \rho\sqrt{d}\lambda_0^{-1} \max_{1 \leq k \leq d} R_{n,k,\gamma}\},$
- (iii) *there exists a constant $c > 0$ such that with probability tending to 1, $\boldsymbol{\mu}_0 \in \underline{\mathcal{C}}_\boldsymbol{\mu} := \{\boldsymbol{\mu} : \|\boldsymbol{\mu} - \tilde{\boldsymbol{\mu}}\|_\infty \leq c\rho \max_{1 \leq k \leq d} R_{n,k,\gamma}\} \subset \mathcal{C}_\boldsymbol{\mu}$ and $\Pi(\underline{\mathcal{C}}_\boldsymbol{\mu} | \mathbf{Y}) \rightarrow 1,$*
- (iv) *the credibility and coverage probability of \mathcal{C}_M for M tend to 1,*
- (v) $\mathcal{C}_M \subset \{M : |M - \tilde{M}| \leq \rho R_{n,0,\gamma}\}.$

In particular, (ii) and (iii) imply that the radius of $\mathcal{C}_\boldsymbol{\mu}$ is of the order of $\max_{1 \leq k \leq d} R_{n,k,\gamma} \asymp \max_{1 \leq k \leq d} (\log n/n)^{\alpha^*(1-\alpha_k^{-1})/(2\alpha^*+d)}$ by the second assertion of Theorem 9.2 in Section 9 Appendix (with $\mathbf{r} = \mathbf{e}_k$); while (v) and the same result (with $\mathbf{r} = \mathbf{0}$) imply that the radius of \mathcal{C}_M is of the order $(\log n/n)^{\alpha^*/(2\alpha^*+d)}$. Note that these radius lengths coincide exactly to the contraction rates of Theorem 4.1 above.

The above result concludes that the induced credible regions for $\boldsymbol{\mu}$ and M , i.e., $\mathcal{C}_\boldsymbol{\mu}$ and \mathcal{C}_M respectively, have adequate frequentist coverages that are of (nearly) optimal sizes. Assertion (iii) also implies that the hyper-cubes $\bar{\mathcal{C}}_\boldsymbol{\mu}$ and $\underline{\mathcal{C}}_\boldsymbol{\mu}$, centered at $\tilde{\boldsymbol{\mu}}$ are both at least $(1 - \gamma)$ -credible and are confidence sets of nearly optimal size. Thus a credible set with guaranteed frequentist coverage can be chosen to be a simple set like a hyper-cube centered at the posterior mean. In practice, it is easier to construct such a hyper-cube than the set $\mathcal{C}_\boldsymbol{\mu}$, as it requires performing function maximization multiple times to obtain the points of $\mathcal{C}_\boldsymbol{\mu}$.

The construction of $\bar{\mathcal{C}}_\boldsymbol{\mu}$ from the data is simple: one finds b such that the credibility of $\{\boldsymbol{\mu} : \|\boldsymbol{\mu} - \tilde{\boldsymbol{\mu}}\|_\infty \leq b\}$ is $1 - \gamma$, and then inflates this around $\tilde{\boldsymbol{\mu}}$ by a large constant factor $\rho > 0$. For this set to serve as the domain for second stage sampling in a two-stage estimation, some modifications are needed, in that we adjust the length of $\bar{\mathcal{C}}_\boldsymbol{\mu}$ in each direction so that it adapts to different smoothness. In other words, we embed $\bar{\mathcal{C}}_\boldsymbol{\mu}$ inside a hyper-rectangle

and do uniform sampling inside this larger set. Clearly, keeping the constant inflation factor as small as possible makes the credible sets smaller, but it will be seen that for optimal contraction rate in the second stage, an inflation factor which goes to infinity at a specific rate will be needed.

6 Two-stage Bayesian estimation and accelerated rates of contraction

In this section we show that by obtaining samples in two stages in an appropriate manner, we can accelerate the posterior contraction rates of $\boldsymbol{\mu}$ and M to the optimal sequential rates. Given a sampling budget of n , we first obtain $n_1 < n$ samples to compute the posterior distribution. The remaining $n_2 = n - n_1$ samples are then obtained by sampling points uniformly from some regularly shaped credible region. Since this is a small region, we can approximate the regression function f by a multivariate polynomial. By further endowing the coefficients with a normal prior, we then use the second stage samples to construct the posterior distribution for f which will be seen to be more concentrated near the truth. By (4.1) and (4.2), this translates to improved posterior distributions for $\boldsymbol{\mu}$ and M .

Let $p \in (0, 1)$. In the first stage, we choose $n_1 \in \mathbb{N}$ design points $\{\tilde{\boldsymbol{x}}_i, i = 1, \dots, n_1\}$ such that $n_1/n \rightarrow p$ as $n \rightarrow \infty$ to obtain data $\mathcal{D}_1 = \{(\tilde{\boldsymbol{x}}_i, \tilde{Y}_i), i = 1, \dots, n_1\}$ for the model in (1.1). Typically, one choose $p = 1/2$ to achieve equal sample splitting but other proportions are possible depending on the sampling configuration (e.g., grid or random sampling) and other practical considerations such as field conditions and financial constraints. By using the B-spline tensor product prior discussed in Section 3, we obtain a first stage posterior for f . This then allows us to construct the set

$$\{\boldsymbol{\mu} : |\mu_k - \tilde{\mu}_k| \leq \delta_{n,k}, k = 1, \dots, d\},$$

where $\tilde{\boldsymbol{\mu}}$ is the mode of the posterior mean of f . We choose $\delta_{n,k}, k = 1, \dots, d$ such that

$$\min_{1 \leq k \leq d} \delta_{n,k} = \rho_n \max_{1 \leq k \leq d} (\log n/n)^{\alpha^*(1-\alpha_k^{-1})/(2\alpha^*+d)} \quad (6.1)$$

for a chosen sequence $\rho_n \rightarrow \infty$, so that this set is a valid credible set, as it contains $\bar{\mathcal{C}}_{\boldsymbol{\mu}}$ for large n . Now sample $n_2 = n - n_1$ locations $\{\boldsymbol{x}_1, \dots, \boldsymbol{x}_{n_2}\}$ uniformly from this credible set and observe the second stage samples $\mathcal{D}_2 = \{(\boldsymbol{x}_i, Y_i), i = 1, \dots, n_2\}$.

Next, we center the second stage design points at the origin by $\boldsymbol{z}_i = \boldsymbol{x}_i - \tilde{\boldsymbol{\mu}}$ for $i = 1, \dots, n_2$. In other words, $\boldsymbol{z}_i, i = 1, \dots, n_2$ are i.i.d uniform samples from the hyper-rectangle $\mathcal{Q} := \{\boldsymbol{x} : |x_k| \leq \delta_{n,k}, k = 1, \dots, d\}$ of sides $\delta_{n,k}, k = 1, \dots, d$. Note that $\delta_{n,k}$ is the size of the second-stage sampling domain at direction $k = 1, \dots, d$, which we shall refer to as the localization parameter at dimension k . Observe also that $\boldsymbol{z}_i, i = 1, \dots, n_2$ are independent from the errors $\boldsymbol{\varepsilon}$ in this sampling scheme. We chose this sampling domain because we need its length at each direction to adapt to different smoothness, and it is operationally more

convenient to construct credible sets in the form of hyper-rectangles and do uniform sampling on it (see Remark 6.3 below for a more thorough discussion).

At the second stage, we put a prior on the regression function by representing $f(\mathbf{z})$ at $\mathbf{z} = (z_1, \dots, z_d)^T \in \mathcal{Q}$ as a multivariate polynomial function of fixed order $\mathbf{m}_\alpha = (\alpha_1 - 1, \dots, \alpha_d - 1)^T$, i.e., for $\mathbf{z}^{\mathbf{i}} = \prod_{k=1}^d z_k^{i_k}$,

$$f_\theta(\mathbf{z}) = \sum_{\mathbf{i} \leq \mathbf{m}_\alpha} \theta_{\mathbf{i}} \mathbf{z}^{\mathbf{i}} = \mathbf{p}(\mathbf{z})^T \boldsymbol{\theta}, \quad (6.2)$$

where $\mathbf{p}(\mathbf{z}) = (\mathbf{z}^{\mathbf{i}} : \mathbf{i} \leq \mathbf{m}_\alpha)^T$ and $\boldsymbol{\theta} = (\theta_{\mathbf{i}} : \mathbf{i} \leq \mathbf{m}_\alpha)^T$ are the corresponding basis coefficients. The elements of $\{\mathbf{i} : \mathbf{i} \leq \mathbf{m}_\alpha\}$ can be enumerated as $\{\mathbf{i}_0, \mathbf{i}_1, \dots, \mathbf{i}_W\}$ where $W + 1 = \prod_{k=1}^d \alpha_k$ with $\mathbf{i}_0 = \mathbf{0}$. Define $\mathbf{Z} = (\mathbf{p}(\mathbf{z}_1), \dots, \mathbf{p}(\mathbf{z}_{n_2}))^T$, and note that for $d = 1$, \mathbf{Z} is a Vandermonde matrix.

We endow $\boldsymbol{\theta}$ with the prior $\boldsymbol{\theta} | \sigma^2 \sim N_{W+1}(\boldsymbol{\xi}, \sigma^2 \mathbf{V})$, where the entries of $\boldsymbol{\xi}$ do not depend on n and $\mathbf{V} = \text{diag} \left\{ \prod_{k=1}^d \delta_{n,k}^{-2(i_j)_k} : j = 0, 1, \dots, W \right\}$. Then it follows that the posterior $\Pi(\boldsymbol{\theta} | \mathbf{Y}, \sigma^2)$ is

$$N_{W+1} \left[(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} (\mathbf{Z}^T \mathbf{Y} + \mathbf{V}^{-1} \boldsymbol{\xi}), \sigma^2 (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \right]. \quad (6.3)$$

The empirical posterior follows by replacing σ^2 with $\tilde{\sigma}_*^2 = (n_1 \tilde{\sigma}_1^2 + n_2 \tilde{\sigma}_2^2)/n$, where $\tilde{\sigma}_1^2 = (\tilde{\mathbf{Y}} - \mathbf{B}\boldsymbol{\eta})^T (\mathbf{B}\boldsymbol{\Omega}\mathbf{B}^T + \mathbf{I}_{n_1})^{-1} (\tilde{\mathbf{Y}} - \mathbf{B}\boldsymbol{\eta})/n_1$ is the empirical estimate of σ based on the first stage samples, and $\tilde{\sigma}_2^2 = n_2^{-1} (\mathbf{Y} - \mathbf{Z}\boldsymbol{\xi})^T (\mathbf{Z}\mathbf{V}\mathbf{Z}^T + \mathbf{I}_{n_2})^{-1} (\mathbf{Y} - \mathbf{Z}\boldsymbol{\xi})$ is the same estimate based on the second stage samples.

For the hierarchical Bayes approach, we use the first stage posterior of σ as prior for the second stage. That is, we equip σ^2 with $\text{IG}(\beta_1/2, \beta_2/2)$ prior at the first stage, where $\beta_1 > 4$ and $\beta_2 > 0$, we then use the resulting posterior $\text{IG}[(\beta_1 + n_1)/2, (\beta_2 + n_1 \tilde{\sigma}_1^2)/2]$ as prior for the second stage, which will further yield $\text{IG}[(\beta_1 + n)/2, (\beta_2 + n \tilde{\sigma}_*^2)/2]$ as the second stage posterior for σ^2 . The proposition below shows that the second stage empirical Bayes estimator and the hierarchical Bayes posterior of σ are consistent, and it is a key step in establishing the main result given in Theorem 6.2 below.

Proposition 6.1 (Second stage error variance). *Uniformly over $\|f_0\|_{\alpha, \infty} \leq R$,*

- (a) *Second stage empirical Bayes estimator $\tilde{\sigma}_*^2$ converges to σ_0^2 in P_0 -probability at the rate $\max\{n^{-1/2}, n^{-2\alpha^*/(2\alpha^*+d)}, \sum_{k=1}^d \delta_{n,k}^{2\alpha_k}\}$.*
- (b) *If inverse gamma posterior from the first stage is used as the prior in the second stage, the second stage posterior of σ^2 contracts to σ_0^2 at the same rate.*

Let $\mathbf{r} = (r_1, \dots, r_d)^T$ be such that $\mathbf{r} \leq \mathbf{m}_\alpha$. Then the \mathbf{r} mixed partial derivative of f_θ is

$$D^{\mathbf{r}} f_\theta(\mathbf{z}) = \sum_{\mathbf{i} \leq \mathbf{m}_\alpha} \theta_{\mathbf{i}} \prod_{k=1}^d \frac{\partial^{r_k}}{\partial z_k^{r_k}} z_k^{i_k} = \sum_{\mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha} \theta_{\mathbf{i}} \frac{\mathbf{i}!}{(\mathbf{i} - \mathbf{r})!} \mathbf{z}^{\mathbf{i} - \mathbf{r}}, \quad (6.4)$$

which is a multivariate polynomial of order $\mathbf{m}_\alpha - \mathbf{r}$. The posterior distributions of $D^r f_\theta$ can then be induced from (6.3).

Define the location of the maximum of f_θ inside the centered region as $\boldsymbol{\mu}_z = \arg \max_{z \in Q} f_\theta(z)$. We then relate this location back to the original domain by $\boldsymbol{\mu} = \tilde{\boldsymbol{\mu}} + \boldsymbol{\mu}_z$ with corresponding maximum value $M = f_\theta(\boldsymbol{\mu}_z)$. Following the same reasoning as in Lemma 3.1, $\boldsymbol{\mu}$ is unique for almost all sample paths of f_θ under the empirical or hierarchical posterior. The following theorem establishes the second stage posterior contraction rates of $\boldsymbol{\mu}$ and M for any smoothness level $\alpha_k > 2, k = 1, \dots, d$, uniformly over $\|f_0\|_{\alpha, \infty} \leq R$.

Theorem 6.2. *For any chosen sequence $\rho_n \rightarrow \infty$, let $\delta_{n,k}, k = 1, \dots, d$, be such that $\min_{1 \leq k \leq d} \delta_{n,k} = \rho_n \max_{1 \leq k \leq d} (\log n/n)^{\alpha^*(1-\alpha_k^{-1})/(2\alpha^*+d)}$. Then under Assumptions 1, 2 and 3, we have uniformly over $\|f_0\|_{\alpha, \infty} \leq R$ and for any $m_n \rightarrow \infty$,*

$$\begin{aligned} \mathbb{E}_0 \Pi \left[\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| > m_n \max_{1 \leq k \leq d} \delta_{n,k}^{-1} \left(n^{-1/2} + \sum_{l=1}^d \delta_{n,l}^{\alpha_l} \right) \middle| \mathbf{Y} \right] &\rightarrow 0, \\ \mathbb{E}_0 \Pi \left[|M - M_0| > m_n \left(n^{-1/2} + \sum_{k=1}^d \delta_{n,k}^{\alpha_k} \right) \middle| \mathbf{Y} \right] &\rightarrow 0. \end{aligned}$$

In particular, if $\alpha_k > 1 + \sqrt{1 + d/2}$ for all $k = 1, \dots, d$, then for the choice $\delta_{n,k} = n^{-1/(2\alpha_k)}$, $k = 1, \dots, d$, the posterior distributions for $\boldsymbol{\mu}$ and M contract at the rates $n^{-(\underline{\alpha}-1)/(2\underline{\alpha})}$ and $n^{-1/2}$ respectively, where $\underline{\alpha} = \min_{1 \leq k \leq d} \alpha_k$.

Let us take $\delta_{n,k} = n^{-1/(2\alpha_k)}$ the optimal choice as suggested above. By comparing this theorem and the single stage contraction rates of Theorem 4.1, we can draw 3 main conclusions:

1. If we perform a Bayesian two-stage procedure, we accelerate contraction rates for estimating $\boldsymbol{\mu}$ and M , with M achieving the parametric rate $n^{-1/2}$.
2. At the same time, we removed extra logarithmic factors that was present in the single stage rates.
3. The second stage rates for $\boldsymbol{\mu}$ and M do not depend on d the dimension of the regression function's domain, and the effect of dimension is mitigated to a lower bound $1 + \sqrt{1 + d/2}$ required on the smoothness at each direction.

The first conclusion says that the second stage posteriors for $\boldsymbol{\mu}$ and M are more concentrated near the truth when compared with their single stage counterparts, and this is evident since the second stage rate of $\boldsymbol{\mu}$ is $n^{-(\underline{\alpha}-1)/(2\underline{\alpha})} \ll (\log n/n)^{\alpha^*(1-\underline{\alpha}^{-1})/(2\alpha^*+d)}$; while for M is $n^{-1/2} \ll (\log n/n)^{\alpha^*/(2\alpha^*+d)}$. As noted above, M has achieved its oracle or the parametric rate. In the isotropic case where $\alpha_k = \alpha, k = 1, \dots, d$, the second stage rate for $\boldsymbol{\mu}$ is $n^{-(\alpha-1)/(2\alpha)}$ and this is precisely the minimax rate for this problem under sequential sampling (Chen [5], Tsybakov [23] and Belitser *et al.* [1] as mentioned in the introduction).

Remark 6.3. There are other ways to construct credible set and obtain the second stage samples. A first attempt would be to simulate f from its first stage posterior and apply the argmax operator on f . However as the posterior of f contracts to f_0 , this approach does not ensure that the second stage samples are sufficiently spread out, i.e., the distance between samples at direction k is at least some constant multiple of $\delta_{n,k}$ for $k = 1, \dots, d$. Another way is to do uniform sampling on \mathcal{C}_μ constructed in (5.1), that is, we envelope \mathcal{C}_μ , which is possibly irregularly shaped, by the smallest hyper-cube, do uniform sampling on this cube and discard points that fall outside of \mathcal{C}_μ . By (ii) and (iii) of Theorem 5.1, $\underline{\mathcal{C}}_\mu \subset \mathcal{C}_\mu \subset \bar{\mathcal{C}}_\mu$, and samples in $\bar{\mathcal{C}}_\mu$ are proportional to those in $\underline{\mathcal{C}}_\mu$ under uniform sampling. Hence, entries of $(\mathbf{Z}^T \mathbf{Z})^{-1}$ arising from uniform sampling on \mathcal{C}_μ or on hypercubes will have the same order, and the second stage posteriors from these two sampling schemes will have the same asymptotic behavior. Operationally, sampling from \mathcal{C}_μ requires an extra step in deciding whether the sampled points fall in the set or not.


Remark 6.4. For an asymptotic analysis, the prior covariance matrix \mathbf{V} plays a minor role as the data “washes” out the prior. However, this is different for finite samples, and the correct specification of \mathbf{V} is crucial for the success of our proposed method in practical applications. Through empirical experiments, we discovered that \mathbf{V} must reflect the scaling of the space by $\delta_{n,k}$ at direction k , and its inverse must have the same structure as $\mathbf{Z}^T \mathbf{Z}$ so that $(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}$ will act as an effective shrinkage factor in (6.3). Let us write $\Delta := \text{diag} \left\{ \prod_{k=1}^d \delta_{n,k}^{-(i_j)_k} : j = 0, 1, \dots, W \right\}$ and we can see that $\mathbf{Z}^T \mathbf{Z} = \Delta \mathbf{A} \Delta$ where \mathbf{A} is a matrix of constants not depending on n (see Lemma 8.1 for more details). Therefore if we centered the second stage design points, then we match the structure of $\mathbf{Z}^T \mathbf{Z}$ by choosing $\mathbf{V} = \Delta^2$, which is our default choice. If the points are not centered, we found that Zellner’s g -prior will work, i.e., $\mathbf{V} = g(\mathbf{Z}^T \mathbf{Z})^{-1}$ where g can be estimated by empirical or hierarchical Bayes methods.

7 Simulation study

We shall compare the performance of our two-stage Bayesian procedure with two other estimation methods: the single-stage Bayesian, and the two-stage frequentist procedure proposed in Belitser *et al.* [1]. Consider the following true regression function defined on $[0, 1]^2$:

$$f_0(x, y) = \left(1 + e^{-5(2x-1)^2 - 2(2y-1)^4} \right) [\cos 4(2x - 1) + \cos 5(2y - 1)]$$

where the true mode is given by $\boldsymbol{\mu}_0 = (0.5, 0.5)^T$. In the first stage, we observed f_0 on a uniform 30×30 grid with i.i.d. errors distributed as $N(0, 0.01)$ (see Figure 1a with black circles as observations). We use tensor product B-splines with normal coefficients as our prior (see Section 3). We choose the pair (J_1, J_2) that maximizes its posterior, i.e., $\Pi(J_1 = j_1, J_2 = j_2 | \mathbf{Y}, \sigma = \tilde{\sigma}_1) \propto \tilde{\sigma}_1^{-n_1} [\det(\mathbf{B} \boldsymbol{\Omega} \mathbf{B}^T + \mathbf{I}_{n_1})]^{-1/2} \Pi(J_1 = j_1, J_2 = j_2)$. If we put independent discrete uniform priors on each J_1, J_2 , we can ignore the second prior factor when doing our comparisons. We find that $J_1 = 7$ and $J_2 = 9$ based on this criterion. At each dimension, the



(a) A plot of f_0 , with the black solid point as $f_0(\boldsymbol{\mu}_0) = M_0$ and black circles as the first stage 900 observations.

(b) Posterior mean based on bivariate B-spline prior, and the black circles are the 864 second stage samples.

Figure 1: Our proposed Bayesian two-stage procedure: first stage on the left and second stage on the right.

B-spline is of order 4 (cubic) with different uniform knot sequence. For the prior parameters, we set $\boldsymbol{\eta} = \mathbf{0}$ and $\boldsymbol{\Omega} = \mathbf{I}$. Figure 1b shows the surface of the posterior mean of f .

We sample 864 points uniformly in $\{\boldsymbol{\mu} : |\mu_k - \tilde{\mu}_k| \leq \delta_k, k = 1, 2\}$ (see black circles in Figure 1b) to obtain second stage data $Y_i, i = 1, \dots, 864$. This number is chosen so as to fulfill the sampling configuration required by the frequentist method explained below, and to ensure that all methods under consideration will use the same amount of samples. We chose $\delta_1 = 0.11$ and $\delta_2 = 0.13$ by minimizing the root mean square error of the second stage posterior mean. We center the 864 sampled design points at the origin by subtracting each of them by $\tilde{\boldsymbol{\mu}}$, and we use tensor product of quadratic polynomials with normal coefficients as prior (see (6.2)). That is for $x, y \in \mathcal{Q}$, $f_{\boldsymbol{\theta}}(x, y) = \theta_0 + \theta_1 x + \theta_2 y + \theta_3 x^2 + \theta_4 y^2 + \theta_5 xy + \theta_6 xy^2 + \theta_7 x^2 y + \theta_8 x^2 y^2$. Since $x, y \in (0, 1)$, we note that the last three columns of the constructed basis matrix \mathbf{Z} (corresponding to the last three terms) are very small in magnitude compared with the remaining terms. Thus for numerical simplicity, we consider

$$f_{\boldsymbol{\theta}}^*(x, y) = \theta_0 + \theta_1 x + \theta_2 y + \theta_3 x^2 + \theta_4 y^2 + \theta_5 xy,$$

where $\boldsymbol{\theta} | \sigma^2 \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{V})$, with $\mathbf{V} = \text{diag}(1, \delta_1^{-2}, \delta_2^{-2}, \delta_1^{-4}, \delta_2^{-4}, \delta_1^{-2} \delta_2^{-2})$. We use the empirical Bayes method to estimate σ by $\tilde{\sigma}_*$, which is the weighted average of first and second stage estimates $\tilde{\sigma}_1, \tilde{\sigma}_2$. However, we note in our simulations that $\tilde{\sigma}_2$ gives a much better estimate than $\tilde{\sigma}_1$ at the current n_1, n_2 samples, and since both are valid independent estimates of σ , we take $\tilde{\sigma}_* = \tilde{\sigma}_2$. Now, to compute $\boldsymbol{\mu}_z = \arg \max_{z \in \mathcal{Q}} f_{\boldsymbol{\theta}}^*(z)$ for a fixed $\boldsymbol{\theta}$, we solve the

following system of equation $\nabla f_{\theta}^*(\boldsymbol{\mu}_z) = \mathbf{0}$, which is equivalent to solving

$$\begin{pmatrix} 2\theta_3 & \theta_5 \\ \theta_5 & 2\theta_4 \end{pmatrix} \begin{pmatrix} \mu_{z,1} \\ \mu_{z,2} \end{pmatrix} = \begin{pmatrix} -\theta_1 \\ -\theta_2 \end{pmatrix} \quad (7.1)$$

for $\boldsymbol{\mu}_z = (\mu_{z,1}, \mu_{z,2})^T$. Therefore, to induce the posterior distribution of $\boldsymbol{\mu}$, we draw samples from $\Pi(\boldsymbol{\theta}|\mathbf{Y})$ by substituting $\sigma = \tilde{\sigma}_2$ in (6.3), solving for $\boldsymbol{\mu}_z$ using (7.1) for each sample, and translating back to $\boldsymbol{\mu} = \tilde{\boldsymbol{\mu}} + \boldsymbol{\mu}_z$.




Figure 2: Root mean square errors (y -axis) for the Bayesian and frequentist procedures with 1000 Monte Carlo replications.

The procedure described is implemented in the statistical software package R. Univariate B-splines are constructed using the `bs` function from the `splines` package. We then use `tensor.prod.model.matrix` from the `crs` package to form their tensor-products. Observe that we have used a total of $30 \times 30 + 864 = 1764$ observations. To show that the two-stage procedure indeed has better accuracy than single-stage procedures, we then compare our two-stage procedure with a Bayesian single-stage method that uses the same amount of samples, i.e., on a uniform 42×42 grid points. As in the previous case, we observed f_0 at these points with i.i.d errors $N(0, 0.01)$. We then use bivariate tensor B-splines with normal coefficients as prior with the same setting. We found that $J_1 = 9, J_2 = 9$ maximizes its posterior.

To compare Bayesian and frequentist procedures, we repeated the same experiment using the two-stage frequentist procedure implemented in Belitser *et al.* [1]. In their procedure, we first fit a locally linear surface by `loess` regression in R on the first stage design points. We chose the corresponding bandwidth or span parameter to be 0.02 by leave-one-out cross validation. The maximum of this surface serves as a preliminary estimator. We construct a rectangle of size $2\delta_1$ by $2\delta_2$ around this estimator and further divide this region into 4 smaller δ_1 by δ_2 rectangles. We chose $\delta_1 = 0.06$ and $\delta_2 = 0.06$ to minimize the root mean square error of the second stage estimator. Then, we take 96 replicated samples at the 9 grid

points to form 864 second stage samples. A quadratic surface is fit through these points and its coefficients are estimated using least squares. We compute the second stage estimator of $\boldsymbol{\mu}_0$ by (7.1) and call it $\widehat{\boldsymbol{\mu}}_2$.

Let $\widetilde{\boldsymbol{\mu}}_1$ and $\widetilde{\boldsymbol{\mu}}_2$ be the single stage and two-stage posterior means respectively. To compare the performance of our proposed Bayesian two-stage method with the other two, we replicate the experiment 1000 times for each of the three methods. For each replicated experiment, we compute $\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\|$ for each $\boldsymbol{\mu} = \widetilde{\boldsymbol{\mu}}_1, \widetilde{\boldsymbol{\mu}}_2, \widehat{\boldsymbol{\mu}}_2$. Figure 2 shows the box-plots of these 1000 computed root mean square errors (RMSE) for all three procedures.

We see that the our proposed Bayesian two-stage procedure has considerably lower RMSE than the corresponding single-stage procedure, and thus supports the conclusion of Theorem 6.2 in finite sample setting. We also observe that our proposed Bayesian two-stage procedure performs slightly better than the frequentist procedure. This is due to our choice of prior covariance matrix \mathbf{V} where it causes $(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}$ in the posterior (see (6.3)) to act as an effective shrinkage factor.

8 Proofs

Recall that posterior mean of $D^r f$ is $D^r \widetilde{f} := \mathbf{A}_r \mathbf{Y} + \mathbf{c}_r \boldsymbol{\eta}$. We write the posterior contraction rate of $\boldsymbol{\mu}$ given in Theorem 4.1 as $\epsilon_n = \max_{1 \leq k \leq d} \epsilon_{n,k}$, where $\epsilon_{n,k} = (\log n/n)^{\alpha^*(1-\alpha_k^{-1})/(2\alpha^*+d)}$.

Proof of Lemma 3.1. For the empirical Bayes case, the reproducing kernel Hilbert space of Gaussian posterior process $\{f(\mathbf{t}) : \mathbf{t} \in [0, 1]^d\}$ given \mathbf{Y} is the J -dimensional space of polynomial splines spanned by elements of $\mathbf{b}_{J,q}(\cdot)$ (see Theorem 4.2 of van der Vaart and van Zanten [32]). This implies that it has continuous sample path at every realization. Observe that $\widetilde{\sigma}_n^2 > 0$ almost surely. Then if

$$\begin{aligned} 0 &= \text{Var}(f(\mathbf{t}) - f(\mathbf{s}) | \mathbf{Y}, \sigma = \widetilde{\sigma}_n) \\ &= \widetilde{\sigma}_n^2 (\mathbf{b}_{J,q}(\mathbf{t}) - \mathbf{b}_{J,q}(\mathbf{s}))^T (\mathbf{B}^T \mathbf{B} + \boldsymbol{\Omega}^{-1})^{-1} (\mathbf{b}_{J,q}(\mathbf{t}) - \mathbf{b}_{J,q}(\mathbf{s})), \end{aligned}$$

this implies that $B_{j_k, q_k}(t_k) = B_{j_k, q_k}(s_k)$ for $j_k = 1, \dots, J_k$, and $k = 1, \dots, d$. Since $q_k \geq \alpha_k > 2$ for $k = 1, \dots, d$, this rule out the possibility that the B-splines are step functions and further implies that $\mathbf{t} = \mathbf{s}$. Thus by Lemma 2.6 of Kim and Pollard [17], $\boldsymbol{\mu}$ is unique for almost all sample paths of $\{f(\mathbf{t}) : \mathbf{t} \in [0, 1]^d\}$ under the empirical posterior distribution given \mathbf{Y} .

For hierarchical Bayes, consider the conditional posterior process $\Pi(f | \mathbf{Y}, \sigma)$ for arbitrary $\sigma > 0$. The reproducing kernel Hilbert space of this Gaussian process is still the space of splines as before, and each realization of f has continuous sample path. Now by substituting σ for $\widetilde{\sigma}_n$ in the preceding display, we see that $\text{Var}(f(\mathbf{t}) - f(\mathbf{s}) | \mathbf{Y}, \sigma) = 0$ implies $\mathbf{t} = \mathbf{s}$. Again by Lemma 2.6 of Kim and Pollard [17], almost every sample path of f given σ has unique maximum for any $\sigma > 0$. Note that f is generated from the following scheme: draw $\sigma^2 \sim \Pi(\sigma^2 | \mathbf{Y})$, and then $f | \sigma \sim \text{GP}(\mathbf{A}\mathbf{Y} + \mathbf{c}\boldsymbol{\eta}, \sigma^2 \boldsymbol{\Sigma})$. Hence almost every draw of f has unique maximum $\boldsymbol{\mu}$ under $\Pi(f | \mathbf{Y})$. \square

Proof of Theorem 4.1. If both $f, f_0 \geq 0$, then (4.2) follows from the reverse triangle inequality by noting that $M = \|f\|_\infty$ and $M_0 = \|f_0\|_\infty$ in this case. If not, we can add a large enough constant $C > 0$ such that $g = f + C \geq 0$ and $g_0 = f_0 + C \geq 0$. Then by the reverse triangle inequality, $|\|g\|_\infty - \|g_0\|_\infty| \leq \|g - g_0\|_\infty$. The right hand side is $\|f - f_0\|_\infty$, while the left hand side is $|M - M_0|$ because $\|g\|_\infty = M + C$, $\|g_0\|_\infty = M_0 + C$.

To prove (4.1), we need to first establish consistency of the induced posterior of $\boldsymbol{\mu}$. Now for any $\epsilon > 0$ and $\delta > 0$, $\Pi(\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| > \epsilon | \mathbf{Y})$ is bounded above by

$$\begin{aligned} & \Pi\left(\sup_{\mathbf{x} \notin \mathcal{B}(\boldsymbol{\mu}_0, \epsilon)} f(\mathbf{x}) > f(\boldsymbol{\mu}_0) \mid \mathbf{Y}\right) \\ & \leq \Pi\left(\sup_{\mathbf{x} \notin \mathcal{B}(\boldsymbol{\mu}_0, \epsilon)} f(\mathbf{x}) > f(\boldsymbol{\mu}_0), \quad f(\boldsymbol{\mu}_0) \geq f_0(\boldsymbol{\mu}_0) - \delta/2 \mid \mathbf{Y}\right) \\ & \quad + \Pi(f(\boldsymbol{\mu}_0) < f_0(\boldsymbol{\mu}_0) - \delta/2 | \mathbf{Y}). \end{aligned}$$

The second term is bounded above by $\Pi(|f(\boldsymbol{\mu}_0) - f_0(\boldsymbol{\mu}_0)| > \delta/2 | \mathbf{Y}) \leq \Pi(\|f - f_0\|_\infty > \delta/2 | \mathbf{Y})$, and this goes to 0 in P_0 -probability by Theorem 9.1 in Section 9 Appendix with $\mathbf{r} = \mathbf{0}$. The well-separation property of Assumption 2 implies that there exists a $\delta > 0$, such that $f_0(\mathbf{x}) < f_0(\boldsymbol{\mu}_0) - \delta$ for $\mathbf{x} \notin \mathcal{B}(\boldsymbol{\mu}_0, \epsilon)$. Hence, for this $\delta > 0$ and appealing again to Theorem 9.1, the first term is bounded above by

$$\Pi\left(\bigcup_{\mathbf{x} \notin \mathcal{B}(\boldsymbol{\mu}_0, \epsilon)} \left\{f(\mathbf{x}) > f_0(\mathbf{x}) + \frac{\delta}{2}\right\} \mid \mathbf{Y}\right) \leq \Pi\left(\|f - f_0\|_\infty > \frac{\delta}{2} \mid \mathbf{Y}\right) \rightarrow 0$$

in P_0 -probability as $n \rightarrow \infty$. Thus the posterior distribution of $\boldsymbol{\mu}$ is consistent at $\boldsymbol{\mu}_0$.

Now by Taylor's theorem, $\nabla f_0(\boldsymbol{\mu}) = \nabla f_0(\boldsymbol{\mu}_0) + \mathbf{H}f_0(\boldsymbol{\mu}^*)(\boldsymbol{\mu} - \boldsymbol{\mu}_0)$ for some $\boldsymbol{\mu}^* = \lambda\boldsymbol{\mu} + (1 - \lambda)\boldsymbol{\mu}_0$ with $\lambda \in [0, 1]$. Since the posterior of $\boldsymbol{\mu}$ is consistent as shown above and $\boldsymbol{\mu}^*$ falls between $\boldsymbol{\mu}$ and $\boldsymbol{\mu}_0$, it must be that as $n \rightarrow \infty$ and for any $\epsilon > 0$, $\Pi(\|\boldsymbol{\mu}^* - \boldsymbol{\mu}_0\| < \epsilon | \mathbf{Y}) \xrightarrow{P_0} 1$. Let us introduce the event $\mathcal{B}_n = \{\lambda_{\max}(\mathbf{H}f_0(\boldsymbol{\mu}^*)) \leq -\lambda_0\}$, and note that under Assumptions 1 and 2, $\Pi(\mathcal{B}_n | \mathbf{Y}) \xrightarrow{P_0} 1$ and $\mathbf{H}f_0(\boldsymbol{\mu}^*)$ is invertible with probability tending to one under the posterior. Therefore, as $n \rightarrow \infty$ and intersecting with \mathcal{B}_n , $\boldsymbol{\mu} - \boldsymbol{\mu}_0 = \mathbf{H}f_0(\boldsymbol{\mu}^*)^{-1}(\nabla f_0(\boldsymbol{\mu}) - \nabla f_0(\boldsymbol{\mu}_0))$. Noting that $\nabla f_0(\boldsymbol{\mu}_0) = \nabla f(\boldsymbol{\mu}) = \mathbf{0}$, then

$$\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| \leq \frac{1}{\lambda_0} \|\nabla f_0(\boldsymbol{\mu}) - \nabla f(\boldsymbol{\mu})\| \leq \frac{\sqrt{d}}{\lambda_0} \max_{1 \leq k \leq d} \|D_k f - D_k f_0\|_\infty,$$

where we have used the sub-multiplicative property of the $\|\cdot\|_{(2,2)}$ -norm, i.e., $\|\mathbf{A}\mathbf{y}\| \leq \|\mathbf{A}\|_{(2,2)}\|\mathbf{y}\|$ for some matrix \mathbf{A} and vector \mathbf{y} .

Theorem 9.1 with $\mathbf{r} = \mathbf{0}$ together with (4.2) now proves the desired contraction rate on M . To derive the rate (4.3) for $\boldsymbol{\mu}$, apply again Theorem 9.1 with $\mathbf{r} = \mathbf{e}_k$, and the L_∞ -contraction rate for $D_k f$ is $\epsilon_{n,k}$, $k = 1, \dots, d$. In view of (4.1), we have for any $m_n \rightarrow \infty$,

$$\mathbb{E}_0 \Pi(\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| > m_n \epsilon_n | \mathbf{Y}) \leq \sum_{k=1}^d \mathbb{E}_0 \Pi\left(\|D_k f - D_k f_0\|_\infty > \frac{\lambda_0}{\sqrt{d}} m_n \epsilon_n \mid \mathbf{Y}\right)$$

approaches 0 uniformly in $\|f_0\|_{\alpha,\infty} \leq R$ and hence establishing the assertion. \square

Proof of Theorem 5.2. By construction, \mathcal{C}_μ contains $\tilde{\boldsymbol{\mu}}$, the mode of the posterior mean \tilde{f} of f . Since $\gamma < 1/2$, $R_{n,k,\gamma}$ is greater than the posterior median of $\|D_k f - D_k \tilde{f}\|_\infty$. For the empirical Bayesian posterior, $D_k f - D_k \tilde{f}$ is a Gaussian process under Assumption 1 and in view of the second assertion of Theorem 9.2, it follows that $R_{n,k,\gamma} \gtrsim \epsilon_{n,k}$ (with $\mathbf{r} = \mathbf{e}_k$). Define $c_{n,k} := \sup_{\mathbf{x} \in [0,1]^d} \text{var}(D_k f(\mathbf{x}) - D_k \tilde{f}(\mathbf{x}) | \mathbf{Y})$ and observe that by repeatedly applying the inequality $\mathbf{y}^T \mathbf{A} \mathbf{y} \leq \lambda_{\max}(\mathbf{A}) \|\mathbf{y}\|^2$, we deduce that for any $\mathbf{x} \in [0,1]^d$,

$$\begin{aligned} \text{var}(D_k f(\mathbf{x}) - D_k \tilde{f}(\mathbf{x}) | \mathbf{Y}) &\leq \sigma_0^2 \lambda_{\max}(\mathbf{B}^T \mathbf{B}) \lambda_{\min}^{-2}(\mathbf{B}^T \mathbf{B} + \boldsymbol{\Omega}^{-1}) \lambda_{\max}(\mathbf{W}_{\mathbf{e}_k} \mathbf{W}_{\mathbf{e}_k}^T) \|\mathbf{b}_{\mathbf{J}, \mathbf{q} - \mathbf{e}_k}(\mathbf{x})\|^2 \\ &\lesssim n^{-1} J_k^2 \prod_{l=1}^d J_l \lesssim n^{-2\alpha^*(1-\alpha_k^{-1})/(2\alpha^*+d)}, \end{aligned}$$

for any $n \geq 2$, and we conclude that $c_{n,k} = o(\epsilon_{n,k}^2)$. Here by using $0 \leq B_{j_l, q_l}(x_l) \leq 1$ and the partition of unity property of B-splines $\sum_j \prod_{l=1}^d B_{j_l, q_l}(x_l) = 1$, we have $\|\mathbf{b}_{\mathbf{J}, \mathbf{q} - \mathbf{e}_k}(\mathbf{x})\|^2 \leq \sum_{j_1=1}^{J_1} \cdots \sum_{j_d=1}^{J_d} \prod_{l=1}^d B_{j_l, q_l - 1\{l=k\}}(x_l) \leq 1$ for any $\mathbf{x} \in [0,1]^d$. The rest of the norms are then bounded by Lemma 9.4 (with $\mathbf{r} = \mathbf{e}_k$) and (9.2), (9.1) of Lemma 9.3. Therefore by Borell's inequality (cf. Proposition A.2.1 of van der Vaart and Wellner [33]) and taking ρ to be large enough, e.g. $\rho > 1$,

$$\begin{aligned} \Pi(\boldsymbol{\mu} \notin \mathcal{C}_\mu | \mathbf{Y}) &\leq \sum_{k=1}^d \Pi(\|D_k f - D_k \tilde{f}\|_\infty > \rho R_{n,k,\gamma} | \mathbf{Y}) \\ &\leq d \max_{1 \leq k \leq d} \exp[-(\rho - 1)^2 R_{n,k,\gamma}^2 / c_{n,k}] \end{aligned}$$

which goes to zero since $R_{n,k,\gamma}^2 / c_{n,k} \rightarrow \infty$. Thus the credibility of \mathcal{C}_μ is at least $1 - \gamma$. For the hierarchical Bayesian posterior, the same conclusion follows since conditionally on σ , the posterior law obeys a Gaussian process and σ lies in a small neighborhood of true σ_0 with high posterior probability (from (c) of Proposition 9.5). To ensure coverage, note that by the construction of \mathcal{C}_μ , it contains $\boldsymbol{\mu}_0$ if $\|D_k f_0 - D_k \tilde{f}\|_\infty \leq \rho R_{n,k,\gamma}$ for all $k = 1, \dots, d$. When f_0 is the true regression function, the P_0 -probability of the last event tends to one for all k by Theorem 9.2 with $\mathbf{r} = \mathbf{e}_k$, and hence the assertion on coverage is established. This proves assertion (i).

Assertion (ii) follows in view of (5.1) and if (4.1) holds also when we replace $\boldsymbol{\mu}_0$ by $\tilde{\boldsymbol{\mu}}$ and f_0 by \tilde{f} . By Remark 5.1, $\nabla \tilde{f}(\tilde{\boldsymbol{\mu}}) = \mathbf{0}$ and the Hessian matrix $\mathbf{H} \tilde{f}(\tilde{\boldsymbol{\mu}})$ is non-negative definite and symmetric. Moreover, since \tilde{f} is a polynomial splines of order $q_k \geq \alpha_k > 2$, $k = 1, \dots, d$ by Assumption 2, $\mathbf{H} \tilde{f}(\mathbf{x})$ is continuous. Now since $\tilde{f} \rightarrow f_0$ uniformly in P_0 -probability as a result of Theorem 9.1 and f_0 has a well-separated maximum by Assumption 2, it follows that $\tilde{\boldsymbol{\mu}}$ is consistent in estimating $\boldsymbol{\mu}_0$ by Theorem 5.7 of van der Vaart [31]. Hence for n large enough, $\mathcal{B}(\tilde{\boldsymbol{\mu}}, \tau/2)$ is contained in $\mathcal{B}(\boldsymbol{\mu}_0, \tau)$ for the same τ appearing in Assumption 3. Now since $D^r \tilde{f}$ converges uniformly to $D^r f_0$ (Theorem 9.1), and using the fact that the maximum eigenvalue is a continuous operation, we have by the continuous mapping theorem that $\sup_{\mathbf{x} \in \mathcal{B}(\tilde{\boldsymbol{\mu}}, \tau/2)} \lambda_{\max}(\mathbf{H} \tilde{f}(\mathbf{x})) \leq -\lambda_0$ for sufficiently large n . Consequently, the proof of

the inequality (4.1) given in Theorem 4.1 will go through even if we replace f_0 with \tilde{f} and $\boldsymbol{\mu}_0$ with $\tilde{\boldsymbol{\mu}}$.

Let $\mathbf{1}_d$ be a d -dimensional vector of ones. For (iii), we have by the mean value theorem, uniformly in $\|f_0\|_{\alpha,\infty} \leq R$ that for $k = 1, \dots, d$,

$$\|D_k \tilde{f}(\cdot - R^{-1} \mathbf{1}_d \rho R_{n,k,\gamma}) - D_k \tilde{f}\|_\infty \leq \|D_{k+1} \tilde{f}\|_\infty \rho R_{n,k,\gamma} / R \leq \rho R_{n,k,\gamma}$$

when n is large enough. The last inequality follows since $\|D_{k+1} \tilde{f}\|_\infty$ converges in probability to $\|D_{k+1} f_0\|_\infty \leq \|f_0\|_{\alpha,\infty}$. Now since the mode of $\tilde{f}(\cdot - R^{-1} \mathbf{1}_d \rho R_{n,k,\gamma})$ is $\tilde{\boldsymbol{\mu}} + R^{-1} \mathbf{1}_d \rho R_{n,k,\gamma}$, it follows immediately that $\tilde{\boldsymbol{\mu}} + R^{-1} \mathbf{1}_d \rho \max_{1 \leq k \leq d} R_{n,k,\gamma} \in \mathcal{C}_\mu$. This shows that with probability tending to one, \mathcal{C}_μ contains a hyper-cube centered at $\tilde{\boldsymbol{\mu}}$ of size $R^{-1} \rho \max_{1 \leq k \leq d} R_{n,k,\gamma}$. The statements about credibility and coverage of \mathcal{C}_μ hold in view of Theorem 9.2 if

$$\|\tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}_0\| \leq \sqrt{d} \lambda_0^{-1} \max_{1 \leq k \leq d} \|D_k \tilde{f} - D_k f_0\|_\infty \quad (8.1)$$

in P_0 -probability. Indeed as shown above, $\tilde{\boldsymbol{\mu}}$ is consistent in estimating $\boldsymbol{\mu}_0$ and we can use the same Taylor's series argument used in proving (4.1) of Theorem 4.1 with $\tilde{\boldsymbol{\mu}}$ and \tilde{f} playing the roles of $\boldsymbol{\mu}$ and f respectively.

The proof of (iv) is similar to that of (i) with $\max_{1 \leq k \leq d} R_{n,k,\gamma}$ replaced by $R_{n,0,\gamma}$. The proof of assertion (v) can be completed by following the arguments used in the proof of assertion (ii) with (4.2) applied to the pair f and \tilde{f} . \square

To prove results in Section 6, we need to first lay out some preliminary details. Define $f_{0,z}(\mathbf{x} - \tilde{\boldsymbol{\mu}}) = f_0(\mathbf{x})$ to be the shifted true function. Let $\boldsymbol{\theta}_0 = (\theta_{0,i} : i \leq \mathbf{m}_\alpha)^T$ be a random vector such that $f_{\boldsymbol{\theta}_0}(\mathbf{x} - \tilde{\boldsymbol{\mu}}) = T_{\boldsymbol{\mu}_0} f_0(\mathbf{x})$, where $f_{\boldsymbol{\theta}}$ is from (6.2) and $T_{\boldsymbol{\mu}_0} f_0(\mathbf{x})$ is the Taylor polynomial of order \mathbf{m}_α by expanding f_0 around $\boldsymbol{\mu}_0$, that is,

$$\sum_{i \leq \mathbf{m}_\alpha} \theta_{0,i} (\mathbf{x} - \tilde{\boldsymbol{\mu}})^i = f_0(\boldsymbol{\mu}_0) + \sum_{i \leq \mathbf{m}_\alpha, |i| \geq 2} \frac{D^i f_0(\boldsymbol{\mu}_0)}{i!} (\mathbf{x} - \boldsymbol{\mu}_0)^i, \quad (8.2)$$

where $\nabla f_0(\boldsymbol{\mu}_0) = \mathbf{0}$ by Assumption 2. Hence $\boldsymbol{\theta}_0$ can be thought of as the true $\boldsymbol{\theta}$ by projecting f_0 onto the space of polynomials of order \mathbf{m}_α . Note that $\boldsymbol{\theta}_0$ is random and depends on $\tilde{\boldsymbol{\mu}}$, $\boldsymbol{\mu}_0$ and f_0 . By applying D^i on both sides of (8.2) and evaluating at $\mathbf{x} = \tilde{\boldsymbol{\mu}}$, we have $i! \theta_{0,i} = D^i T_{\boldsymbol{\mu}_0} f_0(\tilde{\boldsymbol{\mu}})$. Note that since $D^i f_0(\mathbf{x})$, $i \leq \mathbf{m}_\alpha$, are continuous by Assumption 2, they are bounded over $\{\boldsymbol{\mu} : |\mu_k - \tilde{\mu}_k| \leq \delta_{n,k}, k = 1, \dots, d\}$, and this implies that for any $i \leq \mathbf{m}_\alpha$, $|\theta_{0,i}| = O_{P_0}(1)$ uniformly over $\|f_0\|_{\alpha,\infty} \leq R$. The design matrix \mathbf{Z} is generated using i.i.d. uniform samples and by Lemma 8.1, we know that $\mathbf{Z}^T \mathbf{Z}$ is invertible with probability going to 1 as $n \rightarrow \infty$. In actual computation, the invertibility of $\mathbf{Z}^T \mathbf{Z}$ is not an important issue as there is the presence of the prior covariance matrix \mathbf{V} to serve as a regularization factor, and $\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1}$ in (6.3) is always invertible by our choice of \mathbf{V} .

As a consequence of Theorem 9.1, $D_k \tilde{f}$ converges uniformly to $D_k f_0$ at the rate $\epsilon_{n,k}$, and by (8.1), this translates to $\|\tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}_0\| = O_{P_0}(\epsilon_n)$. Let $\mathbf{F}_0 = (f_0(\mathbf{x}_1), \dots, f_0(\mathbf{x}_{n_2}))^T$

where $\{\mathbf{x}_1, \dots, \mathbf{x}_{n_2}\}$ is the original (unshifted) second stage samples. Note that $(\mathbf{Z}\boldsymbol{\theta}_0)_i = f_{\boldsymbol{\theta}_0}(\mathbf{x}_i - \tilde{\boldsymbol{\mu}}) = T_{\boldsymbol{\mu}_0} f_0(\mathbf{x}_i)$ and under the assumption of (2.2), we have

$$\begin{aligned} \|\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0\|_\infty &= \max_{1 \leq i \leq n_2} |f_0(\mathbf{x}_i) - T_{\boldsymbol{\mu}_0} f_0(\mathbf{x}_i)| \lesssim \max_{1 \leq i \leq n_2} \sum_{k=1}^d |x_{ik} - \mu_{0k}|^{\alpha_k} \\ &\lesssim \max_{1 \leq i \leq n_2} \sum_{k=1}^d |x_{ik} - \tilde{\mu}_k|^{\alpha_k} + \sum_{k=1}^d |\tilde{\mu}_k - \mu_{0k}|^{\alpha_k} \lesssim \sum_{k=1}^d \delta_{n,k}^{\alpha_k} \end{aligned} \quad (8.3)$$

uniformly over $\|f_0\|_{\alpha, \infty} \leq R$. The second line follows from the inequality $|x + y|^r \leq \max\{1, 2^{r-1}\}(|x|^r + |y|^r)$, while the last inequality is due to $|x_{ik} - \tilde{\mu}_k| \leq \delta_{n,k}$ almost surely since $x_{ik} - \tilde{\mu}_k \sim \text{Uniform}(-\delta_{n,k}, \delta_{n,k})$; and $|\tilde{\mu}_k - \mu_{0k}| \leq \|\tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}_0\| = O_{P_0}(\epsilon_n) = o_{P_0}(\delta_{n,k})$ as argued above, where $\epsilon_n = o(\delta_{n,k})$, $k = 1, \dots, d$ was by our choice in (6.1).

Proof of Proposition 6.1. By the triangle inequality, $|\tilde{\sigma}_*^2 - \sigma_0^2| \leq |\tilde{\sigma}_1^2 - \sigma_0^2| + |\tilde{\sigma}_2^2 - \sigma_0^2|$. By (a) of Proposition 9.5, the first term is $O_{P_0}(\max\{n^{-1/2}, n^{-2\alpha^*/(2\alpha^*+d)}\})$. To bound the second term, let $\mathbf{U} = (\mathbf{Z}\mathbf{V}\mathbf{Z}^T + \mathbf{I}_n)^{-1}$. By equation (33) of page 355 in Searle [25],

$$\begin{aligned} |\mathbb{E}(\tilde{\sigma}_2^2 | \boldsymbol{\theta}_0) - \sigma_0^2| &= |n^{-1}\sigma_0^2 \text{tr}(\mathbf{U}) - \sigma_0^2| + n^{-1}(\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\xi})^T \mathbf{U} (\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\xi}) \\ &\lesssim n^{-1}[\text{tr}(\mathbf{I}_n - \mathbf{U}) + (\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0)^T \mathbf{U} (\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0) \\ &\quad + (\mathbf{Z}\boldsymbol{\theta}_0 - \mathbf{Z}\boldsymbol{\xi})^T \mathbf{U} (\mathbf{Z}\boldsymbol{\theta}_0 - \mathbf{Z}\boldsymbol{\xi})], \end{aligned} \quad (8.4)$$

where we have used $(\mathbf{x} + \mathbf{y})^T \mathbf{G} (\mathbf{x} + \mathbf{y}) \leq 2\mathbf{x}^T \mathbf{G} \mathbf{x} + 2\mathbf{y}^T \mathbf{G} \mathbf{y}$ for any matrix $\mathbf{G} \geq \mathbf{0}$ (Cauchy-Schwartz and the geometric-arithmetic inequalities). Let $\mathbf{P}_Z = \mathbf{Z}(\mathbf{Z}^T \mathbf{Z})^{-1} \mathbf{Z}^T$ be the orthogonal projection matrix. For matrices $\mathbf{Q}, \mathbf{C}, \mathbf{T}, \mathbf{W}$, the binomial inverse theorem (see Theorem 18.2.8 of Harville [13]) says that

$$(\mathbf{Q} + \mathbf{C}\mathbf{T}\mathbf{W})^{-1} = \mathbf{Q}^{-1} - \mathbf{Q}^{-1} \mathbf{C} (\mathbf{T}^{-1} + \mathbf{W}\mathbf{Q}^{-1} \mathbf{C})^{-1} \mathbf{W}\mathbf{Q}^{-1}.$$

Applying the above twice to \mathbf{U} yields

$$(\mathbf{Z}\mathbf{V}\mathbf{Z}^T + \mathbf{I}_n)^{-1} = \mathbf{I}_n - \mathbf{Z}(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{Z}^T = \mathbf{I}_n - \mathbf{P}_Z + \mathbf{M}, \quad (8.5)$$

where $\mathbf{M} = \mathbf{Z}(\mathbf{Z}^T \mathbf{Z})^{-1} [\mathbf{V} + (\mathbf{Z}^T \mathbf{Z})^{-1}]^{-1} (\mathbf{Z}^T \mathbf{Z})^{-1} \mathbf{Z}^T \geq \mathbf{0}$. Hence the first term in (8.4) is $n^{-1} \text{tr}(\mathbf{P}_Z - \mathbf{M}) \leq n^{-1} \text{tr}(\mathbf{P}_Z) = (W + 1)/n$. Note $\mathbf{U} \leq \mathbf{I}_n$ since $\mathbf{Z}\mathbf{V}\mathbf{Z}^T \geq \mathbf{0}$, and the second term in (8.4) is bounded by

$$n^{-1} \|\mathbf{U}\|_{(2,2)} \|\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0\|^2 \leq \|\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0\|_\infty^2 \lesssim \sum_{k=1}^d \delta_{n,k}^{2\alpha_k},$$

in view of (8.3). By (8.5) and $(\mathbf{I} - \mathbf{P}_Z)\mathbf{Z} = \mathbf{0}$, the last term in (8.4) is $n^{-1}(\boldsymbol{\theta}_0 - \boldsymbol{\xi})^T [\mathbf{V} + (\mathbf{Z}^T \mathbf{Z})^{-1}]^{-1} (\boldsymbol{\theta}_0 - \boldsymbol{\xi}) \leq n^{-1} \sum_{j=0}^W \delta_n^{i_j} (\theta_{0,i_j} - \xi_{i_j})^2 = O_{P_0}(n^{-1})$, since $\delta_{n,k} = o(1)$, $k = 1, \dots, d$, $\theta_{0,i_j} = O_{P_0}(1)$ and $\xi_{i_j} = O(1)$ by assumption on the prior for any $0 \leq j \leq W$. Combining the three bounds established into (8.4), we obtain $|\mathbb{E}(\tilde{\sigma}_2^2 | \boldsymbol{\theta}_0) - \sigma_0^2| \lesssim n^{-1} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k}$.

We write $n\tilde{\sigma}_2^2 = (\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\xi})^T \mathbf{U}(\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\xi}) + 2(\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\xi})^T \mathbf{U}\boldsymbol{\varepsilon} + \boldsymbol{\varepsilon}^T \mathbf{U}\boldsymbol{\varepsilon}$ by substituting $\mathbf{Y} = \mathbf{F}_0 + \boldsymbol{\varepsilon}$. Observe that $\boldsymbol{\varepsilon}$ and $\boldsymbol{\theta}_0$ are independent by definition. Using the inequality $\text{Var}(A_1 + A_2) \leq 2\text{Var}(A_1) + 2\text{Var}(A_2)$ (from Cauchy-Schwartz and geometric-arithmetic inequalities), we conclude that $\text{Var}(\tilde{\sigma}_2^2|\boldsymbol{\theta}_0)$ is bounded up to a constant multiple by

$$n^{-2}[(\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0)^T \mathbf{U}^2(\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0) + (\mathbf{Z}\boldsymbol{\theta}_0 - \mathbf{Z}\boldsymbol{\xi})^T \mathbf{U}^2(\mathbf{Z}\boldsymbol{\theta}_0 - \mathbf{Z}\boldsymbol{\xi}) + \text{Var}(\boldsymbol{\varepsilon}^T \mathbf{U}\boldsymbol{\varepsilon})]. \quad (8.6)$$

In view of (8.3) and $\mathbf{U} \leq \mathbf{I}_n$, the first term is bounded by $n^{-2}\|\mathbf{U}\|_{(2,2)}^2\|\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0\|^2 \leq n^{-1}\|\mathbf{F}_0 - \mathbf{Z}\boldsymbol{\theta}_0\|_\infty^2 \lesssim n^{-1}\sum_{k=1}^d \delta_{n,k}^{2\alpha_k}$. Observe that since $\mathbf{V} \geq \mathbf{0}$,

$$\begin{aligned} \mathbf{Z}^T \mathbf{M}^2 \mathbf{Z} &= [\mathbf{V} + (\mathbf{Z}^T \mathbf{Z})^{-1}]^{-1} (\mathbf{Z}^T \mathbf{Z})^{-1} [\mathbf{V} + (\mathbf{Z}^T \mathbf{Z})^{-1}]^{-1} \\ &\leq [\mathbf{V} + (\mathbf{Z}^T \mathbf{Z})^{-1}]^{-1} \leq \mathbf{Z}^T \mathbf{Z}. \end{aligned} \quad (8.7)$$

Using (8.5), idempotency of $\mathbf{I}_n - \mathbf{P}_Z$ and $(\mathbf{I}_n - \mathbf{P}_Z)\mathbf{Z} = \mathbf{0}$, the second term in (8.6) is $n^{-2}(\boldsymbol{\theta}_0 - \boldsymbol{\xi})^T \mathbf{Z}^T (\mathbf{I}_n - \mathbf{P}_Z + \mathbf{M})^2 \mathbf{Z}(\boldsymbol{\theta}_0 - \boldsymbol{\xi})$, which is

$$n^{-2}(\boldsymbol{\theta}_0 - \boldsymbol{\xi})^T \mathbf{Z}^T \mathbf{M}^2 \mathbf{Z}(\boldsymbol{\theta}_0 - \boldsymbol{\xi}) \leq n^{-2}(\boldsymbol{\theta}_0 - \boldsymbol{\xi})^T \mathbf{Z}^T \mathbf{Z}(\boldsymbol{\theta}_0 - \boldsymbol{\xi}), \quad (8.8)$$

in view of (8.7). By (8.9) in the proof of Lemma 8.1, we can write $\mathbf{Z}^T \mathbf{Z} = n_2 \boldsymbol{\Delta} \mathbf{A} \boldsymbol{\Delta}$ where $\boldsymbol{\Delta} = \text{diag}\{\delta_n^{i_j} : j = 0, \dots, W\}$ and $\mathbf{A} \rightarrow \text{E}\mathbf{U}\mathbf{U}^T$ in probability entry-wise. By the continuous mapping theorem, $\|\mathbf{A}\|_{(2,2)} \rightarrow \|\text{E}\mathbf{U}\mathbf{U}^T\|_{(2,2)}$ in probability. The entries of $\text{E}\mathbf{U}\mathbf{U}^T$ are mixed moments of $\text{Uniform}[-v, v]$ and hence is nonsingular with $\|\text{E}\mathbf{U}\mathbf{U}^T\|_{(2,2)} > 0$. Since $\|\boldsymbol{\Delta}\|_{(2,2)} = 1$ and $n_2 \leq n$, the right hand side of (8.8) is bounded by

$$n_2 n^{-2} \|\mathbf{A}\|_{(2,2)} \|\boldsymbol{\Delta}\|_{(2,2)}^2 \|\boldsymbol{\theta}_0 - \boldsymbol{\xi}\|^2 = O_{P_0}(n^{-1}),$$

because $\|\boldsymbol{\theta}_0 - \boldsymbol{\xi}\| \leq \|\boldsymbol{\theta}_0\| + \|\boldsymbol{\xi}\| = O_{P_0}(1)$. By Lemma A.10 of Yoo and Ghosal [34] with $\|\mathbf{U}\|_{(2,2)} \leq 1$ and Gaussian errors by Assumption 1, the last term in (8.6) is $O(1/n)$. Combining this with the three bounds established above, we obtain $\text{Var}(\tilde{\sigma}_2^2|\boldsymbol{\theta}_0) = O_{P_0}(1/n)$. Therefore, the mean square error is $\text{E}_0(\tilde{\sigma}_2^2 - \sigma_0^2)^2 = \text{E}\{\text{E}[(\tilde{\sigma}_2^2 - \sigma_0^2)^2|\boldsymbol{\theta}_0]\} \lesssim n^{-1} + \sum_{k=1}^d \delta_{n,k}^{4\alpha_k}$.

To prove (b), observe that $\text{E}(\sigma^2|\mathbf{Y}) \lesssim n^{-1} + \tilde{\sigma}_*^2$ and $\text{Var}(\sigma^2|\mathbf{Y}) \lesssim n^{-3} + n^{-1}\tilde{\sigma}_*^4$. Therefore by Markov's inequality, the second stage posterior of σ^2 concentrates around the second stage empirical Bayes estimator $\tilde{\sigma}_*^2$, and thus (b) will inherit the rate from (a) as established above. \square

We break the proof of Theorem 6.2 into a series of steps. First, let us enumerate the elements of $\{\mathbf{i} : \mathbf{i} \leq \mathbf{m}_\alpha\}$ as $\{\mathbf{i}_0, \dots, \mathbf{i}_W\}$ with $W + 1 = \prod_{k=1}^d \alpha_k$. For the rest of this section, we follow this convention and index the entries of vectors $\boldsymbol{\xi}$, $\boldsymbol{\theta}$ and $\boldsymbol{\theta}_0$ by elements of $\{\mathbf{i} : \mathbf{i} \leq \mathbf{m}_\alpha\}$. For matrices $\mathbf{Z}^T \mathbf{Z}$ and \mathbf{V} , we enumerate their rows and columns starting from 0 and ending at W . We note that in our first and second stage sampling plans, $n_1 \asymp n \asymp n_2$.

The first key step is to derive sharp upper bounds for the posterior mean and variance, which will involve upper bounding the entries of $(\mathbf{Z}^T \mathbf{Z})^{-1}$. These calculations are made simpler by centering the design points so that they are uniformly distributed around zero in each co-ordinate, and $(\mathbf{Z}^T \mathbf{Z})^{-1}$ will not depend on $\tilde{\boldsymbol{\mu}}$. The following lemma describes the asymptotic behavior of the entries of $(\mathbf{Z}^T \mathbf{Z})^{-1}$ when the second stage samples are collected under uniform random sampling. In the following, we define $\boldsymbol{\delta}_n = (\delta_{n,1}, \dots, \delta_{n,d})^T$ and write $\boldsymbol{\delta}_n^{\mathbf{i}}$ to mean $\prod_{k=1}^d \delta_{n,k}^{i_k}$.

Lemma 8.1. *As $n \rightarrow \infty$, $\mathbf{Z}^T \mathbf{Z}$ is invertible with probability tending to 1. Moreover for $a, b = 0, \dots, W$, we have $[(\mathbf{Z}^T \mathbf{Z})^{-1}]_{ab} = O_P(n^{-1} \delta_n^{-(i_a+i_b)})$.*

Proof. After centering, second stage samples are distributed as $\mathbf{z}_i = (z_{i1}, \dots, z_{id})^T \stackrel{i.i.d.}{\sim} \text{Uniform}\left(\prod_{k=1}^d [-\delta_{n,k}, \delta_{n,k}]\right)$, $i = 1, \dots, n_2$, and $z_{ik} \sim \text{Uniform}[-\delta_{n,k}, \delta_{n,k}]$. Thus,

$$\mathbf{Z}^T \mathbf{Z} = n_2 \begin{pmatrix} a_{00} & a_{01} \delta_n^{i_1} & \cdots & a_{0W} \delta_n^{i_W} \\ a_{10} \delta_n^{i_1} & a_{11} \delta_n^{2i_1} & \cdots & a_{1W} \delta_n^{i_1+i_W} \\ \vdots & \vdots & \ddots & \vdots \\ a_{W0} \delta_n^{i_W} & a_{W1} \delta_n^{i_W+i_1} & \cdots & a_{WW} \delta_n^{2i_W} \end{pmatrix} := n_2 \mathbf{\Delta} \mathbf{A} \mathbf{\Delta} \quad (8.9)$$

with the (i, j) -entry of matrix \mathbf{A} being $a_{ij} = n_2^{-1} \sum_{k=1}^{n_2} \mathbf{U}_k^{i_i} \mathbf{U}_k^{i_j}$ where $\mathbf{U}_k = (U_{k1}, \dots, U_{kd})^T \stackrel{i.i.d.}{\sim} \text{Uniform}[-1, 1]^d$, and $\mathbf{\Delta} = \text{diag}\{\delta_n^{i_j} : j = 0, \dots, W\}$. Define $\mathbb{U} = (\mathbf{U}^{i_0}, \dots, \mathbf{U}^{i_W})^T$ for $\mathbf{U} = (U, \dots, U)^T \sim \text{Uniform}[-1, 1]^d$. By the law of large numbers, we have that \mathbf{A} converges in probability to $\mathbb{E} \mathbb{U} \mathbb{U}^T$ entry-wise, and hence $\mathbb{E} \mathbb{U} \mathbb{U}^T - \epsilon \mathbf{I} \leq \mathbf{A} \leq \mathbb{E} \mathbb{U} \mathbb{U}^T + \epsilon \mathbf{I}$ for a sufficiently small $\epsilon > 0$. Observe that entries of $\mathbb{E} \mathbb{U} \mathbb{U}^T$ are mixed moments of $U \sim \text{Uniform}[-1, 1]$ and hence is positive definite. Then $\mathbb{E} \mathbb{U} \mathbb{U}^T - \epsilon \mathbf{I}$ is invertible when ϵ is smaller than the minimum eigenvalue of $\mathbb{E} \mathbb{U} \mathbb{U}^T$. Thus for sufficiently small $\epsilon > 0$,

$$n_2^{-1} \mathbf{\Delta}^{-1} (\mathbb{E} \mathbb{U} \mathbb{U}^T + \epsilon \mathbf{I})^{-1} \mathbf{\Delta}^{-1} \leq (\mathbf{Z}^T \mathbf{Z})^{-1} \leq n_2^{-1} \mathbf{\Delta}^{-1} (\mathbb{E} \mathbb{U} \mathbb{U}^T - \epsilon \mathbf{I})^{-1} \mathbf{\Delta}^{-1}.$$

Let u^{ij} be the (i, j) th entry of $(\mathbb{E} \mathbb{U} \mathbb{U}^T)^{-1}$ and recall that $n_2 \geq cn$ for some constant $c > 0$. It then follows that for $a = 0, \dots, W$, $[(\mathbf{Z}^T \mathbf{Z})^{-1}]_{aa}$ is $O_P(n_2^{-1} u^{aa} \delta_n^{-(i_a+i_a)}) = O_P(n^{-1} \delta_n^{-(i_a+i_a)})$. Using the fact that for positive definite \mathbf{G} , $g_{ij} \leq \sqrt{g_{ii} g_{jj}}$ by the Cauchy-Schwartz inequality, $[(\mathbf{Z}^T \mathbf{Z})^{-1}]_{ab}$ with $a, b = 0, \dots, W$, is bounded above by

$$\sqrt{[(\mathbf{Z}^T \mathbf{Z})^{-1}]_{aa} [(\mathbf{Z}^T \mathbf{Z})^{-1}]_{bb}} = O_P(n^{-1} \delta_n^{-(i_a+i_b)}). \quad \square$$

Let $\mathcal{K}_n := [\sigma_0^2 - m_n \nu_n, \sigma_0^2 + m_n \nu_n]$ where m_n is any sequence going to infinity (e.g., slowly varying such as $\log n$) and $\nu_n = \max\{n^{-1/2}, n^{-2\alpha^*/(2\alpha^*+d)}, \sum_{k=1}^d \delta_{n,k}^{2\alpha_k}\}$. Recall that $\mathcal{Q} = \{\mathbf{x} : |x_k| \leq \delta_{n,k}, k = 1, \dots, d\}$ is the centered second stage credible set/sampling region.

Lemma 8.2. *For any $\mathbf{i} \leq \mathbf{m}_\alpha$ and $\sigma^2 \in \mathcal{K}_n$,*

$$\mathbb{E}[(\theta_{\mathbf{i}} - \theta_{0,\mathbf{i}})^2 | \mathbf{Y}, \sigma^2] = O_{P_0} \left[\prod_{k=1}^d \delta_{n,k}^{-2i_k} \left(\frac{1}{n} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right) \right].$$

Proof. Let $0 \leq h \leq W$. Since $\mathbf{V} > \mathbf{0}$ by assumption, we have by Lemma 8.1 that $\sup_{\sigma^2 \in \mathcal{K}_n} \text{Var}(\theta_{i_h} | \mathbf{Y}, \sigma^2)$ is

$$[\sigma_0^2 + o(1)] [(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}]_{hh} \lesssim [(\mathbf{Z}^T \mathbf{Z})^{-1}]_{hh} \lesssim n^{-1} \delta_n^{-2i_h}. \quad (8.10)$$

Now the bias for the conditional posterior mean in vector form is

$$\begin{aligned} \mathbb{E}(\boldsymbol{\theta}|\mathbf{Y}, \sigma^2) - \boldsymbol{\theta}_0 &= (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} (\mathbf{Z}^T \mathbf{Y} + \mathbf{V}^{-1} \boldsymbol{\xi}) - \boldsymbol{\theta}_0 \\ &= (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} [\mathbf{Z}^T \boldsymbol{\varepsilon} + \mathbf{Z}^T (\mathbf{F}_0 - \mathbf{Z} \boldsymbol{\theta}_0) + \mathbf{V}^{-1} (\boldsymbol{\xi} - \boldsymbol{\theta}_0)]. \end{aligned} \quad (8.11)$$

Following the same reasoning as in (8.10), the h th diagonal entry of the covariance matrix $\sigma_0^2 (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{Z}^T \mathbf{Z} (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}$ of $(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \boldsymbol{\varepsilon}$ is

$$\sigma_0^2 [(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{Z}^T \mathbf{Z} (\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}]_{hh} \lesssim n^{-1} \delta_n^{-2i_h}.$$

Since $\mathbb{E}_0(\boldsymbol{\varepsilon}) = \mathbf{0}$, it follows from Markov's inequality that the h th entry of $(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{Z}^T \boldsymbol{\varepsilon}$ is $O_{P_0}(n^{-1/2} \delta_n^{-i_h})$ for $0 \leq h \leq W$.

Let β_{ij} be the (i, j) th element of $(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}$, κ_{ij} be the (i, j) th element of \mathbf{V}^{-1} and γ_i be the i th entry of $\mathbf{F}_0 - \mathbf{Z} \boldsymbol{\theta}_0$. By (8.3), we have uniformly over $1 \leq i \leq n$ that $|\gamma_i| \lesssim \sum_{k=1}^d \delta_{n,k}^{\alpha_k}$. Now using the fact that for positive definite \mathbf{G} , $g_{hj} \leq \sqrt{g_{hh} g_{jj}}$ by the Cauchy-Schwarz inequality, we have for $0 \leq h, j \leq W$,

$$\begin{aligned} [(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}]_{hj} &\leq \sqrt{[(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}]_{hh} [(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1}]_{jj}} \\ &\leq \sqrt{[(\mathbf{Z}^T \mathbf{Z})^{-1}]_{hh} [(\mathbf{Z}^T \mathbf{Z})^{-1}]_{jj}} \lesssim n^{-1} \delta_n^{-(i_h + i_j)}. \end{aligned}$$

Since $\mathbf{z}_j \in \mathcal{Q}$, $j = 1, \dots, n_2$, we have $|\mathbf{z}_j^i| \leq \delta_n^i$ for $i \leq \mathbf{m}_\alpha$. Therefore, since $n_2 \leq n$, $[(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{Z}^T (\mathbf{F}_0 - \mathbf{Z} \boldsymbol{\theta}_0)]_h$ is

$$\beta_{h0} \sum_{j=1}^{n_2} \mathbf{z}_j^{i_0} \gamma_j + \dots + \beta_{hW} \sum_{j=1}^{n_2} \mathbf{z}_j^{i_W} \gamma_j \lesssim \delta_n^{-i_h} \sum_{k=1}^d \delta_{n,k}^{\alpha_k}.$$

It remains to bound each entry of the last term in (8.11). Since $|\theta_{0,i_j} - \xi_{0,i_j}| \leq |\theta_{0,i_j}| + |\xi_{0,i_j}| = O_{P_0}(1)$ for $j = 0, \dots, W$, then Lemma 8.1 and the choice of \mathbf{V} implies that $[(\mathbf{Z}^T \mathbf{Z} + \mathbf{V}^{-1})^{-1} \mathbf{V}^{-1} (\boldsymbol{\xi} - \boldsymbol{\theta}_0)]_h$ is

$$\beta_{h0} \sum_{j=0}^W \kappa_{0j} (\xi_{i_j} - \theta_{0,i_j}) + \dots + \beta_{hW} \sum_{j=0}^W \kappa_{Wj} (\xi_{i_j} - \theta_{0,i_j}) \lesssim n^{-1} \delta_n^{-i_h}.$$

Combining the bounds established back into (8.11), the squared bias $(\mathbb{E}(\theta_{i_h} | \mathbf{Y}, \sigma^2) - \theta_{0,i_h})^2$ is $O_{P_0} \left[n^{-2} \delta_n^{-2i_h} + \delta_n^{-2i_h} \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right]$. The result follows in view of the bounds established and (8.10). \square

Lemma 8.3. *Uniformly over $\|f_0\|_{\alpha, \infty} \leq R$, for any $\mathbf{r} \leq \mathbf{m}_\alpha$, $\mathbf{x} \in \mathcal{Q}$ and $m_n \rightarrow \infty$, $\mathbb{E}_0 \Pi(|D^{\mathbf{r}} f_{\boldsymbol{\theta}}(\mathbf{x}) - D^{\mathbf{r}} f_{0,\mathbf{z}}(\mathbf{x})| > m_n \epsilon_{n,\mathbf{r}} | \mathbf{Y}) \rightarrow 0$, where $\epsilon_{n,\mathbf{r}} := \delta_n^{-\mathbf{r}} (n^{-1/2} + \sum_{k=1}^d \delta_{n,k}^{\alpha_k})$.*

Proof. In view of (6.4),

$$D^{\mathbf{r}} f_{\boldsymbol{\theta}}(\mathbf{x}) - D^{\mathbf{r}} f_{\boldsymbol{\theta}_0}(\mathbf{x}) = \mathbf{r}! (\theta_{\mathbf{r}} - \theta_{0,\mathbf{r}}) + \sum_{\mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha, \mathbf{i} \neq \mathbf{r}} \frac{\mathbf{i}!}{(\mathbf{i} - \mathbf{r})!} (\theta_{\mathbf{i}} - \theta_{0,\mathbf{i}}) \mathbf{x}^{\mathbf{i} - \mathbf{r}}.$$

Observe that for any $\mathbf{x} \in \mathcal{Q}$, $|\mathbf{x}^{i-r}| \leq \delta_n^{i-r}$. Also, by noting that $r_k \leq i_k \leq \alpha_k - 1$ for $k = 1, \dots, d$, we have both $\mathbf{r}!, \mathbf{i}! \leq \prod_{k=1}^d (\alpha_k - 1)$. Using the fact $(\sum_{i=1}^n |b_i|)^p \leq n^{p-1} \sum_{i=1}^n |b_i|^p$ for $p \geq 1$, $|D^r f_\theta(\mathbf{x}) - D^r f_{\theta_0}(\mathbf{x})|^2$ is bounded above up to some constant by

$$|\theta_{\mathbf{r}} - \theta_{0,\mathbf{r}}|^2 + \sum_{\mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha, \mathbf{i} \neq \mathbf{r}} |\theta_{\mathbf{i}} - \theta_{0,\mathbf{i}}|^2 \delta_n^{2i-2r}. \quad (8.12)$$

Therefore, for any $\mathbf{r} \leq \mathbf{m}_\alpha$ and any $\mathbf{x} \in \mathcal{Q}$, we have uniformly over $\|f_0\|_{\alpha,\infty} \leq R$ that $\mathbb{E}_0 \sup_{\sigma^2 \in \mathcal{K}_n} \mathbb{E}(|D^r f_\theta(\mathbf{x}) - D^r f_{\theta_0}(\mathbf{x})|^2 | \mathbf{Y}, \sigma^2)$ is bounded up to a constant multiple by

$$\mathbb{E}_0 \sup_{\sigma^2 \in \mathcal{K}_n} \mathbb{E}[(\theta_{\mathbf{r}} - \theta_{0,\mathbf{r}})^2 | \mathbf{Y}, \sigma^2] + \sum_{\mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha, \mathbf{i} \neq \mathbf{r}} \delta_n^{2i-2r} \mathbb{E}_0 \sup_{\sigma^2 \in \mathcal{K}_n} \mathbb{E}[(\theta_{\mathbf{i}} - \theta_{0,\mathbf{i}})^2 | \mathbf{Y}, \sigma^2]. \quad (8.13)$$

In view of Lemma 8.2, the first term is bounded by $\delta_n^{-2r}(n^{-1} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k})$. By noting that the sum over $\{\mathbf{i} : \mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha, \mathbf{i} \neq \mathbf{r}\}$ has at most $\prod_{k=1}^d \alpha_k$ terms, another application of Lemma 8.2 implies that the second term in (8.13) is bounded above by

$$\sum_{\mathbf{r} \leq \mathbf{i} \leq \mathbf{m}_\alpha, \mathbf{i} \neq \mathbf{r}} \delta_n^{2i-2r} \left[\delta_n^{-2i} \left(\frac{1}{n} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right) \right] \lesssim \delta_n^{-2r} \left(\frac{1}{n} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right).$$

Using the inequality $|a+b|^r \leq \max(1, 2^{r-1})(|a|^r + |b|^r)$, $r > 0$ and (2.2), we have $|D^r f_{\theta_0}(\mathbf{x}) - D^r f_{0,z}(\mathbf{x})|$ is

$$\begin{aligned} |D^r T_{\mu_0}(\mathbf{x} + \tilde{\boldsymbol{\mu}}) - D^r f_0(\mathbf{x} + \tilde{\boldsymbol{\mu}})| &\lesssim \sum_{k=1}^d |x_k + \tilde{\mu}_k - \mu_{0,k}|^{\alpha_k - r_k} \\ &\lesssim \sum_{k=1}^d \delta_{n,k}^{\alpha_k - r_k} + \sum_{k=1}^d |\tilde{\mu}_k - \mu_{0,k}|^{\alpha_k - r_k}, \end{aligned} \quad (8.14)$$

and $\mathbb{E}_0 |D^r f_{\theta_0}(\mathbf{x}) - D^r f_{0,z}(\mathbf{x})|^2 \lesssim \sum_{k=1}^d \delta_{n,k}^{2\alpha_k - 2r_k}$ uniformly in $\|f_0\|_{\alpha,\infty} \leq R$ by (8.3).

Define $P_{n,r}(\mathbf{x}) := \mathbb{E}_0 \sup_{\sigma^2 \in \mathcal{K}_n} \mathbb{E}[(D^r f_\theta(\mathbf{x}) - D^r f_{0,z}(\mathbf{x}))^2 | \mathbf{Y}, \sigma^2]$. Combining the bounds established and (8.13), we have uniformly over $\|f_0\|_{\alpha,\infty} \leq R$,

$$\begin{aligned} P_{n,r}(\mathbf{x}) &\lesssim \mathbb{E}_0 \sup_{\sigma^2 \in \mathcal{K}_n} \mathbb{E}(|D^r f_\theta(\mathbf{x}) - D^r f_{\theta_0}(\mathbf{x})|^2 | \mathbf{Y}, \sigma^2) + \mathbb{E}_0 |D^r f_{\theta_0}(\mathbf{x}) - D^r f_{0,z}(\mathbf{x})|^2 \\ &\lesssim \delta_n^{-2r} \left(\frac{1}{n} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right) + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k - 2r_k} \lesssim \epsilon_{n,r}^2. \end{aligned}$$

By Proposition 6.1, $P_0(\tilde{\sigma}_*^2 \in \mathcal{K}_n) \rightarrow 1$ as $n \rightarrow \infty$ uniformly in $\|f_0\|_{\alpha,\infty} \leq R$. For the empirical Bayes posterior $\Pi(\cdot | \mathbf{Y}) \equiv \Pi_{\tilde{\sigma}_*}(\cdot | \mathbf{Y})$, by Markov's inequality, we have for any $m_n \rightarrow \infty$,

$$\mathbb{E}_0 \Pi_{\tilde{\sigma}_*}(|D^r f_\theta(\mathbf{x}) - D^r f_{0,z}(\mathbf{x})| > m_n \epsilon_{n,r} | \mathbf{Y}) \leq \frac{P_{n,r}(\mathbf{x})}{m_n^2 \epsilon_{n,r}^2} + o(1) \rightarrow 0, \quad (8.15)$$

uniformly over $\|f_0\|_{\alpha,\infty} \leq R$. For the hierarchical Bayes procedure, we have for any $m_n \rightarrow \infty$ that $E_0\Pi(|D^r f_{\boldsymbol{\theta}}(\mathbf{x}) - D^r f_{0,z}(\mathbf{x})| > m_n \epsilon_{n,r} | \mathbf{Y})$ is uniformly over $\|f_0\|_{\alpha,\infty} \leq R$ bounded above by

$$P_{n,r}(\mathbf{x}) / (m_n \epsilon_{n,r})^2 + E_0\Pi(\sigma^2 \notin \mathcal{K}_n | \mathbf{Y}). \quad (8.16)$$

The first term is $o(1)$ since $P_{n,r}(\mathbf{x}) \lesssim \epsilon_{n,r}^2$, while the second term goes to zero by Proposition 6.1. \square

An immediate consequence of the previous lemma is the following:

Corollary 8.4. *Uniformly in $\|f_0\|_{\alpha,\infty} \leq R$, we have for any $\mathbf{r} \leq \mathbf{m}_\alpha$ and $m_n \rightarrow \infty$ that $E_0\Pi(\|D^r f_{\boldsymbol{\theta}} - D^r f_{0,z}\|_\infty > m_n \epsilon_{n,r} | \mathbf{Y}) \rightarrow 0$.*

Proof. By (8.12), we have

$$\begin{aligned} \|D^r f_{\boldsymbol{\theta}} - D^r f_{\boldsymbol{\theta}_0}\|_\infty &= \sup_{\mathbf{x} \in \mathcal{Q}} |D^r f_{\boldsymbol{\theta}}(\mathbf{x}) - D^r f_{\boldsymbol{\theta}_0}(\mathbf{x})| \\ &\lesssim |\theta_{\mathbf{r}} - \theta_{0,\mathbf{r}}| + \sum_{r \leq i \leq \mathbf{m}_\alpha, i \neq r} |\theta_i - \theta_{0,i}| \delta_n^{i-r}. \end{aligned} \quad (8.17)$$

Hence, the upper bound (8.13) is applicable and uniformly over $\|f_0\|_{\alpha,\infty} \leq R$, we will have $E_0 \sup_{\sigma^2 \in \mathcal{K}_n} E[\|D^r f_{\boldsymbol{\theta}} - D^r f_{\boldsymbol{\theta}_0}\|_\infty^2 | \mathbf{Y}, \sigma^2] \lesssim \delta_n^{-2r} (n^{-1} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k})$. Moreover, since the bound in (8.14) is uniform for all $\mathbf{x} \in \mathcal{Q}$, this implies that $E_0 \|D^r f_{\boldsymbol{\theta}_0} - D^r f_{0,z}\|_\infty^2 \lesssim \sum_{k=1}^d \delta_{n,k}^{2\alpha_k - 2r_k}$. Therefore, we conclude that uniformly over $\|f_0\|_{\alpha,\infty} \leq R$,

$$\begin{aligned} E_0 \sup_{\sigma^2 \in \mathcal{K}_n} E[\|D^r f_{\boldsymbol{\theta}} - D^r f_{0,z}\|_\infty^2 | \mathbf{Y}, \sigma^2] \\ &\lesssim E_0 \sup_{\sigma^2 \in \mathcal{K}_n} E[\|D^r f_{\boldsymbol{\theta}} - D^r f_{\boldsymbol{\theta}_0}\|_\infty^2 | \mathbf{Y}, \sigma^2] + E_0 \|D^r f_{\boldsymbol{\theta}_0} - D^r f_{0,z}\|_\infty^2 \\ &\lesssim \delta_n^{-2r} \left(\frac{1}{n} + \sum_{k=1}^d \delta_{n,k}^{2\alpha_k} \right). \end{aligned}$$

The empirical and hierarchical posterior contraction rates then follow from (8.15) and (8.16) with absolute values replaced by sup-norms. \square

We are now ready to prove Theorem 6.2.

Proof of Theorem 6.2. We shall prove only for the empirical Bayes case; the hierarchical Bayes case follows the same steps. Recall that $\boldsymbol{\mu} = \tilde{\boldsymbol{\mu}} + \boldsymbol{\mu}_z$. As a consequence of Theorem 4.1 and our choice of $\delta_{n,k}$, $k = 1, \dots, d$, we have $P_0(\boldsymbol{\mu}_0 - \tilde{\boldsymbol{\mu}} \in \mathcal{Q}) \rightarrow 1$. Therefore by (4.1),

$$\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| = \|\boldsymbol{\mu}_z - (\boldsymbol{\mu}_0 - \tilde{\boldsymbol{\mu}})\| \leq \frac{\sqrt{d}}{\lambda_0} \max_{1 \leq k \leq d} \sup_{\mathbf{x} \in \mathcal{Q}} |D_k f_{\boldsymbol{\theta}}(\mathbf{x}) - D_k f_{0,z}(\mathbf{x})|.$$

Define $\tau_{n,k} := \delta_{n,k}^{-1}(n^{-1/2} + \sum_{k=1}^d \delta_{n,k}^{\alpha_k})$. Using this bound and Corollary 8.4 with $\mathbf{r} = \mathbf{e}_k$, then for any $m_n \rightarrow \infty$, $\mathbb{E}_0 \Pi(\|\boldsymbol{\mu} - \boldsymbol{\mu}_0\| > m_n \max_{1 \leq k \leq d} \tau_{n,k} | \mathbf{Y})$ is bounded above by

$$\sum_{k=1}^d \mathbb{E}_0 \Pi \left(\|D_k f_{\boldsymbol{\theta}} - D_k f_{0,z}\|_{\infty} > \frac{\lambda_0}{\sqrt{d}} m_n \max_{1 \leq k \leq d} \tau_{n,k} \middle| \mathbf{Y} \right) \rightarrow 0.$$

By definition, $M = f_{\boldsymbol{\theta}}(\boldsymbol{\mu}_z)$ and $M_0 = f_{0,z}(\boldsymbol{\mu}_0 - \tilde{\boldsymbol{\mu}})$. Then by (4.2), $|M - M_0| \leq \sup_{\mathbf{x} \in \mathcal{Q}} |f_{\boldsymbol{\theta}}(\mathbf{x}) - f_{0,z}(\mathbf{x})|$ since $P_0(\boldsymbol{\mu}_0 - \tilde{\boldsymbol{\mu}} \in \mathcal{Q}) \rightarrow 1$ as before. Therefore by Corollary 8.4 with $\mathbf{r} = \mathbf{0}$, we have for $m_n \rightarrow \infty$, $\mathbb{E}_0 \Pi[|M - M_0| > m_n(n^{-1/2} + \sum_{k=1}^d \delta_{n,k}^{\alpha_k}) | \mathbf{Y}] \leq \mathbb{E}_0 \Pi[\|f_{\boldsymbol{\theta}} - f_{0,z}\|_{\infty} > m_n(n^{-1/2} + \sum_{k=1}^d \delta_{n,k}^{\alpha_k}) | \mathbf{Y}] \rightarrow 0$, uniformly over $\|f_0\|_{\alpha, \infty} \leq R$.

To prove the last part, note that $\delta_{n,k} = n^{-1/(2\alpha_k)}$, $k = 1, \dots, d$, comes from equating the two terms in the second stage rates for $\boldsymbol{\mu}$ and M , i.e., $n^{-1/2} = \sum_{k=1}^d \delta_{n,k}^{\alpha_k}$. To solve for $\delta_{n,k}$, take $\delta_{n,k} = (d^{-1}\delta_n)^{1/\alpha_k}$, where δ_n is a positive sequence in n that does not depend on k . It follows that $\delta_n = n^{-1/2}$ and hence $\delta_{n,k} = n^{-1/(2\alpha_k)}$. The condition $\min_{1 \leq k \leq d} \delta_{n,k} = \rho_n \epsilon_n$ or equivalently $\epsilon_n = o(\min_{1 \leq k \leq d} \delta_{n,k})$ is fulfilled when $1/(2\underline{\alpha}) < \alpha^*(1 - \underline{\alpha}^{-1})/(2\alpha^* + d)$ for $\underline{\alpha} = \min_{1 \leq k \leq d} \alpha_k$. By rearranging, we need $2\underline{\alpha}\alpha^* - 4\alpha^* - d > 0$. Since $\underline{\alpha} > 2$ by Assumption 2, we have $2\underline{\alpha}\alpha^* - 4\alpha^* - d = 2(\underline{\alpha} - 2)\alpha^* - d \geq 2\underline{\alpha}^2 - 4\underline{\alpha} - d$. Thus, it suffice to find $\underline{\alpha}$ such that $2\underline{\alpha}^2 - 4\underline{\alpha} - d > 0$ and this is satisfied if $\underline{\alpha} > 1 + \sqrt{1 + d/2}$ under the constraint that $\underline{\alpha} > 2$. \square

9 Appendix

In this section, we collect some auxiliary results and technical lemmas that were used in several places to prove the main theorems in the previous section.

The next two results concern contraction rates and credible band coverage for the regression function f and its derivatives, and they correspond to Theorems 4.4 and 5.3 of Yoo and Ghosal [34] respectively.

Theorem 9.1. *Let $J_{n,k} \asymp (n/\log n)^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$, $k = 1, \dots, d$. Then under Assumption 1, we have for any sequence $m_n \rightarrow \infty$,*

$$\sup_{\|f_0\|_{\alpha, \infty} \leq R} \mathbb{E}_0 \Pi(\|D^{\mathbf{r}} f - D^{\mathbf{r}} f_0\|_{\infty} > m_n (\log n/n)^{\alpha^*\{1 - \sum_{k=1}^d (r_k/\alpha_k)\}/(2\alpha^*+d)} | \mathbf{Y}) \rightarrow 0.$$

In particular, contraction rate for f can be recovered by setting $\mathbf{r} = \mathbf{0}$.

Consider the simultaneous credible band $\{f : \|D^{\mathbf{r}} f - D^{\mathbf{r}} \tilde{f}\|_{\infty} \leq \rho R_{n,r,\gamma}\}$, where the quantile $R_{n,r,\gamma}$ is chosen such that $\Pi(\|D^{\mathbf{r}} f - D^{\mathbf{r}} \tilde{f}\|_{\infty} \leq R_{n,r,\gamma} | \mathbf{Y}) = 1 - \gamma$ and $\rho > 0$ is some large enough constant.

Theorem 9.2. *Let $J_{n,k} \asymp (n/\log n)^{\alpha^*/\{\alpha_k(2\alpha^*+d)\}}$, $k = 1, \dots, d$. Then under Assumption 1,*

$$1. \inf_{\|f_0\|_{\alpha, \infty} \leq R} P_0(\|D^{\mathbf{r}} \tilde{f} - D^{\mathbf{r}} f_0\|_{\infty} \leq \rho R_{n,r,\gamma}) \rightarrow 1,$$

2. $R_{n,\mathbf{r},\gamma} \asymp (\log n/n)^{\alpha^* \{1 - \sum_{k=1}^d (r_k/\alpha_k)\} / (2\alpha^* + d)}$ in P_0 .

Credible bands for f is recovered by $\mathbf{r} = \mathbf{0}$ and for $D^{\mathbf{e}_k} f \equiv D_k f$ by $\mathbf{r} = \mathbf{e}_k$, in the latter case we also write the radius as $R_{n,k,\gamma} = R_{n,\mathbf{e}_k,\gamma}$.

The result below was taken from (3.10) and (3.11) of Yoo and Ghosal [34], and it shows that B-splines despite being a non-orthonormal basis, are approximately orthogonal under our assumption on the design points.

Lemma 9.3. *Let $J = \prod_{k=1}^d J_k$. If the design points were chosen such that (2.3) hold, then for some constants $C_1, C_2, c_1, c_2 > 0$,*

$$C_1(n/J) \leq \mathbf{B}^T \mathbf{B} \leq C_2(n/J) \quad (9.1)$$

$$C_1(n/J) + c_2^{-1} \leq \lambda_{\min}(\mathbf{B}^T \mathbf{B} + \mathbf{\Omega}^{-1}) \leq \lambda_{\max}(\mathbf{B}^T \mathbf{B} + \mathbf{\Omega}^{-1}) \leq C_2(n/J) + c_1^{-1} \quad (9.2)$$

Lemma 9.4. *Let $\mathbf{W}_{\mathbf{r}}$ be the finite difference matrix as seen in (3.1). Under the quasi-uniformity of the knot distribution, we have $\lambda_{\max}(\mathbf{W}_{\mathbf{r}} \mathbf{W}_{\mathbf{r}}^T) = O(\prod_{k=1}^d J_k^{2r_k})$.*

Proof. Note that $\lambda_{\max}(\mathbf{W}_{\mathbf{r}} \mathbf{W}_{\mathbf{r}}^T) = \lambda_{\max}(\mathbf{W}_{\mathbf{r}}^T \mathbf{W}_{\mathbf{r}}) \leq \|\mathbf{W}_{\mathbf{r}}^T \mathbf{W}_{\mathbf{r}}\|_{(2,2)} \lesssim \prod_{k=1}^d J_k^{2r_k}$, where the later upper bound was computed in (7.15) of Yoo and Ghosal [34]. \square

The following proposition shows that the single stage or the first stage (in the setting of two stage procedure) empirical or hierarchical Bayes estimator of σ^2 is consistent. Note that this result corresponds to Proposition 4.1 of Yoo and Ghosal [34].

Proposition 9.5 (First stage error variance). *Suppose $J_{n,k} \asymp (n/\log n)^{\alpha^* / \{\alpha_k(2\alpha^* + d)\}}$. Then uniformly over $\|f_0\|_{\alpha,\infty} \leq R$,*

- (a) *First stage empirical Bayes estimator $\tilde{\sigma}_1$ converges to σ_0 at rate $\max\{n^{-1/2}, n^{-2\alpha^* / (2\alpha^* + d)}\}$ under P_0 .*
- (b) *If inverse gamma prior is used, first stage posterior for σ contracts to σ_0 at the same rate.*
- (c) *If the prior used has continuous and positive density on $(0, \infty)$, then the first stage posterior for σ is consistent.*

References

- [1] Belitser, E., Ghosal, S., and van Zanten, H. (2012). Optimal two-stage procedures for estimating location and size of the maximum of a multivariate regression function. *Ann. Statist.*, 40(6):2850–2876.
- [2] Blum, J. R. (1954). Multidimensional stochastic approximation methods. *Ann. Math. Statist.*, 25(4):737–744.

- [3] Castillo, I. and Nickl, R. (2013). Nonparametric Bernstein–von Mises theorems in Gaussian white noise. *Ann. Statist.*, 41(4):1999–2028.
- [4] Castillo, I. and Nickl, R. (2014). On the Bernstein–von Mises phenomenon for nonparametric Bayes procedures. *Ann. Statist.*, 42(5):1941–1969.
- [5] Chen, H. (1988). Lower rate of convergence for locating a maximum of a function. *Ann. Statist.*, 16(3):1330–1334.
- [6] Cox, D. D. (1993). An analysis of Bayesian inference for nonparametric regression. *Ann. Statist.*, 21(2):903–923.
- [7] de Boor, C. (2001). *A Practical Guide to Splines*. Springer-Verlag New York, Inc., revised edition.
- [8] Dippon, J. (2003). Accelerated randomized stochastic optimization. *Ann. Statist.*, 31(4):1260–1281.
- [9] Donoho, D. L. and Liu, R. C. (1991). Geometrizing rates of convergence, III. *Ann. Statist.*, 19(2):668–701.
- [10] Fabian, V. (1967). Stochastic approximation of minima with improved asymptotic speed. *Ann. Math. Statist.*, 38(1):191–200.
- [11] Facer, M. R. and Müller, H.-G. (2003). Nonparametric estimation of the location of a maximum in a response surface. *J. Multivariate Anal.*, 87(1):191–217.
- [12] Freedman, D. (1999). On the Bernstein-von Mises theorem with infinite dimensional parameters. *Ann. Statist.*, 27(4):1119–1140.
- [13] Harville, D. A. (1997). *Matrix Algebra from a Statistician’s Perspective*. Springer-Verlag New York, Inc.
- [14] Hasminskii, R. Z. (1979). Lower bound for the risks of nonparametric estimates of the mode. In *Contributions to Statistics (J. Jureckova, ed.)*, pages 91–97.
- [15] Jørgensen, M., C. T. Nielsen, N. K., and Skakkeback, N. E. (1985). Parametrische und nichtparametrische modelle für wachstumsdaten. In G. C. Pflug, e., editor, *Med. Informatik und Statistik*, volume 60, pages 74–87. Springer, Berlin.
- [16] Kiefer, J. and Wolfowitz, J. (1952). Stochastic estimation of the maximum of a regression function. *Ann. Math. Statist.*, 23(3):462–466.
- [17] Kim, J. and Pollard, D. (1990). Cube root asymptotics. *Ann. Statist.*, 18(1):191–219.
- [18] Knapik, B. T., van der Vaart, A. W., and van Zanten, J. H. (2011). Bayesian inverse problems with Gaussian priors. *Ann. Statist.*, 39(5):2626–2657.

- [19] Lan, Y., Banerjee, M., and Michailidis, G. (2009). Change-point estimation under adaptive sampling. *Ann. Statist.*, 37(4):1752–1791.
- [20] Mokkadem, A. and Pelletier, M. (2007). A companion for the Kiefer-Wolfowitz-Blum stochastic approximation algorithm. *Ann. Statist.*, 35(4):1749–1772.
- [21] Müller, H.-G. (1985). Kernel estimators of zeros and of location and size of extrema of regression function. *Scand. J. Stat.*, 12(3):221–232.
- [22] Müller, H.-G. (1989). Adaptive nonparametric peak estimation. *Ann. Statist.*, 17(3):1053–1069.
- [23] Polyak, B. T. and Tsybakov, A. B. (1990). Optimal orders of accuracy for search algorithm of stochastic optimization. *Probl. Inf. Transm.*, 26(2):126–133.
- [24] Schumaker, L. (2007). *Spline Functions: Basic Theory*. Cambridge University Press, New York, third edition.
- [25] Searle, S. R. (1982). *Matrix Algebra Useful for Statistics*. John Wiley and Sons, Inc.
- [26] Shoung, J.-M. and Zhang, C.-H. (2001). Least squares estimators of the mode of a unimodal regression function. *Ann. Statist.*, 29(3):648–665.
- [27] Silverman, B. W. (1985). Some aspects of the spline smoothing approach to nonparametric regression curve fitting (with discussion). *J. Roy. Statist. Soc. Ser. B*, 47(1):1–52.
- [28] Szabo, B., van der Vaart, A. W., and van Zanten, J. H. (2015). Frequentist coverage of adaptive nonparametric Bayesian credible sets. *Ann. Statist.*, 43(4):1391–1428.
- [29] Tang, R., Banerjee, M., and Michailidis, G. (2011). A two-stage hybrid procedure for estimating an inverse regression function. *Ann. Statist.*, 39(2):956–989.
- [30] Tsybakov, A. B. (1990). Recurrent estimation of the mode of a multidimensional distribution. *Probl. Inf. Transm.*, 26(1):31–37.
- [31] van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge University Press, Cambridge.
- [32] van der Vaart, A. W. and van Zanten, J. H. (2008). *Reproducing kernel Hilbert spaces of Gaussian priors.*, volume 3, pages 200–222. Inst. Math. Stat. Collect.
- [33] van der Vaart, A. W. and Wellner, J. A. (1996). *Weak Convergence and Empirical Process With Applications to Statistics*. Springer-Verlag New York, Inc.
- [34] Yoo, W. W. and Ghosal, S. (2016). Supremum norm posterior contraction and credible sets for nonparametric multivariate regression. *Ann. Statist.*, 44(3):1069–1102.