

# Cosmology of non-minimal derivative coupling to gravity in Palatini formalism

Burin Gumjudpai<sup>1,\*</sup> and Narakorn Kaewkhao<sup>1,†</sup>

<sup>1</sup>*The Institute for Fundamental Study “The Tah Poe Academia Institute”,  
Naresuan University, Phitsanulok 65000, Thailand*

(Dated: June 12, 2019)

We consider, in Palatini formalism, a modified gravity model of which the scalar field derivative term couples to the Einstein tensor. In this scenario, Ricci scalar, Ricci tensor and Einstein tensor are functions of the connection field. Conformal factor is found as function of scalar field derivative,  $f = 1 - \kappa\phi'^\alpha\phi_{,\alpha}/2$ . In FLRW universe, NMDC coupling constant is limited in a range of  $-2/\dot{\phi}^2 < \kappa \leq \infty$  so that the Lorentz signature of the conformal metric is preserved. The slowly-rolling regime provides  $\kappa < 0$  hence forbidding graviton from taking superluminal speed. Effective gravitational coupling and the entropy of blackhole’s apparent horizon are derived. In case of negative coupling, the acceleration could happen even with  $w_{\text{eff}} > -1/3$ . Power-law potentials of chaotic inflation are considered. The negative NMDC coupling could enhance amount of the inflation. For  $V \propto \phi^2$  and  $V \propto \phi^4$ , it is possible to obtain tensor-to-scalar ratio lower than that of GR case so that it lies in the range,  $r < 0.12$  as constrained by Planck 2015 [34]. The  $V \propto \phi^2$  could be viable since it is possible to yield acceptable range of spectrum index and range of  $r$  values. However the quartic potential’s spectrum index is disfavored by the Planck result.

## I. INTRODUCTION

Astrophysical observations strongly convinces us that the space is in the state of accelerating expansion. Results obtained from supernova type Ia (SNIa) [1–10], large-scale structure surveys [11, 12], cosmic microwave background (CMB) anisotropies [13–16] and X-ray luminosity from galaxy clusters [15, 17, 18] are examples of the evidence of the acceleration. If the expansion is to be accelerated, some unknown form of dark energy [19–21] is suggested as a driving force of the dynamics. Typically dark energy is in form of cosmological constant or scalar field [19–22] such as quintessence [23]–scalar with canonical kinetic term, and classes of k-essence type kinetic energy [24–26] which are hypothesized as dark energy. Alternative of Einstein gravity such as braneworlds,  $f(R)$  could as well result in present acceleration (see e.g. [27, 28]). Other the other situation, inflationary expansion [29–33] in the early universe is also strongly supported by the recent CMB anisotropy observations [34]. Scalar field models or modified gravities should provide explanation to either or both inflationary acceleration and present acceleration.

A gravitational theory with non-minimal coupling (NMC) between scalar field’s derivative term and a gravity sector could as well give accelerating expansion. In metric formalism, of which the metric  $g_{\mu\nu}$  is a dynamical variable, the coupling function  $f(\phi, \phi_{,\mu}, \phi_{,\mu\nu}, \dots)$  can be motivated by requirement of scalar quantum electrodynamics to preserve U(1) symmetry or by models with gravitational constant as a function of the mass density [35]. Non-minimal derivative coupling (NMDC) to  $R$  term can be found in lower energy limits of higher dimensional theories and in Weyl anomaly of  $N = 4$  conformal supergravity [36, 37]. Without loss of generality, other possible coupling terms apart from  $R\phi_{,\mu}\phi'^\mu$  and  $R^{\mu\nu}\phi_{,\mu}\phi_{,\nu}$  are not necessary [38]. Hence gravitational theory with NMDC terms,  $R^{\mu\nu}\phi_{,\mu}\phi_{,\nu}$  and  $R\phi_{,\mu}\phi'^\mu$  with a free canonical kinetic term but without  $V(\phi)$  nor  $\Lambda$  term was studied and found cosmologically interesting, i.e. it gives de Sitter expansion [39]. Moreover types of NMDC models with two separated couplings have been investigated in various contexts and with further modifications [40–42].

Considering a special case of  $\kappa_1 R\phi_{,\mu}\phi'^\mu$  and  $\kappa_2 R^{\mu\nu}\phi_{,\mu}\phi_{,\nu}$  term, one can set  $\kappa \equiv \kappa_2 = -2\kappa_1$ . As a result, the two NMDC terms combine into the Einstein tensor,  $G^{\mu\nu}$  coupling to scalar field’s kinetic part as  $\kappa G_{\mu\nu}\phi'^\mu\phi'^\nu$ . The field equations contain terms with second-order derivative in  $g_{\mu\nu}$  and  $\phi$  at most order hence it is a good dynamical theory as Lagrangian contains only divergence-free tensors [47]. In flat FLRW universe, for  $\kappa > 0$ , there is quasi-de Sitter phase at very early stage and there is initial singularity at very early stage for  $\kappa < 0$ . The expansion is  $a \propto t^{1/3}$  at very late time for any sign of  $\kappa$  [47]. When adding  $V = \text{constant}$  and allowing phantom sign of the free kinetic term, it is possible to transit from de-Sitter phase to other types of expansions [48]. If without free kinetic term, the model gives superluminal sound speed [49] and if having both free and coupling  $\kappa G_{\mu\nu}\phi'^\mu\phi'^\nu$  term with  $V(\phi) = 0$ , the model

\* buring@nu.ac.th: corresponding

† narakornk56@email.nu.ac.th

can not have phantom crossing. Therefore potential is added into the theory and the acceptable action is

$$S = \int d^4x \sqrt{-g} \left[ \frac{R}{8\pi G_N} - (\varepsilon g_{\mu\nu} + \kappa G_{\mu\nu}) \phi^{,\mu} \phi^{,\nu} - 2V(\phi) \right] + S_m. \quad (1)$$

It is found that the potential must be less steep than quadratic potential in acquiring inflation [50]. With a constant potential and a matter term in the model, it is able to describe transition from inflation to matter domination epoch without reheating and this description includes transition to late de-Sitter epoch [51]. For positive potential,  $\kappa > 0$  gives unbound  $\dot{\phi}$  value with restricted Hubble parameter [50]. For  $V = \text{constant}$  and  $\kappa > 0$ , inflationary phase is always possible and the inflation depends solely on the value of coupling constant. Gravitational heavy particles are less produced during inflation when coupling to the inflaton field or to the particles gets stronger [52]. Perturbations and inflationary analysis of the model with a cosmological constant (or equivalently, the constant potential) was shown in [53] confronting observational data. Other very interesting studies of slightly different versions of NMDC model are reported, see such as [54–65] and for recent review we refer to [66].

So far the results given above are obtained in metric formalism. Considering Einstein-Hilbert action with matter term, the metric formalism gives equivalent field equation as that of the Palatini formalism. However if GR is modified, the Palatini approach might not give the same field equations as those in metric formalism as there is a coupling between the Ricci scalar to the field and/or having some form of functions of the Ricci scalar. Unlike the metric formalism, the approach with the Palatini formalism considers the metric tensor and the connection field both as independent dynamical variables. The Palatini connection is hence not the usual Levi-Civita connection of metric  $g_{\mu\nu}$  [67–70] but another independent dynamical field. Recently the Palatini approach to the NMDC gravity was explored considering there are two separated coupling constants, i.e. one of the  $R$  coupling term and the other one of the  $R_{\mu\nu}$  term with a non-zero potential. It was found that phantom crossing with oscillating equation of state parameter is possible [71]. In our work here, we consider similar model but with NMDC Lagrangian term in form of one Einstein tensor coupling to the scalar field's kinetic term. Therefore there is only one combined coupling constant as seen in Eq. (1). We approach the action (1) with the Palatini formalism to derive the model's conformal transformation [72] of which the generalized case was introduced by Bekenstien [73], with conformal and disformal factors. Conformal transformation preserves global shape with a change of local units of length between two points hence shrinking or stretching it in all spacetime directions equally, whereas, disformal transformation changes units of length along the direction of scalar field gradient hence distorting global shape [73]. As to be discussed in Sec. II, the transformation for the action (1) is conformal type. We shall investigate cosmological scenario of the model in Sec. III. whereas it is suggested that the field should be slowly-rolling and the coupling constant should be negative in order to preserve Lorentz invariance and to prevent the graviton from traveling faster than light. Cosmological field equations under slow-roll condition are stated in Sec. IV. Inflationary consideration is explored in Sec. V in which slow-roll parameters and spectral index were derived. We consider power-law potential of chaotic inflation in Sec. VI in which we claim the chaotic inflation model could be viable in our theory. We conclude our work in Sec. VII.

## II. PALATINI NMDC GRAVITY AND CONFORMAL TRANSFORMATION

In the metric formalism, Sushkov's NMDC action is [47]

$$S_g = \int d^4x \sqrt{-g} \left\{ R(g) - \left[ \varepsilon g_{\mu\nu} + \kappa \left( R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} R(g) \right) \right] \phi^{,\mu} \phi^{,\nu} - 2V(\phi) \right\} + S_m[g_{\mu\nu}, \Psi], \quad (2)$$

where the  $\Psi$  represents matter fields and we set the unit  $c = 1$  and  $8\pi G_N = 1$ . Canonical scalar field Lagrangian density is  $\mathcal{L}_\phi = -\varepsilon g_{\mu\nu} \phi^{,\mu} \phi^{,\nu} - 2V(\phi)$  where  $\varepsilon = \pm 1$  is for canonical and the phantom cases respectively. The Einstein tensor is conventional,  $G_{\mu\nu}(g) = R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} R(g)$ , as in Eq. (1). Differently, in Palatini formalism, the NMDC action is expressed as [71]

$$S_{\text{Palatini}} = \int d^4x \sqrt{-g} \left\{ \tilde{R}(\Gamma) - \left[ \varepsilon g_{\mu\nu} + \kappa_1 g_{\mu\nu} \tilde{R}(\Gamma) + \kappa_2 \tilde{R}_{\mu\nu}(\Gamma) \right] \phi^{,\mu} \phi^{,\nu} - 2V(\phi) \right\} + S_m[g_{\mu\nu}, \Psi]. \quad (3)$$

Tilde symbol denotes variables that depend on the connection field. Following Sushkov [47], we set  $\kappa = \kappa_2 = -2\kappa_1$  and define the Einstein tensor in Palatini formalism,

$$\tilde{G}_{\mu\nu}(\Gamma) = \tilde{R}_{\mu\nu}(\Gamma) - \frac{1}{2} g_{\mu\nu} \tilde{R}(\Gamma). \quad (4)$$

Hence Eq. (3) is

$$S_{\text{Palatini}} = \int d^4x \sqrt{-g} \left\{ \tilde{R}(\Gamma) - \left[ \varepsilon g_{\mu\nu} + \kappa \tilde{G}_{\mu\nu}(\Gamma) \right] \phi^{,\mu} \phi^{,\nu} - 2V(\phi) \right\} + S_m[g_{\mu\nu}, \Psi]. \quad (5)$$

The Palatini Ricci tensor is defined by the independent dynamical connection,

$$\tilde{R}_{\mu\nu}(\Gamma) = \tilde{R}^{\lambda}_{\mu\lambda\nu}(\Gamma) = \partial_{\lambda}\Gamma^{\lambda}_{\mu\nu} - \partial_{\nu}\Gamma^{\lambda}_{\mu\lambda} + \Gamma^{\lambda}_{\sigma\lambda}\Gamma^{\sigma}_{\mu\nu} - \Gamma^{\lambda}_{\sigma\nu}\Gamma^{\sigma}_{\mu\lambda}, \quad (6)$$

and the Palatini Ricci scalar is  $\tilde{R} = \tilde{R}(\Gamma) = g^{\mu\nu}\tilde{R}_{\mu\nu}(\Gamma)$ . Varying the Palatini NMDC action in Eq. (5) with respect to the metric, we obtain the first field equation,

$$T_{\mu\nu} = \tilde{G}_{\mu\nu}(\Gamma) + \left[ \frac{\kappa}{2}\tilde{G}_{\mu\nu}(\Gamma)\phi_{,\lambda}\phi^{,\lambda} + \frac{\kappa}{2}\tilde{R}_{\alpha\beta}(\Gamma)g_{\mu\nu}\phi^{,\alpha}\phi^{,\beta} - \kappa\tilde{R}_{\nu\lambda}(\Gamma)\phi_{,\mu}\phi^{,\lambda} + \frac{\kappa}{2}\tilde{R}(\Gamma)\phi_{,\mu}\phi_{,\nu} - 2\kappa\tilde{R}_{\mu\lambda}(\Gamma)\phi_{,\nu}\phi^{,\lambda} + \frac{\varepsilon}{2}g_{\mu\nu}\phi_{,\alpha}\phi^{,\alpha} - \varepsilon\phi_{,\mu}\phi_{,\nu} + g_{\mu\nu}V(\phi) \right], \quad (7)$$

whereas the matter energy-momentum tensor is

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta S_{\text{m}}[g_{\kappa\lambda}, \Psi]}{\delta g^{\mu\nu}}. \quad (8)$$

The second Palatini NMDC field equation comes from the second degree of freedom, the independent connection field  $\Gamma^{\lambda}_{\mu\nu}$  and it is

$$\nabla_{\lambda}^{\Gamma} \left\{ \sqrt{-g} \left[ g^{\mu\nu} \left( 1 - \frac{1}{2}\kappa\phi^{,\alpha}\phi_{,\alpha} \right) \right] \right\} = 0, \quad (9)$$

where  $\nabla_{\lambda}^{\Gamma}$  is the covariant derivative with respect to the independent connection. This is written as

$$\nabla_{\lambda}^{\Gamma} (\sqrt{-g}g^{\mu\nu}f) = 0, \quad (10)$$

where

$$f = 1 - \frac{1}{2}\kappa\phi^{,\alpha}\phi_{,\alpha}. \quad (11)$$

Solving Eq. (10), the new metric or the conformal metric  $h_{\mu\nu}$  is related to the metric  $g_{\mu\nu}$  via a transformation factor  $f$  as

$$h_{\mu\nu} = fg_{\mu\nu} = \left( 1 - \frac{1}{2}\kappa\phi^{,\alpha}\phi_{,\alpha} \right) g_{\mu\nu}, \quad (12)$$

of which  $\sqrt{-h} = \sqrt{-g}f^2$  and its inverse is  $h^{\mu\nu} = f^{-1}g^{\mu\nu}$ . In Palatini formalism, conformal transformation connects dual conformal frames [72, 74, 75]. Further generalization of the conformal transformation is the disformal transformation in form of [73, 76, 79],

$$h_{\mu\nu} \equiv \alpha(\phi, X)g_{\mu\nu} + \beta(\phi, X)\phi_{,\mu}\phi_{,\nu}, \quad (13)$$

where  $\alpha(\phi, X)$  and  $\beta(\phi, X)$  represent conformal and disformal factors respectively. In general, the conformal/disformal factors depend on the field kinetic term,  $X = g^{\mu\nu}\nabla_{\mu}\phi\nabla_{\nu}\phi$  however the new metric in Eq. (12) is not disformally transformed. It obeys only the conformal transformation as we shall see. The conformal part of Eq. (13) can be written as

$$h_{\mu\nu} = \alpha(\phi, X)g_{\mu\nu} = [\alpha_1(\phi) + \alpha_2(\phi)\phi^{,\sigma}\phi_{,\sigma}]g_{\mu\nu}. \quad (14)$$

From Eq. (12), one can see that  $\alpha_1(\phi) = 1$  and  $\alpha_2(\phi) = -\kappa/2$  hence the Palatini approach to NMDC gravity obeys only the conformal transformation. Using these relations above, the action (5) in conformal frame is hence

$$S_{\text{Palatini}} = \int d^4x\sqrt{-h} \left\{ \frac{\tilde{R}(h)}{f^2} - \left[ \frac{\varepsilon h_{\mu\nu}}{f^3} + \kappa \frac{\tilde{G}_{\mu\nu}(h)}{f^2} \right] \phi^{,\mu}\phi^{,\nu} - \frac{2V(\phi)}{f^2} \right\} + S_{\text{m}} \left( \frac{h_{\mu\nu}}{f}, \Psi \right). \quad (15)$$

From here we will consider cosmological scenario of the model.

### III. COSMOLOGICAL SCENARIO

The conformal factor in flat FLRW geometry with homogenous scalar field is

$$f(\dot{\phi}) = 1 - \frac{\kappa}{2} g^{00} \frac{d\phi}{dt} \frac{d\phi}{dt} = 1 + \frac{\kappa}{2} \dot{\phi}^2. \quad (16)$$

Note that there is a factor  $8\pi G_N \equiv 1$  multiplying with  $\kappa$  in this equation. With Eq. (14), the new metric  $h_{\mu\nu}$  preserves Lorentz signature  $(-, +, +, +)$  if  $\alpha_2(\dot{\phi}) = -\kappa/2$  and  $-2/\dot{\phi}^2 < \kappa$  due to positivity of the conformal factor  $f(\dot{\phi})$ . For fast-rolling field, the coupling is allowed in positive region or in very small negative-value region. Hence the coupling estimably ranges from  $0 < \kappa$ . For slowly-rolling field, the coupling is permitted in vast negative region. The FLRW conformal metric is therefore

$$h_{\mu\nu} = \begin{pmatrix} -1 - \frac{\kappa}{2} \dot{\phi}^2 & 0 & 0 & 0 \\ 0 & a^2(1 + \frac{\kappa}{2} \dot{\phi}^2) & 0 & 0 \\ 0 & 0 & a^2(1 + \frac{\kappa}{2} \dot{\phi}^2) & 0 \\ 0 & 0 & 0 & a^2(1 + \frac{\kappa}{2} \dot{\phi}^2) \end{pmatrix}, \quad (17)$$

of which graviton speed is modified with the NMDC-Palatini effect of the conformal factor [73]. Slowly-rolling field allows negative  $\kappa$  hence graviton travels under the speed of light. On the other hand, fast-rolling case could result in faster-than-light graviton. We hence restrict our consideration to the slowly-rolling case. The result above enables us to find that  $\nabla_{\lambda}^{\Gamma}(\sqrt{-h}h^{\mu\nu}) = \nabla_{\lambda}^{\Gamma}(\sqrt{-g}g^{\mu\nu}f)$  and the Levi-Civita connection  $\Gamma_{\mu\nu}^{\lambda}(h)$  is constructed with the conformal metric  $h_{\mu\nu}$  as

$$\Gamma_{\mu\nu}^{\lambda}(h) = \frac{1}{2} h^{\lambda\sigma} (\partial_{\mu} h_{\sigma\nu} + \partial_{\nu} h_{\sigma\mu} - \partial_{\sigma} h_{\mu\nu}). \quad (18)$$

Following e.g. [77, 78], the effective gravitational coupling of Palatini NMDC gravity can be expressed as

$$G_{\text{eff}} = \frac{f^2}{8\pi} = \frac{1}{8\pi} \left(1 + \frac{\kappa}{2} \dot{\phi}^2\right)^2, \quad (19)$$

leading to modification of the entropy of blackhole's apparent horizon for this theory,

$$S_{\text{AH}} = \frac{A}{4(1 + \frac{\kappa}{2} \dot{\phi}^2)^2 / 8\pi}. \quad (20)$$

The effective gravitational coupling strength of the model could be tested by observing temporal variation of the effective gravitational coupling,

$$\frac{\dot{G}_{\text{eff}}}{G_{\text{eff}}} = \frac{2\kappa\dot{\phi}\ddot{\phi}}{(1 + \frac{\kappa}{2} \dot{\phi}^2)}. \quad (21)$$

For fast-rolling field,  $\dot{G}_{\text{eff}}/G_{\text{eff}} \simeq 4\ddot{\phi}/\dot{\phi}$  and on the other hand,  $\dot{G}_{\text{eff}}/G_{\text{eff}} \simeq 2\kappa\dot{\phi}\ddot{\phi}$  for slowly-rolling field. Since  $\Gamma = \Gamma(h, \partial h)$  hence the field equation (7) is expressed as function of  $h$  and  $\partial h$ , for example,  $\tilde{R}_{\mu\nu}(\Gamma)$  is  $\tilde{R}_{\mu\nu}(h, \partial h)$  (written  $\tilde{R}_{\mu\nu}(h)$  for brevity). The energy-momentum tensor obeys the relation  $\tilde{T}_{\mu\nu} = f^{-1}T_{\mu\nu}$  (see Appendix A). Considering time-time component of the field equation,

$$T_{00} = \tilde{G}_{00}(h) - \frac{\kappa}{2} \tilde{G}_{00}(h) \dot{\phi}^2 + \frac{5\kappa}{2} \tilde{R}_{00}(h) \dot{\phi}^2 + \frac{\kappa}{2} \tilde{R}(h) \dot{\phi}^2 - \left(\frac{\varepsilon}{2} \dot{\phi}^2 + V(\phi)\right). \quad (22)$$

The Ricci tensor for the conformal metric  $h_{\mu\nu}$  in  $n$  dimensions is related to the usual Ricci tensor by the following formula (see e.g. [80, 81]),

$$\tilde{R}_{\sigma\nu}(h) = R_{\sigma\nu}(g) - [(n-2)\delta_{\sigma}^{\alpha}\delta_{\nu}^{\beta} + g_{\sigma\nu}g^{\alpha\beta}] \frac{1}{\sqrt{f}} (\nabla_{\alpha}^g \nabla_{\beta}^g \sqrt{f}) + [2(n-2)\delta_{\sigma}^{\alpha}\delta_{\nu}^{\beta} - (n-3)g_{\sigma\nu}g^{\alpha\beta}] \frac{1}{f} (\nabla_{\alpha}^g \sqrt{f})(\nabla_{\beta}^g \sqrt{f}), \quad (23)$$

where  $\nabla_{\lambda}^g$  is the usual covariant derivative constructed from  $g_{\mu\nu}$ . Ricci tensor and Ricci scalar for flat FLRW universe are derived,

$$R_{00}(g) = -3(\dot{H} + H^2), \quad R_{ii}(g) = a^2(\dot{H} + 3H^2), \quad R(g) = 6(\dot{H} + 2H^2) \quad (24)$$

and

$$\tilde{R}_{00}(h) = -3(\dot{H} + H^2) - \frac{3}{2} \left( \frac{\ddot{f}}{f} - \frac{2\dot{f}^2}{f^2} \right), \quad \tilde{R}_{ii}(h) = R_{ii}(g) + \frac{a^2 \ddot{f}}{2f}. \quad (25)$$

First and second-order time derivative of the conformal factor are expressed in term of  $\dot{\phi}$ ,  $\ddot{\phi}$  and  $\ddot{\phi}$ , i.e.

$$\dot{f} = \kappa \dot{\phi} \ddot{\phi}, \quad \ddot{f} = \kappa (\ddot{\phi}^2 + \dot{\phi} \ddot{\phi}). \quad (26)$$

The Ricci scalar under the conformal transformation is [81]

$$\tilde{R}(h) = f^{-1}R(g) - 2(n-1)g^{\alpha\beta}f^{-3/2} \left( \nabla_\alpha \nabla_\beta \sqrt{f} \right) - (n-1)(n-4)g^{\alpha\beta}f^{-2} \left( \nabla_\alpha \sqrt{f} \right) \left( \nabla_\beta \sqrt{f} \right). \quad (27)$$

In four dimensions, this is

$$\tilde{R}(h) = \frac{1}{f}R(g) + 3 \left( \frac{\ddot{f}}{f^2} - \frac{\dot{f}^2}{2f^3} \right) = \frac{6}{f} (\dot{H} + 2H^2) + 3 \left( \frac{\ddot{f}}{f^2} - \frac{\dot{f}^2}{2f^3} \right). \quad (28)$$

Using  $\tilde{T}_{\mu\nu} = f^{-1}T_{\mu\nu}$ , the  $T_{00}$  component of NMDC-Palatini field equation is the matter density,

$$\rho_m = \dot{H} \left[ 12f + \frac{6}{f} - 18 \right] + H^2 \left[ 12f + \frac{12}{f} - 21 \right] - \frac{3}{2}(1-f) \left( \frac{4\ddot{f}}{f} - \frac{8\dot{f}^2}{f^2} \right) - \frac{3\ddot{f}}{2f} + \frac{3\dot{f}}{f^2} + \frac{3\dot{f}^2}{f^2} - \frac{3\dot{f}^2}{2f^3} - \rho_\phi, \quad (29)$$

where  $\rho_{\text{tot}} \equiv \rho_m + \rho_\phi$  and  $\rho_\phi = \varepsilon \dot{\phi}^2/2 + V(\phi)$ . The matter pressure found from the  $T_{ii}$  components is

$$p_m = \dot{H} (4f - 6) + H^2 (6f - 9) - \frac{3}{2}(1-f) \left( \frac{\ddot{f}}{f} - \frac{2\dot{f}^2}{f^2} \right) + \frac{\ddot{f}}{f} - \frac{3\dot{f}}{2f} + \frac{3\dot{f}^2}{4f^2} - p_\phi, \quad (30)$$

where  $p_{\text{tot}} \equiv p_m + p_\phi$ . The pressure of scalar field is  $p_\phi = \varepsilon \dot{\phi}^2/2 - V(\phi)$ . For brevity, we define

$$A \equiv 4f - 6, \quad (31)$$

$$B \equiv 6f - 9, \quad (32)$$

$$C \equiv -\frac{3}{2}(1-f) \left( \frac{\ddot{f}}{f} - \frac{2\dot{f}^2}{f^2} \right) + \frac{\ddot{f}}{f} - \frac{3\dot{f}}{2f} + \frac{3\dot{f}^2}{4f^2}, \quad (33)$$

$$D \equiv 12f + \frac{6}{f} - 18, \quad (34)$$

$$E \equiv 12f + \frac{12}{f} - 21, \quad (35)$$

$$F \equiv -\frac{3}{2}(1-f) \left( \frac{4\ddot{f}}{f} - \frac{8\dot{f}^2}{f^2} \right) - \frac{3\ddot{f}}{2f} + \frac{3\dot{f}}{f^2} + \frac{3\dot{f}^2}{f^2} - \frac{3\dot{f}^2}{2f^3}. \quad (36)$$

The effective equation of state parameter is hence

$$w_{\text{eff}} \equiv \frac{p_{\text{tot}}}{\rho_{\text{tot}}} = \frac{A\dot{H} + BH^2 + C}{D\dot{H} + EH^2 + F}. \quad (37)$$

The modified  $\dot{H}$  and modified Friedmann equations are found as

$$\dot{H} = \frac{[(B - Ew_{\text{eff}})H^2 - Fw_{\text{eff}} + C]}{Dw_{\text{eff}} - A}, \quad H^2 = \frac{\rho_{\text{tot}}}{3} \frac{\left[ 1 - \frac{(C - Fw_{\text{eff}})D}{(Dw_{\text{eff}} - A)\rho_{\text{tot}}} - \frac{F}{\rho_{\text{tot}}} \right]}{\left[ \frac{(B - Ew_{\text{eff}})D}{3(Dw_{\text{eff}} - A)} + \frac{E}{3} \right]}. \quad (38)$$

The acceleration equation can be found from the  $\ddot{a}/a = \dot{H} + H^2$ . In the GR limit,  $f = 1$  making  $A = -2$ ,  $B = -3$ ,  $E = 3$  and  $C = D = F = 0$  hence  $p_{\text{tot}}$  and  $w_{\text{eff}}$  reduce to the usual  $p_{\text{tot}} = -2\dot{H} - 3H^2$  and  $w_{\text{eff}} = -1 - 2\dot{H}/3H^2$ . The Friedmann and acceleration equations can as well reduced to standard GR case,  $H^2 = \rho_{\text{tot}}/3$ , and  $\ddot{a}/a =$

$-(1/6)(\rho_{\text{tot}} + 3p_{\text{tot}})$ . Modified Klein-Gordon equation can be obtained from the Euler-Lagrange equation for scalar field (see e.g. [81] for standard result),

$$\ddot{\phi} \left[ -\varepsilon + \frac{\kappa}{2} \left( \tilde{R}(h) - \tilde{R}_{00}(h) \right) \right] - \kappa \dot{\phi} \nabla_0^h \tilde{R}_{00}(h) + \frac{\kappa}{2} \dot{\phi} \nabla_0^h \tilde{R}(h) - 3\varepsilon H \dot{\phi} - V' = 0, \quad (39)$$

where  $V' = dV(\phi)/d\phi$  and  $\nabla_\mu^h \phi = \nabla_\mu^g \phi = \partial_\mu \phi$ , and

$$\nabla_\mu^h \nabla_\nu^h \phi = \nabla_\mu^g \nabla_\nu^g \phi - (\delta_\mu^\alpha \delta_\nu^\beta + \delta_\mu^\beta \delta_\nu^\alpha - g_{\mu\nu} g^{\alpha\beta}) \frac{1}{\sqrt{f}} \left( \nabla_\alpha^g \sqrt{f} \right) \left( \nabla_\beta^g \sqrt{f} \right). \quad (40)$$

The time component of the equation (40) reads,  $\nabla_0^h \nabla_0^h \phi = \ddot{\phi} = \ddot{\phi} - \dot{f}^2/(4f^{3/2})$ . The modified Klein-Gordon equation hence reads

$$\begin{aligned} & \ddot{\phi} \left\{ -\varepsilon + \frac{\kappa}{2} \left[ \left( \frac{6\dot{H} + 12H^2}{f} + 3 \left( \frac{\ddot{f}}{f^2} - \frac{\dot{f}^2}{2f^3} \right) \right) - \left( -3(\dot{H} + H^2) - \frac{3}{2} \left( \frac{\ddot{f}}{f} - \frac{2\dot{f}^2}{f^2} \right) \right) \right] \right\} \\ & + \frac{\kappa}{2} \dot{\phi} \nabla_0^g \left[ \frac{6\dot{H} + 12H^2}{f} + 3 \left( \frac{\ddot{f}}{f^2} - \frac{\dot{f}^2}{2f^3} \right) \right] - 3\varepsilon H \dot{\phi} - V' = 0. \end{aligned} \quad (41)$$

which recovers the usual Klein-Gordon equation,  $\varepsilon \ddot{\phi} + 3\varepsilon H \dot{\phi} + V' = 0$  in the GR limit.

#### IV. SLOW-ROLL REGIME

Slowing-rolling field obeys the condition  $0 < |\dot{\phi}| \ll 1$  and we approximate further that  $|\ddot{\phi}| \ll |\dot{\phi}| \ll |\phi|$ , i.e.  $0 \sim |\dot{f}| \ll |\ddot{f}| \ll |f|$ . This enables us to neglect  $\ddot{\phi}$ ,  $\dot{\phi}^4 \ddot{\phi}^2$ ,  $\dot{\phi} \ddot{\phi}$  terms and to do binomial approximation,  $(1 + \kappa \dot{\phi}^2/2)^{-1} \simeq 1 - \kappa \dot{\phi}^2/2$  hence  $1/f \simeq 2 - f$  in the calculation. As a result,  $A \simeq -2 + 2\kappa \dot{\phi}^2$ ,  $B \simeq -3 + 3\kappa \dot{\phi}^2$ ,  $D \simeq 3\kappa \dot{\phi}^2$ ,  $E \simeq 3$ ,  $C \simeq F \simeq 0$  and  $A \simeq 2B/3$ . We found that

$$\frac{\ddot{a}}{a} \simeq -\frac{1}{6} \rho_{\text{tot}} \left[ 1 + \frac{7}{2} \kappa \dot{\phi}^2 + 3w_{\text{eff}} \left( 1 + \frac{3}{2} \kappa \dot{\phi}^2 \right) \right], \quad (42)$$

giving the acceleration condition,

$$w_{\text{eff}} \lesssim -\frac{1}{3} \left( 1 + 2\kappa \dot{\phi}^2 \right). \quad (43)$$

As discuss earlier, the slowly-rolling field allows negative  $\kappa$  hence the expansion is under acceleration at  $w_{\text{eff}}$  slightly greater than  $-1/3$ . The modified Friedmann equation (38) is approximated as

$$H^2 \simeq \frac{1}{3} \rho_{\text{tot}} \left[ \frac{3(Dw_{\text{eff}} - A)}{BD - EA} \right] \simeq \frac{\rho_{\text{tot}}}{3} \left[ 1 + \frac{3}{2} \kappa \dot{\phi}^2 (1 + w_{\text{eff}}) \right], \quad (44)$$

and the modified Klein-Gordon equation (41) in the slow-roll regime is

$$\ddot{\phi} \left\{ \varepsilon - \frac{9\kappa}{2} \dot{H} (1 - \kappa \dot{\phi}^2) - \frac{3\kappa}{2} H^2 (5 - 6\kappa \dot{\phi}^2) \right\} + 3H \dot{\phi} \left[ \varepsilon - \left( \frac{\ddot{H}}{H} + 4\dot{H} \right) \kappa \left( 1 - \frac{\kappa \dot{\phi}^2}{2} \right) \right] + V' \simeq 0. \quad (45)$$

#### V. SLOW-ROLL INFLATION

Considering the early universe when the scalar field dynamically drives the inflation. Scalar field density dominates the universe and slow-roll regime is plausible during the era. The Friedmann equation (44) reads

$$H^2 \simeq \frac{\rho_\phi}{3M_{\text{P}}^2} \left[ 1 + \frac{3}{2} \kappa \dot{\phi}^2 \left( 1 + \frac{p_\phi}{\rho_\phi} \right) \right] \simeq \frac{1}{3} \frac{V(\phi)}{M_{\text{P}}^2}, \quad (46)$$

as the condition  $\dot{\phi}^2 \ll V(\phi)$  is assumed. Here we restore  $8\pi G_N = M_{\text{P}}^{-2}$  to the equation. The coupling constant can be considered in  $mass^{-2}$  dimension,  $\kappa = M^{-2} \lesssim M_{\text{P}}^{-2}$ . The other useful relation is

$$\dot{H} \simeq \frac{V'\dot{\phi}}{6HM_{\text{P}}^2} \simeq \frac{\sqrt{3}V'\dot{\phi}}{6\sqrt{V}M_{\text{P}}} \quad (47)$$

as the slow-roll approximated Friedmann equation is used. We can derive  $\dot{\phi}$  from Eq. (45) assuming slow-roll regime,  $\ddot{\phi} \simeq 0$  and  $|\ddot{H}| \ll |H\dot{H}| \ll |H^3|$ . This gives

$$\dot{\phi} \simeq -\frac{V'(\phi)}{3H(\varepsilon - 4\kappa\dot{H})}. \quad (48)$$

Using this in (47) hence,

$$\dot{H} \simeq -\frac{(V'(\phi))^2}{18H^2M_{\text{P}}^2(\varepsilon - 4\kappa\dot{H})}. \quad (49)$$

Deriving slow-roll parameters, the first one is

$$\epsilon_v \equiv -\frac{\dot{H}}{H^2} \simeq \frac{M_{\text{P}}^2}{2(\varepsilon - 4\kappa\dot{H})} \left(\frac{V'}{V}\right)^2, \quad (50)$$

with a slow-roll condition,  $\epsilon_v \ll 1$ . The second parameter

$$\delta \equiv \frac{\ddot{\phi}}{H\dot{\phi}} \simeq -\frac{V''(\phi)}{3H^2(\varepsilon - 4\kappa\dot{H})} + \frac{V'(\phi)\dot{H}}{3H^3(\varepsilon - 4\kappa\dot{H})\dot{\phi}} - \frac{4\kappa\ddot{H}V'(\phi)}{3M_{\text{P}}^2H^2(\varepsilon - 4\kappa\dot{H})^2\dot{\phi}}, \quad (51)$$

with a slow-roll condition,  $|\delta| \ll 1$ . The first term on the right-hand side of Eq.(51) is, in fact, the other slow-roll parameter,

$$\eta_v \equiv \frac{M_{\text{P}}^2}{(\varepsilon - 4\kappa\dot{H})} \frac{V''(\phi)}{V(\phi)} \simeq \frac{V''(\phi)}{3H^2(\varepsilon - 4\kappa\dot{H})} \quad (52)$$

with a slow-roll condition,  $|\eta_v| \ll 1$ . The second term on the right-hand side of Eq. (51) is just  $\epsilon_v$  while the third term, with help of Eq. (48), is a new slow-roll parameter,  $\eta_\kappa$ , manifesting another NMDC-Palatini effect,

$$\eta_\kappa \equiv \frac{4\kappa\ddot{H}}{HM_{\text{P}}^2(\varepsilon - 4\kappa\dot{H})} \simeq -\frac{4\kappa\ddot{H}V'(\phi)}{3H^2M_{\text{P}}^2(\varepsilon - 4\kappa\dot{H})^2\dot{\phi}} \simeq \frac{4\kappa}{M_{\text{P}}^2(\varepsilon - 4\kappa\dot{H})^3} \left[ \frac{V''(V')^2}{18V} - \frac{V'^4}{36V^2} \right]. \quad (53)$$

Therefore  $\delta = -\eta_v + \epsilon_v + \eta_\kappa$ . The spectral index of the model can be derived from  $n_s - 1 = -4\epsilon_v - 2\delta$  to obtain [83],

$$n_s - 1 = -6\epsilon_v + 2\eta_v - 2\eta_\kappa \quad (54)$$

or written in full form of  $V(\phi)$  and its field derivatives,

$$n_s - 1 = -\frac{3M_{\text{P}}^2}{(\varepsilon - 4\kappa\dot{H})} \left(\frac{V'}{V}\right)^2 + \frac{2M_{\text{P}}^2}{(\varepsilon - 4\kappa\dot{H})} \frac{V''}{V} - \frac{8\kappa}{M_{\text{P}}^2(\varepsilon - 4\kappa\dot{H})^3} \left[ \frac{V''(V')^2}{18V} - \frac{(V')^4}{36V^2} \right]. \quad (55)$$

The  $e$ -folding number during the inflationary epoch can be found from  $\mathcal{N} = \ln(a_f/a_i) = \int_{t_i}^{t_f} H dt$ , where  $i$  and  $f$  denotes the beginning and the end of inflationary phase. From Eqs. (46) and (48),  $H dt = [-V(\varepsilon - 4\kappa\dot{H})/(V'M_{\text{P}}^2)]d\phi$ . During inflation,  $\dot{H}$  is almost constant due to slowly-rolling field as seen in Eq. (47), the number of  $e$ -folds is hence approximately

$$\mathcal{N} \simeq \frac{(\varepsilon - 4\kappa\dot{H})}{M_{\text{P}}^2} \int_{\phi_f}^{\phi_i} \frac{V(\phi)}{V'(\phi)} d\phi = (\varepsilon - 4\kappa\dot{H})^{1/2} \int_{\phi_f}^{\phi_i} \frac{1}{\sqrt{2\epsilon_{v,\text{GR}}}} \frac{d\phi}{M_{\text{P}}}, \quad (56)$$

where here  $\epsilon_{v,\text{GR}} \equiv (M_{\text{P}}^2/2)(V'/V)^2$ . A known form of the scalar field potential is necessary in order to evaluate  $n_s$  and  $\epsilon_v$ . From now on, we consider only the non-phantom field, i.e.  $\varepsilon = 1$ . If the NMDC-Palatini effect,  $4\kappa\dot{H} > 0$ , it reduces the amount of inflation from that of the GR case. On the other hand, for  $4\kappa\dot{H} < 0$ , the effect is opposite, that is to enhance the amount of inflation. From acceleration condition,  $\dot{H} = \ddot{a}/a - H^2 \gg -H^2$ , the inflationary acceleration makes  $\dot{H} > 0$  for most of the time. Hence,  $\kappa > 0$  case reduces the amount of inflation while the  $\kappa < 0$  case enhances the amount of inflation.

## VI. RESCUING THE CHAOTIC INFLATION

Consider single monomial potential (chaotic inflation [84]) in form of

$$V(\phi) = V_0 \phi^n, \quad (57)$$

where  $V_0 \equiv \lambda(M_{\text{P}}^4/M_{\text{P}}^n)$ . With  $V'/V = n/\phi$  and  $V''/V = n(n-1)/\phi^2$ , the slow-roll parameters and the spectral index can be found,

$$\epsilon_v = \frac{n^2}{2(1-4\kappa\dot{H})} \frac{M_{\text{P}}^2}{\phi^2}, \quad \eta_v = \frac{n(n-1)}{(1-4\kappa\dot{H})} \frac{M_{\text{P}}^2}{\phi^2}, \quad \eta_\kappa = \frac{n^3(n-2)}{9(1-4\kappa\dot{H})^3} \kappa V_0^2 \frac{\phi^{2n-4}}{M_{\text{P}}^2} \quad (58)$$

$$n_s - 1 = -\frac{M_{\text{P}}^2 [n(n+2)]}{(1-4\kappa\dot{H})} \phi^{-2} - \frac{2\kappa V_0^2 [n^3(n-2)]}{9M_{\text{P}}^2(1-4\kappa\dot{H})^3} \phi^{2n-4}. \quad (59)$$

In GR, scalar field rolling in power-law potential (with  $n > \sqrt{2}$ ) is super-Planckian in order to satisfy the slow-roll condition. Considering  $\epsilon_v < 1$ , the GR slow-roll condition is modified with the NMDC-Palatini effect,  $[|n|/(\sqrt{2}\sqrt{1-4\kappa\dot{H}})]M_{\text{P}} < \phi$ . For  $\kappa < 0$  case, the slowly-rolling scalar field can avoid the super-Planckian regime if the coupling takes the value in a range,  $\kappa < -(n^2-2)/(8\dot{H})$ . That is for  $V \propto \phi^{-2}$  or  $V \propto \phi^2$ , it is  $\kappa < -1/(4\dot{H})$  and for  $V \propto \phi^4$ , it is  $\kappa < -7/(4\dot{H})$ . Oppositely if  $\kappa > 0$ , the field takes more super-Planckian value in order to slowly roll. For simplification, we let  $\phi_{\text{f}} = 0$  and the integral (56) is  $\mathcal{N} \simeq (1-4\kappa\dot{H})\phi_{\text{f}}^2/(2nM_{\text{P}}^2)$  so that we define

$$\phi^2 \equiv \phi_{\text{f}}^2(n, \mathcal{N}, \dot{H}) \simeq \frac{2nM_{\text{P}}^2}{(1-4\kappa\dot{H})} \mathcal{N}. \quad (60)$$

The  $e$ -folding number of the GR case is  $\mathcal{N}_{\text{GR}} = \phi^2/(2nM_{\text{P}}^2)$ , therefore  $\mathcal{N} = \mathcal{N}_{\text{GR}}(1-4\kappa\dot{H})$  and  $\mathcal{N} > \mathcal{N}_{\text{GR}}$  for  $\kappa < 0$ . The slow-roll parameters can be expressed in term of  $\mathcal{N}_{\text{GR}}$  as

$$\epsilon_v = \frac{n}{4\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right), \quad \eta_v = \frac{n-1}{2\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right), \quad \eta_\kappa = \frac{\kappa V_0^2 2^{2n-2} M_{\text{P}}^{2n-6}}{9(1-4\kappa\dot{H})^3} (n-2)n^{n+1} \mathcal{N}_{\text{GR}}^{n-2}, \quad (61)$$

where the  $\epsilon_{v,\text{GR}} \equiv n/(4\mathcal{N}_{\text{GR}})$  and  $\eta_{v,\text{GR}} \equiv (n-1)/(2\mathcal{N}_{\text{GR}})$ . The spectral index is hence

$$n_s \simeq 1 - \frac{n+2}{2\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right) - 2^{n-1} \frac{\kappa V_0^2 M_{\text{P}}^{2n-6}}{9(1-4\kappa\dot{H})^3} (n-2)n^{n+1} \mathcal{N}_{\text{GR}}^{n-2}. \quad (62)$$

For the  $V = V_0\phi^2$  case, we have

$$\epsilon_v = \eta_v = \frac{1}{2\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right), \quad \eta_\kappa = 0 \quad \text{and} \quad n_s = 1 - \frac{2}{\mathcal{N}_{\text{GR}}(1-4\kappa\dot{H})}, \quad (63)$$

and for the  $V = V_0\phi^4$  case, they are

$$\epsilon_v = \frac{1}{\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right), \quad \eta_v = \frac{3}{2\mathcal{N}_{\text{GR}}} \left( \frac{1}{1-4\kappa\dot{H}} \right), \quad \eta_\kappa = \frac{8192}{9} \frac{\kappa V_0^2 M_{\text{P}}^2 \mathcal{N}_{\text{GR}}^2}{(1-4\kappa\dot{H})^3} \quad (64)$$

and

$$n_s = 1 - \frac{3}{\mathcal{N}_{\text{GR}}(1-4\kappa\dot{H})} - \frac{16384}{9} \frac{\kappa V_0^2 M_{\text{P}}^2}{(1-4\kappa\dot{H})^3} \mathcal{N}_{\text{GR}}^2. \quad (65)$$

The GR predictions for the  $V \propto \phi^n$  models, given  $\mathcal{N}_{\text{GR}} = 60$  and  $n = 2$ , are the tensor-to-scalar ratio  $r \simeq 16\epsilon_{v,\text{GR}} \simeq 0.13$  and  $n_s \simeq 0.967$ . The  $n = 4$  case has  $r \simeq 0.27$  and  $n_s \simeq 0.95$ . These are disfavored by Planck 2015's results [34] which are  $r < 0.12$  at 95% CL ( $B$ -mode polarization constraint from the BICEP2/Keck Array/Planck joint analysis) and  $n_s = 0.968 \pm 0.006$  (temperature and large angular scale polarization data). Eq. (47) for the power-law potential case is

$$\dot{H} \simeq \frac{\sqrt{3}\sqrt{V_0}}{6} n \phi^{(n-2)/2} \frac{\dot{\phi}}{M_{\text{P}}}. \quad (66)$$



For  $V = V_0\phi^2$ , it is found from Eq. (63) that the range,  $\kappa\dot{H} \lesssim -0.027$  can satisfy the upper bound of  $r < 0.12$ . The Planck 2015 bound of  $n_s = 0.968 \pm 0.006$  corresponds to the range  $-0.071 \lesssim \kappa\dot{H} \lesssim 0.031$ , hence the combined bound is

$$-0.071 \lesssim \kappa\dot{H} \lesssim -0.027. \quad (67)$$

Using  $\dot{H} \simeq \sqrt{V_0/3}(\dot{\phi}/M_{\text{P}})$  (the  $n = 2$  case of Eq. (66)), the combined condition (67) is satisfied when

$$-\frac{0.122}{\dot{\phi}(m/M_{\text{P}})} \lesssim \kappa \lesssim -\frac{0.068}{\dot{\phi}(m/M_{\text{P}})}, \quad (68)$$

where  $V_0 \equiv (1/2)m^2 = \lambda M_{\text{P}}^2$  and hence  $\lambda = (1/2)(m/M_{\text{P}})^2$ . The small value of  $\dot{\phi}$  and the mass  $m > M_{\text{P}}$  counterbalance the value of  $\kappa$ . For  $V = V_0\phi^4$ , using Eq. (64), the range  $\kappa\dot{H} \lesssim -0.306$  satisfies the bound of  $r < 0.12$ . Using  $\dot{H} \simeq 2\sqrt{V_0/3}(\dot{\phi}\dot{\phi}/M_{\text{P}})$  (the  $n = 4$  case of Eq. (66)), the condition  $r < 0.12$  is satisfied as  $\kappa \lesssim -0.264/[\sqrt{\lambda}\dot{\phi}(\phi/M_{\text{P}})]$  where in this case  $V_0 = \lambda$ . As seen in Eq. (65),  $n_s$  of the  $V = V_0\phi^4$  case needs to be much fine-tuned due to the NMDC-Palatini effect of the  $\eta_\kappa$  term, therefore the quartic potential is unlikely as a viable model.

## VII. CONCLUSIONS

We investigate cosmology of a non-minimal derivative coupling (NMDC) to gravity model in Palatini formalism imposing non-minimal constant coupling between the Einstein tensor and the scalar field derivative term. The Lagrangian contains also a free scalar field derivative term and a scalar potential as proposed in [47]. In Palatini formalism, the connection field is a dynamical variable hence Ricci scalar and Ricci tensor are also functions of connection field. As a result, the Einstein tensor is a function of the connection,  $\tilde{G}_{\mu\nu}(\Gamma)$ . Variation of the NMDC action with respect to the independent connection gives conformal transformation factor,  $f = 1 - \kappa\phi^\alpha\phi_{,\alpha}/2$ . In FLRW spacetime the conformal factor takes the form,  $f(\dot{\phi}) = 1 + \kappa\dot{\phi}^2/2$ . The NMDC coupling constant is enforced to be in a range of  $-2/\dot{\phi}^2 < \kappa \leq \infty$  in order to preserve the Lorentz signature of the conformal metric. The coupling needs to be negative in order to prevent graviton traveling with superluminal speed. In conformal frame, the effective gravitational coupling is  $G_{\text{eff}} = (8\pi)^{-1}(1 + \kappa\dot{\phi}^2/2)^2$  (in the unit of  $8\pi G_{\text{N}} = c = 1$ ) which reduces to standard GR case when there is no NMDC coupling. The NMDC-Palatini effective gravitational constant leads to modification of the entropy of blackhole's apparent horizon to  $S_{\text{AH}} = A/[4(1 + \frac{\kappa}{2}\dot{\phi}^2)^2/8\pi]$ . The cosmological field equations found can reduce to standard form in the GR limit. Field equations are approximated in the slow-roll regime. We see that the acceleration condition is modified to  $w_{\text{eff}} \lesssim -(1/3)(1 + 2\kappa\dot{\phi}^2)$ . The NMDC-Palatini effect of the  $2\kappa\dot{\phi}^2$  term, with  $\kappa < 0$ , results in acceleration at  $w_\phi$  value even more than  $-1/3$ . Considering power-law potentials  $V \propto \phi^2$  and  $V \propto \phi^4$  in chaotic inflation, the NMDC-Palatini effect of the  $\kappa < 0$  case can enhance the amount of inflation. The NMDC-Palatini effect results in an extra term  $\eta_\kappa$  in the slow-roll parameter  $\delta = -\eta_v + \epsilon_v + \eta_\kappa$  so that it affects the spectrum index. In case of  $V \propto \phi^2$ , the  $\eta_\kappa$  term vanishes in the expression of  $n_s$ , however in the case of  $V \propto \phi^4$ , it contributes to enormous value of  $n_s$ . Therefore, in this model, the quartic potential is not likely to be viable compared to the Planck 2015's predicted range of spectrum index [34]. Moreover in the  $V \propto \phi^2$  case, the  $\kappa < 0$  coupling could help resolving the Planck 2015's tensor-to-scalar ratio constraint, i.e. it is possible to obtain  $r < 0.12$ . In the GR case, for  $\mathcal{N}_{\text{GR}} = 60$ , the  $V \propto \phi^2$  potential gives  $r \simeq 0.13$  and  $n_s \simeq 0.967$  which are disfavored by Planck 2015. However in the NMDC-Palatini model, the range  $-0.071 \lesssim \kappa\dot{H} \lesssim -0.027$  can satisfy Planck 2015 tensor-to-scalar ratio upper bound ( $r < 0.12$ ) and the constraint  $n_s = 0.968 \pm 0.006$ . The viable range corresponds to  $-0.122 \lesssim \kappa\dot{\phi}(m/M_{\text{P}}) \lesssim -0.068$  of which the small  $\dot{\phi}$  and  $m > M_{\text{P}}$  counterbalance the value of  $\kappa$  so that the chaotic inflation  $V \propto \phi^2$  could be viable in the range.

## ACKNOWLEDGEMENTS

This work is supported by the National Research Council of Thailand (R2558B132). The authors thank Antonio De Felice, Phongsaphat Rangdee and Pichet Vanichchapongjaroen for discussions.

**Appendix A: Proof of relation between  $\tilde{T}_{\mu\nu}$  and  $T_{\mu\nu}$**

$$\begin{aligned}\tilde{T}_{\mu\nu} &= -\frac{2}{\sqrt{-h}} \frac{\delta\mathcal{L}_m(g_{\kappa\lambda}, \Psi)}{\delta h^{\mu\nu}} = -\frac{2}{f^2\sqrt{-g}} \frac{\delta\mathcal{L}_m(g_{\kappa\lambda}, \Psi)}{\delta(f^{-1}g^{\mu\nu})} \\ &= -\frac{2}{f\sqrt{-g}} \frac{\delta\mathcal{L}_m(g_{\kappa\lambda}, \Psi)}{\delta g^{\mu\nu}} = f^{-1}T_{\mu\nu}.\end{aligned}\tag{A1}$$

- [1] Amanullah, R., *et al.*: *Astrophys. J.* **716**, (2010) 712.
- [2] Astier, P., *et al.* (SNLS Collaboration): *Astron. Astrophys.* **447**, (2006) 31.
- [3] Goldhaber, G., *et al.* (The Supernova Cosmology Project Collaboration): *Astrophys. J.* **558**, (2001) 359.
- [4] Perlmutter, S., *et al.* (Supernova Cosmology Project Collaboration): *Nature* **391**, (1998) 51.
- [5] Perlmutter, S., *et al.* (Supernova Cosmology Project Collaboration): *Astrophys. J.* **517**, (1999) 565.
- [6] Riess, A. G., *et al.* (Supernova Search Team Collaboration): *Astron. J.* **116**, (1998) 1009.
- [7] Riess, A. G.: arXiv:astro-ph/9908237.
- [8] Riess, A. G., *et al.* (Supernova Search Team Collaboration): *Astrophys. J.* **607**, (2004) 665.
- [9] Riess, A. G., *et al.*: *Astrophys. J.* **659**, (2007) 98.
- [10] Tonry, J. L., *et al.* (Supernova Search Team Collaboration): *Astrophys. J.* **594**, (2003) 1.
- [11] Scranton, R., *et al.* (SDSS Collaboration): arXiv:astro-ph/0307335.
- [12] Tegmark, K., *et al.* (SDSS Collaboration): *Phys. Rev. D* **69**, (2004) 103501.
- [13] Larson, D., *et al.*: *Astrophys. J. Suppl.* **192**, (2011) 16.
- [14] Komatsu, E., *et al.* (WMAP Collaboration): *Astrophys. J. Suppl.* **192**, (2011) 18.
- [15] Hu, J. W., Cai, R. G., Guo, Z. K., Hu, B.: *JCAP* **1405**, (2014) 020.
- [16] Masi, S., *et al.*: *Prog. Part. Nucl. Phys.* **48**, (2002) 243.
- [17] Allen, S. W., *et al.*: *Mon. Not. Roy. Astron. Soc.* **353**, (2004) 457.
- [18] Rapetti, D., Allen, S. W., Weller, J.: *Mon. Not. Roy. Astron. Soc.* **360**, (2005) 555.
- [19] Copeland, E. J., Sami, M., Tsujikawa, S.: *Int. J. Mod. Phys. D* **15**, (2006) 1753.
- [20] Padmanabhan, T.: *Curr. Sci.* **88**, (2005) 1057.
- [21] Padmanabhan, T.: *AIP Conf. Proc.* **861**, (2006) 179.
- [22] Amendola, L., Tsujikawa, S.: *Dark Energy: Theory and Observations* (Cambridge University Press, 2010).
- [23] Caldwell, R. R., Dave, R., Steinhardt, P. J.: *Phys. Rev. Lett.* **80**, (1998) 1582.
- [24] Chiba, T., Okabe, T., Yamaguchi, M.: *Phys. Rev. D* **62**, (2000) 023511.
- [25] Armendariz-Picon, C., Mukhanov, V. F., Steinhardt, P. J.: *Phys. Rev. Lett.* **85**, (2000) 4438.
- [26] Armendariz-Picon, C., Mukhanov, V. F., Steinhardt, P. J.: *Phys. Rev. D* **63**, (2001) 103510.
- [27] De Felice, A., Tsujikawa, S.: *Liv. Rev. Rel.* **13**, (2010) 3.
- [28] Carroll, S. M., Duvvuri, V., Trodden, M., Turner, M. S.: *Phys. Rev. D* **70**, (2004) 043528.
- [29] Guth, A. H.: *Phys. Rev. D* **23** (1981) 347.
- [30] Sato, K: *Mon. Not. R. Astron. Soc.* **195** (1981) 467; *Phys. Lett. B* **99** (1981) 66.
- [31] Starobinsky, A. A.: *Phys. Lett. B* **91** (1980) 99.
- [32] Linde, A.: *Phys. Lett. B* **108** (1982) 389.
- [33] Albrecht, A. and Steinhardt, P.: *Phys. Rev. Lett.* **48** (1982) 1220.
- [34] Ade, P. A. R., *et al.* [Planck Collaboration], *Astron. Astrophys.* **594** (2016) A20.
- [35] Amendola, L.: *Phys. Lett. B* **301**, (1993) 175.
- [36] Liu, H. and Tseytlin, A. A.: *Nucl. Phys. B* **533**, (1998) 88.
- [37] Nojiri, S. and Odintsov, S. D.: *Phys. Lett. B* **444**, (1998) 92.
- [38] Capozziello, S., Lambiase, G., Schmidt, H. J.: *Annalen Phys.* **9**, (2000) 39.
- [39] Capozziello, S., Lambiase, G.: *Gen. Rel. Grav.* **31**, (1999) 1005.
- [40] Granda, L. N., Cardona, W.: *JCAP* **1007**, (2010) 021.
- [41] Granda, L. N.: *JCAP* **1104**, (2011) 016.
- [42] Granda, L. N.: *Class. Quant. Grav.* **28**, (2011) 025006.
- [43] Granda, L. N.: *Mod. Phys. Lett. A* **27**, (2012) 1250018.
- [44] Granda, L. N., Torrente-Lujan, E. and Fernandez-Melgarejo, J. J.: *Eur. Phys. J. C* **71**, (2011) 1704.
- [45] Granda, L. N.: *JCAP* **1007**, (2010) 006.
- [46] Granda, L. N.: *Rev. Col. Fís.* **42**, (2010) 257.
- [47] Sushkov, S. V.: *Phys. Rev. D* **80**, (2009) 103505.
- [48] Saridakis, E. N. and Sushkov, S. V.: *Phys. Rev. D* **81**, (2010) 083510.
- [49] Gao, C.: *JCAP* **1006**, (2010) 023.
- [50] Skugoreva, M. A., Sushkov S. V. and Toporensky, A. V.: *Phys. Rev. D* **88**, (2013) 083539. [*Phys. Rev. D* **88**, no. 10, (2013) 109906 Erratum]

- [51] Sushkov, S. V.: Phys. Rev. D **85**, (2012) 123520.
- [52] Koutsoumbas, G., Ntrekis K. and Papantonopoulos, E.: JCAP **1308**, (2013) 027.
- [53] Darabi, F. and Parsiya, A.: arXiv:1312.1322 [gr-qc].
- [54] Germani, C. and Kehagias, A.: JCAP **0903**, (2009) 028.
- [55] Germani, C. and Kehagias, A.: Phys. Rev. Lett. **105**, (2010) 011302.
- [56] Germani, C. and Watanabe, Y.: JCAP **1107**, (2011) 031.
- [57] Sadjadi, H. M. and Goodarzi, P.: JCAP **1302**, (2013) 038.
- [58] Tsujikawa, S.: Phys. Rev. D **85**, (2012) 083518.
- [59] Ema, Y., Jinno, R., Mukaida K. and Nakayama, K.: arXiv:1504.07119 [gr-qc].
- [60] Ghalee, A.: arXiv:1402.6798 [astro-ph.CO].
- [61] Yang, N., Gao, Q. and Gong, Y.: arXiv:1504.05839 [gr-qc].
- [62] Myung, Y. S., Moon, T. and Lee, B. H.: arXiv:1505.04027 [gr-qc].
- [63] Sadjadi, H. M.: Phys. Rev. D **83**, (2011) 107301.
- [64] Sadjadi, H. M. and Goodarzi, P.: Phys. Lett. B **732**, (2014) 278.
- [65] Sheikhahmadi, H., Saridakis, E. N., Aghamohammadi, A. and Saaidi, K.: arXiv:1603.03883 [gr-qc].
- [66] Gumjudpai, B. and Rangdee, P.: Gen. Rel. Grav. **47**, (2015) 140.
- [67] Misner, C. W., Thorne, K. S. and Wheeler, J. A.: *Gravitation* (Freeman, 1973).
- [68] Ferraris, M. and Francaviglia, I. and Volovich, I.: Class. Quant. Grav. **11**, (1994) 1505.
- [69] Magnano, G. and Sokolowski, L. M.: Phys. Rev. D **50**, (1994) 5039.
- [70] Capozziello, S. and Faraoni, V.: *Beyond Einstein Gravity* (Springer, 2010).
- [71] Luo, X., Wu, P. and Yu, H.: Astrophys. Space. Sci. **350**, (2014) 831.
- [72] Capozziello, S., Francaviglia, M. and Mercadante, S.: Found. Phys. **39** (2009) 1161.
- [73] Bekenstien, J. D.: Phys. Rev. D **48** (1993) 3641.
- [74] Allemandi, G., Capone, M., Capozziello, S., and Francaviglia, M.: Gen. Rel. Grav. **38**, (2006) 33.
- [75] Santos, J., Santos, C. de Souza.: AIP Conf. Proc. **1471** (2012) 111.
- [76] Zummalacárregui, M., Koivisto, T.S., Mota, D.F. and Ruiz-Lapuente, P.: JCAP **1005** (2010) 038.
- [77] Brustein R., Gorbonos D. and Hadad M.: Phys. Rev. D **79** (2009) 044025.
- [78] Faraoni, V.: Phys. Rev. D **84** (2011) 024003.
- [79] Tsujikawa, S.: JCAP **04** (2015) 043.
- [80] Fujii, Y. and Maeda, K.: *The Scalar-Tensor Theory of Gravitation* (Cambridge, 2003).
- [81] Carroll, S. M.: *Spacetime and Geometry: An Introduction to General Relativity* (Addison-Wesley, 2004).
- [82] Blaschke, D. B. and Dąbrowski, M. P.: Entropy **14**, (2012) 1978.
- [83] Liddle, A. R., Parsons, P. and Barrow, J. D.: Phys. Rev. D **50**, (1994) 7222.
- [84] Linde, A. D., Phys. Lett., B **129** (1983) 177.