

# Global solutions to random 3D vorticity equations for small initial data

Viorel Barbu\*      Michael Röckner<sup>†</sup>

## Abstract

One proves the existence and uniqueness in  $(L^p(\mathbb{R}^3))^3$ ,  $\frac{3}{2} < p < 2$ , of a global mild solution to random vorticity equations associated to stochastic 3D Navier-Stokes equations with linear multiplicative Gaussian noise of convolution type, for sufficiently small initial vorticity. This resembles some earlier deterministic results of T. Kato [15] and are obtained by treating the equation in vorticity form and reducing the latter to a random nonlinear parabolic equation. The solution has maximal regularity in the spatial variables and is weakly continuous in  $(L^3 \cap L^{\frac{3p}{4p-6}})^3$  with respect to the time variable. Furthermore, we obtain the pathwise continuous dependence of solutions with respect to the initial data.

**Keywords:** stochastic Navier-Stokes equation, vorticity, Biot-Savart operator.

**MSC:** 60H15, 35Q30.

## 1 Introduction

Consider the stochastic 3D Navier–Stokes equation

---

\*Octav Mayer Institute of the Romanian Academy and Al.I. Cuza University, Iași, Romania. Email: vb41@uaic.ro

<sup>†</sup>Fakultät für Mathematik, Universität Bielefeld, D-33501 Bielefeld, Germany. Email: roeckner@math.uni-bielefeld.de

$$\begin{aligned}
dX - \Delta X dt + (X \cdot \nabla)X dt &= \sum_{i=1}^N (B_i(X) + \lambda_i X) d\beta_i(t) + \nabla \pi dt \\
&\text{on } (0, \infty) \times \mathbb{R}^3, \\
\nabla \cdot X &= 0 \\
&\text{on } (0, \infty) \times \mathbb{R}^3, \\
X(0) &= x \\
&\text{in } (L^p(\mathbb{R}^3))^3,
\end{aligned} \tag{1.1}$$

where  $\lambda_i \in \mathbb{R}$ ,  $x : \Omega \rightarrow \mathbb{R}^3$  is a random variable. Here  $\pi$  denotes the pressure and  $\{\beta_i\}_{i=1}^N$  is a system of independent Brownian motions on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  with normal filtration  $(\mathcal{F}_t)_{t \geq 0}$ ,  $x$  is  $\mathcal{F}_0$ -measurable, and  $B_i$  are the convolution operators

$$B_i(X)(\xi) = \int_{\mathbb{R}^3} h_i(\xi - \bar{\xi}) X(\bar{\xi}) d\bar{\xi} = (h_i * X)(\xi), \quad \xi \in \mathbb{R}^3, \tag{1.2}$$

where  $h_i \in L^1(\mathbb{R}^3)$ ,  $i = 1, 2, \dots, N$ , and  $\Delta$  is the Laplacian on  $(L^2(\mathbb{R}^3))^3$ .

It is not known whether (1.1) has a probabilistically strong solution in the mild sense for all time. Therefore, we shall rewrite (1.1) in vorticity form and then transform it into a random PDE, which we shall prove, has a global in time solution for  $\mathbb{P}$ -a.e. fixed  $\omega \in \Omega$ , provided the initial condition is small enough.

Consider the vorticity field

$$U = \nabla \times X = \text{curl } X \tag{1.3}$$

and apply the curl operator to equation (1.1). We obtain (see e.g. [4], [8]) the transport vorticity equation

$$\begin{aligned}
dU - \Delta U dt + ((X \cdot \nabla)U - (U \cdot \nabla)X) dt &= \sum_{i=1}^N (h_i * U + \lambda_i U) d\beta_i \\
&\text{in } (0, \infty) \times \mathbb{R}^3,
\end{aligned} \tag{1.4}$$

$$U(0, \xi) = U_0(\xi) = (\text{curl } x)(\xi), \quad \xi \in \mathbb{R}^3.$$

The vorticity  $U$  is related to the velocity  $X$  by the equation

$$X(t, \xi) = K(U(t))(\xi), \quad t \in (0, \infty), \quad \xi \in \mathbb{R}^3, \tag{1.5}$$

where  $K$  is the Biot–Savart integral operator

$$K(u)(\xi) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\xi - \bar{\xi}}{|\xi - \bar{\xi}|^3} \times u(\bar{\xi}) d\bar{\xi}, \quad \xi \in \mathbb{R}^3. \tag{1.6}$$

Then one can rewrite the vorticity equation (1.4) as

$$\begin{aligned} dU - \Delta U dt + ((K(U) \cdot \nabla)U - (U \cdot \nabla)K(U))dt \\ = \sum_{i=1}^N (h_i * U + \lambda_i U) d\beta_i \quad \text{in } (0, \infty) \times \mathbb{R}^3, \\ U(0, \xi) = U_0(\xi), \quad \xi \in \mathbb{R}^3. \end{aligned} \quad (1.7)$$

Equivalently,

$$\begin{aligned} U(t) = e^{t\Delta} U_0 - \int_0^t e^{(t-s)\Delta} ((K(U(s)) \cdot \nabla)U(s) - (U(s) \cdot \nabla)K(U(s))) ds \\ + \int_0^t \sum_{i=1}^N e^{-(t-s)\Delta} (h_i * U(s) + \lambda_i U(s)) d\beta_i(s), \quad t \geq 0. \end{aligned} \quad (1.8)$$

Now, we consider the transformation

$$U(t) = \Gamma(t)y(t), \quad t \in [0, \infty), \quad (1.9)$$

where  $\Gamma(t) : (L^2(\mathbb{R}^3))^3 \rightarrow (L^2(\mathbb{R}^3))^3$  is the linear continuous operator defined by the equations

$$d\Gamma(t) = \sum_{i=1}^N (B_i + \lambda_i I) \Gamma(t) d\beta_i(t), \quad t \geq 0, \quad \Gamma(0) = I, \quad (1.10)$$

where (see (1.2))

$$B_i u = h_i * u, \quad \forall u \in (L^p(\mathbb{R}^3))^3, \quad i = 1, \dots, N, \quad p \in (1, \infty). \quad (1.11)$$

We also set

$$\tilde{B}_i = B_i + \lambda_i I, \quad i = 1, \dots, N, \quad (1.12)$$

where  $I$  is the identity operator.

Since  $B_i B_j = B_j B_i$ , equation (1.10) has a solution  $\Gamma$  and can be equivalently expressed as (see [9], Section 7.4)

$$\Gamma(t) = \prod_{i=1}^N \exp \left( \beta_i(t) \tilde{B}_i - \frac{t}{2} \tilde{B}_i^2 \right), \quad t \geq 0. \quad (1.13)$$

Here (1.10) is meant in the sense that, for every  $z_0 \in (L^2(\mathbb{R}^3))^3$ , the continuous  $(\mathcal{F}_t)$ -adapted  $(L^2(\mathbb{R}^3))^3$ -valued process  $z(t) := \Gamma(t)z_0$ ,  $t \geq 0$ , solves the following SDE on  $H := (L^2(\mathbb{R}^3))^3$ ,

$$dz(t) = \sum_{i=1}^N \tilde{B}_i z(t) d\beta_i(t), \quad z(0) = z_0,$$

where  $H$  is equipped with the usual scalar product  $\langle \cdot, \cdot \rangle$ .

Applying the Itô formula in (1.7) (the justification for this is as in [2]), we obtain for  $y$  the random differential equation

$$\begin{aligned} \frac{dy}{dt}(t) - \Gamma^{-1}(t)\Delta(\Gamma(t)y(t)) + \Gamma^{-1}(t)(K(\Gamma(t)y(t)) \cdot \nabla)(\Gamma(t)y(t)) \\ - (\Gamma(t)y(t) \cdot \nabla)(K(\Gamma(t)y(t))) = 0, \quad t \in [0, \infty), \end{aligned} \quad (1.14)$$

$y(0) = U_0$ .

Taking into account that, for all  $i$ ,  $B_i\Delta = \Delta B_i$  on  $H^2(\mathbb{R}^3)$ , it follows by (1.10), (1.13) that  $\Delta\Gamma(t) = \Gamma(t)\Delta$  on  $H^2(\mathbb{R}^3)$ ,  $\forall t \geq 0$ .

In what follows, equation (1.14) will be taken in the following mild sense

$$y(t) = e^{t\Delta}U_0 + \int_0^t e^{(t-s)\Delta}\Gamma^{-1}(s)M(\Gamma(s)y(s))ds, \quad t \in [0, \infty), \quad (1.15)$$

where

$$(e^{t\Delta}u)(\xi) = \frac{1}{(4\pi t)^{\frac{3}{2}}} \int_{\mathbb{R}^3} \exp\left(-\frac{|\xi - \bar{\xi}|^2}{4t}\right) u(\bar{\xi})d\bar{\xi}, \quad t \in [0, \infty), \quad \xi \in \mathbb{R}^3, \quad (1.16)$$

and  $M$  is defined by

$$M(u) = -[(K(u) \cdot \nabla)(u) - (u \cdot \nabla)(K(u))], \quad t \in [0, \infty). \quad (1.17)$$

We note that  $U(t) = \Gamma(t)y(t)$  is the solution to the equation

$$U(t) = e^{t\Delta}\Gamma(t)U_0 + \int_0^t e^{(t-s)\Delta}\Gamma(t)\Gamma^{-1}(s)M(U(s))ds, \quad (1.18)$$

which may be viewed as the random version of the stochastic vorticity equation (1.8).

Our aim here and the principal contribution of this work is to show that, for every  $\varepsilon \in (0, 1)$ , there exists  $\Omega_\varepsilon \in \mathcal{F}$  such that  $\mathbb{P}(\Omega_\varepsilon) \geq 1 - \varepsilon$  and, for all  $\omega \in \Omega_\varepsilon$ , we have the existence and uniqueness of a solution (in the mild sense) for (1.15) if the vorticity of  $x$ , i.e.,  $U_0 = \text{curl } x$ , is  $\mathbb{P}$ -a.s. sufficiently small in a sense to be made precise in Theorem 1.1 below. We recall that, for a deterministic Navier–Stokes equation, such a result was first established by T. Kato [15] (see also T. Kato and H. Fujita [16]) and extended later to more general initial data by Y. Giga and T. Miyakawa [14], M. Taylor [21], H. Koch and D. Tataru [17]. However, the standard approach [15], [16] cannot be applied in the present case for one reason: the nonlinear inertial term  $(X \cdot \nabla)X$  cannot be conveniently estimated in the space  $C_b([0, \infty); L^p(\Omega \times \mathbb{R}^d))$  and similarly for the nonlinearity arising in (1.7). As regards the stochastic 3D Navier-Stokes equations, to best of our knowledge all global existence results were limited to martingale solutions. Since the fundamental work [11], the literature on (global) martingale solutions for stochastic 3D-Navier-Stokes equations has grown enormously. We refer, e.g., to [6], [10], [12], [13], [18], and the references therein.

In the following, we denote by  $L^p$ ,  $1 \leq p \leq \infty$ , the space  $(L^p(\mathbb{R}^3))^3$  with the norm  $|\cdot|_p$ , by  $W^{1,p}$  the corresponding Sobolev space and by  $C_b([0, \infty); L^p)$  the space of all bounded and continuous functions  $u : [0, \infty) \rightarrow L^p$  with the sup norm. We also set  $D_i = \frac{\partial}{\partial \xi_i}$ ,  $i = 1, 2, 3$ , and denote by  $\nabla \cdot u$  the divergence of  $u$ , while

$$((u \cdot \nabla)v)_j = u_i D_i v_j, \quad j = 1, 2, 3, \quad u = \{u_i\}_{i=1}^3, \quad v = \{v_j\}_{j=1}^3.$$

As usual

$$q' = \frac{q}{q-1} \quad \text{for } q \in (1, \infty).$$

We set for  $p \in (\frac{3}{2}, 3)$

$$\eta(t) = \|\Gamma(t)\|_{L(L^p, L^p)} \|\Gamma(t)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \|\Gamma^{-1}(t)\|_{L^q, L^q}, \quad t \geq 0, \quad (1.19)$$

where for  $q \in (1, \infty)$ ,  $\|\cdot\|_{L(L^q, L^q)}$  is the norm of the space  $L(L^q, L^q)$  of linear continuous operators on  $L^q$ .

For  $p \in [1, \infty)$ , we denote by  $\mathcal{Z}_p$  the space of all functions  $y : (0, \infty) \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$  such that

$$\begin{aligned} t^{1-\frac{3}{2p}} y &\in C_b([0, \infty); L^p), \\ t^{\frac{3}{2}(1-\frac{1}{p})} D_i y &\in C_b([0, \infty); L^p), \quad i = 1, 2, 3. \end{aligned} \quad (1.20)$$

The space  $\mathcal{Z}_p$  is endowed with the norm

$$\|y\|_{p,\infty} = \sup \left\{ t^{1-\frac{3}{2p}} |y(t)|_p + t^{\frac{3}{2}(1-\frac{1}{p})} |D_i y(t)|_p; t \in (0, \infty), i = 1, 2, 3 \right\}. \quad (1.21)$$

In the following, we take  $\lambda_i \in \mathbb{R}$  such that

$$|\lambda_i| > (\sqrt{12} + 3) |h_i|_1, \quad \forall i = 1, 2, \dots, N. \quad (1.22)$$

We note that

$$\|B_i\|_{L(L^q, L^q)} \leq |h_i|_{L^1}, \quad \forall i = 1, \dots, N.$$

Theorem 1.1 is the main result.

**Theorem 1.1.** *Let  $p, q \in (1, \infty)$  such that*

$$\frac{3}{2} < p < 2, \quad \frac{1}{q} = \frac{2}{p} - \frac{1}{3}. \quad (1.23)$$

Let  $\Omega_0 = \{\sup_{t \geq 0} \eta(t) < \infty\}$  and consider (1.15) for fixed  $\omega \in \Omega_0$ . Set  $\Gamma(t) := \Gamma(t)(\omega)$ ,  $\eta(t) := \eta(t, \omega)$ . Then  $\mathbb{P}(\Omega_0) = 1$  and there is a positive constant  $C^*$  independent of  $\omega \in \Omega_0$  such that, if  $U_0 \in L^{\frac{3}{2}}$  is such that

$$\sup_{t \geq 0} \eta(t) |U_0|_{\frac{3}{2}} \leq C^*, \quad (1.24)$$

then the random equation (1.15) has a unique solution  $y \in \mathcal{Z}_p$  which satisfies

$$M(\Gamma(t)y) \in L^1(0, T; L^q). \quad (1.25)$$

Moreover, for each  $\varphi \in L^3 \cap L^{q'}$ , the function  $t \rightarrow \int_{\mathbb{R}^3} y(t, \xi) \varphi(\xi) d\xi$  is continuous on  $[0, \infty)$ . The map  $U_0 \rightarrow y$  is Lipschitz from  $L^{\frac{3}{2}}$  to  $\mathcal{Z}_p$ .

In particular, the random vorticity equation (1.18) has a unique solution  $U$  such that  $\Gamma^{-1}U \in \mathcal{Z}_p$ .

**Remark 1.2.** Concerning condition (1.24), we note that an elementary calculation shows that

$$\eta(t) \leq \prod_{i=1}^N \exp(3|\beta_i(t)|(|h_i|_1 + |\lambda_i|) - t\alpha_i), \quad t \in [0, \infty),$$

where  $\alpha_i := \frac{1}{2} \lambda_i^2 - \frac{3}{2} (|h_i|_1^2 + 2|\lambda_i| |h_i|_1)$ , which is strictly positive by (1.22).

By the law of the iterated logarithm, it follows that

$$\sup_{t \geq 0} \eta(t) < \infty, \quad \mathbb{P}\text{-a.e.},$$

hence for  $\Omega_r := \{\sup_{t \geq 0} \eta(t) \leq r\}$  we have  $\mathbb{P}(\Omega_r^c) \rightarrow 0$  as  $r \rightarrow \infty$ .

But, taking into account that, for each  $r > 0$  and all  $\nu > 0$ ,  $i = 1, \dots, N$ , we have (see Lemma 3.4 in [1])

$$\mathbb{P} \left[ \sup_{t \geq 0} \{\exp(\beta_i(t) - \nu t)\} \geq r \right] = r^{-2\nu},$$

and, more explicitly, we get that

$$\mathbb{P}(\Omega_r^c) \leq 2Nr^{-\frac{N\alpha}{\gamma^2}}, \quad \forall r > 0,$$

where  $\alpha = \min_{1 \leq i \leq N} \alpha_i$ ,  $\gamma = 3 \max\{(|h_i|_1 + |\lambda_i|); i \leq N\}$ . Therefore, if  $\omega \in \Omega_r$  and  $U_0 = U_0(\omega)$  satisfies

$$|U_0|_{\frac{3}{2}} \leq \frac{C^*}{r}, \tag{1.26}$$

then condition (1.24) holds. It is trivial to define such an  $\mathcal{F}$ -measurable function  $U_0 : \Omega \rightarrow L^{\frac{3}{2}}$ , for instance,

$$U_0 := \sum_{n=1}^{\infty} \frac{C^*}{n} \mathbf{1}_{\{n-1 \leq \sup_{t \geq 0} \eta(t) < n\}} u_0,$$

for some  $u_0 \in L^{\frac{3}{2}}$ . But, of course,  $U_0$  is not  $\mathcal{F}_0$ -measurable and so the process  $U(t)$ ,  $t \geq 0$ , given by Theorem 1.1, is not  $(\mathcal{F}_t)_{t \geq 0}$ -adapted. Therefore,  $U = \Gamma(t)y$  is not a solution to the stochastic vorticity equation (1.8). However, it can be viewed as a generalized solution to (1.8).

It should also be mentioned that assumption (1.22) is not necessary for existence of a solution to equation (1.15), but only to make sure that condition (1.24) is not void.

## 2 Proof of Theorem 1.1

To begin with, we note below in Lemma 2.1 a few immediate properties of the operator  $\Gamma$  defined in (1.10)–(1.13).

**Lemma 2.1.** *We have*

$$|\Gamma(t)z|_q + |\Gamma^{-1}(t)z|_q \leq C_t |z|_q, \quad t \in [0, \infty), \quad \forall z \in L^q, \quad \forall q \in [1, \infty), \quad (2.1)$$

and

$$|\nabla(\Gamma(t)z)|_q \leq \|\Gamma(t)\|_{L(L^q, L^q)} |\nabla z|_q, \quad \text{for all } z \in W^{1,q}(\mathbb{R}^3). \quad (2.2)$$

**Proof.** By (1.2), (1.11) and by the Young inequality, we see that

$$|B_i(u)|_q \leq (|h_i|_1 + |\lambda_i|) |u|_q, \quad \forall u \in L^q, \quad i = 1, \dots, N. \quad (2.3)$$

Recalling (1.13), we see by (2.3) that (2.1), (2.2) hold, as claimed.  $\blacksquare$

**Lemma 2.2.** *Let  $\frac{1}{q} = \frac{1}{r_1} + \frac{1}{r_2}$ ,  $\frac{3}{2} < r_1 < \infty$ ,  $r_1^* = \frac{3r_1}{3+r_1}$ ,  $1 < q < \infty$ . Then, for some  $C > 0$  independent of  $\omega$ ,*

$$|M(\Gamma(t)z)|_q \leq C \|\Gamma(t)\|_{L(L^{r_1}, L^{r_1})} \|\Gamma(t)\|_{L(L^{r_2}, L^{r_2})} (|z|_{r_1} |z|_{r_2} + |z|_{r_1^*} |\nabla z|_{r_2}), \quad (2.4)$$

$$t \in [0, \infty),$$

for all  $z \in L^{r_1} \cap L^{r_2} \cap L^{r_1^*}$  with  $\nabla z \in L^{r_2}$ .

**Proof.** We have by (1.17) and (2.1)

$$|M(\Gamma(t)z)|_q \leq |(K(\Gamma(t)z) \cdot \nabla)(\Gamma(t)z)|_q + |(\Gamma(t)z \cdot \nabla)K(\Gamma(t)z)|_q. \quad (2.5)$$

On the other hand, by (2.2) and the Hölder inequality we have

$$\begin{aligned} |(K(\Gamma(t)z) \cdot \nabla)(\Gamma(t)z)|_q &\leq |K(\Gamma(t)z)|_{r_1} |\nabla(\Gamma(t)z)|_{r_2} \\ &\leq \|\Gamma(t)\|_{L(L^{r_1}, L^{r_1})} \|\Gamma(t)\|_{L(L^{r_2}, L^{r_2})} |K(z)|_{r_1} |\nabla z|_{r_2}. \end{aligned} \quad (2.6)$$

Now, we recall the classical estimate for Riesz potentials (see [20], p. 119)

$$\left| \int_{\mathbb{R}^3} \frac{f(\bar{\xi})}{|\xi - \bar{\xi}|^2} d\xi \right|_{\beta} \leq C |f|_{\alpha}, \quad \forall f \in L^{\alpha},$$

where  $\frac{1}{\beta} = \frac{1}{\alpha} - \frac{1}{3}$ ,  $\alpha \in (1, 3)$ . By virtue of (1.6), this yields

$$|K(u)|_{\beta} \leq C |u|_{\alpha}; \quad \forall u \in L^{\alpha}, \quad \frac{1}{\beta} = \frac{1}{\alpha} - \frac{1}{3}, \quad (2.7)$$

and so, for  $\beta = r_1$ ,  $\alpha = \frac{3r_1}{3+r_1} = r_1^*$ , we get by (2.2) and (2.6) the estimate

$$|(K(\Gamma(t)z) \cdot \nabla)(\Gamma(t)z)|_q \leq C \|\Gamma(t)\|_{L(L^{r_1}, L^{r_1})} \|\Gamma(t)\|_{L(L^{r_2}, L^{r_2})} |z|_{r_1^*} |\nabla z|_{r_2}. \quad (2.8)$$



(Here and everywhere in the following,  $|\nabla z|_p$  means  $\sup\{|D_i z|_p; i = 1, 2, 3\}$ .) Taking into account that, by the Calderon–Zygmund inequality (see [7], Theorem 1),

$$|\nabla K(z)|_{\tilde{p}} \leq C|z|_{\tilde{p}}, \quad \forall z \in L^{\tilde{p}}, \quad 1 \leq \tilde{p} < \infty, \quad (2.9)$$

one obtains that

$$|(\Gamma(t)z \cdot \nabla)(K(t)\Gamma(t)z)|_q \leq C\|\Gamma(t)\|_{L(L^{r_1}, L^{r_1})}\|\Gamma(t)\|_{L(L^{r_2}, L^{r_2})}|z|_{r_1}|z|_{r_2}. \quad (2.10)$$

Substituting (2.8), (2.10) in (2.5), one obtains estimate (2.4), as claimed. ■

**Lemma 2.3.** *Let  $r_1 = \frac{3r_2}{3-r_2}$ ,  $\frac{3}{2} < r_2 < 3$ ,  $q = \frac{3r_1}{r_1+6}$ . Then, we have, for some  $C > 0$  independent of  $\omega$ ,*

$$|M(\Gamma(t)z)|_q \leq C\|\Gamma(t)\|_{L(L^{r_1}, L^{r_1})}\|\Gamma(t)\|_{L(L^{r_2}, L^{r_2})}|z|_{r_2}|\nabla z|_{r_2}, \quad \forall z \in W^{1, r_2}. \quad (2.11)$$

**Proof.** We have by the Sobolev–Gagliardo–Nirenberg inequality (see, e.g., [5], p. 278)

$$|z|_{r_1} \leq C|\nabla z|_{r_2}, \quad \forall z \in W^{1, r_2}(\mathbb{R}^3).$$

Substituting in (2.4) and taking into account that  $r_1^* = r_2$ , we obtain (2.11), as claimed. ■

In the following, we fix  $p = r_2$ ,  $r_1$  and  $q$  as in Lemma 2.3, (2.11), that is,

$$\frac{3}{2} < p < 2, \quad r_1 = \frac{3p}{3-p}, \quad \frac{1}{q} = \frac{2}{p} - \frac{1}{3}. \quad (2.12)$$

We write equation (1.15) as

$$y(t) = G(y)(t) = e^{t\Delta}U_0 + F(y)(t), \quad t \in [0, \infty), \quad (2.13)$$

where

$$F(z)(t) = \int_0^t e^{(t-s)\Delta}\Gamma^{-1}(s)M(\Gamma(s)z(s))ds, \quad t \in [0, \infty). \quad (2.14)$$

By (1.16), we have for  $1 < \tilde{q} \leq \tilde{p} < \infty$  the estimates

$$|e^{t\Delta}u|_{\tilde{p}} \leq Ct^{-\frac{3}{2}(\frac{1}{\tilde{q}} - \frac{1}{\tilde{p}})}|u|_{\tilde{q}}, \quad u \in L^{\tilde{q}}, \quad (2.15)$$

$$|D_j e^{t\Delta}u|_{\tilde{p}} \leq Ct^{-\frac{3}{2}(\frac{1}{\tilde{q}} - \frac{1}{\tilde{p}}) - \frac{1}{2}}|u|_{\tilde{q}}, \quad j = 1, 2, 3. \quad (2.16)$$

(Everywhere in the following, we shall denote by  $C$  several positive constants independent of  $\omega$  and  $t \geq 0$ .)

We apply (2.15), with  $\tilde{q} = q$ ,  $\tilde{p} = p$ . By (2.11)–(2.14), we obtain that

$$\begin{aligned} |F(z(t))|_p &\leq C \int_0^t (t-s)^{-\frac{1}{2}(\frac{3}{p}-1)} |\Gamma^{-1}(s)M(\Gamma(s)z(s))|_q ds \\ &\leq C \int_0^t (t-s)^{-\frac{1}{2}(\frac{3}{p}-1)} \|\Gamma(s)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \\ &\quad \|\Gamma(s)\|_{L(L^p, L^p)} \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} |z(s)|_p |\nabla z(s)|_p ds. \end{aligned} \quad (2.17)$$

Similarly, we obtain by (2.16) that

$$\begin{aligned} |D_j F(z(t))|_p &\leq C \int_0^t (t-s)^{-\frac{3}{2p}} \|\Gamma(s)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \\ &\quad \|\Gamma(s)\|_{L(L^p, L^p)} \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} |z(s)|_p |\nabla z(s)|_p ds, \quad j = 1, 2, 3. \end{aligned} \quad (2.18)$$

We consider the Banach space  $\mathcal{Z}_p$  defined by (1.20), that is,

$$\mathcal{Z}_p = \left\{ y; t^{1-\frac{3}{2p}} y \in C_b([0, \infty); L^p), t^{\frac{3}{2}(1-\frac{1}{p})} D_j y \in C_b([0, \infty); L^p), \right. \\ \left. j = 1, 2, 3 \right\}, \quad (2.19)$$

with the norm

$$\|z\|_{p, \infty} = \|z\| = \sup_{t>0} \left\{ \left( t^{1-\frac{3}{2p}} |z(t)|_p + t^{\frac{3}{2}(1-\frac{1}{p})} |D_i z(t)|_p \right), i = 1, 2, 3 \right\}. \quad (2.20)$$

We note that

$$|z(t)|_p |\nabla z(t)|_p \leq C t^{-\frac{5}{2}+\frac{3}{p}} \|z\|^2, \quad \forall z \in \mathcal{Z}_p, \quad t \in (0, \infty). \quad (2.21)$$

By (2.17) and (2.20) we see that, for  $z \in \mathcal{Z}_p$ , we have

$$\begin{aligned} |F(z(t))|_p &\leq \int_0^t (t-s)^{-\frac{1}{2}(\frac{3}{p}-1)} \|\Gamma(s)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \|\Gamma(s)\|_{L(L^p, L^p)} \\ &\quad \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} |z(s)|_p |\nabla z(s)|_p ds \\ &\leq C \int_0^t (t-s)^{-\frac{1}{2}(\frac{3}{p}-1)} |s|^{-\frac{5}{2}+\frac{3}{p}} \|\Gamma(s)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \\ &\quad \|\Gamma(s)\|_{L(L^p, L^p)} \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} ds \|z\|^2 \\ &\leq C t^{\frac{3}{2p}-1} \sup_{0 \leq s \leq t} \eta(s) \int_0^1 (1-s)^{-\frac{1}{2}(\frac{3}{p}-1)} s^{-\frac{5}{2}+\frac{3}{p}} ds \|z\|^2, \quad \forall t > 0, \end{aligned} \quad (2.22)$$

where  $\eta$  is given by (1.19). This yields

$$t^{1-\frac{3}{2p}}|F(z(t))|_p \leq C \sup_{0 \leq s \leq t} \{\eta(s)\} B\left(\frac{3}{2}\left(\frac{2}{p}-1\right), \frac{3}{2}\left(1-\frac{1}{p}\right)\right) \|z\|^2, \quad \forall t > 0, \quad (2.23)$$

where  $B$  is the classical beta function (which is finite by virtue of (1.23)).

Similarly, by (2.16) and (2.21), we have, for  $j = 1, 2, 3$ ,

$$\begin{aligned} |D_j F(z)(t)|_p &\leq C \int_0^t (t-s)^{-\frac{3}{2p}} s^{-\frac{5}{2}+\frac{3}{p}} \|\Gamma(s)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})} \\ &\quad \|\Gamma(s)\|_{L(L^p, L^p)} \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} ds \|z\|^2 \quad (2.24) \\ &\leq C \sup_{0 \leq s \leq t} \{\eta(s)\} t^{-\frac{3}{2}(1-\frac{1}{p})} B\left(3\left(\frac{1}{p}-\frac{1}{2}\right), 1-\frac{3}{2p}\right) \|z\|^2, \quad t > 0. \end{aligned}$$

Hence,

$$t^{\frac{3}{2}(1-\frac{1}{p})} |D_j F(z(t))|_p \leq C \sup_{0 \leq s \leq t} \eta(s) \|z\|^2, \quad \forall z \in \mathcal{Z}_p, \quad t > 0, \quad j = 1, 2, 3. \quad (2.25)$$

By (2.15)–(2.16), we have

$$\begin{aligned} |e^{t\Delta} U_0|_p &\leq C t^{\frac{3}{2p}-1} |U_0|_{\frac{3}{2}}, \quad t > 0, \\ |D_j e^{t\Delta} U_0|_p &\leq C t^{\frac{3}{2p}-\frac{3}{2}} |U_0|_{\frac{3}{2}}, \quad t > 0, \quad j = 1, 2, 3. \end{aligned}$$

Therefore, by (2.20) we get

$$\|e^{t\Delta} U_0\| \leq C |U_0|_{\frac{3}{2}}. \quad (2.26)$$

By (2.20), (2.23), (2.25), (2.26), we get,

$$\|G(z)\| \leq C_1 \left( |U_0|_{\frac{3}{2}} + \sup_{t \geq 0} \eta(t) \|z\|^2 \right), \quad \forall z \in \mathcal{Z}_p, \quad (2.27)$$

where  $C_1 > 0$  is independent of  $\omega$ .

We set

$$\eta_\infty = \sup_{t \geq 0} \eta(t), \quad (2.28)$$

and so (2.27) yields

$$\|G(z)\| \leq C_1 (|U_0|_{\frac{3}{2}} + \eta_\infty \|z\|^2), \quad \forall z \in \mathcal{Z}_p. \quad (2.29)$$

We set

$$\Sigma = \{z \in \mathcal{Z}_p; \|z\| \leq R^*\}$$

and note that, by (2.29), it follows that  $G(\Sigma) \subset \Sigma$  if

$$|U_0|_{\frac{3}{2}} \eta_\infty \leq (4C_1^2)^{-1}, \quad (2.30)$$

(so  $U_0$  must depend on  $\omega$ ) and  $R^* = R^*(\omega)$  is given by

$$R^* = 2C_1 |U_0|_{\frac{3}{2}}. \quad (2.31)$$

(We recall that  $C_1$  is independent of  $\omega$  and  $U_0$ .) Moreover, by (1.17) and (2.14), we have, for all  $z, \bar{z} \in \mathcal{Z}_p$ ,

$$\begin{aligned} G(z)(t) - G(\bar{z})(t) &= - \int_0^t e^{(t-s)\Delta} \Gamma^{-1}(s) [(K\Gamma(s)(z(s) - \bar{z}(s)) \cdot \nabla) \Gamma(s) z(s) \\ &+ (K(\Gamma(s)\bar{z}(s)) \cdot \nabla) \Gamma(s)(z(s) - \bar{z}(s)) - \Gamma(s)(z(s) - \bar{z}(s)) \cdot \nabla K(\Gamma(s)z(s)) \\ &- (\Gamma(s)\bar{z}(s) \cdot \nabla) K(\Gamma(s)(z(s) - \bar{z}(s)))] ds. \end{aligned}$$

Proceeding as above, we get, as in (2.17), (2.22), (2.23), that

$$\begin{aligned} &|G(z)(t) - G(\bar{z})(t)|_p \\ &\leq C \int_0^t (t-s)^{-\frac{1}{2}(\frac{3}{p}-1)} \|\Gamma(s)\|_{L(L^{\frac{3p}{p-3}}, L^{\frac{3p}{p-3}})} \|\Gamma(s)\|_{L(L^p, L^p)} \\ &\quad \|\Gamma^{-1}(s)\|_{L(L^q, L^q)} (|z(s) - \bar{z}(s)|_p (|\nabla z(s)|_p + |\nabla \bar{z}(s)|_p) \\ &\quad + |\nabla z(s) - \nabla \bar{z}(s)|_p (|z(s)|_p + |\bar{z}(s)|_p)) ds \\ &\leq Ct^{-(1-\frac{3}{2p})} \sup_{0 \leq s \leq t} \eta(s) \|z - \bar{z}\| (\|z\| + \|\bar{z}\|), \quad \forall t > 0, \end{aligned} \quad (2.32)$$

and also (see (2.18), (2.24), (2.25))

$$|D_j G(z)(t) - D_j G(\bar{z}(t))|_p \leq Ct^{-\frac{3}{2}(1-\frac{1}{p})} \sup_{0 \leq s \leq t} \eta(s) \|z - \bar{z}\| (\|z\| + \|\bar{z}\|), \quad \forall t > 0,$$

for  $j = 1, 2, 3$ . Hence, by (2.20) and (2.28), we obtain that

$$\|G(z) - G(\bar{z})\| \leq C_2 \eta_\infty R^* \|z - \bar{z}\|, \quad \forall z, \bar{z} \in \Sigma, \quad (2.33)$$

where  $C_2$  is independent of  $\omega$ .

Then, by (2.31), (2.33), it follows that, if (2.30) and

$$2C_1C_2\eta_\infty|U_0|_{\frac{3}{2}} < 1, \quad (2.34)$$

hold, then the operator  $G$  is a contraction on  $\Sigma$  and so there is a unique solution  $U \in \Sigma$  to (1.15) provided (1.24) holds with  $C^* < (2C_1C_2)^{-1}$ .

Now, as seen earlier, by (2.11), (1.15) and (2.21) we have

$$\begin{aligned} |M(\Gamma(t)y(t))|_q &\leq C\|\Gamma(t)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})}\|\Gamma(t)\|_{L(L^p, L^p)}|y(t)|_p|\nabla y(t)|_p \\ &\leq C\|\Gamma(t)\|_{L(L^{\frac{3p}{3-p}}, L^{\frac{3p}{3-p}})}\|\Gamma(t)\|_{L(L^p, L^p)}t^{-\frac{5}{2}+\frac{3}{p}}\|y\|^2, \quad \forall t > 0. \end{aligned} \quad (2.35)$$

On the other hand, we have for all  $\varphi \in C_0^\infty(\mathbb{R}^3)$ ,

$$\begin{aligned} \int_{\mathbb{R}^3} y(t, \xi) \cdot \varphi(\xi) d\xi &= \int_{\mathbb{R}^3} (e^{t\Delta})U_0(\xi) \cdot \varphi(\xi) d\xi \\ &+ \int_0^t \int_{\mathbb{R}^3} \Gamma^{-1}(s)M(\Gamma(s)y(s)) \cdot e^{(t-s)\Delta}\varphi(\xi) d\xi ds. \end{aligned} \quad (2.36)$$

Recalling that, for all  $1 \leq \tilde{p} < \infty$ ,  $|e^{t\Delta}\varphi|_{\tilde{p}} \leq |\varphi|_{\tilde{p}}$ , it follows by (2.35) that

$$\begin{aligned} &\left| \int_0^t \int_{\mathbb{R}^3} \Gamma^{-1}(s)M(\Gamma(s)y(s)) \cdot e^{(t-s)\Delta}\varphi(\xi) d\xi ds \right| \\ &\leq C \sup_{0 \leq s \leq t} \eta(s) \int_0^t s^{-\frac{5}{2}+\frac{3}{p}} ds \|y\|^2 |\varphi|_{q'} \\ &\leq C \sup_{0 \leq s \leq t} \eta(s) t^{\frac{3}{p}-\frac{3}{2}} \|y\|^2 |\varphi|_{q'}, \quad \forall t \in (0, \infty). \end{aligned} \quad (2.37)$$

We also have by (2.26)

$$\left| \int_{\mathbb{R}^3} e^{t\Delta}U_0(\xi)\varphi(\xi) d\xi \right| \leq C|U_0|_{\frac{3}{2}}|\varphi|_3, \quad \forall t \in [0, \infty).$$

Combining the latter with (2.36), (2.37), we obtain that, for  $T > 0$ ,

$$\left| \int_{\mathbb{R}^3} y(t, \xi) \cdot \varphi(\xi) d\xi \right| \leq CT^{\frac{3}{p}-\frac{3}{2}}(|\varphi|_{q'} + |\varphi|_3), \quad \forall \varphi \in L^{q'} \cap L^3, \quad t \in [0, T].$$

Hence, by (2.36) and since  $t \rightarrow e^{t\Delta}U_0$  is continuous on  $L^{\frac{3}{2}}$ , the function  $t \rightarrow y(t)$  is  $L^3 \cap L^{q'}$  weakly continuous on  $[0, \infty)$ , where  $q' = \frac{3p}{4p-6}$ .

If  $U = y(t, U_0) \in \mathcal{Z}_p$  is the solution to (1.15), equivalently (2.13), we have for all  $U_0, \bar{U}_0$  satisfying (1.24) (see (2.26) and (2.33))

$$\begin{aligned} \|y(\cdot, U_0) - y(\cdot, \bar{U}_0)\| &\leq \|e^{t\Delta}(U_0 - \bar{U}_0)\| + \|F(y(\cdot, U_0)) - F(y(\cdot, \bar{U}_0))\| \\ &\leq C|U_0 - \bar{U}_0|_{\frac{3}{2}} + \eta_\infty R^* C_2 \|y(\cdot, U_0) - y(\cdot, \bar{U}_0)\|. \end{aligned}$$

Recalling that by (2.31) and (2.34) we have  $R^* C_2 \eta_\infty < 1$ , this yields

$$\|y(\cdot, U_0) - y(\cdot, \bar{U}_0)\| \leq \frac{C}{1 - R^* C_2 \eta_\infty} |U_0 - \bar{U}_0|_{\frac{3}{2}} \leq C(\omega) |U_0 - \bar{U}_0|_{\frac{3}{2}},$$

and so, the map  $y \rightarrow U(\cdot, U_0)$  is Lipschitz from  $L^{\frac{3}{2}}$  to  $\mathcal{Z}_p$ . This completes the proof of Theorem 1.1.  $\blacksquare$

It should be noted that, by (2.30) and (2.31), we have by the Fernique theorem

$$|U_0|_{\frac{3}{2}}, R^* \in \bigcap_{r \geq 1} L^r(\Omega),$$

and so, taking into account that  $y \in \Sigma$ , we see by (2.19), (2.20) that

$$\sup_{t \geq 0} \left\{ t^{1 - \frac{3}{2p}} |y(t)|_p + t^{\frac{3}{2} \left(1 - \frac{1}{p}\right)} |D_i y(t)|_p \right\} \in \bigcap_{r \geq 1} L^r(\Omega), \quad i = 1, 2, 3. \quad (2.38)$$

We have, therefore, the following completion of Theorem 1.1.

**Corollary 2.4.** *Under the assumptions of Theorem 1.1, the solution  $y = y(t, \omega)$  to the equation (1.15) satisfies (2.38). The same result holds for the solution  $U(t) = \Gamma(t)y(t)$  of the random vorticity equation (1.18).*

### 3 The random version of the 3D Navier-Stokes equation

We fix in (1.1) the initial random variable  $x$  by the formula

$$x = K(U_0), \quad (3.1)$$

where  $U_0 = \text{curl } x$ ,  $U_0 = U_0(\omega)$  satisfies condition (1.24) for all  $\omega \in \Omega_0$  (see Remark 1.2). If  $y$  is the corresponding solution to equation (1.15) given by Theorem 1.1, we define the process  $X$  by formula (1.5), that is,

$$X(t) = K(U(t)) = K(\Gamma(t)y(t)), \quad \forall t \in [0, \infty). \quad (3.2)$$

By (2.7), where  $U$  is the solution to the vorticity equation (1.1), we have

$$|X(t)|_{\frac{3p}{3-p}} \leq C|U(t)|_p, \quad \forall t \in [0, \infty). \quad (3.3)$$

(Everywhere in the following,  $C$  are positive constants independent of  $\omega \in \Omega$ .)

On the other hand, by the Carlderon–Zygmund inequality (2.9), we have

$$|D_i X(t)|_p \leq C|U(t)|_p, \quad i = 1, 2, 3. \quad (3.4)$$

By (3.3) and by Theorem 1.1, it follows that

$$t^{1-\frac{3}{2p}} X \in C_b([0, \infty); L^{\frac{3p}{3-p}}), \quad (3.5)$$

while, by (3.4), we have for  $i = 1, 2, 3$

$$t^{\frac{3}{2}(1-\frac{1}{p})} D_i X \in C_b([0, \infty); L^p). \quad (3.6)$$

Now, if in (2.9) we take  $z = D_j X$ , we get that, for all  $i, j = 1, 2, 3$ ,

$$t^{\frac{3}{2}(1-\frac{1}{p})} |D_i D_j X|_p \leq C t^{\frac{3}{2}(1-\frac{1}{p})} |D_i U|_p \leq C, \quad \forall t \in [0, \infty).$$

This yields

$$t^{\frac{3}{2}(1-\frac{1}{p})} D_{ij}^2 X \in C_b([0, \infty); L^p), \quad i, j = 1, 2, 3. \quad (3.7)$$

Moreover, by Corollary 2.4, we also have

$$t^{1-\frac{3}{2p}} X \in C_b([0, \infty); L^r(\Omega; L^{\frac{3p}{3-p}})), \quad \forall r \geq 1, \quad (3.8)$$

$$t^{\frac{3}{2}(1-\frac{1}{p})} D_i X \in C_b([0, \infty); L^r(\Omega; L^p)), \quad i = 1, 2, 3, \quad (3.9)$$

$$t^{\frac{3}{2}(1-\frac{1}{p})} D_{ij} X \in C_b([0, \infty); L^r(\Omega; L^p)), \quad i, j = 1, 2, 3. \quad (3.10)$$

Now, if in equation (1.18) one applies the Biot–Savart operator  $K$ , we obtain for  $X$  the equation

$$X(t) = K(e^{t\Delta}\Gamma(t)\text{curl } x) + \int_0^t K(e^{(t-s)\Delta}\Gamma(t)\Gamma^{-1}(s)M(\text{curl } X(s)))ds, \quad (3.11)$$

$t \geq 0,$

where  $M$  is given by (1.17). It should be noted that, by virtue of (3.7)-(3.10), the right hand side of (3.11) is well defined.

Equation (3.11) can be viewed as the random version of the Navier-Stokes equation (1.1). However, since, as seen earlier,  $U_0$  is not  $\mathcal{F}_0$ -measurable, the processes  $t \rightarrow y(t)$ ,  $t \rightarrow U(t)$  are not  $(\mathcal{F}_t)_{t \geq 0}$ -adapted, and so  $X$  is not  $(\mathcal{F}_t)_{t \geq 0}$ -adapted, too. Therefore, (3.11) cannot be transformed back into (1.1). By Theorem 1.1, it follows that

**Theorem 3.1.** *Under assumptions (1.24), the random Navier-Stokes equation (3.11) has a unique solution  $X$  satisfying (3.7)-(3.10).*

**Remark 3.2.** As easily seen from the proofs, Theorem 1.1 extends mutatis-mutandis to the noises  $\sum_{i=1}^N \sigma_i(t, X) \dot{\beta}_i(t)$ , where

$$\sigma_i(t, x)(\xi) = \int_{\mathbb{R}^3} h_i(t, \xi - \bar{\xi}) x(\bar{\xi}) d\bar{\xi}, \quad \xi \in \mathbb{R}^3, \quad i = 1, \dots, N,$$

where  $t \rightarrow h_i(t, \xi)$  is continuous and

$$|h_i(t)|_1 \leq C, \quad \forall t \geq 0, \quad i = 1, \dots, N.$$

**Remark 3.3.** The linear multiplicative case  $B_i(X) := \alpha_i X$ ,  $i = 1, \dots, N$ , that is  $h_i := \delta$ , where  $\delta$  is the Dirac measure, can be approximated by taking  $h_i(\xi) = \frac{1}{\varepsilon^d} \rho\left(\frac{\xi}{\varepsilon}\right)$ , where  $\rho \in C_0^\infty(\mathbb{R}^d)$ , support  $\rho \subset \{\xi; |\xi|_d \leq 1\}$ ,  $\int \rho(\xi) d\xi = 1$ .

**Acknowledgements.** Financial support through SFB701 at Bielefeld University is gratefully acknowledged. V. Barbu was also partially supported by a CNCS UEFISCDI (Romania) grant, project PN-II-ID-PCE-2012-4-0156.

## References

- [1] V. Barbu, G. Da Prato, Internal stabilization by noise of the Navier-Stokes equation, *SIAM J. Control Optim.*, 49 (1) (2012), 1-20.
- [2] V. Barbu, M. Röckner, An operatorial approach to stochastic partial differential equations driven by linear multiplicative noise, *J. Eur. Math. Soc.*, 17 (2015), 1789-1815.
- [3] V. Barbu, M. Röckner, D. Zhang, Stochastic nonlinear Schrödinger equations with linear multiplicative noise: rescaling approach, *J. Nonlinear Sci.*, vol. 34 (2014), 383-409.



- [4] G.K. Batchelor, *Fluid Dynamics*, Cambridge University Press, New York, 1988.
- [5] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, New York, Dordrecht, Heidelberg, London, 2011.
- [6] Z. Brzezniak, E. Motyl, Existence and martingale solution of the stochastic Navier-Stokes equations in unbounded  $2D$  and  $3D$  domains, *J. Differential Equations*, 254 (2013), 1627-1642.
- [7] A.P. Calderon, A. Zygmund, On the existence of certain singular integrals, *Acta Mathematica*, 88 (1) (1952), 85-139.
- [8] A.J. Chorin, J.E. Marsden, *A Mathematical Introduction to Fluid Mechanics*, Springer-Verlag, 1975.
- [9] G. Da Prato, J.Zabczyk, *Stochastic Equations in Infinite Dimensions*, Second Edition, Cambridge University Press, Cambridge, 2014.
- [10] A. Debussche, C. Odasso, Markov solutions for  $3D$  stochastic Navier-Stokes equations, with state dependent noise, *J. Evol. Equ.*, 6 (2006), 305-324.
- [11] F. Flandoli, D. Gatarek, Martingale and stationary solutions for stochastic Navier-Stokes equations, *Probab. Theory Related Fields*, 102 (1995), 367-391.
- [12] F. Flandoli, M. Romito, Markov solutions for the  $3D$  stochastic Navier-Stokes equations, *Probab. Theory Related Fields*, 140 (2008), 407-458.
- [13] B. Goldys, M. Röckner, X. Zhang, martingale solutions and Markov selections for stochastic partial differential equations, *Stochastic Process. Appl.*, 119 (2009), 1725-1764.
- [14] Y. Giga, T. Miyakawa, Navier-Stokes flow in  $\mathbb{R}^3$  with measures as initial vorticity and Morrey spaces, *Comm. Partial Differential Equations*, 14 (1988), 577-618.
- [15] T. Kato, Strong  $L^p$ -solutions of the Navier-Stokes equation in  $\mathbb{R}^m$  with applications to weak solutions, *Math. Z.*, 187 (1984), 471-480.

- [16] T. Kato, H. Fujita, On the nonstationary Navier–Stokes system, *Rend. Sem. mat. Univ. Padova*, 32 (1962), 243-260.
- [17] H. Koch, D. Tataru, Well-posedness for the Navier–Stokes equations, *Advances in Mathematics*, 157 (1) (2001), 22–35.
- [18] R. Mikulevicius, B. Rozovskii, Global  $L_2$ -solutions of stochastic Navier-Stokes equations, *Ann. Probab.*, 33 (1) (2005), 137-176.
- [19] R. Mikulevicius, B. Rozovskii, Stochastic Navier-Stokes equations for turbulent flows, *SIAM J. Math. Anal.*, 35 (2004), 1250-1320.
- [20] E.M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [21] M. Taylor, Analysis on Morrey spaces and applications to Navier–Stokes equation, *Comm. PDE*, 17 (1992), 1407–1456.