

Predictive Inference Based on Markov Chain Monte Carlo Output

Fabian Krüger^a Sebastian Lerch^{a,b} Thordis Thorarinsdottir^c
 Tilmann Gneiting^{b,a}

^aKarlsruhe Institute of Technology

^bHeidelberg Institute for Theoretical Studies

^cNorwegian Computing Center

May 8, 2022

Abstract

In Bayesian inference, predictive distributions are typically in the form of samples generated via Markov chain Monte Carlo (MCMC) or related algorithms. In this paper, we conduct a systematic analysis of how to make and evaluate probabilistic forecasts from such simulation output. Based on proper scoring rules, we develop a notion of consistency that allows to assess the adequacy of methods for estimating the stationary distribution underlying the simulation output. We then provide asymptotic results that account for the salient features of Bayesian posterior simulators, and derive conditions under which choices from the literature satisfy our notion of consistency. Importantly, these conditions depend on the scoring rule being used, such that the choices of approximation method and scoring rule are intertwined. While the logarithmic rule requires fairly stringent conditions, the continuous ranked probability score (CRPS) yields consistent approximations under minimal assumptions. These results are illustrated in a simulation study and an economic data example. Overall, mixture-of-parameters approximations which exploit the parametric structure of Bayesian models perform particularly well. Under the CRPS, the empirical distribution function is a simple and appealing alternative option.

1 Introduction

Probabilistic forecasts are predictive probability distributions over quantities or events of interest. They implement an idea that was eloquently expressed already at the beginning of the 20th century in the context of meteorological prediction:

“It seems to me that the condition of confidence or otherwise forms a very important part of the prediction, and ought to find expression.”

(Cooke, 1906, pp. 23–24)

Despite this early acknowledgment of the importance of forecast uncertainty, constructing principled and realistic measures of the latter remains challenging in practice. In this

context, a rapidly growing transdisciplinary literature uses Bayesian inference to produce posterior predictive distributions in a wide range of applications, including economic, ecological, and meteorological problems, among many others. Bayesian posterior predictive distributions naturally account for sources of uncertainty – such as unknown model parameters, or latent variables in state space models – that are not easily captured using frequentist methods; see, e.g., Clark (2005) for an ecological perspective.

Formally, posterior predictive distributions arise as mixture distributions with respect to the posterior distribution of the parameter vector. In the following, we assume that the parameter vector contains all quantities that are subject to Bayesian inference, including also latent state variables, for example. For a real-valued continuous quantity of interest, the posterior predictive distribution, F_0 , can be represented by its cumulative distribution function (CDF) or the respective density. The posterior predictive CDF is then of the generic form

$$F_0(x) = \int_{\Theta} F_c(x|\theta) dP_{\text{post}}(\theta) \quad (1)$$

for $x \in \mathbb{R}$, where P_{post} is the posterior distribution of the parameter, θ , over some parameter space, Θ , and $F_c(\cdot|\theta)$ is the conditional predictive CDF when $\theta \in \Theta$ is the true parameter. Harris (1989) argues that predictive distributions of this form have appeal in frequentist settings as well. Often, the integral in (1) does not admit a solution in closed form, and so the posterior predictive CDF must be approximated or estimated in some way, typically using some form of Markov chain Monte Carlo (MCMC); see, e.g., Gelfand and Smith (1990) and Gilks et al. (1996).

Given a simulated sequence $(\theta_i)_{i=1}^m$ of parameter values from P_{post} , one approach, which we call the *mixture-of-parameters* (MP) technique, is to approximate F_0 by

$$\hat{F}_m^{\text{MP}}(x) = \frac{1}{m} \sum_{i=1}^m F_c(x|\theta_i). \quad (2)$$

However, this method can be used only when the conditional distributions $F_c(\cdot|\theta)$ are available in closed form. An alternative route is to simulate a sequence $(X_i)_{i=1}^m$ where $X_i \sim F_c(\cdot|\theta_i)$, and to approximate F_0 based on this sample, using either nonparametric or parametric techniques. The most straightforward option is to estimate F_0 by the *empirical CDF* (ECDF),

$$\hat{F}_m^{\text{ECDF}}(x) = \frac{1}{m} \sum_{i=1}^m \mathbb{1}\{x \geq X_i\}. \quad (3)$$

Alternatively, one might employ a *kernel density* (KD) estimate of the posterior predictive density, namely,

$$\hat{f}_m^{\text{KD}}(x) = \frac{1}{mh_m} \sum_{i=1}^m K\left(\frac{x - X_i}{h_m}\right), \quad (4)$$

where K is a kernel function, i.e., a symmetric, bounded, and square-integrable probability density, such as the Gaussian or the Epanechnikov kernel, and h_m is a suitable bandwidth (Rosenblatt, 1956; Silverman, 1986). Finally, much extant work employs a *Gaussian approximation* (GA) to F_0 , namely

$$\hat{F}_m^{\text{GA}}(x) = \Phi\left(\frac{x - \hat{\mu}_m}{\hat{\sigma}_m}\right), \quad (5)$$

where Φ is the CDF of the standard normal distribution, and $\hat{\mu}_m$ and $\hat{\sigma}_m$ are the empirical mean and standard deviation of the sample $(X_i)_{i=1}^m$.

Following Rubin (1984) and Little (2006), it is now widely accepted that posterior predictive inference should be evaluated using frequentist principles, without prior information entering at the model evaluation stage. For the comparison and ranking of probabilistic forecasting methods one typically uses a proper scoring rule (Gneiting and Raftery, 2007) that assigns a numerical score or penalty based on the predictive CDF, F , or its density, f , and the corresponding realization, y , such as the logarithmic score (LogS; Good, 1952),

$$\text{LogS}(F, y) = -\log f(y), \quad (6)$$

or the continuous ranked probability score (CRPS; Matheson and Winkler, 1976),

$$\text{CRPS}(F, y) = \int_{\mathbb{R}} (F(z) - \mathbb{1}\{z \geq y\})^2 dz. \quad (7)$$

While the LogS and CRPS are the two most popular scoring rules in applications, they feature interesting conceptual differences which we discuss in Section 2.2. In practice, one finds and compares the mean score over an out-of-sample test set, and the forecasting method with the smaller mean score is preferred. Formal tests of the null hypothesis of equal predictive performance can be employed as well (Diebold and Mariano, 1995; Giacomini and White, 2006; Clark and McCracken, 2013; DelSole and Tippett, 2014).

Table 1 of the Online Supplement summarizes the use of evaluation techniques in recently published comparative studies of probabilistic forecasting methods that use Bayesian inference via MCMC. As shown in the table, the mixture-of-parameters technique has mainly been applied in concert with the logarithmic score, whereas the empirical CDF method can be used in conjunction with the CRPS only. However, to this date, there are few, if any, guidelines to support choices in Table 1 of the Online Supplement, and it is not clear how they affect practical model comparisons. The present paper provides a systematic analysis of this topic. We focus on the following questions. First, what defines reasonable choices of the approximation method and scoring rule? Second, under what conditions do extant choices from the literature satisfy this definition? Third, for a given scoring rule, how accurate are alternative approximation methods in practically relevant scenarios?

We emphasize that the present study — and the use of scoring rules in general — concern the *comparative* assessment of two or more predictive models: The model with the smallest mean score is considered the most appropriate. Comparative assessment is essential in order to choose among a large number of specifications typically available in practice. This task is different from *absolute* assessment, which amounts to diagnosing possible misspecification, using the probability integral transform (Dawid, 1984; Diebold et al., 1998), posterior predictive checks (Gelman et al., 1996; Held et al., 2010; Gelman et al., 2014a, Chapter 6) and related methods.

The remainder of this paper is organized as follows. Section 2 introduces the notion of a consistent approximation to F_0 . This formalizes the idea that, as the size of the simulated sample becomes larger and larger, and in terms of a given scoring rule, the approximation ought to perform as well as the unknown true forecast distribution. In Section 3 we provide theoretical justifications of approximation methods encountered in the literature. Sections 4 and 5 present simulation and empirical evidence on the performance of these methods, and Section 6 concludes with a discussion. Overall, our findings support the use of the mixture-of-parameters estimator at (2) in order to approximate the posterior predictive distribution of interest. If this estimator is unavailable, the ECDF estimator at (3) is a simple and appealing alternative. Technical material and supplementary analyses are

deferred to Appendices A to E. The Online Supplement contains a bibliography of the pertinent applied literature and additional figures.

2 Formal Setting

In this section, we discuss the posterior predictive distribution in Bayesian forecasting, give a brief review of proper scoring rules and score divergences, and introduce the concept of a consistent approximation method based on MCMC output.

As discussed earlier, the posterior predictive cumulative distribution function (CDF) of a Bayesian forecasting model is given by

$$F_0(x) = \int_{\Theta} F_c(x|\theta) dP_{\text{post}}(\theta)$$

where $\theta \in \Theta$ is the parameter, P_{post} is the posterior distribution of the parameter, and $F_c(\cdot|\theta)$ is the predictive distribution *conditional* on a parameter value θ ; see, e.g., Greenberg (2013, p. 33) or Gelman et al. (2014a, p. 7). A generic Markov chain Monte Carlo (MCMC) algorithm designed to sample from F_0 can be sketched as follows.

- Fix $\theta_0 \in \Theta$ at some arbitrary value.
- For $i = 1, 2, \dots$ iterate as follows:
 - Draw $\theta_i \sim \mathcal{K}(\theta_i|\theta_{i-1})$, where \mathcal{K} is a transition kernel that specifies the conditional distribution of θ_i given θ_{i-1} .
 - Draw $X_i \sim F_c(\cdot|\theta_i)$.

We assume throughout that the transition kernel \mathcal{K} is such that the sequence $(\theta_i)_{i=1,2,\dots}$ is stationary and ergodic in the sense of Geweke (2005, Definition 4.5.5) with invariant distribution P_{post} , as holds widely in practice (Craiu and Rosenthal, 2014). Importantly, stationarity and ergodicity of $(\theta_i)_{i=1,2,\dots}$ with invariant distribution P_{post} imply that $(X_i)_{i=1,2,\dots}$ is stationary and ergodic with invariant distribution F_0 (Genon-Catalot et al., 2000, Proposition 3.1).

This generic MCMC algorithm allows for two general options for estimating the posterior predictive distribution F_0 in (1), namely,

- Option A: Based on parameter draws $(\theta_i)_{i=1}^m$,
- Option B: Based on a sample $(X_i)_{i=1}^m$,

where m typically is on the order of a few thousands or ten thousands. Alternatively, some authors, such as Krüger et al. (2017), generate, for each $i = 1, \dots, m$, independent draws $X_{ij} \sim F_c(\cdot|\theta_i)$, where $j = 1, \dots, J$; see also Waggoner and Zha (1999, Section III. B). The considerations below apply in this more general setting as well.

2.1 Approximation methods

In the case of Option A, the sequence $(\theta_i)_{i=1}^m$ of parameter draws is used to approximate the posterior predictive distribution, F_0 , by the mixture-of-parameters estimator \hat{F}_m^{MP} in (2). Under the assumption of ergodicity,

$$\hat{F}_m^{\text{MP}}(x) = \frac{1}{m} \sum_{i=1}^m F_c(x|\theta_i) \longrightarrow \int_{\Theta} F_c(x|\theta) dP_{\text{post}}(\theta) = F_0(x)$$

for $x \in \mathbb{R}$. This estimator was popularized by Gelfand and Smith (1990, Section 2.2), based on earlier work by Tanner and Wong (1987), and is often called a *conditional* or *Rao-Blackwellized* estimator. The latter term hints at variance reduction that may result from conditioning on the parameter draws (see Theorem 4 below). We refer to \hat{F}_m^{MP} as the *mixture-of-parameters* (MP) estimator.

In the case of Option B, the sample $(X_i)_{i=1}^m$ is employed to approximate the posterior predictive distribution F_0 . Various methods for doing this have been proposed and used, including the *empirical CDF* of the sample, denoted \hat{F}_m^{ECDF} in (3), the *kernel density* estimator \hat{f}_m^{KD} in (4), and the *Gaussian* approximation \hat{F}_m^{GA} in (5). Approaches of this type incur ‘more randomness than necessary’, in that the simulation step to draw $(X_i)_{i=1}^m$ can be avoided if Option A is used. That said, Option A requires full knowledge of the model specification, as the conditional distributions must be known in closed form in order to compute \hat{F}_m^{MP} . There are situations where this is restrictive, e.g., when the task is to predict a complicated transformation of the original, possibly vector-valued predictand. The implementation of the approximation methods is straightforward, except for the case of kernel density estimation, for which we discuss implementation choices in Section 3.3.

2.2 Proper scoring rules and score divergences

Let $\Omega \subseteq \mathbb{R}$ denote the set of possible values of the quantity of interest, and let \mathcal{F} denote a convex class of probability distributions on Ω . A *scoring rule* is a function

$$S : \mathcal{F} \times \Omega \longrightarrow \mathbb{R} \cup \{\infty\}$$

that assigns numerical values to pairs of forecasts $F \in \mathcal{F}$ and observations $y \in \Omega$. We typically set $\Omega = \mathbb{R}$, but will occasionally restrict attention to compact subsets.

Throughout this paper, we define scoring rules to be negatively oriented, i.e., a lower score indicates a better forecast. A scoring rule is *proper* relative to \mathcal{F} if the expected score

$$S(F, G) = \mathbb{E}_{Y \sim G} S(F, Y)$$

is minimized for $F = G$, i.e.,

$$\mathbb{E}_{Y \sim G} S(G, Y) = S(G, G) \leq S(F, G) = \mathbb{E}_{Y \sim G} S(F, Y)$$

for all probability distributions $F, G \in \mathcal{F}$. It is *strictly proper* relative to the class \mathcal{F} if, furthermore, equality implies that $F = G$. The *score divergence* associated with the scoring rule S is given by

$$d_S(F, G) = S(F, G) - S(G, G).$$

Clearly, $d_S(F, G) \geq 0$ for all $F, G \in \mathcal{F}$ is equivalent to propriety of the scoring rule S , which is a critically important property in practice.¹

¹See Brier (1950) and Shuford et al. (1966) for early references arguing that scoring rules should be proper, and Gneiting and Raftery (2007) for a review of the statistical implications.

Table 1: Examples of proper scoring rules, along with the associated score divergence and natural domain, \mathcal{F} . For a probability distribution with CDF F , we write μ_F for its mean, σ_F for its standard deviation, and f for its density.

Scoring rule	$S(F, y)$	$d_S(F, G)$	\mathcal{F}
Logarithmic score	$-\log f(y)$	$\int g(z) \log \frac{g(z)}{f(z)} dz$	\mathcal{L}_1
CRPS	$\int (F(z) - \mathbb{1}\{z \geq y\})^2 dz$	$\int (F(z) - G(z))^2 dz$	\mathcal{M}_1
Dawid–Sebastiani score	$\log \sigma_F^2 + \frac{(y - \mu_F)^2}{\sigma_F^2}$	$\frac{\sigma_G^2}{\sigma_F^2} - \log \frac{\sigma_G^2}{\sigma_F^2} + \frac{(\mu_F - \mu_G)^2}{\sigma_F^2} - 1$	\mathcal{M}_2

Table 1 shows frequently used proper scoring rules, along with the associated score divergences and the natural domain. For any given scoring rule S , the associated *natural domain* is the largest convex class of probability distributions F such that $S(F, y)$ is well-defined and finite almost surely under F . Specifically, the natural domain for the popular *logarithmic score* (LogS; eq. (6)) is the class \mathcal{L}_1 of the probability distribution with densities, and the respective score divergence is the Kullback-Leibler divergence. The logarithmic score is local (Bernardo, 1979), that is, it evaluates a predictive model based only on the density value at the realizing outcome. Conceptually, this means that the logarithmic score ignores the model’s predicted probabilities of events that could have happened but did not. For the *continuous ranked probability score* (CRPS; eq. (7)), the natural domain is the class \mathcal{M}_1 of the probability distributions with finite mean. The LogS and CRPS are both strictly proper relative to their respective natural domains. In contrast to the logarithmic score, the CRPS rewards predictive distributions that place mass close to the realizing outcome, a feature that is often called ‘sensitivity to distance’ (e.g. Matheson and Winkler, 1976, Section 2). While various authors have argued in favor of either locality or sensitivity to distance, the choice between these two contrasting features appears ultimately subjective. Finally, the natural domain for the *Dawid–Sebastiani score* (DSS; Dawid and Sebastiani, 1999) is the class \mathcal{M}_2 of the probability distributions with strictly positive, finite variance. This score is proper, but not strictly proper, relative to \mathcal{M}_2 .

2.3 Consistent approximations

To study the combined effects of the choices of approximation method and scoring rule in the evaluation of Bayesian predictive distributions, we introduce the notion of a *consistent* approximation procedure.

Specifically, let $(\theta_i)_{i=1,2,\dots}$ or $(X_i)_{i=1,2,\dots}$, where $X_i \sim F_c(\cdot|\theta_i)$, be output from a generic MCMC algorithm with the following property.

- (A) The process $(\theta_i)_{i=1,2,\dots}$ is stationary and ergodic with invariant distribution P_{post} .

As noted, assumption (A) implies that $(X_i)_{i=1,2,\dots}$ is stationary and ergodic with invariant distribution F_0 . Consider an approximation method that produces, for all sufficiently large positive integers m , an estimate \hat{F}_m that is based on $(\theta_i)_{i=1}^m$ or $(X_i)_{i=1}^m$, respectively. Let S be a proper scoring rule, and let \mathcal{F} be the associated natural domain. Then the approximation method is *consistent relative to the scoring rule S at the distribution $F_0 \in \mathcal{F}$* if $\hat{F}_m \in \mathcal{F}$ for all sufficiently large m , and

$$d_S(\hat{F}_m, F_0) \longrightarrow 0$$

or, equivalently, $S(\hat{F}_m, F_0) \rightarrow S(F_0, F_0)$ almost surely as $m \rightarrow \infty$. This formalizes the idea that under continued MCMC sampling, the approximation ought to perform as well as the unknown true posterior predictive distribution. We contend that this is a highly desirable property in practical work.

Note that \hat{F}_m is a random quantity that depends on the sample $(\theta_i)_{i=1}^m$ or $(X_i)_{i=1}^m$. The specific form of the divergence stems from the scoring rule, which mandates convergence of a certain functional of the estimator or approximation, \hat{F}_m , and the theoretical posterior predictive distribution, F_0 . As we will argue, this aspect has important implications for the choice of scoring rule and approximation method.

Our concept of a consistent approximation procedure is independent of the question of how well a forecast model represents the ‘true’ uncertainty. The definition thus allows to separate the problem of interest, namely, to find a good approximation \hat{F}_m to F_0 , from the distinct task of finding a good probabilistic forecast F_0 .² We further emphasize that we study convergence in the sample size, m , of MCMC output, given a fixed number of observations, say, T , used to fit the model. Our analysis is thus distinct from traditional Bayesian asymptotic analyses that study convergence of the posterior distribution as T becomes larger and larger (see, e.g., Gelman et al., 2014a, Section 4), thereby calling for markedly different technical tools.

2.4 Relation to total variation and Wasserstein distances

Our focus on score divergences (in particular, on d_{LogS} and d_{CRPS}) is motivated by their natural relation to scoring rules, which in turn are popular tools in the applied literature on probabilistic forecasting. As reviewed by Gibbs and Su (2002), many other distance metrics for comparing two probability distributions have been proposed in the literature. Among these metrics, the total variation distance (d_{TV}) has received much attention in theoretical work on MCMC (e.g. Tierney, 1994; Rosenthal, 1995), and is thus particularly relevant in our context. The total variation distance between two absolutely continuous probability measures with densities f and g is defined as

$$d_{\text{TV}}(F, G) = \frac{1}{2} \int_{-\infty}^{\infty} |f(z) - g(z)| \, dz.$$

As $2d_{\text{TV}}(F, G)^2 \leq d_{\text{LogS}}(F, G)$ (e.g., Barron et al., 1992), convergence in terms of d_{LogS} implies convergence in terms of d_{TV} .

The Wasserstein distance is a divergence function motivated by optimal transport problems (Villani, 2008) and has received much attention in statistics and machine learning (Panaretos and Zemel, 2019). Here, we limit our discussion to the Wasserstein distance of order 1, which is most common in practice, and denote the corresponding metric by

$$d_{\text{W}}(F, G) = \int_0^1 |F^{-1}(\alpha) - G^{-1}(\alpha)| \, d\alpha = \int_{-\infty}^{\infty} |F(z) - G(z)| \, dz,$$

where F^{-1} and G^{-1} are the quantile functions of F and G respectively. Bellemare et al. (2017) discuss shortcomings of Wasserstein distances in estimation with stochastic gradient descent methods and suggest d_{CRPS} as a superior alternative. This recommendation

²It is possible for an inconsistent approximation to a *misspecified* posterior predictive distribution F_0 to yield better forecasts than a consistent approximation that approaches the misguided F_0 . However, the misspecification can be detected by diagnostic tools such as probability integral transform histograms; see Dawid (1984) and Diebold et al. (1998). The appropriate remedy thus is to improve the model specification. Once a well-specified model has been found, the use of a consistent approximation improves the predictive performance further.

Table 2: Upper bounds on the computational complexity of approximation methods in terms of the size m of the MCMC sample $(\theta_i)_{i=1}^m$ or $(X_i)_{i=1}^m$, respectively, for pre-processing, and for the exact computation of the CRPS, Dawid–Sebastiani score (DSS) and logarithmic score (LogS).

Approximation method	Pre-processing	CRPS	DSS	LogS
MP	$\mathcal{O}(1)$	$\mathcal{O}(m^2)$	$\mathcal{O}(m^2)$	$\mathcal{O}(m)$
ECDF	$\mathcal{O}(1)$	$\mathcal{O}(m \log m)$	$\mathcal{O}(m)$	
KD	$\mathcal{O}(m)$	$\mathcal{O}(m^2)$	$\mathcal{O}(m)$	$\mathcal{O}(m)$
Gaussian	$\mathcal{O}(m)$	$\mathcal{O}(1)$	$\mathcal{O}(1)$	$\mathcal{O}(1)$

relates to the observation that there is no proper scoring rule with d_W as score divergence (Thorarinsdottir et al., 2013, Theorem 2).

As $d_{\text{CRPS}}(F, G) \leq d_W(F, G)$, convergence in terms of d_W implies convergence in terms of d_{CRPS} . If F and G have densities with support in a common interval of length l , $d_W(F, G) \leq l \cdot d_{\text{TV}}(F, G) \leq l \cdot \sqrt{d_{\text{LogS}}(F, G)/2}$, so in this case consistency relative to the logarithmic score implies consistency relative to the CRPS. For further relations to the Kolmogorov, Lévy, Prohorov and bounded Lipschitz distances see Section 2.4 of Huber and Ronchetti (2009).

3 Consistency results and computational complexity

We now investigate sufficient conditions for consistency of the aforementioned approximation methods, namely, the mixture-of-parameters (MP) estimator \hat{F}_m^{MP} in (2), the empirical CDF (ECDF) method \hat{F}_m^{ECDF} in (3), the kernel density (KD) estimate \hat{f}_m^{KD} in (4), and the Gaussian approximation (GA) \hat{F}_m^{GA} in (5). Table 2 summarizes upper bounds on the computational cost of pre-processing and computing the CRPS, Dawid–Sebastiani score (DSS) and logarithmic score (LogS) under these methods in terms of the size m of the MCMC sample $(\theta_i)_{i=1}^m$ or $(X_i)_{i=1}^m$, respectively.

Consistency requires the convergence of some functional of the approximation, \hat{F}_m , and the true posterior predictive distribution, F_0 . The conditions to be placed on the Bayesian model F_0 , the estimator \hat{F}_m , and the dependence structure of the MCMC output depend on the scoring rule at hand.

3.1 Mixture-of-parameters estimator

We now establish consistency of the mixture-of-parameters estimator \hat{F}_m^{MP} in (2) relative to the CRPS, DSS and logarithmic score. The proofs are deferred to Appendix B.

Theorem 1 (consistency of mixture-of-parameters approximations relative to the CRPS and DSS). *Under assumption (A), the mixture-of-parameters approximation is consistent relative to the CRPS at every distribution F_0 with finite mean, and consistent relative to the DSS at every distribution F_0 with strictly positive, finite variance.*

Theorem 1 is the best possible result of its kind: It applies to every distribution in the natural domain and does not invoke any assumptions on the Bayesian model. In contrast,

Theorem 2 hinges on rather stringent further conditions on the distribution F_0 and the Bayesian model (1), as follows.

- (B) The distribution F_0 is supported on some bounded interval Ω . It admits a density, f_0 , that is continuous and strictly positive on Ω . Furthermore, the density $f_c(\cdot|\theta)$ is continuous for every $\theta \in \Theta$.

Theorem 2 (consistency of mixture-of-parameters approximations relative to the logarithmic score). *Under assumptions (A) and (B), the mixture-of-parameters approximation is consistent relative to the logarithmic score at the distribution F_0 .*

While we believe that the mixture-of-parameters technique is consistent under weaker assumptions, this is the strongest result that we have been able to prove. In particular, condition (B) does not allow for the case $\Omega = \mathbb{R}$. However, practical applications often involve a truncation of the support for numerical reasons, as exemplified in Section 4, and in this sense the assumption may not be overly restrictive.

Computing the logarithmic score and the DSS for a predictive distribution \hat{F}_m^{MP} of the form (2) is straightforward. To compute the CRPS, we note from eq. (21) of Gneiting and Raftery (2007) that

$$\text{CRPS}(\hat{F}_m^{\text{MP}}, y) = \frac{1}{m} \sum_{i=1}^m \mathbb{E}|Z_i - y| - \frac{1}{2m^2} \sum_{i=1}^m \sum_{j=1}^m \mathbb{E}|Z_i - Z_j|, \quad (8)$$

where Z_i and Z_j are independent random variables with distribution $F_c(\cdot|\theta_i)$ and $F_c(\cdot|\theta_j)$, respectively. The expectations on the right-hand side of (8) often admit closed form expressions that can be derived with techniques described by Jordan (2016) and Taillardat et al. (2016), including but not limited to the ubiquitous case of Gaussian variables. Then the evaluation requires $\mathcal{O}(m^2)$ operations, as reported in Table 2. In Appendix A, we provide details and investigate the use of numerical integration in (7), which provides an attractive, computationally efficient alternative.

3.2 Empirical CDF-based approximation

The empirical CDF-based approximation \hat{F}_m^{ECDF} in (3), which builds on a simulated sample $(X_i)_{i=1}^m$, is consistent relative to the CRPS and DSS under minimal assumptions. We prove the following result in Appendix C, which is the best possible of its kind, as it applies to every distribution in the natural domain and does not invoke any assumptions on the Bayesian model.

Theorem 3 (consistency of empirical CDF-based approximations relative to the CRPS and DSS). *Under assumption (A), the empirical CDF technique is consistent relative to the CRPS at every distribution F_0 with finite mean, and consistent relative to the DSS at every distribution F_0 with strictly positive, finite variance.*

As stated in Table 2, the computation of the CRPS under \hat{F}_m^{ECDF} requires $\mathcal{O}(m \log m)$ operations only. Specifically, let $X_{(1)} \leq \dots \leq X_{(m)}$ denote the order statistics of X_1, \dots, X_m , which can be obtained in $\mathcal{O}(m \log m)$ operations. Then

$$\text{CRPS}(\hat{F}_m^{\text{ECDF}}, y) = \frac{2}{m^2} \sum_{i=1}^m (X_{(i)} - y) \left(m \mathbb{1}\{y < X_{(i)}\} - i + \frac{1}{2} \right); \quad (9)$$

see Jordan (2016, Section 6) for details. A special case of eq. (8) suggests another way of computing the CRPS, in that

$$\text{CRPS}(\hat{F}_m^{\text{ECDF}}, y) = \frac{1}{m} \sum_{i=1}^m |X_i - y| - \frac{1}{2m^2} \sum_{i=1}^m \sum_{j=1}^m |X_i - X_j|. \quad (10)$$

The representations in (9) and (10) are algebraically equivalent, but the latter requires $\mathcal{O}(m^2)$ operations and thus is inefficient.

While the consistency results support the use of both \hat{F}_m^{MP} and \hat{F}_m^{ECDF} , Rao-Blackwellization arguments indicate superiority of \hat{F}_m^{MP} .

Theorem 4 (comparison of \hat{F}_m^{MP} and \hat{F}_m^{ECDF}). *Under assumption (A), $\mathbb{E} \hat{F}_m^{\text{MP}}(z) = \mathbb{E} \hat{F}_m^{\text{ECDF}}(z)$ and $\text{Var} \hat{F}_m^{\text{MP}}(z) \leq \text{Var} \hat{F}_m^{\text{ECDF}}(z)$ for any $z \in \Omega$ and $m \in \mathbb{N}$. If furthermore F_0 has finite mean, then $\mathbb{E} d_{\text{CRPS}}(\hat{F}_m^{\text{MP}}, F_0) \leq \mathbb{E} d_{\text{CRPS}}(\hat{F}_m^{\text{ECDF}}, F_0)$ for any $m \in \mathbb{N}$.*

Theorem 4 demonstrates that \hat{F}_m^{MP} outperforms \hat{F}_m^{ECDF} in terms of expected divergence, for every given sample size m . Proposition 5.1 of Bolin and Wallin (2019) shows that if F_0 is a normal location-scale mixture then the CRPS under the mixture-of-parameters estimator additionally has smaller variance than under the empirical CDF-based approximation.

Despite the theoretical superiority of \hat{F}_m^{MP} , \hat{F}_m^{ECDF} may be attractive in practice, especially if the conditional distributions $F_c(\cdot|\theta)$ underlying \hat{F}_m^{MP} are difficult to compute analytically. For example, this may occur if the predictand Y is modeled only indirectly (such as when Y is the maximal element of a vector valued random variable).

3.3 Kernel density estimator

We now discuss conditions for the consistency of the kernel density estimator \hat{f}_m^{KD} . In the present case of dependent samples $(X_i)_{i=1}^m$, judicious choices of the bandwidth h_m in (4) require knowledge of dependence properties of the sample, and the respective conditions are difficult to verify in practice.

The score divergence associated with the logarithmic score is the Kullback-Leibler divergence, which is highly sensitive to tail behavior. Therefore, consistency of \hat{f}_m^{KD} requires that the tail properties of the kernel K in (4) and the true posterior predictive density f_0 be carefully matched, and any results tend to be technical (cf. Hall (1987)). Roussas (1988), Györfi et al. (1989), Yu (1993) and Liebscher (1996), among others, establish almost sure strong uniform consistency of \hat{f}_m^{KD} under α - or β -mixing and other conditions. As noted in Appendix B, almost sure strong uniform consistency then implies consistency relative to the logarithmic score under assumption (B). Based on Hansen (2008) who proves general results we give conditions for consistency of the kernel density estimator \hat{f}_m^{KD} , and summarize the relevant assumptions in the following condition.

- (H) For the kernel function K , the bandwidth h_m , and the dependence properties of $(X_i)_{i=1,2,\dots}$ assumptions 1–3 and the conditions of Theorem 7 of Hansen (2008) are satisfied.

Theorem 5 (consistency of kernel density estimator-based approximations relative to the LogS). *Under assumptions (A), (B), and (H), the kernel density estimator-based approximation technique is consistent relative to the logarithmic score at the distribution F_0 .*

The result is a direct consequence of Hansen (2008, Theorem 7) who further provides optimal convergence rates. However, the respective conditions are stringent and difficult to check in practice. Indeed, Wasserman (2006, p. 57) opines that “Despite the natural role of Kullback-Leibler distance in parametric statistics, it is usually not an appropriate loss function in smoothing problems”.

Under the conditions of Theorem 5, consistency of \hat{F}_m^{KD} relative to the CRPS follows directly, see Section 2.4. Kernel density estimation approximations are generally not consistent relative to the DSS due to the variance inflation induced by typical choices of the bandwidth. However, adaptations based on rescaling or weighting allow for kernel density estimation under moment constraints, see, e.g., Jones (1991) and Hall and Presnell (1999).

As this brief review suggests, the theoretical properties of kernel density estimators depend on the specifics of both the MCMC sample and the estimator. However, under the CRPS and DSS, a natural alternative is readily available: The empirical CDF approach is simpler and computationally cheaper than kernel density estimation, and is consistent under weak assumptions (Theorem 3).

In our simulation and data examples, we use a simple implementation of kernel density estimator-based approximations based on the Gaussian kernel and the Silverman (1986, Section 3.4) plug-in rule for bandwidth selection. This leads to the specific form

$$\hat{F}_m^{\text{KD}}(x) = \frac{1}{m} \sum_{i=1}^m \Phi\left(\frac{x - X_i}{h_m}\right), \quad (11)$$

where Φ denotes the CDF of the standard normal distribution, and

$$h_m = \left(\frac{4}{3} \frac{\hat{\sigma}_m^5}{m}\right)^{1/5} \approx 1.06 \hat{\sigma}_m m^{-1/5}, \quad (12)$$

with $\hat{\sigma}_m$ being the standard deviation of $(X_i)_{i=1}^m$, for pre-processing costs of $\mathcal{O}(m)$, as shown in Table 2. This choice is motivated by simulation evidence in Hall et al. (1995). Using the Sheather and Jones (1991) rule or cross-validation based methods yields slightly inferior results in our experience.³

3.4 Gaussian approximation

A parametric approximation method fits a member of a fixed parametric family, say \mathcal{F}_Γ , of probability distributions to the MCMC sample $(X_i)_{i=1}^m$. The problem of estimating the unknown distribution F_0 is thus reduced to a finite-dimensional parameter estimation problem. The most important case is the *quadratic* or *Gaussian approximation*, which takes \mathcal{F}_Γ to be the Gaussian family, so that

$$\hat{F}_m^{\text{GA}}(x) = \Phi\left(\frac{x - \hat{\mu}_m}{\hat{\sigma}_m}\right),$$

where $\hat{\mu}_m$ and $\hat{\sigma}_m$ are the empirical mean and standard deviation of $(X_i)_{i=1}^m$. If F_0 has a density f_0 that is unimodal and symmetric, the approximation can be motivated by a Taylor series expansion of the log predictive density at the mode, similar to Gaussian

³Sköld and Roberts (2003) and Kim et al. (2016) discuss bandwidth selection rules that are motivated by density estimation in MCMC samples. However, both studies rely on mean integrated squared error criteria that are different from the performance measures we consider here.

approximations of posterior distributions in large-sample Bayesian inference (e.g. Kass and Raftery, 1995; Gelman et al., 2014a, Chapter 4).

If F_0 is not Gaussian, \hat{F}_m^{GA} fails to be consistent relative to the logarithmic score and CRPS. However, the Ergodic Theorem implies that the Gaussian approximation is consistent relative to the Dawid–Sebastiani score under minimal conditions.

Theorem 6 (consistency of Gaussian approximations relative to the DSS). *Under assumption (A), the Gaussian approximation technique is consistent relative to the DSS at every distribution F_0 with strictly positive, finite variance.*

We also note that the logarithmic score for the Gaussian approximation \hat{F}_m^{GA} corresponds to the Dawid–Sebastiani score for the empirical CDF-based approximation \hat{F}_m^{ECDF} , in that

$$\text{LogS}(\hat{F}_m^{\text{GA}}, y) = \frac{1}{2} \left(\log 2\pi + \text{DSS}(\hat{F}_m^{\text{ECDF}}, y) \right)$$

for $y \in \mathbb{R}$. Therefore, the Gaussian approximation under the logarithmic score yields model rankings that are identical to those for the empirical CDF technique under the Dawid–Sebastiani score. From an applied perspective, this equivalence suggests that the inconsistency of the Gaussian approximation may not be overly problematic when the approximation is used in concert with the logarithmic score, an assessment that is in line with empirical findings by Warne et al. (2016). However, researchers should be aware of the fact that they are effectively using a proper, but not strictly proper, scoring rule (namely, the Dawid–Sebastiani score) that focuses on the first two moments of the predictive distribution only.

4 Simulation study

We now investigate the various approximation methods in a simulation study that is designed to emulate realistic MCMC behavior with dependent samples. Here, the posterior predictive distribution F_0 is known by construction, and so we can compare the different approximations to the true forecast distribution.

In order to judge the quality of an approximation \hat{F}_m of F_0 we consider the score divergence $d_S(\hat{F}_m, F_0)$. Note that $d_S(\hat{F}_m, F_0)$ is a random variable, since \hat{F}_m depends on the particular MCMC sample $(\theta_i)_{i=1}^m$ or $(X_i)_{i=1}^m$. In our results below, we therefore consider the distribution of $d_S(\hat{F}_m, F_0)$ across repeated simulation runs. For a generic approximation method producing an estimate \hat{F}_m , we proceed as follows:

- For simulation run $k = 1, \dots, K$:
 - Draw MCMC samples $(\theta_i^{(k)})_{i=1}^m$ and $(X_i^{(k)})_{i=1}^m$
 - Compute the approximation $\hat{F}_m^{(k)}$ and the divergence $d_S(\hat{F}_m^{(k)}, F_0)$ for the approximation methods and scoring rules under consideration.
- For each approximation method and scoring rule, summarize the distribution of $d_S(\hat{F}_m^{(1)}, F_0), \dots, d_S(\hat{F}_m^{(K)}, F_0)$.

In order to simplify notation, we typically suppress the superscript that identifies the Monte Carlo iteration. The results below are based on $K = 1\,000$ replicates.

Table 3: Hyper-parameters for the data generating process in the simulation setting of equations (14) to (17).

Parameter	Main role	Value(s) considered
α	Persistence of θ_i^2	$\{0.1, 0.5, 0.9\}$
s	Unconditional mean of θ_i^2	2
n	Unconditional variance of θ_i^2	$\{12, 20\}$

4.1 Data generating process

We generate sequences $(\theta_i)_{i=1}^m$ and $(X_i)_{i=1}^m$ in such a way that the invariant distribution,

$$F_0(x) = \int_{(0,\infty)} \Phi\left(\frac{x}{\theta}\right) dH_0(\theta^2),$$

where Φ denotes the standard normal CDF, is a compound Gaussian distribution or normal scale mixture. Depending on the measure H_0 , which assumes the role of the posterior distribution P_{post} in the general Bayesian model (1), F_0 can be modeled flexibly, including many well-known parametric distributions (Gneiting, 1997). As detailed below, our choice of H_0 implies that

$$F_0(x) = \mathsf{T}\left(x \left| 0, \frac{ns}{n+2}, n+2 \right.\right) \quad (13)$$

where $\mathsf{T}(\cdot|a, b, c)$ denotes the CDF of a variable Z with the property that $(Z - a)/\sqrt{b}$ is standard Student t distributed with c degrees of freedom. To mimic a realistic MCMC scenario with dependent draws, we proceed as proposed by Fox and West (2011). Given parameter values $n > 0$, $s > 0$ and $\alpha \in (-1, 1)$, let

$$\psi_i \sim \text{IG}\left(\frac{1}{2}(n+3), \frac{1}{2}ns(1-\alpha^2)\right), \quad (14)$$

$$v_i | \psi_i \sim \mathcal{N}\left(\alpha, \frac{\psi_i}{ns}\right), \quad (15)$$

$$\theta_i^2 = \psi_i + v_i^2 \theta_{i-1}^2, \quad (16)$$

$$X_i | \theta_i^2 \sim \mathcal{N}(0, \theta_i^2) \quad (17)$$

where IG is the Inverse Gamma distribution, parametrized such that $Z \sim \text{IG}(a, b)$ when $1/Z \sim \text{G}(a, b)$, with G being the Gamma distribution with shape $a \geq 0$ and rate $b > 0$.

Table 3 summarizes our choices for the parameter configurations of the data generating process. The parameter α determines the persistence of the chain, in that the unconditional mean of v_i^2 , which can be viewed as an average autoregressive coefficient (Fox and West, 2011, Section 2.3), is given by $(n\alpha^2 + 1)/(n + 1)$. We consider three values, aiming to mimic MCMC chains with different persistence properties. The parameter s represents a scale effect, and n governs the tail thickness of the unconditional Student t distribution in (13). We consider values of 12 and 20 that seem realistic for macroeconomic variables, such as the growth rate of the gross domestic product, that feature prominently in the empirical literature.

4.2 Approximation methods

We consider the following approximation methods, which have been discussed in detail in Section 3. The first approximation uses a sequence $(\theta_i)_{i=1}^m$ of parameter draws, and the other three employ an MCMC sample $(X_i)_{i=1}^m$.

- Mixture-of-parameters estimator \hat{F}_m^{MP} in (2), which here is of the form

$$\hat{F}_m^{\text{MP}}(x) = \frac{1}{m} \sum_{i=1}^m \Phi\left(\frac{x}{\theta_i}\right),$$

where θ_i is the predictive standard deviation drawn in MCMC iteration i .

- Empirical CDF-based approximation \hat{F}_m^{ECDF} in (3).
- The nonparametric kernel density estimator \hat{f}_m^{KD} using a Gaussian kernel and the Silverman rule for bandwidth selection, with predictive CDF \hat{F}_m^{KD} of the form (11).
- Gaussian approximation \hat{F}_m^{GA} in (5).

It is interesting to observe that here \hat{F}_m^{MP} is a scale mixture of centered Gaussian distributions, and \hat{F}_m^{KD} is a location mixture of normal distributions, whereas the quadratic approximation \hat{F}_m^{GA} is a single Gaussian.

The conditions for consistency of the mixture-of-parameters and empirical CDF approximations relative to the CRPS in Theorems 1 and 3 are satisfied. Furthermore, one might argue that numerically the support of F_0 and \hat{F}_m^{MP} is bounded (cf. below), and then the assumptions of Theorem 2 also are satisfied. Clearly, the Gaussian approximation fails to be consistent relative to the CRPS or the logarithmic score, as F_0 is not Gaussian.

For each approximation method, scoring rule S , sample size m and replicate k , we evaluate the score divergence $d_S(\hat{F}_m^{(k)}, F_0)$. The divergence takes the form of a univariate integral (cf. Table 1) that is not available in closed form. Therefore, we compute $d_S(\hat{F}_m^{(k)}, F_0)$ by numerical integration as implemented in the R function `integrate`. This is unproblematic if the scoring rule is the CRPS. For the logarithmic score, the integration is numerically challenging, as the logarithm of the densities needs to be evaluated in their tails. We therefore truncate the support of the integral to the minimal and maximal values that yield numerically finite values of the integrand.

4.3 Main results

In the interest of brevity, we restrict attention to results for a single set of parameters of the data generating process, namely $(\alpha, s, n) = (0.5, 2, 12)$. This implies an unconditional Student t distribution with 14 degrees of freedom, and intermediate autocorrelation of the MCMC draws. The results for the other parameter constellations in Table 3 are similar and available in the Online Supplement.

Figure 1 illustrates the performance of the approximation methods under the logarithmic score and the CRPS, by showing the distribution of the score divergence $d_S(\hat{F}_m, F_0)$ as the sample size m grows. The mixture-of-parameters estimator dominates the other methods by a wide margin: Its divergences are very close to zero, and show little variation across replicates. Under the logarithmic score, the performance of the kernel density estimator is highly variable across the replicates, even for large sample sizes. The variability is less under the CRPS, where the kernel density approach using the Silverman (1986) rule




Figure 1: Score divergences in the simulation study with $(\alpha, s, n) = (0.5, 2, 12)$. For a given method and MCMC sample size, the bars range from the 10th to the 90th percentile of the score divergences across 1 000 replicates. The squares mark the respective medians.

of thumb for bandwidth selection performs similar to the empirical CDF-based approximation. Other bandwidth selection rules we have experimented with tend to be inferior, as indicated by slower convergence and higher variability across replicates. Finally, we observe the lack of consistency of the Gaussian approximation.

Figure 2 provides insight into the performance of the mixture-of-parameters approximation for small MCMC samples. Using as few as 150 draws, the method attains a lower median CRPS divergence than the kernel density estimator based on 20 000 draws. The superiority of the mixture-of-parameters estimator is even more pronounced under the logarithmic score, where only 50 draws are required to outperform the kernel density estimator based on 20 000 draws.

4.4 Thinning the MCMC Sample

In Appendix D, we present simulation analyses of the effects of thinning an MCMC sample (i.e., keeping only every τ th draw, where $\tau \in \mathbb{N}$ is the thinning factor), which is often done in practice with the goal of reducing autocorrelation in the MCMC draws. From a practical perspective, the analysis in Appendix D suggests that thinning is justified if, and only if, a small MCMC sample is desired and the mixture-of-parameter estimator is applied. Two arguments in favor of a small sample appear particularly relevant even today. First, storing large amounts of data on public servers (as is often done for replication purposes) may be costly or inconvenient. Second, post-processing procedures such as score computations applied to the MCMC sample may be computationally demanding (cf. Table 2), and therefore may encourage thinning.

5 Economic data example

In real-world uses of Bayesian forecasting methods, the posterior predictive distribution F_0 is typically not available in closed form. Therefore, computing or estimating the object




Figure 2: Performance of the mixture-of-parameters estimator. The design is as in Figure 1, but for smaller sample sizes. For comparison, the blue horizontal line marks the median divergence of the kernel density estimator based on 20 000 draws.

of interest for assessing consistency, i.e., the score divergence $d_S(\hat{F}_m, F_0)$, is not feasible. In the subsequent data example, we thus compare the approximation methods via their out-of-sample predictive performance, and examine the variation of the mean scores across chains obtained by replicates with distinct random seeds. While studying the predictive performance does not allow to assess consistency of the approximations, it does allow us to assess the variability and applicability of the approximations in a practical setting.

5.1 Data

We consider quarterly growth rates of U.S. real gross domestic product (GDP), as illustrated in the Online Supplement. The training sample used for model estimation is recursively expanded as forecasting moves forward in time. We use the real-time data set provided by the Federal Reserve Bank of Philadelphia⁴, which provides historical snapshots of the data vintages available at any given date in the past, and consider forecasts for the period from the second quarter of 1996 to the third quarter of 2014, for a total of $T = 74$ forecast cases. For brevity, we present results for a prediction horizon of one quarter only. The Online Supplement contains results for longer horizons, which are qualitatively similar to the ones presented here.

5.2 Probabilistic forecasts

To construct density forecasts, we consider an autoregressive (AR) model with a single lag and state-dependent error term variance, in that

$$Y_t = \nu + \alpha Y_{t-1} + \varepsilon_t, \quad (18)$$

where $\varepsilon_t \sim \mathcal{N}(0, \eta_{s_t}^2)$ and $s_t \in \{1, 2\}$ is a discrete state variable that switches according to a first-order Markov chain. The model, which is a variant of the Markov switching model proposed by Hamilton (1989), provides a simple description of time-varying heteroscedasticity. The latter is an important stylized feature of macroeconomic time series (see, e.g., Clark and Ravazzolo, 2015).

⁴<https://www.phil.frb.org/research-and-data/real-time-center/real-time-data/>




Figure 3: Mean score in the data example against sample size. The dots represent 16 parallel MCMC chains, and the lines connect averages across chains.

We conduct Bayesian inference via a Gibbs sampler, for which we give details in Appendix E. Let θ_i denote the complete set of latent states and model parameters at iteration i of the Gibbs sampler. Conditional on θ_i , the predictive distribution under the model in (18) is Gaussian with mean $\mu_i = \mu(\theta_i)$ and standard deviation $\sigma_i = \sigma(\theta_i)$, where we suppress time and forecast horizon for simplicity. At each forecast origin date $t = 1, \dots, T = 74$, we produce 10 000 burn-in draws, and use 40 000 draws post burn-in. We construct 16 parallel chains in this way. The (time-averaged) mean score of a given approximation method, based on m MCMC draws within chain $c = 1, \dots, 16$, is

$$\bar{S}_{m,c} = \frac{1}{T} \sum_{t=1}^T S(\hat{F}_{m,c,t}, y_t),$$

where $\hat{F}_{m,c,t}$ is the probabilistic forecast at time t . The variation of $\bar{S}_{m,c}$ across chains c is due to differences in random seeds. From a pragmatic perspective, a good approximation method should be such that the values $(\bar{S}_{m,c})_{c=1}^{16}$ are small and display little variation.

5.3 Results

In Figure 3, the mean score is plotted against the size of the MCMC sample. The mixture-of-parameters approximation outperforms its competitors: Its scores display the smallest variation across chains, for both the CRPS and the logarithmic score, and for all sample sizes. The scores of the mixture-of-parameter estimator also tend to be lower (i.e., better) than the scores for the other methods. The kernel density estimator performs poorly for small sample sizes, with the scores varying substantially across chains. Under the CRPS, the kernel density estimator is dominated by the empirical CDF technique, which can be interpreted as kernel density estimation with a bandwidth of zero.

6 Discussion

We have investigated how to make and evaluate probabilistic forecasts based on MCMC output. The formal notion of consistency allows us to assess the appropriateness of approximation methods within the framework of proper scoring rules. Despite their popularity in the literature, Gaussian approximations generally fail to be consistent. Conditions for consistency depend on the scoring rule of interest, and we have demonstrated that the mixture-of-parameters and empirical CDF-based approximations are consistent relative to the CRPS under minimal conditions. Proofs of consistency relative to the logarithmic score generally rely on stringent assumptions.

In view of these theoretical considerations as well as the practical perspective taken in our simulation and data examples, we generally recommend the use of the mixture-of-parameters estimator, which provides an efficient approximation method and outperforms all alternatives. This can be explained by the fact that it efficiently exploits the parametric structure of the Bayesian model. The empirical CDF-based approximation provides a good alternative if the conditional distributions fail to be available in closed form, or if for some reason the draws are to be made directly from the posterior predictive distribution, as opposed to using parameter draws. The empirical CDF-based approximation is available under the CRPS and DSS but not under the logarithmic score, where a density is required. Under the logarithmic score, the case for the mixture-of-parameters estimator is thus particularly strong. In particular, the score’s sensitivity to the tails of the distribution renders kernel density estimators unattractive from both theoretical and applied perspectives.

Our recommendations have been implemented in the `scoringRules` package for R (R Core Team, 2017); see Jordan et al. (2019) for details. The functions and default choices aim to provide readily applicable and efficient approximations. The mixture-of-parameters estimator depends on the specific structure of the Bayesian model and can therefore not be covered in full generality. However, the implemented analytical solutions of the CRPS and logarithmic score allow for straightforward and efficient computation. The `scoringRules` package further contains functions and data for replicating the simulation and case study, with details provided at https://github.com/FK83/scoringRules/blob/master/KLTG2020_replication.pdf.

Ferro (2014) studies the notion of a fair scoring rule in the context of ensemble weather forecasts. A scoring rule is called *fair* if the expected score is optimal for samples with members that behave as though they and the verifying observation were sampled from the same distribution. While certainly relevant in the context of meteorological forecast ensembles, where the sample size m is typically between 10 and 50, these considerations seem less helpful in the context of MCMC output, where m is on the order of thousands and can be increased at low cost. Furthermore, the proposed small sample adjustments and the characterization of fair scores hold for independent samples only, an assumption that is thoroughly violated in the case of MCMC.

We are interested in evaluating probabilistic forecasts produced via MCMC, so that the predictive performance of a model during an out-of-sample, test or evaluation period can be used to estimate its forecast performance on future occasions. In contrast, information criteria suggest a different route towards estimating forecast performance (Spiegelhalter et al., 2002; Watanabe, 2010; Hooten and Hobbs, 2015). They consider a method’s in-sample performance, and account for model complexity via penalty terms. Preferred ways of doing so have been the issue of methodological debate, and a consensus has not been reached; see, e.g., the comments in Gelman et al. (2014b) and Spiegelhalter et al. (2014). This present analysis does not concern in-sample comparisons, and does not address the

question of whether these are more or less effective than out-of-sample comparisons. However, our results and observations indicate that out-of-sample comparisons of the type considered here yield robust results across a range of implementation choices.

Necessarily, the scope of this paper is restricted along several dimensions. First, our theoretical results focus on consistency but do not cover rates of convergence. Results on the latter tend to rely on theoretical conditions that are hard to verify empirically, and the plausibility of which is likely to depend on the specifics of the MCMC algorithm. In contrast, many of our consistency results require only minimal conditions that hold across a wide range of sampling algorithms in the interdisciplinary applied literature. Second, we have focused on univariate continuous forecast distributions. The corresponding applied literature is large and features a rich variety of implementation variants (cf. Table 1 of the Online Supplement). Nevertheless, there are other empirically relevant setups, notably univariate discrete distributions and multivariate continuous distributions, which we briefly discuss in turn.

In relevant discrete settings, such as predicting probabilities of a binary or categorical outcome, the estimation problem becomes considerably simpler than for the real-valued case. The more complex case of integer-valued count data can be handled using methods similar to the ones we discuss. Instead of probability density functions, the count data case involves probability mass functions to which both the logarithmic score and the CRPS transfer naturally (Czado et al., 2009). Furthermore, all of the approximation methods we discuss can be used in the count data case. For example, the mixture-of-parameters estimator can be used in concert with a Poisson or Negative Binomial specification. Similarly, Shirota and Gelfand (2017, Section 4) consider eq. (10) in a count data context, and kernel-type smoothing methods have been proposed for count data as well (Rajagopalan and Lall, 1995).

The multivariate case features novel challenges. Perhaps most fundamentally, a consensus on practically appropriate choices of the scoring rule is yet to be reached (Gneiting et al., 2008; Scheuerer and Hamill, 2015). Held et al. (2017, Section 4.2) and White et al. (2019, Section 3.3) propose the use of the empirical CDF approximation in concert with the multivariate energy score. In this setting, analogues of our Theorem 3 hold, assuring consistency under weak conditions. For kernel density estimators the ‘curse of dimensionality’ applies, and for the mixture-of-parameters estimator we expect numerical challenges when evaluating, say, a log predictive density in a high-dimensional space. Clearly, there is considerable scope and opportunity for future research in these directions.

References

- Amisano, G. and Giacomini, R. (2007). Comparing density forecasts via weighted likelihood ratio tests. *J. Bus. Econom. Statist.*, 25, 177–190.
- Barron, A. R., Györfi, L. and van der Meulen, E. C. (1992). Distribution estimation consistent in total variation and in two types of information divergence. *IEEE Trans. Inform. Theory*, 38, 1437–1454.
- Bellemare, M. G., Danihelka, I., Dabney, W., Mohamed, S., Lakshminarayanan, B., Hoyer, S. and Munos, R. (2017). The Cramer distance as a solution to biased Wasserstein gradients. Preprint, available at <http://arxiv.org/abs/1705.10743>.
- Bernardo, J. M. (1979). Expected information as expected utility. *Ann. Statist.*, 7, 686–690.
- Bolin, D. and Wallin, J. (2019). Multivariate type-G Matérn stochastic partial differential equation random fields. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* Forthcoming.

- Brier, G. W. (1950). Verification of forecasts expressed in terms of probability. *Mon. Weather Rev.*, 78, 1–3.
- Clark, J. S. (2005). Why environmental scientists are becoming Bayesians. *Ecol. Lett.*, 8, 2–14.
- Clark, T. and McCracken, M. (2013). Advances in forecast evaluation. In *Handbook of Economic Forecasting* (G. Elliott and A. Timmermann, eds.), vol. 2. Elsevier, 1107–1201.
- Clark, T. E. and Ravazzolo, F. (2015). Macroeconomic forecasting performance under alternative specifications of time-varying volatility. *J. Appl. Econometrics*, 30, 551–575.
- Cooke, W. E. . (1906). Forecasts and verifications in Western Australia. *Mon. Weather Rev.*, 34, 23–24.
- Craiu, R. V. and Rosenthal, J. S. (2014). Bayesian computation via Markov chain Monte Carlo. *Annu. Rev. Stat. Appl.*, 1, 179–201.
- Czado, C., Gneiting, T. and Held, L. (2009). Predictive model assessment for count data. *Biometrics*, 65, 1254–1261.
- Dawid, A. P. (1984). Present position and potential developments: Some personal views. Statistical theory: The prequential approach. *J. R. Stat. Soc. Ser. A. Gen.*, 147, 278–290.
- Dawid, A. P. and Sebastiani, P. (1999). Coherent dispersion criteria for optimal experimental design. *Ann. Statist.*, 27, 65–81.
- Dehling, H. and Philipp, W. (2002). Empirical process techniques for dependent data. In *Empirical Process Techniques for Dependent Data* (H. Dehling, T. Mikosch and M. Sørensen, eds.). Birkhäuser, Boston, 3–113.
- DelSole, T. and Tippett, M. K. (2014). Comparing forecast skill. *Mon. Weather Rev.*, 142, 4658–4678.
- Diebold, F. X., Gunther, T. A. and Tay, A. S. (1998). Evaluating density forecasts with applications to financial risk management. *Internat. Econom. Rev.*, 39, 863–883.
- Diebold, F. X. and Mariano, R. S. (1995). Comparing predictive accuracy. *J. Bus. Econom. Statist.*, 13, 253–263.
- Ferro, C. A. T. (2014). Fair scores for ensemble forecasts. *Q. J. Royal Meteorol. Soc.*, 140, 1917–1923.
- Fox, E. B. and West, M. (2011). Autoregressive models for variance matrices: Stationary inverse Wishart processes. Preprint, available at <http://arxiv.org/abs/1107.5239>.
- Gelfand, A. E. and Smith, A. F. M. (1990). Sampling-based approaches to calculating marginal densities. *J. Amer. Statist. Assoc.*, 85, 398–409.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A. and Rubin, D. B. (2014a). *Bayesian Data Analysis*. 3rd ed. Chapman & Hall/CRC, Boca Raton.
- Gelman, A., Hwang, J. and Vehtari, A. (2014b). Understanding predictive information criteria for Bayesian models. *Stat. Comput.*, 24, 997–1016.
- Gelman, A., Meng, X.-L. and Stern, H. (1996). Posterior predictive assessment of model fitness via realized discrepancies. *Statist. Sinica*, 6, 733–760.
- Genon-Catalot, V., Jeantheau, T. and Larédo, C. (2000). Stochastic volatility models as hidden Markov models and statistical applications. *Bernoulli*, 6, 1051–1079.
- Geweke, J. (2005). *Contemporary Bayesian Econometrics and Statistics*. John Wiley & Sons, Hoboken.
- Geyer, C. J. (1992). Practical Markov chain Monte Carlo. *Statist. Sci.*, 7, 473–483.
- Giacomini, R. and White, H. (2006). Tests of conditional predictive ability. *Econometrica*, 74, 1545–1578.
- Gibbs, A. L. and Su, F. E. (2002). On choosing and bounding probability metrics. *Int. Stat. Rev.*, 70, 419–435.
- Gilks, W. R., Richardson, S. and Spiegelhalter, D. J. (eds.) (1996). *Markov Chain Monte*

- Carlo in Practice*. Chapman & Hall/CRC, Boca Raton.
- Gneiting, T. (1997). Normal scale mixtures and dual probability densities. *J. Stat. Comput. Simul.*, 59, 375–384.
- Gneiting, T. and Raftery, A. E. (2007). Strictly proper scoring rules, prediction, and estimation. *J. Amer. Statist. Assoc.*, 102, 359–378.
- Gneiting, T., Stanberry, L. I., Grimit, E. P., Held, L. and Johnson, N. A. (2008). Assessing probabilistic forecasts of multivariate quantities, with an application to ensemble predictions of surface winds (with discussion and rejoinder). *Test*, 17, 211–264.
- Good, I. J. (1952). Rational decisions. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 14, 107–114.
- Greenberg, E. (2013). *Introduction to Bayesian Econometrics*. 2nd ed. Cambridge University Press, New York.
- Grimit, E. P., Gneiting, T., Berrocal, V. J. and Johnson, N. A. (2006). The continuous ranked probability score for circular variables and its application to mesoscale forecast ensemble verification. *Q. J. Royal Meteorol. Soc.*, 132, 2925–2942.
- Györfi, L., Härdle, W., Sarda, P. and Vieu, P. (1989). *Nonparametric Curve Estimation from Time Series*. Springer, Berlin.
- Hall, P. (1987). On Kullback-Leibler loss and density estimation. *Ann. Statist.*, 15, 1491–1519.
- Hall, P., Lahiri, S. N. and Truong, Y. K. (1995). On bandwidth choice for density estimation with dependent data. *Ann. Statist.*, 23, 2241–2263.
- Hall, P. and Presnell, B. (1999). Density estimation under constraints. *J. Comput. Graph. Statist.*, 8, 259–277.
- Hamilton, J. D. (1989). A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57, 357–384.
- Hansen, B. E. (2008). Uniform convergence rates for kernel estimation with dependent data. *Econom. Theory*, 24, 726–748.
- Harris, I. R. (1989). Predictive fit for natural exponential families. *Biometrika*, 76, 675–684.
- Held, L., Meyer, S. and Bracher, J. (2017). Probabilistic forecasting in infectious disease epidemiology: The 13th Armitage lecture. *Stat. Med.*, 36, 3443–3460.
- Held, L., Schrödle, B. and Rue, H. (2010). Posterior and cross-validatory predictive checks: A comparison of MCMC and INLA. In *Statistical Modelling and Regression Structures: Festschrift in Honour of Ludwig Fahrmeir* (T. Kneib and G. Tutz, eds.). Physica-Verlag HD, Heidelberg, 91–110.
- Hooten, M. B. and Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists. *Ecol. Monogr.*, 85, 3–28.
- Huber, P. J. and Ronchetti, E. M. (2009). *Robust Statistics*. 2nd ed. Wiley, Hoboken, New Jersey.
- Jones, M. C. (1991). On correcting for variance inflation in kernel density estimation. *Comput. Stat. Data Anal.*, 11, 3–15.
- Jordan, A. (2016). Facets of forecast evaluation. Ph.D. thesis, Karlsruhe Institute of Technology, available at <https://publikationen.bibliothek.kit.edu/1000063629>.
- Jordan, A., Krüger, F. and Lerch, S. (2019). Evaluating probabilistic forecasts with scoringRules. *J. Stat. Softw.*, 90, 1–37.
- Kass, R. E. and Raftery, A. E. (1995). Bayes factors. *J. Amer. Statist. Assoc.*, 90, 773–795.
- Kim, H. J., MacEachern, S. N. and Jung, Y. (2016). Bandwidth selection for kernel density estimation with a Markov chain Monte Carlo sample. Preprint, available at <http://arxiv.org/abs/1607.08274>.
- Krüger, F., Clark, T. E. and Ravazzolo, F. (2017). Using entropic tilting to combine BVAR forecasts with external nowcasts. *J. Bus. Econom. Statist.*, 35, 470–485.

- Kullback, S. (1959). *Information Theory and Statistics*. John Wiley & Sons.
- Liebscher, E. (1996). Strong convergence of sums of α -mixing random variables with applications to density estimation. *Stochastic Process. Appl.*, 65, 69–80.
- Link, W. A. and Eaton, M. J. (2012). On thinning of chains in MCMC. *Methods Ecol. Evol.*, 3, 112–115.
- Little, R. J. (2006). Calibrated Bayes: A Bayes/frequentist roadmap. *Amer. Statist.*, 60, 213–223.
- MacEachern, S. N. and Berliner, L. M. (1994). Subsampling the Gibbs sampler. *Amer. Statist.*, 48, 188–190.
- Matheson, J. E. and Winkler, R. L. (1976). Scoring rules for continuous probability distributions. *Manag. Sci.*, 22, 1087–1096.
- Panaretos, V. M. and Zemel, Y. (2019). Statistical aspects of Wasserstein distances. *Annu. Rev. Stat. Appl.*, 6, 405–431.
- R Core Team (2017). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Rajagopalan, B. and Lall, U. (1995). A kernel estimator for discrete distributions. *J. Nonparametr. Stat.*, 4, 409–426.
- Rosenblatt, M. (1956). Remarks on some nonparametric estimates of a density function. *Ann. Math. Stat.*, 27, 832–837.
- Rosenthal, J. S. (1995). Minorization conditions and convergence rates for Markov chain Monte Carlo. *J. Amer. Statist. Assoc.*, 90, 558–566.
- Roussas, G. G. (1988). Nonparametric estimation in mixing sequences of random variables. *J. Statist. Plann. Inference*, 18, 135–149.
- Rubin, D. B. (1984). Bayesianly justifiable and relevant frequency calculations for the applied statistician. *Ann. Statist.*, 12, 1151–1172.
- Scheuerer, M. and Hamill, T. M. (2015). Variogram-based proper scoring rules for probabilistic forecasts of multivariate quantities. *Mon. Weather Rev.*, 143, 1321–1334.
- Sheather, S. J. and Jones, M. C. (1991). A reliable data-based bandwidth selection method for kernel density estimation. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 53, 683–690.
- Shirota, S. and Gelfand, A. E. (2017). Space and circular time log Gaussian Cox processes with application to crime event data. *The Annals of Applied Statistics*, 11, 481–503.
- Shuford, E. H., Albert, A. and Massengill, H. E. (1966). Admissible probability measurement procedures. *Psychometrika*, 31, 125–145.
- Silverman, B. W. (1986). *Density Estimation for Statistics and Data Analysis*. Chapman and Hall, London.
- Sköld, M. and Roberts, G. O. (2003). Density estimation for the Metropolis-Hastings algorithm. *Scand. J. Stat.*, 30, 699–718.
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P. and van der Linde, A. (2002). Bayesian measures of model complexity and fit (with discussion and rejoinder). *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 64, 583–639.
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P. and van der Linde, A. (2014). The deviance information criterion: 12 years on. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 76, 485–493.
- Taillardat, M., Mestre, O., Zamo, M. and Naveau, P. (2016). Calibrated ensemble forecasts using quantile regression forests and ensemble model output statistics. *Mon. Weather Rev.*, 144, 2375–2393.
- Tanner, M. A. and Wong, W. H. (1987). The calculation of posterior distributions by data augmentation. *J. Amer. Statist. Assoc.*, 82, 528–540.
- Thorarinsdottir, T. L., Gneiting, T. and Gissibl, N. (2013). Using proper divergence

- functions to evaluate climate models. *SIAM/ASA J. Uncertain. Quantif.*, 1, 522–534.
- Tierney, L. (1994). Markov chains for exploring posterior distributions. *Ann. Statist.*, 22, 1701–1728.
- van der Vaart, A. W. (2000). *Asymptotic Statistics*. Cambridge University Press, Cambridge.
- Villani, C. (2008). *Optimal Transport*. Grundlehren der mathematischen Wissenschaften, 338, Springer, Berlin, Heidelberg.
- Waggoner, D. F. and Zha, T. (1999). Conditional forecasts in dynamic multivariate models. *Rev. Econ. Stat.*, 81, 639–651.
- Warne, A., Coenen, G. and Christoffel, K. (2016). Marginalized predictive likelihood comparisons of linear Gaussian state-space models with applications to DSGE, DSGE-VAR and VAR models. *J. Appl. Econometrics*, 32, 103–119.
- Wasserman, L. (2006). *All of Nonparametric Statistics*. Springer, New York.
- Watanabe, S. (2010). Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory. *J. Mach. Learn. Res.*, 11, 3571–3594.
- White, P. A., Gelfand, A. E., Rodrigues, E. R. and Tzintzun, G. (2019). Pollution state modelling for Mexico City. *J. R. Stat. Soc. Ser. A. Stat. Soc.*, 182, 1039–1060.
- Yu, B. (1993). Density estimation in the L^∞ norm for dependent data with applications to the Gibbs sampler. *Ann. Statist.*, 21, 711–735.

Acknowledgements

The work of Tilmann Gneiting and Fabian Krüger was funded by the European Union Seventh Framework Programme under grant agreement 290976. Sebastian Lerch and Thordis L. Thorarinsdottir acknowledge support by the Volkswagen Foundation through the program “Mesoscale Weather Extremes — Theory, Spatial Modelling and Prediction (WEX-MOP)”. Lerch further acknowledges support by Deutsche Forschungsgemeinschaft (DFG) through RTG 1953 “Statistical Modeling of Complex Systems and Processes” and SFB/TRR 165 “Waves to Weather”. Gneiting, Krüger and Lerch thank the Klaus Tschira Foundation for infrastructural support at the Heidelberg Institute for Theoretical Studies (HITS). Helpful comments by Werner Ehm, Sylvia Frühwirth-Schnatter, Alexander Jordan, as well as seminar and conference participants at HITS, KIT, University of Bern, University of Bonn, University of Oslo, the Extremes 2014 symposium (Hannover), CFE (Pisa, 2014), GPSD (Bochum, 2016), ISBA (Sardinia, 2016), and Deutsche Bundesbank (Workshop on Forecasting, 2017) are gratefully acknowledged. Furthermore, we thank Gianni Amisano for sharing his program code for Bayesian Markov switching models. We further thank an anonymous referee of a previous version of the manuscript for pointing us to the Rao-Blackwellization arguments employed in Theorem 4.

A Computing the CRPS for mixtures of Gaussians

Here we discuss the computation of the CRPS in (7) when the predictive distribution is an equally weighted mixture of normal distributions, say $F = \hat{F}_m^{\text{MP}}$, where $F_c(\cdot|\theta_i)$ is Gaussian with mean μ_i and variance σ_i^2 . Gneiting et al. (2006) note that in this case (8) can be written




Figure 4: CRPS scores for the first quarter of 2011 in the data example, for 16 parallel chains and various MCMC sample sizes. Left: The segments connect the CRPS value obtained using numerical integration (left node) to the score obtained using the exact formula (right node). Right: Computation times in seconds, for numerical integration (dots; solid line) and exact formula (crosses; dashed line).

as

$$\text{CRPS}(\hat{F}_m^{\text{MP}}, y) = \frac{1}{m} \sum_{i=1}^m A(y - \mu_m, \sigma_m^2) - \frac{1}{2m^2} \sum_{i=1}^m \sum_{j=1}^m A(\mu_i - \mu_j, \sigma_i^2 + \sigma_j^2), \quad (19)$$

where $A(\mu, \sigma^2) = 2\sigma\phi(\frac{\mu}{\sigma}) + \mu(2\Phi(\frac{\mu}{\sigma}) - 1)$, with ϕ and Φ denoting the standard normal density and CDF, respectively. The `scoringRules` software package (Jordan et al., 2019) contains R/C++ code for the evaluation of (19), which requires $\mathcal{O}(m^2)$ operations.

A potentially much faster, but not exact, alternative is to evaluate the integral in (7) numerically.⁵ Here we provide some evidence on the viability of this strategy, which we implement via the R function `integrate`, with arguments `rel.tol` and `abs.tol` of `integrate` set to 10^{-6} . As a first experiment, we use numerical integration to re-compute the CRPS scores of the mixture-of-parameters estimator in our data example for the first quarter of 2011. Figure 4 summarizes the results for 16 parallel chains. The left panel shows that the approximate scores are visually identical to the exact ones across all sample sizes and chains. Indeed, the maximal absolute error incurred by numerical integration is 8.0×10^{-8} . The approximation errors are dwarfed by the natural variation of the scores across MCMC chains. The right panel compares the computation time for exact evaluation vs. numerical integration. The latter is much faster, especially for large samples. For a sample of size 40,000 numerical integration requires less than 1.5 seconds, whereas exact evaluation requires about two minutes on an Intel i7 processor.

To obtain broad-based evidence, we next compare exact evaluation vs. numerical integration for all 74 forecast dates, from the second quarter of 1996 to the third quarter of 2014, employing 16 parallel chains for each date. We focus on the two largest MCMC sample sizes, 20 000 and 40 000, and find that across all 2 368 instances (74 dates times 2

⁵Numerical integration could also be based on another representation of the CRPS that has recently been derived by Taillardat et al. (2016, p. 2390, bottom right).

sample sizes times 16 chains), the absolute difference of the two CRPS values never exceeds 6.3×10^{-7} . Therefore, we feel that numerical integration allows for the efficient evaluation of the CRPS for mixtures of normal distributions. The differences to the exact values are practically irrelevant and well in line with the error bounds in R's `integrate` function.

B Consistency of mixture-of-parameters approximations

Proof of Theorem 1

In the case of the CRPS, we prove the stronger result that $\int_{\mathbb{R}} |\hat{F}_m^{\text{MP}}(z) - F_0(z)| dz \rightarrow 0$ almost surely as $m \rightarrow \infty$. Putting $H(z) = 1 - F_0(z) + F_0(-z)$ and $\hat{H}_m(z) = 1 - \hat{F}_m^{\text{MP}}(z) + \hat{F}_m^{\text{MP}}(-z)$ for $z \in \mathbb{R}$, we find that, for every fixed $N > 0$,

$$\begin{aligned} \limsup_{m \rightarrow \infty} \int_{\mathbb{R}} |\hat{F}_m^{\text{MP}}(z) - F_0(z)| dz &\leq \limsup_{m \rightarrow \infty} \int_{-N}^N |\hat{F}_m^{\text{MP}}(z) - F_0(z)| dz \\ &\quad + \int_N^{\infty} H(z) dz + \limsup_{m \rightarrow \infty} \int_N^{\infty} \hat{H}_m(z) dz. \end{aligned} \quad (20)$$

The Ergodic Theorem implies that the first term on the right-hand side of (20) tends to zero, and that

$$\int_N^{\infty} \hat{H}_m(z) dz = \int_N^{\infty} \frac{1}{m} \sum_{i=1}^m (1 - F_c(z|\theta_i) + F_c(-z|\theta_i)) dz \longrightarrow \int_N^{\infty} H(z) dz$$

almost surely as $m \rightarrow \infty$. In view of (20) we conclude that

$$\limsup_{m \rightarrow \infty} \int_{\mathbb{R}} |\hat{F}_m^{\text{MP}}(z) - F_0(z)| dz \leq 2 \int_N^{\infty} H(z) dz \quad (21)$$

almost surely as $m \rightarrow \infty$. As the right-hand side of (21) decreases to zero as N grows without bounds, the proof of the claim is complete.

In the case of the DSS, let $K(z) = 1 - F_0(z) - F_0(-z)$ and $\hat{K}_m(z) = 1 - \hat{F}_m^{\text{MP}}(z) - \hat{F}_m^{\text{MP}}(-z)$ for $z \in \mathbb{R}$. Due to the finiteness of the first moments of F_0 and \hat{F}_m^{MP} , $\int_{\mathbb{R}} z dF_0(z) = \int_0^{\infty} K(z) dz$ and $\int_{\mathbb{R}} z d\hat{F}_m^{\text{MP}}(z) = \int_0^{\infty} \hat{K}_m(z) dz$. For the second moments, we find similarly that $\int_{\mathbb{R}} z^2 dF_0(z) = 2 \int_0^{\infty} zH(z) dz$ and $\int_{\mathbb{R}} z^2 d\hat{F}_m^{\text{MP}}(z) = 2 \int_0^{\infty} z\hat{H}_m(z) dz$. Proceeding as before, the Ergodic Theorem implies almost sure convergence of the first and second moments, and thereby consistency relative to the DSS.

Proof of Theorem 2

By Lemma 2.1 in Chapter 4 of Kullback (1959),

$$\sup_{z \in \mathbb{R}} \left| 1 - \frac{\hat{f}_m^{\text{MP}}(z)}{f_0(z)} \right| \longrightarrow 0$$

almost surely as $m \rightarrow \infty$ implies the desired convergence of the Kullback-Leibler divergence. Let P_m denote the empirical CDF of the parameter draws $(\theta_i)_{i=1}^m$. Under assumption (B) almost sure strong uniform consistency,

$$\sup_{z \in \Omega} \left| \hat{f}_m^{\text{MP}}(z) - f_0(z) \right| = \sup_{z \in \Omega} \left| \int_{\Theta} f_c(z|\theta) [dP_m(\theta) - dP_{\text{post}}(\theta)] \right| \longrightarrow 0$$

almost surely as $m \rightarrow \infty$, yields Kullback's condition. Finally, we establish almost sure strong uniform convergence under assumptions (A) and (B) by applying Theorem 19.4 and Example 19.8 of van der Vaart (2000).

C Consistency of empirical CDF-based approximations

Proof of Theorem 3

In the case of the CRPS, we proceed in analogy to the proof of Theorem 1 and demonstrate the stronger result that $\int_{\mathbb{R}} |\hat{F}_m^{\text{ECDF}}(z) - F_0(z)| dz \rightarrow 0$ almost surely as $m \rightarrow \infty$. Putting $H(z) = 1 - F_0(z) + F_0(-z)$ and $\hat{H}_m(z) = 1 - \hat{F}_m^{\text{ECDF}}(z) + \hat{F}_m^{\text{ECDF}}(-z)$ for $z \in \mathbb{R}$, we see that, for every fixed $N > 0$,

$$\begin{aligned} \limsup_{m \rightarrow \infty} \int_{\mathbb{R}} |\hat{F}_m^{\text{ECDF}}(z) - F_0(z)| dz &\leq \limsup_{m \rightarrow \infty} \int_{-N}^N |\hat{F}_m^{\text{ECDF}}(z) - F_0(z)| dz \\ &\quad + \int_N^\infty H(z) dz + \limsup_{m \rightarrow \infty} \int_N^\infty \hat{H}_m(z) dz. \end{aligned} \quad (22)$$

The Generalized Glivenko-Cantelli Theorem (Dehling and Philipp, 2002, Theorem 1.1) implies that the first term on the right-hand side of (22) tends to zero almost surely as $m \rightarrow \infty$. If Z_0 has distribution F_0 , then $\int_N^\infty H(z) dz = \mathbb{E}(|Z_0| - N)_+$, where $(W)_+ = \max(W, 0)$ denotes the positive part of W . Furthermore, by the Ergodic Theorem

$$\int_N^\infty \hat{H}_m(z) dz = \frac{1}{m} \sum_{i=1}^m (|X_i| - N)_+ \longrightarrow \mathbb{E}(|Z_0| - N)_+$$

almost surely as $m \rightarrow \infty$, which along with (22) implies that

$$\limsup_{m \rightarrow \infty} \int_{\mathbb{R}} |\hat{F}_m^{\text{ECDF}}(z) - F_0(z)| dz \leq 2 \mathbb{E}(|Z_0| - N)_+ \quad (23)$$

almost surely as $m \rightarrow \infty$. As the right-hand side of (23) gets arbitrarily close to zero as N grows without bounds, the proof of the claim is complete.

In the case of the DSS, it suffices to note that the moments of the empirical CDF are the sample moments of $(X_i)_{i=1}^m$, and then to apply the Ergodic Theorem.

Proof of Theorem 4

By the law of total expectation, $\mathbb{E} \hat{F}_m^{\text{ECDF}}(z) = \mathbb{E} \hat{F}_m^{\text{MP}}(z)$ as

$$\begin{aligned} \mathbb{E} \left(\hat{F}_m^{\text{ECDF}}(z) | \theta_1, \dots, \theta_m \right) &= \frac{1}{m} \sum_{i=1}^m \mathbb{P}(X_i \leq z | \theta_1, \dots, \theta_m) \\ &= \frac{1}{m} \sum_{i=1}^m \mathbb{P}(X_i \leq z | \theta_i) \\ &= \hat{F}_m^{\text{MP}}(z). \end{aligned}$$

Further, the law of total variance implies

$$\begin{aligned}\text{Var}\left(\hat{F}_m^{\text{ECDF}}(z)\right) &= \mathbb{E}\left[\text{Var}\left(\hat{F}_m^{\text{ECDF}}(z)|\theta_1, \dots, \theta_m\right)\right] + \text{Var}\left[\mathbb{E}\left(\hat{F}_m^{\text{ECDF}}(z)|\theta_1, \dots, \theta_m\right)\right] \\ &\geq \text{Var}\left(\hat{F}_m^{\text{MP}}(z)\right)\end{aligned}$$

for every $z \in \mathbb{R}$ and $m \in \mathbb{N}$. For a generic estimator \hat{F}_m with finite mean,

$$\begin{aligned}\mathbb{E} d_{\text{CRPS}}(\hat{F}_m, F_0) &= \mathbb{E} \int \left(\hat{F}_m(z) - F_0(z)\right)^2 dz \\ &= \int \mathbb{E} \left(\hat{F}_m(z) - F_0(z)\right)^2 dz \\ &= \int \text{Var} \hat{F}_m(z) dz + \int \left(\mathbb{E} \hat{F}_m(z) - F_0(z)\right)^2 dz.\end{aligned}$$

In this light, the first part of the theorem's statement implies the second part.

D Simulation Study on Thinning an MCMC sample


Using the same simulation setup as in Section 4, we further investigate the effect of thinning the Markov chains. Thinning a chain by a factor of τ means that only every τ th simulated value is retained, and the rest is discarded. Thinning is often applied routinely with the goal of reducing autocorrelation in the draws. Of the articles listed in Table 1 of the Online Supplement, about one in four explicitly reports thinning of the simulation output, with thinning factors ranging from 2 to 100. Here we compare three sampling approaches:

- (S1) 4000 MCMC draws, without thinning
- (S2) 4000 MCMC draws, retaining every 10th draw from a sequence of 40 000 draws
- (S3) 40 000 MCMC draws, without thinning


Note that the samples in S1 and S3 have the same dynamic properties, whereas S2 will typically produce a chain with less autocorrelation. Furthermore, S2 and S3 require the same computing time, which exceeds that of S1 by a factor of ten. Figure 5 summarizes the corresponding simulation results, using parameter values $s = 2$ and $n = 12$, and varying values of the persistence parameter α . We report results for four popular combinations of scoring rules and approximation methods.

As expected, S2 tends to outperform S1: When the sample size is held fixed, less autocorrelation entails more precise estimators. While the difference in performance is modest in most cases, S2 attains large (relative) gains over S1 when the mixture-of-parameters estimator is applied to a very persistent sample with $\alpha = 0.9$. This can be explained by the direct effect of the persistence parameter α on the parameter draws $(\theta_i)_{i=1}^m$, whereas the influence is less immediate for the KDE and ECDF approximation methods, which are based on the sequence $(X_i)_{i=1}^m$ obtained in an additional sampling step. Furthermore, S3 outperforms S2 in all cases covered. While the effects of thinning have not been studied in the context of predictive distributions before, this observation is in line with extant reports of the greater precision of unthinned chains (Geyer, 1992; MacEachern and Berliner, 1994; Link and Eaton, 2012). The performance gap between S3 and S2 is modest for the mixture-of-parameters estimator (top row of Figure 5), but very pronounced for the other estimators.


Logarithmic score, Mixture-of-parameters



CRPS, Mixture-of-parameters



Logarithmic score, Kernel density estimation



CRPS, Empirical CDF




Figure 5: Performance of three sampling strategies: S1: 4000 draws, without thinning; S2: 4000 MCMC draws, retaining every 10th draw from a sequence of 40000 draws; S3: 40000 draws, without thinning. Bars range from the 10th to the 90th percentile of the score divergences across 1000 replicates. The squares mark the respective medians.

E Implementation details for the data example

Here we provide additional information on the Markov switching model for the quarterly U.S. GDP growth rate, Y_t . As described in equation (18) in Section 5, the model is given by $Y_t = \nu + \alpha Y_{t-1} + \varepsilon_t$, where $\varepsilon_t \sim \mathcal{N}(0, \eta_{s_t}^2)$, and $s_t \in \{1, 2\}$ is a discrete state variable that switches according to a Markov chain.

Our implementation follows Amisano and Giacomini (2007, Section 6.3), in that our prior distributions have the same functional forms but possibly different parameter choices, as summarized in Table 4. However, note that we use prior parameters for the residual variances in both latent states, whereas Amisano and Giacomini (2007) assume the residual variance to be constant across states.

Let $\beta = (\nu, \alpha)'$ denote the parameters for the conditional mean equation (18), $\bar{s}_t = (s_1, \dots, s_t)'$ the sequence of latent states up to time t , $h = (\eta_1^{-2}, \eta_2^{-2})'$ the inverses of the state-dependent residual variances, and \mathbf{P} the 2×2 transition matrix for the latent states. Our Gibbs sampler can then be sketched as follows:

- Draw $\beta | h, \bar{s}_t$ from a Gaussian posterior. The mean and variance derive from a generalized least squares problem, with observation t receiving weight $\eta_{s_t}^{-2}$.
- Draw $h | \beta, \bar{s}_t$ from a Gamma posterior. The Gamma distribution parameters for

Table 4: Prior parameters in the Markov switching model.

Symbol in Amisano and Giacomini (2007)	$\underline{\mu}_\delta$	$\underline{\mathbf{H}}_\delta^{-1}$	\underline{s}	$\underline{\nu}$	\mathbf{R}
Parameter choice	$0_{[2,1]}$	$25 \times I_2$	0.3	3	$\begin{bmatrix} 8 & 2 \\ 2 & 8 \end{bmatrix}$
Relation to our eq. (18)	Prior mean for $(\nu, \alpha)'$	Prior variance for $(\nu, \alpha)'$	Prior parameters for $\eta_{s_t}^2$		Dirichlet prior state transitions

$\eta_s^{-2}, s \in \{1, 2\}$, are calculated from the observations t for which $s_t = s$. If necessary, permute the draws such that $\eta_1^2 > \eta_2^2$.

- Draw $\bar{s}_t | \beta, h, \mathbf{P}$ using the algorithm described by Greenberg (2013, pp. 194–195).
- Draw $\mathbf{P} | \bar{s}_t$ from a Dirichlet posterior.

Gianni Amisano kindly provides implementation details and Matlab code via his website (<https://sites.google.com/site/gianniamisanowebiste/home/teaching/istanbul-2014>, last accessed: March 25, 2019). An R implementation of his code is available within the R package `scoringRules` (Jordan et al., 2019), see https://github.com/FK83/scoringRules/blob/master/KLTG2020_replication.pdf for details.

Supplement to Predictive Inference Based on Markov Chain Monte Carlo Output

Fabian Krüger^a Sebastian Lerch^{a,b} Thordis Thorarinsdottir^c
 Tilmann Gneiting^{b,a}

^aKarlsruhe Institute of Technology

^bHeidelberg Institute for Theoretical Studies

^cNorwegian Computing Center

May 8, 2022

S1 Literature survey

In Table 1, we provide a comprehensive survey of articles that consider probabilistic forecasts based on MCMC output, subject to the following further criteria:

- We include published or pre-published articles in scientific journals or proceedings only. In particular, Table 1 does not list working papers and preprints.
- We focus on studies where forecasts based on Bayesian MCMC methods are produced, and evaluated via proper scoring rules, and we restricted our attention to studies of real-valued linear variables, ignoring articles that deal with binary or categorical observations only.
- Furthermore, as we are interested in full probabilistic forecasts based on MCMC output, we disregarded articles where only functionals of the forecast distributions, such as means or medians, are evaluated, and we retained those studies only where the computation of the scores is documented in sufficient detail.
- Finally, Table 1 lists only papers that employ the CRPS or the logarithmic score. Very few studies have used scoring rules other than these: Riebler et al. (2012) use the Dawid–Sebastiani score, and Smith and Vahey (2016) and Tran et al. (2016) use weighted versions of the logarithmic score and the CRPS, as described by Diks et al. (2011) and Gneiting and Ranjan (2011), respectively. Multivariate settings have been discussed in Held et al. (2017) and White et al. (2019a).

If a study reports several choices of approximation method and scoring rule, each choice constitutes a separate entry.

Table 1: Approximation methods and scoring rules in recent studies using probabilistic forecasts based on MCMC output.

Approximation method	Logarithmic score	CRPS
<i>Based on parameter draws</i>		
Mixture-of-parameters	Amisano and Giacomini (2007)	Kallache et al. (2010)
	Gschlößl and Czado (2007)	Trombe et al. (2012)
	Lopes et al. (2008)	Risser and Calder (2015)
	Maheu and Gordon (2008)	Fan et al. (2018)
	Liu and Maheu (2009)	
	Maheu and McCurdy (2009)	
	Geweke and Amisano (2010, 2011)	
	Jensen and Maheu (2010, 2013, 2014)	
	Jochmann et al. (2010)	
	Kallache et al. (2010)	
	Li et al. (2010)	
	Delatola and Griffin (2011)	
	Maheu et al. (2012)	
	Maneesoonthorn et al. (2012)	
	Jin and Maheu (2013, 2016)	
	Bağtürk et al. (2014)	
	Maheu and Song (2014)	
	Risser and Calder (2015)	
	Zhou et al. (2015)	
	Kastner (2016)	
	Maheu and Yang (2016)	
	Warne et al. (2016)	
	Amisano and Geweke (2017)	
	Leão et al. (2017)	
	Bitto and Frühwirth-Schnatter (2019)	
	Liu and Maheu (2018)	
	Maheu and Song (2018)	
	Wijeyakulasuriya et al. (2018)	
<i>Based on a sample</i>		
Empirical CDF		Gschlößl and Czado (2007)
		Lopes et al. (2008)
		Panagiotelis and Smith (2008)
		De la Cruz and Branco (2009)
		Salazar et al. (2011)
		Friederichs and Thorarinsdottir (2012)
		Sigrist et al. (2012)
		Groen et al. (2013)
		Leininger et al. (2013)
		Berrocal et al. (2014)
		Clark and Ravazzolo (2015)
		Krüger et al. (2017)
		Sahu et al. (2015)
		Sigrist et al. (2015)
		Smith and Vahey (2016)
		Berg (2017)
		Aastveit et al. (2017)
		Berk et al. (2018)
		Clark et al. (2018)
		Risser et al. (2019)
Kernel density estimation	Belmonte et al. (2014)	Krüger and Nolte (2016)
	Bauwens et al. (2015)	
	Berg and Henzel (2015)	
	Carriero et al. (2015b,c)	
	Berg (2017)	
Gaussian approximation	Metaxoglou et al. (2019)	
	Adolfson et al. (2007)	Brandt et al. (2014)
	Clark (2011)	Rodrigues et al. (2014)
	Carriero et al. (2015a, 2016)	
	Clark and Ravazzolo (2015)	
	Giannone et al. (2015)	
	Warne et al. (2016)	

S2 Additional figures




Figure 1: Same as Figure 1 in the paper, but for various parameter constellations of the data-generating process (cf. Table 3 in the paper).




Figure 2: Same as Figure 1 in the paper, but for various other parameter constellations of the data-generating process (cf. Table 3 in the paper).




Figure 3: Quarterly growth rate of U.S. real GDP from the second quarter of 1947 to the third quarter of 2014, using the data vintage from the first quarter of 2015.




Figure 4: Same as Figure 3 in the paper, but at a prediction horizon of two quarters.




Figure 5: Same as Figure 3 in the paper, but at a prediction horizon of three quarters.

References for the supplemental material

- Aastveit, K. A., Carriero, A., Clark, T. E. and Marcellino, M. (2017). Have standard VARs remained stable since the crisis? *J. Appl. Econom.*, 32, 931–951.
- Adolfson, M., Linde, J. and Villani, M. (2007). Forecasting performance of an open economy DSGE model. *Econometric Rev.*, 26, 289–328.
- Amisano, G. and Geweke, J. (2017). Prediction using several macroeconomic models. *Rev. Econ. Stat.*, 99, 912–925.
- Amisano, G. and Giacomini, R. (2007). Comparing density forecasts via weighted likelihood ratio tests. *J. Bus. Econom. Statist.*, 25, 177–190.
- Baştürk, N., Cakmakli, C., Ceyhan, S. P. and Van Dijk, H. K. (2014). Posterior-predictive evidence on US inflation using extended new Keynesian Phillips curve models with non-filtered data. *J. Appl. Econometrics*, 29, 1164–1182.
- Bauwens, L., Koop, G., Korobilis, D. and Rombouts, J. V. K. (2015). The contribution of structural break models to forecasting macroeconomic series. *J. Appl. Econometrics*, 30, 596–620.
- Belmonte, M. A. G., Koop, G. and Korobilis, D. (2014). Hierarchical shrinkage in time-varying parameter models. *J. Forecast.*, 33, 80–94.
- Berg, T. O. (2017). Forecast accuracy of a BVAR under alternative specifications of the zero lower bound. *Stud. Nonlinear Dyn. Econom.*, 21, 20150084.
- Berg, T. O. and Henzel, S. R. (2015). Point and density forecasts for the Euro area using Bayesian VARs. *Int. J. Forecast.*, 31, 1067–1095.
- Berk, K., Hoffmann, A. and Müller, A. (2018). Probabilistic forecasting of industrial electricity load with regime switching behavior. *Int. J. Forecast.*, 34, 147–162.
- Berrocal, V. J., Gelfand, A. E. and Holland, D. M. (2014). Assessing exceedance of ozone standards: A space-time downscaler for fourth highest ozone concentrations. *Environmetrics*, 25, 279–291.
- Bitto, A. and Frühwirth-Schnatter, S. (2019). Achieving shrinkage in a time-varying parameter model framework. *J. Econometrics*, 210, 75–97.
- Brandt, P. T., Freeman, J. R. and Schrodtt, P. A. (2014). Evaluating forecasts of political conflict dynamics. *Int. J. Forecast.*, 30, 944–962.
- Carriero, A., Clark, T. E. and Marcellino, M. (2015a). Bayesian VARs: Specification choices and forecast accuracy. *J. Appl. Econometrics*, 30, 46–73.
- Carriero, A., Clark, T. E. and Marcellino, M. (2015b). Realtime nowcasting with a Bayesian mixed frequency model with stochastic volatility. *J. R. Stat. Soc. Ser. A. Stat. Soc.*, 178, 837–862.
- Carriero, A., Clark, T. E. and Marcellino, M. (2016). Common drifting volatility in large Bayesian VARs. *J. Bus. Econom. Statist.*, 34, 375–390.
- Carriero, A., Mumtaz, H. and Theophilopoulou, A. (2015c). Macroeconomic information, structural change, and the prediction of fiscal aggregates. *Int. J. Forecast.*, 31, 325–348.
- Clark, T. E. (2011). Real-time density forecasts from BVARs with stochastic volatility. *J. Bus. Econom. Statist.*, 29, 327–341.
- Clark, T. E., McCracken, M. W. and Mertens, E. (2018). Modeling time-varying uncertainty of multiple-horizon forecast errors. *Rev. Econ. Stat.* Forthcoming.
- Clark, T. E. and Ravazzolo, F. (2015). Macroeconomic forecasting performance under alternative specifications of time-varying volatility. *J. Appl. Econometrics*, 30, 551–575.
- De la Cruz, R. and Branco, M. D. (2009). Bayesian analysis for nonlinear regression model under skewed errors, with application in growth curves. *Biom. J.*, 51, 588–609.
- Delatola, E.-I. and Griffin, J. E. (2011). Bayesian nonparametric modelling of the return

- distribution with stochastic volatility. *Bayesian Anal.*, 6, 901–926.
- Diks, C., Panchenko, V. and van Dijk, D. (2011). Likelihood-based scoring rules for comparing density forecasts in tails. *J. Econometrics*, 163, 215–230.
- Fan, M., Paul, D., Lee, T. C. and Matsuo, T. (2018). A multi-resolution model for non-Gaussian random fields on a sphere with application to ionospheric electrostatic potentials. *Ann. Appl. Stat.*, 12, 459–489.
- Friederichs, P. and Thorarindottir, T. L. (2012). Forecast verification for extreme value distributions with an application to probabilistic peak wind prediction. *Environmetrics*, 23, 579–594.
- Geweke, J. and Amisano, G. (2010). Comparing and evaluating Bayesian predictive distributions of asset returns. *Int. J. Forecast.*, 26, 216–230.
- Geweke, J. and Amisano, G. (2011). Hierarchical Markov normal mixture models with applications to financial asset returns. *J. Appl. Econometrics*, 26, 1–29.
- Giannone, D., Lenza, M. and Primiceri, G. E. (2015). Prior selection for vector autoregressions. *Rev. Econ. Stat.*, 97, 436–451.
- Gneiting, T. and Ranjan, R. (2011). Comparing density forecasts using threshold- and quantile-weighted scoring rules. *J. Bus. Econom. Statist.*, 29, 411–422.
- Groen, J. J. J., Paap, R. and Ravazzolo, F. (2013). Real-time inflation forecasting in a changing world. *J. Bus. Econom. Statist.*, 31, 29–44.
- Gschlößl, S. and Czado, C. (2007). Spatial modelling of claim frequency and claim size in non-life insurance. *Scand. Actuar. J.*, 2007, 202–225.
- Held, L., Meyer, S. and Bracher, J. (2017). Probabilistic forecasting in infectious disease epidemiology: The 13th Armitage lecture. *Stat. Med.*, 36, 3443–3460.
- Jensen, M. J. and Maheu, J. M. (2010). Bayesian semiparametric stochastic volatility modeling. *J. Econometrics*, 157, 306–316.
- Jensen, M. J. and Maheu, J. M. (2013). Bayesian semiparametric multivariate GARCH modeling. *J. Econometrics*, 176, 3–17.
- Jensen, M. J. and Maheu, J. M. (2014). Estimating a semiparametric asymmetric stochastic volatility model with a Dirichlet process mixture. *J. Econometrics*, 178, 523–538.
- Jin, X. and Maheu, J. M. (2013). Modeling realized covariances and returns. *J. Financ. Economet.*, 11, 335–369.
- Jin, X. and Maheu, J. M. (2016). Bayesian semiparametric modeling of realized covariance matrices. *J. Econometrics*, 192, 19–39.
- Jochmann, M., Koop, G. and Strachan, R. W. (2010). Bayesian forecasting using stochastic search variable selection in a VAR subject to breaks. *Int. J. Forecast.*, 26, 326–347.
- Kallache, M., Maksimovich, E., Michelangeli, P.-A. and Naveau, P. (2010). Multimodel combination by a Bayesian hierarchical model: Assessment of ice accumulation over the oceanic Arctic region. *J. Clim.*, 23, 5421–5436.
- Kastner, G. (2016). Dealing with stochastic volatility in time series using the R package stochvol. *J. Stat. Softw.*, 69, 1–30.
- Krüger, F., Clark, T. E. and Ravazzolo, F. (2017). Using entropic tilting to combine BVAR forecasts with external nowcasts. *J. Bus. Econom. Statist.*, 35, 470–485.
- Krüger, F. and Nolte, I. (2016). Disagreement versus uncertainty: Evidence from distribution forecasts. *J. Bank. Finance*, 72, 172–186.
- Leão, W. L., Abanto-Valle, C. A. and Chen, M.-H. (2017). Bayesian analysis of stochastic volatility-in-mean model with leverage and asymmetrically heavy-tailed error using generalized hyperbolic skew Student’s *t*-distribution. *Stat. Its Interface*, 10, 529.
- Leininger, T. J., Gelfand, A. E., Allen, J. M. and Silander Jr, J. A. (2013). Spatial regression modeling for compositional data with many zeros. *J. Agric. Biol. Environ.*

- Stat.*, 18, 314–334.
- Li, F., Villani, M. and Kohn, R. (2010). Flexible modeling of conditional distributions using smooth mixtures of asymmetric Student t densities. *J. Statist. Plann. Inference*, 140, 3638–3654.
- Liu, C. and Maheu, J. M. (2009). Forecasting realized volatility: A Bayesian model-averaging approach. *J. Appl. Econometrics*, 24, 709–733.
- Liu, J. and Maheu, J. M. (2018). Improving markov switching models using realized variance. *J. Appl. Econometrics*, 33, 297–318.
- Lopes, H. F., Salazar, E. and Gamerman, D. (2008). Spatial dynamic factor analysis. *Bayesian Anal.*, 3, 759–792.
- Louzis, D. P. (2019). Steady-state modeling and macroeconomic forecasting quality. *J. Appl. Econometrics*, 34, 285–314.
- Maheu, J. M. and Gordon, S. (2008). Learning, forecasting and structural breaks. *J. Appl. Econometrics*, 23, 553–583.
- Maheu, J. M. and McCurdy, T. H. (2009). How useful are historical data for forecasting the long-run equity return distribution? *J. Bus. Econom. Statist.*, 27, 95–112.
- Maheu, J. M., McCurdy, T. H. and Song, Y. (2012). Components of bull and bear markets: Bull corrections and bear rallies. *J. Bus. Econom. Statist.*, 30, 391–403.
- Maheu, J. M. and Song, Y. (2014). A new structural break model, with an application to Canadian inflation forecasting. *Int. J. Forecast.*, 30, 144–160.
- Maheu, J. M. and Song, Y. (2018). An efficient Bayesian approach to multiple structural change in multivariate time series. *J. Appl. Econometrics*, 33, 251–270.
- Maheu, J. M. and Yang, Q. (2016). An infinite hidden Markov model for short-term interest rates. *Journal of Empirical Finance*, 38, 202–220.
- Maneesoonthorn, W., Martin, G. M., Forbes, C. S. and Grose, S. D. (2012). Probabilistic forecasts of volatility and its risk premia. *J. Econometrics*, 171, 217–236.
- Metaxoglou, K., Pettenuzzo, D. and Smith, A. (2019). Option-implied equity premium predictions via entropic tilting. *J. Financ. Economet.*, 17, 559–586.
- Panagiotelis, A. and Smith, M. (2008). Bayesian density forecasting of intraday electricity prices using multivariate skew t distributions. *Int. J. Forecast.*, 24, 710–727.
- Riebler, A., Held, L. and Rue, H. (2012). Estimation and extrapolation of time trends in registry data — Borrowing strength from related populations. *Ann. Appl. Stat.*, 6, 304–333.
- Risser, M. D. and Calder, C. A. (2015). Regression-based covariance functions for nonstationary spatial modeling. *Environmetrics*, 26, 284–297.
- Risser, M. D., Calder, C. A., Berrocal, V. J. and Berrett, C. (2019). Nonstationary spatial prediction of soil organic carbon: Implications for stock assessment decision making. *Ann. Appl. Stat.*, 13, 165–188.
- Rodrigues, L. R. L., García-Serrano, J. and Doblas-Reyes, F. (2014). Seasonal forecast quality of the West African monsoon rainfall regimes by multiple forecast systems. *J. Geophys. Res. Atmos.*, 119, 7908–7930.
- Sahu, S. K., Bakar, K. S. and Awang, N. (2015). Bayesian forecasting using spatio-temporal models with applications to ozone levels in the eastern United States. In *Geometry Driven Statistics* (I. L. Dryden and J. T. Kent, eds.), chap. 13. John Wiley & Sons, 260–281.
- Salazar, E., Sansó, B., Finley, A. O., Hammerling, D., Steinsland, I., Wang, X. and Delamater, P. (2011). Comparing and blending regional climate model predictions for the American Southwest. *J. Agric. Biol. Environ. Stat.*, 16, 586–605.
- Sigrist, F., Künsch, H. R. and Stahel, W. A. (2012). A dynamic nonstationary spatio-temporal model for short term prediction of precipitation. *Ann. Appl. Stat.*, 6, 1452–

- Sigrist, F., Künsch, H. R. and Stahel, W. A. (2015). Stochastic partial differential equation based modelling of large space-time data sets. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 77, 3–33.
- Smith, M. S. and Vahey, S. (2016). Asymmetric density forecasting of U.S. macroeconomic variables. *J. Bus. Econom. Statist.*, 34, 416–434.
- Tran, M.-N., Pitt, M. K. and Kohn, R. (2016). Adaptive Metropolis–Hastings sampling using reversible dependent mixture proposals. *Stat. Comput.*, 26, 361–381.
- Trombe, P.-J., Pinson, P. and Madsen, H. (2012). A general probabilistic forecasting framework for offshore wind power fluctuations. *Energies*, 5, 621–657.
- Warne, A., Coenen, G. and Christoffel, K. (2016). Marginalized predictive likelihood comparisons of linear Gaussian state-space models with applications to DSGE, DSGE-VAR and VAR models. *J. Appl. Econometrics*, 32, 103–119.
- White, P. A., Gelfand, A. E., Rodrigues, E. R. and Tzintzun, G. (2019a). Pollution state modelling for Mexico City. *J. R. Stat. Soc. Ser. A. Stat. Soc.*, 182, 1039–1060.
- White, P. A. and Porcu, E. (2019). Nonseparable covariance models on circles cross time: A study of Mexico City ozone. *Environmetrics*, 30, e2558.
- White, P. A., Reese, C. S., Christensen, W. F. and Rupper, S. (2019b). A model for Antarctic surface mass balance and ice core site selection. *Environmetrics*, 30, e2579.
- Wijeyakulasuriya, D. A., Hanks, E. M., Shaby, B. A. and Cross, P. C. (2018). Extreme value-based methods for modeling elk yearly movements. *J. Agric. Biol. Environ. Stat.*, 24, 73–91.
- Zhou, Z., Matteson, D. S., Woodard, D. B., Henderson, S. G. and Micheas, A. C. (2015). A spatio-temporal point process model for ambulance demand. *J. Amer. Statist. Assoc.*, 110, 6–15.