Analyzing multilevel experiments in the presence of peer effects*

Guillaume Basse Harvard Avi Feller UC Berkeley

October 15, 2018

Abstract

Multilevel or two-stage randomization is a powerful design for estimating treatment effects in the presence of social interactions. Our motivating example is a multilevel randomized trial evaluating an intervention to reduce student absenteeism in the School District of Philadelphia. In that experiment, households with multiple students were first assigned to treatment or control; then, in treated households, one student was randomly assigned to treatment. Using this example, we highlight key considerations for analyzing multilevel experiments in practice. Our first contribution is to address additional complexities that arise when household sizes vary; in this case, researchers must decide between assigning equal weight to households or equal weight to individuals. We propose unbiased estimators for a broad class of individual- and household-weighted estimands, with corresponding theoretical and estimated variances. Our second contribution is to connect two common approaches for analyzing multilevel designs: linear regression and randomization inference. We show that, with suitably chosen standard errors, these two approaches yield identical point and variance estimates, which is somewhat surprising given the complex randomization scheme. Finally, we explore options for incorporating covariates to improve precision and confirm our analytic results via simulation studies. We apply these methods to the attendance study and find large, substantively meaningful spillover effects.

Key Words: Multilevel randomization; randomization inference; causal inference under interference; student attendance.

1 Introduction

Multilevel randomization is a powerful design for estimating causal effects in the presence of social interactions. A typical multilevel design has two stages. First, whole clusters (e.g., households, schools, or graph partitions) are assigned to treatment or control. Second, units within each treated cluster are randomly assigned to treatment or control, as if each treated cluster were a separate, individually-randomized experiment. This design allows researchers to assess peer effects either by comparing untreated units in treated clusters with pure control units in control clusters or by comparing units across clusters with different proportions assigned to treatment (Hudgens and Halloran, 2008).

^{*}Email: afeller@berkeley.edu. We thank Peter Aronow, Peng Ding, Winston Lin, Joel Middleton, Caleb Miles, Luke Miratrix, James Pustejovsky, Todd Rogers, Shruthi Subramanyam, John Ternovksi, and Elizabeth Tipton for helpful comments and discussion. We also thank the excellent research partners at the School District of Philadelphia, especially Adrienne Reitano and Tonya Wolford.

Our motivating example is a large randomized evaluation of an intervention targeting student absenteeism among elementary and high school students in the School District of Philadelphia (Rogers and Feller, 2016). In the original study, parents of at-risk students were randomly assigned to a direct mail intervention with tailored information about their students' attendance over the course of the year. In treated households with multiple eligible students, one student was selected at random to be the subject of the mailings, following a multilevel randomization. Substantively, this is a rare opportunity to study intra-household dynamics around student behavior. Methodologically, this presents a rich test case for understanding how to analyze multilevel experiments in practice.

There has been substantial interest in multilevel randomization in recent years, with prominent examples in economics (Crépon et al., 2013), education (Somers et al., 2010), political science (Sinclair et al., 2012), and public health (Hudgens and Halloran, 2008), as well as closely related variants in the context of large-scale social networks (Ugander et al., 2013). Such designs have become especially common in development economics; see Baird et al. (2014) and Angelucci and Di Maro (2016). There is also a small but growing methodological literature on analyzing multilevel experiments, including Hudgens and Halloran (2008), Liu and Hudgens (2014), and Rigdon and Hudgens (2015) in statistics; and Sinclair et al. (2012) and Baird et al. (2014) in the social sciences.

We build on this literature by addressing three practical issues that arise in analyzing the attendance study. First, school districts are typically interested in the intervention's overall impact on students rather than on households; that is, districts give equal weight to each individual rather than equal weight to each household. Existing approaches, however, focus either on equal weights for households (e.g., Hudgens and Halloran, 2008) or side-step the issue by assuming households are of equal size (e.g., Baird et al., 2014). We propose unbiased estimators for a broad class of individual- and household-weighted estimands, with corresponding theoretical and estimated variances. We also derive the bias of a simple difference in means for estimating individual-weighted estimands.

Second, we connect two common approaches for analyzing multilevel designs: linear regression, which is more common in the social sciences, and randomization inference, which is more common in epidemiology and public health. We show that, with suitably chosen standard errors, regression and randomization inference yield identical point and variance estimates. These results hold for a broad class of weighted estimands. We believe this equivalence will be important in practice, since the vast majority of applied papers in this area take a "regression first" approach to analysis (e.g., Baird et al., 2014) that can obfuscate key inferential

¹One important exception is Sinclair et al. (2012), who briefly address this issue.

issues.

Lastly, we explore options for incorporating covariates to improve precision, with a focus on poststratification and model-assisted estimation. We then confirm our analytic results via simulation studies and apply these methods to the attendance project. Overall, we find strong evidence of a spillover effect that is (depending on the scale of the outcome) roughly 60 to 80 percent as large as the primary effect. This holds across different estimands as well as with and without covariate adjustment.

This paper proceeds as follows. Section 2 defines the multilevel randomization, sets up the notation, and discusses the relevant assumptions. Section 3 defines estimands and corresponding estimators and variances in the case of equal-sized households. Section 4 addresses additional complications that arise when household sizes vary. Section 5 demonstrates how we can use regression with appropriate standard errors to obtain the randomization-based estimators. Section 6 explores covariate adjustment. Section 7 reports the results of extensive simulation studies. Section 8 analyzes the student attendance experiment. Section 9 concludes and offers directions for future work. The supplementary materials contains additional technical material and all proofs.

2 Setup and assumptions

We now review the setup and assumptions for a multilevel experiment in the presence of peer effects. The discussion closely follows Hudgens and Halloran (2008). We modify their terminology slightly to better fit our applied example and to recognize some small differences in emphasis with the social science literature. We begin with a description of multilevel designs and then turn to the relevant assumptions. For additional reviews on causal inference under interference, see, among others, Sinclair et al. (2012); Aronow and Samii (2013); Bowers et al. (2013); VanderWeele et al. (2014); Athey and Imbens (2016).

2.1 Defining multilevel experimental designs

Multilevel randomization is a special case of a multi-stage, nested randomization (see, e.g., Hudgens and Halloran, 2008; Hinkelmann and Kempthorne, 2012). First, a multilevel design requires a nested structure of at least two levels, such as individuals nested within households. Second, random assignment in a multilevel design proceeds sequentially, starting with the highest level; that is, the second-stage random assignment (at the lower level) depends on the realized first-stage random assignment (at the higher level).

For the sake of exposition and unless otherwise noted, we restrict our attention to a special case of a




Figure 1: Schematic of the multilevel randomized design we consider.

multilevel design that both highlights key features of the problem and reflects our motivating example. In particular, we consider a setting with only two nested levels, which we generally refer to as individuals nested within households. We assume that there are N households, i = 1, ..., N, with n_i individuals within each household and $n^+ \equiv \sum n_i$ individuals overall. To be consistent with the existing literature on multilevel experiments, we use the double-index notation, such that \cdot_{ij} denotes the individual j in household i.

Figure 1 gives a schematic overview of the multilevel design we consider. First, N_1 of N households are randomly assigned to treatment via complete randomization. Formally, let $\mathbf{H} = (H_1, \dots, H_N)$ be the vector of treatment assignments at the household level, such that $H_i = 1$ if household i is assigned to treatment and $H_i = 0$ otherwise. Second, separately for each household assigned to treatment, that is, among the set $\{i: H_i = 1\}$, randomly assign one of out n_i individuals to treatment via complete randomization within each household. For household i, let $\mathbf{Z}_i = (Z_{i1}, \dots, Z_{in_i})$ denote the assignment vector for the n_i units in that household, where Z_{ij} denotes whether the j^{th} individual in household i is assigned to treatment. Similarly, define $\mathbf{Z}_{i,-j}$ as the sub vector of \mathbf{Z}_i that exclude the j^{th} value. Finally, we aggregate all household-level assignments via $\mathbf{Z} = \{\mathbf{Z}_1, \dots, \mathbf{Z}_N\}$. Note that all individuals assigned to treatment are in households assigned to treatment; that is, if $H_i = 0$ then $Z_{ij} = 0$ for all individuals in household i.

2.2 Potential outcomes and relaxing SUTVA

We use the potential outcomes framework (Neyman, 1923; Rubin, 1974) to define our causal effects of interest. A key complication is that we cannot simply invoke the Stable Unit Treatment Value Assumption (SUTVA; Rubin, 1980). This assumption has two main parts: there are no hidden versions of the treatment, and there is no interference between units. We retain the first assumption, but relax the second. Let Y_{ij}^{obs} denote the observed outcome for individual j in household i. In general, let $Y_i(Z) = (Y_{i1}(Z), ..., Y_{in_i}(Z))$ be the vector of potential outcomes for household i, and $Y(Z) = \{Y_1(Z), ..., Y_N(Z)\}$ be the list of potential outcome vectors for all households.

At this stage, practical inference is infeasible without imposing additional restrictions on the structure of potential outcomes (Sobel, 2006; Hudgens and Halloran, 2008). Many such restrictions are possible. Aronow and Samii (2013) address the general case; Toulis and Kao (2013) focus on estimation with specific network structures (see also Tchetgen Tchetgen and VanderWeele, 2012). Given the data structure in our motivating example, we focus on two types of interference: between-household and within-household interference.

2.2.1 Between-household interference

First, we assume that there is no between-household interference. Sobel (2006) calls this partial interference; Athey and Imbens (2016) refer to this setting as a randomized experiment "with interactions in subpopulations."

Assumption 1 (No Interference Across Household). Interference occurs only within a household. That is, $Y_i(\mathbf{Z}) = Y_i(\mathbf{Z}_i)$.

This is effectively a "between household SUTVA" assumption and greatly reduces the complexity of the problem. At the same time, Assumption 1 remains quite strong in practice, especially in settings where spillover is an important consideration. In the context of the attendance study, this assumption states that students in different households do not affect each others' attendance. This assumption is violated if, for instance, friends skip school together. In theory, we could collect additional data on student social networks and other relationships and impose a network structure on the between-cluster interference; in practice, this was not possible. Nonetheless, we view spillovers within households as far more important than spillovers between households and thus consider Assumption 1 to be a useful approximation. See Sobel (2006), Hudgens and Halloran (2008), and Baird et al. (2014) for additional discussion.

2.2.2 Within-household interference

Even with the simplifying Assumption 1, the possible potential outcomes remain maddeningly complex. To illustrate this point, consider three siblings in the same household: Alice, Bob, and Carl. We are interested in Alice's potential outcome if one of her brothers (i.e., either Bob or Carl) is randomized to treatment. With no additional assumptions on interference, Alice has two different potential outcomes depending on which brother is treated, Y_{Alice} (Bob treated) or Y_{Alice} (Carl treated). That is, the notation allows for Alice to have different responses depending on the precise identity of the treated brother.

Sobel (2006) and Hudgens and Halloran (2008) introduced the idea of an *individual-level average potential* outcome in this setting. Define Alice's individual-level average potential outcome if one of her brothers is treated as

$$Y_{\text{Alice}}(\text{either brother treated}) = \frac{1}{2}Y_{\text{Alice}}(\text{Bob treated}) + \frac{1}{2}Y_{\text{Alice}}(\text{Carl treated}).$$

Importantly, individual-level average potential outcomes are functions both of identity-specific potential outcomes and of the exact randomization scheme; the above definition implicitly assumes that each brother has equal probability of receiving treatment. This raises both philosophical and practical issues. As we note below, calculating variances for such individual-level average potential outcomes proves especially difficult. For additional discussion, see Hudgens and Halloran (2008); Tchetgen Tchetgen and VanderWeele (2012); Aronow and Samii (2013); Manski (2013a).

As with between-household interference, one solution is to impose additional structure on the problem. For example, Paluck et al. (2016) collect detailed social network data for students within schools; they assume no interference between schools but leverage the within-school network structure for inference. This approach, however, is infeasible in settings without detailed data. In the attendance study, it is difficult to imagine collecting data on which individuals within the household are more influential than others; in other words, we cannot simply go ask Alice. Instead, we implicitly assume that each individual is equally influential for Alice's potential outcome.

A sensible restriction is therefore to assume that potential outcomes only depend on the number (or, equivalently, proportion) of individuals assigned to treatment within each household. Hudgens and Halloran (2008) call this the *stratified interference* assumption, which states that the precise identity of the treated individual in the treated cluster does not matter for untreated individuals in the same cluster (see also Manski, 2013b). That is, Alice's potential outcome is the same if Bob is treated or if Carl is treated, $Y_{\text{Alice}}(\text{Bob treated}) = Y_{\text{Alice}}(\text{Carl treated})$.

Assumption 2 (Stratified Interference).

$$Y_{ij}(\mathbf{Z}_{i,-j}, Z_{ij} = 0) = Y_{ij}(\mathbf{Z}'_{i,-j}, Z_{ij} = 0) \quad \forall \mathbf{Z}_{i,-j}, \mathbf{Z}'_{i,-j} \quad s.t. \quad \sum Z_{ij} = \sum Z'_{ij} = 1$$
 (1)

This assumption has two important ramifications. First, we no longer require individual-level average potential outcomes, since the potential outcomes that correspond to Bob or Carl received treatment are the same. This simplifies some conceptual challenges and breaks the dependence of the potential outcomes on the precise experimental design. Second, this greatly simplifies variance calculations, as we discuss below. This assumption is violated if Alice behaves differently when her parents receive information about Bob rather than Carl.

2.2.3 Connecting observed and potential outcomes

Given these assumptions, we now define potential outcomes for each individual j in household i, $Y_{ij}(H_i = h, Z_{ij} = z)$. There are three possible combinations: $Y_{ij}(1,1)$, $Y_{ij}(1,0)$, and $Y_{ij}(0,0)$. We regard these potential outcomes as fixed and define the observed outcome as a deterministic function of the treatment assignment and potential outcomes:

$$Y_{ij}^{\text{obs}} = H_i Z_{ij} Y_{ij}(1,1) + H_i (1 - Z_{ij}) Y_{ij}(1,0) + (1 - H_i) Y_{ij}(0,0),$$

where the randomness is entirely due to H and Z. That is, unless otherwise noted, all expectations and variances are with respect to the randomization distribution; inference is fully justified by the randomization itself (Fisher, 1935). Finally, we introduce the sets $\mathcal{T}_{hz} = \{(i,j) : H_i = h \text{ and } Z_{ij} = z\}$ to denote the set of households and individuals who are assigned to $H_i = h$ and $Z_{ij} = z$.

3 Inference with equal-sized households

Next, we review existing results for inference when all households or clusters are of equal size, turning to the more general case in Section 4. Starting with this simplified setting allows us to clarify the key points before addressing the additional complications of varying household size. These results essentially follow Hudgens and Halloran (2008). We begin with a discussion of the estimands of interest, the primary and spillover effects, and derive the corresponding unbiased estimators. We then give both their variances as well as conservative estimators of these variances. Readers familiar with this literature can skip to Section 4.

3.1 Estimands

We first assume that all households are of the same size. That is, $n_i = n$ for all households i = 1, ..., N, where N_1 and N_0 denote the number of households assigned to treatment and control, respectively. Unless otherwise noted, all estimands we consider here are finite sample estimands; that is, they are defined for the units in our sample. For discussion, see Imbens and Rubin (2015).

We define two main estimands, one for the primary effect, and one for the spillover effect. First, we define two average potential outcomes (averaging across units):

$$\overline{Y}_i(h,z) = \frac{1}{n} \sum_{i=1}^n Y_{ij}(h,z), \qquad \overline{Y}(h,z) = \frac{1}{N} \sum_{i=1}^N \overline{Y}_i(h,z) = \frac{1}{nN} \sum_{i=1}^N \sum_{j=1}^n Y_{ij}(h,z),$$

the average potential outcome for $H_i = h$ and $Z_{ij} = z$ for cluster i and for the entire finite population, respectively.

Definition 1 (Estimands with equal-sized households). Define the average primary effect as follows:

$$\tau^{P} = \frac{1}{Nn} \sum_{i}^{N} \sum_{j}^{n} (Y_{ij}(1,1) - Y_{ij}(0,0)) = \frac{1}{N} \sum_{i}^{N} (\overline{Y}_{i}(1,1) - \overline{Y}_{i}(0,0)) = \overline{Y}(1,1) - \overline{Y}(0,0), \tag{2}$$

and the average spillover effect as:

$$\tau^{S} = \frac{1}{Nn} \sum_{i}^{N} \sum_{j}^{n} (Y_{ij}(1,0) - Y_{ij}(0,0)) = \frac{1}{N} \sum_{i}^{N} (\overline{Y}_{i}(1,0) - \overline{Y}_{i}(0,0)) = \overline{Y}(1,0) - \overline{Y}(0,0).$$
 (3)

Note that these estimands have various names in the literature. We take the terminology primary effect from Toulis and Kao (2013). Hudgens and Halloran (2008) call these quantities the total and indirect effects, respectively. Baird et al. (2014) use the term treatment on the uniquely treated for an analogous quantity to the primary effect. Finally, we note that many other estimands are possible. Halloran et al. (1991); Hudgens and Halloran (2008) define the direct and overall effects, which are essentially the impact of the within-household randomization and the impact of randomization on the entire household, respectively. We do not explore these estimands further here.

3.2 Estimators and their variances

We now turn to estimating these quantities and deriving their sampling variances. All proofs in this section closely follow Hudgens and Halloran (2008) and are special cases of the more general results from Section 4.

See supplementary materials for additional details.

First, we show that the difference-in-means estimators are unbiased.

Proposition 1 (Unbiasedness of difference-in-means estimators). The primary and spillover effect estimators $\hat{\tau}^P$ and $\hat{\tau}^S$,

$$\widehat{\tau}^{P} = \frac{1}{N_{1}} \sum_{(i,j) \in \mathcal{T}_{11}} Y_{ij}^{obs}(1,1) - \frac{1}{nN_{0}} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{obs}(0,0),$$

$$\widehat{\tau}^{S} = \frac{1}{(n-1)N_{1}} \sum_{(i,j) \in \mathcal{T}_{10}} Y_{ij}^{obs}(1,0) - \frac{1}{nN_{0}} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{obs}(0,0),$$

are unbiased for their corresponding estimands with respect to the randomization distribution. That is $\mathbb{E}[\hat{\tau}^P] = \tau^P$ and $\mathbb{E}[\hat{\tau}^S] = \tau^S$.

Next, we determine the theoretical variance of these estimators with respect to the randomization distribution. We first define several useful terms, effectively decomposing the overall variance of the potential outcomes into a within- and between-household variance. Let $\sigma_{i,hz}^2 = 1/n \sum_j (Y_{ij}(h,z) - \overline{Y}_i(h,z))^2$ be the within-household potential outcome variances for $Y_{ij}(h,z)$, and let $\Sigma_{11} = 1/N \sum_i \sigma_{i,11}^2$ and $\Sigma_{10} = 1/N \sum_i \frac{1}{(n-1)^2} \sigma_{i,10}^2$ be the (re-scaled) average within-cluster variances for $Y_{ij}(1,1)$ and $Y_{ij}(1,0)$ respectively. Finally, define the between-cluster variance of cluster-level averages:

$$V_{hz} = \frac{1}{N-1} \sum_{i}^{N} (\overline{Y}_{i}(h,z) - \overline{Y}(h,z))^{2},$$

$$V_{P} = \frac{1}{N-1} \sum_{i}^{N} ([\overline{Y}_{i}(1,1) - \overline{Y}_{i}(0,0)] - [\overline{Y}(1,1) - \overline{Y}(0,0)])^{2},$$

$$V_{S} = \frac{1}{N-1} \sum_{i}^{N} ([\overline{Y}_{i}(1,0) - \overline{Y}_{i}(0,0)] - [\overline{Y}(1,0) - \overline{Y}(0,0)])^{2},$$

where V_{hz} is the between-cluster variance of the average cluster-level potential outcome, $\overline{Y}_i(h,z)$. V_P and V_S are the (unidentifiable) cluster-level treatment effect variation for the primary and spillover effects, respectively.

Proposition 2 (Theoretical variance of the estimators). The estimators have the following variance under the randomization distribution:

$$Var(\widehat{\tau}^P) = \frac{\Sigma_{11} + V_{11}}{N_1} + \frac{V_{00}}{N_0} - \frac{V_P}{N}$$

and

$$Var(\widehat{\tau}^S) = \frac{\Sigma_{10} + V_{10}}{N_1} + \frac{V_{00}}{N_0} - \frac{V_S}{N}.$$

This variance has the same form as the standard Neymanian variance. However, the increased variance due to the two-level randomization is reflected in the first numerator, which has two terms instead of one. Intuitively, this is a decomposition of the marginal variance of potential outcomes into Σ_{hz} , the average of the within-household variances, and V_{hz} , the variance of the household-level average potential outcomes. It is important to note that the theoretical variance becomes much more complex without the stratified interference assumption. See Tchetgen Tchetgen and VanderWeele (2012), Aronow and Samii (2013), and Rigdon and Hudgens (2015).

Finally, we can obtain an estimated variance that is a "conservative" estimate for the true variance (in the sense of being too wide in expectation) with respect to the randomization distribution. Let s_{hz}^2 be the cluster-level sample variance for the cluster-level average potential outcomes, $\overline{Y}_i^{\text{obs}}(h,z)$. That is,

$$s_{hz}^2 = \frac{1}{N_h - 1} \sum_{i=1}^{N} \mathbb{1}(H_i = h) \left(\overline{Y}_i^{\text{obs}}(h, z) - \overline{Y}^{\text{obs}}(h, z) \right)^2,$$

where $\overline{Y}^{\text{obs}}(h,z)$ is the average observed outcome for the set \mathcal{T}_{hz} , and where $\overline{Y}_{i}^{\text{obs}}(h,z)$ is the average observed outcome for the set \mathcal{T}_{hz} in household i.

Theorem 3.1 (Estimated variance of the estimators). Consider the variance estimators $\widehat{Var}(\widehat{\tau}^P)$ and $\widehat{Var}(\widehat{\tau}^S)$:

$$\widehat{Var}(\widehat{\tau}^P) = \frac{s_{11}^2}{N_1} + \frac{s_{00}^2}{N_0}, \tag{4}$$

$$\widehat{Var}(\widehat{\tau}^S) = \frac{s_{10}^2}{N_1} + \frac{s_{00}^2}{N_0}.$$
 (5)

The proposed estimators are conservative estimates of their respective estimands. That is, $\mathbb{E}(\widehat{Var}(\widehat{\tau}^P)) \geq Var(\widehat{\tau}^P)$ and $\widehat{Var}(\widehat{\tau}^P)$ and $\widehat{Var}(\widehat{\tau}^S)$ are unbiased if $V_P = 0$ and $V_S = 0$, respectively.

This result is due to Hudgens and Halloran (2008). As they show, the estimated variances are unbiased if the treatment effects are constant. There are several approaches to practical inference in this setting. For binary outcomes, Rigdon and Hudgens (2015) obtain exact confidence intervals via test inversion. For the general case, Liu and Hudgens (2014) demonstrate that an asymptotic regime in which the number of households grows large motivates the usual Wald-type confidence interval based on a Normal approximation.

We rely on that approach here, though note that the asymptotic approximation might have poor performance in small samples (see, for example, Pustejovsky and Tipton, 2016).

4 General results for varying household size

We now generalize these results to allow for varying household size. Doing so introduces both conceptual and technical complications. Broadly, there are now two types of estimands, household-weighted estimands ('HW') that assign equal weight to households, regardless of the number of individuals in each household; and individual-weighted estimands ('IW') that assign equal weight to individuals, regardless of the distribution across households. A substantial literature on cluster-randomized trials addresses related questions; see, among others, Donner and Klar (2000); Imai et al. (2009); Schochet (2013); Middleton and Aronow (2015).

There are many settings in which researchers might be more interested in IW estimands than HW estimands. In the attendance study we consider, district administrators are generally more concerned with raising the overall attendance rate than with raising the average attendance rate across households. Public health researchers administering treatment to villages of different sizes might similarly be interested in the overall effect on the population rather than on village-level averages, especially if the treatment is more effective in larger villages. And so on.

One important consideration is that researchers can typically estimate HW estimands more precisely than IW estimands. For example, consider the case of one massive household and many small households (e.g., Athey and Imbens, 2016). In that setting, inference for the IW estimand is difficult because units in the large cluster always receive the same assignment; inference for the HW estimand, however, is likely much more precise. We therefore recommend that researchers interested in IW estimands report both HW and IW estimates in practice.

We next describe results for a general class of weighted estimands, of which both HW and IW estimands are a special case. We then discuss several approaches for estimating IW estimands.

4.1 Inference for general weighted estimands

We now consider a class of weights we call *multilevel weights*; household and individual weights are both special cases. We sketch the key ideas here and defer the details to the supplementary materials.

Definition 2 (Multilevel weights). Define multilevel weights $w_i^{(00)}, w_i^{(10)}, w_i^{(11)}$ and w_i^* as any positive real

number such that:

$$w_i^{(11)} = \frac{1}{P(H_i = 1)} \frac{1}{P(Z_{ij} = 1 | H_i = 1)} w_i^*,$$

$$w_i^{(10)} = \frac{1}{P(H_i = 1)} \frac{1}{P(Z_{ij} = 0 | H_i = 1)} w_i^*,$$

$$w_i^{(00)} = \frac{1}{P(H_i = 0)} w_i^*.$$

These are standard Horvitz-Thompson weights modified for our two-stage randomization. Thus the multilevel weighted estimand and estimator for the primary effect are:

$$\tau_W^P = \sum_{i=1}^N w_i^* \sum_{j=1}^{n_i} (Y_{ij}(1,1) - Y_{ij}(0,0)) \text{ and } \widehat{\tau}_W^P = \sum_{(i,j) \in \mathcal{T}_{11}} w_i^{(11)} Y_{ij}^{\text{obs}}(1,1) - \sum_{(i,j) \in \mathcal{T}_{00}} w_i^{(00)} Y_{ij}^{\text{obs}}(0,0),$$

with analogous quantities for the spillover effect. In the supplementary materials, we show that these estimators are unbiased, derive their theoretical variances, and give conservative estimated variances. These are essentially the weighted generalization of the results in Section 3.

Finally, we show in Supplementary Material that $w_i^* = \frac{1}{Nn_i}$ corresponds to household-weighted estimands and $w_i^* = \frac{1}{n^+}$ corresponds to individual-weighted estimands. Reassuringly, plugging in the household weights exactly recovers the results of Hudgens and Halloran (2008). These estimands and estimators have a particularly simple form, corresponding to the natural difference-in-means estimators:

$$\tau_{HW}^{P} = \frac{1}{N} \sum_{i=1}^{N} \frac{1}{n_i} \sum_{j=1}^{n_i} (Y_{ij}(1,1) - Y_{ij}(0,0)) \quad \text{and} \quad \widehat{\tau}_{HW}^{P} = \frac{1}{N_1} \sum_{i \in \mathcal{T}_{11}} \overline{Y}_i^{\text{obs}}(1,1) - \frac{1}{N_0} \sum_{i \in \mathcal{T}_{00}} \overline{Y}_i^{\text{obs}}(0,0), \quad (6)$$

where $\overline{Y}_i^{\text{obs}}(h,z)$ are the household-level averages of the observed outcomes with $H_i=h$ and $Z_{ij}=z$.

4.2 Inference for individual-weighted estimands

Inference for the individual-weighted estimand requires some more care. The estimands have the following form:

$$\tau_{IW}^P = \frac{1}{n^+} \sum_{i=1}^{N} \sum_{j=1}^{n_i} (Y_{ij}(1,1) - Y_{ij}(0,0)) \quad \text{and} \quad \tau_{IW}^S = \frac{1}{n^+} \sum_{i=1}^{N} \sum_{j=1}^{n_i} (Y_{ij}(1,0) - Y_{ij}(0,0)),$$

where n^+ is the total number of individuals. We outline three approaches for estimating the quantities: unbiased estimation; simple difference; and stratification or post-stratification by household size. Finally, we suggest that researchers can mix-and-match these methods in practice.

4.2.1 Unbiased estimation

Applying Horvitz-Thompson weights yields unbiased estimators for τ_{IW}^P and τ_{IW}^S :

$$\widehat{\tau}_{IW}^{P} = \frac{1}{n^{+}} \frac{N}{N_{1}} \sum_{(i,j) \in \mathcal{T}_{11}} \frac{n_{i}}{1} Y_{ij}^{\text{obs}}(1,1) \qquad - \frac{1}{n^{+}} \frac{N}{N_{0}} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{\text{obs}}(0,0),$$

$$\widehat{\tau}_{IW}^S = \frac{1}{n^+} \frac{N}{N_1} \sum_{(i,j) \in \mathcal{T}_{10}} \frac{n_i}{n_i - 1} Y_{ij}^{\text{obs}}(1,0) - \frac{1}{n^+} \frac{N}{N_0} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{\text{obs}}(0,0),$$

with household-level assignment probabilities N_1/N and N_0/N and (conditional) individual-level probabilities $1/n_i$ and $(n_i-1)/n_i$. Thus, the primary effect estimator up-weights larger households while the spillover effect estimator down-weights larger households. As is common with Horvitz-Thompson estimators, unbiased estimation typically comes at the price of additional variance. In practice, researchers can often reduce this variance by first normalizing the weights (i.e., Hájek weights), which introduces some small bias. See the supplementary materials for more details on the Hájek estimator in this context.

4.2.2 Simple difference estimator

Contrast the unbiased estimator with a simple difference estimator; that is, the difference-in-means across individuals, ignoring households.

$$\widehat{\tau}_{sd}^{P} = \frac{1}{n_{11}^{+}} \sum_{(i,j) \in \mathcal{T}_{11}} Y_{ij}^{\text{obs}}(1,1) - \frac{1}{n_{00}^{+}} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{\text{obs}}(0,0)$$

$$\tag{7}$$

$$\widehat{\tau}_{sd}^{S} = \frac{1}{n_{10}^{+}} \sum_{(i,j) \in \mathcal{T}_{10}} Y_{ij}^{\text{obs}}(1,0) - \frac{1}{n_{00}^{+}} \sum_{(i,j) \in \mathcal{T}_{00}} Y_{ij}^{\text{obs}}(0,0), \tag{8}$$

where
$$n_{11}^+ = \sum_i \mathbb{1}(H_i = 1) \sum_j \mathbb{1}(Z_{ij} = 1)$$
, $n_{10}^+ = \sum_i \mathbb{1}(H_i = 1) \sum_j \mathbb{1}(Z_{ij} = 0)$, and $n_{00}^+ = \sum_i \mathbb{1}(H_i = 0)n_i$.

Despite its intuitive appeal, this estimator can be biased in practice. There are two main sources of bias. First, echoing results from Middleton and Aronow (2015), when household sizes vary, the quantities n_{11}^+ , n_{10}^+ , and n_{00}^+ are themselves random variables. Thus, both the numerator and denominator of each group average are random; and the mean of a ratio is not, in general, equal to the ratio of means. Second, individual-level treatment probabilities vary by household size; in the design we consider here, the probability of treatment assignment conditional on being in a treated household is $\mathbb{P}\{Z_{ij}=1 \mid H_i=1\}=1/n_i$. Thus, ignoring n_i —

and, by extension, the varying treatment probability—can lead to biased estimates. This is analogous to the possible bias when analyzing stratified randomized designs in which treatment probabilities vary across strata (see, e.g., Imbens and Rubin, 2015). We derive the exact form of the bias in the following proposition.

Proposition 3. The simple difference estimators, $\hat{\tau}_{sd}^P$ and $\hat{\tau}_{sd}^S$, defined in Equation 7 and 8, have the following bias for their respective estimands.

$$bias\left(\widehat{\tau}_{sd}^{P}\right) = \frac{1}{N\overline{n}} \sum_{i} \left(\frac{\overline{n}}{n_{i}} - 1\right) \sum_{i} Y_{ij}(1, 1) + \frac{1}{N_{0}\overline{n}} cov\left(\frac{\sum_{\mathcal{T}_{00}} Y_{ij}(0, 0)}{n_{00}^{+}}, n_{00}^{+}\right)$$
(9)

and

$$bias\left(\widehat{\tau}_{sd}^{S}\right) = \frac{1}{N\overline{n}} \sum_{i} \left(\frac{\frac{\overline{n}}{\overline{n}-1}}{\frac{n_{i}}{n_{i}-1}} - 1\right) \sum_{j} Y_{ij}(1,0) + \left(\frac{1}{N_{0}\overline{n}} cov\left(\frac{\sum_{\mathcal{T}_{00}} Y_{ij}(0,0)}{n_{00}^{+}}, n_{00}^{+}\right) - \frac{1}{N_{1}(\overline{n}-1)} cov\left(\frac{\sum_{\mathcal{T}_{10}} Y_{ij}(1,0)}{n_{10}^{+}}, n_{10}^{+}\right)\right).$$

$$(10)$$

If household size is constant, all of these terms are zero. If the covariance between household size and potential outcomes is zero, only the first term of each equation remains. In simulations in Section 7, we show that the overall bias can be large if household sizes vary and treatment effects also vary by household size. Lastly, researchers might choose to correct for only one of these sources of bias, for example by only adjusting for the varying probability of treatment within each treated household. Given the performance of the post-stratified estimator in Section 7, this seems unnecessary in our setting.

4.2.3 Stratification and post-stratification by household size

Finally, we consider stratification and post-stratification by household size. If household-level randomization is stratified by household size, inference for the individual-weighted estimand is immediate. In particular, let τ_k^P and τ_k^S be the stratum-specific estimands for the stratum with household size $n_i = k$,

$$\tau_k^P = \frac{1}{N^{(k)}} \sum_i^N \mathbbm{1}(n_i = k) \frac{1}{k} \sum_j^k (Y_{ij}(1, 1) - Y_{ij}(0, 0)) \quad \text{and} \quad \tau_k^S = \frac{1}{N^{(k)}} \sum_i^N \mathbbm{1}(n_i = k) \frac{1}{k} \sum_j^k (Y_{ij}(1, 0) - Y_{ij}(0, 0)),$$

where $N^{(k)}$ is the number of households of size k. Since household size is constant within each stratum, the corresponding household- and individual-weighted estimands are equivalent. We can therefore re-write the

overall individual-weighted estimands as weighted averages of the stratum-specific effects,

$$\tau_{IW}^P = \sum_{k=2}^K \frac{n^{(k)+}}{n^+} \tau_k^P \quad \text{ and } \quad \tau_{IW}^S = \sum_{k=2}^K \frac{n^{(k)+}}{n^+} \tau_k^S,$$

where $n^{(k)+} = \sum_{i: n_i = k} n_i$ and where we assume (without essential loss of generality) that household sizes range from k = 2, ..., K. To simplify variance calculations, we further assume that there are at least two households of each size. Due to stratification, we can regard each household size as a separate "mini experiment." The results of Section 3 therefore carry through essentially unchanged. That is, for the primary effect (with analogous results for the spillover effect):

$$\widehat{\tau}_{IW}^P = \sum_{k=2}^K \frac{n^{(k)+}}{n^+} \widehat{\tau}_k^P, \quad \text{and} \quad \operatorname{Var}(\widehat{\tau}_{IW}^P) = \sum_{k=2}^K \left(\frac{n^{(k)+}}{n^+}\right)^2 \operatorname{Var}(\widehat{\tau}_k^P), \tag{11}$$

where $\hat{\tau}_{IW}^P$ is an unbiased estimate of τ_{IW}^P , $\operatorname{Var}(\hat{\tau}_{IW}^P)$ is its sampling variance, and $\widehat{\operatorname{Var}}(\hat{\tau}_{IW}^P)$ is the corresponding conservative variance estimate. To modify the above results for household-weighted estimates, simply replace the weight $n^{(k)+}/n^+$ with $N^{(k)}/N$. In other words, weight each stratum by the number of households in that stratum, rather than the number of individuals. While stratification is not necessary to obtain unbiased estimates of household-weighted estimands, stratification will improve precision so long as household size is predictive of the outcome.

In practice, it is not always possible or feasible to stratify randomization by household size. This is especially true if there are important blocking factors other than size. For example, the multilevel randomization in the attendance study was merely one part of a broader experiment in which randomization was stratified by school and grade. Fortunately, researchers can often post-stratify by household size; that is, the researcher can analyze the experiment as if randomization had been stratified by size. In the supplementary materials, we extend the theoretical guarantees from Miratrix et al. (2013) to multilevel randomization, and include additional discussion of the technical details.

4.2.4 "Mix and match"

If there are relatively small samples or the distribution of household sizes varies widely, researchers might want to "mix and match" among possible strategies. As we discuss in Section 8, there are 3,109 households size of $n_i = 2$ but only two households of size $n_i = 7$. Thus, it is unreasonable to post-stratify precisely on household size. Instead, we post-stratify by dividing household size into $n_i \in \{2, 3, 4 - 7\}$, using the

unbiased IW estimator for households of size four to seven. This is inherently a bias-variance tradeoff and will depend on the particular context. Finally, we could also adjust for n_i via regression, as discussed in Section 6 (see also Middleton and Aronow, 2015). Of course, researchers should pre-specify such procedures whenever possible.

5 Regression-based estimation

We now connect these randomization-based results with more familiar regression-based methods. Our key result is that, with the appropriate standard errors, conventional linear regression estimates are equivalent to the randomization-based estimates. Importantly, this approach regards regression as a convenient tool (sometimes known as a *derived linear model*; see Hinkelmann and Kempthorne, 2012), and does not equate the regression with a specific generative model. In other words, this approach does not impose a model for the potential outcomes (Imbens and Rubin, 2015).

We consider two basic regression approaches, an individual-level regression and a household-level regression. For simplicity, we start with the equal-sized household case and then show that these results generalize to any multilevel weights. We then demonstrate the dangers of using standard errors that ignore the multilevel structure. While regression estimators of other estimands are possible, especially the direct and overall effects (Hudgens and Halloran, 2008), we do not pursue those here. Finally, these results build on recent advances in estimating robust and cluster-robust standard errors, beginning with McCaffrey et al. (2001) and Bell and McCaffrey (2002). See also Imbens and Kolesar (2012); Cameron and Miller (2015); Pustejovsky and Tipton (2016).

5.1 Individual-level regression

First, we construct the individual-level linear model,

$$Y_{ij}^{\text{obs}} = \alpha + \beta^P H_i Z_{ij} + \beta^S H_i (1 - Z_{ij}) + \varepsilon_{ij}. \tag{12}$$

Importantly, the uncertainty in ε_{ij} is entirely due to randomization. It is straightforward to show that standard OLS estimates for β^P and β^S are identical to the randomization-based estimators in Section 3.2. Estimating the variance, however, requires more care. We now show that the cluster-robust generalization of HC2 standard errors are equivalent to the randomization-based standard errors derived above.

Theorem 5.1. For equal-sized households, let Y be the vector containing all the observed outcomes Y_{ij}^{obs} ; let X denote the appropriate design matrix (formally defined in the supplementary materials) with columns corresponding to the intercept, HZ, and H(1-Z); and let $\beta = (\alpha, \beta^P, \beta^S)$. The linear model in Equation 12 can therefore be re-written as $Y = X\beta + \varepsilon$, with corresponding least squares estimate, $\widehat{\beta}^{ols}$. These estimates are unbiased for their corresponding estimands. Further, define the cluster-robust generalization of HC2 standard errors as:

$$\widehat{Var}_{hc2}^{clust}(\widehat{\beta}^{ols}) = (X^t X)^{-1} \sum_{s=1}^{S} X_s^t (I_{N_s} - P_{ss})^{-1/2} \widehat{\varepsilon}_s \widehat{\varepsilon}_s (I_{N_s} - P_{ss})^{-1/2} X_s (X^t X)^{-1}$$

where X_s and $\widehat{\varepsilon}_s$ are the subsets of X and $\widehat{\varepsilon}$ corresponding to household s, and P_{ss} is defined as $P_{ss} = X_s(X^tX)^{-1}X_s^t$. Then

$$\widehat{Var}_{hc2}^{clust}(\widehat{\beta}^{P,ols}) = \widehat{Var}(\widehat{\tau}^{P}) \quad \ and \quad \ \widehat{Var}_{hc2}^{clust}(\widehat{\beta}^{S,ols}) = \widehat{Var}(\widehat{\tau}^{S})$$

$$where \ \widehat{Var}_{hc2}^{\ clust}(\widehat{\beta}^{P,ols}) = (\widehat{Var}_{hc2}^{\ clust}(\widehat{\beta}^{ols}))_{22} \ \ and \ \widehat{Var}_{hc2}^{\ clust}(\widehat{\beta}^{S,ols}) = \widehat{Var}_{hc2}^{\ clust}(\widehat{\beta}^{ols}))_{33}.$$

In short, Theorem 5.1 confirms that we can obtain the same randomization-based point- and variance-estimators via the individual-level linear model in Equation 12 with HC2 cluster-robust standard errors. This is similar to results obtained with heteroskedastic-robust standard errors in simpler designs (e.g., Samii and Aronow, 2012; Imbens and Rubin, 2015). In Section 5.4, we demonstrate the effect of failing to account for clustering on standard errors. Researchers can estimate these standard errors directly in R via, for example, the clubSandwich package. See Pustejovsky and Tipton (2016) for additional discussion on the performance of clustered standard errors with a relatively small number of clusters.

5.2 Household-level regression

We now consider a regression at the household level. This is a common strategy in cluster-randomized trials (e.g., Athey and Imbens, 2016) and yields identical inference to individual-level regression with clustered standard errors (see, for example Cameron and Miller, 2015).

Aggregation here requires some care in that we separately aggregate treated and control units within each treated household. Thus, we consider three types of household-level aggregates for household i. Each treated household has two household-level averages, $\overline{Y}_i^{\text{obs}}(1,1)$ and $\overline{Y}_i^{\text{obs}}(1,0)$; each control household has one household-level average, $\overline{Y}_i^{\text{obs}}(0,0)$. We can therefore assemble a vector of household-average outcomes,

 $\overline{Y}_k^{\text{obs}}$ of length $2N_1 + N_0$. We introduce the indicators $H^{(11)}$ and $H^{(10)}$; $H_k^{(11)} = 1$ if the aggregate is over the treated units in treated households and $H_k^{(11)} = 0$ otherwise; $H_k^{(10)} = 1$ if the aggregate is over the control units in treated households and $H_k^{(10)} = 0$ otherwise. We then consider the following linear model:

$$\overline{Y}_{k}^{\text{obs}} = \alpha + \beta^{P} H_{k}^{(11)} + \beta^{S} H_{k}^{(10)} + \varepsilon_{k}'. \tag{13}$$

We now show that we can obtain the randomization-based point and variance estimates via the linear model estimates with standard (i.e., non-cluster) heteroskedastic-robust standard errors.

Theorem 5.2. For equal-sized households, the OLS estimates for β^P and β^S in Equation 13 are unbiased estimators of the corresponding estimands. Define the heteroskedastic-robust estimator of the variances:

$$\widehat{Var}_{hc2}(\widehat{\tau}^P) \equiv \widehat{Var}_{hc2}(\widehat{\beta}^P) = \frac{\sum_{k:H_k^{(11)}=0} \widehat{\varepsilon}_k^2 (H_k^{(10)} - \overline{H})^2}{(\sum_{k:H_k^{(10)}=0} (H_k^{(11)} - \overline{H}^{(11)})^2)^2}$$

and

$$\widehat{Var}_{hc2}(\widehat{\tau}^S) \equiv \widehat{Var}_{hc2}(\widehat{\beta}^S) = \frac{\sum_{k:H_k^{(11)}=0} \widehat{\varepsilon}_k^2 (H_k^{(10)} - \overline{H})^2}{(\sum_{k:H_k^{(11)}=0} (H_k^{(10)} - \overline{H}^{(10)})^2)^2}$$

We have:

$$\widehat{Var}_{hc2}(\widehat{\tau}^P) = \widehat{Var}(\widehat{\tau}^P) \quad and \quad \widehat{Var}_{hc2}(\widehat{\tau}^S) = \widehat{Var}(\widehat{\tau}^S),$$

where the $\hat{\varepsilon}_k$'s are the HC2 residuals (see supplementary materials for exact definition).

In short, Theorem 5.2 states that we can aggregate to the household level and proceed as if this were a standard completely randomized trial (Imbens and Rubin, 2015). Intuitively, the aggregation at the household level is another way of accounting for the household structure in the multilevel randomization scheme. Note that the definition for the heteroskedastic-robust estimator of the variance is entirely standard; it is straightforward to fit Equation 13 and the corresponding variance with common software, for example, the vcovHC function with the HC2 option in the sandwich package.

5.3 Results for weighted estimands

Finally, we can modify the household-level regression to estimate any multilevel weighted estimand. Thus, we can use this approach to obtain the HW and IW results in Section 4. The key idea is to run an unweighted regression on transformed outcomes.

Theorem 5.3. Let $w_i^{(00)}$, $w_i^{(10)}$, $w_i^{(11)}$ and w_i be multilevel weights. Let the transformed observed outcomes be $\tilde{Y}_{ij}^{obs}(1,1) = N_1 w_i^{(11)} Y_{ij}^{obs}(1,1)$, $\tilde{Y}_{ij}^{obs}(1,0) = N_1 (n_i - 1) w_i^{(10)} Y_{ij}^{obs}(1,0)$, and $\tilde{Y}_{ij}^{obs}(0,0) = N_0 n_i w_i^{(00)} Y_{ij}^{obs}(0,0)$. Then the results of Theorem 5.2 hold. That is: $\hat{\beta}^P = \hat{\tau}_W^P$, $\hat{\beta}^S = \hat{\tau}_W^S$, $\widehat{Var}_{hc2}(\hat{\beta}^P) = \widehat{Var}(\hat{\tau}_W^P)$, and $\widehat{Var}_{hc2}(\hat{\beta}^S) = \widehat{Var}(\hat{\tau}_W^S)$.

This approach is subtly different from using Weighted Least Squares, which would also reweight the design matrix.

5.4 Failing to account for clustering

It is instructive to consider the consequences of a "naive" analysis of a multilevel experiment that does not directly address the two-stage randomization. While there are many ways that such an analysis could fail, we consider the case of incorrectly analyzing a multilevel experiment as if it were a completely randomized experiment; that is, the researcher ignores the household structure. With equal-sized households, the point estimates will be the same as for the appropriate analysis, but the standard errors will differ. In particular, let $\widehat{\text{Var}}_{\text{hc2}}^{\text{het}}(\widehat{\tau}^P)$ be the (non-cluster) HC2 robust standard error for the primary effect; that is, these are the variances from Equation 13 for the household aggregates incorrectly applied to the individual level. We show in the supplementary materials that,

$$\mathbb{E}\left[\widehat{\operatorname{Var}}_{\operatorname{hc2}}^{\operatorname{het}}(\widehat{\tau}^{P})\right] - \operatorname{Var}(\widehat{\tau}^{P}) = (\Sigma_{00} + V_{00}) \left\{ \frac{1}{nN_{0} - 1} \left(1 - n\rho_{00}\right) - \frac{1}{N} \frac{V_{p}}{\Sigma_{00} + V_{00}} \right\},\,$$

where V_{00} and Σ_{00} are the between- and within-household variances, respectively, of the control potential outcomes, $Y_{ij}(0,0)$, and $\rho_{00} \equiv V_{00}/(\Sigma_{00} + V_{00})$ is the intraclass correlation. This quantity is negative—that is, the variance is anti-conservative in expectation—if:

$$\rho_{00} > \frac{1}{n} - \left(\frac{nN_0 - 1}{nN}\right) \left(\frac{V_P}{\Sigma_{00} + V_{00}}\right).$$

Since the last term is non-negative, we can build intuition by setting that term to zero. For example, consider the special case of $V_P = 0$, which would occur if there is a constant additive effect. Under these conditions, the estimated variance is anti-conservative if $\rho_{00} > 1/n$. In the social sciences, typical values of ICC range from 0.1 to 0.3 (e.g., Gelman and Hill, 2006). Thus, even with households of size 4 or 5, the estimated variance could be anti-conservative. We see this behavior in the simulations in Section 7.

6 Covariate adjustment

Finally, we explore how to incorporate individual- and cluster-level covariates in a multilevel experiment. There is an extensive literature on the use of covariates in randomized trials (see, for example, Imbens and Rubin, 2015). In this section, we briefly address stratification, post-stratification, and model-assisted estimation. In ongoing research, we are working to extend results from covariate adjustment in completely randomized designs (e.g., Lin, 2013) to cluster-randomized and multilevel experiments.

Stratification and post-stratification. As with household size in Section 4.2.3, the simplest way to account for covariates is to incorporate them into the randomization by stratifying on them. In general, the researcher can partition households into discrete strata, regard each stratum as a separate "mini experiment," and estimate an overall effect by averaging across strata. This will improve the precision of the treatment effect estimate so long as the stratifying covariate is predictive of the outcome. Note that researchers cannot stratify by a covariate that varies within household, as this could destroy the nested structure in the data by assigning different individuals in the same household to different "household-level" treatments. For example, we cannot stratify the multilevel randomization by gender, as some houses will have both boys and girls; instead, we could stratify by whether all students in the household are boys, which is an aggregate version of the individual-level covariate. Finally, as with household size, researchers can also post-stratify on household-level covariates. Of course, it is possible to combine stratification and post-stratification; for example, first stratifying by household size and then, for each household size, post-stratifying by whether the household speaks English as the primary language.

Model-assisted estimation. First developed in survey sampling (Cochran, 1977), model-assisted estimation has seen increased use in causal inference (Rosenbaum et al., 2002; Hansen and Bowers, 2009; Aronow and Middleton, 2013). The main idea is to modify a standard estimator by incorporating a so-called "working model." The researcher typically chooses this model such that, if certain conditions hold (e.g., if the linear model is the actual data generating process), the overall estimator enjoys optimality properties, such as efficiency; if these conditions do not hold, the estimator nonetheless has other guarantees, such as consistency. Rather than present an extensive discussion of model-assisted estimation, we outline a straightforward application here and refer interested readers to the above references. We use this approach in the analysis in Section 8.

Following the setup in Hansen and Bowers (2009), consider K covariates $x^{(1)}, \ldots, x^{(K)}$ (which typically

include a constant) with corresponding coefficient vector, $\gamma = (\gamma_1, \dots, \gamma_K)$, such that $r(\gamma) = \{r_{ij}(\{x^{(k)}\}_k, \gamma)\}$ for $i = 1, \dots, N$ and $j = 1, \dots, n_i$ is a function mapping covariates to predictions. To simplify notation, we will let $x = \{x^{(k)}\}_k$. In practice, the vector γ are typically coefficients from a linear regression. In this case, let

$$r_{ij}(x,\gamma) = \sum_{k=1}^{K} x_{ij}^{(k)} \gamma_k,$$

where x_{ij} is a vector of covariates associated with unit j in cluster i. Importantly, we regard $r_{ij}(x, \gamma)$ as fixed and known for all units, rather than estimated from the data. We then define the (residualized) potential outcome as:

$$e_{ij}^{\gamma}(h,z) = Y_{ij}(h,z) - r_{ij}(x,\gamma). \tag{14}$$

As with the corresponding potential outcomes, $Y_{ij}(h,z)$, the residualized potential outcomes, $e_{ij}^{\gamma}(h,z)$, are assumed to be fixed and are only observed if $H_i = h$ and $Z_{ij} = z$. We can then substitute $e_{ij}^{\gamma}(h,z)$ for $Y_{ij}(h,z)$ in defining the primary effect:

$$\begin{split} \tau^P &= \overline{Y}(1,1) - \overline{Y}(0,0) \\ &= \left[\overline{e}^{\gamma}(1,1) + \overline{r}(x,\gamma) \right] - \left[\overline{e}^{\gamma}(0,0) + \overline{r}(x,\gamma) \right] = \overline{e}^{\gamma}(1,1) - \overline{e}^{\gamma}(0,0), \end{split}$$

re-writing τ^S in an analogous way.

Given $r_{ij}(\gamma)$, model-assisted estimation of τ^P is immediate via substituting the observed values of the residualized outcomes, $e_{ij}^{\gamma,\text{obs}}$, in place of the unadjusted outcomes, Y_{ij}^{obs} . Importantly, the resulting difference-in-means estimator is unbiased regardless of the exact values of $r_{ij}(\gamma)$; that is, there is no need to appeal to a "correctly specified" linear model to obtain the particular coefficient vector, γ . So long as the covariates are predictive of the outcome, the variance of $e_{ij}^{\gamma,\text{obs}}$ will generally be smaller than the variance of Y_{ij}^{obs} , and the resulting model-assisted estimator will also have smaller estimated variance. This will not necessarily hold for extreme values of γ . See, Hansen and Bowers (2009) and Aronow and Middleton (2013) for additional discussion.

Finally, the above derivations assume that γ is fixed and known; in practice, we must find some way to determine γ . The most straightforward approach is to generate a random hold-out sample, and estimate γ via a regression of Y on X for this group.² While not always possible, the motivating example effectively has a hold-out sample that we can use for this purpose. An alternative approach is to use the same data

²This is sometimes known as the Williams approach. See Aronow and Middleton (2013).

twice; once for estimating the working model and once for estimating the effects of interest. This obviously introduces additional complications, including possible bias, and, if all the models are linear, is effectively the same as standard regression adjustment (Lin, 2013). We are exploring this approach in ongoing work.

7 Simulations

We now turn to the simulations, in which we investigate quantitatively two elements discussed in the paper. First we show that failing to account for the cluster structure lead to confidence intervals that potentially severely under-cover. Second, we compare the three estimators for the IW estimand discussed in Section 4.2.

7.1 Failing to account for the cluster structure

We start by assessing quantitatively the importance of accounting for the cluster structure. First, we show that the proposed randomization-based methods indeed have nominal, conservative coverage. Second, we show that methods that fail to account for clustering dramatically under-cover in a range of settings. To clarify exposition, we focus on the case with equal-sized households.

We generate the potential outcomes in two stages. First, we simulate household-level average potential outcomes via:

$$\overline{Y}_{i}(0,0) \sim N\left(\mu_{00}, \sigma_{\mu}^{2}\right)$$

$$\tau_{i}^{P} \sim N\left(\overline{\tau}^{P}, \sigma_{\tau^{P}}^{2}\right)$$

$$\tau_{i}^{S} \sim N\left(\overline{\tau}^{S}, \sigma_{\tau^{S}}^{2}\right),$$

where $\overline{Y}_i(1,1) = \overline{Y}_i(0,0) + \tau_i^P$ and $\overline{Y}_i(1,0) = \overline{Y}_i(0,0) + \tau_i^S$. Then we generate individual-level potential outcomes via $Y_{ij}(h,z) \sim N\left(\overline{Y}_i(h,z),\sigma_y^2\right)$, for each h and z. Across all simulations, we fix the mean potential outcomes, with $\mu_{00} = 2$, $\bar{\tau}^S = 0.7$, and $\bar{\tau}^P = 1.5$, and fixed household size at $n_i = 4$, constant across all households. For convenience, we also restrict the household-level variance terms to be equal to each other, such that $\sigma_{\mu} = \sigma_{\tau^P} = \sigma_{\tau^S} = \sigma_c$, where σ_c is the common standard deviation. Thus, we vary three main parameters, $N \in \{50, 100, 500, 1000\}$ (we always set $N_1 = N/2$) and $\sigma_c, \sigma_y \in \{0.1, 0.2, 0.3, 0.4, 0.5\}$. For each combination of parameters, we consider three methods: cluster-robust standard errors; non-cluster, robust standard errors; and nominal standard errors. For each, we compute the coverage of the associated 95% confidence interval, averaging over 4000 randomizations of the assignment vector.

Table 1: Average coverage of 95% confidence intervals across all simulations

	$\widehat{ au}^P$	$\widehat{ au}^S$
Cluster robust SEs	0.97	0.98
Non-cluster robust SEs	0.93	0.87
Nominal SEs	0.95	0.86




Figure 2: Coverage for 95% Confidence Intervals for clustered SEs, non-cluster robust SEs, and nominal SEs, with respect to the intraclass correlation of control potential outcomes, $\sigma_c^2/(\sigma_c^2 + \sigma_y^2)$.

Table 1 shows the overall coverage for 95% confidence intervals, averaged across all values of the simulation parameters. As expected, coverage with the cluster-robust standard errors is slightly larger than 95% coverage. By contrast, the non-cluster and nominal standard errors have below 95% coverage. First, this coverage pattern is quite stable across different sample sizes. At the same time, coverage strongly depends on the within- and between-household variances. Figure 2 shows the relationship between coverage and the intraclass correlation among control potential outcomes, defined as $\sigma_c^2/(\sigma_c^2 + \sigma_y^2)$ in this simulation. Consistent with the results in Section 5.4, the coverage for non-clustered standard errors grows increasingly poor as the ICC increases.

Table 2: Bias and SE for different estimators for the IW estimand over 2,000 replications. '—' denotes $< 10^{-3}$.

	hh size unco	rrelated with effect	hh size correlated with effect			
Estimator	Avg. bias	Monte Carlo SE	Avg. bias	Monte Carlo SE		
Primary effect						
Unbiased	_	0.11		0.07		
Simple difference	_	0.04	0.18	0.12		
Post-stratified	_	0.04	_	0.04		
Spillover effect						
Unbiased	_	0.12		0.07		
Simple difference	_	0.04	0.12	0.13		
Post-stratified		0.04	_	0.04		

7.2 Comparing the three estimators for the IW estimand

We now focus on the IW estimand and compare the unbiased estimator, the difference-in-means estimator, and the post-stratified estimator. We consider two scenarios: (a) when the treatment effects are uncorrelated with household size; and (b) when treatment effects are correlated with household size. The data generating process is the same as above, with a balanced household-level randomization with N=200 and $N_1=100$, and with fixed $\sigma_c=\sigma_y=0.3$. We generate households of size 2, 3 and 4 with equal probability, and introduce the parameters $\overline{\tau}_k^P, \overline{\tau}_k^S, \mu_{00}^{(k)}$ for k=2,3,4. For scenario (a), we set $\overline{\tau}_k^P=1.5, \overline{\tau}_k^S=0.7, \mu_{00}^{(k)}=2$ for all k=2,3,4. For scenario (b), we allow the effects to vary by household size, as follows: $\overline{\tau}_2^P=1.5, \overline{\tau}_3^P=0.75, \overline{\tau}_4^P=0.37, \overline{\tau}_2^S=0.7, \overline{\tau}_3^S=0.35, \overline{\tau}_4^S=0.17$, and $\mu_{00}^{(2)}=2, \mu_{00}^{(3)}=1, \mu_{00}^{(4)}=0.5$.

The results are presented in Table 2. We see that when the treatment effect is uncorrelated with household size, the bias of all three estimators is negligible, but the Monte Carlo standard error of the unbiased estimator is an order of magnitude larger than that of the other two estimators. When the treatment effect is correlated with household size, the biases of the unbiased and the post-stratified estimators are still very small, but the bias of the simple difference estimator is substantial—roughly the same size as the standard error. Again, the standard errors are smallest for the post-stratified estimator; overall, the post-stratified estimator clearly dominates in terms of RMSE.

8 Student absenteeism in the School District of Philadelphia

8.1 Overview

Student absenteeism in the United States is astonishingly high. More than 10 percent of public school students—around 6.5 million students—are chronically absent each year, defined as missing 18 or more days of the roughly 180-day school year (ED Office for Civil Rights, 2016). The rate is substantially higher in large, urban school districts: over one-third of the experimental sample in the School District of Philadelphia is chronically absent. High student absenteeism is predictive of a broad range of negative students outcomes, from dropping out of school to crime to drug and alcohol use. It is also an important performance metric for schools and districts, and, in many states, is tied directly to school funding. Policymakers have redoubled their efforts to reduce student absence from school, such as in the newly enacted Every Student Succeeds Act (PL 114-95) and in a recent Obama Administration initiative that aims to reduce chronic absenteeism by ten percent each year (Rogers and Feller, 2016). However, it will be challenging to meet these goals as existing interventions either have limited impacts or are difficult to scale.

Rogers and Feller (2016) recently conducted the first randomized evaluation of an intervention aimed at reducing student absenteeism. This intervention delivered targeted information to parents of at-risk students in the School District of Philadelphia via five pieces of direct mail over the 2014–2015 School Year. The mailing clearly stated the student's number of absences that year ("Alice has been absent 16 days this school year"), included a simple bar chart showing the same information graphically, and gave additional text on the importance of attending school.³ Based on random assignment of 28,080 households, Rogers and Feller (2016) find that the treatment reduces chronic absenteeism by over 10 percent relative to control. The approach is extremely cost-effective, costing around \$5 per additional day of student attendance—more than an order of magnitude more cost-effective than the current best-practice intervention.

A key practical challenge in implementing the original study was that the mailings were explicitly designed to provide information about a single student. Thus, for households with multiple eligible students, one student was randomly selected to be the focal student. Rogers and Feller (2016) addressed this issue only briefly in the original study, largely because the focus was on the overall effect of the intervention and because households with multiple eligible students were a small fraction of the overall sample.

Nonetheless, random assignment of one student within each household presents a rare opportunity to as-

³The original study included three treatment arms and one control arm. The impact of the first treatment arm was relatively weak. The impacts of the second and third treatment arms were large and virtually identical to each other. Based on these results, and for the sake of exposition, we therefore drop the first treatment arm and combine the second and third treatment arms. This yields a much simpler, two-arm trial while preserving the important substantive question.

sess intra-household spillovers. There is substantial evidence across fields that such intra-household spillovers are meaningful in magnitude. For example, several voter mobilization studies have found spillover effects that are between one-third and two-thirds as large as the primary effect (Nickerson, 2008; Sinclair et al., 2012). We are interested in spillover in the attendance study for two key reasons. First, ignoring the spillover effect under-states the overall impact of the intervention. For example, an important metric is the cost of each additional student day; ignoring spillover artificially lowers the corresponding cost-effectiveness estimates. Second, the research team faced a practical question of whether to implement a distinct intervention for households with multiple eligible students, which would be costly to implement and test. If the spillover effect is comparable in magnitude to the primary effect, such development is unnecessary. This is similar to decisions around interventions targeting infectious diseases (Hudgens and Halloran, 2008). Baird et al. (2014) discuss related substantive issues in economics.

8.2 Multilevel randomization and covariate balance

We consider a subset of N=3,804 households with between $n_i=2$ and $n_i=7$ eligible students in each household and $n^+=8,496$ total students.⁴ Table 3 shows the distribution. The vast majority of these households (82 percent) have only two eligible students; only one percent (35 households) have five or more eligible students. We are broadly interested in days absent as the outcome of interest. However, the distribution of absences has a long right tail; for example, several students in the sample are absent over half the time. As this greatly increases the variance, we consider two transformed outcomes of interest. First, we consider an indicator for whether a student is chronically absent, defined as missing 18 or more days during the school year, i.e., $1(\text{days} \ge 18)$; among students in the control group, 36 percent are chronically absent. Second, we consider log-absences, defined as $\log(\text{days} + 1)$, to allow for a continuous outcome without the very heavy right tail; baseline absences among students in the control group are around 13 days or $\log(13+1) \approx 2.6$.

Our goal is to estimate the primary and spillover effects for these two outcomes for the finite sample of either N=3,804 households and $n^+=8,496$ students. For comparison, we report both household- and individual-weighted estimands. As a policy matter, we focus on individual-weighted estimands, as school districts are typically interested in the impact of the intervention on overall student attendance.

Since the original experiment was not directly designed to estimate spillover, we first assess the quality of randomization on the subset of multi-student households. Of the N = 3,804 total households, $N_1 = 2,521$ (66

⁴The original experiment included 24,276 households with only one eligible student.

Table 3: Number of eligible students in each household. Individual-level balance is restricted to households assigned to treatment.

	2	3	4-7
Overall N	3,109	547	148
Proportion in treatment	0.66	0.66	0.66

Table 4: Covariate balance by stage of randomization

	Household-level avg.			Individual-level			
	$\overline{\bar{x}_i}_1$	$\overline{\bar{x}_i}_0$	Δ		\overline{x}_1	\overline{x}_0	Δ
Female	0.53	0.54	-0.03		0.53	0.53	-0.01
Black/African-American	0.51	0.50	0.03		0.52	0.52	-0.01
English spoken at home	0.84	0.83	0.04		0.84	0.83	0.02
Limited English Proficiency	0.07	0.07	-0.01		0.07	0.08	-0.04
Free or Reduced Price Lunch	0.78	0.79	-0.03		0.78	0.79	-0.02
Prior year absences (log-days)	2.74	2.72	0.03		2.73	2.75	-0.04
Start-of-year absences (log-days)	0.57	0.55	0.04		0.58	0.57	0.02
Individuals per household (n_i)	2.2	2.2	-0.01				

percent) were assigned to treatment and $N_0 = 1,283$ (34 percent) were assigned to control. While householdlevel randomization was not stratified by household size (see Section 4.2.3), the balance by household size is excellent, as shown in Table 3. Table 4 shows the covariate balance for each stage of randomization. The left bank shows balance for covariates averaged to the household level (denoted \bar{x}_i), corresponding to the household-level randomization. The right bank shows balance for individual-level covariates among households assigned to treatment. Statistically, Table 4 shows that covariate balance is excellent for both stages of randomization, with all normalized differences (Imbens and Rubin, 2015) below 0.05 in absolute value. Substantively, Table 4 emphasizes that the students come from largely disadvantaged households. Over three-quarters of these students qualify for Free or Reduced Price Lunch, which is only available to families at or near the Federal Poverty Line. Over 15 percent of households speak a language other than English at home, with 7 percent of students designated as Limited English Proficiency (LEP). Moreover, this is a very high-absence group, with an average of around 13 days absent in the previous school year (out of roughly 180 possible days). Note that we also include number of absences prior to randomization in early October. Finally, we observe the grade for each student, which we treat as a discrete covariate and which ranges from first grade to high school senior. While we do not show balance by grade to conserve space, there is excellent balance across this covariate as well.

8.3 Results

Figure 3 shows the estimated impacts and corresponding 95% confidence intervals for the primary and spillover effects for both household- and individual-weighted estimands. In terms of chronic absenteeism (i.e., the binary outcome), the estimates for the HW and IW estimands are nearly identical: the unbiased estimates for the primary effects are around -4 percentage points (SE of 1.5 percentage points) for both τ_{HW}^P and τ_{IW}^P ; the unbiased estimates for the spillover effects are around -3 percentage points (SE of 1.5 percentage points) for both τ_{HW}^S and τ_{IW}^S . These results are virtually unchanged when post-stratifying by household size, using the (conditionally) unbiased estimator within each post-stratification cell, defined by $n_i = 2, n_i = 3$, and $n_i \in \{4, ..., 7\}$.

The results are somewhat more variable for the impact on log-absences (i.e., the continuous outcome). The point estimates are quite close for the household-weighted and individual-weighted estimands: $\hat{\tau}_{HW}^P = -0.085$ log-days and $\hat{\tau}_{IW}^P = -0.093$ log-days for the primary effect, and $\hat{\tau}_{HW}^S = -0.051$ log-days and $\hat{\tau}_{IW}^P = -0.058$ log-days for the spillover effect. The point estimates are similarly close for the post-stratified estimator. The standard errors, however, are considerably larger for the unadjusted IW estimates: roughly 0.033 log-days for the IW estimands compared to 0.023 log-days for the HW estimands. Thus, the corresponding confidence intervals are roughly 50 percent larger for the IW estimands than for the HW estimands. Post-stratification greatly reduces the standard errors for the IW estimand: for both IW and HW estimates, the standard errors are roughly 0.023 log-days, comparable to the standard errors for the HW estimate.

While it is instructive to explore differences between the different estimates, the overall pattern is clear. In general, we find that the spillover effect is between 60 and 80 percent as large as the primary effect, depending on the outcome. We also find few differences between the HW and IW estimands, which suggests that there is not meaningful treatment effect variation by household size.

Next, Figure 4 shows covariate-adjusted estimates for individual-weighted estimands. First, we take advantage of the fact that there is a natural holdout sample in the experiment as analyzed.⁵ To obtain $\hat{r}_{ij}(\gamma)$, we regress the outcome on covariates listed in Table 4 as well as student grade (categorical). Results do not appear sensitive to the particular choice of model. The resulting point estimates in Figure 4 are largely unchanged, if slightly larger in magnitude than the unadjusted estimates. The standard errors, however, are meaningfully smaller, especially for the continuous outcome: 0.018 log-days for the model-

⁵The original design included a treatment arm in which the intervention merely reminded parents of the importance of attendance and did not provide any student-specific information. Rogers and Feller (2016) found minimal impact of this intervention relative to control; thus, we exclude that arm here as we are not interested in measuring spillover for a weak effect. The practical upshot is that we can use households assigned to this weak condition as a holdout sample for estimating the covariate adjustment model.




Figure 3: Treatment effect estimates and 95% confidence intervals for primary (filled-in circles) and spillover (open circles) effects, for household- and individual-weighted estimands with and without post-stratification (PS) by household size.




Figure 4: Treatment effect estimates and 95% confidence intervals for individual-weighted estimands with model-assisted estimation and post-stratification.

assisted estimator versus 0.033 log-days for the unadjusted estimator. Next, we can combine model-assisted estimation with post-stratification, though the results are essentially identical. Finally, while do not have theoretical guarantees for covariate adjustment in a multilevel experiment, classical regression adjustment is nearly identical to the model-assisted estimation with the holdout sample.

Overall, we find strong evidence of intra-household spillover for the attendance intervention. This pattern holds with and without covariate adjustment, though the covariate-adjusted estimates are more precise. This underscores the fact that merely focusing on the primary effect significantly under-estimates the impact of the intervention. Moreover, these results suggest that there are limited gains from introducing an intervention that is specific to multi-student households, since the spillover effects are quite large already.

9 Discussion

Multilevel randomizations are increasingly important designs in settings with interactions between units. This paper addresses important issues that arise when analyzing such designs in practice. First, we address issues that arise when household sizes vary. Second, we demonstrate that regression can yield identical point- and variance-estimates to those derive from fully randomization-based methods. Methodologically, we believe that this is a useful addition to the literatures on both causal inference with interference and randomization-based inference. Substantively, we find important insights into the intra-household dynamics of student behavior.

There are several directions for future work. First, we are actively exploring covariate adjustment in this and other settings with more complex randomization schemes. The model-assisted approach is one such option, but many are possible (Lin, 2013; Aronow and Middleton, 2013). Second, there is an open question of how to separately test the null hypotheses for no primary and no spillover effects in this type of design. Recent work from Athey et al. (2015) offers one promising direction. Third, it will be useful to extend these results to other, related designs. For example, Weiss et al. (2016) discuss an interesting setting in which random assignment occurs at the individual level but individuals are then administered treatment in groups (such as in group therapy). Kang and Imbens (2016) propose a "peer encouragement" design, which extends the multilevel randomization considered here to consider noncompliance. Finally, we anticipate additional connections with non-randomized studies that mimic a multilevel randomized design, such as Hong et al. (2006) and Perez-Heydrich et al. (2014). Overall, we hope that the results we give here will lead to increased use of multilevel designs in practice.

References

- Angelucci, M. and V. Di Maro (2016). Programme evaluation and spillover effects. *Journal of Development Effectiveness* 8(1), 22–43.
- Aronow, P. M. and J. A. Middleton (2013). A class of unbiased estimators of the average treatment effect in randomized experiments. *Journal of Causal Inference* 1(1), 135–154.
- Aronow, P. M. and C. Samii (2013). Estimating average causal effects under interference between units. arXiv:1305.6156.
- Athey, S., D. Eckles, and G. W. Imbens (2015). Exact p-values for network interference. arXiv:1506.02084.
- Athey, S. and G. W. Imbens (2016). The Econometrics of Randomized Experiments. *Handbook of Field Experiments*.
- Baird, S., J. A. Bohren, C. McIntosh, and B. Ozler (2014). Designing experiments to measure spillover effects.
- Bell, R. M. and D. F. McCaffrey (2002). Bias reduction in standard errors for linear regression with multistage samples. Survey Methodology 28(2), 169–181.
- Bowers, J., M. M. Fredrickson, and C. Panagopoulos (2013). Reasoning about interference between units: A general framework. *Political Analysis* 21(1), 97–124.
- Cameron, A. C. and D. L. Miller (2015). A practitioner's guide to cluster-robust inference. *Journal of Human Resources* 50(2), 317–372.
- Cochran, W. G. (1977). Sampling Techniques (3rd ed.). New York: John Wiley & Sons.
- Crépon, B., E. Duflo, M. Gurgand, R. Rathelot, and P. Zamora (2013). Do labor market policies have displacement effects? evidence from a clustered randomized experiment. The Quarterly Journal of Economics 128(2), 531–580.
- Donner, A. and N. Klar (2000). Design and analysis of cluster randomization trials in health research. New York: Oxford University Press.
- ED Office for Civil Rights (2016). Civil Rights Data Collection. Available at http://www2.ed.gov/about/offices/list/ocr/data.html.

- Fisher, R. A. (1935). The design of experiments. Oliver & Boyd.
- Gelman, A. and J. Hill (2006). Data analysis using regression and multilevel/hierarchical models. Cambridge University Press.
- Halloran, M. E., M. Haber, I. M. Longini, and C. J. Struchiner (1991). Direct and indirect effects in vaccine efficacy and effectiveness. *American Journal of Epidemiology* 133(4), 323–331.
- Hansen, B. B. and J. Bowers (2009). Attributing effects to a cluster-randomized get-out-the-vote campaign.

 Journal of the American Statistical Association 104 (487), 873–885.
- Hinkelmann, K. and O. Kempthorne (2012). Design and Analysis of Experiments, Special Designs and Applications, Volume 3. John Wiley & Sons.
- Hong, G., S. W. Raudenbush, et al. (2006). Evaluating kindergarten retention policy: A case study of causal inference for multilevel observational data. Journal of the American Statistical Association 101, 901–910.
- Hudgens, M. G. and M. E. Halloran (2008). Toward causal inference with interference. Journal of the American Statistical Association 103(482), 832–842.
- Imai, K., G. King, C. Nall, et al. (2009). The essential role of pair matching in cluster-randomized experiments, with application to the mexican universal health insurance evaluation. *Statistical Science* 24(1), 29–53.
- Imbens, G. W. and M. Kolesar (2012). Robust standard errors in small samples: Some practical advice.

 Review of Economics and Statistics (0).
- Imbens, G. W. and D. B. Rubin (2015). Causal Inference in Statistics, Social, and Biomedical Sciences.

 Cambridge University Press.
- Kang, H. and G. Imbens (2016). Peer encouragement designs in causal inference with interference. Presentation at 2016 Joint Statistical Meetings.
- Lin, W. (2013). Agnostic notes on regression adjustments to experimental data: Reexamining freedman's critique. The Annals of Applied Statistics 7(1), 295–318.
- Liu, L. and M. G. Hudgens (2014). Large sample randomization inference of causal effects in the presence of interference. *Journal of the American Statistical Association* 109(505), 288–301.

- Manski, C. F. (2013a). Identification of treatment response with social interactions. *The Econometrics Journal* 16(1), S1–S23.
- Manski, C. F. (2013b). Public policy in an uncertain world: analysis and decisions. Harvard University Press.
- McCaffrey, D. F., R. M. Bell, and C. H. Botts (2001). Generalizations of biased reduced linearization. In *Proceedings of the Annual Meeting of the American Statistical Association*, Number 1994.
- Middleton, J. and P. M. Aronow (2015). Unbiased Estimation of the Average Treatment Effect in Cluster Randomized Experiments. *Statistics, Politics and Policy* 6(1–2), 39–75.
- Miratrix, L. W., J. S. Sekhon, and B. Yu (2013). Adjusting treatment effect estimates by post-stratification in randomized experiments. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 75(2), 369–396.
- Neyman, J. (1923). On the application of probability theory to agricultural experiments. Essay on principles. Section 9. Statistical Science 5, 465–472.
- Nickerson, D. W. (2008). Is voting contagious? evidence from two field experiments. American Political Science Review 102(01), 49–57.
- Paluck, E. L., H. Shepherd, and P. M. Aronow (2016). Changing climates of conflict: A social network experiment in 56 schools. *Proceedings of the National Academy of Sciences* 113(3), 566–571.
- Perez-Heydrich, C., M. G. Hudgens, M. E. Halloran, J. D. Clemens, M. Ali, and M. E. Emch (2014).

 Assessing effects of cholera vaccination in the presence of interference. *Biometrics* 70(3), 731–741.
- Pustejovsky, J. E. and E. Tipton (2016). Small sample methods for cluster-robust variance estimation and hypothesis testing in fixed effects models. arXiv preprint arXiv:1601.01981.
- Rigdon, J. and M. G. Hudgens (2015). Exact confidence intervals in the presence of interference. Statistics and Probability Letters 105, 130–135.
- Rogers, T. and A. Feller (2016). Intervening through influential third parties: Reducing student absences at scale via parents.
- Rosenbaum, P. R. et al. (2002). Covariance adjustment in randomized experiments and observational studies. Statistical Science 17(3), 286–327.

- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies.

 Journal of educational Psychology 66(5), 688.
- Rubin, D. B. (1980). Comment. Journal of the American Statistical Association 75 (371), 591–593.
- Samii, C. and P. M. Aronow (2012). On equivalencies between design-based and regression-based variance estimators for randomized experiments. Statistics & Probability Letters 82(2), 365–370.
- Schochet, P. Z. (2013). Estimators for clustered education rcts using the neyman model for causal inference.

 Journal of Educational and Behavioral Statistics 38(3), 219–238.
- Sinclair, B., M. McConnell, and D. P. Green (2012). Detecting spillover effects: Design and analysis of multilevel experiments. *American Journal of Political Science* 56(4), 1055–1069.
- Sobel, M. E. (2006). What do randomized studies of housing mobility demonstrate? causal inference in the face of interference. *Journal of the American Statistical Association* 101(476), 1398–1407.
- Somers, M.-A., W. Corrin, S. Sepanik, T. Salinger, J. Levin, and C. Zmach (2010). The enhanced reading opportunities study final report: The impact of supplemental literacy courses for struggling ninth-grade readers. ncee 2010-4021. *National Center for Education Evaluation and Regional Assistance*.
- Tchetgen Tchetgen, E. and T. VanderWeele (2012). Estimation of causal effects in the presence of interference. Stat. Methods Med. Res. 21, 55–75.
- Toulis, P. and E. Kao (2013). Estimation of causal peer influence effects. In *Proceedings of The 30th International Conference on Machine Learning*, pp. 1489–1497.
- Ugander, J., B. Karrer, L. Backstrom, and J. Kleinberg (2013). Graph cluster randomization: Network exposure to multiple universes. In *Proceedings of the 19th ACM SIGKDD international conference on Knowledge discovery and data mining*, pp. 329–337. ACM.
- VanderWeele, T. J., E. J. Tchetgen Tchetgen, and M. E. Halloran (2014). Interference and Sensitivity Analysis. Statistical Science 29(4), 687–706.
- Weiss, M. J., J. R. Lockwood, and D. F. McCaffrey (2016). Estimating the Standard Error of the Impact Estimator in Individually Randomized Trials With Clustering. *Journal of Research on Educational Effectiveness* 9(3), 421–444.