

WEIERSTRASS SIGMA FUNCTION COEFFICIENTS DIVISIBILITY HYPOTHESIS

ELENA YU. BUNKOVA

ABSTRACT. We consider the coefficients in the series expansion at zero of the Weierstrass sigma function

$$\sigma(z) = z \sum_{i,j \geq 0} \frac{a_{i,j}}{(4i+6j+1)!} \left(\frac{g_2 z^4}{2} \right)^i (2g_3 z^6)^j.$$

We have $a_{i,j} \in \mathbb{Z}$. We present the divisibility Hypothesis for the integers $a_{i,j}$

$$\nu_2(a_{i,j}) = \nu_2((4i+6j+1)!) - \nu_2(i!) - \nu_2(j!) - 3i - 4j,$$

$$\nu_3(a_{i,j}) = \nu_3((4i+6j+1)!) - \nu_3(i!) - \nu_3(j!) - i - j.$$

If this conjecture holds, then $\sigma(z)$ is a Hurwitz series over the ring $\mathbb{Z}[\frac{g_2}{2}, 6g_3]$.

1. EXPOSITION

An *elliptic function* is a meromorphic function on the complex torus $T = \mathbb{C}/\Gamma$, where $\Gamma \subset \mathbb{C}$ is a lattice of rank 2. That is, an elliptic function is a meromorphic function in \mathbb{C} with the periodicity property

$$f(z + \omega) = f(z) \quad \text{for } \omega \in \Gamma.$$

The *Weierstrass sigma function* $\sigma(z) = \sigma(z; g_2, g_3)$ is an entire function in \mathbb{C} determined by the system of equations $Q_0\sigma(z) = 0$, $Q_2\sigma(z) = 0$, where

$$Q_0 = 4g_2 \frac{\partial}{\partial g_2} + 6g_3 \frac{\partial}{\partial g_3} - z \frac{\partial}{\partial z} + 1, \quad Q_2 = 6g_3 \frac{\partial}{\partial g_2} + \frac{1}{3}g_2^2 \frac{\partial}{\partial g_3} - \frac{1}{2} \frac{\partial^2}{\partial z^2} - \frac{1}{24}g_2 z^2, \quad (1)$$

and initial conditions $\sigma(0) = 0$, $\sigma'(0) = 1$. See [1].

The *Weierstrass functions* $\zeta(z)$ and $\wp(z)$ are determined by the expressions

$$\zeta(z) = \frac{\partial \ln \sigma(z)}{\partial z} \quad \text{and} \quad \wp(z) = -\frac{\partial \zeta(z)}{\partial z}.$$

Set $\Delta = g_2^3 - 27g_3^2$. For $\Delta \neq 0$ the function $\wp(z)$ is elliptic. Denote it's lattice of periods by Γ . Any elliptic function with periods Γ is a rational function in $\wp(z)$ and $\frac{\partial}{\partial z}\wp(z)$.

The functions $\wp(z)$ and $\wp'(z)$ satisfy the Weierstrass equation

$$\wp'(z)^2 = 4\wp(z)^3 - g_2\wp(z) - g_3. \quad (2)$$

2. WEIERSTRASS RECURSION

Consider $\sigma(z)$ as a series expansion at 0. Then $\sigma(z) \in \mathbb{Q}[g_2, g_3][[z]]$. From (1) we have

$$\sigma(z) = z - \frac{g_2 z^5}{2 \cdot 5!} - 6g_3 \frac{z^7}{7!} - 9 \frac{g_2^2 z^9}{4 \cdot 9!} - 18g_2g_3 \frac{z^{11}}{11!} + \dots$$

Set

$$\sigma(z) = z \sum_{i,j \geq 0} \frac{a_{i,j}}{(4i+6j+1)!} \left(\frac{g_2 z^4}{2} \right)^i (2g_3 z^6)^j, \quad a_{i,j} \in \mathbb{Q}.$$

Supported in part by Young Russian Mathematics award and RFBR grant 16-51-55017.

Theorem 2.1 (Weierstrass recursion, see [1]). *We have*

$$a_{i,j} = 3(i+1)a_{i+1,j-1} + \frac{16}{3}(j+1)a_{i-2,j+1} - \frac{1}{3}(4i+6j-1)(2i+3j-1)a_{i-1,j} \quad (3)$$

for $i \geq 0, j \geq 0, (i,j) \neq (0,0)$, and

$$a_{0,0} = 1; \quad a_{i,j} = 0 \text{ for } i < 0 \text{ or } j < 0.$$

This theorem follows from (1). Let us note that it determines all the coefficients $a_{i,j}$.

3. HURWITZ SERIES EXPANSION RING

We say that $\varphi(z)$ is a Hurwitz series over the ring R and write $\varphi(z) \in HR[[z]]$ if $\varphi(z) \in R \otimes \mathbb{Q}[[z]]$ and

$$\varphi(z) = \sum_{k=0}^{\infty} \varphi_k \frac{z^k}{k!}, \quad \varphi_k \in R.$$

Problem 3.1. *Find the minimal ring \mathcal{R} , such that $\sigma(z) \in H\mathcal{R}[[z]]$.*

We have $\mathbb{Z}[\frac{g_2}{2}, 6g_3] \subset \mathcal{R} \subset \mathbb{Q}[g_2, g_3]$.

Corollary 3.2 (from 2.1). $\sigma(z) \in H\mathbb{Z}[\frac{1}{3}][\frac{g_2}{2}, 2g_3][[z]]$.

Theorem 3.3 ([3]). $\sigma(z) \in H\mathbb{Z}[\frac{1}{2}][g_2, g_3][[z]]$.

Proof. The article [3] states that $\sigma(z) \in H\mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6][[z]]$ for the sigma-function of the general Weierstrass model of the elliptic curve $y^2 + (\mu_1 x + \mu_3)y = x^3 + \mu_2 x^2 + \mu_4 x + \mu_6$. Set $\mu_1 = \mu_2 = \mu_3 = 0, \mu_4 = -\frac{g_2}{4}, \mu_6 = -\frac{g_3}{4}$. We get the curve (2). Thus $\sigma(z) \in H\mathbb{Z}[\frac{g_2}{4}, \frac{g_3}{4}][[z]]$ and therefore $\sigma(z) \in H\mathbb{Z}[\frac{1}{2}][g_2, g_3][[z]]$. \square

Corollary 3.4 (from 3.2, 3.3). $\sigma(z) \in H\mathbb{Z}[\frac{g_2}{2}, 2g_3][[z]]$.

Therefore in (3) we have $a_{i,j} \in \mathbb{Z}$.

4. DIVISIBILITY HYPOTHESIS

Let $\nu_p(\cdot)$ denote p-adic valuation. Set

$$b_{i,j} = \frac{(4i+6j+1)!}{2^{3i+4j} 3^{i+2j} i! j!}.$$

Lemma 4.1. $b_{i,j} \in \mathbb{Z}$.

Proof. As $\frac{(4i+6j+1)!}{i! j! (3i+5j+1)!} \in \mathbb{Z}$ we have $\nu_p(b_{i,j}) \geq 0$ for prime $p > 3$.

By Legendre's formula $\nu_p(n!) = \sum_{k=1}^{\infty} \left\lfloor \frac{n}{p^k} \right\rfloor$, so for $p = 2$ we have

$$\begin{aligned} \nu_2((4i+6j+1)!) &= \sum_{k=1}^{\infty} \left\lfloor \frac{4i+6j+1}{2^k} \right\rfloor = 2i+3j + \sum_{k=1}^{\infty} \left\lfloor \frac{2i+3j}{2^k} \right\rfloor \geq \\ &\geq 2i+3j+i+j + \sum_{k=1}^{\infty} \left\lfloor \frac{i}{2^k} \right\rfloor + \sum_{k=1}^{\infty} \left\lfloor \frac{j}{2^k} \right\rfloor = 3i+4j + \nu_2(i!) + \nu_2(j!), \end{aligned}$$

and for $p = 3$ we have

$$\begin{aligned} \nu_3((4i + 6j + 1)!) &= \sum_{k=1}^{\infty} \left\lfloor \frac{4i + 6j + 1}{3^k} \right\rfloor = i + 2j + \left\lfloor \frac{i + 1}{3} \right\rfloor + \sum_{k=1}^{\infty} \left\lfloor \frac{i + 2j + \frac{i+1}{3}}{3^k} \right\rfloor \geq \\ &\geq i + 2j + \sum_{k=1}^{\infty} \left\lfloor \frac{i}{3^k} \right\rfloor + \sum_{k=1}^{\infty} \left\lfloor \frac{j}{3^k} \right\rfloor = i + 2j + \nu_3(i!) + \nu_3(j!). \end{aligned}$$

□

Hypothesis 4.2. $\nu_2(a_{i,j}) = \nu_2(b_{i,j}), \quad \nu_3(a_{i,j}) = j + \nu_3(b_{i,j})$.

Corollary 4.3. *If Hypothesis 4.2 holds, the ring $\mathbb{Z}[\frac{q_2}{2}, 6g_3]$ solves problem 3.1.*

5. NUMERICAL CALCULATIONS

The coefficients $a_{i,j}$ grow fast. We have, for example, the prime factorisation

$$a_{10,10} = -2^{11} \cdot 3^{20} \cdot 5^2 \cdot 7 \cdot 11 \cdot 19 \cdot 415516114672128127554409484207124689335643.$$

So far Hypothesis 4.2 has no proof. Numerical calculations using mathematical software (Maple 2015) show that Hypothesis 4.2 holds for all $i \leq 100, j \leq 100$.

The following tables give $\nu_2(a_{i,j})$ for $0 \leq i \leq 20, 0 \leq j \leq 20$ and $\nu_3(a_{i,j}), \nu_3(a_{i,j}) - j$ and $\nu_5(a_{i,j})$ for $0 \leq i \leq 20, 0 \leq j \leq 10$.

Table 4 suggests there is no formula like Hypothesis 4.2 for $\nu_p(a_{i,j})$ for prime $p > 3$.

TABLE 1. $\nu_2(a_{i,j})$

0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
0	1	0	2	0	1	0	3	0	1	0	2	0	1	0	4	0	1	0	2	0
1	3	2	3	1	4	3	4	1	3	2	3	1	5	4	5	1	3	2	3	1
3	3	3	3	4	4	4	4	3	3	3	3	5	5	5	5	3	3	3	3	4
3	3	5	5	4	4	5	5	3	3	6	6	5	5	6	6	3	3	5	5	4
3	6	5	6	4	6	5	6	3	7	6	7	5	7	6	7	3	6	5	6	4
6	7	6	8	6	7	6	10	7	8	7	9	7	8	7	10	6	7	6	8	6
7	7	8	8	7	7	10	10	8	8	9	9	8	8	10	10	7	7	8	8	7
7	7	7	7	9	9	9	9	8	8	8	8	9	9	9	9	7	7	7	7	10
7	8	7	11	9	10	9	11	8	9	8	11	9	10	9	11	7	8	7	12	10
8	12	11	12	10	12	11	12	9	12	11	12	10	12	11	12	8	13	12	13	11
12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	13	13	13	13	13
12	12	13	13	12	12	14	14	12	12	13	13	12	12	16	16	13	13	14	14	13
12	14	13	14	12	15	14	15	12	14	13	14	12	17	16	17	13	15	14	15	13
14	15	14	17	15	16	15	17	14	15	14	19	17	18	17	19	15	16	15	18	16
15	15	17	17	16	16	17	17	15	15	19	19	18	18	19	19	16	16	18	18	17
15	15	15	15	15	15	15	15	17	17	17	17	17	17	17	17	16	16	16	16	16
15	16	15	17	15	16	15	20	17	18	17	19	17	18	17	20	16	17	16	18	16
16	18	17	18	16	21	20	21	18	20	19	20	18	21	20	21	17	19	18	19	17
18	18	18	18	21	21	21	21	20	20	20	20	21	21	21	21	19	19	19	19	21
18	18	22	22	21	21	22	22	20	20	22	22	21	21	22	22	19	19	22	22	21

TABLE 2. $\nu_3(a_{i,j})$

0	0	2	1	1	3	2	4	5	4	4	6	5	5	8	7	8	9	8	8	12
1	2	3	3	3	6	5	5	7	5	6	7	8	8	10	9	9	11	9	12	13
3	3	5	4	6	7	7	7	9	7	7	10	9	10	11	11	11	15	13	13	15
4	4	7	6	6	8	7	8	9	9	9	11	10	10	12	11	14	15	14	14	16
5	7	8	8	8	10	9	9	12	10	11	12	12	12	16	15	15	17	15	16	17
8	8	10	9	10	11	12	12	14	12	12	14	13	16	17	17	17	19	17	17	20
9	9	11	10	10	13	12	13	14	13	13	17	16	16	18	17	18	19	19	19	21
10	11	12	13	13	15	14	14	16	14	17	18	18	18	20	19	19	22	20	21	22
12	12	15	14	15	16	16	16	20	18	18	20	19	20	21	22	22	24	22	22	24
13	13	15	14	14	16	15	18	19	18	18	20	19	19	22	21	22	23	22	22	25
14	15	16	16	16	20	19	19	21	19	20	21	22	22	24	23	23	25	23	25	26

TABLE 3. $\nu_3(a_{i,j}) - j$

0	0	2	1	1	3	2	4	5	4	4	6	5	5	8	7	8	9	8	8	12
0	1	2	2	2	5	4	4	6	4	5	6	7	7	9	8	8	10	8	11	12
1	1	3	2	4	5	5	5	7	5	5	8	7	8	9	9	9	13	11	11	13
1	1	4	3	3	5	4	5	6	6	6	8	7	7	9	8	11	12	11	11	13
1	3	4	4	4	6	5	5	8	6	7	8	8	8	12	11	11	13	11	12	13
3	3	5	4	5	6	7	7	9	7	7	9	8	11	12	12	12	14	12	12	15
3	3	5	4	4	7	6	7	8	7	7	11	10	10	12	11	12	13	13	13	15
3	4	5	6	6	8	7	7	9	7	10	11	11	11	13	12	12	15	13	14	15
4	4	7	6	7	8	8	8	12	10	10	12	11	12	13	14	14	16	14	14	16
4	4	6	5	5	7	6	9	10	9	9	11	10	10	13	12	13	14	13	13	16
4	5	6	6	6	10	9	9	11	9	10	11	12	12	14	13	13	15	13	15	16

TABLE 4. $\nu_5(a_{i,j})$

0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	1	0	0	0	0	1	0	0	0	0	1	0	0	0	0	1	0
0	0	1	1	1	0	0	1	1	1	0	0	1	1	1	0	0	1	1	1	0
0	2	1	1	1	0	1	1	1	1	0	1	1	1	1	0	1	1	1	1	0
2	1	1	1	1	2	1	1	1	1	2	1	1	1	1	2	1	1	1	1	2
1	1	1	1	1	1	1	1	1	1	2	1	1	1	2	1	1	2	1	2	2
1	1	1	1	2	1	2	1	1	2	1	1	1	1	3	1	1	2	2	2	3
1	1	3	2	3	3	1	2	2	2	1	1	2	3	2	1	2	2	3	3	2
1	3	2	2	2	1	2	2	2	3	1	2	5	2	2	2	2	4	3	3	2
3	2	2	2	2	3	2	2	3	2	3	3	2	2	2	3	3	3	3	4	4
2	2	2	2	3	2	2	2	3	2	2	5	2	2	4	3	3	4	3	3	3

REFERENCES

- [1] K. Weierstrass. *Zur Theorie der elliptischen Funktionen*, Mathematische Werke, Bd. **2** (1894), Berlin, Teubner, p. 245–255.
- [2] E. T. Whittaker, G. N. Watson, *A Course of Modern Analysis*, Reprint of 4th (1927) ed., Vol 2. *Transcendental functions*, Cambridge Univ. Press, Cambridge, 1996.
- [3] Y. Ônishi, *Hurwitz integrality of power series expansion of the sigma function for a plane curve*, arXiv:1510.03002

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
E-mail address: `bunkova@mi.ras.ru`