Image: Display information of the problem in the tors, etc.) play a crucial role in many different areas of Mathematics. These operators (are studied extensively but somehow less attention is devoted to operators which are <u>non-compact</u> but close to the class of compact operators. In this work we will focus on two such operators.

 $\sim$  First, consider the Volterra operator V, given by

$$Vf(t) = \int_0^t f(s) \,\mathrm{d}s, \quad (0 \le t \le 1), \quad \text{for } f \in L^1(0,1).$$
 (1.1)

When V is regarded as an operator from  $L^p$  into  $L^q$ ,  $(1 < p, q < \infty)$ , it is a compact experator, however in the limiting case, when V maps  $L^1$  into the space C of continuous functions on the closed unit interval, the operator is bounded, with the operator norm ||V|| = 1, but non-compact. It is worth mentioning that, despite being non-compact or even weakly non-compact, this operator possesses some good properties as being strictly singular (follows from [BG89], or see [Lef17]). This makes Volterra operator, in the above-mentioned limiting case, an interesting example of a non-compact operator "close" to the class of compact operators. The focus of our paper will be on obtaining exact values of strict s-numbers for this operator.

Volterra operator was already extensively studied. Let us briefly recall those results related to our work. The first credit goes to V. I. Levin [Lev38], who computed explicitly the norm of V between two  $L^p$  spaces (1 and described the extremalfunction which is connected with the function  $\sin_n$ . Later on, E. Schmidt in [Sch40] extended this result for  $V: L^p \to L^q$ , where  $1 < p, q < \infty$ . This operator was also studied in the context of Approximation theory [Pin85b, Kol36, Tih60, TB67, BS67, Pin85a].

Date: March 29, 2018.

<sup>2010</sup> Mathematics Subject Classification. Primary 47B06, Secondary 47G10.

Key words and phrases. Integral operator, summation operator, s-numbers.

This research was partly supported by the United States – India Educational Foundation (USIEF) and by the grant P201-13-14743S of the Grant Agency of the Czech Republic.

Later a weighted version of this operator was studied in connection with Brownian motion [LL02], Spectral theory [EE04, EE87] and Approximation theory [EL11].

Recently, sharp estimates for Bernstein-numbers of V in the limiting case were obtained in [Lef17, Theorem 2.2], more specifically,

$$b_n(V) = \frac{1}{2n-1} \quad \text{for } n \in \mathbb{N}, \tag{1.2}$$

and also the estimates for the essential norm can be found in recent preprint [AHLM16].

In our paper, we will compute exact values of all the remaining strict *s*-numbers, i.e., Approximation, Gelfand, Kolmogorov, Mityagin and Isomorphism numbers denoted by  $a_n$ ,  $c_n$ ,  $d_n$ ,  $m_n$  and  $i_n$  respectively (for the exact definitions see Section 2). Our main result reads as follows.

**Theorem 1.1.** Let  $V: L^1(0,1) \to C[0,1]$  be defined as in (1.1). Then

$$a_n(V) = c_n(V) = d_n(V) = \frac{1}{2} \quad \text{for } n \ge 2$$
 (1.3)

and

$$m_n(V) = i_n(V) = \frac{1}{2n-1} \quad \text{for } n \in \mathbb{N}.$$
 (1.4)

If we also include the result (1.2) concerning the Bernstein numbers, we see that all the strict *s*-numbers of *V* split between two groups. The upper half (1.3) remains bounded from below while the lower half (1.4) converges to zero. This phenomenon for this operator was already observed; for instance compare [BS67] and [BG89], and for weighted version see [EL07] and [EL06].

The similar results continue to hold for the sequence spaces and for the discrete analogue of V, namely for the operator  $\sigma: \ell^1 \to c$ , defined as

$$\sigma(\mathbf{x})_k = \sum_{j=1}^k x_j, \quad (k \in \mathbb{N}), \quad \text{for } \mathbf{x} \in \ell^1,$$
(1.5)

where we denoted  $\mathbf{x} = \{x_j\}_{j=1}^{\infty}$  for brevity. The operator is well-defined and bounded with the operator norm  $\|\sigma\| = 1$ . It is shown in [Lef17, Theorem 3.2] that

$$b_n(\sigma) = \frac{1}{2n-1}$$
 for  $n \in \mathbb{N}$ .

We have the next result.

**Theorem 1.2.** Let  $\sigma: \ell^1 \to c$  be the operator from (1.5). Then

$$a_n(\sigma) = c_n(\sigma) = d_n(\sigma) = \frac{1}{2} \quad \text{for } n \ge 2$$

and

$$m_n(\sigma) = i_n(\sigma) = \frac{1}{2n-1}$$
 for  $n \in \mathbb{N}$ .

The proofs are provided at the end of Section 3.

## 2. BACKGROUND MATERIAL

We shall fix the notation in this section, although we mostly work with standard notions from functional analysis.

2.1. Normed linear spaces. For normed linear spaces X and Y, we denote by B(X, Y) the set of all bounded linear operators acting between X and Y. For any  $T \in B(X, Y)$ , we use just ||T|| for its operator norm, since the domain and target spaces are always clear from the context. By  $B_X$ , we mean the closed unit ball of X and, similarly,  $S_X$  stands for the unit sphere of X. It is well-known fact that  $B_X$  is compact if and only if X is finite-dimensional.

Let Z be a closed subspace of the normed space X. The quotient space X/Z is the collection of the sets  $[x] = x + Z = \{x + z; z \in Z\}$  equipped with the norm

$$||[x]||_{X/Z} = \inf\{||x - z||_X; z \in Z\}.$$

We sometimes adopt the notation  $||x||_{X/Z}$  when no confusion is likely to happen. Recall the notion of canonical map  $Q_Z \colon X \to X/Z$ , given by  $Q_Z(x) = [x]$ .

By the Lebesgue space  $L^1$ , we mean the set of all real-valued, Lebesgue integrable functions on (0, 1) identified almost everywhere and equipped with the norm

$$||f||_1 = \int_0^1 |f(s)| \, \mathrm{d}s.$$

The space of real-valued, continuous functions on [0, 1], denoted by C, enjoys the norm

$$||f||_{\infty} = \sup_{0 \le t \le 1} |f(t)|.$$

The discrete counterpart to  $L^1$  is the space of all summable sequences,  $\ell^1$ , where

$$\|\mathbf{x}\|_1 = \sum_{j=1}^{\infty} |x_j|$$

and, similarly to the space C, we denote by c the space of all convergent sequences endowed with the norm

$$\|\mathbf{x}\|_{\infty} = \sup_{j \in \mathbb{N}} |x_j|.$$

Here and in the latter, we use the abbreviation  $\mathbf{x} = \{x_j\}_{j=1}^{\infty}$  for the sequences and we write them in bold font. Note that we also consider only real-valued sequence spaces.

All the above-mentioned spaces are complete, i.e., they form Banach spaces.

2.2. s-numbers. Let X and Y be Banach spaces. To every operator  $T \in B(X, Y)$ , one can attach a sequence of non-negative numbers  $s_n(T)$  satisfying for every  $n \in \mathbb{N}$  the following conditions

- (S1)  $||T|| = s_1(T) \ge s_2(T) \ge \dots \ge 0$ ,
- (S2)  $s_n(T+S) \leq s_n(T) + ||S||$  for every  $S \in B(X, Y)$ ,
- (S3)  $s_n(B \circ T \circ A) \le ||B|| s_n(T) ||A||$  for every  $A \in B(X_1, X)$  and  $B \in B(Y, Y_1)$ ,
- (S4)  $s_n(\operatorname{Id} \colon \ell_n^2 \to \ell_n^2) = 1,$
- (S5)  $s_n(T) = 0$  whenever rank T < n.

The number  $s_n(T)$  is then called the *n*-th *s*-number of the operator *T*. When (S4) is replaced by a stronger condition

(S6)  $s_n(\text{Id}: E \to E) = 1$  for every Banach space E, dim E = n, we say that  $s_n(T)$  is the *n*-th strict *s*-number of T.

Note that the original definition of s-numbers, which was introduced by Pietsch in [Pie74], uses the condition (S6) which was later modified to accommodate wider class of s-numbers (like Weyl, Chang and Hilbert numbers). For a detailed account of s-numbers, one is referred for instance to [Pie07], [CS90] or [EL11].

We shall briefly recall some particular strict s-numbers. Let  $T \in B(X, Y)$  and  $n \in \mathbb{N}$ . Then the *n*-th Approximation, Gelfand, Kolmogorov, Isomorphism, Mityagin and Bernstein numbers of T are defined by

$$a_n(T) = \inf_{\substack{F \in B(X,Y) \\ \operatorname{rank} F < n}} \|T - F\|,$$
  

$$c_n(T) = \inf_{\substack{M \subseteq X \\ \operatorname{codim} M < n}} \sup_{x \in B_M} \|Tx\|_Y,$$
  

$$d_n(T) = \inf_{\substack{N \subseteq Y \\ \dim N < n}} \sup_{x \in B_X} \|Tx\|_{Y/N},$$
  

$$i_n(T) = \sup \|A\|^{-1} \|B\|^{-1},$$

where the supremum is taken over all Banach spaces E with dim  $E \ge n$  and  $A \in B(Y, E), B \in B(E, X)$  such that  $A \circ T \circ B$  is the identity map on E,

$$m_n(T) = \sup_{\substack{\rho \ge 0}} \sup_{\substack{N \subseteq Y \\ \operatorname{codim} N > n}} Q_N T B_X \supseteq \rho B_{Y/N},$$

and

$$b_n(T) = \sup_{\substack{M \subseteq X \\ \dim M \ge n}} \inf_{x \in S_M} \|Tx\|_Y,$$

respectively.

### 3. Proofs

**Lemma 3.1.** Let  $n \in \mathbb{N}$ . Then we have the following lower bounds of the Isomorphism numbers of V and  $\sigma$ .

(i)

$$i_n(V) \ge \frac{1}{2n-1};$$

(ii)

$$i_n(\sigma) \ge \frac{1}{2n-1}.$$

**Proof.** (i) Let  $n \in \mathbb{N}$  be fixed. We shall construct a pair of maps A and B such that the chain

$$\ell^1_{w,n} \xrightarrow{B} L^1 \xrightarrow{V} C \xrightarrow{A} \ell^1_{w,n}$$

forms the identity on  $\ell^1_{w,n}$ . Here  $\ell^1_{w,n}$  is the *n*-dimensional weighted space  $\ell^1$  with the norm given by

$$\|\mathbf{x}\|_{\ell^1_{w,n}} = \sum_{k=1}^n w_k |x_k|$$

For the purpose of this proof, we choose  $w_k = 2$  for  $1 \le k \le n-1$  and  $w_n = 1$ . Now, define  $A: C \to \ell^1_{w,n}$  by

$$(Af)_k = (2n-1) f\left(\frac{2k-1}{2n-1}\right), \quad (1 \le k \le n), \text{ for } f \in C.$$

Obviously, A is bounded with the operator norm

$$||A|| = (2n-1)^2.$$

In order to construct the mapping B, consider the partition of the unit interval into subintervals  $I_1, I_2, \ldots, I_{2n-1}$  of the same length, i.e.

$$I_k = \left[\frac{k-1}{2n-1}, \frac{k}{2n-1}\right] \quad \text{for } 1 \le k \le 2n-1,$$

and define

$$B(\mathbf{x}) = \sum_{k=1}^{n-1} x_k \left( \chi_{I_{2k-1}} - \chi_{I_{2k}} \right) + x_n \chi_{I_{2n-1}}.$$

Clearly  $B(\mathbf{x})$  is integrable for every  $\mathbf{x} \in \ell^1_{w,n}$ , B is bounded and the operator norm satisfies

$$||B|| = \frac{1}{2n-1}.$$

One can observe that the composition  $A \circ V \circ B$  is the identity mapping on  $\ell^1_{w,n}$  and by the very definition of the *n*-th Isomorphism number we have

$$i_n(V) \ge ||A||^{-1} ||B||^{-1} = \frac{1}{2n-1}$$

which completes the proof.

As for the discrete case (ii), we consider the chain

$$\ell_n^\infty \xrightarrow{B} \ell^1 \xrightarrow{\sigma} c \xrightarrow{A} \ell_n^\infty$$

where  $\ell_n^{\infty}$  stands for the *n*-dimensional space  $\ell^{\infty}$ , A is given by

$$A(\mathbf{x})_k = x_{2k-1}, \quad (1 \le k \le n), \quad \text{for } \mathbf{x} \in c$$

and B satisfies

$$B(\mathbf{y}) = (y_1, -y_1, \dots, y_{n-1}, -y_{n-1}, y_n, 0, 0, \dots) \text{ for } \mathbf{y} \in \ell_n^{\infty}.$$

Both A and B are bounded, the composition  $A \circ \sigma \circ B$  forms the identity on  $\ell_n^{\infty}$  and thus

$$i_n(\sigma) \ge ||A||^{-1} ||B||^{-1} = \frac{1}{2n-1},$$

since ||A|| = 1 and ||B|| = 2n - 1.

**Lemma 3.2.** Let  $n \in \mathbb{N}$  then we have the following estimates of the Mityagin numbers of V and  $\sigma$ .

(i)

$$m_n(V) \le \frac{1}{2n-1};$$

(ii)

$$m_n(\sigma) \le \frac{1}{2n-1}.$$

**Proof.** (i) Fix some  $n \in \mathbb{N}$  and any  $0 < \rho < m_n(V)$ . By the definition of the *n*-th Mityagin number, there is a subspace of C, say N, such that  $\operatorname{codim} N \ge n$  and

$$Q_N V B_{L^1} \supseteq \varrho B_{C/N}. \tag{3.1}$$

 $\Box$ 

Define

$$E = \left\{ f \in L^1; \operatorname{span}\{Vf\} \cap N = \{0\} \right\}$$

and observe that, since V is injective, E is a subspace of dimension at most n and satisfies

$$Q_N V B_{L^1} = \{ [Vf]; f \in B_E \}.$$
(3.2)

Thus, by (3.1) and (3.2), we have

 $||Vf||_{\infty} \ge ||Vf||_{C/N} \ge \varrho \quad \text{for } f \in S_E,$ 

and hence

 $||Vf||_{\infty} \ge \varrho ||f||_1 \text{ for } f \in E.$ 

Therefore, thanks to [Lef17, Lemma 2.4],

$$\varrho \le \frac{1}{2n-1} \tag{3.3}$$

and the lemma follows by taking the limit  $\rho \to m_n(V)$ .

(ii) The exact analogy holds in the discrete case, as for every  $0 < \rho < m_n(\sigma)$  one can find a *n*-th dimensional subspace E in  $\ell^1$ , such that

$$\|\sigma(\mathbf{x})\|_{\infty} \ge \varrho \|\mathbf{x}\|_1 \text{ for } \mathbf{x} \in E.$$

The assertion (3.3) then also holds by [Lef17, Lemma 2.4] and its discrete modification in the proof of [Lef17, Theorem 3.2].

**Lemma 3.3.** Let  $n \ge 2$  then the following lower bounds of the Kolmogorov numbers of V and  $\sigma$  hold.

(i)

$$d_n(V) \ge \frac{1}{2};$$

(ii)

$$d_n(\sigma) \ge \frac{1}{2}.$$

**Proof.** (i) Fix arbitrary  $n \ge 2$  and  $\varepsilon > 0$ . By the very definition of the *n*-th Kolmogorov number, there exists a subspace of C, say N, such that dim N < n and

$$d_n(V) + \varepsilon \ge \sup_{f \in B_{L^1}} \|Vf\|_{C/N}.$$
(3.4)

Let us define the trial functions

$$f_k = 2^{k+1} \chi_{(2^{-k-1}, 2^{-k})}, \quad (k \in \mathbb{N}).$$
(3.5)

There is  $||f_k||_1 = 1$  for every  $k \in \mathbb{N}$ . Now, by the definition of the quotient norm, to every  $Vf_k$  one can attach a function  $g_k \in N$  in a way that

$$\|Vf_k - g_k\|_{\infty} \le \|Vf_k\|_{C/N} + \varepsilon.$$
(3.6)

Observe that the set of all the functions  $g_k$  is bounded in N. Indeed, by (3.4) and (3.6),

$$||g_k||_{\infty} \le ||Vf_k - g_k||_{\infty} + ||Vf_k||_{\infty} \le ||Vf_k||_{C/N} + \varepsilon + ||V|| ||f_k||_1 \le d_n(V) + 2\varepsilon + 1$$

for every  $k \in \mathbb{N}$ . Thus, since N is finite-dimensional, there is a convergent subsequence of  $\{g_k\}$  which we denote  $\{g_k\}$  again. Hence  $g_k$  converges to, say,  $g \in N$ , i.e. there is an index  $k_0$  such that  $||g_k - g||_{\infty} < \varepsilon$  for every  $k \ge k_0$ . The limiting function g then satisfies

$$\|Vf_k - g\|_{\infty} \le \|Vf_k - g_k\|_{\infty} + \|g_k - g\|_{\infty} \le \|Vf_k\|_{C/N} + 2\varepsilon$$

for  $k \geq k_0$  and thus

$$\sup_{k \ge k_0} \|Vf_k - g\|_{\infty} \le d_n(V) + 3\varepsilon.$$
(3.7)

Next, we estimate the left hand side of (3.7) by taking the value attained at zero, i.e.

$$\sup_{k \ge k_0} \|Vf_k - g\|_{\infty} \ge \sup_{k \ge k_0} |Vf_k(0) - g(0)| = |g(0)|,$$
(3.8)

or at the points  $2^{-k}$ , i.e.

$$\sup_{k \ge k_0} \|Vf_k - g\|_{\infty} \ge \sup_{k \ge k_0} |Vf_k(2^{-k}) - g(2^{-k})| \ge |1 - g(0)|,$$
(3.9)

where we used that g is continuous. Combining (3.7), (3.8) and (3.9), we get

$$d_n(V) + 3\varepsilon \ge \max\{|g(0)|, 1 - |g(0)|\} \ge \frac{1}{2}$$

Therefore  $d_n(V) \ge 1/2$  by the arbitrariness of  $\varepsilon$ .

(ii) The proof needs just slight modifications here. Instead of functions  $f_k$ , one can use the canonical vectors  $\mathbf{e}^k = (0, \dots, 0, 1, 0, \dots)$ , where the element 1 is placed at the *k*-th coordinate. If we follow the previous lines, we find close points  $\mathbf{y}^k \in N$ , their limit  $\mathbf{y}$  in *c* and we end up with

$$\sup_{k \ge k_0} \|\sigma(\mathbf{e}^k) - \mathbf{y}\|_{\infty} \le d_n(\sigma) + 3\varepsilon.$$
(3.10)

Now, we estimate the supremum in (3.10) by taking the k-th coordinate, i.e.

$$\sup_{k\geq k_0} \|\sigma(\mathbf{e}^k) - \mathbf{y}\|_{\infty} \geq \sup_{k\geq k_0} |\sigma(\mathbf{e}^k)_k - y_k| \geq |1 - \lim_{k\to\infty} y_k|,$$

or by taking the coordinate k - 1, i.e.

$$\sup_{k\geq k_0} \|\sigma(\mathbf{e}^k) - \mathbf{y}\|_{\infty} \geq \sup_{k\geq k_0} |\sigma(\mathbf{e}^k)_{k-1} - y_{k-1}| \geq |\lim_{k\to\infty} y_k|,$$

and the conclusion follows.

**Lemma 3.4.** Let  $n \ge 2$  then the estimates of Gelfand numbers of V and  $\sigma$  read as (i)

$$c_n(V) \ge \frac{1}{2};$$

(ii)

$$c_n(\sigma) \ge \frac{1}{2}$$

**Proof.** (i) Let  $n \ge 2$  and  $\varepsilon > 0$  be fixed. By the definition of the *n*-th Gelfand number, we can find a subspace M in  $L^1$  having codim M < n and satisfying

$$c_n(V) + \varepsilon \ge \sup_{f \in B_M} \|Vf\|_{\infty}.$$
(3.11)

The proof will be finished once we show that the supremum in (3.11) is at least one half. We make use the step functions  $f_k$ ,  $(k \in \mathbb{N})$ , defined by (3.5) in the proof of Lemma 3.3. Recall that  $||f_k||_1 = 1$ ,  $||f_k - f_l||_1 = 2$  and  $||Vf_k - Vf_l||_{\infty} = 1$  for  $k \neq l$ . Note that the quotient space  $L^1/M$  is of finite dimension thus, the projected sequence  $\{[f_k]\}$  is bounded and hence there is a Cauchy subsequence, which we denote  $\{[f_k]\}$ again. Now, let  $\eta > 0$  be fixed. We have

$$\|f_k - f_l\|_{L^1/M} < \eta \tag{3.12}$$

for k and l sufficiently large. Let us denote  $f = \frac{1}{2}(f_k - f_l)$  for these k and l. Thanks to (3.12) and the definition of quotient norm, one can find a function  $g \in M$  such that

$$\|f - g\|_1 \le \eta.$$

On setting

$$h = \frac{g}{1+\eta}$$

we have

$$\|h\|_{1} \leq \frac{1}{1+\eta} (\|f\|_{1} + \|f-g\|_{1}) \leq 1,$$

whence  $h \in B_M$ . Next

$$\|Vh\|_{\infty} \ge \frac{1}{1+\eta} (\|Vf\|_{\infty} - \|V(f-g)\|_{\infty})$$
$$\ge \frac{1}{1+\eta} \left(\frac{1}{2} - \|V\|\|f-g\|_{1}\right)$$
$$\ge \frac{1}{1+\eta} \left(\frac{1}{2} - \eta\right),$$

thus

$$\sup_{f \in B_M} \|Vf\|_{\infty} \ge \frac{1}{1+\eta} \left(\frac{1}{2} - \eta\right)$$

and the lemma follows, since  $\eta > 0$  was arbitrarily chosen.

The proof of the discrete counterpart (ii) is completely analogous, once we consider the canonical vectors  $\mathbf{e}^k$  instead of  $f_k$ , hence we omit it.

**Lemma 3.5.** Let  $n \ge 2$ . Then we have the following upper bounds of the Approximation numbers of V and  $\sigma$ .

(i)

(ii) 
$$a_n(V) \le \frac{1}{2};$$
$$a_n(\sigma) \le \frac{1}{2}.$$

**Proof.** (i) Consider the one-dimensional operator  $F: L^1 \to C$  given by

$$Ff(t) = \frac{1}{2} \int_0^1 f(s) \, \mathrm{d}s, \quad (0 \le t \le 1), \quad \text{for } f \in L^1.$$

Then F is a sufficient approximation of V. Indeed,

$$\begin{split} \|Vf - Ff\|_{\infty} &= \sup_{0 \le t \le 1} \left| \int_{0}^{t} f(s) \, \mathrm{d}s - \frac{1}{2} \int_{0}^{1} f(s) \, \mathrm{d}s \right| \\ &= \sup_{0 \le t \le 1} \left| \frac{1}{2} \int_{0}^{t} f(s) \, \mathrm{d}s - \frac{1}{2} \int_{t}^{1} f(s) \, \mathrm{d}s \right| \\ &\le \sup_{0 \le t \le 1} \frac{1}{2} \int_{0}^{t} |f(s)| \, \mathrm{d}s + \frac{1}{2} \int_{t}^{1} |f(s)| \, \mathrm{d}s \\ &= \frac{1}{2} \|f\|_{1} \end{split}$$

and therefore

$$a_n(V) \le ||V - F|| \le \frac{1}{2}.$$

In order to show (ii), choose the operator

$$\varrho(\mathbf{x})_k = \frac{1}{2} \sum_{j=1}^{\infty} x_j, \quad (k \in \mathbb{N}), \text{ for } \mathbf{x} \in \ell^1.$$

This is a well-defined one-dimensional operator and, by the calculations similar to above,  $\|\sigma - \varrho\| \leq 1/2$ . The proof is complete.

Now, we are at the position to prove the main results.

**Proof of Theorem 1.1.** Let  $n \ge 2$  be fixed. Since  $a_n(V)$  is the largest between all *s*-numbers, we immediately obtain the inequalities  $a_n(V) \ge c_n(V)$  and  $a_n(V) \ge d_n(V)$ . Using Lemma 3.5 with Lemma 3.4 and Lemma 3.3 we obtain

 $\frac{1}{2} \ge a_n(V) \ge c_n(V) \ge \frac{1}{2} \qquad \text{and} \qquad \frac{1}{2} \ge a_n(V) \ge d_n(V) \ge \frac{1}{2}$ 

respectively. This gives (1.3).

Next, let *n* be arbitrary. Due to  $i_n(V)$  being the smallest strict *s*-number, we have that  $i_n(V) \leq b_n(V)$  and also  $i_n(V) \leq m_n(V)$ . For the lower bound, we use Lemma 3.1, while for the upper, we make use of Lemma 3.2 and the result of Lefèvre, [Lef17]. This gives (1.4).

**Proof of Theorem 1.2.** The proof follows along exactly the same lines as that of Theorem 1.1 and hence omitted.  $\Box$ 

# References

- [AHLM16] I. A. Alam, G. Habib, P. Lefèvre, and F. Maalouf. Essential norms of Volterra and Cesàro operators on Müntz spaces. arXiv:1612.03218, 2016.
  - [BG89] J. Bourgain and M. Gromov. Estimates of Bernstein widths of Sobolev spaces. In *Geometric aspects of functional analysis (1987–88)*, volume 1376 of *Lecture Notes in Math.*, pages 176–185. Springer, Berlin, 1989.
  - [BS67] M. S. Birman and M. Z. Solomjak. Piecewise polynomial approximations of functions of classes W<sup>α</sup><sub>p</sub>. Mat. Sb. (N.S.), 73 (115):331–355, 1967.
  - [CS90] B. Carl and I. Stephani. Entropy, compactness and the approximation of operators, volume 98 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1990.
  - [EE87] D. E. Edmunds and W. D. Evans. Spectral theory and differential operators. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1987. Oxford Science Publications.
  - [EE04] D. E. Edmunds and W. D. Evans. Hardy operators, function spaces and embeddings. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004.
  - [EL06] D. E. Edmunds and J. Lang. Approximation numbers and Kolmogorov widths of Hardy-type operators in a non-homogeneous case. *Math. Nachr.*, 279(7):727–742, 2006.
  - [EL07] D. E. Edmunds and J. Lang. Bernstein widths of Hardy-type operators in a non-homogeneous case. J. Math. Anal. Appl., 325(2):1060–1076, 2007.
  - [EL11] D. E. Edmunds and J. Lang. Eigenvalues, embeddings and generalised trigonometric functions, volume 2016 of Lecture Notes in Mathematics. Springer, Heidelberg, 2011.

- [Kol36] A. Kolmogoroff. über die beste Annäherung von Funktionen einer gegebenen Funktionenklasse. Ann. of Math. (2), 37(1):107–110, 1936.
- [Lef17] P. Lefèvre. The Volterra operator is finitely strictly singular from  $L^1$  to  $L^{\infty}$ . J. Approx. Theory, 214:1–8, 2017.
- [Lev38] V. I. Levin. On a class of integral inequalities. Recueil Mathématiques, 4(46):309–331, 1938.
- [LL02] M. A. Lifshits and W. Linde. Approximation and entropy numbers of Volterra operators with application to Brownian motion. *Mem. Amer. Math. Soc.*, 157(745):viii+87, 2002.
- [Pie74] A. Pietsch. s-numbers of operators in Banach spaces. Studia Math., 51:201– 223, 1974.
- [Pie07] A. Pietsch. History of Banach spaces and linear operators. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [Pin85a] A. Pinkus. n-widths in approximation theory, volume 7 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1985.
- [Pin85b] A. Pinkus. n-widths of Sobolev spaces in L<sup>p</sup>. Constr. Approx., 1(1):15–62, 1985.
  - [Sch40] E. Schmidt. über die Ungleichung, welche die Integrale über eine Potenz einer Funktion und über eine andere Potenz ihrer Ableitung verbindet. Math. Ann., 117:301–326, 1940.
  - [TB67] V. M. Tihomirov and S. B. Babadžanov. Diameters of a function class in an  $L_p$ -space ( $p \ge 1$ ). Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk, 11(2):24-30, 1967.
  - [Tih60] V. M. Tihomirov. Diameters of sets in functional spaces and the theory of best approximations. *Russian Math. Surveys*, 15(3):75–111, 1960.

E-mail address, Ö. Bakşi: baksi@yildiz.edu.tr

E-mail address, T. Khan: taqi.khan91@gmail.com

E-mail address, J. Lang: lang@math.osu.edu

ORCiD: 0000-0003-1582-7273

 $E\text{-}mail\ address,\ V.\ Musil:\ \texttt{musil@karlin.mff.cuni.cz}$ 

*ORCiD*: 0000-0001-6083-227X

<sup>1</sup>Department of Mathematics, Ohio State University, Columbus OH, 43210-1174 USA

 $^2 {\rm Yildiz}$  Technical University, Faculty of Art and Science, Department of Mathematics, Istanbul, Turkey

<sup>3</sup>Aligarh Muslim University, Aligarh, U.P.-202002, India

<sup>4</sup>Department of Mathematical Analysis, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75 Praha 8, Czech Republic