Pentavalent symmetric graphs admitting vertex-transitive non-abelian simple groups

Jia-Li Du, Yan-Quan Feng, Jin-Xin Zhou

Department of Mathematics, Beijing Jiaotong University, Beijing 100044, China

Abstract

A graph Γ is said to be *symmetric* if its automorphism group $\operatorname{Aut}(\Gamma)$ is transitive on the arc set of Γ . Let G be a finite non-abelian simple group and let Γ be a connected pentavalent symmetric graph such that $G \leq \operatorname{Aut}(\Gamma)$. In this paper, we show that if G is transitive on the vertex set of Γ , then either $G \subseteq \operatorname{Aut}(\Gamma)$ or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that $G \subseteq T$ and G is one of 58 possible pairs of non-abelian simple groups. In particular, if G is arc-transitive, then G is one of 17 possible pairs, and if G is regular on the vertex set of Γ , then G is one of 13 possible pairs, which improves the result on pentavalent symmetric Cayley graph given by Fang, Ma and Wang in 2011.

Keywords: Symmetric graph, Cayley graph, coset graph, simple group.

2010 Mathematics Subject Classification: 05C25, 20B25.

1 Introduction

Let G be a permutation group on a set Ω and let $\alpha \in \Omega$. Denote by G_{α} the stabilizer of α in G, that is, the subgroup of G fixing the point α . We say that G is regular on Ω if for any two points there is a unique element of G mapping one to the other. Denote by \mathbb{Z}_n , D_n , A_n and S_n the cyclic group of order n, the dihedral group of order 2n, the alternating group and the symmetric group of degree n, respectively. For a subgroup H of a group G, denote by $C_G(H)$ the centralizer of H in G and by $N_G(H)$ the normalizer of H in G.

Throughout this paper, all groups and graphs are finite, and all graphs are simple and undirected. For a graph Γ , we denote its vertex set and automorphism group by $V(\Gamma)$ and $\operatorname{Aut}(\Gamma)$, respectively. A graph Γ is said to be G-vertex-transitive for $G \leq \operatorname{Aut}(\Gamma)$ if G acts transitively on $V(\Gamma)$, G-regular if G acts regularly on $V(\Gamma)$, and G-symmetric if G acts transitively on the arc set of Γ (an arc is an ordered pair of adjacent vertices). In particular, Γ is vertex-transitive or symmetric if it is $\operatorname{Aut}(\Gamma)$ -vertex-transitive or $\operatorname{Aut}(\Gamma)$ -symmetric, respectively. The graph Γ is a Cayley graph on G if G is regular on the vertex set of Γ , and the Cayley graph is normal if G is a normal subgroup of $\operatorname{Aut}(\Gamma)$.

^{*}Corresponding author. E-mails: JiaLiDu@bjtu.edu.cn, yqfeng@bjtu.edu.cn, jxzhou@bjtu.edu.cn

Let G be a non-abelian simple group and let Γ be a pentavalent symmetric G-vertex-transitive graph. In this paper, we show that either $G \subseteq \operatorname{Aut}(\Gamma)$ or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and (G,T) is one of 58 possible pairs of non-abelian simple groups. The motivation of this investigation comes from the two extreme cases, that is, G-symmetric and G-regular.

There are two steps to study a symmetric graph Γ — the first step is to investigate the normal quotient graph for a normal subgroup of an arc-transitive group of automorphisms (see Section 2 for the definition of quotient graph), and the second step is to reconstruct the original graph Γ from the normal quotient by using covering techniques. usually done by taking a maximal normal subgroup such that the normal quotient graph has the same valency as the graph Γ , and in this case, the normal quotient graph is called a basic graph of Γ . The situation seems to be somewhat more promising with 2-arctransitive graphs (a 2-arc is a directed path of length 2), and the strategy for the structural analysis of these graphs, based on taking normal quotients, was first laid out by Praeger (see [23, 24]). The strategy works for locally primitive graphs, that is, vertex-transitive graphs with vertex stabilizers acting primitively on the corresponding neighbors sets (see [25, 26]). For more results, refer to [9, 20] for example. For a non-abelian simple group G, the G-symmetric graphs are important basic graphs and have received wide attention. For example, Fang and Praeger [6, 7] classified G-symmetric graphs admitting G as Suzuki simple groups or Ree simple groups acting transitively on the set of 2-arcs of the graphs. For cubic G-symmetric graph, it was proved by Li [19] that either G is a normal in $Aut(\Gamma)$, or $(G, \text{Aut}(\Gamma)) = (A_7, A_8), (A_7, S_8), (A_7, 2.A_8), (A_{15}, A_{16})$ or (GL(4, 2), AGL(4, 2)). Fang et al [3] proved that none of the above five pairs can happen, that is, G is always normal in $\operatorname{Aut}(\Gamma)$. In this paper, we show that if Γ be a connected pentavalent G-symmetric graph, then either $G \triangleleft \operatorname{Aut}(\Gamma)$ or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and (G, T) is one of 17 possible pairs of non-abelian simple groups.

Investigation of Cayley graphs on a non-abelian simple group is currently a hot topic in algebraic graph theory. One of the most remarkable achievements is the complete classification of connected trivalent symmetric non-normal Cayley graphs on finite non-abelian simple groups. This work was began in 1996 by Li [19], and he proved that a connected trivalent symmetric Cayley graph Γ on a non-abelian simple group G is either normal or $G = A_5$, A_7 , PSL(2, 11), M_{11} , A_{11} , A_{15} , M_{23} , A_{23} or A_{47} . In 2005, Xu et al [32] proved that either Γ is normal or $G = A_{47}$, and two years later, Xu et al [33] further showed that if $G = A_{47}$ and Γ is not normal, then Γ must be 5-arc-transitive and up to isomorphism there are exactly two such graphs.

Let Γ be a connected symmetric Cayley graph Γ on a non-abelian simple group G. Fang, Praeger and Wang [8] developed a general method to investigate the automorphism group $\operatorname{Aut}(\Gamma)$, and using this, the symmetry of Cayley graphs of valency 4 or 5 has been investigated in [4, 5]. Now let Γ be non-normal and of valency 5. Fang $\operatorname{et} \operatorname{al}[5]$ proved that if $\operatorname{Aut}(\Gamma)$ is quasiprimitive then $(G,\operatorname{soc}(\operatorname{Aut}(\Gamma)))=(\operatorname{A}_{n-1},\operatorname{A}_n)$, where either $n=60\cdot k$ with $k\mid 2^{15}\cdot 3$ and $k\neq 3,4,6,8$ or $n=10\cdot m$ with $m\mid 8$, and if $\operatorname{Aut}(\Gamma)$ is not quasiprimitive then there is a maximal intransitive normal subgroup K of $\operatorname{Aut}(\Gamma)$ such that the socle of $\operatorname{Aut}(\Gamma)/K$, denoted by \bar{L} , is a simple group containing $\bar{G}=GK/K\cong G$ properly, where $(\bar{G},\bar{L})=(\Omega^-(8,2),\operatorname{PSp}(8,2)), (A_{14},A_{16}), (A_{n-1},A_n)$ with $n\geq 6$ such that n is a

divisor of $2^{17} \cdot 3^2 \cdot 5$ or $\bar{G} = \bar{L}$ is isomorphic to an irreducible subgroup of $\mathrm{PSL}(d,2)$ for $4 \leq d \leq 17$ and the 2-part $|G|_2 > 2^d$. In this paper, we prove that $\mathrm{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and (G,T) is one of 13 possible pairs of non-abelian simple groups. The following is the main result of this paper.

Theorem 1.1 Let G be a non-abelian simple group and Γ a connected pentavalent symmetric G-vertex-transitive graph. Then either $G \subseteq \operatorname{Aut}(\Gamma)$, or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and $(G,T) = (\Omega_8^-(2), \operatorname{PSp}(8,2))$, (A_{14}, A_{16}) , $(\operatorname{PSL}(2,8), A_9)$ or (A_{n-1}, A_n) with $n \ge 6$ and $n \mid 2^9 \cdot 3^2 \cdot 5$.

Corollary 1.2 Let G be a non-abelian simple group and Γ a connected pentavalent Gsymmetric graph. Then either $G \subseteq \operatorname{Aut}(\Gamma)$, or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple
normal subgroup T such that G < T and $(G,T) = (A_{n-1},A_n)$ with $n = 2 \cdot 3$, $2^2 \cdot 3$, 2^4 , $2^3 \cdot 3$, 2^5 , $2^2 \cdot 3^2$, $2^4 \cdot 3$, $2^3 \cdot 3^2$, $2^5 \cdot 3$, $2^4 \cdot 3^2$, $2^6 \cdot 3$, $2^5 \cdot 3^2$, $2^7 \cdot 3$, $2^6 \cdot 3^2$, $2^7 \cdot 3^2$, $2^8 \cdot 3^2$ or $2^9 \cdot 3^2$. Moreover, if $T = A_6$, then $\operatorname{Aut}(\Gamma) = S_6$ and if $T = A_{2^5}$, $A_{2^7 \cdot 3^2}$, $A_{2^8 \cdot 3^2}$ or $A_{2^9 \cdot 3^2}$, then $\operatorname{Aut}(\Gamma) = T$.

Corollary 1.3 Let G be a non-abelian simple group and Γ a connected pentavalent symmetric G-regular graph. Then either $G \subseteq \operatorname{Aut}(\Gamma)$, or $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and $(G,T) = (A_{n-1}, A_n)$ with $n = 2 \cdot 5$, $2^2 \cdot 5$, $2^3 \cdot 5$, $2 \cdot 3 \cdot 5$, $2^4 \cdot 5$, $2^3 \cdot 3 \cdot 5$, $2^4 \cdot 3^2 \cdot 5$, $2^6 \cdot 3 \cdot 5$, $2^5 \cdot 3^2 \cdot 5$, $2^7 \cdot 3 \cdot 5$, $2^6 \cdot 3^2 \cdot 5$, $2^7 \cdot 3^2 \cdot 5$ or $2^9 \cdot 3^2 \cdot 5$.

For connected symmetric cubic Cayley graphs on non-abelian simple groups, similar to Corollary 1.3 there are six possible pairs $(G,T)=(A_{47},A_{48})$, $(PSL(2,11),M_{11})$, (M_{11},M_{12}) , (A_{11},A_{12}) , (M_{23},M_{24}) or (A_{23},A_{24}) (see [19, Theorem 7.1.3]), and Xu et al [32, 33] proved that only the pair $(G,T)=(A_{47},A_{48})$ can happen and there are exactly two connected non-normal symmetric cubic Cayley graphs on A_{47} with automorphism groups A_{48} . Based on the method given in [32, 33], we may show that $(G,T)=(A_{9},A_{10})$ or (A_{19},A_{20}) cannot happen in Corollary 1.3. For $(G,T)=(A_{39},A_{40})$, with MAGMA [1] we find a connected non-normal pentavalent Cayley graph on A_{39} (see Example 3.8).

2 Preliminaries

In this section, we describe some preliminary results which will be used later. First we describe vertex stabilizers of connected pentavalent symmetric graphs.

Proposition 2.1 [11, Theorem 1.1] Let Γ be a connected pentavalent G-symmetric graph with $v \in V(\Gamma)$. Then $G_v \cong \mathbb{Z}_5$, D_5 , D_{10} , F_{20} , $F_{20} \times \mathbb{Z}_2$, $F_{20} \times \mathbb{Z}_4$, A_5 , S_5 , $A_4 \times A_5$, $S_4 \times S_5$, $(A_4 \times A_5) \rtimes \mathbb{Z}_2$, ASL(2, 4), AGL(2, 4), AΣL(2, 4), AΓL(2, 4) or $\mathbb{Z}_2^6 \rtimes \Gamma L(2, 4)$, where F_{20} is the Frobenius group of order 20, $A_4 \rtimes \mathbb{Z}_2 = S_4$ and $A_5 \rtimes \mathbb{Z}_2 = S_5$. In particular, $|G_v| = 5$, $2 \cdot 5$, $2^2 \cdot 5$, $2^3 \cdot 5$, $2^4 \cdot 5$, $2^2 \cdot 3 \cdot 5$, $2^3 \cdot 3 \cdot 5$, $2^4 \cdot 3^2 \cdot 5$, $2^6 \cdot 3^2 \cdot 5$, $2^5 \cdot 3^2 \cdot 5$, $2^6 \cdot 3 \cdot 5$, $2^7 \cdot 3 \cdot 5$, $2^7 \cdot 3^2 \cdot 5$ or $2^9 \cdot 3^2 \cdot 5$, respectively.

Let Γ be a graph and $N \leq \operatorname{Aut}(\Gamma)$. The quotient graph Γ_N of Γ relative to N is defined as the graph with vertices the orbits of N on $V(\Gamma)$ and with two orbits adjacent if there is an edge in Γ between these two orbits.

Proposition 2.2 [21, Theorem 9] Let Γ be a connected G-symmetric graph of prime valency, and let $N \subseteq G$ have at least three orbits on $V(\Gamma)$. Then N is the kernel of G on $V(\Gamma_N)$, and semiregular on $V(\Gamma)$. Furthermore, Γ_N is G/N-symmetric with $G/N \subseteq \operatorname{Aut}(\Gamma_N)$.

For a group G and a prime p, denote by $O_p(G)$ the largest normal p-subgroup of G and by $\Phi(G)$ the Frattini subgroup of G, that is, the intersection of all maximal subgroups of G. By [29, Lemma 2.7], we have the following proposition.

Proposition 2.3 For a group G and a prime p, let $H = O_p(G)$ and $V = H/\Phi(H)$. Then G has a natural action on V, induced from the action of G on H by conjugation. If $C_G(H) \leq H$, then H is the kernel of this action of G on V, that is, $G/H \leq GL(V)$.

The following result follows from the classification of three-factor simple groups.

Proposition 2.4 [17, Theorem I] Let G be a non-abelian simple $\{2, 3, 5\}$ -group. Then $G = A_5$, A_6 or PSU(4, 2).

Given a group G, its factorization G = HD is said to be maximal if both H and D are maximal subgroups of G.

Proposition 2.5 [5, Lemma 3.3] Let T be a simple group and G a non-abelian simple subgroup such that |T:G|=n>1 and $n=2^a\cdot 3^b\cdot 5^c$ for some $0\leq a\leq 9,\ 0\leq b\leq 2$ and $0\leq c\leq 1$. If T=HD is a maximal factorization with $G\leq H$, then T, G and |T:G| are listed in Table 1.

T	G	T:G	T	G	T:G
M_{11}	PSL(2,11)	$2^2 \cdot 3$	M_{12}	PSL(2,11)	$2^4 \cdot 3^2$
M_{24}	M_{23}	$2^3 \cdot 3$	M_{12}	M_{11}	$2^2 \cdot 3$
$P\Omega_8^+(2)$	PSp(6,2)	$2^3 \cdot 3 \cdot 5$	PSp(6,2)	A_8	$2^3 \cdot 3^2$
$P\Omega_8^+(2)$	A_9	$2^6 \cdot 3 \cdot 5$	PSp(6,2)	A_7	$2^6 \cdot 3^2$
PSp(8,2)	$\Omega_8^-(2)$	$2^4 \cdot 3 \cdot 5$	PSU(4,2)	A_6	$2^3 \cdot 3^2$
PSU(3,3)	PSL(2,7)	$2^2 \cdot 3^2$	${}^{2}F_{4}(2)'$	PSL(2,25)	$2^8 \cdot 3^2$
PSp(4,4)	PSL(2, 16)	$2^4 \cdot 3 \cdot 5$	PSL(3,4)	PSL(2,7)	$2^3 \cdot 3 \cdot 5$
A_{16}	A_{14}	$2^4 \cdot 3 \cdot 5$	A_{10}	A_8	$2 \cdot 3^2 \cdot 5$
A_{10}	A_7	$2^4 \cdot 3^2 \cdot 5$	A_9	A_7	$2^3 \cdot 3^2$
A_9	PSL(2,8)	$2^3 \cdot 3^2 \cdot 5$	A_8	PSL(3,2)	$2^3 \cdot 3 \cdot 5$
A_7	PSL(2,7)	$3 \cdot 5$	PSp(6,2)	PSL(2,8)	$2^6 \cdot 3^2 \cdot 5$
A_n	A_{n-1}	$n = 2^a \cdot 3^b \cdot 5^c$			

Table 1: Non-abelian simple group pairs of index $2^a \cdot 3^b \cdot 5^c$

Let G and E be two groups. We call an extension E of G by N a central extension of G if E has a central subgroup N such that $E/N \cong G$, and if further E is perfect, that is,

the derived group E' equal to E, we call E a covering group of G. A covering group E of G is called a double cover if |E| = 2|G|. Schur [28] proved that for every non-abelian simple group G there is a unique maximal covering group M such that every covering group of G is a factor group of E (see [15, Kapitel V, §23]). This group E is called the full covering group of E, and the center of E is the Schur multiplier of E, denoted by E Mult(E).

Proposition 2.6 For $n \geq 5$, the alternating group A_n has a unique double cover $2.A_n$, and for $n \geq 7$, all subgroups of index n of $2.A_n$ are conjugate and isomorphic to $2.A_{n-1}$.

Proof: By Kleidman and Liebeck [18, Theorem 5.1.4], $\operatorname{Mult}(A_n) \cong \mathbb{Z}_2$ for $n \geq 5$ with $n \neq 6, 7$, and $\operatorname{Mult}(A_n) \cong \mathbb{Z}_6$ for n = 6 or 7. This implies that A_n has a unique double cover for $n \geq 5$, denoted by $2.A_n$. Since A_n has no proper subgroup of index less than n, all subgroups of index n of $2.A_n$ contain the center of $2.A_n$. Let $n \geq 7$. By [31, 2.7.2], $2.A_n$ contains a subgroup $2.A_{n-1}$ of index n, and since all subgroups of index n of A_n are conjugate, all subgroups of index n of $2.A_n$ are conjugate and isomorphic to $2.A_{n-1}$. \square

Let G be a group. For $H \leq G$, let D be a union of some double cosets of H in G such that $D^{-1} = D$. The coset graph $\Gamma = \operatorname{Cos}(G, H, D)$ on G with respect to H and D is defined to have vertex set $V(\Gamma) = [G:H]$, the set of right cosets of H in G, and edge set $E(\Gamma) = \{\{Hg, Hdg\} \mid g \in G, d \in D\}$. The graph Γ has valency |D|/|H| and it is connected if and only if $G = \langle D, H \rangle$, that is, D and H generate G. The action of G on $V(\Gamma)$ by right multiplication induces a transitive group of automorphisms, and this group is symmetric if and only if D is a single double coset. Moreover, this action is faithful if and only if $H_G = 1$, where H_G is the largest normal subgroup of G contained in H.

Conversely, let Γ be a G-vertex-transitive graph with $G \leq \operatorname{Aut}(\Gamma)$. By [27], the graph Γ is isomorphic to a coset graph $\operatorname{Cos}(G, H, D)$, where $H = G_v$ is the vertex stabilizer of $v \in V(\Gamma)$ in G and D consists of all elements of G mapping v to one of its neighbors. It is easy to show that $H_G = 1$ and D is a union of some double cosets of H in G satisfying $D^{-1} = D$. Assume that Γ is G-symmetric and $g \in G$ interchanges v and one of its neighbors. Then $g^2 \in H$ and D = HgH. Furthermore, g can be chosen as a 2-element in G, and the valency of Γ is $|D|/|H| = |H: H \cap H^g|$. For more details regarding coset graph, referee to [6, 21, 22, 27].

Proposition 2.7 Let Γ be a connected G-symmetric graph of valency k, and let $\{u,v\}$ be an edge of Γ . Then Γ is isomorphic to a coset graph $Cos(G, G_v, G_vgG_v)$, where g is a 2-element in G such that $G_{uv}^g = G_{uv}$, $g^2 \in G_v$, $\langle G_v, g \rangle = G$, and $k = |G_v : G_v \cap G_v^g|$.

In Proposition 2.7, the 2-element g such that $g^2 \in G_v$, $\langle G_v, g \rangle = G$ and $k = |G_v|$: $G_v \cap G_v^g$, is called *feasible* to G and G_v . Feasible g can be computed by MAGMA [1] when the order |G| is not too large, and this is often used in Section 1.1.

3 Proof of Theorem 1.1

In this section, we always assume that G is a non-abelian simple group. Let us begin by proving a series of lemmas which will be used in the proof of Theorem 1.1.

Lemma 3.1 Let Γ a connected pentavalent symmetric G-vertex-transitive graph with $G = A_5$. Then either $G \subseteq \operatorname{Aut}(\Gamma)$, or $\Gamma = K_6$ and $\operatorname{Aut}(\Gamma) = S_6$.

Proof: Set $A = \operatorname{Aut}(\Gamma)$. Assume $G \not \triangleq A$. Since $|A_5| = 2^2 \cdot 3 \cdot 5$ and Γ is a connected pentavalent symmetric G-vertex-transitive graph, we have $|V(\Gamma)| = 2^2 \cdot 3 \cdot 5$, $2 \cdot 3 \cdot 5$, $2^2 \cdot 5$, $2 \cdot 5$, $2^2 \cdot 3$ or $2 \cdot 3$. If $|V(\Gamma)| = 2^2 \cdot 3 \cdot 5$, by [12, Theorem 4.1], $\Gamma = \mathcal{G}_{60}$ and $A = A_5 \rtimes D_5$, implying that $G \subseteq A$, contrary to the assumption $G \not \triangleq A$. By [14, Theorems 4.1 and 4.2], there is no connected pentavalent symmetric graph of order $2 \cdot 3 \cdot 5$ or $2^2 \cdot 5$. If $|V(\Gamma)| = 2 \cdot 5$, then $\Gamma = K_{5,5}$ with $A = (S_5 \times S_5) \rtimes \mathbb{Z}_2$ where $S_5 \times S_5$ fixes the bipartite sets of Γ . Since $G = A_5 \leq S_5 \times S_5$, Γ is not G-vertex-transitive, contrary to the hypothesis that Γ is G-vertex-transitive. If $|V(\Gamma)| = 2^2 \cdot 3$, by [14, Theorem 4.1], Γ is the Icosahedron graph I_{12} , or $K_{6,6} - 6K_2$, the complete bipartite graph of order 12 minus a one factor. If $\Gamma = K_{6,6} - 6K_2$ then $A = S_6 \times \mathbb{Z}_2$ and S_6 fixes the bipartite sets of Γ . Thus, $G = A_5 \leq A_6 \leq S_6$ and Γ is not G-vertex-transitive, a contradiction. If $\Gamma = I_{12}$ then $A = A_5 \times \mathbb{Z}_2$, implying that $G \subseteq A$, a contradiction. Thus, $|V(\Gamma)| = 6$. It follows that $\Gamma = K_6$ and $A = S_6$.

Lemma 3.2 Let Γ be a connected pentavalent symmetric G-vertex-transitive graph with $v \in V(\Gamma)$. Let $H \leq \operatorname{Aut}(\Gamma)$ and $GH \leq \operatorname{Aut}(\Gamma)$. Then $|(HG)_v|/|G_v| = |H|/|H \cap G|$ is a divisor of $2^9 \cdot 3^2 \cdot 5$.

Proof: Since Γ is G-vertex-transitive, the Frattini argument implies $GH = G(GH)_v$ and $|H||G|/|H \cap G| = |HG| = |G(GH)_v| = |G||(GH)_v|/|G \cap (GH)_v| = |G||(GH)_v|/|G_v|$, that is, $|H|/|H \cap G| = |(GH)_v|/|G_v|$. Since $GH \leq \operatorname{Aut}(\Gamma)$, Proposition 2.1 implies $|(GH)_v| \mid 2^9 \cdot 3^2 \cdot 5$. It follows $|H|/|H \cap G| \mid 2^9 \cdot 3^2 \cdot 5$.

The radical of a finite group is the largest solvable normal subgroup of the group.

Lemma 3.3 Let Γ be a connected pentavalent symmetric G-vertex-transitive graph and R the radical of $\operatorname{Aut}(\Gamma)$. Let $G \neq A_5$ and $5 \mid |R|$. Then $RG = R \times G$.

Proof: Let $v \in V(\Gamma)$ and set B = RG. Since $G \cap R \subseteq G$, we have $G \cap R = 1$ because the simplicity of G implies $G \not \leq R$. It follows |B| = |R||G|, and by Lemma 3.2, $|R| = |B_v|/|G_v| \mid 2^9 \cdot 3^2 \cdot 5$. Since $5 \mid |R|$, we have $5 \mid |B_v|$, that is, Γ is B-symmetric.

Since R is solvable, R has a Hall $\{2,3\}$ -subgroup, say H. Set $\Omega = \{H^r | r \in R\}$. By [13], all Hall $\{2,3\}$ -subgroups of R are conjugate and so the conjugate action of B is transitive on Ω . Since $|R| \mid 2^9 \cdot 3^2 \cdot 5$, we have $|\Omega| = |R: N_R(H)| = 1$ or S, and since $G \neq A_S$, the conjugate action of S on S is trivial. This means that S normalizes S implying S is S set S is S in S in S in S is S in S in

Since $|G||H|/|G \cap H| = |GH| = |Y| = |G||Y_v|$, Y_v is a $\{2,3\}$ -group as H is a $\{2,3\}$ -group, and hence $(Y_B)_v$ is a $\{2,3\}$ -group. Recall that Γ is B-symmetric and has valency 5. If $(Y_B)_v \neq 1$, then B_v is primitive on the neighborhood $\Gamma(v)$ of v in Γ , and then $Y_B \leq B$ implies $5 \mid |(Y_B)_v|$, a contradiction. It follows $(Y_B)_v = 1$ and $Y_B = G(Y_B)_v = G$. In particular, $G \leq B$ and so $B = R \times G$.

Lemma 3.4 Let Γ a connected pentavalent G-symmetric graph with $G \leq S_9$ and $G \neq A_5$. Then $G \leq \operatorname{Aut}(\Gamma)$.

Proof: Set $A = \operatorname{Aut}(\Gamma)$. By the simplicity of G, $G \leq S_9$ implies $G \leq A_9$. By the Atlas [2, p. 37], $G = A_9$ or $G \leq A_8$, A_7 , A_6 , A_5 , $\operatorname{PSL}(2,8)$. Similarly, if $G \leq A_8$ then $G = A_8$ or $G \leq A_7$, A_6 , $\operatorname{PSL}(3,2)$, A_5 ; if $G \leq A_7$ then $G = A_7$ or $G \leq A_6$, A_5 , $\operatorname{PSL}(2,7)$; if $G \leq A_6$ then $G = A_6$ or A_5 . Since Γ is G-symmetric with $G \neq A_5$, we have $G = A_9$, A_8 , A_7 or A_6 .

By Proposition 2.7, $\Gamma = \text{Cos}(G, G_v, G_v g G_v)$ for some feasible g, that is, a 2-element g in G satisfying $g^2 \in G_v$, $\langle G_v, g \rangle = G$ and $5 = |G_v : G_v \cap G_v^g|$. By Proposition 2.1, g has order at most 8.

For $G = A_9$, by the Atlas [2, p. 37] and Proposition 2.1, we have $G_v \cong \mathbb{Z}_5$, D_5 , D_{10} , F_{20} , $F_{20} \times \mathbb{Z}_2$, A_5 , S_5 , $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$. By MAGMA, G has two conjugacy classes of subgroups isomorphic to D_5 , D_{10} , F_{20} , $F_{20} \times \mathbb{Z}_2$, A_5 or S_5 and one conjugacy class of subgroups isomorphic to \mathbb{Z}_5 , $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$.

Let $G_v \cong A_5$ or S_5 . By taking a given G_v in each conjugacy class, computation with MAGMA shows that there is no feasible g.

Let $G_v \cong D_5$, D_{10} , F_{20} or $F_{20} \times \mathbb{Z}_2$. Then G_v has two conjugacy classes, and computation shows that there is no feasible g in one of the two classes. Take a given G_v in the other class. By MAGMA, if $G_v \cong D_5$ then there are 80 feasible g, and for each of them, the corresponding coset graph Γ is a connected pentavalent G-symmetric graph with $A = A_9 \times \mathbb{Z}_2^2$. Similarly, if $G_v \cong D_{10}$, F_{20} or $F_{20} \times \mathbb{Z}_2$ then in each case, there are 80 feasible g, and for each of them, the corresponding coset graph Γ is a connected pentavalent G-symmetric graph with $A = A_9 \times \mathbb{Z}_2$, $A_9 \rtimes \mathbb{Z}_4$ or S_9 respectively. It follows $G \subseteq A$.

Let $G_v \cong \mathbb{Z}_5$, $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$. Then G_v has one conjugacy class and take a given G_v in the class. By MAGMA, if $G_v \cong \mathbb{Z}_5$ then there are 120 feasible g, and for each of them, the corresponding coset graph is a connected pentavalent G-symmetric graph with $A = A_9 \times \mathbb{Z}_4$. Similarly, if $G_v \cong A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$ then in each case, there are 480 feasible g, and for each of them, the corresponding coset graph is a connected pentavalent G-symmetric graph with $A = A_9 \times \mathbb{Z}_2$ or S_9 respectively. It follows $G \subseteq A$.

For $G = A_8$, A_7 or A_6 , similar arguments as the case $G = A_9$ give rise to $G \subseteq A$. \square

Lemma 3.5 Let Γ be a connected pentavalent G-symmetric graph and R the radical of $\operatorname{Aut}(\Gamma)$. Then $RG = R \times G$.

Proof: Set B = RG and $A = \operatorname{Aut}(\Gamma)$. Since $G \cap R \subseteq G$, we have $G \cap R = 1$ because the simplicity of G implies $G \not\leq R$. It follows |B| = |R||G|. Since Γ is G-symmetric, we have $5 \mid |G_v|$ for $v \in V(\Gamma)$. By Lemma 3.2, $|R| = |B_v|/|G_v|$, and by Proposition 2.1, $|R| \mid 2^9 \cdot 3^2$.

If $G \leq A_9$, Lemmas 3.1 and 3.4 imply that either $G \subseteq A$, or $G = A_5$ and $\Gamma = K_6$ (for this case R = 1). Thus, $RG = R \times G$ for $G \leq A_9$. In what follows we may assume $G \nleq S_9$.

Let $\mathrm{Syl}_2(R)$ be the set of Sylow 2-subgroups of R. Recall that $R \subseteq A$ and $|R| \mid 2^9 \cdot 3^2$. By Sylow's Theorem, $|\mathrm{Syl}_2(R)| = 1, 3$ or 9. Since $R \subseteq B$, G has a natural action on $\mathrm{Syl}_2(R)$ by conjugation, and since $G \nleq S_9$, G fixes every element in $\mathrm{Syl}_2(R)$. Thus, G normalizes P for any $P \in \mathrm{Syl}_2(R)$. Clearly, $|P| = 2^r$ for some $0 \le r \le 9$.

Now we claim $GP = G \times P$. Set D = PG. Then $P \subseteq D$ and Γ is D-symmetric. The simplicity of G implies $P \cap G = 1$. By Lemma 3.2, $|P| = |P|/|P \cap G| = |D_v|/|G_v| = 2^r$ and hence $|D_v| = 2^r \cdot |G_v|$, which implies that if $G_v \cong \mathbb{Z}_5$ then by Proposition 2.1, $D_v \cong D_5$, D_{10} , F_{20} , $F_{20} \times \mathbb{Z}_2$ or $F_{20} \times \mathbb{Z}_4$. Thus, $r \leq 5$. This is always true for other G_v by checking all other possible cases in Proposition 2.1. In particular, r = 5 if and only if $(G_v, D_v) \cong (A_5, A\Sigma L(2, 4))$ or $(A_4 \times A_5, \mathbb{Z}_2^6 \rtimes \Gamma L(2, 4))$.

Let $C = C_D(P)$. Then $C \subseteq D$ and $G \cap C = G$ or 1. Suppose $G \cap C = 1$. Since P is 2-group and $D/P \cong G$ is simple, we have $P = O_2(D)$, the largest normal 2-subgroup of D. On the other hand, since D = PG and $G \cap C = 1$, C is 2-group and hence $C \subseteq P$. Since $r \subseteq S$, Proposition 2.3 implies $G \cong D/P \subseteq GL(5,2) \cong PSL(5,2)$, and since $G \not\subseteq S_9$, the Atlas [2, p. 70] implies G = PSL(5,2) because $PSL(4,2) \cong A_8$. It follows that r = 5 and $(G_v, D_v) \cong (A_5, A\Sigma L(2,4))$ or $(A_4 \times A_5, \mathbb{Z}_2^6 \rtimes \Gamma L(2,4))$. However, by MAGMA, G has no subgroup isomorphic to $A_4 \times A_5$. Thus, $G_v \cong A_5$, and by Proposition 2.7, $\Gamma = Cos(G, G_v, G_v g G_v)$ for a 2-element g such that $g^2 \in G_v$, $\langle G_v, g \rangle = G$ and $|G_v : G_v \cap G_v^g| = 5$. Again by MAGMA, G has four conjugacy classes of subgroups isomorphic to A_5 and by taking a G_v in each class, computation shows that there is no feasible g for G_v . It follows $G \cap C = G$, that is, $G \subseteq C$. Since $G \cap P = 1$, we have $D = P \times G$, as claimed.

Set $\Omega = \{Db \mid b \in B\}$. Then $|\Omega| = 1, 3$ or 9. By considering the right multiplication action of B on Ω , we have $B/D_B \leq S_9$, where D_B is the largest normal subgroup of B contained in D. Since $G \nleq S_9$, we have $G \leq D_B$. Thus, G is characteristic in D_B because $D = P \times G$ implies that G is characteristic in D. It follows that $G \subseteq B$ as $D_B \subseteq B$. Since $R \subseteq B$ and $R \cap G = 1$, we have $B = RG = R \times G$.

Lemma 3.6 Let Γ be a connected pentavalent X-symmetric G-vertex-transitive graph with $G \leq X$. Assume X has trivial radical. Then either $G \subseteq X$, or X has a non-abelian simple normal subgroup T such that G < T and $(G,T) = (\Omega_8^-(2), \operatorname{PSp}(8,2)), (A_{14}, A_{16}), (\operatorname{PSL}(2,8), A_9)$ or (A_{n-1}, A_n) with $n \geq 6$ and $n \mid 2^9 \cdot 3^2 \cdot 5$.

Proof: Let $A = \operatorname{Aut}(\Gamma)$. Let N be a minimal normal subgroup of X. Since X has trivial radical, $N = T^s$ for a positive integer s and a non-abelian simple group T. Then $NG \leq X$, and by Lemma 3.2, $|N|/|N \cap G| \mid 2^9 \cdot 3^2 \cdot 5$.

If $G = A_5$, by Lemma 3.1, either $G \subseteq X$ or $\Gamma = K_6$ and $A = S_6$. For the latter, $(G, T) = (A_5, A_6)$. In what follows we may assume $G \neq A_5$ and $G \not\subseteq X$. Thus, $G \not\subseteq A$.

Suppose $N \cap G = 1$. Then $|N| = |N|/|N \cap G| \mid 2^9 \cdot 3^2 \cdot 5$, |K| = |NG| = |N||G| and $K/N \cong G$. Write K = NG. For $v \in V(\Gamma)$, we have $K = GK_v$ and $|K| = |GK_v| = |G||K_v|/|G_v|$. It follows $|K_v| = |G_v||N|$. Note that $|PSU(4,2)| = 2^6 \cdot 3^4 \cdot 5$. By Proposition 2.4, $N = A_5$ or A_6 . Thus, $5 \mid |K_v|$, forcing that Γ is K-symmetric. Let $C = C_K(N)$.

Assume C = 1. Then $G \leq K = K/C \leq \operatorname{Aut}(N) \leq S_6$, implying that $5 \mid |G|$ and $5^2 \mid |K|$. On the other hand, $5^2 \mid |N||G| = |K|$, a contradiction.

Assume $C \neq 1$. Then $C \cap N = Z(N) = 1$, and $C = C/C \cap N \cong CN/N \subseteq K/N \cong G$. Since |K| = |N||G| = |N||C|, we have $K = C \times N$. First let $C \cap G = 1$. Then |CG| = |C||G|. Since $CG \subseteq K$, |CG| is a divisor of |K| = |N||C|, implying $|G| \mid |N|$. Since $N \cong A_5$ or A_6 and $G \neq A_5$, we have $C \cong G \cong N = A_6$ and hence $K = C \times N \cong A_6 \times A_6$. If $C_v = 1$ then C is regular on $V(\Gamma)$ because Γ is G-vertex-transitive. In this case, $|C||N| = |K| = |CK_v| = |C||K_v|$ and $|K_v| = |N| = |A_6| = 2^3 \cdot 3^2 \cdot 5$, which is impossible by Proposition 2.1. Thus, $C_v \neq 1$, and similarly, $N_v \neq 1$. Since Γ is K-symmetric and has valency 5, K_v is primitive on the neighborhood N(v), and since $C_v \subseteq K_v$ and $N_v \subseteq K_v$, both C_v and N_v are transitive on N(v). It follows $5 \mid |N_v|$ and $5 \mid |C_v|$, and hence $5^2 \mid |K_v|$, contrary to Proposition 2.1. Now let $C \cap G \neq 1$. Then $G \subseteq C$ as G is simple. Since $C \cong G$, we have C = G, and so $K = G \times N$. Set $L = C_X(N)$ and M = LN. Then $L \subseteq X$, $G \subseteq L$ and $M = L \times N$ because $L \cap N = Z(N) = 1$. Since $G \not\subseteq X$, G is a proper subgroup of L and hence $L_v \neq 1$. It follows $5 \mid |L_v|$ because Γ is X-symmetric. Since $|L||N| = |M| = |LM_v| = |L||M_v|/|L_v|$, we have $|M_v| = |L_v||N|$, which implies $5^2 \mid |M_v|$, contrary to Proposition 2.1.

The above contradictions imply $N \cap G \neq 1$, and so $G \leq N$. Thus, $|N|/|G| = |N|/|N \cap G| \mid 2^9 \cdot 3^2 \cdot 5$. Since $G \nleq X$, G is a proper subgroup of N, yielding $5 \mid |N_v|$. It follows that Γ is N-symmetric. Recall that $N = T^s$ for some $s \geq 1$.

Suppose $s \geq 2$. We claim $|T| \mid 2^9 \cdot 3^2 \cdot 5$. It suffices to show $|T| \mid |N|/|G|$. Note that $T \cap G = G$ or 1. If $T \cap G = G$ then $G \leq T$ and so $|G| \mid |T|$. Since $s \geq 2$, we have $|T| \mid |N|/|G|$. If $G \cap T = 1$ then |TG| = |T||G|. Since $TG \leq N$, we have $|G| \mid |T|^{s-1}$ and hence $|T| \mid |N|/|G|$, as required. The proof also implies that if s = 2 then $|G| \mid |T|$.

Since $G \leq N$, the above claim means that both G and T are non-abelian simple $\{2,3,5\}$ -groups. Since $G \neq A_5$, Proposition 2.4 implies $G = A_6$ or PSU(4,2) and also $T = A_6$ or PSU(4,2). Since $5^2 \nmid |N|/|G|$, we have s = 2 and so $|G| \mid |T|$. Note that $N = T \times T$. If both direct factors T of N are not semiregular, then $5 \mid |T_v|$ for each T because Γ is N-symmetric, yielding $5^2 \mid |N_v|$, contrary to Proposition 2.1. Thus, N has a direct factor T with $T_v = 1$. Since $|G| \mid |T|$ and Γ is G-vertex-transitive, T is regular on $V(\Gamma)$ and so $|N_v| = |T| = |A_6|$ or |PSU(4,2)|, that is, $2^3 \cdot 3^2 \cdot 5$ or $2^6 \cdot 3^4 \cdot 5$, of which both are impossible by Proposition 2.1.

The above contradictions imply s=1, that is, N=T is a non-abelian simple group. Thus, $G \leq T \leq X$ and $|T:G|=|N|/|G| \mid 2^9 \cdot 3^2 \cdot 5$. Since Γ is X-symmetric, it is also T-symmetric, that is, $5 \mid |T_v|$. Furthermore, $|T|/|G|=|T_v|/|G_v|$ and $G \neq T$ as $G \nleq X$. If $|T|/|G| \mid 2^9 \cdot 3^2$, then $5 \mid |T_v|$, implying that Γ is G-symmetric. By Proposition 2.7, $\Gamma = \text{Cos}(T, T_v, T_v t T_v)$ for some feasible t, and if Γ is G-symmetric then $\Gamma = \text{Cos}(G, G_v, G_v g G_v)$ for some feasible g.

By the Frattini argument, $T = GT_v$. Let H and D be maximal subgroups of T containing G and T_v , respectively. Then T = HD is a maximal factorization of T. By Proposition 2.5, (G,T) is listed in Table 1. Clearly, $(G,T) \neq (PSL(2,7), PSU(3,3))$ as $5 \nmid |PSU(3,3)|$. To finish the proof, we only need to show $(G,T) \neq (A_6, PSU(4,2))$, $(A_7, PSp(6,2)), (A_7, A_9), (A_8, PSp(6,2)), (PSL(2,11), M_{11}), (PSL(2,11), M_{12}), (M_{11}, M_{12}), (M_{23}, M_{24}), (PSL(2,25), {}^2F_4(2)'), (PSL(2,16), PSp(4,4)), (PSL(2,7), PSL(3,4)), (PSL(2,7), A_7), (PSL(3,2), A_8), (PSL(2,8), PSp(6,2)), (A_8, A_{10}), (A_7, A_{10}), (A_9, P\Omega^+(8,2))$ or $(PSp(6,2), P\Omega^+(8,2))$.

Suppose $(G,T) = (A_6, \mathrm{PSU}(4,2)), (A_7, \mathrm{PSp}(6,2)), (A_7, A_9)$ or $(A_8, \mathrm{PSp}(6,2)).$ Then $|T_v|/|G_v| = |T|/|G| \mid 2^6 \cdot 3^2$ and Γ is G-symmetric. By Lemma 3.4, $G \subseteq A$, a contradiction. Suppose $(G,T) = (\mathrm{PSL}(2,11), M_{11})$ or $(\mathrm{PSL}(2,11), M_{12}).$ Then $|T_v|/|G_v| = |T|/|G| \mid 2^4 \cdot 3^2$ and Γ is G-symmetric. By the Atlas [2, p. 7] and Proposition 2.1, $G_v \cong \mathbb{Z}_5, D_5$

or A_5 . If $G_v \cong A_5$ then $|V(\Gamma)| = 11$, which is impossible because there is no pentavalent graph of odd order. It follows $G_v \cong \mathbb{Z}_5$ or D_5 , and hence $|V(\Gamma)| = 132$ or 66. By [12, Theorem 4.1] and [14, Theorem 4.2], $\Gamma = \mathcal{G}_{132}^1$, \mathcal{G}_{132}^2 , \mathcal{G}_{132}^3 or \mathcal{G}_{66} : $A = \mathrm{PSL}(2, 11) \times \mathbb{Z}_2$ for $\Gamma = \mathcal{G}_{132}^1$ and $A = \mathrm{PGL}(2, 11)$ for other cases. In all cases, $G \subseteq A$, a contradiction.

Suppose $(G,T)=(M_{11},M_{12})$. Then $|T_v|/|G_v|=|T|/|G|$ | $2^2 \cdot 3$ and Γ is G-symmetric. By the Atlas [2, p. 18] and Proposition 2.1, $G_v \cong \mathbb{Z}_5$, D_5 , F_{20} , A_5 or S_5 , and $|T_v|=|G_v||T|/|G|=2^2 \cdot 3 \cdot 5$, $2^3 \cdot 3 \cdot 5$, $2^4 \cdot 3 \cdot 5$, $2^4 \cdot 3^2 \cdot 5$ or $2^5 \cdot 3^2 \cdot 5$, respectively. By Proposition 2.1, $(G_v, T_v) \cong (\mathbb{Z}_5, A_5)$, (D_5, S_5) , $(A_5, A_4 \times A_5)$ or $(S_5, (A_4 \times A_5) \rtimes \mathbb{Z}_2)$. By MAGMA, T has no subgroups isomorphic to $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$, and G has one conjugacy class of subgroups isomorphic to \mathbb{Z}_5 or D_5 . Take a given G_v in each class: for $G_v \cong D_5$, there is no feasible g for G_v , and for $G_v \cong \mathbb{Z}_5$, there are 40 feasible g for G_v . For every such feasible g, the corresponding coset graph Γ is a connected pentavalent G-symmetric graph with $A = M_{11}$. Thus, $G \subseteq A$, a contradiction.

Suppose $(G,T)=(M_{23},M_{24})$. Then $|T_v|/|G_v|=|T|/|G|$ | $2^3 \cdot 3$ and Γ is G-symmetric. By the Atlas [2, p. 71] and Proposition 2.1, $G_v \cong \mathbb{Z}_5$, D_5 , F_{20} , A_5 , S_5 , ASL(2,4), AGL(2,4), A Σ L(2,4) or A Γ L(2,4), and $|T_v|=|G_v||T|/|G|=2^3 \cdot 3 \cdot 5$, $2^4 \cdot 3 \cdot 5$, $2^5 \cdot 3 \cdot 5$, $2^5 \cdot 3^2 \cdot 5$, $2^6 \cdot 3^2 \cdot 5$, $2^9 \cdot 3^2 \cdot 5$, $2^9 \cdot 3^3 \cdot 5$, $2^{10} \cdot 3^2 \cdot 5$ or $2^{10} \cdot 3^3 \cdot 5$, respectively. By Proposition 2.1, $(G_v, T_v) \cong (\mathbb{Z}_5, S_5)$, $(A_5, (A_4 \times A_5) \rtimes \mathbb{Z}_2)$, $(S_5, AGL(2,4))$, $(S_5, S_4 \times S_5)$ or $(ASL(2,4), \mathbb{Z}_2^6 \rtimes \Gamma L(2,4))$. By MAGMA, T has no subgroup isomorphic to $S_4 \times S_5$, AGL(2,4) has no subgroup isomorphic to S_5 , and $\mathbb{Z}_2^6 \rtimes \Gamma L(2,4)$ has no subgroup isomorphic to ASL(2,4) (both are subgroups of M_{24}). Furthermore, T has five conjugacy classes of subgroups isomorphic to S_5 and one conjugacy class of subgroups isomorphic to $(A_4 \times A_5) \rtimes \mathbb{Z}_2$. Take a given T_v in each class, and computation shows that T has no feasible t for each T_v , a contradiction.

Suppose $(G,T) = (\mathrm{PSL}(2,25), {}^2F_4(2)')$. Then $|T_v|/|G_v| = |T|/|G| \mid 2^8 \cdot 3^2$ and Γ is G-symmetric. By the Atlas [2, p. 16] and Proposition 2.1, $G_v \cong \mathbb{Z}_5$, D_5 , F_{20} , A_5 or S_5 , and $|T_v| = |G_v||T|/|G| = 2^8 \cdot 3^2 \cdot 5$, $2^9 \cdot 3^2 \cdot 5$, $2^{10} \cdot 3^2 \cdot 5$, $2^{10} \cdot 3^3 \cdot 5$ or $2^{11} \cdot 3^3 \cdot 5$, respectively. By Proposition 2.1, $(G_v, T_v) \cong (D_5, \mathbb{Z}_2^6 \rtimes \Gamma L(2, 4))$, and by MAGMA, G contains two conjugacy classes of subgroups isomorphic to D_5 . By taking a given T_v in each conjugacy class, computation shows that for each G_v there are 40 feasible G, and for every such G, the corresponding coset graph G is a connected pentavalent G-symmetric graph with G = PSL(2, 25) × G Thus, $G \subseteq G$, a contradiction.

Suppose $(G,T)=(\mathrm{PSL}(2,16),\mathrm{PSp}(4,4))$. Then $|T_v|/|G_v|=|T|/|G|=2^4\cdot 3\cdot 5 \mid |T_v|$. Since $|T|=2^8\cdot 3^2\cdot 5^2\cdot 17$, Proposition 2.1 implies $T_v\cong A_4\times A_5$, $(A_4\times A_5)\rtimes \mathbb{Z}_2$, $S_4\times S_5$, $\mathrm{ASL}(2,4)$, $\mathrm{AGL}(2,4)$, $\mathrm{ASL}(2,4)$, $\mathrm{AFL}(2,4)$. However, by MAGMA, $\mathrm{PSp}(4,4)$ has no subgroup isomorphic to $(A_4\times A_5)\rtimes \mathbb{Z}_2$, $S_4\times S_5$, $\mathrm{ASL}(2,4)$, $\mathrm{AGL}(2,4)$, $\mathrm{ASL}(2,4)$ or $\mathrm{AFL}(2,4)$. For $T_v\cong A_4\times A_5$, T_v has two conjugacy classes in T. By taking a given T_v in each conjugacy class, computation shows that there is no feasible t, a contradiction.

Suppose (G,T) = (PSL(2,7), PSL(3,4)). Then $|T| = 2^6 \cdot 3^2 \cdot 5 \cdot 7$ and $|T_v|/|G_v| = |T|/|G| = 2^3 \cdot 3 \cdot 5 \mid |T_v|$. Note that there is no pentavalent graph of odd order. By Proposition 2.1, $T_v \cong S_5$, $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$. By the Atlas [2, p. 23], T has no subgroup isomorphic to S_5 , $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$, a contradiction.

Suppose $(G,T) = (PSL(2,7), A_7)$. Then $|T_v|/|G_v| = |T|/|G| = 3 \cdot 5 \mid |T_v|$. By the Atlas [2, p. 10] and Proposition 2.1, $T_v \cong A_5$. By MAGMA, $T_v \cong A_5$ has two conjugacy classes in T and by taking a given T_v in each conjugacy class, computation shows that

there is no feasible t, a contradiction.

Suppose $(G, T) = (PSL(3, 2), A_8)$. Then $|T_v|/|G_v| = |T|/|G| = 2^3 \cdot 3 \cdot 5 \mid |T_v|$. By the Atlas [2, p. 22] and Proposition 2.1, $T_v \cong S_5$. By MAGMA, $T_v \cong S_5$ has two conjugacy classes and by taking a given T_v in each conjugate class, computation shows that there is no feasible t, a contradiction.

Suppose (G, T) = (PSL(2, 8), PSp(6, 2)). Then $|T_v|/|G_v| = |T|/|G| = 2^6 \cdot 3^2 \cdot 5 \mid |T_v|$. Since $|T| = 2^9 \cdot 3^4 \cdot 5 \cdot 7$, Proposition 2.1 implies $T_v \cong S_4 \times S_5$, AGL(2, 4) or AFL(2, 4). However, by the Atlas [2, p. 46], PSp(6, 2) has no subgroup isomorphic to $S_4 \times S_5$, AGL(2, 4) or AFL(2, 4), a contradiction.

Suppose $(G,T)=(A_8,A_{10})$ or (A_7,A_{10}) . Then $3^2 \mid |T|/|G| = |T_v|/|G_v|$, and by the Atlas [2, p. 48] and Proposition 2.1, $T_v \cong A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$. For each case, by MAGMA, T_v has two conjugacy classes. By taking a given T_v in each conjugacy class, computation shows that there is no feasible t, a contradiction.

Suppose $(G,T)=(A_9, P\Omega^+(8,2))$ or $(PSp(6,2), P\Omega^+(8,2))$. Then $|T_v|/|G_v|=|T|/|G|=2^6\cdot 3\cdot 5$ or $2^3\cdot 3\cdot 5$. By Proposition 2.1, $T_v\cong S_5$, $A_4\times A_5$, $(A_4\times A_5)\rtimes \mathbb{Z}_2$, $S_4\times S_5$, ASL(2,4), AGL(2,4), $A\Sigma L(2,4)$, $A\Gamma L(2,4)$ or $\mathbb{Z}_2^6\rtimes \Gamma L(2,4)$. By MAGMA, $P\Omega^+(8,2)$ has no subgroup isomorphic to $S_4\times S_5$, ASL(2,4), AGL(2,4), $A\Sigma L(2,4)$, $A\Gamma L(2,4)$ or $\mathbb{Z}_2^6\rtimes \Gamma L(2,4)$. Thus, $T_v\cong S_5$, $A_4\times A_5$ or $(A_4\times A_5)\rtimes \mathbb{Z}_2$, and the computation shows that T_v has 15, 3, 3 conjugacy classes, respectively. By taking a given T_v in each conjugacy class, there is no feasible t respectively, a contradiction.

Lemma 3.7 Let Γ be a connected pentavalent symmetric G-vertex-transitive graph and let $\operatorname{Aut}(\Gamma)$ have a non-trivial radical R with at least three orbits. Assume $G \not \supseteq \operatorname{Aut}(\Gamma)$ and $RG = R \times G$. Then $\operatorname{Aut}(\Gamma)$ contains a non-abelian simple normal subgroup T such that G < T and $(G,T) = (\Omega_8^-(2), \operatorname{PSp}(8,2)), (A_{14}, A_{16}), (\operatorname{PSL}(2,8), A_9)$ or (A_{n-1}, A_n) with $n \geq 8$ and $n \mid 2^9 \cdot 3^2 \cdot 5$.

Proof: Set $A = \operatorname{Aut}(\Gamma)$ and $B = RG = R \times G$. Since $R \neq 1$ has at least three orbits, by Proposition 2.2, the quotient graph Γ_R is a connected pentavalent A/R-symmetric graph with $A/R \leq \operatorname{Aut}(\Gamma_R)$. Furthermore, Γ_R is B/R-vertex-transitive and G is characteristic in B. It follows that $B \not \supseteq A$ because $G \not \supseteq A$, and hence $G \cong B/R \not \supseteq A/R$. Since R is the largest solvable normal subgroup of A, A/R has trivial radical. By Lemma 3.6, A/R has a non-abelian simple normal subgroup I/R such that $B/R \leq I/R \cong T$ and $(B/R, I/R) \cong (G, T) = (\Omega_8^-(2), \operatorname{PSp}(8, 2)), (A_{14}, A_{16}), (\operatorname{PSL}(2, 8), A_9)$ or (A_{n-1}, A_n) with $n \geq 6$ and $n \mid 2^9 \cdot 3^2 \cdot 5$. If $(G, T) \neq (A_5, A_6)$ then $A = S_6$, contrary to the fact that A has a non-trivial radical. Thus, $n \geq 8$

Let $C = C_I(R)$. Then $G \leq C$ as $B = R \times G \leq I$. Clearly, $C \leq I$ and $C \cap R \leq Z(C)$. Since $G \cong R \times G/R \leq CR/R$ and $C/C \cap R \cong CR/R \leq I/R \cong T$, we have $C \cap R = Z(C)$, $C/Z(C) \cong T$ and I = CR. Thus, $C'/C' \cap Z(C) \cong C'Z(C)/Z(C) = (C/Z(C))' = C/Z(C) \cong T$, and so $Z(C') = C' \cap Z(C)$, C = C'Z(C) and $C'/Z(C') \cong T$. Furthermore, C' = (C'Z(C))' = C'', implying that C' is a covering group of T. It follows $G \leq C'$ because C/C' is abelian.

By [18, Theorem 5.1.4] and the Atlas [2, p. 123], $\operatorname{Mult}(A_n) = \mathbb{Z}_2$ for $n \geq 8$ and $\operatorname{Mult}(\operatorname{PSp}(8,2)) = 1$. Since C' is a covering group of T, we have Z(C') = 1 or \mathbb{Z}_2 .

Suppose $Z(C') = \mathbb{Z}_2$. Then $(G,T) = (A_{14}, A_{16})$, $(\operatorname{PSL}(2,8), A_9)$, or (A_{n-1}, A_n) with $n \geq 8$ and $n \mid 2^9 \cdot 3^2 \cdot 5$. Furthermore, C' is the unique double cover of A_n with $n \geq 8$, that is, $C' = 2.A_n$. Since $Z(C') \leq I$ and $I/R \cong T$ is simple, we have $Z(C') \leq R$, and hence $G \times Z(C') \leq C'$ as $B = G \times R$.

Let $(G,T)=(A_{14},A_{16})$. Then $C'=2.A_{16}$. Clearly, A_{14} has no proper subgroup of index less than 13, and by MAGMA, it also has no subgroup of index 15 or 16. This implies that each subgroup of A_{16} isomorphic to A_{14} fixes exactly two vertices in $\{1,2,\cdots,16\}$, and so lies in a subgroup of index 16 of A_{16} . It follows that $G\times Z(C')$ lies in a subgroup of index 16 of C', that is, $G\times Z(C')\leq L$ and |C':L|=16 for some $L\leq C'$. By Proposition 2.6, $L\cong 2.A_{15}$ and $G\times Z(C')\cong \mathbb{Z}_2\times A_{14}$, which is impossible because $G\times Z(C')\cong A_{14}\times \mathbb{Z}_2$.

Let $(G,T) = (\operatorname{PSL}(2,8), A_9)$. Then $C' = 2.A_9 = G \cdot C'_v$, and so $3^2 \mid |C'_v|$ as $|C'| = 2^7 \cdot 3^4 \cdot 5 \cdot 7$. By Proposition 2.1, $C'_v \cong A_4 \times A_5$, $(A_4 \times A_5) \rtimes \mathbb{Z}_2$, $S_4 \times S_5$, $\operatorname{ASL}(2,4)$ or $\operatorname{AGL}(2,4)$. Note that C' has a subgroup H isomorphic to $2.A_6$ with $Z(C') \leq H$. Since $H \cong \operatorname{SL}(2,9)$, Z(C') is the unique subgroup of order 2 in H. Since $Z(C') \leq C'$, we have $Z(C') \nleq C'_v$. Suppose that $|H_w|$ is odd for any $w \in V(\Gamma)$. Then $|w^H| = |H|/|H_w| = 2^4 \cdot (3^2 \cdot 5/|H_w|)$, implying that the orbit of H containing w has length a multiple of 2^4 , and so $|V(\Gamma)|$ is a multiple of 2^4 , contrary to the fact $|V(\Gamma)| \mid |\operatorname{PSL}(2,8)| = 2^3 \cdot 3^2 \cdot 7$. Thus, we may assume that H_v is even. Since $H_v = H \cap C'_v$ and Z(C') is the unique subgroup of order 2 in H, we have $Z(C') \leq H_v$ and hence $Z(C') \leq C'_v$, a contradiction.

Let $(G,T)=(A_{n-1},A_n)$ with $n \geq 8$. Then $G \times Z(C')$ is a subgroup of index n of C' isomorphic to $A_{n-1} \times \mathbb{Z}_2$, which is impossible by Proposition 2.6.

The above contradictions give rise to Z(C') = 1. Then $C' \cong T$ and $G < C' \unlhd I$. Since |I| = |I/R||R| = |T||R| = |C'||R| and $C' \cap R = 1$, we have $I = C' \times R$. Then C' is characteristic in I, and so $C' \unlhd A$ because $I \unlhd A$. It follows that A has a non-abelian simple normal subgroup C' such that G < C' and $G(G,C') \cong G(G,T) = G(G_8^-(2), PSp(8,2))$, G(G,G) = G(G,G) or G(G,G) = G(G,G) o

Now, we are ready to prove Theorem 1.1.

The proof of Theorem 1.1: Let G be a non-abelian simple group and Γ a connected pentavalent symmetric G-vertex-transitive graph with $v \in V(\Gamma)$. Let $A = \operatorname{Aut}(\Gamma)$ and R the radical of A. To prove the theorem, we may assume $G \not \supseteq A$. If R = 1, the theorem is true by Lemma 3.6. In what follows we assume $R \neq 1$. If $G = A_5$, by Lemma 3.1, $\Gamma = K_6$ and $A = S_6$, which is impossible because A has trivial radical. Thus, $G \neq A_5$.

Set B = RG. Then $G \cap R = 1$ and |B| = |R||G|. By Lemma 3.2, $|R| = |B_v|/|G_v| | 2^9 \cdot 3^2 \cdot 5$.

Suppose that R has one or two orbits on $V(\Gamma)$. Since $|G| = |V(\Gamma)||G_v|$, we have $|R| = |v^R||R_v| = |G||R_v|/|G_v|$ or $|G||R_v|/(2 \cdot |G_v|)$, and since R and G_v are $\{2, 3, 5\}$ -groups, G is a non-abelian $\{2, 3, 5\}$ -simple group. By Proposition 2.4, $G = A_6$ or PSU(4, 2) as $G \neq A_5$, and since $|R| = |B_v|/|G_v|$, we have $|B_v| = |G||R_v|$ or $|G||R_v|/2$. Note that $5 \mid |G|$, and so $5 \mid |B_v|$, that is, Γ is B-symmetric. If $R_v = 1$, then $|B_v| = |G|$ or |G|/2, that is, $|B_v| = 2^3 \cdot 3^2 \cdot 5$, $2^2 \cdot 3^2 \cdot 5$, $2^6 \cdot 3^4 \cdot 5$ or $2^5 \cdot 3^4 \cdot 5$, of which all are impossible by Proposition 2.1. If $R_v \neq 1$ then $5 \mid |R_v|$ as $R \leq A$, and in this case, $5^2 \mid |B_v| = |G||R_v|$, a contradiction.

Now we have shown that R has at least three orbits. By Lemma 3.7, if $B = R \times G$ then the theorem is true. To finish the proof, we only need to show $B = R \times G$.

By Proposition 2.2, R is semiregular on $V(\Gamma)$ and the quotient graph Γ_R is a connected pentavalent A/R-symmetric graph with $A/R \leq \operatorname{Aut}(\Gamma_R)$. Moreover, Γ_R is B/R-vertextransitive. Since R is the largest solvable normal subgroup of A, A/R has trivial radical, by Lemma 3.6, $B/R \leq A/R$ or A/R has a non-abelian simple normal subgroup I/R such that $B/R \leq I/R$ and $(B/R, I/R) \cong (G, T)$ with $(G, T) = (\Omega_8^-(2), \operatorname{PSp}(8, 2)), (A_{14}, A_{16}), (\operatorname{PSL}(2, 8), A_9)$ or (A_{n-1}, A_n) with $n \geq 6$ and $n \mid 2^9 \cdot 3^2 \cdot 5$.

Suppose $B/R \subseteq A/R$. Then $B \subseteq A$ and $5 \mid |B_v|$. Note that $1 \neq |R| = |B_v|/|G_v|$. Then $5 \mid |R|$ or $5 \mid |G_v|$. By Lemmas 3.3 and 3.5, $B = R \times G$, that is, G is characteristic in B, and hence $G \subseteq A$, contrary to the assumption $G \not \supseteq A$. Thus, $B/R \subseteq I/R \subseteq A/R$. It follows $(I/R)_{\alpha} \neq 1$ for $\alpha \in V(\Gamma_R)$ and hence $5 \mid |(I/R)_{\alpha}|$. Thus, Γ_R is I/R-symmetric. Since $|V(\Gamma_R)| = |V(\Gamma)|/|R| = |G|/(|R||G_v|)$ and $|T| = |I/R| = |V(\Gamma_R)||(I/R)_{\alpha}| = |G|/(|R||G_v|) \cdot |(I/R)_{\alpha}|$, we have $|(I/R)_{\alpha}| = |R||T||G_v|/|G|$, and by Proposition 2.1, $|R||T||G_v|/|G|$ is a divisor of $2^9 \cdot 3^2 \cdot 5$.

Since $|R| \mid 2^9 \cdot 3^2 \cdot 5$, we may write $|R| = 2^m \cdot 3^n \cdot 5^k$, where $0 \le m \le 9$, $0 \le n \le 2$ and $0 \le k \le 1$. Since R is solvable, there exists a series of subgroups of B:

$$B > R = R_s > \cdots > R_1 > R_0 = 1$$

such that $R_i \subseteq B$ and R_{i+1}/R_i is an elementary abelian r-group with $0 \le i \le s-1$, where r=2, 3 or 5. Clearly, $G \le B$ has a natural action on R_{i+1}/R_i by conjugation.

Suppose to the contrary $B \neq R \times G$. Then there exists some $0 \leq j \leq s-1$ such that $GR_i = G \times R_i$ for any $0 \leq i \leq j$, but $GR_{j+1} \neq G \times R_{j+1}$. If G acts trivially on R_{j+1}/R_j by conjugation, then $[GR_j/R_j, R_{j+1}/R_j] = 1$. Since $GR_j/R_j \cong G$ is simple, we have $(GR_j/R_j) \cap (R_{j+1}/R_j) = 1$. Note that $|GR_{j+1}/R_j| = |GR_{j+1}/R_{j+1}||R_{j+1}/R_j| = |G||R_{j+1}/R_j| = |GR_j/R_j||R_{j+1}/R_j|$. Then $GR_{j+1}/R_j = GR_j/R_j \times R_{j+1}/R_j$. In particular, $GR_j \subseteq GR_{j+1}$ and so $G \subseteq GR_{j+1}$ because $GR_j = G \times R_j$ implies that G is characteristic in GR_j . It follows that $GR_{j+1} = G \times R_{j+1}$, a contradiction. Thus, G acts non-trivially on R_{j+1}/R_j , and the simplicity of G implies that G acts faithfully on R_{j+1}/R_j .

Set $|R_{j+1}/R_j| = r^{\ell}$. Since $|R| = 2^m \cdot 3^n \cdot 5^k$, we have $\ell \leq m \leq 9$ for r = 2, $\ell \leq 2$ for r = 3 and $\ell \leq 1$ for r = 5. Recall that G acts faithfully on R_{j+1}/R_j and R_{j+1}/R_j is elementary abelian. Then $G \leq \operatorname{GL}(\ell, r)$. If r = 3 or 5 then $G \leq \operatorname{GL}(2, 3)$ or $\operatorname{GL}(1, 5)$, yielding that G is solvable, a contradiction. Thus, $G \leq \operatorname{GL}(m, 2) = \operatorname{PSL}(m, 2)$ with $m \leq 9$. Recall that $(G, T) = (\Omega_8^-(2), \operatorname{PSp}(8, 2)), (A_{14}, A_{16}), (\operatorname{PSL}(2, 8), A_9)$ or (A_{n-1}, A_n) with $n \geq 6$ and $n \mid 2^9 \cdot 3^2 \cdot 5$.

Let $G = A_{14}$ or A_{n-1} with $n \mid 2^9 \cdot 3^2 \cdot 5$ and $n \geq 12$. Since $11 \mid |A_{n-1}|$ and $11 \nmid |PSL(9,2)|$, we have $G \nleq PSL(9,2)$, a contradiction.

Let $G = \operatorname{PSL}(2,8)$ or A_7 , since $|G| = |V(\Gamma)||G_v|$ and R is semiregular on $V(\Gamma)$, we have $|R| \mid |G|/|G_v|$, and so $m \leq 2$ because Γ_R cannot be of odd order. It follows that $A_7 \leq \operatorname{PSL}(2,2)$ or $\operatorname{PSL}(2,8) \leq \operatorname{PSL}(2,2)$, a contradiction. Similarly, for $G = A_9$, we have $m \leq 5$ and $A_9 \leq \operatorname{PSL}(5,2)$, which is also impossible because $3^4 \mid |A_9|$ and $3^4 \nmid |\operatorname{PSL}(5,2)|$.

Since $G \neq A_5$, the left pairs are $(G, T) = (A_8, A_9)$ or $(\Omega_8^-(2), \operatorname{PSp}(8, 2))$. For $(G, T) = (A_8, A_9)$, since $|(I/R)_{\alpha}| = |R||T||G_v|/|G|$ and $5 \mid |(I/R)_{\alpha}|, |R|$ or $|G_v|$ is divisible by 5, by Lemmas 3.3 and 3.5, $B = R \times G$, a contradiction. Thus, $(G, T) = (\Omega_8^-(2), \operatorname{PSp}(8, 2))$.

Since $|T|/|G| = 2^4 \cdot 3 \cdot 5$ and $|R||T||G_v|/|G|$ is a divisor of $2^9 \cdot 3^2 \cdot 5$, |R| is a divisor of $2^5 \cdot 3$. Then $m \leq 5$ and $G \leq \mathrm{PSL}(5,2)$, which is impossible because 17 $|\Omega_8^-(2)|$ and $17 \nmid |\mathrm{PSL}(5,2)|$.

The above contradictions imply $B = R \times G$, as required. This completes the proof of Theorem 1.1.

The proof of Corollary 1.2: Let G be a non-abelian simple group and Γ a connected pentavalent G-symmetric graph with $v \in V(\Gamma)$. Let $A = \operatorname{Aut}(\Gamma)$ and let R the radical of A. To prove the corollary, we may assume $G \not \supseteq A$. By Theorem 1.1, A has a non-abelian simple normal subgroup T such that G < T and G > T is as given in Theorem 1.1. Since G is symmetric and $G < T \unlhd A$, $|G_v|$ and $|T_v|$ are divisible by 5, and since $|T|/|G| = |T_v|/|G_v|$, by Proposition 2.1 $(G,T) = (A_{n-1},A_n)$ with $n \ge 6$, $n \mid 2^9 \cdot 3^2$ and $n \ne 2^9, 2^8, 2^7, 2^6, 2^9 \cdot 3, 2^8 \cdot 3$.

Let $(G,T) = (A_{2^3-1}, A_{2^3})$ or (A_{3^2-1}, A_{3^2}) . By Lemma 3.4, $G \subseteq A$, a contradiction.

Let $(G,T) = (A_{2\cdot 3^2-1}, A_{2\cdot 3^2})$. By Proposition 2.1, $(G_v, T_v) \cong (F_{20} \times \mathbb{Z}_2, A_4 \times A_5)$ or $(F_{20} \times \mathbb{Z}_4, (A_4 \times A_5) \rtimes \mathbb{Z}_2)$. However, by MAGMA, $A_4 \times A_5$ or $(A_4 \times A_5) \rtimes \mathbb{Z}_2$ has no subgroup isomorphic to $F_{20} \times \mathbb{Z}_2$ or $F_{20} \times \mathbb{Z}_4$, respectively, a contradiction.

Let $T = A_6$. Then $G = A_5$ and by Lemma 3.4, $\Gamma = K_6$ and $A = S_6$. To finish the proof, we only need to show that for $n = 2^5, 2^7 \cdot 3, 2^7 \cdot 3^2, 2^8 \cdot 3^2$ and $2^9 \cdot 3^2$, we have A = T. To do this, it suffices to show $A_v = T_v$ because $A = TA_v$.

Let $n=2^9\cdot 3^2$ or $2^8\cdot 3^2$. By Proposition 2.1, $(G_v,T_v)\cong (\mathbb{Z}_5,\mathbb{Z}_2^6\rtimes \Gamma \mathrm{L}(2,4))$ or $(D_5,\mathbb{Z}_2^6\rtimes \Gamma \mathrm{L}(2,4))$, respectively. This forces $A_v=T_v$ because $T_v\leq A_v$, as required.

Let $n = 2^7 \cdot 3^2$. By Proposition 2.1, $(G_v, T_v) \cong (\mathbb{Z}_5, \mathrm{A}\Gamma\mathrm{L}(2,4))$, $(F_{20}, \mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4))$ or $(D_{10}, \mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4))$. Clearly, if $(G_v, T_v) \cong (F_{20}, \mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4))$ or $(D_{10}, \mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4))$, then $T_v = A_v$. Assume $(G_v, T_v) \cong (\mathbb{Z}_5, \mathrm{A}\Gamma\mathrm{L}(2,4))$. If $T_v \neq A_v$, then by Proposition 2.1 we have $A_v = \mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4)$, and since $T \subseteq A$, we have $T_v \subseteq A_v$. This implies that $\mathbb{Z}_2^6 \rtimes \Gamma\mathrm{L}(2,4)$ has a normal subgroup isomorphic to $\mathrm{A}\Gamma\mathrm{L}(2,4)$, which is impossible by [30, Theorem 1.1] and MAGMA. Thus, $T_v = A_v$, as required.

Let $n=2^7\cdot 3$. By Proposition 2.1, $(G_v,T_v)\cong (\mathbb{Z}_5,\mathrm{A}\Sigma\mathrm{L}(2,4))$ or $(A_5,\mathbb{Z}_2^6\rtimes\Gamma\mathrm{L}(2,4))$. For the latter, $A_v=T_v$. Assume $(G_v,T_v)\cong (\mathbb{Z}_5,\mathrm{A}\Sigma\mathrm{L}(2,4))$. If $A_v\neq T_v$ then by Proposition 2.1 we have $A_v=\mathbb{Z}_2^6\rtimes\Gamma\mathrm{L}(2,4)$ or $\mathrm{A}\Gamma\mathrm{L}(2,4)$, which are impossible because $\mathbb{Z}_2^6\rtimes\Gamma\mathrm{L}(2,4)$ and $\mathrm{A}\Gamma\mathrm{L}(2,4)$ have no normal subgroup isomorphic to $\mathrm{A}\Sigma\mathrm{L}(2,4)$ by [30, Theorem 1.1] and MAGMA. Thus, $T_v=A_v$, as required.

Let $n = 2^5$. By Proposition 2.1, $(G_v, T_v) \cong (A_5, A\Sigma L(2, 4))$ or $(A_4 \times A_5, \mathbb{Z}_2^6 \rtimes \Gamma L(2, 4))$. A similar argument to the above paragraph implies $T_v = A_v$, as required. This completes the proof of Corollary 1.2.

The proof of Corollary 1.3: Let G be a non-abelian simple group and Γ a connected pentavalent G-regular graph with $v \in V(\Gamma)$. Let $A = \operatorname{Aut}(\Gamma)$ and R the radical of A. To prove the corollary, we may assume $G \not \supseteq A$. By Theorem 1.1, A contains non-abelian simple normal subgroup T containing G such that (G,T) is given in Theorem 1.1. Since G is regular and $G < T \subseteq A$, $G_v = 1$ and T is symmetric. Since $|T|/|G| = |T_v|/|G_v| = |T_v|$, by Proposition 2.1, $(G,T) = (A_{n-1},A_n)$ with $n = 2 \cdot 5$, $2^2 \cdot 5$, $2^3 \cdot 5$, $2 \cdot 3 \cdot 5$, $2^4 \cdot 5$, $2^3 \cdot 3 \cdot 5$,

$$2^4 \cdot 3^2 \cdot 5, \ 2^6 \cdot 3 \cdot 5, \ 2^5 \cdot 3^2 \cdot 5, \ 2^7 \cdot 3 \cdot 5, \ 2^6 \cdot 3^2 \cdot 5, \ 2^7 \cdot 3^2 \cdot 5 \text{ or } 2^9 \cdot 3^2 \cdot 5.$$

To end the paper, we give an example to show that the pair $(G,T) = (A_{39}, A_{40})$ in Corollary 1.3 can happen.

Example 3.8 Let $G = A_{39}$ and $T = A_{40}$. Define $x, y, z, g \in T$ as following:

```
x = (1\ 2\ 7\ 18\ 5)(3\ 11\ 8\ 17\ 14)(4\ 9\ 22\ 33\ 15)(6\ 20\ 19\ 10\ 21)\ (12\ 28\ 26\ 31\ 23) (13\ 27\ 24\ 32\ 30)(16\ 35\ 34\ 25\ 36)(29\ 39\ 38\ 40\ 37),
```

- $y = (1\ 3\ 12\ 6)(2\ 8\ 23\ 10)(4\ 13\ 29\ 16)(5\ 17\ 28\ 19)(7\ 14\ 31\ 20)(9\ 24\ 37\ 25)$ $(11\ 26\ 21\ 18)(15\ 32\ 39\ 34)\ (22\ 30\ 40\ 35)(27\ 38\ 36\ 33),$
- $z = (1\ 4)(2\ 9)(3\ 13)(5\ 15)(6\ 16)(7\ 22)(8\ 24)(10\ 25)(11\ 27)(12\ 29)(14\ 30)$ (17\ 32) (18\ 33)(19\ 34)(20\ 35)(21\ 36)(23\ 37)(26\ 38)(28\ 39)(31\ 40),
- $g = (1\ 7)(2\ 24)(3\ 14)(4\ 40)(5\ 11)(6\ 20)(8\ 37)(9\ 10)(12\ 31)(13\ 35)(15\ 36)(16\ 30)$ $(17\ 26)(18\ 19)(21\ 28)(22\ 29)(23\ 25)(27\ 39)(32\ 33)(34\ 38).$

By MAGMA[1], $H = \langle x, y, z \rangle \cong F_{20} \times \mathbb{Z}_2$, $T = \langle H, g \rangle$, $|H : H \cap H^g| = 5$ and H is regular on $\{1, 2, \dots, 40\}$. Thus, T = GH with $G \cap H = 1$ and the coset graph $\Gamma = \text{Cos}(T, H, HgH)$ is a connected pentavalent T-symmetric G-regular graph, where G and T are viewed as groups of automorphisms of Γ by right multiplication.

By Corollary 1.3, $T \subseteq \operatorname{Aut}(\Gamma)$. Again by MAGMA [1], $\operatorname{Aut}(T, H, HgH) \cong H$, where \tilde{H} is the automorphism group of T induced by conjugation of elements in H. Thus, $\operatorname{Aut}(\Gamma) = T$ by [29, Lemma 2.10].

Acknowledgement: This work was supported by the National Natural Science Foundation of China (11571035, 11231008, 11271012) and by the 111 Project of China (B16002).

References

- [1] W. Bosma, J. Cannon, and C. Playoust, The MAGMA algebra system I: The user language, J. Symbolic Comput. 24 (1997), 235-265.
- [2] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, Atlas of Finite Group, Clarendon Press, Oxford, 1985.
- [3] X.G. Fang, L.J. Jia, J. Wang, On the automorphism groups of symmetric graphs admitting an almost simple group, Europ. J. Combin. 29 (2008) 1467-1472.
- [4] X. G. Fang, C. H. Li, and M. Y. Xu, On edge-transitive Cayley graphs of valency four, Europ. J. Combin. 25 (2004), 1107-1116.
- [5] X. G. Fang, X. S. Ma, and J. Wang, On locally primitive Cayley graphs of finite simple groups, J. Combin. Theory Ser. A 118 (2011), 1039-1051.

- [6] X. G. Fang and C. E. Praeger, Finite two-arc transitive graphs admitting a Suzuki simple group, Comm. Algebra 27 (1999), 3727-3754.
- [7] X.G. Fang and C.E. Praeger, Finite two-arc-transitive graphs admitting a Ree simple group, Comm. Algebra 27 (1999) 3755-3769.
- [8] X. G. Fang, C. E. Praeger, and J. Wang, On the automorphism group of Cayley graphs of finite simple groups, J. London Math. Soc. 66 (2002), 563-578.
- [9] Y.-Q. Feng, C.H. Li, J.-X. Zhou, Symmetric cubic graphs with sovable automorphism groups, Europ. J. Combin. 45 (2015) 1-11.
- [10] C. D. Godsil, On the full automorphism group of a graph, Combinatorica 1 (1981), 243-256.
- [11] S.-T. Guo and Y.-Q. Feng, A note on pentavalent s-transitive graphs, Discrete Math. 312 (2012), 2214-2216.
- [12] S.-T. Guo, J.-X. Zhou, and Y.-Q. Feng, Pentavalent symmetric graphs of order 12p, Electronic J. Combin. 18 (2011), 1-13 \$\pm\$P233.
- [13] P. Hall, A note on soluble groups, J. London Math. Soc 3 (1928), 98-105.
- [14] X.-H. Hua, Y.-Q. Feng, and J. Lee, Pentavalent symmetric graphs of order 2pq, Discrete Math. 311 (2011), 2259-2267.
- [15] B. Huppert, Eudiche Gruppen I, Springer-Verlag, 1967.
- [16] B. Huppert and N. Blackburn, Finite Groups III, Springer-Verlag, 1982.
- [17] B. Huppert and W. Lempken, Simple groups of order divisible by at most four primes, Proc. of the F. Scorina Gemel State University 16 (2000), 64-75.
- [18] P. Kleidman and M. Liebeck, The Subgroup Structure of The Finite Classical Groups, Cambridge Univ. Press, Cambridge, 1990.
- [19] C. H. Li, Isomorphisms of finite Cayley graphs, Ph.D. Thesis, The University of Western Australia, 1996.
- [20] C. H. Li, Z. P. Lu, H. Zhang, Tetravalent edge-transitive Cayley graphs with odd number of vertices, J. Combin. Theory Ser. B 96 (2006), 164-181.
- [21] P. Lorimer, Vertex-transitive graphs: symmetric graphs of prime valency, J. Graph Theory 8 (1984), 55-68.
- [22] R. C. Miller, The trivalent symmetric graphs of girth at most six, J. Combin. Theory Ser. B 10 (1971), 163-182.
- [23] C.E. Praeger, On a reduction theorem for finite, bipartite, 2-arc-transitive graphs, Australas. J. Combin. 7 (1993), 21-36.

- [24] C.E. Praeger, An O'Nan-Scott theorem for finite quasiprimitive permutation groups and an application to 2-arc transitive graphs, J. London Math. Soc. 47 (1993), 227-239.
- [25] C.E. Praeger, Finite transitive permutation groups and finite vertex-transitive graphs, in: Graph Symmetry: Algebraic Methods and Applications, in: NATO Adv. Sci. Inst. Ser. C, 497 (1997), 277-318.
- [26] C.E. Praeger, Imprimitive symmetric graphs, Ars Combin. A 19 (1985), 149-163.
- [27] B. O. Sabidussi, Vertex-transitive graphs, Monash Math. 68 (1964), 426-438.
- [28] J. Schur, Über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen, J. Reine Angew. Math. 127 (1904), 20-50.
- [29] Y. Wang, Y.-Q. Feng, J.-X. Zhou, Automorphism of Cayley digraph of 2-genetic groups of prime-power order, J. Combin Theory Ser. A, 143 (2016), 88-106.
- [30] R.M. Weiss, Presentations for (G, s)-transitive graphs of small valency, Math. Proc. Cambridge Philos. Soc. 101 (1987), 7-20.
- [31] R. A. Wilson, The Finite Simple Groups, Springer-Verlag, London, 2009.
- [32] S. J. Xu, X. G. Fang, J. Wang, and M. Y. Xu, On cubic s-arc transitive Cayley graphs of finite simple groups, Europ. J. Combin. 26 (2005), 133-143.
- [33] S. J. Xu, X. G. Fang, J. Wang, and M. Y. Xu, 5-arc transitive cubic Cayley graphs on finite simple groups, Europ. J. Combin. 28 (2007), 1023-1036.