

SOME ALGEBRAIC INVARIANTS OF EDGE IDEAL OF CIRCULANT GRAPHS

GIANCARLO RINALDO

ABSTRACT. Let G be the circulant graph $C_n(S)$ with $S \subseteq \{1, \dots, \lfloor \frac{n}{2} \rfloor\}$ and let $I(G)$ be its edge ideal in the ring $K[x_0, \dots, x_{n-1}]$. Under the hypothesis that n is prime we : 1) compute the regularity index of $R/I(G)$; 2) compute the Castelnuovo-Mumford regularity when $R/I(G)$ is Cohen-Macaulay; 3) prove that the circulant graphs with $S = \{1, \dots, s\}$ are sequentially S_2 . We end characterizing the Cohen-Macaulay circulant graphs of Krull dimension 2 by n and S and computing their Cohen-Macaulay type and Castelnuovo-Mumford regularity.

INTRODUCTION

Let $S \subseteq \{1, 2, \dots, \lfloor \frac{n}{2} \rfloor\}$. The *circulant graph* $G := C_n(S)$ is a graph with vertex set $\mathbb{Z}_n = \{0, \dots, n-1\}$ and edge set $E(G) := \{\{i, j\} \mid |j - i|_n \in S\}$ where $|k|_n = \min\{|k|, n - |k|\}$.

Let $R = K[x_0, \dots, x_{n-1}]$ be the polynomial ring on n variables over a field K . The *edge ideal* of G , denoted by $I(G)$, is the ideal of R generated by all square-free monomials $x_i x_j$ such that $\{i, j\} \in E(G)$. Edge ideals of a graph have been introduced by Villarreal [11] in 1990, where he studied the Cohen-Macaulay property of such ideals. Many authors have focused their attention on such ideals (see [8], [6]). A known fact about Cohen-Macaulay edge ideals is that they are well-covered.

A graph G is said *well-covered* if all the maximal independent sets of G have the same cardinality. Recently well-covered circulant graphs have been studied (see [1], [2], [9]). In [13] and [4] the authors studied well-covered circulant graphs that are Cohen-Macaulay.

In this article we put in relation the values n, S of a circulant graph $C_n(S)$ and algebraic invariants of $R/I(G)$. In particular we study the regularity index, the Castelnuovo-Mumford regularity, the Cohen-Macaulayness and Serre's condition of $R/I(G)$.

In the first section we recall some concepts and notations and preliminary notions.

In the second section under the hypothesis that n is prime we observe that the regularity index of $R/I(G)$ is 1 obtaining as a by-product the

Castelnuovo-Mumford regularity of the ring when it is Cohen-Macaulay. Moreover we prove that each k -skeleton of the simplicial complex of the independent set of G is connected. As an application we prove that the circulant graphs $C_n(\{1, \dots, s\})$ (studied in [1], [2], [4], [9], [11],[13]) are sequentially S_2 (see [7]).

In the third section we characterize the Cohen-Macaulay circulant graphs of Krull dimension 2 by n and S and compute their Cohen-Macaulay type and Castelnuovo-Mumford regularity. We observe, as in the previous section, the depth of $R/I(G)$ is equal to its Castelnuovo-Mumford regularity.

1. PRELIMINARIES

In this section we recall some concepts and notations on graphs and on simplicial complexes that we will use in the article. Let G be a simple graph with vertex set $V(G)$ and the edge set $E(G)$. A subset C of $V(G)$ is called a *clique* of G if for all i and j belonging to C with $i \neq j$ one has $\{i, j\} \in E(G)$. A subset A of $V(G)$ is called an *independent set* of G if no two vertices of A are adjacent. The *complement graph* \bar{G} of G is the graph with vertex set $V(\bar{G}) = V(G)$ and edge set $E(\bar{G}) = \{\{u, v\} \in V(G)^2 \mid \{u, v\} \notin E(G)\}$.

Set $V = \{x_1, \dots, x_n\}$. A *simplicial complex* Δ on the vertex set V is a collection of subsets of V such that

- (i) $\{x_i\} \in \Delta$ for all $x_i \in V$;
- (ii) $F \in \Delta$ and $G \subseteq F$ imply $G \in \Delta$.

An element $F \in \Delta$ is called a *face* of Δ . A maximal face of Δ with respect to inclusion is called a *facet* of Δ .

If Δ is a simplicial complex with facets F_1, \dots, F_q , we call $\{F_1, \dots, F_q\}$ the facet set of Δ and we denote it by $\mathcal{F}(\Delta)$. The dimension of a face $F \in \Delta$ is $\dim F = |F| - 1$, and the dimension of Δ is the maximum of the dimensions of all facets in $\mathcal{F}(\Delta)$. If all facets of Δ have the same dimension, then Δ is called *pure*. Let $d-1$ the dimension of Δ and let f_i be the number of faces of Δ of dimension i with the convention that $f_{-1} = 1$. Then the f -vector of Δ is the d -tuple $f(\Delta) = (f_{-1}, f_0, \dots, f_{d-1})$. The h -vector of Δ is $h(\Delta) = (h_0, h_1, \dots, h_d)$ with

$$h_k = \sum_{i=0}^k (-1)^{k-i} \binom{d-i}{k-i} f_{i-1}.$$

The sum

$$\tilde{\chi}(\Delta) = \sum_{i=-1}^{d-1} (-1)^i f_i$$

is called the reduced Euler characteristic of Δ and $h_d = (-1)^{d-1} \tilde{\chi}(\Delta)$. Given any simplicial complex Δ on V , we can associate a monomial ideal

I_Δ in the polynomial ring R as follows:

$$I_\Delta = (\{x_{j_1}x_{j_2}\cdots x_{j_r} : \{x_{j_1}, x_{j_2}, \dots, x_{j_r}\} \notin \Delta\}).$$

R/I_Δ is called Stanley-Reisner ring and its Krull dimension is d . If G is a graph we call the *independent complex* of G by

$$\Delta(G) = \{A \subset V(G) : A \text{ is an independent set of } G\}.$$

The *clique complex* of a graph G is the simplicial complex whose faces are the cliques of G . Let \mathbb{F} be the minimal free resolution of the quotient ring $R/I(G)$. Then

$$\mathbb{F} : 0 \rightarrow F_p \rightarrow \cdots \rightarrow F_{p-1} \rightarrow \cdots \rightarrow F_0 \rightarrow R/I(G) \rightarrow 0$$

with $F_i = \bigoplus_j R(-j)^{\beta_{ij}}$. The numbers β_{ij} are called the Betti numbers of \mathbb{F} .

The Castelnuovo–Mumford regularity of $R/I(G)$, denoted by $\text{reg } R/I(G)$, is defined by

$$\text{reg } R/I(G) = \max\{j - i : \beta_{ij} \neq 0\}.$$

A graph G is said Cohen-Macaulay if the ring $R/I(G)$, or equivalently $R/I_{\Delta(G)}$ is Cohen-Macaulay (over the field K) (see [3], [10], [16]). The Cohen-Macaulay type of $R/I(G)$ is equal to the last total Betti number in the minimal free resolution \mathbb{F} .

We end this section with the following

Remark 1.1. Let $T = \{1, 2, \dots, \lfloor \frac{n}{2} \rfloor\}$ and G be a circulant graph on $S \subseteq T$ with $s = |S|$, then:

- (1) \bar{G} is a circulant graph on $\bar{S} = T \setminus S$;
- (2) The clique complex of \bar{G} is the independent complex of G , $\Delta(G)$;
- (3)

$$|E(G)| = \begin{cases} ns - \frac{n}{2} & \text{if } n \text{ is even and } \frac{n}{2} \in S \\ ns & \text{otherwise.} \end{cases}$$

2. REGULARITY AND CONNECTEDNESS OF THE INDEPENDENT COMPLEX OF CIRCULANT GRAPHS OF PRIME ORDER

We recall some basic facts about the regularity index (see also [14]). Let R be standard graded ring and I be a homogeneous ideal. The *Hilbert function* $H_{R/I} : \mathbb{N} \rightarrow \mathbb{N}$ is defined by

$$H_{R/I}(k) := \dim_K(R/I)_k$$

and the Hilbert-Poincaré series of R/I is given by

$$\text{HP}_{R/I}(t) := \sum_{k \in \mathbb{N}} H_{R/I}(k)t^k.$$

By Hilbert-Serre theorem, the Hilbert-Poincaré series of R/I is a rational function, that is

$$\text{HP}_{R/I}(t) = \frac{h(t)}{(1-t)^n}.$$

There exists a unique polynomial such that $H_{R/I}(k) = P_{R/I}(k)$ for all $k \gg 0$. The minimum integer $k_0 \in \mathbb{N}$ such that $H_{R/I}(k) = P_{R/I}(k) \forall k \geq k_0$ is called *regularity index* and we denote it by $\text{ri}(R/I)$.

Remark 2.1. *Let R/I_Δ be a Stanley-Reisner ring. Then*

$$\text{ri}(R/I_\Delta) = \begin{cases} 0 & \text{if } h_d = 0 \\ 1 & \text{if } h_d \neq 0 \end{cases}$$

Proof. By the hypothesis the Hilbert series can be represented by the reduced rational function

$$\frac{h(t)}{(1-t)^d}$$

where d is the Krull dimension of R/I_Δ and $h(t) = \sum_{i=0}^d h_i t^i$ where h_i are the entries of the h -vector of Δ . We observe that $\text{ri}(R/I) = \max(0, \deg h(t) - d + 1)$. If $\text{ri}(R/I_\Delta) > 0$ then $\deg h(t) > d - 1$. But since $\deg h(t) \leq d$ we have $\deg h(t) = d$. Therefore $h_d \neq 0$ and $\text{ri}(R/I_\Delta) = 1$. The other case follows by the same argument. \square

Lemma 2.2. *Let G be a circulant graph on S with n prime. Then the entries of the f -vector of $\Delta(G)$ are*

$$f_i = n f'_i$$

with $0 \leq i \leq d - 1$ and $f'_i = f_{i,0}/(i + 1) \in \mathbb{N}$ where $f_{i,0}$ is the number of faces of dimension i containing the vertex 0.

Proof. Call $\mathcal{F}_i \subset \Delta$ the set of faces of dimension i , that is

$$\mathcal{F}_i = \{F_1, \dots, F_{f_i}\}.$$

Let $f_{i,j}$, number of faces in \mathcal{F}_i containing a given vertex $j = 0, \dots, n - 1$. Since G is circulant

$$f_{i,j} = f_{i,0} \text{ for all } j \in \{0, \dots, n - 1\}.$$

Let $A \in \mathbb{F}_2^{f_i \times n} = (a_{jk})$ be the incidence matrix with $a_{jk} = 1$ if the vertex $k - 1$ belongs to the facet F_j and 0 otherwise. We observe that each row has exactly $i + 1$ 1-entries. Hence summing the entries of the matrix we have $(i + 1)f_i$. Moreover each column has exactly $f_{i,j}$ non zero entries. That is

$$n f_{i,0} = (i + 1) f_i.$$

Since n is prime the assertion follows. \square

Theorem 2.3. *Let G be a circulant graph on S with n prime. Then*

$$\text{ri}(R/I(G)) = 1.$$

Proof. By Remark 2.1 it is sufficient to show that h_d is different from 0. Since

$$|h_d| = \left| \sum_{i=0}^d (-1)^i f_{i-1} \right| \neq 0,$$

it is sufficient to show that the reduced Euler formula is different from 0, that is

$$\sum_{i=1}^d (-1)^i f_{i-1} \neq 1.$$

By Lemma 2.2 we obtain

$$\sum_{i=1}^d (-1)^i f_{i-1} = n \sum_{i=1}^d (-1)^i f'_{i-1}$$

since n is prime and the assertion follows. \square

Remark 2.4. *In the proof of Theorem 2.3 we are giving a partial positive answer to the Conjecture 5.8 of [9] that states that for all circulant graphs $\tilde{\chi}(\Delta) \neq 0$.*

Corollary 2.5. *Let G be a circulant graph on S with n prime that is Cohen-Macaulay. Then $\text{reg } R/I(G) = \text{depth } R/I(G)$.*

Proof. By Corollary 4.8 of [5] since $\text{ri}(R/I) = 1$ the assertion follows. \square

We end this section studying nice properties of the independent complex $\Delta(G)$ of a circulant graph G that have prime order. We start by the following

Definition 2.6. *Let Δ be a simplicial complex then we define the pure simplicial complexes $\Delta^{[k]}$ whose facets are*

$$\mathcal{F}(\Delta^{[k]}) = \{F \in \Delta : \dim(F) = k\}, \quad 0 \leq k \leq \dim(\Delta).$$

One interesting property of Cohen-Macaulay ring R/I_Δ is that the each simplicial complex $\Delta^{[k]}$ is connected. Hence the following Lemma is of interest.

Lemma 2.7. *Let G be a circulant graph on S with n prime. Then the k -skeleton of the simplicial complex Δ , $\Delta^{[k]}$ is connected for every $k \geq 1$.*

Proof. To prove the claim we find a Hamiltonian cycle connecting all the vertices in $V = \{0, \dots, n-1\}$ of the 1-skeleton of $\Delta^{[k]}$. Then it follows that since the 1-skeleton is connected then $\Delta^{[k]}$ is connected, too.

We assume without loss of generality that $F_0 = \{v_0, v_1, \dots, v_k\} \in \Delta^{[k]}$ such that $v_0 = 0$, $v_1 = s \in S$. We define the set

$$F_j = \{v_{0,j}, v_{1,j}, \dots, v_{k,j}\}$$

with $v_{i,j} = v_i + js \pmod n$. It is easy to observe that since F_0 is in $\Delta^{[k]}$ and G is circulant, F_j is in $\Delta^{[k]}$, too.

Moreover if we focus on the first two vertices of F_j we obtain that

$$v_{1,j} = v_{0,j-1} \text{ for all } j = 1, \dots, n-1,$$

and $v_{0,n-1} = v_{1,0}$. Since

$$v_{0,j} = js \pmod n$$

the set $\{v_0, \dots, v_{n-1}\}$, by the primality of n , is equal to V . Hence the cycle with vertices

$$v_{0,0}, v_{0,1}, \dots, v_{0,n-1}$$

and edges

$$\{v_{0,0}, v_{0,1}\}, \dots, \{v_{0,n-2}, v_{0,n-1}\}, \{v_{0,n-1}, v_{0,0}\}$$

is a Hamiltonian cycle and the assertion follows. \square

Recall that a finitely generated graded module M over a Noetherian graded K -algebra R is said to satisfy the Serre's condition S_r if

$$\text{depth } M_{\mathfrak{p}} \geq \min(r, \dim M_{\mathfrak{p}}),$$

for all $\mathfrak{p} \in \text{Spec}(R)$.

Definition 2.8. *Let M be a finitely generated \mathbb{Z} -graded module over a standard graded K -algebra R where K is a field. For a positive integer r we say that M is sequentially S_r if there exists a finite filtration of graded R -modules*

$$0 = M_0 \subset M_1 \subset \dots \subset M_r = M$$

such that each M_i/M_{i-1} satisfies the S_r condition and the Krull dimensions of the quotients are increasing:


$$\dim(M_1/M_0) < \dim(M_2/M_1) < \dots < \dim(M_t/M_{t-1}).$$

A nice characterization of sequentially S_2 simplicial complexes is the following:

Theorem 2.9 ([7]). *Let Δ be a simplicial complex with vertex set V . Then Δ is sequentially S_2 if and only if the following conditions hold:*

- (1) $\Delta^{[i]}$ is connected for all $i \geq 1$;
- (2) $\text{link}_{\Delta}(x)$ is sequentially S_2 for all $x \in V$.

Example 2.10. *Let G be the circulant graph $C_6(\{1\})$. Then its simplicial complex Δ is connected, but $\Delta^{[2]}$ is not (in the picture from left to right are $C_6(\{1\})$, Δ and $\Delta^{[2]}$).*



Theorem 2.11. *Let G be the circulant graph $C_n(\{1, \dots, s\})$ with n prime. Then G is sequentially S_2 .*

Proof. By Lemma 2.7 the first condition of Theorem 2.9 is satisfied. To check the second condition of Theorem 2.9 we prove that $\text{link}_\Delta(x_0)$ is sequentially Cohen Macaulay. Localizing in x_0 we obtain

$$R' = K[\text{link}_\Delta x_0] \cong K[x_{s+1}, \dots, x_{n-s-1}]/I(G')$$

where $I(G') = I(G) \cap K[x_{s+1}, \dots, x_{n-s-1}]$. This fact follows observing that the vertices adjacent to 0 are $\{1, \dots, s\} \cup \{n-s, \dots, n-1\}$. We claim that $I(G')$ is chordal, hence it is sequentially Cohen-Macaulay by Theorem 3.2 of [6]. To prove the claim we observe that the labelling on the vertices of G'

$$s+1, s+2, \dots, n-s-1$$

induces a perfect elimination ordering, that is $N^+(i) = \{j : \{i, j\} \in E(G'), i < j\}$ is a clique. Let $j, k \in N^+(i)$. That is $\{i, j\}$ and $\{i, k\}$ are two edges with $i < j$ and $i < k$ and assume $j < k$. Then $|j-i|_n = j-i \leq s$ and $|k-i|_n = k-i \leq s$. Moreover

$$0 < j-i < k-i \leq s.$$

Hence it follows $|k-j| = k-j < s$. Therefore $\{j, k\} \in E(G')$ and $N^+(i)$ is a clique. \square

Example 2.12. *If a ring is Cohen-Macaulay it is pure and sequentially S_n for all n . The circulant graph of prime order with minimum number of vertices that is Cohen-Macaulay and has Krull dimension greater than 2 is $C_{13}(\{1, 5\})$ (see [4]).*

3. COHEN-MACAULAY CIRCULANT GRAPHS OF DIMENSION 2 AND THEIR CASTELNUOVO-MUMFORD REGULARITY

We start this section by the following

Theorem 3.1. *Let G be the circulant graph $C_n(S)$ with $S \subset \{1, \dots, \lfloor \frac{n}{2} \rfloor\}$. The following conditions are equivalent:*

- (1) G is Cohen-Macaulay of dimension 2;
- (2) $\Delta(G)$ is connected of dimension 1;
- (3) $\gcd(n, \bar{S}) = 1$ and $\forall a, b, c \in \bar{S} \ a + b + c \neq n$.

Proof. (1) \Leftrightarrow (2). Known fact. See also [9] Corollary 4.54.

(2) \Rightarrow (3). If $\Delta(G)$ is connected then there is a path in $\bar{G} \cong \Delta(G)$ connecting the vertices 0 and 1 (see Remark 1.1) whose vertices are

$$0 = v_0, v_1, \dots, v_r = 1$$

and edges

$$\{0, s_1\}, \{s_1, s_1 + s_2\}, \dots, \left\{ \sum_{i=1}^{r-1} s_i, \sum_{i=1}^r s_i \equiv 1 \pmod{n} \right\}$$

with $s_i \in \bar{S}$. Hence there exists a relation

$$\sum a_i s_i \equiv 1 \pmod{n}, \text{ with } a_i \in \mathbb{N}, s_i \in \bar{S}.$$

By the Euclidean algorithm we have that $\gcd(n, \bar{S}) = 1$.

Suppose there exist $a, b, c \in \bar{S}$ with $a + b + c = n$. We claim that $\{0, a, a+b\}$ is a clique in $\Delta(G)$, that is $\dim \Delta(G) \geq 2$. In fact since $a, c \in \bar{S}$ then $\{0, a\}, \{n-c, (n-c)+c\} = \{0, n-c\} \in \Delta(G)$. It is sufficient to show that $\{a, n-c\} \in \Delta(G)$. The claim follows since $b \in \bar{S}$ and $n-c = a+b$ and the graph is circulant. The other direction follows by the same argument. \square

Corollary 3.2. *Let G be the circulant graph $C_n(S)$ with $\{1, \dots, d\} \subseteq S \subset \{1, \dots, \lfloor \frac{n}{2} \rfloor\}$ with $n \leq 3d + 2$ and $\gcd(n, \bar{S}) = 1$. Then G is Cohen-Macaulay.*

Proof. It is sufficient to observe $\min \bar{S} \geq d+1$ and $3(d+1) \geq n+1$ satisfying condition (3) in Theorem 3.1. \square

Theorem 3.3. *Let G be a Cohen-Macaulay circulant graph $C_n(S)$ of dimension 2. Then $\text{reg } R/I(G) = 2$.*

Proof. It is sufficient to prove that $h_2 \neq 0$ (see Remark 2.1 and the proof of Corollary 2.5). We need to compute $h_2 = f_1 - f_0 + f_{-1}$. We observe that f_1 is the number of edges of \bar{G} . By Remark 1.1 one of the two cases to study is

$$\binom{n}{2} - ns,$$

with $h_2 = \binom{n}{2} - n(s+1) + 1$. The only roots $n \in \mathbb{N}$ of the quadratic equation

$$\binom{n}{2} - n(s+1) + 1 = 0$$

are 1 and 2 with $s = 0$. Absurd. The other case follows by the same argument. \square

Theorem 3.4. *Let G be a Cohen-Macaulay circulant graph $C_n(S)$ of dimension 2. Then its Cohen-Macaulay type is*

$$h_2 = \begin{cases} \binom{n}{2} - n(s + \frac{1}{2}) + 1 & \text{if } n \text{ is even and } \frac{n}{2} \in S \\ \binom{n}{2} - n(s + 1) + 1 & \text{otherwise.} \end{cases}$$

Proof. By Auslander-Buchsbaum Theorem (Theorem 1.3.3, [3]) and since the depth $R/I(G) = 2$ we need to compute the Betti number in position $\beta_{i,j}$ when $i = n - 2$. By Theorem 3.3 and the definition of Castelnuovo-Mumford regularity, the Betti numbers $\beta_{n-2,j}$ to check are in the degrees $j \in \{n - 1, n\}$. We recall the Hochster's formula (see [10], Corollary 5.1.2)

$$\beta_{i,\sigma}(R/I_\Delta) = \dim_K \tilde{H}_{|\sigma|-i-1}(\Delta_{|\sigma}; K)$$

where $\tilde{H}(\cdot)$ is the simplicial homology and $\sigma \in \Delta$ is interpreted as squarefree degree in the minimal free resolution and it induces a restriction in Δ defined by

$$\Delta_{|\sigma} = \{F \in \Delta : F \subseteq \sigma\}.$$

We observe that in the squarefree degree σ having total degree $n - 1$

$$\beta_{i,\sigma} = \dim_K \tilde{H}_0(\Delta_{|\sigma}; K) = 0.$$

In fact $\Delta \cong \bar{G}$ is connected and the same happens removing one of the vertices of the circulant graph \bar{G} since circulant graphs are biconnected. Now, if we consider the squarefree degree σ having total degree n , again, by Hochster formula, we obtain

$$\beta_{i,\sigma} = \dim_K \tilde{H}_1(\Delta_{|\sigma}; K).$$

In this case $\Delta_{|\sigma} \cong \Delta \cong \bar{G}$ and the chain complex of Δ

$$\mathcal{C} : 0 \rightarrow C_1 \xrightarrow{\partial_1} C_0 \xrightarrow{\partial_0} C_{-1} \rightarrow 0,$$

has the two homologies $\tilde{H}_0 = \tilde{H}_{-1} = 0$. Therefore

$$\dim_K \tilde{H}_1(\bar{G}; K) = \beta_{i,\sigma} = f_1 - f_0 + f_{-1}$$

and the assertion follows by Remark 1.1. \square

Remark 3.5. *We observe that the rings satisfying Theorem 3.4 are level. For a description of level algebras see Chapter 5.4 and 5.7 of [3].*

Corollary 3.6. *Let G be the circulant graph $C_n(S)$ with $S \subset \{1, \dots, \lfloor \frac{n}{2} \rfloor\}$ and $s = |S|$. The following conditions are equivalent:*

- (1) G is Gorenstein of dimension 2;
- (2) $S = \{1, \dots, \hat{i}, \dots, n\}$ and $\gcd(n, i) = 1$ with $n \geq 4$;
- (3) $\Delta(G) \cong \bar{G}$ is a n -gon with $n \geq 4$.

Proof. (1) \Rightarrow (2). G is Gorenstein if and only if G is Cohen-Macaulay of type 1. Hence by Theorem 3.1 $\Delta(G)$ is connected that is $\gcd(n, \bar{S}) = 1$. Moreover by Theorem 3.4 $h_2 = 1$ and solving the two quadratic equations

$$\binom{n}{2} - n(s + \frac{1}{2}) + 1 = 1, \quad \binom{n}{2} - n(s + 1) + 1 = 1,$$

we obtain respectively

$$n = 2s + 2 \text{ and } n = 2s + 3.$$

In both cases $s = \lfloor \frac{n}{2} \rfloor - 1$. Hence $\bar{S} = i$ with $\gcd(i, n) = 1$ and the assertion follows.

(2) \Rightarrow (3). Let $\bar{S} = \{i\}$ with $\gcd(n, i) = 1$. We easily observe that the vertices

$$0, i, \dots, (n-1)i \bmod n$$

and edges

$$\{0, i\}, \{i, 2i\}, \dots, \{(n-1)i, (n)i \equiv 0 \bmod n\}$$

define a Hamiltonian cycle that is \bar{G} itself.

(3) \Rightarrow (1). Since $\Delta(G)$ is a simplicial 1-sphere is Gorenstein of Krull dimension 2 (see Corollary 5.6.5 of [3]). \square

We observe that in Theorem 4.1 of [4] the Cohen-Macaulayness of the graphs described in Corollary 3.6 has been studied by a different point of view.

REFERENCES

- [1] J. Brown, R. Hoshino, *Independence polynomials of circulants with an application to music*, Discrete Mathematics, **309**, 2009, 2292–2304.
- [2] J. Brown, R. Hoshino, *Well-covered circulant graphs*, Discrete Mathematics, **311**, 2011, 244–251.
- [3] W. Bruns, J. Herzog, *Cohen-Macaulay rings*, Cambridge Univ. Press, Cambridge, 1997.
- [4] J. Earl, K. N. Vander Meulen, A. Van Tuyl, *Independence Complexes of Well-Covered Circulant Graphs*, Experimental Mathematics, **25**, 2016, 441–451.
- [5] D. Eisenbud, *The Geometry of Syzygies*, Graduate texts in Mathematics, Springer, 2005.
- [6] C. A. Francisco, A. Van Tuyl, *Sequentially Cohen-Macaulay edge ideals*, Proc. Amer. Math. Soc., **135**, 2007, 2327–2337.
- [7] H. Haghighi, N. Terai, S. Yassemi, R. Zaare-Nahandi, *Sequentially S_r simplicial complexes and sequentially S_2 graphs*, Proc. Amer. Math. Soc., **139**, 2011, 1993–2005.
- [8] J. Herzog, T. Hibi, *Distributive lattices, bipartite graphs and Alexander duality*, J. Alg. Combin., **22**, 2005, 289–302.
- [9] R. Hoshino, *Independence polynomials of circulant graphs*, PhD Thesis, Dalhousie University, 2008, 1–280.
- [10] E. Miller, B. Sturmfels, *Combinatorial Commutative Algebra*, Springer-Verlag, Berlin, 2005.
- [11] A. Mousivand, *Circulant S_2 graphs*, preprint arXiv:1512.08141v1, 2015, 1–11.

- [12] R. P. Stanley, *Combinatorics and Commutative Algebra, Second Edition*, Birkhäuser, Boston/Basel/Stuttgart, 1996.
- [13] K. N. Vander Meulen, A. Van Tuyl, C. Watt, *Cohen-Macaulay circulant graphs*, *Communications in Algebra*, **42**, 2014, 1896–1910.
- [14] W. Vasconcelos, *Computational methods in commutative algebra and algebraic geometry*, Springer Science & Business Media, **2**, 2004.
- [15] R. H. Villarreal, *Cohen–Macaulay graphs*, *Manuscripta Math.*, **66** 1990, 277–293.
- [16] R. Villarreal, *Monomial algebras*, Marcel Dekker, New-York, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, VIA SOMMARIVE, 14,
38123 POVO (TRENTO), ITALY