

Conditional Lyapunov Exponent Criteria in terms of Ergodic Theory

Masaru Shintani* and Ken Umeno†
Department of Applied Mathematics and Physics,
Graduate School of Informatics, Kyoto University,
Yoshida Honmachi Sakyo-ku, Kyoto, Japan, 606-8501
(Dated: February 10, 2019)

The conditional Lyapunov exponent is defined for investigating chaotic synchronization, in particular complete synchronization and generalized synchronization. We find that the conditional Lyapunov exponent is expressed as a formula in terms of ergodic theory. Dealing with this formula, we find what factors characterize the conditional Lyapunov exponent in chaotic systems.

PACS numbers: 05.45.-a,02.60.Cb,05.45.Xt,05.40.Ca

I. INTRODUCTION

The conditional Lyapunov exponent is defined for investigating chaotic synchronization [1–12], in particular Complete synchronization (CS) and Generalized synchronization (GS). Transitions from desynchronization to synchronization of trajectories occur when the conditional Lyapunov exponent changes from positive to negative [13–15]. Although it is widely known that the chaotic synchronization occurs in many systems, it is not clearly known why the conditional Lyapunov exponent changes. For example, it has not completely been clarified why the CS occurs in chaotic systems. A report [10] showed that an external forcing input in CS may change the dynamical system to another one. The forcing input changes the balance between phase contraction or expansion, and the CS occurs when such contraction dominates. Although the explanation is well considered, there could be another reason why the conditional Lyapunov exponent may change. Furthermore, they have focused on the mean of external forcing inputs for the CS. The study [11] said that they get the transversal Lyapunov exponent with changing the noise distribution. These may be important. However we expect that more precise information on phase space can lead to more relevant analysis. Therefore, we would like to clarify the relation between the conditional Lyapunov exponent and chaotic synchronization in CS.

We have two claims through this research. The first one is about a formula to express the conditional Lyapunov exponent in terms of ergodic theory, and the second is what factors influence the conditional Lyapunov exponent in chaotic dynamical systems. For simplicity we only focus on the CS.

In Section 2, we describe the definition of conditional Lyapunov exponent in chaotic systems. Furthermore, we describe two main claims in our research.

In Sections 3 - 5, we construct solvable chaotic dynamical systems, and confirm the our claims by analysis for such systems.

In Appendix A, we summarize a short introduction of ergodic theory.

II. DEFINITION AND MAIN CLAIM

To discuss the conditional Lyapunov exponent, we consider the following simple one-dimensional unidirectional coupling system:

$$x(t+1) = f(x(t)) + \xi(t) \equiv \psi(x(t), \xi(t)), \quad (1)$$

where t is time, $x(t)$ is a response f is a chaotic mapping, and $\xi(t)$ is an external forcing driver. We define $P_X(X)$ as the probability distribution of variables X in the dynamical system, and also define \mathbb{R}_X as the range in which variables X are defined.

We define the CS of the system as follows [8–11, 13]. We consider two different trajectories $x_1(t)$ and $x_2(t)$ in Eq. (1) with different initial points. To judge whether the response system exhibit CS, we introduce the conditional Lyapunov exponent. The occurrence of CS implies that the difference between $x_1(t)$ and $x_2(t)$ decreases as time increases. The CS is said to occur when the following condition about synchronization error $e'(t) \equiv |x_2(t) - x_1(t)|$:

$$\lim_{t \rightarrow \infty} e'(t) = 0, \quad (2)$$

is satisfied for almost all initial points. Here $|\dots|$ expresses Euclidean norm of the argument. The interpretation of (2) is that the state in the large t limit is independent of any initial state for almost all initial points. The infinitesimal synchronization error $e(T)$ in system (1) is defined as:

$$e(T) := \prod_{t=1}^T \left| \frac{\partial \psi(x, \xi)}{\partial x} \right|_{x=x(t)} e(0). \quad (3)$$

When CS occurs, the equation $\lim_{t \rightarrow \infty} e'(t) = \lim_{T \rightarrow \infty} |e(T)| = 0$ is satisfied. It is noted here that $\xi(t)$ in Eq. (1) affects the time-evolution of $x(t)$ and contributes to $e(T)$.

* shintani.masaru.28a@kyoto-u.jp

† umeno.ken.8z@kyoto-u.ac.jp

The conditional Lyapunov exponent for variables x is defined by

$$\lambda := \lim_{T \rightarrow \infty} \frac{1}{T} \ln \frac{|e(T)|}{|e(0)|}.$$

Clearly λ expresses the stability of the state $x_1(t) = x_2(t)$. In this paper, the CS is said to occur when $\lambda < 0$, and we do not use Eq. (2) directly. The conditional Lyapunov exponent exists when the system (1) has the absolutely continuous invariant measure according to Osledelets' multiplicative ergodic theorem.

When the system (1) is ergodic with respect to the absolutely continuous invariant measure, λ is expressed as the ensemble average:

$$\begin{aligned} \lambda &= \lim_{T \rightarrow \infty} \frac{1}{T} \ln \frac{|e(T)|}{|e(0)|} \\ &= \int_{\mathbb{R}_x} \int_{\mathbb{R}_\xi} P(x, \xi) \ln \left| \frac{\partial \psi(x, \xi)}{\partial x} \right| d\xi dx, \end{aligned} \quad (4)$$

where, $P(x, \xi)$ is the invariant distribution of the system (1). In what follows we assume that the system $x(t+1) = f(x(t))$ has a mixing property (thus ergodic) with the absolutely continuous invariant ergodic measure $\mu(dx) = P_x(x)dx$, ($x \in \mathbb{R}_x$) with some P_x , and $P(x, \xi)$ is a continuous density function in x and ξ satisfying $\int_{\mathbb{R}_x} \int_{\mathbb{R}_\xi} P(x, \xi) d\xi dx = 1$.

Then, we give our two main claims. Firstly, the conditional Lyapunov exponent is expressed by the following equation.

$$\lambda = \tilde{\lambda}, \quad (5)$$

where

$$\lambda = \int_{\mathbb{R}_x} \int_{\mathbb{R}_\xi} P(x, \xi) \ln \left| \frac{\partial \psi(x, \xi)}{\partial x} \right| d\xi dx \quad (6)$$

$$\tilde{\lambda} = \int_{\mathbb{R}_\xi} P_\xi(\xi_0) \lambda_\Psi(\xi_0) d\xi_0. \quad (7)$$

Note that although ξ_0 is originally a constant value, we consider the distribution of ξ_0 in the above equation. Here how to obtain $\lambda_\Psi(\xi_0)$ is as follows. First, by calculating the Lyapunov exponent $\lambda_\Psi(\xi_0)$ for the auxiliary system:

$$y(t+1) = \Psi_{\xi_0}(y(t)) \quad (8)$$

with $\Psi_{\xi_0}(y) = f(y) + \xi_0$. Then, by changing ξ_0 continuously, we have the set $\{\lambda_\Psi(\xi_0) | \xi_0 \in \mathbb{R}_\xi\}$. Note that the function Ψ_{ξ_0} is an one-dimensional mapping with the parameter ξ_0 , while $\psi(y, \xi_0)$ is a two-dimensional function.

Equation (5) implies that the conditional Lyapunov exponent λ is expressed as *the ensemble average of the set of unique Lyapunov exponents* $\{\lambda_\Psi(\xi_0) | \xi_0 \in \mathbb{R}_\xi\}$ provided the ergodicity in the system (8). This is the first main claim in our research.

Then, we show how we get this claim provided the ergodicity in the system (1). The Lyapunov exponent

$\lambda_\Psi(\xi_0)$ can be obtained by the following equation with the ergodicity in the system (8),

$$\lambda_\Psi(\xi_0) = \int_{\mathbb{R}_y} P_y(y|\xi_0) \ln \left| \frac{d\Psi_{\xi_0}(y)}{dy} \right| dy, \quad (9)$$

where $P(y|\xi_0)$ is the unique *conditional probability distribution* in dynamical system (8) with each constant value ξ_0 . To obtain Eq. (7) we use the marginalization about random variables a and b in the probability theory:

$$P(a, b) = P(b)P(a|b). \quad (10)$$

Then, substituting (10) into (6), we have

$$\begin{aligned} \lambda &= \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} P_\xi(\xi_0) P_y(y|\xi_0) \ln \left| \frac{d\Psi_{\xi_0}(y)}{dy} \right| d\xi_0 dy \\ &= \int_{\mathbb{R}_\xi} P_\xi(\xi_0) \left(\int_{\mathbb{R}_y} P_y(y|\xi_0) \ln \left| \frac{d\Psi_{\xi_0}(y)}{dy} \right| dy \right) d\xi_0 \\ &= \int_{\mathbb{R}_\xi} P_\xi(\xi_0) \lambda_\Psi(\xi_0) d\xi_0 = \tilde{\lambda}. \end{aligned} \quad (11)$$

Then, $\lambda = \tilde{\lambda}$ is satisfied when the invariant distribution $P_y(y|\xi)$ exists. This is satisfied when the system (8) is ergodic for any ξ_0 . Therefore, our first claim Eq. (5) is derived from the mixing property of (1) and the ergodicity of ξ and the system (8). Figure 1 illustrates this concept. Here, $\langle \dots \rangle$ expresses the ensemble average with respect to the ergodic invariant measure.




FIG. 1. Our approach for the conditional Lyapunov exponent

As for the second claim, with the first one, we can derive that the conditional Lyapunov exponent in a system (1) is characterized by only two factors, a dynamical system and a distribution of external forcing input.

III. EXAMPLE1 FOR THE FIRST CLAIM

In order to confirm the first claim, we would like to show its relevance by analyzing solvable chaotic synchronization [16, 17]. Here, a solvable chaotic synchronizing

system is such that an invariant measure, the conditional Lyapunov exponent and the threshold between synchronization and desynchronization are *analytically* obtained for the unidirectional coupling system.

A. Definition of our system

As for the first claim, we need three steps to show it. Firstly, we analytically calculate the conditional Lyapunov exponent λ in our solvable chaotic dynamical system in accordance with the definition (4). Secondly, we analyze the auxiliary dynamical system like the Eq. (8), and get the exponent λ accordance with Eqs. (8) - (11). Thirdly, we confirm that the results obtained by the two types of analytical methods coincide.

We study the following dynamical system.

$$x(t+1) = f(x(t)) + \varepsilon\zeta(t), \quad (12)$$

where ε is a coupling parameter, and $\varepsilon\zeta(t)$ correspond to an external forcing input $\xi(t)$. Firstly we prepare the following solvable chaotic dynamical system [18–20]:

$$\begin{aligned} x(t+1) &= f(x(t)) + \varepsilon\zeta(t) \equiv \phi(x(t), \zeta(t)), \\ f(x(t)) &= g(x(t)) \equiv \frac{1}{2} \left(x(t) - \frac{1}{x(t)} \right), \\ \zeta(t+1) &= g(\zeta(t)), \end{aligned} \quad (13)$$

where the function g is associated with the double-angle formula given by $\cot 2\theta = \frac{1}{2} \left(\cot \theta - \frac{1}{\cot \theta} \right)$ [17]. The mapping associated with g is a chaotic mapping which has the mixing property [17], and its Lyapunov exponent is $\ln 2$. The invariant measure of the system $x(t+1) = g(x(t))$ is the standard Cauchy distribution as follows:

$$\mu(dx) = C(x; 0, 1)dx,$$

where $C(x)$ is Cauchy distribution defined as:

$$C(x; c, \gamma) \equiv \frac{\gamma}{\pi \{ (x-c)^2 + \gamma^2 \}},$$

with c being a median and γ a scale parameter. Therefore, the distribution of variable x in dynamical system $x(t+1) = g(x(t))$ follows $C(x; 0, 1)$ [17], and the distribution of external forcing ξ follows also $C(x; 0, 1)$. This is our first solvable chaotic dynamical system. We see that the conditional Lyapunov exponent changes when the coupling parameter is varied. We calculate the conditional Lyapunov exponent of this dynamical system by the definition (4) in accordance with the first step.

B. Conventional ergodic theoretical approach

We express the conditional Lyapunov exponent of the system (13) as $\lambda_g(\varepsilon)$ since it will turn out that the conditional Lyapunov exponent crucially depends on ε . The

conditional Lyapunov exponent $\lambda_g(\varepsilon)$ is expressed as:

$$\begin{aligned} \lambda_g(\varepsilon) &= \int_{\mathbb{R}_x} \int_{\mathbb{R}_\xi} P(x, \xi) \ln \left| \frac{\partial \phi(x, \xi)}{\partial x} \right| d\xi dx \\ &= \int_{\mathbb{R}_x} \int_{\mathbb{R}_\xi} P_{x,\xi}(x) P_\xi(\xi) d\xi \ln \left| \frac{1}{2} \left(1 + \frac{1}{x^2} \right) \right| dx \\ &= \int_{\mathbb{R}_x} P_{x,\xi}(x) \ln \left| \frac{1}{2} \left(1 + \frac{1}{x^2} \right) \right| dx \\ &= \int_{\mathbb{R}_x} P_{x,\varepsilon}(x) \ln \left| \frac{1}{2} \left(1 + \frac{1}{x^2} \right) \right| dx, \end{aligned}$$

where $P_{x,\varepsilon}(x)$ is the invariant distribution for the variable x in the superposed dynamical system $x(t+1) = g(x(t)) + \varepsilon\zeta(t)$ with a parameter ε . The external forcing ξ is dependent on the coupling parameter ε .

Then we firstly calculate the invariant distribution $P_{x,\varepsilon}(x)$, secondly the conditional Lyapunov exponent $\lambda_g(\varepsilon)$, and thirdly the threshold between synchronization and desynchronization analytically.

We can calculate the invariant distribution $P_{x,\varepsilon}(x)$ by using the following three properties, the mixing property for the mapping g (Property 1), the property of preserving the form of Cauchy distributions in terms of the Perron-Frobenius equation (PF equation) for g (Property 2), and property of the stable property for Lévy distributions (Property 3). Then showing these three properties, we explain how these properties play roles in order to get $P_{x,\varepsilon}(x)dx$.

As to Property 1, it is already known that the mapping g has mixing property [17].

As for Property 2, this property implies that g is the mapping which changes an input Cauchy distribution into *another* Cauchy distribution with a *different* median and scale parameter. We consider the PF equation for the equation $z = g(x)$ where the input variables x follows $C(x; c, \gamma)$. Since the g is a two-to-one mapping, $P_z(z)$ satisfies the following PF equation (Probability Preservation Relation):

$$P_z(z)|dz| = C(x_1; c, \gamma)|dx_1| + C(x_2; c, \gamma)|dx_2|$$

where x_1 and x_2 ($x_1 > x_2$) are the solutions of the quadratic equation $z = g(x)$. They satisfy the following,

$$\begin{cases} x_1 + x_2 = 2z \\ x_1 x_2 = -1. \end{cases}$$

With these relations, the probability distribution $P_z(z)$ is obtained, as the following rescaled Cauchy distribution:

$$P_z(z) = C(z; c', \gamma'),$$

where $c' = \frac{c(\gamma^2 + c^2 - 1)}{2(\gamma^2 + c^2)}$ and $\gamma' = \frac{\gamma(\gamma^2 + c^2 + 1)}{2(\gamma^2 + c^2)}$.

As to Property 3, this stable property has already widely been known. When distributions of two independent variables X_1 and X_2 obey a Lévy distribution family, the distribution of the variable $aX_1 + bX_2$ ($a, b \in \mathbb{R}$)

also obeys the same family. These three properties play roles in order to get the invariant distribution $P_{x,\varepsilon}(x)$.

We can prove that variables x and $\varepsilon\zeta$ in the system (13) obey Cauchy distributions respectively because of the three properties. In Appendix B, we describe these three properties in more detail. By taking into account the Properties 2 and 3, the distribution of the variable x is changed to a Cauchy distribution with a different median and scale parameter in every iteration if the initial input follows a Cauchy distribution. Hence, we get the following self-consistent recurrence equations about a median c and a scale parameter γ per iteration, as

$$\begin{cases} c(t+1) = \frac{c(t)(\gamma(t)^2 + c(t)^2 - 1)}{2(\gamma(t)^2 + c(t)^2)} \\ \gamma(t+1) = \frac{\gamma(t)(\gamma(t)^2 + c(t)^2 + 1)}{2(\gamma(t)^2 + c(t)^2)} + |\varepsilon|. \end{cases} \quad (14)$$

We get the following convergence values c^* and γ^* as the stable fixed point of Eq. (14) for $t \rightarrow \infty$:

$$\begin{aligned} c_g^* &= 0 \\ \gamma_g^* &= |\varepsilon| + \sqrt{\varepsilon^2 + 1}. \end{aligned}$$

Hence, we analytically obtain $P_{x,\varepsilon}(x)$ as the fixed point of the recurrence equation Eq. (14):

$$P_{x,\varepsilon}(x) = C(x; 0, \gamma_g^*)$$

We should emphasize the following. Although we assume the initial input follows a Cauchy distribution in the above analysis, we do not need any restriction for a distribution of initial point for the system (1) because of the mixing property (Property 1). Therefore, the invariant distribution $P_{x,\varepsilon}(x)$ is always obtained regardless of any initial points.

With $P_{x,\varepsilon}(x)$, we can calculate the conditional Lyapunov exponent $\lambda_g(\varepsilon)$ as follows:

$$\begin{aligned} \lambda_g(\varepsilon) &= \int_{\mathbb{R}_x} P_{x,\varepsilon}(x) \ln \left| \frac{1}{2} \left(1 + \frac{1}{x^2} \right) \right| dx \\ &= \int_{\mathbb{R}} \frac{\gamma_g^*}{\pi(x^2 + \gamma_g^{*2})} \ln \left| \frac{1}{2} \left(1 + \frac{1}{x^2} \right) \right| dx \\ &= 2 \ln \left(\frac{\gamma_g^* + 1}{\gamma_g^*} \right) - \ln 2 \\ &= 2 \ln (\sqrt{\varepsilon^2 + 1} - |\varepsilon| + 1) - \ln 2. \end{aligned}$$

Note that $\lambda_g(\varepsilon = 0) = \ln 2$. We can also get the threshold ε_g^* between the synchronization and desynchronization. This is the solution of $\lambda_g(\varepsilon_g^*) = 0$, which is

$$\varepsilon_g^* = 1.$$

Figure 2 illustrates that these analytical results coincide with the results of numerical simulation with the initial condition $x_0 = \sqrt{2}$ and $\xi_0 = \sqrt{3}$.




FIG. 2. Conditional Lyapunov exponent $\lambda_g(\varepsilon)$

C. Proposing ergodic theoretical approach

As above, we can calculate $\lambda_g(\varepsilon)$ by Eq. (4). The invariant distribution $P(x, \xi)$ is expressed as:

$$P(x, \xi) = \frac{\gamma_g^*}{\pi(x^2 + \gamma_g^{*2})} \frac{|\varepsilon|}{\pi(\xi^2 + \varepsilon^2)},$$

because $P_\xi(\xi)$ is derived from Property 3 for the variable ζ that obey $C(\zeta; 0, 1)$. Then in accordance with the second step, we would like to calculate $\tilde{\lambda}_g(\varepsilon)$ by Eqs. (8) - (11).

We prepare the following dynamical system by Eq. (8):

$$\begin{aligned} y(t+1) &= G_{\xi_0}(y(t)) \\ G_{\xi_0}(y) &= g(y) + \xi_0. \end{aligned}$$

We firstly need to confirm that this system is ergodic for every ξ_0 , which is the sufficient condition for that the conditional distribution $P_y(y|\xi_0)$ exists for every ξ_0 . Hence, we sufficiently calculate three factors, the conditional distribution $P_y(y|\xi_0)$, and the unique Lyapunov exponents $\lambda_\Psi(\xi_0)$, and the distribution of external forcing inputs $P_\xi(\xi_0)$.

Firstly, we calculate the conditional distribution $P_y(y|\xi_0)$. The invariant measure $P_y(y|\xi_0)dx$ is also obtained with Properties 1 - 3.

In the same way of getting $P_{x,\varepsilon}(x)$, we can prove that $P_y(y|\xi_0)$ also follows a Cauchy distribution. Hence, we get the following self-consistent recurrence equations about a median c and a scale parameter γ , as

$$\begin{cases} c(t+1) = \frac{c(t)(\gamma(t)^2 + c(t)^2 - 1)}{2(\gamma(t)^2 + c(t)^2)} + \xi_0 \\ \gamma(t+1) = \frac{\gamma(t)(\gamma(t)^2 + c(t)^2 + 1)}{2(\gamma(t)^2 + c(t)^2)}. \end{cases}$$

We get the following convergence values \hat{c} and $\hat{\gamma}$ as the

stable fixed point for $t \rightarrow \infty$:

$$\begin{cases} \hat{c} = \xi_0 \\ \hat{\gamma} = \sqrt{1 - \xi_0^2} \end{cases} \quad (\text{if } |\xi_0| < 1),$$

$$\begin{cases} \hat{c} = \xi_0 + \text{sgn}(\xi_0)\sqrt{\xi_0^2 - 1} \\ \hat{\gamma} = 0 \end{cases} \quad (\text{if } |\xi_0| \geq 1),$$

and $\text{sgn}(\xi_0) = \begin{cases} 1 & \text{if } \xi_0 > 0. \\ -1 & \text{if } \xi_0 < 0. \end{cases}$

Thus, we obtain the conditional distribution $P(y|\xi_0)$ as follows:

$$P_y(y|\xi_0) = \begin{cases} C(y; \xi_0, \sqrt{1 - \xi_0^2}) & (|\xi_0| < 1). \\ C(y; \xi_0 + \text{sgn}(\xi_0)\sqrt{\xi_0^2 - 1}, 0) & (|\xi_0| \geq 1). \end{cases}$$

As for unique Lyapunov exponents $\lambda_\Psi(\xi_0)$, we can calculate them as follows:

$$\begin{aligned} \lambda_G(\xi_0) &= \int_{\mathbb{R}_y} P_y(y|\xi_0) \ln \left| \frac{1}{2} \left(1 + \frac{1}{y^2} \right) \right| dy \\ &= \begin{cases} \ln(1 + \sqrt{1 - \xi_0^2}) & (|\xi_0| < 1) \\ \ln \left\{ 1 + \frac{1}{(\xi_0 + \text{sgn}(\xi_0)\sqrt{\xi_0^2 - 1})^2} \right\} - \ln 2 & (|\xi_0| \geq 1), \end{cases} \end{aligned}$$

Thus, we can calculate the exponent $\tilde{\lambda}_g(\varepsilon)$ according to Eq. (11):

$$\begin{aligned} \tilde{\lambda}_g(\varepsilon) &= \int_{\mathbb{R}_\xi} P_\xi(\xi_0) \lambda_G(\xi_0) d\xi_0 \\ &= \int_{\mathbb{R}} \frac{|\varepsilon|}{\pi(\xi_0^2 + \varepsilon^2)} \lambda_G(\xi_0) d\xi_0 \\ &= 2 \ln(\sqrt{\varepsilon^2 + 1} - |\varepsilon| + 1) - \ln 2 \\ &= \lambda_g(\varepsilon). \end{aligned}$$

As above, we confirm that our first main claim is certainly satisfied. This claim signifies that a conditional Lyapunov exponent is expressed as the ensemble average of the set of unique Lyapunov exponents of the auxiliary dynamical system. Figure 3 illustrates that two different trajectories with different initial points ($x_1(0) = 0.46, x_2(0) = 1.3$) synchronize for $t > 50$. The coupling parameter ε is 0.8.

IV. EXAMPLE2 FOR THE FIRST CLAIM

Our analysis is not restricted to the above example. We consider another solvable chaotic dynamical system, and can also confirm that the conditional Lyapunov exponent in the system is obtained by Eq. (5). This dy-




FIG. 3. Chaotic synchronization in the system (13)

namical system is defined as follows:

$$\begin{aligned} x(t+1) &= f(x(t)) + \varepsilon\zeta(t), \\ f(x(t)) &= h(x(t)) \equiv \frac{2x(t)}{1 - x(t)^2}, \\ \zeta(t+1) &= h(\zeta(t)) \\ H_{\xi_0}(y) &= h(y) + \xi_0. \end{aligned}$$

The chaotic mapping h is associated with the double formula $\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$ [17]. Its Lyapunov exponent is $\ln 2$, and the invariant measure obeys the standard Cauchy distribution. Then, the invariant distribution $P_{x,\xi}(x)$ and the conditional Lyapunov exponent $\lambda_h(\varepsilon)$, and the threshold ε_h^* are respectively given as:

$$\begin{aligned} P_{x,\xi}(x) &= C(x; 0, \gamma_h^*), \\ \lambda_h(\varepsilon) &= 2 \ln \left(\frac{\gamma_h^* + 1}{\gamma_h^{*2} + 1} \right) + \ln 2, \\ \varepsilon_h^* &= 0.78 \dots, \end{aligned}$$

where γ_h^* is the positive number which satisfies the following equation:

$$\gamma_h^* - |\varepsilon| \gamma_h^{*2} - \gamma_h^* - |\varepsilon| = 0.$$

V. EXAMPLE FOR THE SECOND CLAIM

Here, as for the second claim, the conditional Lyapunov exponent for a system (1) is characterized by only two factors, a dynamical system f and a distribution of external forcing input ξ .

In order to show this claim we consider six types of different dynamical systems. Each of them has a different chaotic mapping f or different system ξ as follows:

$$f(x) = \begin{cases} g(x) = \frac{1}{2} \left(x - \frac{1}{x} \right) \\ h(x) = \frac{2x}{1 - x^2} \end{cases}$$

$$\begin{cases} \zeta(t+1) = g(\zeta(t)) \\ \zeta(t+1) = h(\zeta(t)) \\ \zeta = \text{Crand}(0, 1), \end{cases}$$

where $\text{Crand}(c, \gamma)$ is a set of random numbers which follow $C(\zeta; c, \gamma)$. The algorithm to get $\text{Crand}(c, \gamma)$ follows:

$$\text{Crand}(0, 1) = \tan\left(\frac{\pi}{2}(U(-1 : 1))\right),$$

where $U(-1 : 1)$ are uniform random numbers on the interval $(-1 : 1)$.

We compare the conditional Lyapunov exponent of these systems with Table I. As we can see from the Table I, we can find that some factors do not influence the conditional Lyapunov exponent such as the property of time series in external forcing input. Then, we show that the conditional Lyapunov exponent is characterized by the original mapping f and the distribution of external forcing input ξ . The original dynamical system is changed to the system which can have an attracting fixed point because of the external forcing input. The conditional Lyapunov exponent is changed by the existence of attracting fixed points. Figures 4 and 5 illustrate that the maps G and H are changed to ones that have attracting fixed points by external forcing input respectively. Attractors of systems depend on how attracting




FIG. 4. Attracting fixed point is generated by changing ξ_0 in $G_{\xi_0}(y)$




FIG. 5. Attracting fixed point is generated by changing ξ_0 in $H_{\xi_0}(y)$

fixed points are generated. The created attractor leads to the ξ -dependence of $\lambda_{\Psi}(\xi_0)$. Figures 6 and 7 each show the set of Lyapunov exponents $\lambda_G(\xi_0)$ and $\lambda_H(\xi_0)$ respectively. Hence, the original mappings determine how

attracting fixed points are generated. Furthermore, with Eq. (5) from the first claim, the conditional Lyapunov exponent is calculated by the set of Lyapunov exponents $\{\lambda_{\Psi}(\xi_0) | \xi_0 \in \mathbb{R}_{\xi}\}$ and the distribution of the external forcing input P_{ξ} . As above, the conditional Lyapunov




FIG. 6. Unique Lyapunov exponents of $G_{\xi_0}(y)$




FIG. 7. Unique Lyapunov exponents of $H_{\xi_0}(y)$

exponent in system (1) is uniquely characterized by only two factors, the original dynamical system and the distribution of external forcing input.

VI. CONCLUSION

We have two main claims. First, the conditional Lyapunov exponent is expressed as Eq. (5) provided the ergodicity in dynamical systems. This yields that the conditional Lyapunov exponent is expressed as the ensemble average of the set of unique Lyapunov exponents of the auxiliary dynamical system (8). Second, the conditional Lyapunov exponent is characterized only two factors, an original dynamical system and a distribution of external forcing input. Then, although we consider CS in

TABLE I. Relations among the combination of f , ξ and the conditional Lyapunov exponents

| f | Generation Mechanism of ζ | Lyapunov exponent of f | The invariant distribution of the system $x(t+1) = f(x(t))$ | $P_\xi(\xi)$ | $P_{x,\xi}(x)$ | the property of time series in external forcing input ξ | λ |
|-----|---------------------------------|--------------------------|---|----------------------------|-----------------------|---|--------------------------|
| g | g | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_g^*)$ | chaotic | $\lambda_g(\varepsilon)$ |
| g | h | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_g^*)$ | chaotic | $\lambda_g(\varepsilon)$ |
| g | Crand(0,1) | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_g^*)$ | random | $\lambda_g(\varepsilon)$ |
| h | g | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_h^*)$ | chaotic | $\lambda_h(\varepsilon)$ |
| h | h | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_h^*)$ | chaotic | $\lambda_h(\varepsilon)$ |
| h | Crand(0,1) | $\ln 2$ | $C(x; 0, 1)$ | $C(\xi; 0, \varepsilon)$ | $C(x; 0, \gamma_h^*)$ | random | $\lambda_h(\varepsilon)$ |

one-dimensional unidirectionally coupled dynamical systems only, this claim will also hold in multi-dimensional chaotic systems in terms of ergodic theory.

ACKNOWLEDGMENTS

The authors thank Shin-itiro Goto (Kyoto University) for stimulating discussions.

-
- [1] H. Fujisaka, T. Yamada, Prog. Theor. Phys. **69**, 32 (1983)
- [2] L. M. Pecora, T. L. Carroll, Phys. Rev. Lett. **64**, 821 (1990)
- [3] L. Kocarev, U. Parlitz, Phys. Rev. Lett. **74**, 5028 (1995)
- [4] L. Kocarev, U. Parlitz, Phys. Rev. Lett. **76**, 1816 (1996)
- [5] L. Yu, E. Ott, Q. Chen, Phys. Rev. Lett. **65**, 2935 (1990).
- [6] A. S. Pikovsky, J. Kurths, Phys. Rev. Lett. **78**, 775 (1997)
- [7] A. S. Pikovsky, M. G. Rosenblum, G. V. Osipov, J. Kurths, Physica D: Nonlinear Phenomena **104**, 219 (1997)
- [8] A. Pikovsky, M. Rosenblum, J. Kurths, *Synchronization A Universal Concept in Nonlinear Sciences*, (Cambridge University Press, Cambridge, 2001)
- [9] S. Boccaletti, J. Bragard, F. T. Arecchi, H. Mancini, Phys. Rev. Lett. **83**, 536 (1999)
- [10] S. Boccaletti, J. Kurths, G. Osipov, D. Valladares, and C. Zhou, Physics reports **366**, 1 (2002).
- [11] S. Rim, D. U. Hwang, I. Kim, C. M. Kim, Phys. Rev. Lett. **85**, 2304 (2000).
- [12] A. Balanov, N. Janson, D. Postnov, O. Sonsnovtseva, *Synchronization: From Simple to Complex*, (2008)
- [13] S. Goto, K. Yoshimura, P Davis, IEICE Trans. Fundamentals, **91-A**, 2535 (2008)
- [14] S. Sunada, K. Arai, K. Yoshimura, M. Adachi, Phys. Rev. Lett. **112**, 204101 (2014)
- [15] A. Uchida, K. Umeno, National Congress of Theoretical and Applied Mechanics, **55**, 79 (2006) (In Japanese)
- [16] K. Umeno, Phys. Rev. E **55**, 5280 (1997)
- [17] K. Umeno, Phys. Rev. E **58**, 2644 (1998)
- [18] K. Umeno and K. Okubo, Prog. Theor. Phys. **2016**, 2, 021A01 (2016)
- [19] M. Shintani, K. Umeno, IEICE Technical Report, CCS, **115(178)**, 11 (2015) (In Japanese)
- [20] K. Umeno, IEICE-NOLTA, **7**, 14 (2016)

Appendix A: Appendix

In this Appendix a short introduction of ergodic theory is provided. Since this appendix aims at providing clear definitions of terminologies used in the main text, this appendix is not comprehensive. For a comprehensive review from a mathematical viewpoint see Refs.[1, 2], also from a physical viewpoint see Ref.[3].

1. Ergodic dynamical systems

Definition A.1. (*Dynamical system, [1]*): A dynamical system $(\mathcal{M}, \mu, \varphi_t)$ is a measure-space (\mathcal{M}, μ) equipped with a one-parameter group φ_t of automorphisms (except for spaces of measure-zero) of (\mathcal{M}, μ) , φ_t depending measurably of t . Here the parameter t denotes an integer.

Given a dynamical system $(\mathcal{M}, \mu, \varphi_t)$, it follows that $\mu(\varphi_t A) = \mu(A)$, where A is a measurable set, and that φ_t is a measurable in $\mathcal{M} \times \mathbb{R}$. In what follows $\mu(\mathcal{M}) = 1$ is assumed.

The following average often appears in physics.

Definition A.2. (*Time-average, [1]*): Let $(\mathcal{M}, \mu, \varphi_t)$ be a dynamical system, and f a complex-valued function defined on \mathcal{M} . If there exists the quantity

$$f^*(x) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(\varphi^n x), \quad x \in \mathcal{M}, \quad n \in \mathbb{Z}$$

then $f^*(x)$ is called the time-average of f .

Also the following average often appears and is related to the time-average for some dynamical systems.

Definition A.3. (*Space-average, [1]*): Let $(\mathcal{M}, \mu, \varphi_t)$ be a dynamical system, and f a complex-valued function defined on \mathcal{M} . If there exists the quantity

$$\bar{f} = \int_{\mathcal{M}} f(x) \mu(dx) \quad x \in \mathcal{M}, \quad n \in \mathbb{Z}$$

then \bar{f} is called the space-average of f .

Space-average defined above is called ensemble average in the main text. The following is used in the main text.

Definition A.4. (*Absolutely continuous with respect to the Lebesgue measure*): In Definition A.3, if $\mu(dx)$ is of the form

$$\mu(dx) = \rho(x) dx,$$

then ρ is called an absolutely continuous with respect to the Lebesgue measure dx .

We are now ready to state the definition of ergodic system.

Definition A.5. (*Ergodic system, [1]*): Let $(\mathcal{M}, \mu, \varphi_t)$ be a dynamical system. If the following condition is satisfied

$$f^*(x) = \bar{f},$$

for any integrable function f in the sense of $f \in L_1(\mathcal{M}, \mu)$, then the dynamical system is called an ergodic system.

This states that for an ergodic dynamical system, one can replace the time-average of f with the space-average of it. An example of how to apply this property is Eq. (4) in the main text.

The following property is a stronger property for dynamical systems.

Definition A.6. (*Mixing system, [1]*): Let $(\mathcal{M}, \mu, \varphi_t)$ be a dynamical system. If the condition

$$\lim_{t \rightarrow \infty} \mu[\varphi_t A \cap B] = \mu(A) \mu(B),$$

is satisfied for all measurable set A and B , then the dynamical system is called a mixing system.

Appendix B: Appendix

In this Appendix, we describe the Property 2 and 3 in more detail. About the Property 2, We get $P_z(z)$ in Section III-A as follows:

$$\begin{aligned} P_z(z) &= \frac{\gamma}{\pi((x_1 - c)^2 + \gamma^2)} \left| \frac{1}{\frac{dz}{dx_1}} \right| + \frac{\gamma}{\pi((x_2 - c)^2 + \gamma^2)} \left| \frac{1}{\frac{dz}{dx_1}} \right| \\ &= C(x_1; c, \gamma) \frac{2x_1^2}{x_1^2 + 1} + C(x_2; c, \gamma) \frac{2x_2^2}{x_2^2 + 1} \\ &= \frac{\frac{\gamma(\gamma^2 + c^2 + 1)}{2(\gamma^2 + c^2)}}{\pi \left\{ \left(z - \frac{c(\gamma^2 + c^2 - 1)}{2(\gamma^2 + c^2)} \right)^2 + \left(\frac{\gamma(\gamma^2 + c^2 + 1)}{2(\gamma^2 + c^2)} \right)^2 \right\}} \\ &= C(z; c', \gamma') \end{aligned}$$

This shows the mapping g rescales the median c and the scale parameter γ of the input Cauchy distribution as:

$$\begin{aligned} \text{median} &: c \rightarrow \frac{c(\gamma^2 + c^2 - 1)}{2(\gamma^2 + c^2)} \quad (\equiv c') \\ \text{scale parameter} &: \gamma \rightarrow \frac{\gamma(\gamma^2 + c^2 + 1)}{2(\gamma^2 + c^2)} \quad (\equiv \gamma'). \end{aligned}$$

We utilize the Properties 1-3 to get Eq: (14). When the variables $x(t)$ which follow a Cauchy distribution $C(c(t), \gamma(t))$ in the dynamical system Eq: (12), the variables $x(t+1)$ which also follow a Cauchy distribution as Figure 8. Hence, we get the self-consistent recurrence equations Eq. (14) about a median c and a scale parameter γ per iteration. This idea is also utilized to get $P_y(y|\xi_0)$ in Section III-C.

[1] V.I. Arnold and A. Avez, "Ergodic problems of classical mechanics", Benjamin, (1968).
[2] A. Boyarsky, and P. Góra, "Laws of chaos", Birkhäuser, (1997).

[3] J.-P. Eckmann and D. Ruelle, "Ergodic theory of chaos and strange attractors", Rev. Mod. Phys. **57**, 617–656, (1985).

$$\underbrace{x(t+1)}_{C(c(t+1), \gamma(t+1))(\cdot, \cdot \text{ Property 3})} = \underbrace{\underbrace{x(t)}_{C(c(t), \gamma(t))} \downarrow}_{C(c'(t), \gamma'(t))(\cdot, \cdot \text{ Property 2})} \underbrace{f(x(t))}_{C(c'(t), \gamma'(t))} + \underbrace{\underbrace{\zeta(t)}_{C(0,1)} \downarrow}_{C(0, |\varepsilon|)} \underbrace{\varepsilon \zeta(t)}_{C(0, |\varepsilon|)}$$

FIG. 8. Image of propagating Cauchy distributions from $x(t)$ and $\zeta(t)$ to $x(t+1)$