# arXiv:1701.02599v1 [q-bio.NC] 10 Jan 2017

# Issues in data expansion in understanding criticality in biological systems

Vaibhav Wasnik<sup>1</sup>

<sup>1</sup>Department of Physics, Saarland University, Saarbrucken, Germany (Dated: May 30, 2022)

(Dated: May 50, 2022

At the point of a second order phase transition also termed as a critical point, systems display long range order and their macroscopic behaviors are independent of the microscopic details making up the system. Due to these properties, it has long been speculated that biological systems that show similar behavior despite having very different microscopics, may be operating near a critical point. Recent methods in neuroscience are making it possible to explore whether criticality exists in neural networks. Despite being large in size, many data sets are still only a minute sample of the neural system and methods towards expanding these data sets have to be considered in order to study the existence of criticality. In this work we develop an analytical method of expanding a dataset to the large N limit so that statements about the critical nature of the data set could be made. We also show using a particular dataset analyzed computationally in literature that expanding data sets keeping the moments of the original data set need not lead to unique values of the critical temperature when the large N limit is considered analytically, despite the mirage of them appearing to do so when analyzed computationally. This suggests that not all available data sets from experiments are amenable for understanding the critically of the underlying system.

PACS numbers:

### INTRODUCTION

Many biological systems display self organization, arising from specific local interactions between the various constituents. The fact that global behaviors emerge because of local interactions has led to question whether biological systems are poised at criticality [1]. Recent work has highlighted such a possibility in systems ranging from gene expression [2], evolutionary ecology [3] to neural networks [4].

Neural networks are biological systems with inherently many body interactions. Criticality of retinal neurons have been studied by [5], [6]. In [6] the spiking of retinal neurons was recorded when it was subject to external stimulus. The spikes were binned in appropriate time intervals, leading to patterns made up of binary bits, akin to the up and down state of spin, with neuron firing corresponding to  $s_i = 1$  and non-firing to  $s_i = -1$ . The frequencies of occurrences of these patterns were then fit with an Ising like model. With the model fit, criticality was then explored in the retinal system by calculating its specific heat. Critical points are identifiable at divergences in the specific heat and for their finite sized data, they identified a possible divergence at the operating temperature of the network, namely  $k_BT = 1$ .

Since, criticality is only observed in statistical mechanical systems in the large N limit, there was a need to expand the data sets. In [5] the construction of larger data sets from smaller one's involved sampling from a distribution of the average spiking  $\langle s_i \rangle$  and the correlation between the spikes  $\langle s_i s_j \rangle$ . This then guaranted the larger data set has the same distribution as the smaller data set. Since this expansion of data was done computationally the expanded data sets even though large did not approach the large N limit. However, it was noticed that as the size of the constructed data set increased, the specific heat peaked closer and closer to  $k_B T = 1$ . This suggested a possible divergence in the specific heat at  $k_B T = 1$  as the system size increased, re-enforcing the possibility that the retinal network was operating at or near the critical point.

In this work we develop a methodology to expand the data sets analytically and show that the evaluated critical temperature could be very much dependent on how the data set is expanded, despite the mirage of them having a unique critical temperature when expanded computationally. We analyze the salamander retinal neurons data set used by [5], to show that the values of the critical temperature are very much dependent on how the dataset is expanded, implying that not all data sets available from experiments are amenable for expansion in order to study criticality.

Recently [7] have proposed a formalism to understand criticality by taking in to account the temporal dynamics of biological systems. [8] have talked about the Zipfs law distributions and hence criticality arising naturally when one of the fluctuating variables in the system is hidden. [9] constructed models that are consistent with distribution of global network activity. [10], [11] have proposed new ideas in modeling efforts to understand criticality in vertebrate retina. The work by [12] tried to model higher order correlations within cortical microcolumns. [13] have suggested an alternative way of understanding criticality by linking criticality to the exact inference of the probability distributions describing the data set. However, available experimental data sets are made up of a finite number of observables

and hence methods to extend available data sets to infinite number of observables are needed so that their critical behavior can be assessed. The work done in this paper addresses this crucial task of analytically extending the data set to the large N limit.

## THE CONSTRUCTION

[5] tried to understand whether biological systems are at a critical point by looking at the data coming from a smaller subsystem of a large system. The procedure they followed was to fit the average values  $\langle s_i \rangle$  and  $\langle c_{ij} \rangle$  for the smaller subsystem, by using a Boltzmann distribution with a Hamiltonian  $H = h_i s_i + J_{ij} s_i s_j$ . Critical properties could only be ascertained by a knowledge of the larger system of which the system is a subsystem of. Assume that the Hamiltonian of the larger system is labeled as  $H_{large}$ . Since the larger system should have the same  $\langle s_i \rangle$  and  $\langle c_{ij} \rangle$  distribution as the smaller subsystem, one could evaluate  $H_{large}$  by choosing the neurons of the larger system, such that one gets the same distribution for  $\langle s_i \rangle$  and  $\langle c_{ij} \rangle$  for the larger system as the smaller subsystem.[5] constructed the larger system in this way and found that the position of the specific heat peaks closer to  $k_BT = 1$  with increasing system size. Because their expansion of data set was done computationally, they could only conjecture the larger N limit.

The aim of our paper is to show that the evaluated value of the critical temperature is very much dependent on how the original data set is expanded. In this paper we consider a particular way of expanding the data set which allows for an exactly solvable large N limit. In order to get the large N limit to be exactly solvable we resort to replicating each neuron  $N \to \infty$  times as is elaborated below. We could have replicated each neuron in the original data set N times and then taken  $N \to \infty$  as a possible way of expanding the data set. This would have kept the same distribution of the  $\langle s^i \rangle$  and  $\langle c_{ij} \rangle$  as the original system. But this would only give us just one realization of expanding the original data set using the method of replication. In order to get a large number of realizations of the expanded data with the same distributions of the  $\langle s^i \rangle$  and  $\langle c_{ij} \rangle$  as the original data set, we instead first expand the original data set to an ensemble of 'middle subsystems' which have the same distributions of the  $\langle s^i \rangle$  and  $\langle c_{ij} \rangle$  as the original data set and then replicate each neuron in each of the 'middle subsystems' N times, so that we get an exact solution of the large N limit. This would then produce an ensemble of large N data sets, which have the same distribution of  $\langle s^i \rangle$  and  $\langle c_{ij} \rangle$  as the original data set, so that a statistical study of the critical temperatures could be carried out.

With this in mind, let us first construct a 'middle subsystem' by choosing  $\langle s^i \rangle$  and  $\langle c_{ij} \rangle$  from the distribution of smaller subsystem. This would then imply that the 'middle subsystem' has the same distribution as the smaller subsystem. Let us then construct the larger subsystem as talked below. We construct  $H_{large}$  by replicating each neuron from the 'middle subsystem' N times. Now, consider the Hamiltonian

$$H_{large} = \sum_{i=1,M} h_i S_i + \sum_{ij=1,M} \frac{J_{ij}}{2N} S_i S_j$$
(1)

Here  $S_i = s_1^i + s_2^i \dots s_N^i$  is the sum of all  $s_N^i$  which are the replicas neuron  $s_i$  in the 'middle subsystem'. M is the number of neurons in the middle subsystem. Because of the form of the above Hamiltonian we have the relations

$$\langle S_i \rangle = N \langle s_i \rangle = N \langle s_1^i \rangle = N \langle s_2^i \rangle \dots$$
<sup>(2)</sup>

This implies that we have N copies of  $\langle s_i \rangle$ , the first moments of the 'middle subsystem' in the larger system, as well as

$$\langle S_i S_j \rangle = N^2 \langle s^i s^j \rangle = N^2 \langle s^i_m s^j_n \rangle \tag{3}$$

for  $i \neq j$ . Next,

$$H_{large} = \sum_{i=1,M} h_i S_i + \sum_{ij=1,M} \frac{J_{ij}}{2N} S_i S_j$$
(4)

can be written as

$$H_{large} = N[\sum_{i=1,M} h_i m_i + \sum_{ij=1,M} J_{ij} m_i m_j]$$
(5)

where

$$m_i = \frac{s_i^1 + s_i^2 \dots + s_i^N}{N}$$
(6)

Hence the partition function can be written down as

$$Z = \sum_{s_i^j i \in [1,M], j \in [1,N]} e^{-\beta H_{large}} = \sum_{m_i} f(m_i) e^{-N\beta [\sum_{i=1,M} h_i m_i + \sum_{ij=1,M} J_{ij} m_i m_j]}$$
(7)

where we are summing over all possible values taken by  $m_i$ .  $f(m_i)$  is the number of ways of getting the value  $m_i$  by all possible combinations of  $s_i^j$  for  $j \in [1, N]$ . This is well known and the answer goes as

$$f(m_i) = \frac{\frac{N}{2}!}{\left(\frac{N}{2}(1+m_i)\right)!\left(\frac{N}{2}(1-m_i)!\right)}$$
(8)

which for  $N \to \infty$  becomes

$$f(m_i) = e^{Nm_i \ln m_i + N(1-m_i) \ln(1-m_i)}$$
(9)

and hence the partition function becomes

$$Z = \sum_{m_i} f(m_i) \sum_{m_i} e^{N[-\beta[\sum_{i=1,M} h_i m_i + \sum_{ij=1,M} J_{ij} m_i m_j] + \frac{1}{2} \sum_{i=1,M} (1+m_i) \ln(1+m_i) + \sum_{i=1,M} (1-m_i) \ln(1-m_i)]}$$
(10)

In the large N approximation the partition function is dominated by the saddle point and hence the solution is

$$\frac{\partial}{\partial m_i} \left[-\beta \left[\sum_{i=1,M} h_i m_i + \sum_{ij=1,M} J_{ij} m_i m_j\right] + \frac{1}{2} \sum_{i=1,M} (1+m_i) \ln(1+m_i) + \sum_{i=1,M} (1-m_i) \ln(1-m_i)\right] = 0 \quad (11)$$

which then gives us

$$m_i = \tanh\beta(\sum_{j=1,M} J_{ij}m_j + h_i) \tag{12}$$

The aim in constructing the larger data set is to make sure that the distribution of the correlations between the neurons  $c_{ij} = \langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle$  in the 'middle subsystem' is replicated into the larger data set. However since one cannot have correlations between a neuron and itself we should have that  $i \neq j$ .

In our larger data set the correlations between neurons have the form  $c_{ij}^{mn} = \langle s_m^i s_n^j \rangle - \langle s_m^i \rangle \langle s_n^j \rangle$ 

The way the Hamiltonian is written would imply that  $N^2$  of the expectation values given by  $\langle s_m^i s_n^j \rangle$  for particular values of  $i, j \in [1, M]$  and the  $N^2$  possible combinations of  $m, n \in [1, N]$  are equal to each other. Or,  $\langle S_i S_j \rangle = N^2 \langle s_m^i s_n^j \rangle$ .

Also, all the  $\langle s_m^i \rangle$  for a particular value of  $i \in [1, M]$  and all values of  $m \in [1, N]$  are equal to each other. Or  $\langle S_i \rangle = N \langle s_m^i \rangle$ .

This would immediately imply that  $c_{ij}^{mn} = \frac{\langle S_i S_j \rangle - \langle S_i \rangle \langle S_j \rangle}{N^2}$  for particular values of with  $i, j \in [1, M]$  with  $i \neq j$ and the  $N^2$  possible combinations of  $m, n \in [1, N]$  are equal to each other. And hence we would have a replication of the  $c_{ij}$  from the 'middle subsystem' in to the larger data set, if we set  $c_{ij}$  from the 'middle subsystem' into  $c_{ij}^{mn} = \frac{\langle S_i S_j \rangle - \langle S_i \rangle \langle S_j \rangle}{N^2}$  in the expanded data set.

However, the larger data set also comes up with additional correlations between neurons given by  $c_{ii}^{mn} = \langle s_{ii}^{i} s_{in}^{i} \rangle - \langle s_{im}^{i} \rangle \langle s_{in}^{i} \rangle = \frac{\langle S_{ii} S_{in} \rangle - \langle S_{ii} \rangle \langle S_{in} \rangle}{N^2}$ , for  $i \in [1, M]$  and  $m, n \in [1, N]$ . Again by construction for each value of i, all the  $c_{ii}^{mn}$ 's are equal to each other. Hence, if we want to keep the distribution of the  $c_{ij}$  from the 'middle subsystem' into the larger data set we should pick the  $c_{ii}^{mn} = \frac{\langle S_{ii} S_{i} \rangle - \langle S_{ii} \rangle \langle S_{ii} \rangle}{N^2}$  from the  $c_{ij}$  distribution of the 'middle subsystem'. Now the mean field solution can be written down as

$$\tanh^{-1} m_i = \beta(\sum_j J_{ij}m_j + h_i) \tag{13}$$

The above equation implies that

$$J_{il} = -[C^{-1}]_{il} \tag{14}$$

if  $i \neq l$  and

$$J_{ii} = \frac{1}{1 - m_i^2} - [C^{-1}]_{ii} \tag{15}$$

Keeping in line with the logic elaborated above we hence choose  $C_{ij} = c_{ij}$  from the 'middle subsystem' for  $i \neq j$  and  $C_{ii}$ 's are chosen from the distribution of  $c_{ij}$  with  $i \neq j$  from the 'middle subsystem'.

As shown in the appendix the critical temperature is then given by the largest eigenvalue of the matrix  $J_{ij}$ .

### ANALYSIS OF SALAMANDER RETINAL DATA

We analyzed the the neuronal firing data from a salamander retina by using the methodology explained above. This data was earlier analyzed in [5], [6]. The data from 39 neurons was binned in 20ms bins. [5] suggested that  $k_BT = 1$  was the critical point for the system when extrapolated to the large N limit. In fig. we plot the results of our evaluations using the methodology outlined above. The x-axis plots the size M of the middle subsystem, while the y-axis plots the logarithm of the critical temperature  $k_BT$ . As we can see the value of the critical temperature is nowhere close to  $k_BT = 1$ , but the mean value of  $k_BT$  varies as a function of the value of M from  $k_BT = 10^{2.5}$  for M = 10, to  $k_BT = 10^{3.5}$  for M = 1000. This shows that the value of the critical temperature is very much dependent on how the data set is expanded. Since the evaluations involved estimating the largest eigenvalue of the matrix  $J_{ij}$ , we could only computationally work with values of M which are bounded. However, the fact that the  $m_i$ 's and  $c_{ij}$ 's are chosen from a distribution implies that the evaluation of the eigenvalues of  $J_{ij}$  for  $M \to \infty$  turns in to a problem in random matrix theory. This hence becomes a problem with potential for future research.

### APPENDIX

The mean field solution Eq. (12) can be derived by the minimization of the free energy functional

$$F = -\beta \sum_{i} h_{i}m_{i} - \beta \sum_{ij} \frac{J_{ij}m_{i}m_{j}}{2} + \sum_{i} \frac{1}{2}(1-m_{i})ln(1-m_{i}) + \sum_{i} \frac{1}{2}(1+m_{i})ln(1+m_{i}).$$
(16)

Since  $m_i$  is smaller than one, we can expand the logarithms in a power series, that gives

$$F = -\beta \sum_{i} h_{i}m_{i} - \beta \sum_{ij} \frac{J_{ij}m_{i}m_{j}}{2} + \sum_{ij} \frac{m_{i}m_{j}I_{ij}}{2} - \sum_{i} \frac{m_{i}^{4}}{12}$$
(17)

where  $I_{ij}$  is the identity matrix. We can then construct a basis in which  $Q_{ij} = \beta J_{ij} - I_{ij}$  is diagonal. In this basis we have  $m_i = \sum_j \Lambda_{ij} M_j$ . Here  $\Lambda$  is a matrix that diagonalizes Q, i.e  $\Lambda^{-1}Q\Lambda$  is diagonal. Substituting for  $m_i$  the above equation becomes

$$F = -\beta \sum_{i} h_{i} \sum_{j} \Lambda_{ij} M_{j} - \beta \sum_{i} \frac{J_{i} M_{i}^{2}}{2} - \sum_{i} \frac{[\sum_{j} \Lambda_{ij} M_{j}]^{4}}{12}$$
(18)

where  $J_i$  are the eigenvalues of  $\beta J_{ij} - I_{ij}$ . Now, if we make the  $M_i$ 's heterogeneous we have to include a energy cost that can be written as  $(\nabla M_i)^2$ , leading to the minimum that is a homogeneous solution. Talking about correlation lengths in field theory requires the presence of a gradient term and hence the motivation behind its introduction here. The free energy becomes

$$F = \frac{1}{V} \int_{V} d^{d}x \left[\sum_{i} c(\nabla M_{i})^{2} - \beta \sum_{i} \frac{J_{i}M_{i}^{2}}{2} - \sum_{i} \frac{\left[\sum_{j} \Lambda_{ij}M_{j}\right]^{4}}{12} - \beta \sum_{i} h_{i} \sum_{j} \Lambda_{ij}M_{j}\right]$$

$$(19)$$

where V is the volume  $\int_V d^d x$ , c is a constant with dimension of  $[L^2]$  and c is related to the energy scale  $\lambda$  of the field theory above as  $c \sim \frac{1}{\lambda^2}$ . We can redefine our fields  $M \to \sqrt{cM}$  so that M has a dimension of [L]. The quadratic terms in the above expression would then look like

$$\frac{1}{V} \int_{V} d^{d}x \left[\sum_{i} (\nabla M_{i})^{2} - \beta \sum_{i} \frac{J_{i}M_{i}^{2}}{2c}\right]$$

$$\tag{20}$$

implying that  $\frac{2c}{\beta J_i}$  is the correlation length for the field  $M_i$ . The values of  $k_B T$  for which a particular  $J_i$  becomes equal to zero hence corresponds to the limit in which the correlation length for that  $M_i$ 's diverges. This is also the critical temperature for parameter  $M_i$  implying that it changes continuously from zero to a finite value around this temperature when the h's are zero. The largest value of  $k_B T$  for which the correlation length of any of the  $M_i$  diverges is then the critical temperature of the system because this is the temperature below which any linear combination of the  $M_i$ 's starts becoming non-zero. This value of  $k_B T$  is the largest eigenvalue of  $J_{ij}$ .

### CONCLUSION

What we have seen in this work through an analysis of the system of retinal neurons in [1] is that the evaluation of the critical temperature of a system by expanding a system such that the distribution of the moments of the original system are preserved does not guarantee a unique value of the critical temperature, but the evaluated value of the critical temperature is very much dependent on the details in which the data is expanded. In order to understand the possible reasons behind this, let us note that first of all the expanding the dataset keeping the moments of the data fixed is only an expression about the intution that a system could be essentially a sum of its parts. Hence systems which are homogenous could be studied by analyzing a part of the system. This was the logic given behind analysis of the salamander retina in [1]. This has nothing to do with the criticality of the systems. If the properties of the system that are evaluated are dependent on how the system is expanded it could imply many things. One possible implication is that the subsystem from which the data set is taken is too small and hence repeating this subsystem does not lead to the production of the entire system. Another possible implication is that the larger system is not plainly a repetition of smaller subunits. These have nothing to do with the problem of expanding the data set, but only are issues related to the constraints imposed by the system one is analyzing. A system whose subsystem from which data is taken is essentially a replica of copies of this subsystem would then give a critical temperature which is unchanged by how the data set is expanded. However even then the system may not have its critical temperature  $k_BT = 1$ . It is only systems that are poised at criticality which would have their critical temperature at  $k_BT = 1$ . It is in order to answer this question whether such systems are poised at criticality that our method of expanding the dataset obtained from a subsystem to its large N limit will be valuable.



FIG. 1: The x-axis plots the size M of the middle subsystem, while the y-axis plots the logarithm of the critical temperature  $k_BT$ . As we can see the value of the critical temperature is nowhere close to  $k_BT = 1$ , but varies as a function of the value of M from  $k_BT = 10^{2.5}$  for M = 10, to  $k_BT = 10^{3.5}$  for M = 1000.

# ACKNOWLEDGEMENT

Vaibhav Wasnik would like to thank Dr. Lukas Janssen for discussions on criticality in statistical systems.

### REFERENCES

- [1] T. Mora and W. Bialek, J. Stat. Phys, 144, 268, (2011).
- [2] M. Nykter, N. D. Price, M. Aldana, S. A. Ramsey, S. A. Kauffman, L. E. Hood, O. Yli-Harja, and I. Shmulevich, PNAS 105, 1897 (2008).
- [3] R.V. Sol, S. C. Manrubia, M. Benton, S. Kauffman and P. Bak, Trends Ecol. Evol. 14, 156 (1999).
- [4] M.G. Kitzbichler, M.L. Smith, S.R. Christensen and E. Bullmore, PLoS Comput. Biol. 5, 1000314 (2009).
- [5] G.Tkacik, E. Schneidman, M.J.Berry II and W. Bialek, arXiv preprint q-bio/0611072, (2006).
- [6] E. Schneidman, M. Berry II, R. Segev, W. Bialek, Nature. 440, 1007 (2006).
- [7] T. Mora, S. Deny, O. Marre, Physical review Letters. 114, 07815 (2015).
- [8] D. Schwab, I. Nemenman, P. Mehta, Physical review Letters. 113, 068102 (2014)
- [9] G. Tkačik, O. Marre, T. Mora, D. Amodel, M. Berry II, W. Bialek, Journal of Statistical Mechanics: Theory and Experiment. P03011 (2013).
- [10] G. Tkačik, O. Marre, T. Mora, D. Amodel, M. Berry II, W. Bialek, arXiv:1407.5946 [q-bio.NC].
- [11] G. Tkačik, O. Marre, E. Schneidman, D. Amodel, M. Berry II, W. Bialek, Plos Computational Biology. DOI: 10.1371/journal.pcbi.1003408 (2014)
- [12] U. K ö ster, J. Sohl-Dickstein, C.M. Gray, B.A. Olshausen, Plos Computational Biology. DOI:10.1371/journal.pcbi.1003684 (2014)
- [13] I. Mastromatteo and M. Marsili, Journal of Statistical Mechanics: Theory and Experiment.10, P10012 (2011).
- [14] D. Ackley, G. Hinton and T. Sejnowski, Cogn. Sci. 9, 147 (1985).
- [15] B. Caracheo, E. Emberly, S. Hadizadeh, J.M. Hyman and J. K. Seamans, Front Neurosci. 7, 74 (2013).
- [16] T. Tanaka, Phys. Rev. E 58, 2302 (1998).
- [17] H. Nishimori, G. Ortiz *Elements of Phase Transitions and Critical Phenomena* (Oxford Graduate Texts January 22, 2011).