

Entropy facilitated active transport

J. M. Rubí^{†, ‡}, A. Lervik^{*‡}, D. Bedeaux[‡], and S. Kjelstrup[‡]

[†]Statistical and Interdisciplinary Physics Section, Department de Física de la Matèria Condensada, Universitat de Barcelona, Martí i Franquès 1, 08028, Barcelona, Spain

[‡]Department of Chemistry, Faculty of Natural Sciences and Technology, Norwegian University of Science and Technology, Trondheim, Norway

Abstract

We show how active transport of ions can be interpreted as an entropy facilitated process. In this interpretation, the pore geometry through which substrates are transported can give rise to a driving force. This gives a direct link between the geometry and the changes in Gibbs energy required. Quantifying the size of this effect for several proteins we find that the entropic contribution from the pore geometry is significant and we discuss how the effect can be used to interpret variations in the affinity at the binding site.

Active transport is of major importance in biology; meaning transport of a compound against its chemical potential, driven by a chemical reaction. A notable example is the large P-type ATPase protein family which functions as ion or lipid pumps, crucial for a wide range of processes in almost all forms of life [1]. The P-type ATPases share a common topology and their operation can be described using a Post-Albers cycle with four key conformations: E1, E1P, EP2 and E2 [2]. The transitions $E1P \leftrightarrow E2P$ and $E2 \leftrightarrow E1$ in the catalytic cycle are associated with large conformational changes [2] and it is clear

*anders.lervik@ntnu.no

that the energy released by ATP hydrolysis drives the enzyme between these states.

One may ask about the meaning of the conformational changes. In particular, the formation of a wide funnel-shaped outlet channel can be observed in several P-type ATPases [3–6] when the enzyme changes from the E1P state to the E2P state. The occluded ion is then eventually exposed to the lumen; but why does the ion leave its binding site when the chemical potential is so much larger in the direction of transport than in the opposite direction? Can this be influenced by the observed pore geometries? How can the change in Gibbs energy by the chemical reaction, which may take place relatively far away from the binding site (see e.g. Ref. [7]), be transferred and used at the ion binding site? These questions, which have been of major interest since the discovery of the pumps, are still not fully answered and this is the topic we discuss here.

We shall argue that active transport can be better understood by shifting focus from an energy to an entropy barrier in a specific part of the translocation process. When the outlet channel is formed in the E2P state, the shape of this channel directly facilitates the transport of ions by allowing for an entropy increase, or equivalently, a chemical potential decrease in the E2P state as we will demonstrate. The idea that the actual pore geometry may play a role in active transport, stems from previous studies which have shown how entropic barriers can play a major role for separation purposes [8–10].

To exemplify the impact of the pore geometry, we will take the Ca^{2+} transporting Ca^{2+} -ATPase of sarcoplasmic reticulum (SR) but we note that the ideas presented may apply more generally. For the Ca^{2+} -ATPase, we illustrate the variation of the chemical potential as ions are transported from the cytosol to the lumen of the SR in Fig. 1. It is known that the binding of Ca^{2+} is fast [11] and at equilibrium, the chemical potential of Ca^{2+} in the E1 state, $\mu_{\text{Ca2.E1}}$, is equal to the chemical potential in the cytosol, μ_{out} . During operation of the pump, there is probably a small difference between μ_{out} and $\mu_{\text{Ca2.E1}}$ but the chemical potential in the final state, μ_{in} , is clearly larger than μ_{out} as depicted in Fig. 1. There is a large uncertainty related to the chemical potential of Ca^{2+} when the enzyme is in the state E2P, $\mu_{\text{Ca2.E2P}}$. In Fig 1 it is assumed to be close to μ_{in} (which is the lower bound resulting in a positive Ca^{2+} flux), enabling the ion to pass to the lumen. The variation in binding energy inherent in this picture has been attributed to the enthalpic part of the chemical potential, as the enthalpy gives a measure of the bond strength: Repulsive forces can raise the chemical potential $\mu_{\text{Ca2.E2P}}$, which




Figure 1: (Color on-line.) Illustration of the variation in the chemical potential μ of a Ca^{2+} -ion as it is transported by the Ca^{2+} -ATPase from the cytosol to the lumen of SR. The coordinate x denotes the extent of the transport inside funnel-shaped pore from the binding site at $x = 0$ to the lumen at $x = 1$. The chemical potential is indicated for four states; in the cytosol/outside (μ_{out}), lumen/inside (μ_{in}), the E1 ($\mu_{\text{Ca2.E1}}$) and the E2P ($\mu_{\text{Ca2.E2P}}$) states. For a concentration ratio of 10^4 at 37°C , the energy requirement is 24 kJ per mol ion transported.

results in a low affinity binding site. In such a situation the ions can move spontaneously to the lumen. Alternatively, this change in chemical potential can be attributed to entropic effects. The leap from $\mu_{\text{Ca2.E1}}$ to $\mu_{\text{Ca2.E2P}}$ is due to the hydrolysis of ATP and it can arise from increasing the enthalpy of the ion, or by lowering the entropy of the ion. The latter can be brought about by a change in the actual shape of the channel where the ion is transported. A particle moving in a pore with a varying cross-sectional area will be subjected to entropic forces due to the change in area and this can in fact *facilitate* the transport.

The outlet channels observed in the P-type ATPases are narrow but widens towards the exit of the channel, as illustrated in Fig 2. We assume that the ion transport is effectively one-dimensional, directed along a spatial coordinate x . The ion density along the transport direction may be approximated by [8]

$$\rho(x) = \rho_0 \frac{A(x)}{A_0} \quad (1)$$

where ρ_0 and A_0 are the ion density and cross-sectional area at a reference

point, say $x = 0$. The size of the particles has been shown to influence the transport properties [12] and here, both $A(x)$ and A_0 measures the available area, e.g. for a circular geometry: $A(x) = \pi (r(x) - a)^2$ where $r(x)$ is the radius of the pore at position x and a the radius of the ion. The density given by Eq. 1 can also be viewed as the canonical distribution function

$$\rho(x) = \rho_0 \exp\left(\frac{\Delta S_{\text{pore}}(x)}{R}\right) \quad (2)$$

where we have introduced the change in entropy, $\Delta S_{\text{pore}}(x)$, associated with the change in cross-sectional area. From these equations, we can conclude that change in the entropy during ion motion along the pore is directly related to the cross sectional area

$$\Delta S_{\text{pore}} = R \ln \frac{A(x)}{A_0} \quad (3)$$

and that the position dependent entropy along the ion trajectory is giving rise to a (thermodynamic) force of entropic nature, F_{ent} , acting on the ions

$$F_{\text{ent}} = T \frac{\partial \Delta S_{\text{pore}}}{\partial x} = RT \frac{1}{A(x)} \frac{\partial A(x)}{\partial x} \quad (4)$$

The direction of the force directly depends on the slope of the channel. For the conical structure shown in Fig. 2 the entropic force is positive, and will facilitate translocation in direction of the wider opening. The overall entropy change (ΔS) for the ion translocation will thus have two contributions: the normal contribution from the ion concentration difference ($\Delta S_{\text{Ca}} = -R \ln(c_{\text{in}}/c_{\text{out}}) = -\Delta\mu_{\text{Ca}}/T$), and a special contribution from the pore shape. Per mol of ion we have $\Delta S = \Delta S_{\text{Ca}} + \Delta S_{\text{pore}}$ and by introducing Eq. 3 for $x = 1$, we obtain

$$\Delta S = R \ln \frac{c_{\text{out}} A_{\text{in}}}{c_{\text{in}} A_0} \quad (5)$$

The entropic force depends on the concentration ratio, but also on the ratio of the cross-sectional areas at the two sides of the channel. A large $c_{\text{in}}/c_{\text{out}}$ ratio may be counteracted by a smaller A_0/A_{in} ratio, meaning that translocation may take place against the gradient in concentration. In other words, it may be facilitated by entropic forces induced by the formation of a conical pore and the larger the ratio A_{in}/A_0 , the larger is the co-acting entropic force.




Figure 2: (Color on-line.) The E2P state for the Ca²⁺-ATPase (Protein Data Bank ID: 3B9B [3]) showing the funnel shaped channel. The inset shows a conical funnel where the area increases from the start of the channel (A_0) to the exit (A_{in}) where $0 \leq x \leq 1$ indicates the position along the channel.

The translocation rate, J_{Ca} , can be shown to be

$$J_{Ca} = -L \left[1 - \left(\frac{c_{out} A_{in}}{c_{in} A_0} \right) \exp \left(-\frac{\Delta H}{RT} \right) \right] \quad (6)$$

where L is the backward reaction rate, and ΔH is the change in enthalpy. The equation holds when the activity coefficients inside and outside the membrane are the same. In the outset ΔH is unknown. In the case that the pump operates far from equilibrium the last term in the parenthesis in Eq. 6 dominates, giving

$$J_{Ca} = L \left(\frac{c_{out} A_{in}}{c_{in} A_0} \right) \exp \left(-\frac{\Delta H}{RT} \right) \quad (7)$$

The equation gives an Arrhenius behavior of the current through the dependence on ΔH . Assuming that the entropy change alone drives the translocation, we have

$$J_{Ca} = L \left(\frac{c_{out} A_{in}}{c_{in} A_0} \right) \quad (8)$$

where the flux is given by the ratio of concentrations and cross-sectional areas.

Assuming a circular cross-sectional area, the entropic driving force is,

$$-T\Delta S = RT \ln \left[\left(\frac{c_{\text{in}}}{c_{\text{out}}} \right) \left(\frac{r_0 - a}{r_{\text{in}} - a} \right)^2 \right] \quad (9)$$

where r_0 is the radius at the start of the channel, r_{in} at the exit of the channel and a the radius of the ion. In Fig. 3 we show the entropic contribution to the driving force for several enzymes. As the results show, the entropic contribution can be sizable. It is not sufficient to explain active transport by itself for all the enzymes as can be seen by the results for the Ca^{2+} -ATPase. In this case, the entropic contribution is on the order of 10 kJ/mol ion transported at 37°C. This accounts for 40% of the chemical potential difference which indicate that other contributions, for instance enthalpic effects, may also be important for this protein. For the Na^+/K^+ -ATPase we find that, due to the narrowness of the pore, the entropic contribution (20 kJ/mol) is relatively large compared to the other enzymes. This is sufficient to overcome the concentration ratio of Na^+ at physiological conditions, however, in this case, the Na^+ ions are transported against the resting membrane potential. Assuming a resting potential of -60 mV and a concentration ratio on the order of 10, the required Gibbs energy is 12 kJ/mol ion transported. This means that the entropic force estimated in Fig. 3 is still sufficient for this enzyme. The entropic contribution for the two other enzymes is similar to the Ca^{2+} -ATPase. We have neglected the possibility of complexing the ions with other species such as water which could increase the effective radius and the entropic force.

We have shown that the actual pore geometry observed in several transport enzymes may give rise to an entropic force, facilitating transport. This entropic effect can be used, together with enthalpic effects, to explain how conformational changes influences the binding site, allowing transport of compounds against their concentration gradients. In particular, the entropic force gives a direct link between the energy released by a reaction at the active site and changes in the chemical potential at the binding site: By modifying the conformation and the pore geometry, a low affinity binding site can be created.




Figure 3: (Color on-line.) Entropic contribution, ΔS_{pore} , (solid black line) to the driving force as a function of the ratio of cross sectional areas. Contributions for several enzymes are also shown: Zn^{2+} -ATPase [2] P-glycoprotein [13], Ca^{2+} -ATPase [3] and Na^+/K^+ -ATPase [4]. For the Zn^{2+} -ATPase, the widths of the channel were estimated using the crystal structure (Protein Data Bank ID: 4UMV). We have considered the size of the ions using the ionic radii reported by Marcus [14] and for the Na^+/K^+ -ATPase we only considered the smallest ion (Na^+). As P-glycoprotein may transport many different compounds, we have omitted the size of the transported compound and the estimate represents a lower bound.

Author contributions

JMR, AL, DB, and SK developed the theory, performed the analysis, and wrote the manuscript. JMR initiated the work.

Acknowledgements

The Norwegian University of Science and Technology is thanked for supporting the stay of JMR.

References

- [1] Palmgren M. G., and P. Nissen. 2011. P-type ATPases. *Annu. Rev. Biophys.*, 40:243–266.
- [2] Sitsel O., C. Grønberg, H. E. Autzen, K. Wang, G. Meloni, P. Nissen, and P. Gourdon. 2015. Structure and function of Cu(I)- and Zn(II)-ATPases. *Biochemistry*, 54:5673–5683.
- [3] Olesen C., M. Picard, A.-M. L. Winther, C. Gyruup, J. P. Morth, C. Oxvig, J. V. Møller, and P. Nissen. 2007. The structural basis of calcium transport by the calcium pump. *Nature*, 450:1036–1042.
- [4] Takeuchi A., N. Reyes, P. Artigas, and D. C. Gadsby. 2009. Visualizing the mapped ion pathway through the Na,K-ATPase pump. *Channels*, 3:383–386.
- [5] Wang K., O. Sitsel, G. Meloni, H. E. Autzen, M. Andersson, T. Klymchuk, A. M. Nielsen, D. C. Rees, P. Nissen, and P. Gourdon. 2014. Structure and mechanism of Zn²⁺-transporting P-type ATPases. *Nature*, 514:518–522.
- [6] Abe K., K. Tani, T. Nishizawa, and Y. Fujiyoshi. 2009. Inter-subunit interaction of gastric H⁺,K⁺-ATPase prevents reverse reaction of the transport cycle. *EMBO J.*, 28:1637–1643.
- [7] Inesi G. 2011. Calcium and copper transport ATPases: analogies and diversities in transduction and signaling mechanisms. *J. Cell. Commun. Signal.*, 5:227–237.

- [8] Reguera D., and J. M. Rubí. 2001. Kinetic equations for diffusion in the presence of entropic barriers. *Phys. Rev. E*, 64:061106.
- [9] Rubí J. M., and D. Reguera. 2010. Thermodynamics and stochastic dynamics of transport in confined media. *Chem. Phys.*, 375:518–522.
- [10] Reguera D., A. Luque, P. S. Burada, G. Schmid, J. M. Rubí, and P. Hänggi. 2012. Entropic splitter for particle separation. *Phys. Rev. Lett.*, 108:020604.
- [11] Peinelt C., and H.-J. Apell. 2005. Kinetics of ca^{2+} binding to the SR Ca-ATPase in the E_1 state. *Biophys. J.*, 89:2427–2433.
- [12] Riefler W., G. Schmid, P. S. Burada, and P. Hänggi. 2010. Entropic transport of finite size particles. *J. Phys. Condens. Matter*, 22:454109.
- [13] Loo T. W., and D. M. Clarke. 2001. Determining the dimensions of the drug-binding domain of human P-glycoprotein using thiol cross-linking compounds as molecular rulers. *J. Biol. Chem.*, 276:36877–36880.
- [14] Marcus Y. 1988. Ionic radii in aqueous solutions. *Chem. Rev.*, 88:1475–1498.