

Braid groups in handlebodies and corresponding Hecke algebras

Valeriy G. Bardakov

Abstract In this paper we study the kernel of the homomorphism $B_{g,n} \rightarrow B_n$ of the braid group $B_{g,n}$ in the handlebody \mathcal{H}_g to the braid group B_n . We prove that this kernel is semi-direct product of free groups. Also, we introduce an algebra $H_{g,n}(q)$, which is some analog of the Hecke algebra $H_n(q)$, constructed by the braid group B_n .

1 Introduction

Let \mathcal{H}_g be a handlebody of genus g . The braid group $B_{g,n}$ on n stings in the handlebody \mathcal{H}_g was introduced by A. B. Sossinsky [12] and independently S. Lambropoulou [7]. Properties of this group are studied by V. V. Vershinin [13, 14] and by S. Lambropoulou [7, 8].

The motivation for studying braids from $B_{g,n}$ comes from studying oriented knots and links in knot complements in compact connected oriented 3-manifolds and in handlebodies, since these spaces may be represented by a fixed braid or a fixed integer-framed braid in \mathbb{S}^3 [9, 7, 4, 10]. Then knots and links in these spaces may be represented by elements of the braid group $B_{g,n}$ or an appropriate cosets of these group [7]. In particular, if M denotes the complements of the g -unlink or a connected sum of g lens spaces of type $L(p, 1)$ or a handlebody of genus g , then knots and links in these spaces may be represented precisely by the braids in $B_{g,n}$ for $n \in \mathbb{N}$. In the case $g = 1$, $B_{1,n}$ is the Artin group of type \mathcal{B} .

The group $B_{g,n}$ can be considered as a subgroup of the braid group B_{g+n} on $g+n$ strings such that the braids in $B_{g,n}$ leave the first g strings identically fixed. Using this fact V. V. Vershinin [13] and S. Lambropoulou [7] defined a epimorphism of $B_{g,n}$ onto the symmetric group S_n and prove that the kernel of this epimorphism: $P_{g,n}$ is a subgroup of the pure braid group P_{g+n} and is semi-direct products of free

Valeriy G. Bardakov
Sobolev Institute of Mathematics, Novosibirsk State University, Novosibirsk State Agrarian University, Novosibirsk 630090, Russia, e-mail: bardakov@math.nsc.ru

groups. On the other side, V. V. Vershinin [13] noted that there is a decomposition $B_{g,n} = R_{g,n} \rtimes B_n$ for some group $R_{g,n}$ and the braid group B_n . S. Lambropoulou proved in [8] that $R_{1,n}$ is isomorphic to a free group F_n of rank n .

The Hecke algebra of type A, $H_n(q)$ was used by V. F. R. Jones [3] for the construction of polynomial invariant for classical links, the well known HOMFLYPT polynomial. S. Lambropoulou [8] defined a generalization of the Hecke algebra of type B to construct a HOMFLYPT-type polynomial invariant for links in the solid torus, which was then used in [2] for extending the study to the lens spaces $L(p, 1)$.

In the present paper we define some decomposition of $P_{g,n}$ as semi-direct product of free groups, which is different from the decomposition defined in [7, 13]. Also, we study the group $R_{g,n}$ and prove that this group is semi-direct products of free groups. Using this decomposition and the decomposition $B_{g,n} = R_{g,n} \rtimes B_n$ we define some algebra, which is a generalization of the Hecke algebra $H_n(q)$.

The paper is organized as follows. In Section 2, we remind some facts on the braid group B_n . In particular, we define vertical and horizontal decompositions of the pure braid group P_n . In Section 3, we recall some facts on the group $B_{g,n}$, we describe the vertical decomposition of $P_{g,n}$ and we define the horizontal decomposition of $P_{g,n}$. In Section 4, we study $R_{g,n}$ and we construct vertical and horizontal decompositions for this group. In Section 5 we introduce some algebra $H_{g,n}(q)$ which is a generalization of the Hecke algebra $H_n(q)$ and we find the quotient of $B_{g,n}$ by the relations $\sigma_i^2 = 1$.

Acknowledgements.

The author gratefully acknowledges Prof. Lambropoulou and her students and colleagues: Neslihan Gügümcü, Stathis Antoniou, Dimos Goundaroulis, Ioannis Diamantis, Dimitrios Kodokostas for the kind invitation to the Athens, where this paper was written, for conversation and interesting discussions.

This work was supported by the Russian Foundation for Basic Research (project 16-01-00414).

2 Braid group

In this section we recall some facts on the braid groups (see [1, 11]).

The braid group B_m , $m \geq 2$, on m strings is generated by elements

$$\sigma_1, \sigma_2, \dots, \sigma_{m-1},$$

and is defined by relations

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i, & \text{for } |i - j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, & \text{for } i = 1, 2, \dots, m - 2. \end{aligned}$$

A subgroup of B_m which is generated by elements

$$a_{ij} = \sigma_{j-1}\sigma_{j-2}\dots\sigma_{i+1}\sigma_i^2\sigma_{i+1}^{-1}\dots\sigma_{j-2}^{-1}\sigma_{j-1}^{-1}, \quad 1 \leq i < j \leq m,$$

is called the *pure braid group* and is denoted P_m . This group is defined by the relations

$$a_{ik}a_{ij}a_{kj} = a_{kj}a_{ik}a_{ij}, \quad (1)$$

$$a_{nj}a_{kn}a_{kj} = a_{kj}a_{nj}a_{kn}, \quad \text{for } n < j, \quad (2)$$

$$(a_{kn}a_{kj}a_{kn}^{-1})a_{in} = a_{in}(a_{kn}a_{kj}a_{kn}^{-1}), \quad \text{for } i < k < n < j, \quad (3)$$

$$a_{kj}a_{in} = a_{in}a_{kj}, \quad \text{for } k < i < n < j \text{ or } n < k. \quad (4)$$

The subgroup P_m is normal in B_m , and the quotient B_m/P_m is the symmetric group S_m . The generators of B_m act on the generator $a_{ij} \in P_m$ by the rules:

$$\sigma_k^{-1}a_{ij}\sigma_k = a_{ij}, \quad \text{for } k \neq i-1, i, j-1, j, \quad (5)$$

$$\sigma_i^{-1}a_{i,i+1}\sigma_i = a_{i,i+1}, \quad (6)$$

$$\sigma_{i-1}^{-1}a_{ij}\sigma_{i-1} = a_{i-1,j}, \quad (7)$$

$$\sigma_i^{-1}a_{ij}\sigma_i = a_{i+1,j}[a_{i,i+1}^{-1}, a_{ij}^{-1}], \quad \text{for } j \neq i+1 \quad (8)$$

$$\sigma_{j-1}^{-1}a_{ij}\sigma_{j-1} = a_{i,j-1}, \quad (9)$$

$$\sigma_j^{-1}a_{ij}\sigma_j = a_{ij}a_{i,j+1}a_{ij}^{-1}, \quad (10)$$

where $[a, b] = a^{-1}b^{-1}ab$.

Denote by

$$U_i = \langle a_{1i}, a_{2i}, \dots, a_{i-1,i} \rangle, \quad i = 2, \dots, m,$$

a subgroup of P_m . It is known that U_i is a free group of rank $i-1$. One can rewrite the relations of P_m as the following conjugation rules (for $\varepsilon = \pm 1$):

$$a_{ik}^{-\varepsilon}a_{kj}a_{ik}^{\varepsilon} = (a_{ij}a_{kj})^{\varepsilon}a_{kj}(a_{ij}a_{kj})^{-\varepsilon}, \quad (11)$$

$$a_{kn}^{-\varepsilon}a_{kj}a_{kn}^{\varepsilon} = (a_{kj}a_{nj})^{\varepsilon}a_{kj}(a_{kj}a_{nj})^{-\varepsilon}, \quad \text{for } n < j, \quad (12)$$

$$a_{in}^{-\varepsilon}a_{kj}a_{in}^{\varepsilon} = [a_{ij}^{-\varepsilon}, a_{nj}^{-\varepsilon}]^{\varepsilon}a_{kj}[a_{ij}^{-\varepsilon}, a_{nj}^{-\varepsilon}]^{-\varepsilon}, \quad \text{for } i < k < n, \quad (13)$$

$$a_{in}^{-\varepsilon}a_{kj}a_{in}^{\varepsilon} = a_{kj}, \quad \text{for } k < i < n < j \text{ or } n < k. \quad (14)$$

From these rules it follows that U_m is normal in P_m and hence P_m has the following decomposition: $P_m = U_m \rtimes P_{m-1}$, where the action of P_{m-1} on U_m is defined by the rules (11-14). By induction on m , P_m is the semi-direct product of free groups:

$$P_m = U_m \rtimes (U_{m-1} \rtimes (\dots \rtimes (U_3 \rtimes U_2) \dots)).$$

We will call this decomposition *vertical decomposition*.

Let $U_m^{(k)}$, $k = 1, 2, \dots, m$, be the subgroup of P_m which is generated by a_{ij} , where $k < j$. Then $U_m^{(k)} = U_m \rtimes (U_{m-1} \rtimes (\dots \rtimes (U_{k+2} \rtimes U_{k+1}) \dots))$. By definition $U_m^{(1)} = P_m$ and these groups form the normal series

$$1 = U_m^{(m)} \leq U_m^{(m-1)} \leq \dots \leq U_m^{(2)} \leq U_m^{(1)} = P_m,$$

where

$$U_m^{(r)} / U_m^{(r+1)} \cong F_r, \quad r = 1, 2, \dots, m-1.$$

On the other side, define the following subgroups of P_m :

$$V_k = \langle a_{k,k+1}, a_{k,k+2}, \dots, a_{k,m} \rangle, \quad k = 1, 2, \dots, m-1.$$

This group is free of rank $m - k$. Using the defining relation of P_m , it is not difficult to prove the following:

Lemma 1. *In P_n hold the following conjugation rules (for $\varepsilon = \pm 1$):*

- 1) $a_{kj}^{-\varepsilon} a_{ik} a_{kj}^{\varepsilon} = (a_{ik} a_{ij})^{\varepsilon} a_{ik} (a_{ik} a_{ij})^{-\varepsilon}$, where $i < k < j$;
- 2) $a_{jk}^{-\varepsilon} a_{ik} a_{jk}^{\varepsilon} = (a_{ij} a_{ik})^{\varepsilon} a_{ik} (a_{ij} a_{ik})^{-\varepsilon}$, where $i < j < k$;
- 3) $a_{kn}^{-\varepsilon} a_{ij} a_{kn}^{\varepsilon} = [a_{ik}^{-\varepsilon}, a_{in}^{-\varepsilon}]^{\varepsilon} a_{ij} [a_{ik}^{-\varepsilon}, a_{in}^{-\varepsilon}]^{-\varepsilon}$, where $i < k < j < n$;
- 4) $a_{in}^{-\varepsilon} a_{kj} a_{in}^{\varepsilon} = a_{kj}$, where $k < i$ and $n < j$;
- 5) $a_{kj}^{-\varepsilon} a_{in} a_{kj}^{\varepsilon} = a_{in}$, where $n < k$.

From this lemma it follows that V_1 is normal in P_m and we have decomposition $P_m = V_1 \rtimes P_{m-1}$. By induction on m , P_m is the semi-direct products of free groups:

$$P_m = V_1 \rtimes (V_2 \rtimes (\dots \rtimes (V_{m-2} \rtimes V_{m-1}) \dots)).$$

We will call this decomposition *horizontal decomposition*. Let $V_m^{(k)}$ be a subgroup of P_m which is generated by a_{ij} for $i < k$. Then we have the normal series

$$1 = V_m^{(1)} \leq V_m^{(2)} \leq \dots \leq V_m^{(m-1)} \leq V_m^{(m)} = P_m,$$

where

$$V_m^{(r)} / V_m^{(r-1)} \cong F_{m-r+1}, \quad r = 2, 3, \dots, m.$$

A motivation for the terms vertical and horizontal is as follows. If we put the generators of P_m in the following table

$$\begin{array}{cccc} a_{12}, & a_{13}, & \dots & a_{1,m-1}, & a_{1,m}, \\ & a_{23}, & \dots & a_{2,m-1}, & a_{2,m}, \\ \dots & \dots & \dots & \dots & \dots \\ & & & a_{m-2,m-1}, & a_{m-2,m}, \\ & & & & a_{m-1,m}, \end{array}$$

then the generators from the k th row generate V_k and the generators from the k th column generate U_{k+1} . The group $U_m^{(r)}$ is generated by the last $m-r$ columns of this table and the group $V_m^{(r)}$ is generated by the first $r-1$ rows of this table.

3 Braid groups in handlebodies

Recall some facts on the braid group $B_{g,n}$ on n strings in the handlebody \mathcal{H}_g (see [7, 12, 13]). The group $B_{g,n}$ is generated by elements

$$\tau_1, \tau_2, \dots, \tau_g, \sigma_{g+1}, \sigma_{g+2}, \dots, \sigma_{g+n-1},$$

and is defined by the following list of relations

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i, & \text{for } |i-j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, & \text{for } i = g+1, \dots, g+n-2, \\ \tau_k \sigma_i &= \sigma_i \tau_k, & \text{for } k \geq 1, i \geq g+2, \\ \tau_k (\sigma_{g+1} \tau_k \sigma_{g+1}) &= (\sigma_{g+1} \tau_k \sigma_{g+1}) \tau_k, & \text{for } k = 1, \dots, g, \\ \tau_k (\sigma_{g+1}^{-1} \tau_{k+l} \sigma_{g+1}) &= (\sigma_{g+1}^{-1} \tau_{k+l} \sigma_{g+1}) \tau_k, & \text{for } k = 1, \dots, g-1, l = 1, \dots, g-k. \end{aligned}$$

The group $B_{g,n}$ can be considered as a subgroup of the classical braid group B_{g+n} on $n+g$ strings such that the braids from $B_{g,n}$ leave the first g strings unbraided. Then $\tau_k = a_{k,g+1}$, in B_{g+n} , i. e.

$$\tau_k = \sigma_g \sigma_{g-1} \dots \sigma_{k+1} \sigma_k^2 \sigma_{k+1}^{-1} \dots \sigma_{g-1}^{-1} \sigma_g^{-1}, \quad k = 1, 2, \dots, g.$$

The elements τ_k , $k = 1, 2, \dots, g$, generate a free group of rank g which is isomorphic to $U_{g+1} = \langle a_{1,g+1}, a_{2,g+1}, \dots, a_{g,g+1} \rangle$ in B_{g+n} . Also, we see that some other generators of P_{g+n} lie in $B_{g,n}$. Put them in the table bellow:

$$\begin{array}{cccccc} a_{1,g+1}, & a_{1,g+2}, & \dots & a_{1,g+n-1}, & a_{1,g+n}, & \\ a_{2,g+1}, & a_{2,g+2}, & \dots & a_{2,g+n-1}, & a_{2,g+n}, & \\ \dots & \dots & \dots & \dots & \dots & \\ a_{g-1,g+1}, & a_{g-1,g+2}, & \dots & a_{g-1,g+n-1}, & a_{g-1,g+n}, & \\ a_{g,g+1}, & a_{g,g+2}, & \dots & a_{g,g+n-1}, & a_{g,g+n}, & \\ \hline & a_{g+1,g+2}, & \dots & a_{g+1,g+n-1}, & a_{g+1,g+n}, & \\ & \dots & \dots & \dots & \dots & \\ & & & a_{g+n-2,g+n-1}, & a_{g+n-2,g+n}, & \\ & & & & a_{g+n-1,g+n}. & \end{array}$$

We will denote by \tilde{B}_n the subgroup of $B_{g,n}$ which is generated by $\sigma_{g+1}, \sigma_{g+2}, \dots, \sigma_{g+n-1}$. It is evident that \tilde{B}_n is isomorphic to B_n . The corresponding pure braid group of \tilde{B}_n will be denote \tilde{P}_n . This group is isomorphic to P_n and is generated by elements from the above table which lie under the horizontal line. The group \tilde{P}_n has

the following vertical decomposition

$$\tilde{P}_n = \tilde{U}_n \wr (\tilde{U}_{n-1} \wr (\dots \wr (\tilde{U}_3 \wr \tilde{U}_2) \dots)),$$

where

$$\tilde{U}_i = \langle a_{g+1,g+i}, a_{g+2,g+i}, \dots, a_{g+i-1,g+i} \rangle, \quad i = 2, 3, \dots, n.$$

There is an epimorphism:

$$\psi_n : B_{g,n} \longrightarrow S_n,$$

which is defined by the rule

$$\psi_n(\tau_k) = 1, \quad k = 1, 2, \dots, g, \quad \psi_n(\sigma_i) = (i, i+1), \quad i = g+1, g+2, \dots, g+n-1.$$

This epimorphism is induced by the standard epimorphism $B_{g+n} \longrightarrow S_{g+n}$. Let $P_{g,n} = \text{Ker}(\psi_n)$. Then $P_{g,n}$ is generated by the element from the table above. In [7, 13] it was proved that there exists the following short exact sequence:

$$1 \longrightarrow P_{g,n} \longrightarrow B_{g,n} \longrightarrow S_n \longrightarrow 1,$$

and was found the vertical decomposition of $P_{g,n}$:

$$P_{g,n} = U_{g+n} \wr (U_{g+n-1} \wr (\dots \wr (U_{g+2} \wr U_{g+1}) \dots)).$$

If we let $U_{g+n}^{(g+n-i)} = U_{g+n} \wr (U_{g+n-1} \wr (\dots \wr (U_{g+n-i+2} \wr U_{g+n-i+1}) \dots))$, then we get the normal series:

$$1 = U_{g+n}^{(g+n)} \leq U_{g+n}^{(g+n-1)} \leq \dots \leq U_{g+n}^{(g)} = P_{g,n},$$

where

$$U_{g+n}^{(r)} / U_{g+n}^{(r+1)} \cong F_r, \quad r = g, g+1, \dots, g+n-1.$$

The homomorphism which is induced by the embedding $B_{g,n} \longrightarrow B_{g+n}$ sends this normal series to the corresponding normal series for P_{g+n} .

Let us construct the horizontal decomposition for $P_{g,n}$. To do this, define the following subgroups in $P_{g,n}$:

$$\begin{aligned} V_{g,1} &= \langle a_{1,g+1}, a_{1,g+2}, \dots, a_{1,g+n} \rangle, \\ V_{g,2} &= \langle a_{2,g+1}, a_{2,g+2}, \dots, a_{2,g+n} \rangle, \\ &\dots\dots\dots \\ V_{g,g} &= \langle a_{g,g+1}, a_{g,g+2}, \dots, a_{g,g+n} \rangle, \\ V_{g,g+1} &= \langle a_{g+1,g+2}, a_{g+1,g+3}, \dots, a_{g+1,g+n} \rangle, \\ V_{g,g+2} &= \langle a_{g+2,g+3}, a_{g+2,g+4}, \dots, a_{g+2,g+n} \rangle, \\ &\dots\dots\dots \\ V_{g,g+n-1} &= \langle a_{g+n-1,g+n} \rangle. \end{aligned}$$

We see that $V_{g,g+i} = V_{g+i}$ for all $i = 1, 2, \dots, n-1$, these subgroups lie in \tilde{P}_n and as we know

$$\tilde{P}_n = V_{g,g+1} \times (V_{g,g+2} \times (\dots \times (V_{g,g+n-2} \times V_{g,g+n-1}) \dots))$$

is the horizontal decomposition of \tilde{P}_n .

We see that the vertical decomposition of $P_{g,n}$ is a part of the vertical decomposition for P_{g+n} . For the horizontal decomposition situation is more complicated.

Example 1. The horizontal decomposition of P_4 has the form

$$P_4 = V_1 \times (V_2 \times V_3),$$

where

$$V_1 = \langle a_{12}, a_{13}, a_{14} \rangle, \quad V_2 = \langle a_{23}, a_{24} \rangle, \quad V_3 = \langle a_{34} \rangle,$$

and V_1 is normal in P_4 , V_2 is normal in $V_2 \times V_3$. The group $P_{2,2}$ contains subgroups

$$V_{2,1} = \langle a_{13}, a_{14} \rangle, \quad V_{2,2} = V_2 = \langle a_{23}, a_{24} \rangle, \quad V_{2,3} = V_3 = \langle a_{34} \rangle,$$

but in this case $V_{2,1}$ is not normal in $P_{2,2}$. Indeed, from Lemma 1 we have the following relations

$$a_{23}^{-\varepsilon} a_{13} a_{23}^{\varepsilon} = (a_{12} a_{13})^{\varepsilon} a_{13} (a_{12} a_{13})^{-\varepsilon}, \quad a_{24}^{-\varepsilon} a_{14} a_{24}^{\varepsilon} = (a_{12} a_{14})^{\varepsilon} a_{14} (a_{12} a_{14})^{-\varepsilon}.$$

Hence, $P_{2,2}$ contains not only $V_{2,1}$ but its normal closure in $P_{2,2}$ and we get the horizontal decomposition

$$P_{2,2} = \bar{V}_{2,1} \times (V_2 \times V_3),$$

where $\bar{V}_{2,1} = \langle V_{2,1} \rangle^{P_{2,2}}$ is the normal closure of $V_{2,1}$ in $P_{2,2}$.

In the general case, let $\bar{V}_{g,i}$ be the normal closure of $V_{g,i}$ in the subgroup $\langle V_{g,i}, V_{g,i+1}, \dots, V_{g,g}, \tilde{P}_n \rangle$, i.e.

$$\bar{V}_{g,i} = \langle V_{g,i} \rangle^{\langle V_{g,i}, V_{g,i+1}, \dots, V_{g,g}, \tilde{P}_n \rangle}.$$

Lemma 2. 1) $\bar{V}_{g,g} = V_{g,g} \cong F_n$.

2) $\bar{V}_{g,i}$ is a subgroup of V_i for every $i = 1, 2, \dots, g$ and, in particular, is a free group.

Proof. 1) We see that $V_{g,g} = V_g$ and from Lemma 1 it follows that V_g is normal in $\langle V_{g,g}, \tilde{P}_n \rangle$.

2) The fact that $\bar{V}_{g,i}$ is a subgroup of V_i follows from the conjugation rules of Lemma 1. The fact that $\bar{V}_{g,i}$ is free follows from the fact that V_i is free.

Since $\tilde{P}_n \cong P_n$, it has the horizontal decomposition:

$$\tilde{P}_n = V_{g+1} \times (V_{g+2} \times (\dots \times (V_{g+n-2} \times V_{g+n-1}) \dots)).$$

Using Lemma 2, one can construct the horizontal decomposition of $P_{g,n}$.

Theorem 1. *The group $P_{g,n}$ is the semi-direct products of groups:*

$$P_{g,n} = \bar{V}_{g,1} \times (\bar{V}_{g,2} \times (\dots \times (\bar{V}_{g,g} \times \tilde{P}_n) \dots)).$$

Proof. Use induction on g . If $g = 1$, then $P_{1,n} = P_{n+1}$ and the horizontal decomposition for P_{n+1} gives the horizontal decomposition: $P_{1,n} = V_g \times \tilde{P}_n$. As follows from Lemma 2, $V_g = \bar{V}_{g,g}$.

Let $g > 1$. Define a homomorphism of $P_{g,n}$ onto the group $P_{g-1,n}$ which sends all generators of $V_{g,1}$ to the unit and keeps all other generators. The kernel of this homomorphism is the normal closure of $V_{g,1}$ in $P_{g,n}$. Denote this kernel by $\bar{V}_{g,1} = \langle V_{g,1} \rangle^{P_{g,n}}$. Since, $V_{g,1}$ is a subgroup in V_1 and V_1 is normal in P_{g+n} we get that $\bar{V}_{g,1}$ lies in V_1 and hence is a free. We have the decomposition $P_{g,n} = \bar{V}_{g,1} \times P_{g-1,n}$. Using the induction hypothesis we get the required decomposition.

4 The kernel of the epimorphism $B_{g,n} \rightarrow \tilde{B}_n$

As was noted in [13] there is an epimorphism

$$\varphi_n : B_{g,n} \longrightarrow \tilde{B}_n,$$

where the subgroup $\tilde{B}_n = \langle \sigma_{g+1}, \sigma_{g+2}, \dots, \sigma_{g+n-1} \rangle$ is isomorphic to B_n . This endomorphism is defined by the rule

$$\varphi_n(\tau_k) = 1, \quad k = 1, 2, \dots, g, \quad \varphi_n(\sigma_i) = \sigma_i, \quad i = g+1, g+2, \dots, g+n-1.$$

If we denote $R_{g,n} = \text{Ker}(\varphi_n)$, then $B_{g,n} = R_{g,n} \times \tilde{B}_n$. The purpose of this section is the description of the group $R_{g,n}$.

Considering the table with the generators of $P_{g,n}$ (see Section 3), we see that all generators, which lie in the first g rows of this table, are elements of $R_{g,n}$. Denote by $Q_{g,n}$ the subgroup of $R_{g,n}$ that is generated by these elements, i.e. $Q_{g,n} = \langle V_{g,1}, V_{g,2}, \dots, V_{g,g} \rangle$. Then $R_{g,n} = \langle Q_{g,n} \rangle^{B_{g,n}}$ is the normal closure of $Q_{g,n}$ in $B_{g,n}$.

On the other side, if we denote

$$U_{g,g+i} = \langle a_{1,g+i}, a_{2,g+i}, \dots, a_{g,g+i} \rangle \leq U_{g+i}, \quad i = 1, 2, \dots, n,$$

then $Q_{g,n} = \langle U_{g,g+1}, U_{g,g+2}, \dots, U_{g,g+n} \rangle$. Note that $U_{g,g+1} = U_{g+1}$. Let $\bar{U}_{g,g+i}$ be the normal closure of $U_{g,g+i}$ in U_{g+i} , $i = 1, 2, \dots, n$. In this notations it holds:

Theorem 2. *The group $R_{g,n}$ has the following decompositions:*

1) $R_{g,n} = \overline{U}_{g,g+n} \wedge (\overline{U}_{g,g+n-1} \wedge (\dots \wedge (\overline{U}_{g,g+2} \wedge \overline{U}_{g,g+1}) \dots))$,
where $\overline{U}_{g,g+1} = U_{g+1}$.

2) $R_{g,n} = \overline{V}_{g,1} \wedge (\overline{V}_{g,2} \wedge (\dots \wedge (\overline{V}_{g,g-1} \wedge \overline{V}_{g,g}) \dots))$,
where $\overline{V}_{g,g} = V_g$.

Proof. As we know, $R_{g,n}$ is the normal closure of $Q_{g,n}$ in $B_{g,n}$. To find this closure, at first, consider conjugations of the generators of $Q_{g,n}$ by σ_{g+k} , $k = 1, 2, \dots, n-1$.

Conjugating generators from the k th column of our table by σ_{g+k} , we have

$$\sigma_{g+k}^{-1} a_{i,g+k} \sigma_{g+k} = a_{i,g+k} a_{i,g+k+1} a_{i,g+k}^{-1}, \quad i = 1, 2, \dots, g.$$

Conjugating generators from the $(k+1)$ st column of our table by σ_{g+k} , we have

$$\sigma_{g+k}^{-1} a_{i,g+k+1} \sigma_{g+k} = a_{i,g+k}, \quad i = 1, 2, \dots, g.$$

Generators from all other columns commute with σ_{g+k} .

Hence, for every generator a_{ij} of $Q_{g,n}$ and every σ_k , $k = g+1, g+2, \dots, g+n-1$, the element $\sigma_k^{-1} a_{ij} \sigma_k$ lies in $Q_{g,n}$.

For the group \tilde{B}_n there exists the following exact sequence

$$1 \longrightarrow \tilde{P}_n \longrightarrow \tilde{B}_n \longrightarrow S_n \longrightarrow 1.$$

Let $m_{kl} = \sigma_{k-1} \sigma_{k-2} \dots \sigma_l$ for $l < k$ and $m_{kl} = 1$ in other cases. Then the set

$$\Lambda_n = \left\{ \prod_{k=g+2}^{g+n} m_{k,j_k} \mid 1 \leq j_k \leq k \right\}$$

is a Schreier set of coset representatives of \tilde{P}_n in \tilde{B}_n .

From the previous observations it follows that if $\alpha \in \Lambda$, then for every generator a_{ij} of $Q_{g,n}$ the element $\alpha^{-1} a_{ij} \alpha$ lies in $Q_{g,n}$.

Now we will consider the conjugations of the generators of $Q_{g,n}$ by the generators of $P_{g,n}$.

1) To prove the first decomposition, take some element $h \in B_{g,n}$ and using the vertical decomposition, write it in the normal form:

$$h = u_{g+1} u_{g+2} \dots u_{g+n} \alpha, \quad \text{where } u_k \in U_k, \quad \alpha \in \Lambda_n.$$

In every group U_k , $k = g+1, g+2, \dots, g+n$, we have two subgroups:

$$U_{g,k} = \langle a_{1k}, a_{2k}, \dots, a_{gk} \rangle, \quad \tilde{U}_k = \langle a_{g+1,k}, a_{g+2,k}, \dots, a_{k-1,k} \rangle \leq \tilde{P}_n.$$

We see that $U_{g,k}$ is a subgroup of $P_{g,n}$, \tilde{U}_k is a subgroup of \tilde{P}_n and $U_k = \langle U_{g,k}, \tilde{U}_k \rangle$ is a free group. Define the projection $\pi_k : U_k \longrightarrow \tilde{U}_k$ by the rules

$$\pi_k(a_{ik}) = 1, \quad i = 1, 2, \dots, g;$$

$$\pi_k(a_{jk}) = a_{jk}, \quad j = g+1, g+2, \dots, k-1.$$

The kernel $\text{Ker}(\pi_k) = \overline{U}_{g,k}$ is the normal closure $\langle U_{g,k} \rangle^{U_k}$ of $U_{g,k}$ into U_k . Hence, element u_k can be written in the form $u_k = \overline{u}_k \pi_k(u_k)$ for some $\overline{u}_k \in \overline{U}_{g,k}$. Denote for simplicity $w_{g+i} = \pi_{g+1}(u_{g+i})$, we can rewrite h in the form

$$h = (\overline{u}_{g+1} w_{g+1}) (\overline{u}_{g+2} w_{g+2}) \dots (\overline{u}_{g+n} w_{g+n}) \alpha.$$

Shifting all w_{g+i} to the right we get

$$h = \overline{u}_{g+1} \overline{u}_{g+2}^{w_{g+1}^{-1}} \dots \overline{u}_{g+n}^{w_{g+n-1}^{-1} w_{g+n-2}^{-1} \dots w_{g+1}^{-1}} w_{g+1} w_{g+2} \dots w_{g+n} \alpha.$$

The image $\varphi_n(h) = w_{g+1} w_{g+2} \dots w_{g+n} \alpha$. Hence, the element

$$\overline{u}_{g+1} \overline{u}_{g+2}^{w_{g+1}^{-1}} \dots \overline{u}_{g+n}^{w_{g+n-1}^{-1} w_{g+n-2}^{-1} \dots w_{g+1}^{-1}}$$

lies in the kernel $\text{Ker}(\varphi_n) = R_{g,n}$. We proved that any element from $R_{g,n}$ lies in the product

$$\overline{U}_{g,g+n} \wr (\overline{U}_{g,g+n-1} \wr (\dots \wr (\overline{U}_{g,g+2} \wr \overline{U}_{g,g+1}) \dots)).$$

On the other side, this product contains $Q_{g,n}$ and is normal in $B_{g,n}$. Hence, $R_{g,n}$ has the required decomposition.

2) Prove the second decomposition. Denote $G = \overline{V}_{g,1} \wr (\overline{V}_{g,2} \wr (\dots \wr (\overline{V}_{g,g-1} \wr \overline{V}_{g,g}) \dots))$ the group from the right side of the decomposition of $R_{g,n}$. Take an arbitrary element $h \in B_{g,n}$. By the theorem on the horizontal decomposition of $B_{g,n}$ it has the following normal form:

$$h = v_1 v_2 \dots v_g b, \quad \text{where } v_i \in V_{g,i}, b \in \widetilde{B}_n$$

and $v_1 v_2 \dots v_g$ lies in $R_{g,n}$. Under the endomorphism $\varphi_n : B_{g,n} \longrightarrow \widetilde{B}_n$ element h goes to the element b . Hence, G lies in $R_{g,n}$.

On the other side we must prove that $R_{g,n}$ lies in G . We know that $R_{g,n}$ is the normal closure of $Q_{g,n}$ in $B_{g,n}$. As was shown before, if $\alpha \in \Lambda_n$ is a coset representative of \widetilde{P}_n into \widetilde{B}_n and a_{ij} is some generator of $Q_{g,n}$, then a_{ij}^α lies in $Q_{g,n}$. Considering the conjugation of $a_{ij} \in Q_{g,n}$ by generators of $P_{g,n}$ and using the Lemma 1 we get an element which lies in G .

In the case $g = 1$ we have $P_{1,n} = P_{n+1}$ and $R_{1,n} = \langle a_{12}, a_{13}, \dots, a_{1,n+1} \rangle \cong F_n$ and we get the decomposition which was found in [8].

Corollary 1. $B_{1,n} \cong F_n \wr B_n$.

5 Some analog of the Hecke algebra for the braid group in the handlebody

Let q be some complex number. Recall that the Hecke algebra $H_n(q)$ is an associative \mathbb{C} -algebra with unit, which is generated by elements

$$s_1, s_2, \dots, s_{n-1},$$

and is defined by the relations

$$\begin{aligned} s_i s_j &= s_j s_i, & \text{for } |i - j| > 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & \text{for } 1 \leq i \leq n - 2, \\ s_i^2 &= (q - 1)s_i + q, & \text{for } i = 1, \dots, n - 1. \end{aligned}$$

The algebra $H_n(q)$ has the following linear basis:

$$S = \{(s_{i_1} s_{i_1-1} \dots s_{i_1-k_1})(s_{i_2} s_{i_2-1} \dots s_{i_2-k_2}) \dots (s_{i_p} s_{i_p-1} \dots s_{i_p-k_p})\}$$

for $1 \leq i_1 < \dots < i_p \leq n - 1$. The basis S is used in the construction of the Markov trace, leading to the HOMFLYPT or 2-variable Jones polynomial (see [3]).

The braid group in the solid torus $B_{1,n}$ is the Artin group of the Coxeter group of type \mathcal{B} , which is related to the Hecke algebra of type \mathcal{B} . The generalized Hecke algebra of type \mathcal{B} , $H_{1,n}(q)$ is defined by S. Lambropoulou in [6]. $H_{1,n}(q)$ is isomorphic to the affine Hecke algebra of type \mathcal{A} , $\tilde{H}_n(q)$. A unique Markov trace is constructed on the algebras $H_{1,n}(q)$ that leads to an invariant for links in the solid torus, the universal analogue of the HOMFLYPT polynomial for the solid torus.

The algebra $H_{1,n}(q)$ is generated by elements

$$t, t^{-1}, s_1, s_2, \dots, s_{n-1},$$

and is defined by the relations

$$\begin{aligned} s_i s_j &= s_j s_i, & \text{for } |i - j| > 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & \text{for } 1 \leq i \leq n - 2, \\ s_i^2 &= (q - 1)s_i + q, & \text{for } i = 1, 2, \dots, n - 1, \\ s_1 t s_1 t &= t s_1 t s_1, \\ t s_i &= s_i t, & \text{for } i > 1. \end{aligned}$$

Hence,

$$H_{1,n}(q) = \frac{\mathbb{C}[B_{1,n}]}{\langle \sigma_i^2 - (q - 1)\sigma_i - q \rangle}.$$

Note that in $H_{1,n}(q)$ the generator t satisfies no polynomial relation, making the algebra $H_{1,n}(q)$ infinite dimensional. If we set $t = 0$ in $H_{1,n}(q)$, we obtain the Hecke algebra $H_n(q)$.

In $H_{1,n}(q)$ are defined in [8] the elements

$$t_i = s_i s_{i-1} \dots s_1 t s_1 \dots s_{i-1} s_i, \quad t'_i = s_i s_{i-1} \dots s_1 t s_1^{-1} \dots s_{i-1}^{-1} s_i^{-1}.$$

It was then proved that the following sets form linear bases for $H_{1,n}(q)$:

$$\Sigma_n = t_{i_1}^{k_1} t_{i_2}^{k_2} \dots t_{i_r}^{k_r} \sigma, \quad \text{where } 1 \leq i_1 < \dots < i_r \leq n-1,$$

$$\Sigma'_n = (t'_{i_1})^{k_1} (t'_{i_2})^{k_2} \dots (t'_{i_r})^{k_r} \sigma, \quad \text{where } 1 \leq i_1 < \dots < i_r \leq n-1,$$

where $k_1, k_2, \dots, k_r \in \mathbb{Z}$ and σ a basis element in $H_n(q)$. The basis Σ'_n is used in [6, 8] for constructing a Markov trace on $\bigcup_{n=1}^{\infty} H_{1,n}(q)$ and a universal HOMFLYPT-type invariant for oriented links in the solid torus.

Definition 1. Let $q \in \mathbb{C}$. The algebra $H_{g,n}(q)$ is an associative algebra over \mathbb{C} with unit that is generated by

$$t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_g^{\pm 1}, s_{g+1}, s_{g+2}, \dots, s_{g+n-1}$$

and is defined by the following relations

$$\begin{aligned} s_i s_j &= s_j s_i, & \text{for } |i-j| > 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & \text{for } i = g+1, g+2, \dots, g+n-2, \\ s_i^2 &= (q-1)s_i + q, & \text{for } i = g+1, g+2, \dots, g+n-1, \\ t_k s_i &= s_i t_k, & \text{for } k \geq 1, i \geq g+2, \\ t_k (s_{g+1} t_k s_{g+1}) &= (s_{g+1} t_k s_{g+1}) t_k, & \text{for } k = 1, 2, \dots, g, \\ t_k (s_{g+1} t_{k+l} s_{g+1}) &= (s_{g+1} t_{k+l} s_{g+1}) t_k, & \text{for } k = 1, 2, \dots, g-1; l = 1, 2, \dots, g-k. \end{aligned}$$

Remark 1. More natural to consider the generators s_1, s_2, \dots, s_{n-1} instead $s_{g+1}, s_{g+2}, \dots, s_{g+n-1}$, but for technical reasons we will use our notation.

We see that

$$H_{g,n}(q) = \frac{\mathbb{C}[B_{g,n}]}{\langle \sigma_i^2 - (q-1)\sigma_i - q \rangle}.$$

If we consider the algebra $H_n(q)$ as a vector space over \mathbb{C} , then it is isomorphic to the vector space $\mathbb{C}[S_n]$. Thus to study $H_{g,n}(q)$, we define a group $G_{g,n}$ as the quotient of $B_{g,n}$ by the relations

$$\sigma_i^2 = 1, \quad i = 1, 2, \dots, n-1.$$

Denote the natural homomorphism $B_{g,n} \longrightarrow G_{g,n}$ by ψ and let $\psi(a_{ij}) = b_{ij}$.

Theorem 3. *The group $G_{g,n}$ is the semi-direct product $G_{g,n} = F_g^n \rtimes S_n$ of the direct product of n copies of free group F_g of rank g and the symmetric group S_n .*

Proof. We know that $B_{g,n} = R_{g,n} \rtimes \tilde{B}_n$ and

$$R_{g,n} = \overline{U}_{g,g+n} \rtimes (\overline{U}_{g,g+n-1} \rtimes (\dots \rtimes (\overline{U}_{g,g+2} \rtimes \overline{U}_{g,g+1}) \dots)).$$

On the other side, U_k , $k = g+1, g+2, \dots, g+n$, is generated by two subgroups:

$$U_{g,k} = \langle a_{1k}, a_{2k}, \dots, a_{gk} \rangle \text{ and } \tilde{U}_k = \langle a_{g+1,k}, a_{g+2,k}, \dots, a_{k-1,k} \rangle \leq \tilde{P}_n.$$

Note that under the homomorphism ψ the subgroups \tilde{U}_k go to 1. Hence, $\psi(\overline{U}_{g,g+k}) = \psi(U_{g,g+k}) \cong F_g$ and

$$\psi(R_{g,n}) = \psi(U_{g,g+n}) \rtimes (\psi(U_{g,g+n-1}) \rtimes (\dots \rtimes (\psi(U_{g,g+2}) \rtimes \psi(U_{g,g+1})) \dots))$$

is the semi-direct product of n copies of free group F_g . Also, $G_{g,n} = \psi(R_{g,n}) \rtimes \psi(\tilde{B}_n) \cong \psi(R_{g,n}) \rtimes S_n$.

Let us show that in fact $\psi(R_{g,n})$ is the direct product of n copies of free group F_g . To do it it is enough to prove that

$$\psi(R_{g,n}) = \psi(U_{g,g+n}) \times \psi(R_{g,n-1}).$$

Denote $b_{ij} = \psi(a_{ij})$, consider the relations (11)-(14) and find their images under the homomorphism ψ . The relation (11) goes to the relation

$$b_{i,g+j}^{-\varepsilon} b_{g+j,g+n} b_{i,g+j}^{\varepsilon} = (b_{i,g+n} b_{g+j,g+n})^{\varepsilon} b_{g+j,g+n} (b_{i,g+n} b_{g+j,g+n})^{-\varepsilon},$$

but $b_{g+j,g+n} = 1$ and we have the trivial relations.

The relation (12) goes to the relation

$$b_{l,g+j}^{-\varepsilon} b_{l,g+n} b_{i,g+j}^{\varepsilon} = (b_{l,g+n} b_{g+j,g+n})^{\varepsilon} b_{l,g+n} (b_{l,g+n} b_{g+j,g+n})^{-\varepsilon},$$

but $b_{g+j,g+n} = 1$ and we have the relation

$$b_{l,g+j}^{-\varepsilon} b_{l,g+n} b_{i,g+j}^{\varepsilon} = b_{l,g+n}.$$

The relation (13) goes to the relation

$$b_{i,g+j}^{-\varepsilon} b_{l,g+n} b_{i,g+j}^{\varepsilon} = [b_{i,g+n}^{-\varepsilon}, b_{g+j,g+n}^{-\varepsilon}]^{\varepsilon} b_{l,g+n} [b_{i,g+n}^{-\varepsilon}, b_{g+j,g+n}^{-\varepsilon}]^{-\varepsilon}, \quad i < l < g + j,$$

but $b_{g+j,g+n} = 1$ and we have the relation

$$b_{i,g+j}^{-\varepsilon} b_{l,g+n} b_{i,g+j}^{\varepsilon} = b_{l,g+n}, \quad i < l < g + j.$$

The relation (14) goes to the relation

$$b_{i,g+j}^{-\varepsilon} b_{l,g+n} b_{i,g+j}^{\varepsilon} = b_{l,g+n}, \quad l < i < g + j < g + n.$$

The image of $\tilde{B}_n \leq B_{g,n}$ is isomorphic to S_n . Thus from the decomposition $B_{g,n} = R_{g,n} \rtimes \tilde{B}_n$ we get the required decomposition for $G_{g,n}$.

From this theorem we have

Corollary 2. *Every element $h \in G_{g,n}$ has the unique normal form*

$$h = h_1 h_2 \dots h_n \alpha,$$

where h_i is a reduced word in the free group

$$\psi(U_{g,g+i}) = \langle b_{1,g+i}, b_{2,g+i}, \dots, b_{g,g+i} \rangle \cong F_g, \quad i = 1, 2, \dots, n,$$

$\alpha \in \Lambda_n$ is a coset representative of $\psi(R_{g,n})$ in $G_{g,n}$.

Using this normal form and the fact that these normal forms form a linear basis of $\mathbb{C}[G_{g,n}]$, we can try to define a basis of $H_{g,n}(q)$. In the algebra $H_{g,n}(q)$ define the following elements

$$\begin{aligned} t_{1,g+1} &= t_1, & t_{1,g+2}, & \dots, & t_{1,g+n}, \\ t_{2,g+1} &= t_2, & t_{2,g+2}, & \dots, & t_{2,g+n}, \\ & \dots & & \dots & \dots \\ t_{g,g+1} &= t_g, & t_{g,g+2}, & \dots, & t_{g,g+n}, \end{aligned}$$

where

$$t_{ij} = s_{j-1}s_{j-2}\dots s_1 t_{i,g+1} s_1 \dots s_{j-2} s_{j-1}, \quad 1 \leq i \leq g, \quad g+2 \leq j \leq g+n.$$

These elements correspond to the elements b_{ij} from $G_{g,n}$, which were defined in 3 as the images of the elements a_{ij} under the map ψ . It is not difficult to see that the elements

$$t_{1,g+i}, t_{2,g+i}, \dots, t_{g,g+i}$$

generate a free group of rank g .

The algebra $H_{g,n}(q)$ contains the subalgebra with the set of generators $s_{g+1}, s_{g+2}, \dots, s_{g+n-1}$, which is isomorphic to $H_n(q)$. Let $\Sigma_{g,n}$ be the following set in $H_{g,n}(q)$:

$$u_1 u_2 \dots u_n \sigma,$$

where u_i is a reduced word in the free group

$$\langle t_{1,g+i}, t_{2,g+i}, \dots, t_{g,g+i} \rangle \cong F_g, \quad i = 1, 2, \dots, n,$$

σ is a basis element in $H_n(q)$.

Conjecture 1. There is an isomorphism $H_{g,n}(q) \cong \mathbb{C}[G_{g,n}]$ as \mathbb{C} -modules. In particular, the set $\Sigma_{g,n}$ is a basis of the algebra $H_{g,n}(q)$.

The algebra $H_{2,n}(q)$ is the subject of study in [5], where one set of elements in $H_{2,n}(q)$, different from our $\Sigma_{2,n}$, is proved to be a spanning set for the algebra.

At the end, we formulate the following question for further investigation.

Question 1. Is it possible to define a Markov trace on the algebra $\bigcup_{n=1}^{\infty} H_{g,n}(q)$ and construct some analogue of the HOMLYPT polynomial that is an invariant of links in the handlebody \mathcal{H}_g ?

References

1. J. S. Birman, *Braids, Links, and Mapping Class Groups*, *Annals of Math. Studies* 82, Princeton University Press, 1974.
2. I. Diamantis, S. Lambropoulou, J. Przytycki, *Topological steps toward the HOMFLYPT skein module of the lens spaces $L(p, 1)$ via braids*, *J. Knot Theory Ramifications*, **25**, N 14 (2016), 1650084 (26 pages). See arXiv:1604.06163.
3. V. F. R. Jones, *Hecke algebra representations of braid groups and links polynomials*, *Ann. Math.*, 1984, **126**, 335–388.
4. R. Haring-Oldenburg, S. Lambropoulou *Knot theory in handlebodies*, *J. Knot Theory Ramifications*, **11**, N 6 (2002), 921–943.
5. D. Kodokostas and S. Lambropoulou, *A spanning set and potential basis of the mixed Hecke algebra on two fixed strands*, arXiv: 1704.03676.
6. S. Lambropoulou, *Solid torus links and Hecke algebras of B-type*, Proc. of the Conf. of Quantum Topology, D. N. Yetter ed., World Scientific Press, 1994.
7. S. Lambropoulou, *Braid structure in knot complements, handlebodies and 3-manifold*, Knots in Hellas '98 (Delphi), Proc. of International Conference Knot Theory and its Ramifications, WS, 274–289.
8. S. Lambropoulou, *Knot theory related to generalized and cyclotomic Hecke algebras of type B*, *J. Knot Theory Ramifications*, **8**, N 5 (1999), 621–658.
9. S. Lambropoulou, C.P. Rourke *Markov's theorem in 3-manifolds*, *Topology and its Applications*, 1997, **78**, 95–122.
10. S. Lambropoulou, C.P. Rourke *Algebraic Markov equivalence for links in 3-manifolds*, *Compositio Mathematica*, 2006, **142**, 1039–1062.
11. A. A. Markov, *Foundations of the algebraic theory of braids*, *Trudy Mat. Inst. Steklov [Proc. Steklov Inst. Math.]*, 1945, **16**, 1–54.
12. A. B. Sossinsky, *Preparation theorem for isotopy invariants of links in 3-manifold*, Quantum Groups. Proc. of the Conf. on Quantum Groups. Berlin a.: Springer-Verl., 1992, 354–362 (Lecture Notes in Math.; N 1510).
13. V. V. Vershinin, *On braid groups in handlebodies*, *Siberian Math. J.*, **39**, N 4 (1998), 645–654. Translation from *Sibirsk. Mat. Zh.*, **39**, N 4 (1998), 755–764.
14. V. V. Vershinin, *Generalization of braids from a homological point of view*, *Sib. Adv. Math.*, **9**, N 2 (1999), 109–139.