# Cycle packings of the complete multigraph

Rosalind A. Cameron
Department of Mathematics and Statistics
Memorial University of Newfoundland
St John's, NL
rahoyte@outlook.com

#### Abstract

Bryant, Horsley, Maenhaut and Smith recently gave necessary and sufficient conditions for when the complete multigraph can be decomposed into cycles of specified lengths  $m_1, m_2, \ldots, m_{\tau}$ . In this paper we characterise exactly when there exists a packing of the complete multigraph with cycles of specified lengths  $m_1, m_2, \ldots, m_{\tau}$ . While cycle decompositions can give rise to packings by removing cycles from the decomposition, in general it is not known when there exists a packing of the complete multigraph with cycles of various specified lengths.

#### 1 Introduction

A decomposition of a multigraph G is a collection  $\mathcal{D}$  of submultigraphs of G such that each edge of G is in exactly one of the multigraphs in  $\mathcal{D}$ . A packing of a multigraph G is a collection  $\mathcal{P}$  of submultigraphs of G such that each edge of G is in at most one of the multigraphs in  $\mathcal{P}$ . The leave of a packing  $\mathcal{P}$  is the multigraph obtained by removing the edges in multigraphs in  $\mathcal{P}$  from G. A cycle packing of a multigraph G is a packing  $\mathcal{P}$  of G such that each submultigraph in  $\mathcal{P}$  is a cycle. For positive integers  $\lambda$  and v,  $\lambda K_v$  denotes the complete multigraph with  $\lambda$  parallel edges between each pair of v distinct vertices. Here we give a complete characterisation of when there exists a packing of  $\lambda K_v$  with cycles of specified lengths  $m_1, m_2, \ldots, m_{\tau}$ . Note that for  $\lambda \geqslant 2$ ,  $\lambda K_v$  contains 2-cycles (pairs of parallel edges).

**Theorem 1.** Let  $m_1, m_2, \ldots, m_{\tau}$  be a nondecreasing list of integers and let  $\lambda$  and v be positive integers. Then there exists a packing of  $\lambda K_v$  with cycles of lengths  $m_1, m_2, \ldots, m_{\tau}$  if and only if

- (i)  $2 \leqslant m_1 \leqslant m_2, \ldots, m_{\tau} \leqslant v;$
- (ii)  $m_1 + m_2 + \cdots + m_{\tau} = \lambda {v \choose 2} \delta$ , where  $\delta$  is a nonnegative integer such that  $\delta \neq 1$  when  $\lambda(v-1)$  is even,  $\delta \neq 2$  when  $\lambda = 1$ , and  $\delta \geqslant \frac{v}{2}$  when  $\lambda(v-1)$  is odd;
- (iii)  $\sum_{m_i=2} m_i \leqslant \begin{cases} (\lambda 1)\binom{v}{2} 2 & \text{if } \lambda \text{ and } v \text{ are odd and } \delta = 2, \\ (\lambda 1)\binom{v}{2} & \text{if } \lambda \text{ is odd; and} \end{cases}$
- (iv)  $m_{\tau} \leqslant \begin{cases} \frac{\lambda}{2} {v \choose 2} \tau + 2 & \text{if } \lambda \text{ is even and } \delta = 0, \\ \frac{\lambda}{2} {v \choose 2} \tau + 1 & \text{if } \lambda \text{ is even and } 2 \leqslant \delta < m_{\tau}. \end{cases}$

Bryant, Horsley, Maenhaut and Smith [5] recently characterised exactly when there exists a decomposition of the complete multigraph  $\lambda K_v$  into cycles of specified lengths  $m_1, m_2, \ldots, m_{\tau}$  (see also [4, 12]). Since a decomposition of a multigraph is a packing whose leave contains no edges, many instances of the cycle packing problem can be solved by removing cycles from a cycle decomposition  $\lambda K_v$ . However there are cases which cannot be solved in this manner. These cases occur when  $\lambda(v-1)$  is odd and there are  $\frac{v}{2}+1$  or  $\frac{v}{2}+2$  edges in the leave of the required packing.

In the case of the complete graph  $K_v$  (with  $\lambda = 1$ ), it had previously been found exactly when there exist decompositions into cycles of specified lengths [6]. Furthermore, Horsley [10] found conditions for the existence of packings of the complete graph with uniform length cycles. These results built on earlier results for cycle decompositions and packings of the complete graph [1, 2, 9, 11] (see [7] for a survey). However, even in the  $\lambda = 1$  case, a complete characterisation of when there exists a packing of  $K_v$  with cycles of lengths  $m_1, m_2, \ldots, m_\tau$  had not previously been obtained.

We will show that the necessity of conditions (i)–(iv) in Theorem 1 follows from known results for cycle decompositions of  $\lambda K_v$ . The sufficiency of these conditions is proved by first decomposing  $\lambda K_v$  into cycles (and a 1-factor if  $\lambda(v-1)$  is even) and then removing cycles and modifying the resulting packing to obtain the one that we require. The existence of these cycle decompositions of  $\lambda K_v$  was obtained by Bryant et al [5] and the exact result is stated as Theorem 5 in Section 3. Section 2 contains the results required for modifying cycle decompositions.

The following definitions and notation will be used throughout this paper. An  $(m_1, m_2, \ldots, m_{\tau})$ -decomposition of  $\lambda K_v$  is a decomposition of  $\lambda K_v$  into  $\tau$  cycles of lengths  $m_1, m_2, \ldots, m_{\tau}$ . Similarly, an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$  is a packing of  $\lambda K_v$  with  $\tau$  cycles of lengths  $m_1, m_2, \ldots, m_{\tau}$ . We shall write  $(m_1^{\ell_1}, m_2^{\ell_2}, \ldots, m_{\tau}^{\ell_{\tau}})$  to denote the list of integers  $(m_1, \ldots, m_1, m_2, \ldots, m_2, \ldots, m_{\tau}, \ldots, m_{\tau})$ .

For vertices x and y in a multigraph G, the multiplicity of xy is the number of edges in G which have x and y as their endpoints, denoted  $\mu_G(xy)$ . If  $\mu_G(xy) \leq 1$  for all pairs of vertices in V(G) then we say that G is a simple graph. A multigraph is said to be even if every vertex has even degree and is said to be odd if every vertex has odd degree.

Given a permutation  $\pi$  of a set V, a subset S of V and a multigraph G with  $V(G) \subseteq V$ ,  $\pi(S)$  is defined to be the set  $\{\pi(x) : x \in S\}$  and  $\pi(G)$  is defined to be the multigraph with vertex set  $\pi(V(G))$  and edge set  $\{\pi(x)\pi(y) : xy \in E(G)\}$ . The m-cycle with vertices  $x_0, x_1, \ldots, x_{m-1}$  and edges  $x_i x_{i+1}$  for  $i \in \{0, \ldots, m-1\}$  (with subscripts modulo m) is denoted by  $(x_0, x_1, \ldots, x_{m-1})$  and the n-path with vertices  $y_0, y_1, \ldots, y_n$  and edges  $y_j y_{j+1}$  for  $j \in \{0, 1, \ldots, n-1\}$  is denoted by  $[y_0, y_1, \ldots, y_n]$ .

A chord of a cycle is an edge which is incident with two vertices of the cycle but is not in the cycle. Note that a chord may be an edge parallel to an edge of the cycle. For integers  $p \ge 2$  and  $q \ge 1$ , a (p,q)-lasso is the union of a p-cycle and a q-path such that the cycle and the path share exactly one vertex and that vertex is an end-vertex of the path. A (p,q)-lasso with cycle  $(x_1, x_2, \ldots, x_p)$  and path  $[x_p, y_1, y_2, \ldots, y_q]$  is denoted by  $(x_1, x_2, \ldots, x_p)[x_p, y_1, y_2, \ldots, y_q]$ . The order of a (p,q)-lasso is p+q.

# 2 Modifying cycle packings of $\lambda K_v$

The aim of this section is to prove Lemmas 3 and 4. These results are useful tools for modifying cycle packings of the complete multigraph. The simple graph versions of Lemmas 3 and 4 are due to Bryant and Horsley [8] and have been applied to prove the maximum packing result of the simple complete graph with uniform length cycles [10].

We require the following cycle switching lemma for cycle packings of multigraphs. Lemma 2 is similar to [4, Lemma 2.1] and is also closely related to the cycle switching method which has been applied to simple graphs (see for example [3]).

**Lemma 2.** Let v and  $\lambda$  be positive integers, let M be a list of integers, let  $\mathcal{P}$  be an (M)packing of  $\lambda K_v$ , let L be the leave of  $\mathcal{P}$ , let  $\alpha$  and  $\beta$  be distinct vertices of L, and let  $\pi$  be the
transposition  $(\alpha\beta)$ . Let E be a subset of E(L) such that, for each vertex  $x \in V(L) \setminus \{\alpha, \beta\}$ , E contains precisely  $\max(0, \mu_L(x\alpha) - \mu_L(x\beta))$  edges with endpoints x and  $\alpha$ , and precisely  $\max(0, \mu_L(x\beta) - \mu_L(x\alpha))$  edges with endpoints x and  $\beta$  (so E may contain multiple edges with
the same endpoints), and E contains no other edges. Then there exists a partition of E into
pairs such that for each pair  $\{x_1y_1, x_2y_2\}$  of the partition, there exists an (M)-packing  $\mathcal{P}'$  of  $\lambda K_v$  with leave  $L' = (L - \{x_1y_1, x_2y_2\}) + \{\pi(x_1)\pi(y_1), \pi(x_2)\pi(y_2)\}$ .

Furthermore, if  $\mathcal{P} = \{C_1, \ldots, C_t\}$ , then  $\mathcal{P}' = \{C'_1, \ldots, C'_t\}$  where for  $i \in \{1, \ldots, t\}$ ,  $C'_i$  is a cycle of the same length as  $C_i$  such that for  $i \in \{1, \ldots, t\}$ 

- If neither  $\alpha$  nor  $\beta$  is in  $V(C_i)$ , then  $C'_i = C_i$ ;
- If exactly one of  $\alpha$  and  $\beta$  is in  $V(C_i)$ , then  $C'_i = C_i$  or  $C'_i = \pi(C_i)$ ; and
- If both  $\alpha$  and  $\beta$  are in  $V(C_i)$ , then  $C'_i = Q_i \cup Q_i^*$  where  $Q_i = P_i$  or  $\pi(P_i)$ ,  $Q_i^* = P_i^*$  or  $\pi(P_i^*)$ , and  $P_i$  and  $P_i^*$  are the two paths from  $\alpha$  to  $\beta$  in  $C_i$ .

**Proof.** When  $\lambda(v-1)$  is even, Lemma 2 reduces to [4, Lemma 2.1] so suppose  $\lambda(v-1)$  is odd. Note that  $\mathcal{P}$  is a cycle packing of  $\lambda K_v$  regardless of the parity of  $\lambda(v-1)$ , whereas when  $\lambda(v-1)$  is odd [4, Lemma 2.1] concerns a cycle packing of  $\lambda K_v - I$ , where I is a 1-factor of  $\lambda K_v$ . Nevertheless, the proof of Lemma 2 follows from similar arguments to those used in the corresponding case of the proof in [4].

In applying Lemma 2 we say that we are performing the  $(\alpha, \beta)$ -switch with origin x and terminus y (where  $\{x_1, y_1, x_2, y_2\} \subseteq \{\alpha, \beta, x, y\}$ ). Note that  $x_1y_1$  and  $x_2y_2$  may be parallel edges, in which case x = y.

**Lemma 3.** Let v, s and  $\lambda$  be positive integers such that  $s \ge 3$ , and let M be a list of integers. Suppose there exists an (M)-packing  $\mathcal{P}$  of  $\lambda K_v$  whose leave contains a lasso of order at least s+2 and suppose that if s is even then the cycle of the lasso has even length. Then there exists an (M,s)-packing of  $\lambda K_v$ .

**Proof.** Let L be the leave of  $\mathcal{P}$ . Suppose that L contains a (p,q)-lasso  $(x_1, x_2, \ldots, x_p)[x_p, y_1, y_2, \ldots, y_q]$  such that  $p+q \ge s+2$  and p is even if s is even. If L contains an s-cycle then we add it to the packing to complete the proof, so assume L does not contain an s-cycle and hence  $p \ne s$ .

Case 1. Suppose  $2 \le p < s$  and either p = 2 or  $p \equiv s \pmod{2}$ . We can assume that p+q = s+2 since L contains a (p, s+2-p)-lasso.

Let L' be the leave of the packing  $\mathcal{P}'$  obtained from  $\mathcal{P}$  by applying the  $(x_1, y_{q-1})$ -switch with origin  $x_2$  (note that  $\mu_L(x_2y_{q-1}) = 0$  for otherwise L contains an s-cycle). If the terminus of the switch is not  $y_{q-2}$  then L' contains an s-cycle which completes the proof (recall that s = p + q - 2). Otherwise, the terminus of the switch is  $y_{q-2}$  and L' contains a (q, p)-lasso  $(x'_1, x'_2, \ldots, x'_q)[x'_q, y'_1, y'_2, \ldots, y'_p]$ . If p = 2 then L' contains an s-cycle which completes the proof, so assume L' contains no s-cycle and  $p \geqslant 3$ .

Let L'' be the leave of the packing  $\mathcal{P}''$  obtained from  $\mathcal{P}'$  by applying the  $(x'_2, y'_p)$ -switch with origin  $x'_3$  (note that  $\mu_{L'}(x'_3y'_p) = 0$  for otherwise L' contains an s-cycle). If the terminus of this switch is not  $y'_{p-1}$  then L'' contains an s-cycle which completes the proof (recall that s = p+q-2). Otherwise, the terminus of the switch is  $y'_{p-1}$  and L'' contains a (p+2, q-2)-lasso, so since p < s and  $p \equiv s \pmod{2}$ , the result follows by repeating the procedure described in this case.

Case 2. Suppose  $3 \le p < s$  and  $p \not\equiv s \pmod{2}$ . As above, assume p + q = s + 2. Then s is odd,  $p \geqslant 4$  is even and q is odd by our hypotheses.

Let L' be the leave of the packing  $\mathcal{P}'$  obtained from  $\mathcal{P}$  by applying the  $(x_2, y_q)$ -switch with origin  $x_3$  (note that  $\mu_L(x_3y_q)=0$  for otherwise L contains an s-cycle). If the terminus of the switch is not  $y_{q-1}$  then L' contains an s-cycle which completes the proof. Otherwise, the terminus of the switch is  $y_{q-1}$  and L' contains a (q+2, p-2)-lasso. Note that  $q+2 \leq s$  (because p+q=s+2 and  $p \geq 4$ ) and  $q+2 \equiv s \pmod{2}$ . If q+2=s then this completes the proof, otherwise we can proceed as in Case 1.

Case 3. Suppose  $3 \le s < p$ . Let L' be the leave of the packing  $\mathcal{P}'$  obtained from  $\mathcal{P}$  by applying the  $(x_{p-s+1}, y_1)$ -switch with origin  $x_{p-s+2}$  (note that  $\mu_L(x_{p-s+2}y_1) = 0$  for otherwise L contains an s-cycle). If the terminus of the switch is not  $x_p$  then L' contains an s-cycle which completes the proof. Otherwise, L' contains a (p-s+2, q+s-2)-lasso. By repeating this process we obtain an (M)-packing of  $\lambda K_v$  whose leave contains a (p', p+q-p')-lasso such that  $2 \le p' \le s$  and  $p' \equiv p \pmod{(s+2)}$ . If p' = s then this completes the proof, otherwise we can proceed as in Case 1 or Case 2.

**Lemma 4.** Let v, s and  $\lambda$  be positive integers with  $s \geq 3$ , and let M be a list of integers. Suppose there exists an (M)-packing of  $\lambda K_v$  whose leave L has a component H containing an (s+1)-cycle with a chord. Then there exists an (M)-packing of  $\lambda K_v$  with a leave L' such that  $E(L') = (E(L) \setminus E(H)) \cup E(H')$ , where H' is a graph with V(H') = V(H) and |E(H')| = |E(H)| which contains an (s,1)-lasso. Furthermore,  $\deg_{H'}(x) \geq \deg_H(x)$  for each vertex x in the scycle of this lasso.

**Proof.** Let  $(x_1, \ldots, x_{s+1})$  be an (s+1)-cycle in H with chord  $x_1x_e$  for some  $e \in \{2, 3, \ldots, s-1\}$  (note that L is not necessarily a simple graph). If H contains an (s, 1)-lasso then we are finished immediately, so suppose otherwise. If e = 2, then perform the  $(x_3, x_2)$ -switch with origin  $x_4$  (note that  $\mu_L(x_2x_4) = 0$  because H contains no (s, 1)-lasso). The leave of the resulting packing contains the (s, 1)-lasso  $(x_4, \ldots, x_{s+1}, x_1, x_2)[x_2, x_3]$ , and  $\deg_{H'}(x_i) \geqslant \deg_H(x_i)$  for  $i \in \{1, \ldots, s+1\} \setminus \{3\}$ . If e = 3, then H contains an (s, 1)-lasso which completes the proof.

So suppose  $e \ge 4$  and let  $\mathcal{P}^*$  be the packing with leave  $L^*$  obtained from  $\mathcal{P}$  by applying the  $(x_{e-1}, x_e)$ -switch with origin  $x_{e-2}$  (note that  $\mu_L(x_{e-2}x_e) = 0$  for otherwise L contains an (s, 1)-lasso). If the terminus of the switch is not  $x_{e+1}$  then  $E(L^*) = (E(L) \setminus E(H)) \cup E(H^*)$ , where  $H^*$  is a graph with  $V(H^*) = V(H)$  and  $|E(H^*)| = |E(H)|$  which contains the (s, 1)-lasso  $(x_{e+1}, \ldots, x_{s+1}, x_1, \ldots, x_{e-2}, x_e)[x_e, x_{e-1}]$ . Also note that  $\deg_{H^*}(x_e) \ge \deg_H(x_e)$  and  $\deg_{H^*}(x_i) = \deg_H(x_i)$  for  $i \in \{1, \ldots, s+1\} \setminus \{e, e-1\}$ . Otherwise, the terminus of the

switch is  $x_{e+1}$  and  $E(L^*) = (E(L) \setminus E(H)) \cup E(H^*)$ , where  $H^*$  is a graph with  $V(H^*) = V(H)$  and  $|E(H^*)| = |E(H)|$  which contains an (s+1)-cycle  $(x_1^*, \ldots, x_{s+1}^*)$  with chord  $x_1^* x_{e-1}^*$ . Furthermore, the degree of each vertex in this (s+1)-cycle remains unchanged in H'. The result follows by repeating this process.

## 3 Main result

This section contains the proof of Theorem 1. We first use Theorem 5 (stated below) to prove Lemma 6 which shows the necessity of the conditions in Theorem 1. The sufficiency of these conditions is then established for  $\lambda$  odd and  $\lambda$  even in Lemmas 7 and 8 respectively. Lemmas 7 and 8 rely on using Lemmas 3 and 4 to modify cycle packings of  $\lambda K_v$  obtained via Theorem 5.

**Theorem 5** ([5]). Let  $m_1, m_2, \ldots, m_{\tau}$  be a nondecreasing list of integers and let  $\lambda$  and v be positive integers. There is an  $(m_1, m_2, \ldots, m_{\tau})$ -decomposition of  $\lambda K_v$  if and only if

- $\lambda(v-1)$  is even;
- $2 \leqslant m_1 \leqslant m_2, \ldots, m_\tau \leqslant v;$
- $m_1 + m_2 + \cdots + m_{\tau} = \lambda \binom{v}{2}$ ;
- $m_{\tau} + \tau 2 \leqslant \frac{\lambda}{2} \binom{v}{2}$  when  $\lambda$  is even; and
- $\sum_{m_i=2} m_i \leqslant (\lambda-1)\binom{v}{2}$  when  $\lambda$  is odd.

There is an  $(m_1, m_2, ..., m_{\tau})$ -decomposition of  $\lambda K_v - I$ , where I is a 1-factor in  $\lambda K_v$ , if and only if

- $\lambda(v-1)$  is odd:
- $2 \leqslant m_1 \leqslant m_2, \ldots, m_{\tau} \leqslant v$ ;
- $m_1 + m_2 + \cdots + m_{\tau} = \lambda \binom{v}{2} \frac{v}{2}$ ; and
- $\sum_{m_i=2} m_i \leqslant (\lambda-1)\binom{v}{2}$ .

The necessity of conditions Theorem 1(i)–(iv) follows from Theorem 5 as we now show.

**Lemma 6.** Let  $m_1, m_2, \ldots, m_{\tau}$  be a nondecreasing list of integers and let  $\lambda$  and v be positive integers. If there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$  then

- (i)  $2 \leqslant m_1 \leqslant m_2, \ldots, m_\tau \leqslant v;$
- (ii)  $m_1 + m_2 + \cdots + m_{\tau} = \lambda {v \choose 2} \delta$ , where  $\delta$  is a nonnegative integer such that  $\delta \neq 1$  when  $\lambda(v-1)$  is even,  $\delta \neq 2$  when  $\lambda = 1$ , and  $\delta \geqslant \frac{v}{2}$  when  $\lambda(v-1)$  is odd;
- (iii)  $\sum_{m_i=2} m_i \leqslant \begin{cases} (\lambda 1)\binom{v}{2} 2 & \text{if } \lambda \text{ and } v \text{ are odd and } \delta = 2, \\ (\lambda 1)\binom{v}{2} & \text{if } \lambda \text{ is odd; and} \end{cases}$
- (iv)  $m_{\tau} \leqslant \begin{cases} \frac{\lambda}{2} \binom{v}{2} \tau + 2 & \text{if } \lambda \text{ is even and } \delta = 0, \\ \frac{\lambda}{2} \binom{v}{2} \tau + 1 & \text{if } \lambda \text{ is even and } 2 \leqslant \delta < m_{\tau}. \end{cases}$

**Proof.** Suppose there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing  $\mathcal{P}$  of  $\lambda K_v$  with leave L. Condition (i) is obvious. The degree of each vertex in  $\lambda K_v$  is  $\lambda(v-1)$ , so if  $\lambda(v-1)$  is even then L is an even multigraph and if  $\lambda(v-1)$  is odd then L is an odd multigraph. Hence (ii) follows because an even graph cannot have a single edge, an even simple graph cannot have two edges, and an odd graph on v vertices has at least  $\frac{v}{2}$  edges. To see that condition (iii) holds, note that there are at most  $\lfloor \frac{\lambda}{2} \rfloor \binom{v}{2}$  edge-disjoint 2-cycles in  $\lambda K_v$ . Furthermore, note that if  $\lambda$  and v are both odd and  $\delta = 2$  then L is a 2-cycle (because L is an even multigraph and has two edges). If  $\lambda$  is even and  $\delta = 0$  then (iv) follows directly from Theorem 5, so suppose  $\lambda$  is even and  $2 \leq \delta < m_{\tau}$ . Then L contains at least one cycle so there exists an  $(m_1, m_2, \ldots, m_{\tau}, M)$ -decomposition of  $\lambda K_v$  for some list M containing at least one entry. So (iv) follows from Theorem 5.

It remains to prove the sufficiency of Theorem 1(i)–(iv) for the existence of cycle packings of  $\lambda K_v$ .

**Lemma 7.** Let  $m_1, m_2, \ldots, m_{\tau}$  be a nondecreasing list of integers and let  $\lambda$  and v be positive integers with  $\lambda$  odd. Then there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$  if and only if

- (i)  $2 \leq m_1 \leq m_2, \dots, m_{\tau} \leq v$ ;
- (ii)  $m_1 + m_2 + \cdots + m_{\tau} = \lambda {v \choose 2} \delta$ , where  $\delta$  is a nonnegative integer such that  $\delta \neq 1$ ,  $(\lambda, \delta) \neq (1, 2)$ , and if v is even then  $\delta \geqslant \frac{v}{2}$ ; and

(iii) 
$$\sum_{m_i=2} m_i \leqslant \begin{cases} (\lambda - 1)\binom{v}{2} - 2 & \text{if } v \text{ is odd and } \delta = 2, \\ (\lambda - 1)\binom{v}{2} & \text{otherwise.} \end{cases}$$

**Proof.** If there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing  $\mathcal{P}$  of  $\lambda K_v$ , then conditions (i)–(iii) hold by Lemma 6. So it remains to show that if  $\lambda$ , v and  $m_1, m_2, \ldots, m_{\tau}$  satisfy (i)–(iii), then there is an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$ .

Let  $\varepsilon = \delta$  if v is odd, and  $\varepsilon = \delta - \frac{v}{2}$  if v is even. If  $\varepsilon = 0$  then the result follows by Theorem 5. If v = 2, then  $\varepsilon$  is even by (i) and (ii) and there exists a 2-cycle decomposition of  $\lambda K_2 - I$ , where I is a 1-factor of  $\lambda K_2$ , so the result follows. So suppose  $\varepsilon \geqslant 1$  and  $v \geqslant 3$ , and note that if v is odd then  $\varepsilon \neq 1$  and  $(\lambda, \varepsilon) \neq (1, 2)$ .

Case 1. Suppose v is odd or  $\varepsilon \geqslant 3$ . Note that if v is odd and  $\varepsilon = 2$  then  $2 + \sum_{m_i=2} m_i \leqslant (\lambda - 1) {v \choose 2}$  by (iii).

We show that there exists a list N such that  $2 \leq n \leq v$  for all  $n \in N$ ,  $\sum N = \varepsilon$  and  $\sum_{n \in N, n=2} n + \sum_{m_j=2} m_j \leq (\lambda - 1) {v \choose 2}$ . If this list exists, then by Theorem 5 there exists an  $(m_1, m_2, \ldots, m_\tau, N)$ -decomposition  $\mathcal{D}$  of  $\lambda K_v$  (if v is odd) or  $\lambda K_v - I$  (if v is even), where I is a 1-factor of  $\lambda K_v$ . We obtain the required packing by removing cycles of lengths N from  $\mathcal{D}$ .

We first consider v=3. If v=3 and  $\varepsilon$  is even, then  $m_i=3$  for some  $i\in\{1,\ldots,\tau\}$  by (i) and (ii). Then  $\varepsilon+\sum_{m_i=2}m_i\leqslant (\lambda-1)\binom{v}{2}$  by (ii) and we take  $N=(2^{\varepsilon/2})$ . If v=3 and  $\varepsilon$  is odd then  $\varepsilon-3+\sum_{m_i=2}m_i\leqslant (\lambda-1)\binom{v}{2}$  by (ii) and we take  $N=(2^{(\varepsilon-3)/2},3)$ . In each of these cases we can see that there exists an  $(m_1,m_2,\ldots,m_\tau,N)$ -decomposition of  $\lambda K_v$  since the assumptions of Theorem 5 are satisfied by (i)–(iii).

Now assume  $v \ge 4$  and let q and r be nonnegative integers such that  $\varepsilon = vq + r$  and  $0 \le r < v$ . If q = 0 or  $r \not\in \{1,2\}$  then we take  $N = (r, v^q)$ . If  $q \ge 1$  and  $r \in \{1,2\}$  then  $N = (3, v - 3 + r, v^{q-1})$  (note that either  $v - 3 + r \ge 3$ , or v = 4 and r = 1). If  $\varepsilon = 2$  or (v, r) = (4, 1), then N contains exactly one entry equal to 2 and otherwise  $n \ge 3$  for all  $n \in N$ . By (iii) and the assumptions of this case, if  $\varepsilon = 2$  then  $2 + \sum_{m_i=2} m_i \le (\lambda - 1)\binom{v}{2}$ .

Further, if v=4 and  $\varepsilon=4q+1$  for some  $q\geqslant 1$  then (i) and (ii) imply that  $m_i=3$  for some  $i\in\{1,\ldots,\tau\}$  so again  $2+\sum_{m_i=2}m_i\leqslant (\lambda-1)\binom{v}{2}$ . We can therefore see that there exists an  $(m_1,m_2,\ldots,m_\tau,N)$ -decomposition of  $\lambda K_v$  (or  $\lambda K_v-I$ ) since the assumptions of Theorem 5 are satisfied by (i)–(iii) and the fact that  $\sum N=\varepsilon$ .

Case 2. Suppose v is even and  $\varepsilon \in \{1, 2\}$ . Let  $M = m_1, m_2, \ldots, m_{\tau}$  and let m be the least odd entry in M if M contains an odd entry, otherwise let m be the least entry in M such that  $m \ge 4$  (such an entry exists by (iii)). Note that  $v \ge 4$  and if  $\varepsilon = 1$  then it follows from (ii) that M contains an odd entry and m is odd.

Case 2a. Suppose  $m + \varepsilon \leq v$ . By Theorem 5 there exists an  $(M \setminus (m), m + \varepsilon)$ -decomposition  $\mathcal{D}$  of  $\lambda K_v - I$ , where I is a 1-factor of  $\lambda K_v$ . Let  $\mathcal{P}$  be the  $(M \setminus (m))$ -packing of  $\lambda K_v$  that is obtained by removing an  $(m + \varepsilon)$ -cycle from  $\mathcal{D}$ . Let L be the leave of  $\mathcal{P}$  and note that L consists of an  $(m + \varepsilon)$ -cycle and the 1-factor I.

If L contains an  $(m + \varepsilon, 1)$ -lasso then we apply Lemma 3 to  $\mathcal{P}$  with s = m to complete the proof. The assumptions of Lemma 3 are satisfied because  $\varepsilon + 1 \ge 2$ , and if m is even then M contains no odd entries so  $\varepsilon = 2$  by (ii).

So suppose L does not contain an  $(m + \varepsilon, 1)$ -lasso. Then  $m + \varepsilon$  is even and L contains a component H such that H is the union of an  $(m + \varepsilon)$ -cycle and a 1-factor on V(H). We apply Lemma 4 to  $\mathcal{P}$  with  $s = m + \varepsilon - 1$  to obtain an  $(M \setminus (m))$ -packing  $\mathcal{P}'$  of  $\lambda K_v$  whose leave L' contains a component H' on  $m + \varepsilon$  vertices that has  $\frac{3}{2}(m + \varepsilon)$  edges and contains an  $(m + \varepsilon - 1, 1)$ -lasso. If  $\varepsilon = 1$  then we can add the m-cycle of this lasso to  $\mathcal{P}'$  to complete the proof. Otherwise  $\varepsilon = 2$  and H' contains an (m + 1)-cycle with a chord because  $m \ge 3$  and any vertex in this cycle has degree at least 3 (note that  $\deg_H(x) = 3$  for each  $x \in V(H)$ ). Then we can apply Lemma 4 with s = m to  $\mathcal{P}'$  to obtain an  $(M \setminus (m))$ -packing  $\mathcal{P}''$  of  $\lambda K_v$  whose leave contains an (m, 1)-lasso. We add the m-cycle of this lasso to  $\mathcal{P}''$  to complete the proof.

Case 2b. Suppose  $m + \varepsilon > v$ . Then  $m \ge v - 1$  and  $\varepsilon = 2$  (note that  $\varepsilon$  is even if m = v).

If m = v then  $m_i \in \{2, v\}$  for all  $i \in \{1, ..., \tau\}$ , so  $\lambda {v \choose 2} - \frac{v}{2} \equiv 2 + \sum_{m_i=2} m_i \pmod{v}$  by (ii) and hence  $2 + \sum_{m_i=2} m_i \leqslant (\lambda - 1) {v \choose 2}$  by (iii). Then by Theorem 5 there exists an (M, 2)-decomposition  $\mathcal{D}$  of  $\lambda K_v - I$ . We remove a 2-cycle from  $\mathcal{D}$  to complete the proof.

So suppose that m = v - 1. Since  $\varepsilon$  is even, M contains an even number of odd entries, so at least two entries of M are equal to v - 1. Let  $\mathcal{D}_0$  be an  $(M \setminus ((v-1)^2), v^2)$ -decomposition of  $\lambda K_v - I$  which exists by Theorem 5. Let  $\mathcal{P}_0$  be the  $(M \setminus ((v-1)^2), v)$ -packing of  $\lambda K_v$  formed by removing a v-cycle from  $\mathcal{D}_0$ . The leave  $L_0$  of  $\mathcal{P}_0$  is the union of a v-cycle and the 1-factor I. Let  $\mathcal{P}_1$  be the packing obtained by applying Lemma 4 to  $\mathcal{P}_0$  with s = v - 1. Then the leave of  $\mathcal{P}_1$  contains a (v - 1, 1)-lasso. We add the (v - 1)-cycle of this lasso to  $\mathcal{P}_1$  and remove a v-cycle to obtain an  $(M \setminus (v - 1))$ -packing  $\mathcal{P}_2$  of  $\lambda K_v$ . The leave of  $\mathcal{P}_2$  has size  $3\frac{v}{2} + 1$ .

By applying Lemma 4 to  $\mathcal{P}_2$  with s = v - 1 we obtain an  $(M \setminus (v - 1))$ -packing  $\mathcal{P}_3$  of  $\lambda K_v$  whose leave contains a (v - 1, 1)-lasso. We add the (v - 1)-cycle of this lasso to  $\mathcal{P}_3$  to complete the proof.

**Lemma 8.** Let  $m_1, m_2, \ldots, m_{\tau}$  be a nondecreasing list of integers and let  $\lambda$  and v be positive integers with  $\lambda$  even. Then there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$  if and only if

- (i)  $2 \leqslant m_1 \leqslant m_2, \ldots, m_{\tau} \leqslant v$ ;
- (ii)  $m_1 + m_2 + \cdots + m_{\tau} = \lambda \binom{v}{2} \delta$ , where  $\delta$  is a nonnegative integer such that  $\delta \neq 1$ ; and

(iii) 
$$m_{\tau} \leqslant \begin{cases} \frac{\lambda}{2} \binom{v}{2} - \tau + 2 & \text{if } \delta = 0, \\ \frac{\lambda}{2} \binom{v}{2} - \tau + 1 & \text{if } 2 \leqslant \delta < m_{\tau}. \end{cases}$$

**Proof.** If there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing  $\mathcal{P}$  of  $\lambda K_v$  with leave L, then conditions (i)-(iii) hold by Lemma 6. So it remains to show that if  $\lambda$ , v and  $m_1, m_2, \ldots, m_{\tau}$  satisfy (i)-(iii), then there exists an  $(m_1, m_2, \ldots, m_{\tau})$ -packing of  $\lambda K_v$ . If  $\delta = 0$  then the result follows immediately from Theorem 5, so suppose  $\delta \geq 2$ .

Let

$$N = \begin{cases} (\delta) & \text{if } 2 \leqslant \delta < m_{\tau}, \\ (2^{(\delta - m_{\tau})/2}, m_{\tau}) & \text{if } \delta \geqslant m_{\tau} \text{ and } \delta \equiv m_{\tau} \pmod{2}, \\ (2^{(\delta - m_{\tau} + 1)/2}, m_{\tau} - 1) & \text{if } \delta \geqslant m_{\tau} \text{ and } \delta \not\equiv m_{\tau} \pmod{2}. \end{cases}$$

Note that in each case  $\sum N = \delta$ . We show that there exists an  $(m_1, m_2, \ldots, m_{\tau}, N)$ -decomposition  $\mathcal{D}$  of  $\lambda K_v$  because the assumptions of Theorem 5 are satisfied by (i)–(iii) and the definition of N. The required packing is then obtained by removing cycles of lengths N from  $\mathcal{D}$ .

Let s be the number of entries in N. Let  $M = m_1, m_2, \ldots, m_{\tau}$ . First observe that  $\sum M + \sum N = \lambda \binom{v}{2}$  by (ii) and since  $\sum N = \delta$ . By (i) and the definition of N it also holds that  $2 \leqslant n \leqslant m_{\tau} \leqslant v$  for all  $n \in N$ . If  $2 \leqslant \delta < m_{\tau}$ , then  $m_{\tau} \leqslant \frac{\lambda}{2} \binom{v}{2} - \tau - s + 2$  by (iii) and since s = 1. If  $\delta \geqslant m_{\tau}$ , then because  $\sum M \geqslant m_{\tau} + 2(\tau - 1)$  and  $\sum N \geqslant m_{\tau} - 1 + 2(s - 1)$ , it follows that

$$\frac{\lambda}{2} \binom{v}{2} - \tau - s + 2 = \frac{1}{2} (\sum M + \sum N) - \tau - s + 2 
\geqslant \frac{1}{2} (m_{\tau} + 2(\tau - 1) + m_{\tau} - 1 + 2(s - 1)) - \tau - s + 2 
= m_{\tau} - \frac{1}{2}.$$

Therefore  $\max(N, M) = m_{\tau} \leq \frac{\lambda}{2} \binom{v}{2} - \tau - s + 2$  because  $\frac{\lambda}{2} \binom{v}{2} - \tau - s + 2$  is an integer. So by Theorem 5 we can see that there exists an (M, N)-decomposition of  $\lambda K_v$  which completes the proof.

## Acknowledgements

The author was supported by a Monash University Faculty of Science Postgraduate Publication Award.

#### References

- [1] B. Alspach and H. Gavlas. Cycle decompositions of  $K_n$  and  $K_n I$ . J. Combin. Theory Ser. B, 81 (2001), 77–99.
- [2] P. N. Balister. On the Alspach conjecture. Combin. Probab. Comput., 10 (2001), 95–125.
- [3] D. E. Bryant and D. Horsley. Decompositions of complete graphs into long cycles. *Bull. Lond. Math. Soc.*, 41 (2009), 927–934.
- [4] D. E. Bryant, D. Horsley, B. M. Maenhaut, and B. R. Smith. Cycle decompositions of complete multigraphs. *J. Combin. Des.*, 19 (2011), 42–69.
- [5] D. E. Bryant, D. Horsley, B. M. Maenhaut, and B. R. Smith. Decompositions of complete multigraphs into cycles of varying lengths. (preprint), (2015) arxiv:1508.00645 [math.CO].
- [6] D. E. Bryant, D. Horsley, and W. Pettersson. Cycle decompositions V: Complete graphs into cycles of arbitrary lengths. *Proc. Lond. Math. Soc.* (3), 108 (2014), 1153–1192.

- [7] D. E. Bryant and C. A. Rodger. Cycle decompositions. *The CRC Handbook of Combinatorial Designs*, C. J. Colbourn and J. H. Dinitz (editors), CRC Press, Boca Raton, 2nd edition (2007), pp. 373–382.
- [8] D. E. Bryant and Daniel Horsley. Packing cycles in complete graphs. *J. Combin. Theory Ser. B*, 98 (2008), 1014–1037.
- [9] S. I. El-Zanati. Maximum packings with odd cycles. Discrete Math., 131 (1994), 91–97.
- [10] D. Horsley. Maximum packings of the complete graph with uniform length cycles. *J. Graph Theory*, 68 (2011), 1–7.
- [11] M. Śajna. Cycle decompositions III. Complete graphs and fixed length cycles. *J. Combin. Des.*, 10 (2002), 27–78.
- [12] B. R. Smith. Cycle decompositions of complete multigraphs. *J. Combin. Des.*, 18 (2010), 85–93.