

A Baer-Krull theorem for quasi-ordered groups

Salma Kuhlmann, Gabriel Lehericy

June 22, 2021

Abstract

We give group analogs of two important theorems of real algebra concerning convex valuations, one of which is the Baer-Krull theorem. We do this by using quasi-orders, which gives a uniform approach to valued and ordered groups. We also recover the classical Baer-Krull theorem from its group analog.

Introduction

The theories of field ordering and field valuation present some strong similarities but are classically treated as separate subjects. However, in [3], Fakhruddin found a way of unifying these two theories by using quasi-orders. He defined a quasi-ordered field as a field K endowed with a total quasi-order (q.o) \lesssim satisfying the following axioms:

$$(Q_1) \quad \forall x(x \sim 0 \Rightarrow x = 0)$$

$$(Q_2) \quad \forall x, y, z(x \lesssim y \wedge z \Rightarrow x + z \lesssim y + z)$$

$$(Q_3) \quad \forall x, y, z(x \lesssim y \wedge 0 \lesssim z) \Rightarrow xz \lesssim yz$$

He then showed that valued and ordered fields are particular instances of q.o fields, and even showed that they are the only ones, so that the theory of q.o fields is very convenient to unify the theories of valued fields with the theory of ordered fields. Two different notions of quasi-ordered groups have been studied in [7] and [8], both of which can be seen as a generalization of ordered and valued groups. The group analog of Fakhruddin's theory was studied in [7].

The notion of compatibility between a valuation and an order plays an important role in the theory of valued fields. One can give several characterizations of compatibility of a valuation v with respect to an order. The notion of coarsening of a valuation also plays an important role in valuation theory and is in some sense analogous to the notion of compatibility. In [6], the authors used Fakhruddin's theory to give a uniform approach to the notions of compatible valuations with respect to an order and the notion of coarsening of a valuation. We recall their result:

Theorem 0.1

Let (K, \lesssim) be a quasi-ordered field and v a valuation on K . The following are equivalent:

- 1) v is compatible with \lesssim
- 2) R_v is \lesssim -convex in K

- 3) I_v is \preceq -convex in K
- 4) $I_v < 1$
- 5) \preceq induces a q.o on the residue field R_v/I_v .

In the special case where \preceq is an order, Theorem 0.1 gives conditions for a valuation v to be compatible with \leq . If \preceq comes from a valuation w then Theorem 0.1 gives conditions for v to be a coarsening of w .

An important result concerning valuation-compatible orders is the Baer-Krull Theorem. Given a valued field (K, v) , the Baer-Krull Theorem describes the set of v -compatible orderings of K modulo the set of orderings of the residue field. We recall the Baer-Krull Theorem as it appears in [2]:

Theorem 0.2 (Baer-Krull)

Let $v : K \rightarrow G \cup \{\infty\}$ be a valuation and let $(\pi_i)_{i \in I}$ be a family of elements of K such that $(v(\pi_i) + 2G)_{i \in I}$ is an \mathbb{F}_2 -basis of $G/2G$. Let $\mathcal{O}(K)$ be the set of field orderings on K with which v is compatible and $\mathcal{O}(Kv)$ the set of orders of the residue field Kv . There is a bijection: $\phi : \mathcal{O}(K) \longleftrightarrow \{-1, 1\}^I \times \mathcal{O}(Kv)$ defined as follows: $\phi(\leq) = (\eta, \leq_{Kv})$, where $\eta(i) = 1$ if and only if $0 \leq \pi_i$ and \leq_{Kv} is the order induced by \leq on the residue field.

The notion of valuation also exists for groups: a valuation on a group G is a map v from G to a totally ordered set $(\Gamma \cup \{\infty\}, \leq)$, where ∞ denotes the maximum of $\Gamma \cup \{\infty\}$, so that the following is satisfied: $v(g) = \infty \Leftrightarrow g = 0$, $v(g - h) \geq \min(v(g), v(h))$ and $v(g) \leq v(h) \Rightarrow v(x + g - x) \leq v(x + h - x)$ for all $g, h, x \in G$. Valued groups have been studied in [10] and [5]; however, the notions of compatible valuation on groups and of coarsening of group valuations still haven't been developed. The goal of this paper is to introduce these notions using the theories of quasi-ordered groups developed in [7] and [8] and to find group analogs of Theorems 0.1 and 0.2. In Section 1 we give some preliminaries on quasi-orders and prove the lemmas which we will need in the next sections. In Section 2 we state an analog of Theorem 0.1 in the framework of the compatible quasi-ordered groups defined in [7], but conclude that there is no analog of the Baer-Krull Theorem in that case. In Section 3 we consider the C-quasi-ordered groups defined in [8] and prove analogs of theorems 0.1 and 0.2 in that setting (Theorems 3.3 and 3.4). In Section 4 we explain how we can recover the classical Baer-Krull Theorem from its group analog.

1 Quasi-orders on groups

We recall that a quasi-order is a binary relation which is reflexive and transitive. For us, a q.o group is just a group endowed with a q.o with no extra assumption. A q.o \preceq naturally induces an equivalence relation which we denote by \sim and defined as $a \sim b \Leftrightarrow a \preceq b \preceq a$. We denote by $cl(a)$ the class of a for this equivalence relation. For any $a, b \in A$, the notation $a \preceq b$ means $a \preceq b \wedge a \not\prec b$. *Every q.o and every order considered in this paper*

is total. If (A, \lesssim) is a quasi-ordered set and $B \subseteq A$, we say that B is **\lesssim -convex** (in A) if for any $a \in A$ and $b, c \in B$, $b \lesssim a \lesssim c$ implies $a \in B$. We say that B is an **initial segment** (respectively, a **final segment**) of A if for any $a \in A$ and $b \in B$, $a \lesssim b$ (respectively, $b \lesssim a$) implies $a \in B$. If $v : G \rightarrow \Gamma \cup \{\infty\}$ is a valuation on a group G , then for any $\gamma \in \Gamma$ we denote by G^γ the set $\{g \in G \mid v(g) \geq \gamma\}$ and by G_γ the set $\{g \in G \mid v(g) > \gamma\}$. Note that if v is a valuation on G , then the relation $g \lesssim h \Leftrightarrow v(g) \geq v(h)$ is a total q.o, which we call the q.o induced by v . We say that a q.o is **valuational** if it is induced by a valuation. If (G, \lesssim) is a q.o group, we say that $g \in G$ is **v-type** if $g \sim -g$ and **o-type** if $g \not\sim -g \vee g = 0$. The choice of this terminology will be made clear later (see Proposition 2.1). We say that **the q.o \lesssim is o-type** (respectively **v-type**) if every element of (G, \lesssim) is o-type (respectively v-type). In order to have a Baer-Krull theorem, we need to define notions of q.o induced on a quotient as well as notions of “lifting” q.o’s from quotients. Given a quasi-order \lesssim on G and a subgroup H , we say that \lesssim induces a q.o on G/H if the following relation \lesssim defined on G/H is transitive: $g_1 + H \lesssim g_2 + H \Leftrightarrow \exists h_1, h_2 \in H g_1 + h_1 \lesssim g_2 + h_2$. In that case we call this relation the **q.o induced by \lesssim on G/H** . Now assume we are given a valuation $v : G \rightarrow \Gamma$ and a family $(\lesssim_\gamma)_{\gamma \in \Gamma}$ of q.o’s, each defined on a quotient G^γ/G_γ . We define **the lifting of (\lesssim_γ) to G** as the q.o given by the formula: $g \lesssim h \Leftrightarrow g + G_\gamma \lesssim_\gamma h + G_\gamma$ where $\gamma = \min(v(g), v(h))$. It is the only q.o such that for any γ , \lesssim induces the q.o \lesssim_γ on G^γ/G_γ . Finally, if (G, \lesssim) is a q.o group and v a valuation on G , we say that v is **compatible with \lesssim** , or that \lesssim is a **v -compatible q.o**, if for any $g, h \in G$, $0 \lesssim g \lesssim h$ implies $v(g) \geq v(h)$.

Towards an analog of Theorem 0.1, we will need the following lemmas. We fix a group G , an arbitrary total q.o \lesssim and a valuation $v : G \rightarrow \Gamma \cup \{\infty\}$.

Lemma 1.1

The following statements are equivalent:

- (1) $\forall \gamma \in \Gamma, G^\gamma$ is \lesssim -convex.
- (2) $\forall \gamma \in \Gamma, G_\gamma$ is \lesssim -convex.

Proof. (1) \Rightarrow (2): let $\gamma \in \Gamma, g, h \in G_\gamma$ and $f \in G$ with $h \lesssim f \lesssim g$. Set $\delta := \min(v(g), v(h))$. We have $\delta > \gamma$ so $G^\delta \subseteq G_\gamma$. Moreover, $g, h \in G^\delta$, which by \lesssim -convexity of G^δ implies $f \in G^\delta \subseteq G_\gamma$. (2) \Rightarrow (1): Assume $g, h \in G^\gamma$. If $f \notin G^\gamma$ then $v(f) < \gamma \leq v(g), v(h)$ hence $g, h \in G_{v(f)}$, which by assumption means we cannot have $h \lesssim f \lesssim g$. \square

Lemma 1.2

v is compatible with \lesssim if and only if $(G_\gamma)^{\gtrsim 0}$ is \lesssim -convex for all $\gamma \in \Gamma$.

Proof. If v is compatible with \lesssim , then for any $0 \lesssim g \lesssim h \in G_\gamma$ we have $v(g) \geq v(h) > \gamma$ so $g \in G_\gamma$. Conversely, assume each $(G_\gamma)^{\gtrsim 0}$ is \lesssim -convex. If $0 \lesssim g \lesssim h$ and $v(h) > v(g)$, then $h \in G_{v(g)}$ by definition and $g \in G_{v(g)}$ by \lesssim -convexity of $(G_\gamma)^{\gtrsim 0}$, which is a contradiction. Thus, $0 \lesssim g \lesssim h \Rightarrow v(g) \geq v(h)$ and v is compatible. \square

Lemma 1.3

Let \lesssim be a q.o such that:

$$(*) \left\{ \begin{array}{l} cl(0) = \{0\} \\ g \lesssim h \not\sim -f \Rightarrow g + f \lesssim h + f \text{ for any } g, h, f \in G. \end{array} \right.$$

and let H be a subgroup of G . Then H is \lesssim -convex if and only if \lesssim induces a total q.o on G/H such that $cl(0 + H) = \{0 + H\}$. Moreover, if H is \lesssim -convex, then for every $g \in G \setminus H$, $g \lesssim -g$ if and only if $g + H \lesssim -g + H$; in particular, g is v-type (resp. o-type) if and only if $g + H$ is v-type (resp. o-type).

Proof. Assume H is \lesssim -convex. Let f, g, h with $f + H \lesssim g + H$ and $g + H \lesssim h + H$. There are $a, b, b', c \in H$ such that $f + a \lesssim g + b$ and $g + b' \lesssim h + c$. Assume first that $g, h \in H$. If $h \lesssim f + a$ then by \lesssim -convexity of H we have $f \in H$ and $f \lesssim h + f - h$. Assume $h \notin H$, so $h + c \notin H$. By \lesssim -convexity, $h + c \not\sim b - b'$, so by assumption $g + b' \lesssim h + c$ implies $f + a \lesssim g + b = g + b' + b - b' \lesssim h + c + b - b'$. If $g \notin H$ then the same reasoning gives $f + a + b' - b \lesssim g + b' \lesssim h + c$. In any case, we have $f + H \lesssim h + H$, so the relation \lesssim on G/H is transitive. This proves that \lesssim induces a q.o on G/H . If $g + H \sim H$ then there are $a, b, c \in H$ with $a \lesssim g + b \lesssim c$ which implies $g \in H$, so $cl(H) = \{H\}$. Conversely, assume \lesssim induces a q.o on G/H such that $cl(0 + H) = \{0 + H\}$. If $g, h \in H$ and $f \in G$ are such that $g \lesssim f \lesssim h$ then $0 = g + H \lesssim f + H \lesssim h + H = 0$ hence by assumption $f \in H$.

For the second statement, let $g \in G \setminus H$. Assume $g \not\lesssim -g$. Since \lesssim is total, we have $-g \lesssim g$. Note that $0 \lesssim g$: for if $0 \not\lesssim g$, then since \lesssim is total we have $g \lesssim 0 \not\sim g = -(-g)$ and (*) implies $0 \lesssim -g$; so since $-g \lesssim g$ we must have $0 \lesssim g$. Because $g \not\lesssim -g$, we have in particular $g \not\sim -g$, and so by (*),

$$g = 0 + g \lesssim g + g = 2g.$$

Thus $0 \lesssim g \lesssim 2g$. If $g + H \lesssim -g + H$, then there are $a, b \in H$ with $g + a \lesssim (-g) + b$. Since H is \lesssim -convex, and $a \in H$, and $-g + b \notin H$, we have $-(-a) = a \not\sim -g + b$, so by (*):

$$g \lesssim -g + b - a.$$

If $-g \sim -g + b - a$, then $g \lesssim -g + b - a \lesssim -g$, so we must have $-g \not\sim -g + b - a$, and by (*) again

$$2g \lesssim b - a.$$

Thus $0 \lesssim g \lesssim 2g \lesssim b - a$, and $g \in H$ because H is \lesssim -convex. This contradiction shows that $g + H \not\lesssim -g + H$ if $g \not\lesssim -g$. \square

Remark 1.4

The reader should pay attention to the fact that the second condition of (*) is not

equivalent to axiom (Q_2) of compatible q.o's given below: example 3.1(a) of [8] satisfies $(*)$ but not (Q_2) . However, it will later turn out that (Q_1) and (Q_2) imply $(*)$. The reason why we choose to state Lemma 1.3 with $(*)$ instead of (Q_1) and (Q_2) is the following: the class of q.o's satisfying $(*)$ contains both the class of compatible q.o's (see Section 2) and the class of C-q.o's (see Section 3), so that Lemma 1.3 can be applied to both cases.

2 Compatible q.o's

A first idea to achieve the goal stated in the introduction is to take inspiration from [3]. We say that a q.o \lesssim is compatible (with the group operation) if it satisfies:

$$(Q_1) \quad cl(0) = \{0\}$$

$$(Q_2) \quad \forall x, y, z (x \lesssim y \wedge z \Rightarrow x + z \lesssim y + z)$$

Group orders and valational q.o's are particular cases of compatible q.o's. More precisely, we proved the following in [7] (Proposition 2.8 and 2.15):

Proposition 2.1

Let \lesssim be a compatible q.o on an abelian group G . Then \lesssim is a group order if and only if every element is o-type, and \lesssim is valational if and only if every element is v-type.

Compatible q.o's have been studied in [7]. We proved there (Theorem 2.21) that if \lesssim is a compatible q.o on an abelian group G , then there is a subgroup G^o of G which is an initial segment of G such that \lesssim is a group order on G^o and \lesssim corresponds to a valuation outside of G^o . It is easy to show that compatible q.o's satisfy the condition $(*)$ of lemma 1.3 and that a subgroup H of a compatible q.o group (G, \lesssim) is \lesssim -convex if and only if $H^{\geq 0}$ is \lesssim -convex. Using the lemmas from section 1, we get:

Theorem 2.2

Let \lesssim be a compatible q.o on an abelian group G and $v : G \rightarrow \Gamma \cup \{\infty\}$ a valuation. The following statements are equivalent:

1. $\forall \gamma \in \Gamma, G^\gamma$ is \lesssim -convex.
2. $\forall \gamma \in \Gamma, G_\gamma$ is \lesssim -convex.
3. $\forall \gamma \in \Gamma, \lesssim$ induces a compatible q.o \lesssim_γ on G^γ/G_γ .
4. v is compatible with \lesssim .

Moreover, \lesssim is valational (respectively, an order) if and only if each \lesssim_γ is valational (respectively, an order).

Proof. All that remains to check is that the induced q.o on a quotient satisfies axiom (Q_2) ; for this we refer to the end of the proof of 2.11 in [7]. The last part of the theorem is a consequence of Proposition 2.1 and Lemma 1.3. \square

This deals with our first problem, i.e finding an analog of theorem 0.1 for groups. However, the class of compatible q.o's is not appropriate for a Baer-Krull theorem. This is due to the fact that the set of o-type elements of a compatible q.o must be an initial segment of G . If we lift an arbitrary family of compatible q.o's to G , then the set of o-type elements of G won't be an initial segment in general, which means that the class of compatible q.o's is not stable under lifting. In order to state an analog of the Baer-Krull theorem, we need to consider another class of q.o's.

3 C-q.o's and the Baer-Krull theorem

We recall that a C-relation is a ternary relation satisfying the universal closure of the following axioms: $C(x, y, z) \Rightarrow C(x, z, y) \wedge \neg C(y, x, z) \wedge (C(w, y, z) \vee C(x, w, z))$ and $x \neq y \Rightarrow C(x, y, y)$. A C-relation on a group $(G, +)$ is said to be **compatible** (with the group operation) if $C(f, g, h) \Rightarrow C(x + f + y, x + g + y, x + h + y)$ holds for every $f, g, h, x, y \in G$. See [1] for more information on C-relations. A C-q.o on a group G is a q.o induced by a compatible C-relation, i.e \lesssim is a C-q.o if and only if there exists a compatible C-relation C on G such that $g \lesssim h \Leftrightarrow \neg C(g, h, 0)$ for every $g, h \in G$. A structure theorem for groups endowed with a C-q.o was given in [8] (Theorem 3.41). The original purpose of C-q.o's is to study C-groups. This is possible thanks to a bijective correspondence between the class of compatible C-relations on a group and the class of C-q.o's. In particular, C-q.o's can be seen as a generalization of both orders and valuations on groups.

Proposition 3.1

Valuational q.o's are C-q.o's. Moreover, a C-q.o \lesssim is valuational if and only if every element of (G, \lesssim) is v-type.

In [4], the author showed that group orders are characterized by their positive cones: if we are given a subset P of G such that $P \cap -P = \{0\}$, $P \cup -P = G$ and $P + P \subseteq P$, then P is the set of positive elements of the group order defined by $g \leq h \Leftrightarrow h - g \in P$. An order is not a C-q.o, but there is a natural connection between orders and o-type C-q.o's. More precisely, we have the following:

Proposition 3.2

Assume (G, \lesssim) is an o-type C-q.o. Then $P := \{g \in G \mid -g \lesssim g\}$ is a positive cone and it is stable under " \sim ".

Proof. We use the following result which was proved in [8] (Lemma 3.6.(i)): For any $g, h \in G$, $h \lesssim -g$ implies $g \sim g + h$. Lemma 3.15 of [8] implies that P is stable under " \sim ". We clearly have $P \cap -P = \{0\}$ and $P \cup -P = G$. We just have to show $P + P \subseteq P$. We actually show that $-P$ is closed under addition. Let $g, h \in -P$. If h or g is 0 then obviously $g + h \in -P$, so we may assume that g and h are both non-zero. If $g \lesssim h$ then $g \lesssim -h$ which implies $g + h \sim h$ so $g + h \in -P$. Otherwise we have $h \lesssim -g$ which means $g + h \sim g$ so $g + h \in -P$. \square

Proposition 3.2 gives us a map $\Omega : \{\text{o-type C-q.o's on } G\} \rightarrow \{\text{group orders on } G\}$, where $\Omega(\lesssim)$ is the order corresponding to the positive cone $\{g \in G \mid -g \lesssim g\}$. This map is surjective: let \leq be an order of G with positive cone P . Then we can define a q.o on G as follows: declare that $0 \lesssim g \lesssim h$ holds for every $g \in -P \setminus \{0\}$ and $h \in P \setminus \{0\}$; declare then that $g \sim h$ whenever $g, h \in -P \setminus \{0\}$ and that $g \lesssim h \Leftrightarrow g \leq h$ whenever g, h are both in $P \setminus \{0\}$. One can easily check that this is a C-q.o and that it is o-type. By construction it obviously is a pre-image of \leq under Ω . The fact that Ω is surjective allows us to see orders as special case of C-q.o's. Through Ω we will be able to transform certain statements concerning C-q.o's into statements about orders. In other words, we can use C-q.o's as a uniform approach to ordered and valued groups.

We proved in [8] (Proposition 3.9) that C-q.o's satisfy the condition (*) of Lemma 1.3. Moreover, we know that if \lesssim is a C-q.o then a subgroup H is \lesssim -convex if and only if $H^{\geq 0}$ is \lesssim -convex. This allows us to apply the lemmas from section 1 to the case of C-q.o's:

Theorem 3.3

Let \lesssim be a C-q.o on G and $v : G \rightarrow \Gamma \cup \{\infty\}$ a valuation. The following statements are equivalent:

1. $\forall \gamma \in \Gamma, G^\gamma$ is \lesssim -convex.
2. $\forall \gamma \in \Gamma, G_\gamma$ is \lesssim -convex.
3. $\forall \gamma \in \Gamma, \lesssim$ induces a C-q.o \lesssim_γ on G^γ/G_γ .
4. v is compatible with \lesssim .

Moreover, \lesssim is v-type (respectively o-type) if and only if each \lesssim_γ is v-type (respectively o-type).

Proof. We just have to show that the induced q.o on the quotient G^γ/G_γ is a C-q.o. For this, we refer to the end of the proof of Proposition 3.11 in [8]. \square

Unlike compatible q.o's, we proved in [8] (Proposition 3.38) that the class of C-q.o's is stable under lifting. This allows us to state a Baer-Krull theorem for C-q.o's:

Theorem 3.4 (Baer-Krull for C-q.o's)

Let (G, v) be a valued group. The map:

$$\begin{array}{ccc} \{\text{C-q.o } \lesssim \mid v \text{ is compatible with } \lesssim\} & \longleftrightarrow & \{\text{family of C-q.o's } (\lesssim_\gamma)_{\gamma \in \Gamma}\} \\ \lesssim & \longmapsto & (\lesssim_\gamma)_{\gamma \in \Gamma}, \end{array}$$

where \lesssim_γ denotes the q.o induced by \lesssim on the quotient G^γ/G_γ , is a bijection. Moreover, \lesssim is v-type (respectively, o-type) if and only if each \lesssim_γ is v-type (respectively o-type).

Proof. We know from Proposition 3.38 of [8] that we can lift a family $(\lesssim_\gamma)_{\gamma \in \Gamma}$ of C-q.o's to obtain a C-q.o \lesssim on G , and \lesssim is obviously compatible with v by definition of the lifting. Conversely, if \lesssim is a C-q.o such that v is compatible with \lesssim , then by Theorem 3.3, we know that \lesssim induces a C-q.o \lesssim_γ on each G^γ/G_γ . It is easy to see that these operations are inverse to each other. \square

Since valuational q.o's are in particular C-q.o's, we have as an immediate corollary:

Theorem 3.5

Let (G, v) be a valued group. The map:

$$\left\{ \begin{array}{l} \text{valuations } w \mid v \text{ is a coarsening of } w \\ w \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{family of valuations } (w_\gamma)_{\gamma \in \Gamma} \\ (w_\gamma)_{\gamma \in \Gamma}, \end{array} \right.$$

where $w_\gamma : G^\gamma/G_\gamma \rightarrow v(G^\gamma \setminus G_\gamma) \cup \{\infty\}$ is defined by $w_\gamma(g + G_\gamma) := v(g)$ for any $g \in G^\gamma \setminus G_\gamma$ and $w_\gamma(0) := \infty$, is a bijection.

Thanks to Lemma 1.3, we see that if \preceq is o-type and if H is a \preceq -convex subgroup of G , then the image under Ω of the q.o induced on G/H by \preceq is exactly the order induced by $\Omega(\preceq)$ on G/H . This leads to another corollary of Theorem 3.4:

Theorem 3.6

Let (G, v) be a valued group. The map:

$$\left\{ \begin{array}{l} \text{orders } \leq \mid v \text{ is compatible with } \leq \\ \leq \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{family of orders } (\leq_\gamma)_{\gamma \in \Gamma} \\ (\leq_\gamma)_{\gamma \in \Gamma}, \end{array} \right.$$

where each \leq_γ is the order induced by \leq on the quotient G^γ/G_γ , is a bijection.

Proof. For each γ , let $\preceq_\gamma \in \Omega^{-1}(\leq_\gamma)$. By Theorem 3.4, we can lift $(\preceq_\gamma)_\gamma$ to G and obtain a C-q.o \preceq . Since each \preceq_γ is o-type, so is \preceq , so we can define \leq as $\Omega(\preceq)$. Then the order induced by \leq on each G^γ/G_γ is \leq_γ . □

4 q-sections and the classical Baer-Krull Theorem

Finally, we want to show how one can recover the classical Baer-Krull theorem from Theorem 3.6. We fix a valued field (K, v) with value group (G, \leq) . Note that if $(\leq_g)_{g \in G}$ is a family of group orders on the quotients K^g/K_g , then Theorem 3.6 only tells us that this family lifts to a group order on $(K, +)$, but there is no reason to think that this lifting is a field order in general. In order to understand the connection between Theorem 3.6 and the classical Baer-Krull theorem, we need to characterize the families $(\leq_g)_{g \in G}$ whose lifting to K is a field order.

We can achieve this by using the the notion of q-section developed in [9]. A q-section of the valued field (K, v) is a map $s : G \rightarrow K$ such that $s(0) = 1$, $v(s(g)) = g$ and $s(g + h) \equiv s(g)s(h) \pmod{K^2}$. It was proved in [9] that every valued field admits a q-section. We now fix a q-section s of (K, v) . Then for any $g \in G$, the map $\phi_g : Kv \rightarrow K^g/K_g, a + K_0 \mapsto s(g)a + K_g$ defines an isomorphism from Kv to K^g/K_g . If we take a family $(\leq_g)_{g \in G}$ of orders on the quotients K^g/K_g then the behavior of the ϕ_g 's with respect to \leq_g 's will tell us if the lifting of $(\leq_g)_g$ is a field order:

Proposition 4.1

Let $(\leq_g)_{g \in G}$ be a family of group orders on the quotients K^g/K_g . Then the lifting of $(\leq_g)_g$ to K is a field order if and only if the following conditions are satisfied:

1. \leq_0 is a field order of Kv

2. there exists a group homomorphism $\epsilon : G \rightarrow \{-1, 1\}$ such that for any $g \in G$, ϕ_g is order-preserving when $\epsilon(g) = 1$ and ϕ_g is order-reversing when $\epsilon(g) = -1$.

Proof. Denote by \leq the lifting. If \leq is a field order, then \leq_0 must be a field order because it is the order induced by \leq on Kv ; moreover, we can define $\epsilon(g) = 1$ if $0 < s(g)$ and $\epsilon(g) = -1$ if $s(g) < 0$, and one easily sees that ϵ has the desired property. Assume now that conditions (1) and (2) are satisfied. We already know from Theorem 3.6 that \leq is a group order of $(K, +)$, so \leq is a field order if and only if the set of positive elements of (K, \leq) is stable under multiplication. By definition of \leq , this is equivalent to saying that for any $a, b \in K$ with $g = v(a)$ and $h = v(b)$, $0 \leq_g a + K_g$ and $0 \leq_h b + K_h$ imply $0 \leq_{g+h} ab + K_{g+h}$. Note that since s is a q-section, we have $ab + K_{g+h} = d^2 \phi_{g+h}(\phi_g^{-1}(a + K_g)\phi_h^{-1}(b + K_h))$ for some $d \in K$; in particular, $ab + K_{g+h}$ has the same sign as $\phi_{g+h}(\phi_g^{-1}(a + K_g)\phi_h^{-1}(b + K_h))$. Now assume for example that ϕ_g is order-preserving and ϕ_h order-reversing. If $a + K_g$ and $b + K_h$ are both positive, we then have $\phi_g^{-1}(a + K_g) \leq_0 0$ and $0 \leq_0 \phi_h^{-1}(b + K_h)$. Since \leq_0 is a field order, this implies that $\phi_g^{-1}(a + K_g)\phi_h^{-1}(b + K_h) \leq_0 0$. Since ϵ is a group homomorphism, then ϕ_{g+h} is order-reversing, hence $0 \leq_{g+h} \phi_{g+h}(\phi_g^{-1}(a + K_g)\phi_h^{-1}(b + K_h))$. The other cases are treated similarly. \square

As a consequence of proposition 4.1 we have the following variant of the Baer-Krull theorem:

Theorem 4.2 (Baer-Krull, variant)

Let \mathcal{O} be the set of field orders of Kv and \mathcal{E} the set of group homomorphisms from G to $\{-1, 1\}$. Then $\mathcal{O} \times \mathcal{E}$ is in bijection with the set of families $(\leq_g)_{g \in G}$ whose lifting to K is a field order.

Proof. Assume that $\leq_0 \in \mathcal{O}$ and $\epsilon \in \mathcal{E}$ are given. Then define \leq_g on K^g/K_g as follows: If $\epsilon(g) = 1$, define $a + K_g \leq_g b + K_g \Leftrightarrow \phi_g^{-1}(a + K_g) \leq_0 \phi_g^{-1}(b + K_g)$; if $\epsilon(g) = -1$, define $a + K_g \leq_g b + K_g \Leftrightarrow \phi_g^{-1}(a + K_g) \geq_0 \phi_g^{-1}(b + K_g)$. By the previous proposition, this family of orders lifts to a field order on K . \square

Now assume that $(\pi_i)_{i \in I}$ is a family of elements of K such that $(v(\pi_i) + 2G)_{i \in I}$ is an \mathbb{F}_2 -Basis of $G/2G$. In order to recover Theorem 0.2 from Theorem 4.2, we need to show that the set of homomorphisms from G to $\{-1, 1\}$ is in bijection with the set of maps from I to $\{-1, 1\}$. First note that we can see I as a subset of G if we identify $i \in I$ with $v(\pi_i)$. Thus, any homomorphism $\epsilon : G \rightarrow \{-1, 1\}$ canonically induces a map $I \rightarrow \{-1, 1\}$ (just take $\epsilon|_I$). For the converse, note that every $g \in G$ has a decomposition $g = \sum_{i \in I} n_i v(\pi_i) + 2h$, where $h \in G$, $n_i \in \{0, 1\}$ and $n_i = 1$ only for finitely many i . If $\epsilon : I \rightarrow \{-1, 1\}$ is given, we can extend ϵ to a homomorphism $G \rightarrow \{-1, 1\}$ as follows: declare that $\epsilon(g) = 1$ if the number of $i \in I$ such that $n_i = 1$ in the decomposition of g is even, and declare $\epsilon(g) = -1$ if it is odd.

References

- [1] Françoise Delon : *C-minimal structures without the density assumption*, In Raf Cluckers, Johannes Nicaise et Julien Sebag, éditeurs : *Motivic Integration and its Interactions with Model Theory and Non-Archimedean Geometry*. Cambridge University Press, Berlin, 2011
- [2] A. J. Engler and A. Prestel, *Valued Fields*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, Zbl 1128.12009. MR 2183496. 286, 2005
- [3] S.M.Fakhruddin, *Quasi-ordered fields*, Journal of Pure and Applied Algebra 45, 207-210, 1987
- [4] L. Fuchs, *Partially ordered algebraic systems*, Pergamon Press, 1963
- [5] Salma Kuhlmann, *Ordered exponential fields*, The Fields Institute Monograph Series, vol 12, 2000
- [6] S. Kuhlmann, M. Matusinski, F. Point, *The valuation difference rank of a quasi-ordered difference field*, Groups, Modules and Model Theory - Surveys and Recent Developments in Memory of Rüdiger Göbel, Springer Verlag, 2017
- [7] G. Lehericy, *A structure theorem for abelian quasi-ordered groups*, preprint, arXiv number 1606.07710, 2016, submitted.
- [8] G. Lehericy, *On the structure of groups endowed with a compatible C-relation*, preprint, arXiv number 1609.01909, 2016, submitted.
- [9] A. Prestel, *Lectures on Formally Real Fields*, Springer Lecture Notes 1093, Springer-Verlag, 1984
- [10] S. Priess-Crampe, *Angeordnete Strukturen: Gruppen, Körper, projektive Ebenen*, Springer-Verlag, 1983

FACHBEREICH MATHEMATIK UND STATISTIK,
UNIVERSITÄT KONSTANZ,
78457, GERMANY.
Email address: salma.kuhlmann@uni-konstanz.de

FACHBEREICH MATHEMATIK UND STATISTIK,
UNIVERSITÄT KONSTANZ,
78457, GERMANY.
Email address: gabriel.lehericy@uni-konstanz.de