

**AN INVOLUTION ON DYCK PATHS THAT PRESERVES THE
RISE COMPOSITION AND INTERCHANGES THE NUMBER OF
RETURNS AND THE POSITION OF THE FIRST DOUBLE FALL**

MARTIN RUBEY

ABSTRACT. Motivated by a recent paper of Adin, Bagno and Roichman, we present an involution on Dyck paths that preserves the rise composition and interchanges the number of returns and the position of the first double fall.

1. INTRODUCTION

In a recent paper of Adin, Bagno and Roichman [1] a new symmetry property of 321-avoiding permutations is used to demonstrate the Schur positivity of the quasi-symmetric function associated with a certain set of permutations. More precisely, the authors show using a recursively defined bijection¹ that

$$\sum_{\pi \in \mathfrak{S}_n(321)} \mathbf{x}^{\text{LRMAX}(\pi)} q^{\text{bl}(\pi)} = \sum_{\pi \in \mathfrak{S}_n(321)} \mathbf{x}^{\text{LRMAX}(\pi)} q^{n - \text{lides}(\pi^{-1})}, \text{ where}$$

- $\mathfrak{S}_n(321)$ is the set of 321-avoiding permutations of $\{1, \dots, n\}$,
- $\text{LRMAX}(\pi)$ is the *set* of positions of the left-to-right maxima of π , and $\mathbf{x}^L = \prod_{i \in L} x_i$,
- $\text{bl}(\pi) = |\{i : \forall j \leq i : \pi(j) \leq i\}|$ is the number of blocks of π and
- $\text{lides}(\pi) = \max\{0\} \cup \{i : \pi(i) > \pi(i+1)\}$ is the position of the last descent of π .

We exhibit an iterative involution that proves the joint symmetry of the two statistics linked by Adin, Bagno and Roichman.

Let us first transform the statistics involved into statistics on Dyck paths, using a bijection due to Krattenthaler [3]². To do so, it is convenient to define a Dyck path as a path in \mathbb{N}^2 consisting of $(1, 0)$ and $(0, 1)$ steps beginning at the origin, ending at (n, n) , and never going below the diagonal $y = x$. We refer to $(1, 0)$ -steps also as *east steps* or *falls*, and to $(0, 1)$ -steps also as *north steps* or *rises*, and the number n as the *semilength* of the path. Krattenthaler's bijection maps

- $\mathfrak{S}_n(321)$ to the set of Dyck paths \mathfrak{D}_n of semilength n ,
- $\text{LRMAX}(\pi)$ to $\text{RISES}(D)$, the set of x -coordinates (plus 1) of the north steps of the Dyck path,
- $\text{bl}(\pi)$ to $\text{ret}(D)$, the number of *returns* of the path, that is, the number of east steps that end on the main diagonal,
- $\text{lides}(\pi^{-1})$ to $\text{ldr}(D)$, the y -coordinate of the middle point of the last double rise, that is, the last pair of two consecutive north steps. For the path without double rises, we set $\text{ldr}(D) = 0$.

¹<http://www.findstat.org/MapsDatabase/Mp00124>

²<http://www.findstat.org/MapsDatabase/Mp00119>

The purpose of this note is to provide an involution Φ that implies

$$(1) \quad \sum_{\pi \in \mathfrak{D}_n} \mathbf{x}^{\text{RISES}(\pi)} p^{\text{ret}(D)} q^{n-\text{ldr}(D)} = \sum_{\pi \in \mathfrak{D}_n} \mathbf{x}^{\text{RISES}(\pi)} p^{n-\text{ldr}(D)} q^{\text{ret}(D)}.$$

2. AN INVOLUTION ON DYCK PATHS

In the spirit of [2]³, let us first introduce an invertible map ϕ from Dyck paths that do not end with a single east step and satisfy $\text{ldr}(D) < n - 1$ (that is, $\text{ldr}(D)$ is not maximal), such that

- $\text{RISES}(\phi(D)) = \text{RISES}(D)$
- $\text{ret}(\phi(D)) = \text{ret}(D) + 1$
- $\text{ldr}(\phi(D)) = \text{ldr}(D) + 1$

Iterating this map (or its inverse) we obtain a map Φ that proves Equation (1).

To define ϕ , decompose D as the concatenation of three paths P , Q and R where

- R is maximal such that R contains only single rises, begins with a rise and Q ends with an east step, and
- QR is a prime Dyck path, that is, its only return is the final east step.

Let $n_0 Q_1 n_1 \dots n_k Q_k$ be the prime decomposition of Q , that is, all the Q_i are prime Dyck paths and the n_i are sequences of north steps. Note that n_0 is non-empty, because Q does not return to the diagonal. We now distinguish three cases:

- (1) If $n_1 \neq \emptyset$, define $\phi(D) = P Q_1 n_1 \dots n_k Q_k n_0 R$.
- (2) If $n_1 = \dots = n_k = \emptyset$, let n be the sequence of north steps of the final rise in Q except one, and let Q' be Q with n_0 and n erased. Then define $\phi(D) = P n Q' n_0 R$.
- (3) Otherwise, let n be the sequence of north steps of the final rise in Q except one, and let Q' be Q with n_0 and Q_1 erased. Then define $\phi(D) = P Q_1 n Q' n_0 R$.

To show that ϕ is invertible we provide an inverse for each of these three cases. Let D be a Dyck path that does not end with a single east step and has at least two returns. Decompose D as the concatenation of three paths P , Q and R where

- R is maximal such that R contains only single rises and begins with a rise, and
- QeR is a Dyck path with precisely two returns, e is a maximal sequence of east steps and n is a maximal sequence of north steps.

Let $n_0 Q_1 n_1 \dots Q_k n_k$ be the prime decomposition of Q .

- (1) If Q ends with a double rise, we have $\phi^{(-1)}(D) = P n Q e R$.
- (2) Otherwise, if Q has no returns, we have $\phi^{(-1)}(D) = P n Q_1 n_1 \dots Q_k n_k n_0 e R$.
- (3) Otherwise, $\phi^{(-1)}(D) = P n Q_1 Q_2 n_2 \dots Q_k n_k n_1 e R$.

REFERENCES

- [1] R. M. Adin, E. Bagno, and Y. Roichman. “Block decomposition of permutations and Schur-positivity”. In: *ArXiv e-prints* (Nov. 2016). arXiv:1611.06979 [math.CO].
- [2] Sergi Elizalde and Martin Rubey. “Symmetries of statistics on lattice paths between two boundaries”. In: *Adv. Math.* 287 (2016), pp. 347–388. ISSN: 0001-8708. DOI: 10.1016/j.aim.2015.09.025. arXiv:1305.2206 [math.CO].

³<http://www.findstat.org/MapsDatabase/Mp00118>




FIGURE 1. A path and its image for case (1)




FIGURE 2. A path and its image for case (2)




FIGURE 3. A path and its image for case (3)

- [3] Christian Krattenthaler. “Permutations with restricted patterns and Dyck paths”. In: *Adv. in Appl. Math.* 27.2-3 (2001). Special issue in honor of Dominique Foata’s 65th birthday (Philadelphia, PA, 2000), pp. 510–530. ISSN: 0196-8858. DOI: 10.1006/aama.2001.0747. arXiv:math/0002200 [math.CO].

E-mail address: Martin.Rubey@tuwien.ac.at

FAKULTÄT FÜR MATHEMATIK UND GEOINFORMATION, TU WIEN, AUSTRIA