

Multi-locus data distinguishes between population growth and multiple merger coalescents

Jere Koskela

`koskela@math.tu-berlin.de`

Institut für Mathematik

Technische Universität Berlin

Straße des 17. Juni 136, 10623 Berlin

Germany

May 14, 2022

Abstract

We introduce a low dimensional function of the site frequency spectrum that is tailor-made for distinguishing coalescent models with multiple mergers from Kingman coalescent models with population growth, and use this function to construct a hypothesis test between these two model classes. The null and alternative sampling distributions of our statistic are intractable, but its low dimensionality renders these distributions amenable to Monte Carlo estimation. We construct kernel density estimates of the sampling distributions based on simulated data, and show that the resulting hypothesis test dramatically improves on the statistical power of a current state-of-the-art method. A key reason for this improvement is the use of multi-locus data, in particular averaging observed site frequency spectra across unlinked loci to reduce sampling variance. We also demonstrate the robustness of our method to nuisance and tuning parameters. Finally we demonstrate that the same kernel density estimates can be used to conduct parameter estimation, and argue that our method is readily generalisable for applications in model selection, parameter inference and experimental design.

1 Introduction

The Kingman coalescent [Kingman, 1982a,b,c, Hudson, 1983a,b, Tajima, 1983] describes the random ancestral relations among DNA sequences sampled from large populations, and is a prominent model with which to make predictions about genetic diversity. This popularity derives from its robustness: a wide class of genealogical models all have the Kingman coalescent, or a variant of it, as their limiting process when the population size is large [Möhle, 1998]. Together, the Kingman coalescent and the infinitely-many-sites (IMS) model also form a tractable model of genetic evolution [Watterson, 1975]. Hence, many inference methods based on the Kingman coalescent have been developed; see e.g. Donnelly and Tavaré [1995], Hudson [1990], Nordborg [2001], Hein et al. [2005] or Wakeley [2007] for reviews.

The site frequency spectrum (SFS) at a given locus is an important and popular statistic by which to summarise genetic data under the IMS model and a coalescent process. Quantities of interest, such as the expectations and covariances of the SFS, are easily computed under a Kingman coalescent [Fu, 1995].

Despite its robustness, many evolutionary histories can also lead to significant deviations from the Kingman coalescent. A variety of statistical tools is available for detecting such deviations, e.g. Tajima's D [Tajima, 1989], Fu and Li's D [Fu and Li, 1993] or Fay and Wu's H [Fay and Wu,

2000], which are all functions of the SFS. However, they do not always allow the identification of the actual evolutionary mechanisms leading to the deviation from the Kingman coalescent. Eldon et al. [2015] investigated the ability of a single-locus SFS to distinguish two different deviations from the Kingman coalescent:

1. population growth, in particular exponential or algebraic population growth, and
2. gene genealogies described by so-called Λ -coalescents [Sagitov, 1999, Pitman, 1999, Donnelly and Kurtz, 1999] featuring multiple mergers. There is growing evidence that such coalescents are an appropriate model for organisms with high fecundity coupled with a skewed offspring distribution [Beckenbach, 1994, Árnason, 2004, Eldon and Wakeley, 2006, Sargsyan and Wakeley, 2008, Hedgecock and Pudovkin, 2011, Birkner et al., 2011, Steinrücken et al., 2013, Tellier and Lemaire, 2014].

Both scenarios lead to an excess of singletons in the SFS compared to the Kingman coalescent, leading e.g. to a negative value of Tajima's D [Durrett and Schweinsberg, 2005]. Eldon et al. [2015] showed that a single-locus SFS can distinguish between scenarios 1 and 2 with moderate statistical power, at least when the rate of mutation was sufficiently high and the deviation from the Kingman coalescent, the population growth rate in the scenario 1 and the prevalence of multiple mergers in scenario 2, was sufficiently large.

The present work builds on the result of Eldon et al. [2015] by proposing a new statistical test to distinguish between scenarios 1 and 2. A key development is that the proposed tests are based on multiple site frequency spectra corresponding to multiple unlinked loci. Modern sequencing technologies have made sequence data from multiple linkage groups commonplace, and we show that making use of multi-locus data in this way greatly improves the power of statistical tests. We assume that no recombination takes place within loci.

In addition, we extend the work of Eldon et al. [2015] by explicitly including diploidy into the models under consideration. Specifically, we consider a diploid, biparental population with symmetric mating, and with each parent contributing a single chromosome to their offspring. This results in ancestries modeled as time-changed Kingman coalescents in scenario 1, and Ξ -coalescents, incorporating up to four simultaneous mergers of groups of multiple lineages, in scenario 2 [Birkner et al., 2013a]. In brief, the presence of up to four simultaneous mergers arises because genetic material at given locus can coalesce at either copy in either diploid parent.

While our new statistical tests contain the population rescaled mutation rate, $\theta/2$, as a nuisance parameter, we show empirically that the statistical power achieved by our method is extremely robust to misspecification of this parameter. Hence, in practice relying on a generalised Watterson estimator for the mutation rate can be expected to produce reliable inferences.

In addition to selecting the model class which best explains the data, it is also of interest to infer those parameters within that class which result in the best fit. Hence, parameter inference for Λ - and Ξ -coalescents has been a growing area of research in recent years [Eldon and Wakeley, 2006, Birkner and Blath, 2008, Sargsyan and Wakeley, 2008, Eldon and Wakeley, 2009, Birkner et al., 2011, Eldon, 2011, Birkner et al., 2013b, Steinrücken et al., 2013, Koskela et al., 2015, Zhu et al., 2015, Blath et al., 2016, Koskela et al., 2017]. We contribute to this body of work by demonstrating that the statistic used in our hypothesis test is also able to distinguish between different parameter values within the alternative hypothesis multiple merger class. Our method is scalable to large data sets in contrast to many of those cited above, and we demonstrate its robustness and unbiasedness empirically via simulations.

2 Summary statistic and approximate likelihood

Consider a sample of n DNA sequences taken at a given genetic locus, and assume that derived mutations can be distinguished from ancestral states. For $n \in \mathbb{N}$ let $[n] := \{1, \dots, n\}$, and let $\xi_i^{(n)}$ denote the total number of sites at which the mutant base appears $i \in [n-1]$ times. Then

$$\boldsymbol{\xi}^{(n)} := \left(\xi_1^{(n)}, \dots, \xi_{n-1}^{(n)} \right)$$

is referred to as the *unfolded* site-frequency spectrum based on the n DNA sequences. If mutant and ancestral type cannot be distinguished, the *folded* spectrum $\boldsymbol{\eta}^{(n)} := (\eta_1^{(n)}, \dots, \eta_{\lfloor n/2 \rfloor}^{(n)})$ [Fu, 1995] is often considered instead, where

$$\eta_i^{(n)} := \frac{\xi_i^{(n)} + \xi_{n-i}^{(n)}}{1 + \delta_{i, n-i}}, \quad 1 \leq i \leq \lfloor n/2 \rfloor,$$

and $\delta_{i,j} = 1$ if $i = j$, and is zero otherwise. Define $\boldsymbol{\zeta}^{(n)} = (\zeta_1^{(n)}, \dots, \zeta_{n-1}^{(n)})$ as the normalised unfolded SFS, whose entries are given by $\zeta_i^{(n)} := \xi_i^{(n)} / |\boldsymbol{\xi}^{(n)}|$, where $|\boldsymbol{\xi}^{(n)}| := \xi_1^{(n)} + \dots + \xi_{n-1}^{(n)}$ is the total number of segregating sites. We adopt the convention that $\boldsymbol{\zeta}^{(n)} = \mathbf{0}$ if there are no segregating sites.

Even though both scenarios 1 and 2 predict an excess of singletons, Eldon et al. [2015] showed that the expected tail of the normalised SFS varies between the two when the singletons have been matched [Eldon et al., 2015, Figure 1]. Hence we define the lumped tail

$$\bar{\zeta}_k^{(n)} := \sum_{j=k}^{n-1} \zeta_j^{(n)}, \quad 3 \leq k \leq n-1 \quad (1)$$

and consider the summary statistic $(\zeta_1^{(n)}, \bar{\zeta}_k^{(n)})$ for some fixed k , to be specified. This two-dimensional summary of the SFS can be expected to distinguish scenarios 1 and 2, while exploiting a lower dimensionality to reduce sampling variance (see the discussion on the effect of lumping in [Eldon et al., 2015] for an account of the same phenomenon in the context of approximate Bayesian computation) and reduce the number of Monte Carlo simulations needed to robustly characterise its sampling distribution.

We make use of multi-locus data by computing an SFS independently for each locus, and averaging over all available loci to reduce variance. To this end, let $(\zeta_1^{(n)}(j), \bar{\zeta}_k^{(n)}(j))$ denote the singleton class and lumped tail corresponding to the j^{th} locus, and

$$(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)}) := \frac{1}{L} \sum_{j=1}^L (\zeta_1^{(n)}(j), \bar{\zeta}_k^{(n)}(j)) \quad (2)$$

denote the mean SFS when there are L loci.

Remark 1. In scenario 1, and for binary, Kingman-type coalescents in general, (2) converges to its expected value as $L \rightarrow \infty$ due to the strong law of large numbers. In contrast, in scenario 2, and for multiple merger coalescents in general, unlinked loci have positively correlated coalescence times and hence a law of large numbers does not hold. Sampling variance will still be reduced by averaging, but by less than in scenario 1, and an $L \rightarrow \infty$ limit will be random if it exists.

Let Π denote an arbitrary coalescent process, $\theta/2$ denote a population-rescaled mutation rate, and $P^{\Pi, \theta}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})$ denote the sampling distribution of (2) under the coalescent Π and mutation

rate $\theta/2$. This distribution is intractable, but if it were available then the likelihood ratio test statistic

$$\frac{\sup_{\Pi \in \Theta_1, \theta > 0} P^{\Pi, \theta}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0, \theta > 0} P^{\Pi, \theta}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})} \quad (3)$$

would be a desirable tool for distinguishing between coalescent models in the class Θ_0 , the null hypothesis, from an alternative class Θ_1 , based on an observed value of (2).

Eldon et al. [2015] proposed a Poisson approximation to a test akin to (3), based on the whole unnormalised SFS, by assuming that the hidden coalescent ancestry of their sample coincided with the expected coalescent branch lengths. These expected branch lengths were then approximated by simulation. The number of mutations was also assumed to always coincide exactly with the number of observed segregating sites, in what is known as the “fixed- s ” method [see e.g. Depaulis and Veuille, 1998, Ramos-Onsins and Rozas, 2002]. In this paper we approximate $P^{\Pi, \theta}$ by simulation directly. This avoids the unquantifiable bias introduced by the Poisson and fixed- s approximations, and is scalable because only a two-dimensional sampling distribution needs to be approximated, regardless of the sample size or the number of alleles. Of course replacing the sampling distributions in (3) with estimators will itself introduce bias, but provided that the estimators are close to the truth then it is reasonable to expect the bias to be small in comparison to an ad-hoc Poisson approximation. In particular, Figure 9 in Section 4 shows no evidence of bias when using our approximate sampling distributions to infer parameters.

Remark 2. A similar method based on kernel density estimation of an intractable likelihood function from simulated data was developed by Diggle and Gratton [1984], albeit with the goal of obtaining point estimators by maximising their approximation of the likelihood. Our method is also reminiscent of the approximate Bayesian computation paradigm [Beaumont, 2010], in that likelihood evaluations are replaced by simulations from a family of models which is assumed to contain that generating the data.

In order to produce an approximate sampling distribution $\hat{P}^{\Pi, \theta}$, a simulation algorithm was developed for producing samples from a specified coalescent Π with mutation rate $\theta/2$ for a desired sample size and number of unlinked alleles, within which we assume no recombination takes place. The method is described in detail in the appendix, and a C++ implementation is available at <https://github.com/JereKoskela/Beta-Xi-Sim>. Three model classes have been implemented under both haploid and diploid models:

1. Beta($2 - \alpha, \alpha$)- Ξ -coalescents for $\alpha \in [1, 2]$ derived from Λ -coalescents with $\Lambda = \text{Beta}(2 - \alpha, \alpha)$ by randomly splitting all merging lineages into four groups, and coalescing each group separately as in [Birkner et al., 2013a, equation (29)] and [Blath et al., 2016, equation (15)],
2. the Kingman coalescent with exponential population growth at rate $\gamma > 0$,
3. the Kingman coalescent with algebraic population growth with exponent $\gamma > 0$.

In the haploid case the first class would more accurately be described as consisting of Beta($2 - \alpha, \alpha$)- Λ -coalescents [Tellier and Lemaire, 2014, equations (A1) and (A4)], because simultaneous multiple mergers are only assumed to appear due to diploidy. Implementations for further model classes, such as Ξ -coalescents driven by measures other than the Beta($2 - \alpha, \alpha$) class or derived from more general models of mating patterns, such as those described in [Birkner et al., 2017], could be easily added. Further evolutionary forces such as natural selection could also be incorporated, though naturally both of these generalisations come at increased computational cost.

Once a sample of simulated realisations of (2) has been produced under the desired null and alternative hypotheses (or suitable discretisations in the case of interval hypotheses), kernel

density estimators were fitted to these samples using the `kde` function in the `ks` package (version 1.10.4) in R under default settings. For a comprehensive introduction to kernel density estimation, see e.g. [Scott, 1992]. In particular, the bandwidth is determined using the SAMSE estimator of [Duong and Hazelton, 2003, equation (6)] as implemented in the `Hpi` function of the aforementioned `ks` package. This package allows kernel density estimation in up to six dimensions, so that higher dimensional statistics could be used to distinguish model classes that are not well separated by just singletons and the lumped tail. However, this increase in dimensionality would come at the cost of increasingly intensive simulation time as more samples are needed to accurately represent a higher dimensional sampling distribution, and was not necessary for our purposes.

Once these kernel density estimators have been produced, they can be substituted into (3) to obtain an approximation

$$\frac{\sup_{\Pi \in \Theta_1, \theta > 0} \hat{P}^{\Pi, \theta}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0, \theta > 0} \hat{P}^{\Pi, \theta}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}.$$

However, simulating samples and producing kernel density estimators for combinations of coalescent processes and mutation rates also incurs a significant computational cost. To alleviate it, we assume that a computationally cheap estimator $\hat{\theta}$ is available, e.g. the generalised Watterson estimator

$$\hat{\theta} = |\xi^{(n)}| / \mathbb{E}^{\Pi}[T^{(n)}],$$

where $\mathbb{E}^{\Pi}[T^{(n)}]$ denotes the expected tree length from n leaves under coalescent mechanism Π . We then consider the test statistic

$$\frac{\sup_{\Pi \in \Theta_1} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})} \quad (4)$$

instead. Because our method makes use of the normalised SFS, it is highly insensitive to misspecification of the mutation rate, unless the mutation rate is small enough that normalisation of the SFS itself becomes a source of variance. This robustness will be confirmed by simulation in the next section.

The statistic (4) is straightforward to evaluate pointwise, and further simulated data can be used to obtain an empirical quantile ρ_{α} such that

$$\sup_{\Pi \in \Theta_0} \mathbb{P}^{\Pi, \hat{\theta}} \left(\frac{\sup_{\Pi \in \Theta_1} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})} \geq \rho_{\alpha} \right) \leq \alpha,$$

where $\mathbb{P}^{\Pi, \hat{\theta}}$ denotes the law of the coalescent process under coalescent mechanism Π and mutation rate $\hat{\theta}/2$. Given such a quantile, a hypothesis test of size α for an observed pair $(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})$ is given by

$$\Phi(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)}) = \begin{cases} 0 & \text{if } \frac{\sup_{\Pi \in \Theta_1} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})} \leq \rho_{\alpha} \\ 1 & \text{if } \frac{\sup_{\Pi \in \Theta_1} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})}{\sup_{\Pi \in \Theta_0} \hat{P}^{\Pi, \hat{\theta}}(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)})} > \rho_{\alpha} \end{cases}, \quad (5)$$

where $\Phi(\zeta_{1,L}^{(n)}, \bar{\zeta}_{k,L}^{(n)}) = 1$ corresponds to rejecting the null hypothesis. In the next section we demonstrate the performance and robustness of this test, and compare it to the Poisson-fixed- s test of Eldon et al. [2015].

Remark 3. Hypothesis tests only provide information on the model fit of the alternative hypothesis relative to the null. Hence, if the null hypothesis is a poor fit to the data, a high value

of (4) does not necessarily indicate that the alternative hypothesis is a good fit, but only a better one than the null hypothesis. The empirical sampling distributions constructed in order to evaluate (4) can be used to address this ambiguity to some extent, e.g. by plotting observed data along with simulated realisations in scatter plots akin to Figure 1 in the next section. This ability to overcome a limitation of hypothesis tests arises out of the low dimensionality of the statistic (2), and the ease with which its sampling distribution can hence be visualised.

3 Results: model selection

In this section we present simulation studies demonstrating the statistical power and robustness of (5). For comparison, the same simulation study will also be undertaken using the Poisson-fixed- s approximate likelihood ratio test of [Eldon et al., 2015]. For ease of computation we discretise the null and alternative hypotheses, and take them to be

$$\begin{aligned} \Theta_0 &:= \{\text{Kingman coalescent with exponential growth at rate } \gamma \in \{0, 0.1, 0.2, \dots, 0.9, 1, 1.5, 2, \\ &\quad 2.5, 3, 3.5, 4, 5, 6, \dots, 19, 20, 25, 30, 35, 40, 50, 60, \dots, 990, 1000\}\} \cup \{\text{Kingman coalescent} \\ &\quad \text{with algebraic growth with exponent } \gamma \in \{0, 0.05, 0.1, \dots, 1.25, 1.3, 1.4, \dots, 3.9, 4.0\}\}, \\ \Theta_1 &:= \{\text{Beta}(2 - \alpha, \alpha)\text{-}\Xi\text{-coalescents with } \alpha \in \{1, 1.025, \dots, 1.975, 2\}\}, \end{aligned}$$

respectively. The somewhat irregular discretisation has been chosen to yield comprehensive coverage of the sampling distribution of the continuous hypotheses based on trial runs, as indicated in Figure 1 in which the sampling distributions appear continuous. Note that all three model classes coincide with the Kingman coalescent at the far left end of the cloud of points, which corresponds to $\gamma = 0$ in scenario 1 and $\alpha = 2$ in scenario 2.




Figure 1: Scatter plot of the joint distribution of the summary statistic (2) with $L = 23$ loci, lumping from $k = 15$ and sample size $n = 100$ on the left and $n = 500$ on the right. Red dots denote realisations from a $\text{Beta}(2 - \alpha, \alpha)\text{-}\Xi$ -coalescent, green dots from an algebraic growth coalescent and blue dots from an exponential growth coalescent. The discontinuities are due to discretisation of the hypotheses.

Note that Eldon et al. [2015] considered the two classes in our null hypothesis as separate hypotheses. Hence the power function of their test is expected to be lower than in their results, since our null hypothesis is a strictly larger model class. Also note that we have significantly

reduced the scope of the algebraic growth rate γ from the $\{0, 1, \dots, 999, 1000\}$ considered by Eldon et al. [2015], because even the singleton class could not be matched to higher rates of algebraic growth by any coalescent in Θ_1 . The size of tests is set at $\alpha = 0.01$ throughout this section.

We emphasize that distinguishing a $\text{Beta}(2 - \alpha, \alpha)$ - Ξ -coalescent from a class of binary coalescents is harder than distinguishing the corresponding $\text{Beta}(2 - \alpha)$ - Λ -coalescent from the same binary class. This is because randomly assigning mergers into four groups can split multiple mergers into either outright binary mergers, or smaller multiple mergers. Thus, the SFS of the resulting Ξ -coalescent tree will resemble the SFS of a binary coalescent more closely than the same Λ -coalescent tree would have done. Thus, our test faces a twofold increase in difficulty in comparison to the scenarios considered by Eldon et al. [2015]: a richer null hypothesis, as well as a less distinct alternative hypothesis.

We simulated 1000 replicates of samples of size $n = 100$ and $n = 500$, with $L = 23$ unlinked loci and with mutation rate $\theta/2 = N_e^{\alpha-1}\mu$, with effective population size $N_e = 1000$ and per-locus-per-generation mutation rate set at both $\mu = 2.6$ and $\mu = 0.26$, for each of the coalescent processes in Θ_0 and Θ_1 . The scaling parameter α in the mutation rate denotes the parameter of the Θ_1 class, with $\alpha = 2$ corresponding to the Kingman coalescent as per the scaling results of Schweinsberg [2003]. The numbers of loci and mutation parameters have been chosen as reasonable approximations to Atlantic cod, which has 23 linkage groups averaging roughly 26 million base pairs in length [Tørresen et al., 2017, Supplementary Table 3]. The mutation rate was set at 10^{-8} per generation per base pair. Atlantic cod is a species for which multiple mergers have frequently been suggested as an important evolutionary mechanism [see e.g. Steinrücken et al., 2013, Tellier and Lemaire, 2014, and references therein]. Simulation times per replicate sample ranged from thousandths of a second to 10 seconds on a single Intel i5-2520M 2.5 GHz processor.

We also simulated independent realisations of the “fixed- s ” data sets of Eldon et al. [2015] to facilitate a comparison, keeping the total number of segregating sites at $s = 50$ as in the original article. We extend their Poisson-fixed- s approximate likelihood test to our multi-locus setting in two ways: by treating multiple loci as independent, or averaging over them as in the KDE case. A comparison of the empirical power as a function of the α -parameter in the $\text{Beta}(2 - \alpha, \alpha)$ -distribution is presented in Figure 2.

Figure 2 shows that (5) substantially improves on the test of [Eldon et al., 2015] in terms of the power to distinguish multiple merger coalescents from population growth. The results for $n = 100$ are already promising, and samples of size $n = 500$ yield statistical power estimates of 1 for a significant range of parameter values.

In Figure 3 we investigate the effect of the lumping threshold k in (1). Lumping corresponding to $k = 2$ was not included, as it yields an effectively one dimensional statistic corresponding to just the singleton class. Lumping from a very low level, say $k \in \{3, 4, 5, 6\}$, can diminish statistical power as mid-range entries of the SFS confound the contribution of the bulk, but beyond these low numbers the good performance of our method is highly insensitive to this tuning parameter.

Figure 4 investigates the effect of averaging across loci. When averaging over $l < L = 23$ loci, we treat the the 1000 simulated replicates of L loci as $1000[L/l]$ replicates to keep the total amount of data as constant as possible. For example, the $l = 1$ curve corresponds to treating the data as 23×1000 independent replicates, the $l = 2$ curve corresponds to treating the data as 11×1000 independent replicates, each being the average of two loci, and so on until the final, $l = 23$ curve corresponds to 1000 replicates, each being the average of all 23 loci. Figure 4 shows diminishing returns as the number of loci being averaged increases, but does not reach a saturation point in the same way as Figure 3; for the hypotheses and parameters considered here, adding more loci always improves power.




Figure 2: Comparison of the Poisson-fixed- s approximate likelihood ratio test with $s = 50$ and assuming either independent loci (“fixed- s ”) or applied to a single SFS obtained by averaging 23 loci (“Mean fixed- s ”), as well as the KDE approximation proposed in this article with $L = 23$ loci and lumping from $k = 15$. Sample size $n = 100$ on the left and $n = 500$ on the right. The expected branch lengths for the Poisson-fixed- s methods were estimated from 100 000 realisations.




Figure 3: Effect of the lumping cutoff in (2) from $k = 3$ (dark green) to $k = 20$ (pink). Sample size $n = 100$ on the left and $n = 500$ on the right.

Finally, we investigate the effect of misspecification of the mutation rate under which inference is conducted. Figures 5 and 6 depict the statistical power resulting from our test when the mutation rate is either too small or too large by a factor of 10. It can be seen that the power curves are nearly indistinguishable from those obtained using the true mutation rate in earlier figures. This confirms empirically the heuristic that statistics based on the normalised SFS should be highly robust to misspecification of the mutation rate.




Figure 4: Effect of the number of loci over which (2) is averaged from $L = 1$ (dark green) to $L = 23$ (pink). Sample size $n = 100$ on the left and $n = 500$ on the right.

We conclude this section with a final simulation study using the reduced null hypothesis

$$\tilde{\Theta}_0 := \{\text{Kingman coalescent with exponential growth at rate } \gamma \in \{0, 0.1, 0.2, \dots, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 5, 6, \dots, 19, 20, 25, 30, 35, 40, 50, 60, \dots, 990, 1000\}\}.$$

This is to facilitate a more direct comparison with the results of Eldon et al. [2015], though we still focus on diploid, multi-locus coalescent models as opposed to the haploid, single-locus coalescents analysed in that work. Figure 7 presents an empirical power function for testing $\tilde{\Theta}_0$ versus Θ_1 using (5). In this case averaging over 23 loci results in extremely good performance even from the “fixed- s ” method, though our KDE-based approximation still outperforms it when multiple mergers are small and infrequent.

4 Results: parameter inference

The kernel density estimators produced in order to approximate the hypothesis test (5) can also be used to perform parameter inference. In this section we investigate the ability of the statistic (2) to distinguish between Beta($2 - \alpha$, α)- Ξ -coalescents by treating the kernel density estimators $\hat{P}^{\Pi, \hat{\theta}}$ as the true likelihood functions.

Figure (8) shows 1000 replicates of simulated likelihood functions with parameter space Θ_1 from Section 3. These have been produced by computing kernel density estimators $\hat{P}^{\Pi, \theta}$ for the likelihood functions as outlined in Section 3, and using these estimators in place of the true, intractable likelihood. The curves are centered around the true values of α on average, showing little evidence of bias from the use of an approximate likelihood function. However, it is clear that realisations resulting in a substantial mismatch between the truth and the maximiser of the approximate likelihood are possible, especially for lower values of α . Note, however, that the Kingman coalescent, $\alpha = 2$, can be excluded with high confidence as soon as $\alpha \lesssim 1.9$. We emphasize that this comparison is different to the hypothesis tests in Section 3 because the comparison is between $\alpha = 2$ and $\alpha < 2$, and not between the two model classes Θ_0 and Θ_1 .

Figure 9 demonstrates the performance of the resulting approximate maximum likelihood estimator for α . The plot clearly centres on a straight, diagonal line confirming that the using an




Figure 5: A repeat of the sensitivity analyses of Figures 3 (top row) and 4 (bottom row) when the true mutation rate is 10 times higher than that used in the simulation algorithm.

approximate likelihood results in no discernible bias. As suggested by Figure 8, the distribution of the maximum likelihood estimator clearly has heavy tails, but Figure 9 also shows that the two central quantiles very tightly around the true parameter values.

Finally, figure 10 demonstrates that parameter inference is also robust to misspecification of the mutation rate. The distribution of maximum likelihood estimators is essentially unchanged when the mutation rate is misspecified by a factor of 10, and in particular the use of an approximate likelihood still results in no bias. Hence, use of the Watterson estimator is justifiable with real data sets in this context, as well as for model selection, when the true mutation is unknown.

5 Discussion

We have shown that directly modelling the sampling distribution of a well chosen function of the SFS based on simulated realisations can result in statistical tests which can reliably distinguish




Figure 6: A repeat of the sensitivity analyses of Figures 3 (top row) and 4 (bottom row) when the true mutation rate is 10 times lower than that used in the simulation algorithm.

multiple merger coalescents from population growth. The same statistics can also be used for parameter inference, albeit with a somewhat lower degree of confidence. The low dimensionality of (2), our function of choice, ensures that a simulation based approach is scalable to non-trivial data sets. Our tests focused on hundreds of samples and tens of loci, and the total run time for both simulating the various data sets and computing all the power curves in Sections 3 and 4 was under a week on a single Intel i5-2520M 2.5 GHz processor. Since much of this time was spent producing independent simulation replicates, it is obvious that parallel computing can yield substantial improvement.

Robustness of both our model selection and parameter inference to misspecification of the mutation rate helps alleviate the computational cost of our method, as otherwise data sets corresponding to a grid of mutation rates would have to be simulated. This results in either faster results, or enables the analysis of more complex coalescent families such as the Beta(β_1, β_2)- Ξ -coalescents for arbitrary $\beta_1 > 0$ and $\beta_2 > 0$. However, it is not clear whether such coalescents can be derived from a biologically relevant individual based models, and what the corresponding




Figure 7: Comparison of the Poisson-fixed- s approximate likelihood ratio test with $s = 50$ and assuming either independent loci (“fixed- s ”) or applied to a single SFS obtained by averaging 23 loci (“Mean fixed- s ”), as well as the KDE approximation proposed in this article with $L = 23$ loci and lumping from $k = 15$, using the reduced null hypothesis Θ_0 . Sample size $n = 100$ on the left and $n = 500$ on the right. The expected branch lengths for the Poisson-fixed- s methods were estimated from 100 000 realisations.

real time embeddings are. Other more complex hypotheses, such as (weak) selection under a Kingman coalescent, or spatial structure, could also be incorporated into the tests, provided that SFS data can be simulated under the corresponding models. Off-the-shelf methods for kernel density estimation in up to six dimensions exist (c.f. the `ks` package in R), which further suggests that more complex scenarios can be handled within our framework as it is not restricted to two dimensional summary statistics.

In addition to hypothesis testing and inference, our method also lends itself to design of experiments by providing estimates of likelihood functions and statistical power under various scenarios. In particular, we have been able to show that 500 samples of 23 unlinked loci results tests with very high power to distinguish $\text{Beta}(2 - \alpha, \alpha)$ - Ξ -coalescents from simple population growth models across most of the $\alpha \in [1, 2]$ parameter range. By way of example, previous estimates have identified $\alpha = 1.5$ as a plausible value for Atlantic cod [Birkner et al., 2011, Table 2], for which our two hypotheses can be distinguished with near certainty. The dramatic improvement in power yielded by increasing the number of loci being averaged in Figure 4 indicates that it is advisable to sequence as many unlinked loci as possible, in which case sample sizes in the hundreds are sufficient. More complex questions of experimental design will, of course, require bespoke simulations run on a case by case basis, but our method serves as a clear guide for how such simulations can be conducted.

Acknowledgements

The author is grateful to Bjarki Eldon, Jochen Blath and Matthias Birkner for their help and suggestions concerning summary statistics for multiple merger detection, and supported by Deutsche Forschungsgemeinschaft (DFG) grant BL 1105/3-2. Part of this work was completed while visiting Matthias Birkner at the Johanner Gutenberg Universität Mainz, funded by DFG grant BI 1058/2-2. Both grants are part of SPP Priority Programme 1590.




Figure 8: 1000 simulated likelihood curves for various values of α , averaging over $L = 23$ loci with lumping at $k = 15$ and sample size $n = 500$. The red dots along the x-axis denote the true, data generating values.

Appendix

A Λ -coalescent for m unlinked loci with $n_i(t)$ lineages at locus $i \in [m]$ can be simulated naively by setting $N := \sum_{i=1}^m n_i(t)$, sampling a merger size K from $\{2, \dots, N\}$ proportional to weights

$$\binom{N}{j} \int_0^1 x^{j-2} (1-x)^{N-j} \Lambda(dx), \quad (6)$$

selecting K lineages uniformly at random without replacement from among the N possible segments, and merging all groups of lineages which occupy the same locus, and incrementing time by a step drawn from an exponential distribution with rate given by the sum of the weights (6). The remaining sample size N can then be updated, and the process iterated until the most recent common ancestor (MRCA) is reached, at which point the algorithm terminates. Diploid, biparental Ξ -coalescent models developed in [Birkner et al., 2013a] can also be incorporated by further randomly assigning all lineages into 4 groups, and only merging lineages which occupy




Figure 9: Box plots of 1000 realisations of maximum likelihood estimators as a function of the true α when $L = 23$, $k = 15$ and $n = 500$ as in Figure 8.




Figure 10: The analysis in Figure 9 repeated using a mutation rate that is too low (left) or high (right) by a factor of ten.

the same locus and the same group.

However, normalising the weights (6) is an $O(N)$ operation, which is expensive when the sample size and number of loci are large. Moreover, the resulting merger is unlikely to sample two or more lineages occupying the same locus unless $\Lambda(dx)$ is concentrated near 1 (resulting in very large mergers), so that most steps of the algorithm result in no progress towards the MRCA. In practice, this naive method was infeasibly slow for our simulation-based model selection. Here we present a superior alternative based on rejection sampling.

The following is a modification of the Poisson construction of Ξ -coalescents given in [Birkner et al., 2009, Section 1.4] to the present multi-locus, biparental, diploid Ξ -coalescents. Consider

a Poisson-Point-Process N on $[0, \infty) \times [0, 1] \times [0, 1]^{\mathbb{N} \times [m]}$ with intensity

$$dt \otimes \frac{\Lambda(dx)}{x^2} \otimes U[0, 1]^{\otimes ([m] \times \mathbb{N})}$$

At each point

$$(t_i, y_i, u_i) \in [0, \infty) \times [0, 1] \times [0, 1]^{[m] \times \mathbb{N}},$$

that is, at time t_i , each of the $n_i(t) \leq n_i$ active lineages at each locus checks whether $u_i(j, k) \leq y_i$ for each $k \in [m], j \in [n_i(t)]$. All lineages for which this is the case take part in a merger within the locus. These merging lineages are the uniformly at random assigned into one of four groups, and any groups with more than one member are merged into a common ancestor.

A technical problem from the simulation point of view is that the intensity measure of our Poisson-Point-Process N is infinite for $\Lambda(dx)$ corresponding to a Beta(β_1, β_2)-coalescent, and hence also for a Beta($2 - \alpha, \alpha$)- Ξ -coalescent for any $\alpha \geq 0$. However, if we restrict N to the set where $u_i(j, k) \leq y_i$ for at least one $j \in [m]$ and two distinct k s from $[n_j(t)]$, the restricted intensity is finite, and given by

$$\int_0^1 f(x; n_1(t), \dots, n_m(t)) \Lambda(dx),$$

where

$$f(x; n_1(t), \dots, n_m(t)) := \frac{1 - \prod_{j=1}^m ((1-x)^{n_j(t)} + n_j(t)x(1-x)^{n_j(t)-1})}{x^2} \quad (7)$$

This restricted intensity corresponds to thinning N by only retaining those events in which there is at least one locus in which two lineages successfully participate in the event.

An alternative simulation mechanism is thus as follows: Given $n_1(t), \dots, n_m(t)$, consider a new *thinned* Poisson-Point-Process \tilde{N} with intensity measure

$$dt \otimes \tilde{\Lambda}(dx), \quad (8)$$

where $\tilde{\Lambda}(dx) := f(x; n_1(t), \dots, n_m(t)) \Lambda(dx)$ determines the merging times and success probabilities (t_i, y_i) . Given (t_i, y_i) , which now appear with finite rate, independently and uniformly choose a *pair* of lineages within a locus to merge out of the

$$\binom{n_1(t)}{2} + \dots + \binom{n_m(t)}{2} =: c(t) \quad (9)$$

within-locus pairs. Further mergers or merging lineages can then be added according to independent uniform coin flips with success probability y_i . As before, all merging lineages within a locus have to further be grouped into four uniform classes in the diploid case, and only lineages within the same locus and class can merge. If no class contains at least two lineages, then nothing happens.

It remains to specify a way of sampling the pairs (t_i, y_i) with intensity (8). Note that the function $f(x; n_1(t), \dots, n_m(t))$ is maximised at $x = 0+$ for fixed $(n_1(t), \dots, n_m(t))$, where it takes the value $c(t)$ from (9). Thus, pairs with the correct intensity can be generated by sampling a time $T^{(1)} \sim \text{Exp}(c(t))$, an independent $U^{(1)} \sim U[0, 1]$ and a further independent $Y^{(1)} \sim \Lambda$, and by checking whether

$$U^{(1)} \leq \frac{f(Y^{(1)}; n_1(t), \dots, n_m(t))}{c(t)}.$$

If yes, accept $(T^{(1)}, Y^{(1)})$ for (t_1, y_1) . If not, repeat this procedure, generating independent tuples $(T^{(2)}, U^{(2)}, Y^{(2)})$ etc. If the first success requires k steps, accept $(T^{(1)} + \dots + T^{(k)}, Y^{(k)})$ for (t_1, y_1) , at which point the merger event can be simulated and $n_1(t_1), \dots, n_m(t_1), f(\cdot; n_1(t), \dots, n_m(t))$

and $c(t)$ updated accordingly. This procedure is then repeated to obtain (t_2, y_2) , and so on, until the most recent common ancestor is reached.

For x close to 0, the function $f(x; n_1(t), \dots, n_m(t))$ is numerically unstable because of the very small number in the denominator in (7), rendering the procedure outlined above unreliable. Note also that it is in fact a polynomial in x and for small arguments, readily represented via

$$f(x; n_1(t), \dots, n_m(t)) = \sum_{j=1}^m \binom{n_j(t)}{2} - 2 \sum_{j=1}^m \binom{n_j(t)}{3} x + O(x^2), \quad (10)$$

as can be verified as follows. For $n \in \mathbb{N}$ we have

$$\begin{aligned} (1-x)^n + nx(1-x)^{n-1} &= 1 + \sum_{k=1}^n \binom{n}{k} (-x)^k + \sum_{j=0}^{n-1} n \binom{n-1}{j} (-1)^j x^{j+1} \\ &= 1 + \sum_{k=2}^n (-x)^k \left[\binom{n}{k} - n \binom{n-1}{k-1} \right] = 1 - \sum_{k=2}^n (k-1) \binom{n}{k} (-x)^k \end{aligned} \quad (11)$$

where we have used the fact that the terms for $k=1$ and for $j=0$ in the first line cancel, then replaced $j=k-1$ in the second sum in the first line and used

$$\binom{n}{k} - n \binom{n-1}{k-1} = \frac{n!}{k!(n-k)!} - \frac{n(n-1)!}{(k-1)!(n-k)!} = \binom{n}{k} (1-k)$$

in the second line. Thus

$$\begin{aligned} \prod_{j=1}^m \left((1-x)^{n_j(t)} + n_j(t)x(1-x)^{n_j(t)-1} \right) &= \prod_{j=1}^m \left(1 - \binom{n_j(t)}{2} x^2 + 2 \binom{n_j(t)}{3} x^3 + O(x^4) \right) \\ &= 1 - \sum_{j=1}^m \binom{n_j(t)}{2} x^2 + 2 \sum_{j=1}^m \binom{n_j(t)}{3} x^3 + O(x^4) \end{aligned}$$

and subtracting from 1 before dividing by x^2 to obtain $f(x; n_1(t), \dots, n_m(t))$ yields (10).

Numerical experiments were performed to determine that the polynomial approximation (10) needed to be used whenever $x \leq 10^{-4}$. The form of higher order terms in (11) shows that the constant of proportionality of the $O(x^k)$ term in (10) contains factors that are $O(n^k)$. Hence, the algorithm cannot remain accurate for samples of size $n = 10^4$, as then approximation errors are $O(1)$ for a polynomial approximation of *any* order, short of the trivial case of using the full, finite expansion which recovers the function $f(x; n_1(t), \dots, n_m(t))$ exactly. We remark that using the full polynomial expansion of $f(x; n_1(t), \dots, n_m(t))$ to compute it exactly is also prone to numerical instability due to alternating signs. However, the first order truncation shown in (10) was found to be fast, accurate and stable for the sample sizes considered in this paper, which we have also shown to be sufficient for our aim of distinguishing multiple merger coalescents from exponential or algebraic population growth models.

References

- E Árnason. Mitochondrial cytochrome *b* variation in the high-fecundity Atlantic cod: trans-Atlantic clines and shallow gene genealogy. *Genetics*, 166:1871–1885, 2004.
- M A Beaumont. Approximate Bayesian computation in evolution and ecology. *Annual Review of Ecology, Evolution, and Systematics*, 41:379–406, 2010.

- A T Beckenbach. Mitochondrial haplotype frequencies in oysters: neutral alternatives to selection models. In B Golding, editor, *Non-Neutral Evolution*, pages 188–198. Chapman & Hall, New York, 1994.
- M Birkner and J Blath. Computing likelihoods for coalescents with multiple collisions in the infinitely many sites model. *J Math Biol*, 57:435–465, 2008.
- M Birkner, J Blath, M Möhle, M Steinrücken, and J Tams. A modified lockdown construction for the Xi-Fleming-Viot process with mutation and populations with recurrent bottlenecks. *ALEA Lat. Am. J. Probab. Math. Stat.*, 6:25–61, 2009.
- M Birkner, J Blath, and M Steinrücken. Importance sampling for Lambda-coalescents in the infinitely many sites model. *Theor Popul Biol*, 79:155–173, 2011.
- M Birkner, J Blath, and B Eldon. An ancestral recombination graph for diploid populations with skewed offspring distribution. *Genetics*, 193:255–290, 2013a.
- M Birkner, J Blath, and B Eldon. Statistical properties of the site-frequency spectrum associated with Lambda-coalescents. *Genetics*, 195:1037–1053, 2013b.
- M Birkner, H Liu, and A Sturm. A note on coalescent results for diploid exchangeable population models. Preprint, 2017.
- J Blath, M C Cronjäger, B Eldon, and M Hammer. The site-frequency spectrum associated with Ξ -coalescents. *Theor. Popul. Biol.*, 110:36–50, 2016.
- F Depaulis and M Veuille. Neutrality tests based on the distribution of haplotypes under an infinite-site model. *Molecular Biology and Evolution*, 15(12):1788, 1998.
- P J Diggle and R J Gratton. Monte Carlo methods of inference for implicit statistical models. *J. R. Stat. Soc. B*, 46(2):193–227, 1984.
- P Donnelly and T G Kurtz. Particle representations for measure-valued population models. *Ann Probab*, 27:166–205, 1999.
- P Donnelly and S Tavaré. Coalescents and genealogical structure under neutrality. *Ann Rev Genet*, 29:401–421, 1995.
- T Duong and M L Hazelton. Plug-in bandwidth matrices for bivariate kernel density estimation. *J. Nonparametr. Statist.*, 15:17–30, 2003.
- R Durrett and J Schweinsberg. A coalescent model for the effect of advantageous mutations on the genealogy of a population. *Stoch Proc Appl*, 115:1628–1657, 2005.
- B Eldon. Estimation of parameters in large offspring number models and ratios of coalescence times. *Theor Popul Biol*, 80:16–28, 2011.
- B Eldon and J Wakeley. Coalescent processes when the distribution of offspring number among individuals is highly skewed. *Genetics*, 172:2621–2633, 2006.
- B Eldon and J Wakeley. Coalescence times and F_{ST} under a skewed offspring distribution among individuals in a population. *Genetics*, 181:615–629, 2009.
- B Eldon, M Birkner, J Blath, and F Freund. Can the site frequency spectrum distinguish exponential population growth from multiple-merger coalescents. *Genetics*, 199(3):841–856, 2015.
- J C Fay and C-I Wu. Hitchhiking under positive Darwinian selection. *Genetics*, 155(3):1405–1413, 2000.
- Y X Fu. Statistical properties of segregating sites. *Theor Popul Biol*, 48:172–197, 1995.

- Y X Fu and W H Li. Statistical tests of neutrality of mutations. *Genetics*, 133:693–709, 1993.
- D Hedgecock and A I Pudovkin. Sweepstakes reproductive success in highly fecund marine fish and shellfish: a review and commentary. *Bull Mar Sci*, 87:971–1002, 2011.
- J Hein, M H Schierup, and Carsten Wiuf. *Gene genealogies, variation and evolution*. Oxford University Press, Oxford, UK, 2005.
- R R Hudson. Properties of a neutral allele model with intragenic recombination. *Theor Popul Biol*, 23:183–201, 1983a.
- R R Hudson. Testing the constant-rate neutral allele model with protein sequence data. *Evolution*, 37:203–217, 1983b.
- R R Hudson. Gene genealogies and the coalescent process. In D J Futuyma and J Antonovics, editors, *Oxford surveys in evolutionary biology*, volume 7, pages 1–44. Oxford University Press, Oxford, 1990.
- J F C Kingman. The coalescent. *Stoch Proc Appl*, 13:235–248, 1982a.
- J F C Kingman. Exchangeability and the evolution of large populations. In G Koch and F Spizzichino, editors, *Exchangeability in probability and statistics*, pages 97–112. North-Holland, Amsterdam, 1982b.
- J F C Kingman. On the genealogy of large populations. *J Appl Probab*, 19A:27–43, 1982c.
- J Koskela, P Jenkins, and D Spanò. Computational inference beyond Kingman’s coalescent. *J. Appl. Probab.*, 52(2):519–537, 2015.
- J Koskela, P Jenkins, and D Spanò. Bayesian non-parametric inference for Λ -coalescents: posterior consistency and a parametric method. *Preprint*, arXiv:1512.00982, 2017.
- M Möhle. Robustness results for the coalescent. *J Appl Probab*, 35:438–447, 1998.
- M Nordborg. Coalescent theory. In D J Balding, M J Bishop, and C Cannings, editors, *Handbook of statistical genetics*, chapter 25, pages 179–212. John Wiley & Sons, Chichester, UK, 2nd edition, 2001.
- J Pitman. Coalescents with multiple collisions. *Ann Probab*, 27:1870–1902, 1999.
- S E Ramos-Onsins and J Rozas. Statistical properties of new neutrality tests against population growth. *Mol Biol Evol*, 19:2092–2100, 2002.
- S Sagitov. The general coalescent with asynchronous mergers of ancestral lines. *J Appl Probab*, 36:1116–1125, 1999.
- O Sargsyan and J Wakeley. A coalescent process with simultaneous multiple mergers for approximating the gene genealogies of many marine organisms. *Theor Popul Biol*, 74:104–114, 2008.
- J Schweinsberg. Coalescent processes obtained from supercritical Galton-Watson processes. *Stoch Proc Appl*, 106:107–139, 2003.
- D W Scott. *Multivariate density estimation: theory, practice and visualization*. John Wiley & Sons, 1992.
- M Steinrücken, M Birkner, and J Blath. Analysis of DNA sequence variation within marine species using beta-coalescents. *Theor Popul Biol*, 87:15–24, 2013.
- F Tajima. Evolutionary relationship of DNA sequences in finite populations. *Genetics*, 105: 437–460, 1983.

- F Tajima. The effect of change in population size on DNA polymorphism. *Genetics*, 123: 597–601, 1989.
- A Tellier and C Lemaire. Coalescence 2.0: a multiple branching of recent theoretical developments and their applications. *Mol Ecol*, 23:2637–2652, 2014.
- O K Tørresen, B Star, S Jentoft, W B Reinart, H Grove, J R Miller, B P Walenz, J Knight, J M Ekholm, P Peluso, R B Edvardsen, A Tooming-Klunderud, M Skage, S Lien, K S Jakobsen, and A J Nederbragt. An improved genome assembly uncovers prolific tandem repeats in Atlantic cod. *bioRxiv*, preprint, 2017.
- J Wakeley. *Coalescent theory*. Roberts & Co, Greenwood Village, 2007.
- G A Watterson. On the number of segregating sites in genetical models without recombination. *Theor Pop Biol*, 7:1539–1546, 1975.
- S Zhu, J H Degnan, S J Goldstein, and B Eldon. Hybrid-Lambda: simulation of multiple merger and Kingman gene genealogies in species networks and species trees. *BMC Bioinformatics*, 16(292), 2015.