# CIRCULANT q-BUTSON HADAMARD MATRICES

TREVOR HYDE AND JOSEPH KRAISLER

ABSTRACT. If  $q = p^n$  is a prime power, then a *d*-dimensional *q*-Butson Hadamard matrix *H* is a  $d \times d$  matrix with all entries *q*th roots of unity such that  $HH^* = dI_d$ . We use algebraic number theory to prove a strong constraint on the dimension of a circulant *q*-Butson Hadamard matrix when  $d = p^m$  and then explicitly construct a family of examples in all possible dimensions. These results relate to the long-standing circulant Hadamard matrix conjecture in combinatorics.

#### 1. INTRODUCTION

For a prime power  $q = p^n$ , a q-Butson Hadamard matrix (q-BH) of dimension d is a  $d \times d$  matrix H with all entries qth roots of unity such that

$$HH^* = dI_d$$

where  $H^*$  is the conjugate transpose of H. A  $d \times d$  matrix H is said to be *circulant* if

 $H_{ij} = f(i-j)$ 

for some function f defined modulo d. In this paper we investigate circulant q-BH matrices of dimension  $d = p^m$ .

**Theorem 1.** If  $q = p^n$  is a prime power and  $d = p^m$ , then there exists a  $d \times d$  circulant q-Butson Hadamard matrix if and only if  $m \le 2n$ , with one exception when (m, n, p) = (1, 1, 2).

Our analysis of circulant q-BH matrices led us to the useful notion of fibrous functions.

**Definition 2.** Let  $d \ge 0$  and  $q = p^n$  be a prime power,

- (1) If X is a finite set, we say a function  $g: X \to \mathbb{Z}/(q)$  is **fibrous** if the cardinality of the fiber  $|g^{-1}(b)|$  depends only on b mod  $p^{n-1}$ .
- (2) We say a function  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$  is  $\delta$ -fibrous if for each  $k \not\equiv 0 \mod d$  the function  $\delta_k(x) = f(x+k) f(x)$  is fibrous.

When q = d = p are both prime,  $\delta$ -fibrous functions coincide with the concept of *planar functions*, which arise in the study of finite projective planes [2] and have applications in cryptography [6]. Circulant q-BH matrices of dimension d are equivalent to  $\delta$ -fibrous functions  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$ .

**Theorem 3.** Let  $q = p^n$  be a prime power and  $\zeta$  a primitive qth root of unity. There is a correspondence between  $d \times d$  circulant q-Butson Hadamard matrices H and  $\delta$ -fibrous functions  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$  given by

$$(H_{ij}) = (\zeta^{f(i-j)}).$$

We restate our main result in the language of  $\delta$ -fibrous functions.

Date: March 14th, 2017.

**Corollary 4.** If  $q = p^n$  is a prime power, then there exist  $\delta$ -fibrous functions  $f : \mathbb{Z}/(p^m) \to \mathbb{Z}/(q)$  if and only if  $m \leq 2n$  with one exception when (m, n, p) = (1, 1, 2).

It would be interesting to know if  $\delta$ -fibrous functions have applications to finite geometry or cryptography.

When q = 2, q-BH matrices are called *Hadamard matrices*. Hadamard matrices are usually defined as  $d \times d$  matrices H with all entries  $\pm 1$  such that  $HH^t = dI_d$ . Buston Hadamard matrices were introduced in [1] as a generalization of Hadamard matrices.

Circulant Hadamard matrices arise in the theory of difference sets, combinatorial designs, and synthetic geometry [7, Chap. 9]. There are arithmetic constraints on the possible dimension of a circulant Hadamard matrix. When the dimension  $d = 2^m$  is a power of two we have:

**Theorem 5** (Turyn [9]). If  $d = 2^m$  is the dimension of a circulant Hadamard matrix, then m = 0 or m = 2.

Turyn's proof uses algebraic number theory, more specifically the fact that 2 is totally ramified in the  $2^n$ th cyclotomic extension  $\mathbb{Q}(\zeta_{2^n})/\mathbb{Q}$ ; an elementary exposition may be found in Stanley [8]. Conjecturally this accounts for all circulant Hadamard matrices [7].

#### **Conjecture 6.** There are no d-dimensional circulant Hadamard matrices for d > 4.

Circulant q-Butson Hadamard matrices provide a natural context within which to consider circulant Hadamard matrices. A better understanding of the former could lead to new insights on the latter. For example, our proof of Theorem 1 shows that the two possible dimensions for a circulant Hadamard matrix given by Turyn's theorem belong to larger family of circulant q-BH matrices with the omission of d = 2 being a degenerate exception. Circulant q-BH matrices have been studied in some specific dimensions [3], but overall seem poorly understood. We leave the existence of q-BH matrices when the dimension d is not a power of p for future work.

Acknowledgements. The authors thank Padraig Cathain for pointers to the literature, in particular for bringing the work of de Launey [4] to our attention. We also thank Jeff Lagarias for helpful feedback on an earlier draft.

## 2. MAIN RESULTS

We always let  $q = p^n$  denote a prime power. First recall the definitions of q-Butson Hadamard and circulant matrices.

**Definition 7.** A d-dimensional q-Butson Hadamard matrix H (q-BH) is a  $d \times d$  matrix all of whose entries are qth roots of unity satisfying

$$HH^* = dI_d,$$

where  $H^*$  is the conjugate transpose of H.

A d-dimensional circulant matrix C is a  $d \times d$  matrix with coefficients in a ring R such that

$$C_{ij} = f(i-j)$$

for some function  $f : \mathbb{Z}/(d) \to R$ .

**Example 8.** Hadamard matrices are the special case of q-BH matrices with q = 2. The q-Fourier matrix  $(\zeta^{ij})$  where  $\zeta$  is a primitive qth root of unity is an example of a q-BH matrix (which may

also be interpreted as the character table of the cyclic group  $\mathbb{Z}/(q)$ .) When q = 3 and  $\omega$  is a primitive 3rd root of unity this is the matrix:

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega^4 \end{pmatrix}$$

This example is not a *circulant* 3-BH matrix. The following is a circulant 3-BH matrix:

$$\begin{pmatrix} 1 & \omega & \omega \\ \omega & 1 & \omega \\ \omega & \omega & 1 \end{pmatrix}$$

The remainder of the paper is divided into two sections: first we prove constraints on the dimension d of a circulant q-BH matrix when d is a power of p; next we introduce the concept of  $\delta$ -fibrous functions and construct examples of circulant q-BH matrices in all possible dimensions.

**Constraints on dimension.** Theorem 9 uses the ramification of the prime *p* in the *q*th cyclotomic extension  $\mathbb{Q}(\zeta)/\mathbb{Q}$  to deduce strong constraints on the dimension of a *q*-BH matrix.

**Theorem 9.** If  $q = p^n$  is a prime power and H is a circulant q-Butson Hadamard matrix of dimension  $d = p^{m+n}$ , then  $m \le n$ .

Note that our indexing of m has changed from the introduction; this choice was made to improve notation in our proof. We use this indexing for the rest of the paper.

*Proof of Theorem 9.* Suppose  $H = (a_{i-j})$  is a circulant q-Butson Hadamard matrix of dimension  $d = p^{m+n}$ . From H being q-BH of dimension d we have

$$HH^* = dI_d$$

hence  $det(H) = \pm d^{d/2}$  and each eigenvalue  $\alpha$  of H has absolute value  $|\alpha| = \sqrt{d} = p^{(m+n)/2}$ . On the other hand, since  $H = (a_{i-j})$  is circulant, it has eigenvalues

$$\alpha_k = \sum_{j < d} a_j \zeta^{jk}.$$
(1)

for  $\zeta$  a primitive dth root of unity with corresponding eigenvector

$$u_k^t = (1, \zeta^k, \zeta^{2k}, \dots, \zeta^{(d-1)k}).$$

These observations combine to give two ways of computing det(H).

$$\prod_{k} \alpha_k = \det(H) = \pm p^{(m+n)d/2}.$$
(2)

The identity (2) is the essential interaction between the circulant and *q*-BH conditions on *H*. The prime *p* is totally ramified in  $\mathbb{Q}(\zeta)$ , hence there is a unique prime ideal  $\mathfrak{p} \subseteq \mathbb{Z}[\zeta]$  over  $(p) \subseteq \mathbb{Z}$ . Since all  $\alpha_k \in \mathbb{Z}[\zeta]$ , it follows from (2) that  $(\alpha_k) = \mathfrak{p}^{v_k}$  as ideals of  $\mathbb{Z}[\zeta]$  for some  $v_k \ge 0$  and for each *k*. So either  $\alpha_0/\alpha_1$  or  $\alpha_1/\alpha_0$  is an element of  $\mathbb{Z}[\zeta]$ . Say  $\alpha_0/\alpha_1$  is the integral quotient. We noted  $|\alpha_k| = p^{(m+n)/2}$  for each *k*, hence  $|\alpha_0/\alpha_1| = 1$ . The only integral elements of  $\mathbb{Z}[\zeta]$  with absolute value 1 are roots of unity, hence  $\alpha_0/\alpha_1 = \pm \zeta^r$  for some  $r \ge 0$ , hence

$$\alpha_0 = \pm \zeta^r \alpha_1. \tag{3}$$

By (1) we have

$$\alpha_0 = \sum_{j < d} a_j \qquad \qquad \alpha_1 = \sum_{j < d} a_j \zeta^j.$$

Each  $j < d = p^{m+n}$  has a unique expression as  $j = j_0 + j_1 p^m$  where  $j_0 < p^m$  and  $j_1 < p^n$ . Let  $\omega = \zeta^{p^m}$  be a primitive *q*th root of unity. Then

$$\zeta^{j} = \zeta^{j_0 + j_1 p^m} = \omega^{j_1} \zeta^{j_0}.$$

Writing  $\alpha_1$  in the linear basis  $\{1, \zeta, \zeta^2, \dots, \zeta^{p^m-1}\}$  of  $\mathbb{Q}(\zeta)/\mathbb{Q}(\omega)$  we have

$$\alpha_1 = \sum_{j < d} a_j \zeta^j = \sum_{j' < p^m} b_{j'} \zeta^{j'},$$

where  $b_{j'}$  is a sum of  $p^n$  complex numbers each with absolute value 1. Now (3) says  $\alpha_0 = \pm \omega^{j_0} \zeta^{-j_1} \alpha_1$  for some  $j_0$  and  $j_1$ , thus

$$\sum_{j < d} a_j = \alpha_0 = \pm \omega^{j_0} \zeta^{-j_1} \alpha_1 = \sum_{j' < p^m} \pm \omega^{j_0} b_{j'} \zeta^{j'-j_1}$$

Comparing coefficients we conclude that

$$\alpha_0 = \pm \omega^{j_0} b_{j_1},$$

which is to say that  $\alpha_0$  is the sum of  $p^n$  complex numbers each with absolute value 1, hence  $|\alpha_0| \leq p^n$ . On the other hand we have  $|\alpha_0| = p^{(m+n)/2}$ . Thus  $m + n \leq 2n \implies 0 \leq m \leq n$  as desired.

*Remark.* The main impediment to extending this result from  $q = p^n$  to a general integer q is that we no longer have the total ramification of the primes dividing the determinant of H. It may be possible to get some constraint in certain cases from a closer analysis of the eigenvalues  $\alpha_k$  and ramification, but we do not pursue this.

 $\delta$ -Fibrous functions and construction of circulant q-BH matrices. Recall the notion of *fibrous functions* from the introduction:

**Definition 10.** Let  $d \ge 0$  and  $q = p^n$  be a prime power,

- (1) If X is a finite set, we say  $g: X \to \mathbb{Z}/(q)$  is fibrous if the cardinality of the fibers  $|g^{-1}(b)|$  depends only on b mod  $p^{n-1}$ .
- (2) We say a function  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$  is  $\delta$ -fibrous if for each  $k \not\equiv 0 \mod d$  the function  $x \mapsto f(x+k) f(x)$  is fibrous.

Lemma 11 is a combinatorial reinterpretation of the cyclotomic polynomials  $\Phi_{p^n}(x)$ .

**Lemma 11.** Let p be a prime and  $\zeta$  a primitive  $p^n$ th root of unity.

- (1) If  $\sum_{k < p^n} b_k \zeta^k = 0$  with  $b_k \in \mathbb{Q}$ , then  $b_k$  depends only on  $k \mod p^{n-1}$ .
- (2) If X is a finite set and  $g: X \to \mathbb{Z}/(p^n)$  is a function, then g is fibrous iff

$$\sum_{x \in X} \zeta^{g(x)} = 0$$

*Proof.* (1) Suppose  $\sum_{k < p^n} b_k \zeta^k = 0$  for some  $b_k \in \mathbb{Q}$ . Then  $r(x) = \sum_{k < p^n} b_k x^k \in \mathbb{Q}[x]$  is a polynomial with degree  $< p^n$  such that  $r(\zeta) = 0$ . So there is some  $s(x) \in \mathbb{Q}[x]$  such that  $r(x) = s(x)\Phi_{p^n}(x)$  where

$$\Phi_{p^n}(x) = \sum_{j < p} x^{jp^{n-1}}$$

is the  $p^n$ th cyclotomic polynomial—the minimal polynomial of  $\zeta$  over  $\mathbb{Q}$ . Since deg  $\Phi_{p^n}(x) = p^n - p^{n-1}$ , it follows that deg  $s(x) < p^{n-1}$ . Let

$$s(x) = \sum_{i < p^{n-1}} a_i x^i$$

for some  $a_i \in \mathbb{Q}$ . Expanding  $s(x)\Phi_{p^n}(x)$  we have

$$r(x) = s(x)\Phi_{p^n}(x) = \sum_{\substack{i < p^{n-1} \\ j < p}} a_i x^{i+jp^{n-1}}.$$

Comparing coefficients yields

$$b_k = b_{i+jp^{n-1}} = a_i,$$

which is to say,  $b_k$  depends only  $i \equiv k \mod p^{n-1}$ .

(2) Suppose g is fibrous. For each  $i < p^{n-1}$ , let  $a_i = |g^{-1}(i)|$ . Then

$$\sum_{x \in X} \zeta^{g(x)} = \sum_{i < p^{n-1}} \sum_{j < p} a_i \zeta^{i+jp^{n-1}} = \Phi_{p^n}(\zeta) \sum_{i < p^{n-1}} a_i \zeta^i = 0.$$

Conversely, for each  $k < p^n$  let  $c_k = |g^{-1}(k)|$ . Then

$$0 = \sum_{x \in X} \zeta^{g(x)} = \sum_{k < p^n} c_k \zeta^k,$$

and (1) implies  $c_k$  depends only on  $k \mod p^{n-1}$ . Hence g is fibrous.

Theorem 12 establishes the equivalence between q-BH matrices of dimension d and  $\delta$ -fibrous functions  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$ .

**Theorem 12.** Let  $q = p^n$  be a prime power. There is a correspondence between circulant *q*-Butson Hadamard matrices H of dimension d and  $\delta$ -fibrous functions  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$  given by

$$(H_{i,j}) = (\zeta^{f(i-j)}).$$

*Proof.* Suppose f is  $\delta$ -fibrous. Define the matrix  $H = (H_{ij})$  by  $H_{ij} = \zeta^{f(i-j)}$  where  $\zeta$  is a primitive qth root of unity. H is plainly circulant and has all entries qth roots of unity. It remains to show that  $HH^* = dI_d$ , which is to say that the inner product  $r_{j,j+k}$  of column j and column j + k is 0 for each j and each  $k \neq 0 \mod d$ . For each  $k \neq 0 \mod d$  the function  $\delta_k(x) = f(x+k) - f(x)$  is fibrous. Then we compute

$$r_{j,j+k} = \sum_{i < d} \zeta^{f(i-j+k) - f(i-j)} = \sum_{i < d} \zeta^{\delta_k(i-j)} = 0,$$

where the last equality follows from Lemma 11 (2).

Conversely, suppose  $H = (H_{i,j})$  is a circulant q-Butson Hadammard matrix. Then  $H_{i,j} = \zeta^{f(i-j)}$  for some function  $f : \mathbb{Z}/(d) \to \mathbb{Z}/(q)$ . Since  $HH^* = dI_d$  we have for each  $k \not\equiv 0 \mod d$ ,

$$0 = r_{j,j+k} = \sum_{i < d} \zeta^{f(i-j+k) - f(i-j)}.$$

Lemma 11 (2) then implies  $\delta_k(x) = f(x+k) - f(x)$  is fibrous. Therefore f is  $\delta$ -fibrous.

Lemma 13 checks that affine functions are fibrous. We use this in our proof of Theorem 14.

**Lemma 13.** If  $q = p^n$  is a prime power, then for all  $a \not\equiv 0 \mod q$  and arbitrary b, the function f(x) = ax + b is fibrous.

*Proof.* Since  $a \not\equiv 0 \mod q$ ,

$$\sum_{j < q} \zeta^{f(j)} = \sum_{j < q} \zeta^{aj+b} = \zeta^b \sum_{j < q} (\zeta^a)^j = 0.$$

Thus, by Lemma 11 (2) we conclude that f is fibrous.

**Theorem 14.** If  $q = p^n$  is a prime power, then there exists a circulant q-Butson Hadamard matrix of dimension  $d = p^{m+n} = p^m q$  for each  $m \le n$  unless (m, n, p) = (0, 1, 2).

Our construction in the proof of Theorem 14 misses the family (m, n, p) = (0, n, 2) for each  $n \ge 1$ . Lemma 15 records a quick observation that circumvents this issue for n > 1, as our construction does give circulant  $2^{n-1}$ -BH matrices of dimension  $2^n$ .

**Lemma 15.** If  $q = p^n$  is a prime power and there exists a circulant q-BH matrix of dimension d, then there exists a circulant  $p^k q$ -BH matrix of dimension d for all  $k \ge 0$ .

*Proof.* Every qth root of unity is also a  $p^k q$ th root of unity, hence we may view a circulant q-BH matrix H of dimension d as a circulant  $p^k q$ -BH for all  $k \ge 0$ .

Proof of Theorem 14. Our strategy is to first construct a sequence of functions

$$\delta_k : \mathbb{Z}/(p^m) \times \mathbb{Z}/(q) \to \mathbb{Z}/(q)$$

which are fibrous for each  $k < p^m q$ . If  $i : \mathbb{Z}/(p^m) \times \mathbb{Z}/(q) \to \mathbb{Z}/(p^m q)$  is the bijection  $i(x, y) = x + p^m y$ , we define  $f : \mathbb{Z}/(p^m q) \to \mathbb{Z}/(q)$  such that  $f(z + k) - f(z) = \delta_k(x, y)$  when z = i(x, y). Hence f is  $\delta$ -fibrous and Theorem 12 implies the existence of a corresponding circulant q-BH matrix of dimension  $p^m q$ .

Now for each  $k \ge 0$  define  $\delta_k$  by

$$\delta_k(x,y) = ky + \sum_{j < k} S_j(x), \tag{4}$$

where

$$S_j(x) = \sum_{i < j} \chi(x+i), \qquad \chi(x) = \begin{cases} 1 & x \equiv -1 \mod p^m, \\ 0 & \text{otherwise.} \end{cases}$$

Observe that  $S_j(x)$  counts the integers in the interval [x, x + j) congruent to  $-1 \mod p^m$  (which only depends on  $x \mod p^m$ .) Any interval of length  $p^m j_1$  contains precisely  $j_1$  integers congruent to  $-1 \mod p^m$ . For each  $j < p^m q$ , write  $j = j_0 + p^m j_1$  with  $j_0 < p^m$  and  $j_1 < q$ , then

$$S_j(x) = \sum_{i < j_0 + p^m j_1} \chi(x+i) = \sum_{i < j_0} \chi(x+i) + j_1 = S_{j_0}(x) + j_1.$$
(5)

We show that  $\delta_k$  is fibrous when  $k < p^m q$ . If  $k \not\equiv 0 \mod q$ , then for each  $x = x_0$  the function  $\delta_k(x_0, y)$  is affine hence fibrous by Lemma 13. So  $\delta_k(x, y)$  is fibrous. Now suppose k = k'q for some  $k' < p^m$ . Using (5) we reduce (4) to

$$\delta_k(x,y) = \sum_{j < k'q} S_j(x) = \sum_{j_0 + p^m j_1 < k'q} S_{j_0}(x) + j_1 = \ell \sum_{j_0 < p^m} S_{j_0}(x) + p^m \binom{\ell}{2}, \tag{6}$$

where  $k'q = p^m(k'p^{n-m}) = p^m\ell$ . Here we use our assumption  $m \le n$ . The definition of  $\chi$  implies

$$S_{j_0}(x) = \begin{cases} 0 & j_0 < p^m - x \\ 1 & j_0 \ge p^m - x, \end{cases} \implies \sum_{j_0 < p^m} S_{j_0}(x) = x,$$

whence  $\delta_k(x, y) = \ell x + p^m \binom{\ell}{2}$ . Since  $k' < p^m$  it follows that  $\ell = k' p^{n-m} < p^n = q$ , so  $\delta_k(x, y)$  is affine hence fibrous by Lemma 13.

Define  $f : \mathbb{Z}/(p^m q) \to \mathbb{Z}/(q)$  by  $f(k) = \delta_k(0,0)$ . For this to be well-defined, it suffices to check that  $\delta_{k+p^m q}(x,y) = \delta_k(x,y)$  with arbitrary k. By (4),

$$\delta_{k+p^{m}q}(x,y) = (k+p^{m}q)y + \sum_{j < k+p^{m}q} S_{j}(x)$$
  
=  $ky + \sum_{j < k} S_{j}(x) + \sum_{j < p^{m}q} S_{j+k}(x)$   
=  $\delta_{k}(x,y) + \sum_{j < p^{m}q} S_{j+k}(x).$ 

The argument leading to (6) gives

$$\sum_{j < p^m q} S_{j+k}(x) = qx + p^m \binom{q}{2} = p^m \binom{q}{2}.$$

Finally,  $p^m \binom{q}{2} \equiv 0 \mod q$  unless (m, n, p) = (0, n, 2). Lemma 15 implies that constructing an example for (1, n - 1, 2) implies the existence of example for (0, n, 2), hence we proceed under the assumption that either  $p \neq 2$  or p = 2 and m > 0. The case (m, n, p) = (0, 1, 2)is an exception as one can check explicitly that there are no 2-dimensional Hadamard matrices. Hence it follows that f is well-defined. Let  $i : \mathbb{Z}/(p^m) \times \mathbb{Z}/(q) \to \mathbb{Z}/(p^m q)$  be the bijection  $i(x, y) = x + p^m y$ . To finish the construction we suppose z = i(x, y), show

$$f(z+k) - f(z) = \delta_k(x,y),$$

and then our proof that  $\delta_k$  is fibrous for all  $k < p^m q$  implies f is  $\delta$ -fibrous. Theorem 12 translates this into the existence of a circulant q-BH matrix of dimension  $p^{m+n}$ . Now,

$$f(z+k) - f(z) = \delta_{z+k}(0,0) - \delta_z(0,0) = \sum_{z \le j < z+k} S_j(0) = \sum_{j-z < k} \sum_{i-z < j-z} \chi(i)$$
$$= \sum_{j' < k} \sum_{i' < j'} \chi(x+i') = \sum_{j' < k} S_{j'}(x) = \delta_k(x,y).$$

*Remark.* Padraig Cathain brought the work of de Launey [4] to our attention after reading an initial draft. There one finds a construction of circulant q-Butson Hadamard matrices of dimension  $q^2$  for all prime powers q which appears to be closely related to our construction in Theorem 14.

**Example 16.** We provide two low dimensional examples to illustrate our construction. First we have an 8 dimensional circulant 4-BH matrix.

( 1	-1	i	1	1	1	i	-1
-1	1	-1	i	1	1	1	i
i	-1	1	-1	i	1	1	1
1	i	-1	1	-1	i	1	1
1	1	i	-1	1	-1	i	1
1	1	1	i	-1	1	-1	i
i	1	1	1	i	-1	1	-1
$\setminus -1$	i	1	1	1	i	-1	1,

Let  $\omega$  be a primitive 3rd root of unity. The following is a 9 dimensional circulant 3-BH matrix.

(1	$\omega^2$	$\omega$	1	1	1	1	$\omega$	$\omega^{2}$
$\omega^2$	1	$\omega^2$	$\omega$	1	1	1	1	$\omega$
$\omega$	$\omega^2$	1	$\omega^2$	$\omega$	1	1	1	1
1	$\omega$	$\omega^2$	1	$\omega^2$	$\omega$	1	1	1
1	1	$\omega$	$\omega^2$	1	$\omega^2$	$\omega$	1	1
1	1	1	$\omega$	$\omega^2$	1	$\omega^2$	$\omega$	1
1	1	1	1	$\omega$	$\omega^2$	1	$\omega^2$	$\omega$
ω	1	1	1	1	$\omega$	$\omega^2$	1	$\omega^2$
$\omega^2$	$\omega$	1	1	1	1	$\omega$	$\omega^2$	1

Corollary 17 is an immediate consequence of our main results by Theorem 12.

**Corollary 17.** If  $q = p^n$  is a prime power and  $d = p^{m+n}$ , then there exists a  $\delta$ -fibrous function  $f : \mathbb{Z}/(p^{m+n}) \to \mathbb{Z}/(q)$  iff  $m \leq n$ , with the one exception of (m, n, p) = (0, 1, 2).

**Closing remarks.** Our analysis focused entirely on the existence of circulant  $p^n$ -BH matrices with dimension d a power of p. The number theoretic method of Theorem 9 cannot be immediately adapted to the case where d is not a power of p, although as we noted earlier, it may be possible to get some constraint with a closer analysis of the eigenvalues of a circulant matrix and the ramification over the primes dividing d in the dth cyclotomic extension  $\mathbb{Q}(\zeta)/\mathbb{Q}$ .

The family of examples constructed in Theorem 14 was found empirically. It would be interesting to know if the construction extends to any dimensions which are not powers of p.

### REFERENCES

- [1] Butson, A.T. "Generalized Hadamard matrices" Proc. Amer. Math. Soc., 13 (1962), pp. 894-898
- [2] Dembowski, Peter, and Thedore G. Ostrom. "Planes of order n with collineation groups of order  $n^2$ ." *Mathematische Zeitschrift* 103.3 (1968): 239-258.
- [3] Hiranandani, Gaurush, and Jean-Marc Schlenker. "Small circulant complex Hadamard matrices of Butson type." *European Journal of Combinatorics* 51 (2016): 306-314.
- [4] de Launey, Warwick. "Circulant  $GH(p^2; \mathbb{Z}_p)$  exist for all primes p." Graphs Combin. 8, no. 4, (1992): 317-321.
- [5] Leung, Ka Hin, and Bernhard Schmidt. "New restrictions on possible orders of circulant Hadamard matrices." *Designs, Codes and Cryptography* 64.1 (2012): 143-151.
- [6] Nyberg, Kaisa, and Lars Ramkilde Knudsen. "Provable security against differential cryptanalysis." Annual International Cryptology Conference. Springer Berlin Heidelberg, 1992.
- [7] Ryser, Herbert John. "Combinatorial mathematics." Vol. 14. Washington, DC: Mathematical Association of America, 1963.
- [8] Stanley, Richard P. "Algebraic combinatorics." Springer 20 (2013): 22.
- [9] Turyn, Richard. "Character sums and difference sets." Pacific Journal of Mathematics 15.1 (1965): 319-346.

DEPT. OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043, *E-mail address*: tghyde@umich.edu

DEPT. OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043, *E-mail address*: jkrais@umich.edu