

# Factors of alternating sums of powers of $q$ -Narayana numbers

Victor J. W. Guo<sup>1</sup> and Qiang-Qiang Jiang<sup>2</sup>

<sup>1</sup>School of Mathematical Sciences, Huaiyin Normal University, Huai'an, Jiangsu 223300, People's Republic of China  
jwguo@hytc.edu.cn

<sup>2</sup>Department of Mathematics, East China Normal University, Shanghai 200241, People's Republic of China  
972126862@qq.com

**Abstract.** The  $q$ -Narayana numbers  $N_q(n, k)$  and  $q$ -Catalan numbers  $C_n(q)$  are respectively defined by

$$N_q(n, k) = \frac{1-q}{1-q^n} \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} n \\ k-1 \end{bmatrix} \quad \text{and} \quad C_n(q) = \frac{1-q}{1-q^{n+1}} \begin{bmatrix} 2n \\ n \end{bmatrix},$$

where  $\begin{bmatrix} n \\ k \end{bmatrix} = \prod_{i=1}^k \frac{1-q^{n-i+1}}{1-q^i}$ . We prove that, for any positive integers  $n$  and  $r$ , there holds

$$\sum_{k=-n}^n (-1)^k q^{jk^2 + \binom{k}{2}} N_q(2n+1, n+k+1)^r \equiv 0 \pmod{C_n(q)},$$

where  $0 \leq j \leq 2r-1$ . We also propose several related conjectures.

*Keywords:*  $q$ -binomial coefficients,  $q$ -Narayana numbers,  $q$ -Catalan numbers

*AMS Subject Classifications:* 11A07, 05A30

## 1 Introduction

The Catalan numbers  $C_n = \frac{1}{n+1} \binom{2n}{n}$  play an important role in combinatorics (see [14]). It is well known that for any positive integer  $n$ ,

$$C_n = \sum_{k=1}^n N(n, k),$$

where  $N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k-1}$  are called Narayana numbers (see [9]).

In the past few years, many congruences on sums or alternating sums of binomial coefficients and combinatorial numbers, such as Catalan numbers, Apéry numbers, central Delannoy numbers, Schröder numbers, Franel numbers, have been obtained by Z.-W. Sun [16–20] and other authors [4–6, 8, 11, 12, 21, 22].

In this paper, motivated mainly by Z.-W. Sun's work, we shall prove the following congruence on alternating sums of powers of Narayana numbers.

**Theorem 1.1.** *Let  $n$  and  $r$  be positive integers. Then*

$$\sum_{k=-n}^n (-1)^k N(2n+1, n+k+1)^r \equiv 0 \pmod{C_n}. \quad (1.1)$$

We know that some congruences may have nice  $q$ -analogues (see, for example, [7, 10, 13]). This is also the case for the congruence (1.1). Recall that the  $q$ -shifted factorials (see [1]) are defined by  $(a; q)_0 = 1$  and  $(a; q)_n = (1-a)(1-aq)\cdots(1-aq^{n-1})$  for  $n = 1, 2, \dots$ , and the  $q$ -binomial coefficients are defined as

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}, & \text{if } 0 \leq k \leq n, \\ 0, & \text{otherwise.} \end{cases}$$

For convenience, we let  $[n] = \frac{1-q^n}{1-q}$  be a  $q$ -integer. It is natural to define the  $q$ -Narayana numbers  $N_q(n, k)$  and the  $q$ -Catalan numbers  $C_n(q)$  as follows:

$$N_q(n, k) = \frac{1}{[n]} \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} n \\ k-1 \end{bmatrix}, \quad C_n(q) = \frac{1}{[n+1]} \begin{bmatrix} 2n \\ n \end{bmatrix}.$$

It is not difficult to see that both  $q$ -Narayana numbers and  $q$ -Catalan numbers are polynomials in  $q$  with nonnegative integer coefficients (see [2, 3]). Note that, the definition of  $N_q(n, k)$  here differs by a factor  $q^{k(k-1)}$  from that in [2]. We have the following  $q$ -analogue of Theorem 1.1.

**Theorem 1.2.** *Let  $n$  and  $r$  be positive integers and let  $0 \leq j \leq 2r-1$ . Then*

$$\sum_{k=-n}^n (-1)^k q^{jk^2 + \binom{k}{2}} N_q(2n+1, n+k+1)^r \equiv 0 \pmod{C_n(q)}. \quad (1.2)$$

It is easily seen that when  $q \rightarrow 1$  the  $q$ -congruence (1.2) reduces to (1.1). It seems that (1.2) also holds for  $j \geq 2r$  (see (3.1) in Conjecture 3.4 for a more general form).

## 2 Proof of Theorem 1.2

Noticing that

$$\begin{bmatrix} 2n+1 \\ n+k \end{bmatrix} \begin{bmatrix} 2n+1 \\ n+k+1 \end{bmatrix} = \frac{(q; q)_{2n+1}^2}{(q; q)_{2n} (q; q)_{2n+2}} \begin{bmatrix} 2n \\ n+k \end{bmatrix} \begin{bmatrix} 2n+2 \\ n+k+1 \end{bmatrix}, \quad (2.1)$$

we can rewrite Theorem 1.2 in the following equivalent form.

**Theorem 2.1.** *Let  $n$  and  $r$  be positive integers and let  $0 \leq j \leq 2r-1$ . Then*

$$\begin{aligned} & \frac{(q; q)_{2n+1}^{2r}}{(q; q)_{2n}^r (q; q)_{2n+2}^r} \sum_{k=-n}^n (-1)^k q^{jk^2 + \binom{k}{2}} \begin{bmatrix} 2n \\ n+k \end{bmatrix}^r \begin{bmatrix} 2n+2 \\ n+k+1 \end{bmatrix}^r \\ & \equiv 0 \pmod{\begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1}}. \end{aligned} \quad (2.2)$$

In the paper [4, Theorem 4.7], Guo, Jouhet and Zeng proved the following result.

**Theorem 2.2.** *For all positive integers  $n_1, \dots, n_m$  and  $0 \leq j \leq m - 1$ , the alternating sum*

$$(q; q)_{n_1} \prod_{i=1}^m \frac{(q; q)_{n_i+n_{i+1}}}{(q; q)_{2n_i}} \sum_{k=-n_1}^{n_1} (-1)^k q^{jk^2+\binom{k}{2}} \prod_{i=1}^m \begin{bmatrix} 2n_i \\ n_i + k \end{bmatrix},$$

where  $n_{m+1} = 0$ , is a polynomial in  $q$  with nonnegative integer coefficients.

In what follows, we will show that the congruence (2.2) can be deduced from combining two special cases of Theorem 2.2.

Denote the left-hand side of (2.2) by  $S(n, r, j)$ . By the relation (2.1), it is clear that  $S(n, r, j)$  is a polynomial in  $q$  with integer coefficients. Letting  $m = 2r$ ,  $n_1 = n_3 = \dots = n_{2r-1} = n$ , and  $n_2 = n_4 = \dots = n_{2r} = n + 1$  in Theorem 2.2, we see that

$$\frac{(q; q)_n (q; q)_{n+1} (q; q)_{2n+1}^{2r-1}}{(q; q)_{2n}^r (q; q)_{2n+2}^r} \sum_{k=-n}^n (-1)^k q^{jk^2+\binom{k}{2}} \begin{bmatrix} 2n \\ n+k \end{bmatrix}^r \begin{bmatrix} 2n+2 \\ n+k+1 \end{bmatrix}^r \in \mathbb{Z}[q],$$

which can also be written as

$$\begin{bmatrix} 2n+1 \\ n \end{bmatrix}^{-1} S(n, r, j) \in \mathbb{Z}[q].$$

Namely,  $S(n, r, j) \equiv 0 \pmod{\begin{bmatrix} 2n+1 \\ n \end{bmatrix}}$ , or

$$[2n+1]^{r-1} S(n, r, j) \equiv 0 \pmod{\begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1}}. \quad (2.3)$$

On the other hand, letting  $m = 2r$ ,  $n_1 = n_2 = \dots = n_r = n$ , and  $n_{r+1} = n_{r+2} = \dots = n_{2r} = n + 1$  in Theorem 2.2, we have

$$\frac{(q; q)_n (q; q)_{n+1} (q; q)_{2n+1} (q; q)_{2n}^{r-1} (q; q)_{2n+2}^{r-1}}{(q; q)_{2n}^r (q; q)_{2n+2}^r} \sum_{k=-n}^n (-1)^k q^{jk^2+\binom{k}{2}} \begin{bmatrix} 2n \\ n+k \end{bmatrix}^r \begin{bmatrix} 2n+2 \\ n+k+1 \end{bmatrix}^r \in \mathbb{Z}[q],$$

which, by the relation  $(q; q)_{2n} (q; q)_{2n+2} = (q; q)_{2n+1}^2 \frac{[2n+2]}{[2n+1]}$ , can be rewritten as

$$\begin{bmatrix} 2n+1 \\ n \end{bmatrix}^{-1} \frac{[2n+2]^{r-1}}{[2n+1]^{r-1}} S(n, r, j) \in \mathbb{Z}[q].$$

Namely,

$$[2n+2]^{r-1} S(n, r, j) \equiv 0 \pmod{\begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1}}. \quad (2.4)$$

It is easy to see that the polynomials  $[2n+1]^{r-1}$  and  $[2n+2]^{r-1}$  are relatively prime. Therefore, by the Euclid algorithm for polynomials, there exist polynomials  $P(q)$  and  $Q(q)$  in  $q$  with rational coefficients such that

$$P(q)[2n+1]^{r-1} + Q(q)[2n+2]^{r-1} = 1. \quad (2.5)$$

It follows from (2.3)–(2.5) that the congruence

$$S(n, r, j) \equiv 0 \pmod{\begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1}}.$$

holds in the ring  $\mathbb{Q}[q]$ . In other words, there exists a polynomial  $R(q) \in \mathbb{Q}[q]$  such that

$$S(n, r, j) = \begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1} R(q). \quad (2.6)$$

Since the polynomials  $S(n, r, j)$  and  $\begin{bmatrix} 2n+1 \\ n \end{bmatrix} [2n+1]^{r-1}$  are in  $\mathbb{Z}[q]$ , and the leading coefficient of the latter is one, the identity (2.6) means that  $R(q) \in \mathbb{Z}[q]$ . This completes the proof.

### 3 Some open problems

It seems that Theorems 1.1 and 1.2 can be further generalized as follows.

**Conjecture 3.1.** *Let  $n_1, \dots, n_m$  be positive integers. Then*

$$\begin{aligned} & \sum_{k=-n_1}^{n_1} (-1)^k \prod_{i=1}^m \binom{n_i + n_{i+1} + 1}{n_i + k} \binom{n_i + n_{i+1} + 1}{n_i + k + 1} \\ & \equiv 0 \pmod{\binom{n_1 + n_m + 1}{n_1} \prod_{i=1}^{m-1} (n_i + n_{i+1} + 1)}, \end{aligned}$$

where  $n_{m+1} = n_1$ .

**Conjecture 3.2.** *Let  $n$  and  $r$  be positive integers and let  $0 \leq j \leq 2r - 1$ . Then*

$$\frac{1}{C_n(q)} \sum_{k=-n}^n (-1)^k q^{jk^2 + \binom{k}{2}} N_q(2n+1, n+k+1)^r$$

is a polynomial in  $q$  with nonnegative integer coefficients.

Note that the upper bound  $2r - 1$  of  $j$  in Conjecture 3.2 seems to be the best possible. Numerical calculation implies that Conjecture 3.2 does not hold when  $j \geq 2r$ . Furthermore, we have the following generalization of Conjecture 3.2.

**Conjecture 3.3.** *For all positive integers  $n_1, \dots, n_m$  and  $0 \leq j \leq 2m - 1$ , the expression*

$$\begin{bmatrix} n_1 + n_m + 1 \\ n_1 \end{bmatrix}^{-1} \prod_{i=1}^{m-1} \frac{1}{[n_i + n_{i+1} + 1]} \sum_{k=-n_1}^{n_1} (-1)^k q^{jk^2 + \binom{k}{2}} \prod_{i=1}^m \begin{bmatrix} n_i + n_{i+1} + 1 \\ n_i + k \end{bmatrix} \begin{bmatrix} n_i + n_{i+1} + 1 \\ n_i + k + 1 \end{bmatrix},$$

where  $n_{m+1} = n_1$ , is a polynomial in  $q$  with nonnegative integer coefficients.

We end the paper with the following  $q$ -analogue of Conjecture 3.1.

**Conjecture 3.4.** Let  $n_1, \dots, n_m$  be positive integers, and let  $f(k)$  be a polynomial in  $k$  with integer coefficients. Then

$$\sum_{k=-n_1}^{n_1} (-1)^k q^{f(k)+\binom{k}{2}} \prod_{i=1}^m \begin{bmatrix} n_i + n_{i+1} + 1 \\ n_i + k \end{bmatrix} \begin{bmatrix} n_i + n_{i+1} + 1 \\ n_i + k + 1 \end{bmatrix} \\ \equiv 0 \pmod{\begin{bmatrix} n_1 + n_m + 1 \\ n_1 \end{bmatrix} \prod_{i=1}^{m-1} [n_i + n_{i+1} + 1]},$$

where  $n_{m+1} = n_1$ . In particular, for any positive integer  $n$ , we have

$$\sum_{k=-n}^n (-1)^k q^{f(k)+\binom{k}{2}} N_q(2n+1, n+k+1)^r \equiv 0 \pmod{C_n(q)}. \quad (3.1)$$

**Acknowledgments.** This work was partially supported by the National Natural Science Foundation of China (grant 11371144), the Natural Science Foundation of Jiangsu Province (grant BK20161304), and the Qing Lan Project of Education Committee of Jiangsu Province.

## References

- [1] G.E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [2] P. Brändén,  $q$ -Narayana numbers and the flag  $h$ -vector of  $J(2 \times n)$ , *Discrete Math.* 281 (2004), 67–81.
- [3] J. Förlinger and J. Hofbauer,  $q$ -Catalan numbers, *J. Combin. Theory, Ser. A* 2 (1985), 248–264.
- [4] V.J.W. Guo, F. Jouhet, J. Zeng, Factors of alternating sums of products of binomial and  $q$ -binomial coefficients, *Acta Arith.* 127 (2007), 17–31.
- [5] V.J.W. Guo and J. Zeng, New congruences for sums involving Apéry numbers or central Delannoy numbers, preprint, *Int. J. Number Theory* 8 (2012), 2003–2016.
- [6] V.J.W. Guo and J. Zeng, Proof of some conjectures of Z.-W. Sun on congruences for Apéry polynomials, *J. Number Theory* 132 (2012), 1731–1740.
- [7] V.J.W. Guo and J. Zeng, Some  $q$ -analogues of supercongruences of Rodriguez-Villegas, *J. Number Theory* 145 (2014), 301–316.
- [8] J.-C. Liu, A supercongruence involving Delannoy numbers and Schröder numbers, *J. Number Theory* 168 (2016), 117–127.
- [9] T.V. Narayana, A partial order and its application to probability, *Sankhyā* 21 (1959), 91–98.
- [10] H. Pan, A  $q$ -analogue of Lehmer’s congruence, *Acta Arith.* 128 (2007), 303–318.
- [11] H. Pan, On divisibility of sums of Apéry polynomials, *J. Number Theory* 143 (2014), 214–223.

- [12] H. Pan and Z.-W. Sun, A combinatorial identity with application to Catalan numbers, *Discrete Math.* 306 (2006), 1921–1940.
- [13] L.-L. Shi and H. Pan, A  $q$ -analogue of Wolstenholme’s harmonic series congruence, *Amer. Math. Monthly* 114 (2007), 529C531.
- [14] R.P. Stanley, *Enumerative Combinatorics, Vol. II*, Cambridge University Press, Cambridge, 1999.
- [15] Z.-W. Sun, On Delannoy numbers and Schröder numbers, *J. Number Theory* 131 (2011), 2387–2397.
- [16] Z.-W. Sun, On sums of Apéry polynomials and related congruences, *J. Number Theory*, 132 (2012), 2673–2699.
- [17] Z.-W. Sun, Connections between  $p = x^2 + 3y^2$  and Franel numbers, *J. Number Theory* 133 (2013), 2914–2928.
- [18] Z.-W. Sun, Supercongruences involving products of two binomial coefficients, *Finite Fields Appl.* 22 (2013), 24–44.
- [19] Z.-W. Sun, Congruences for Franel numbers, *Adv. Appl. Math.* 51 (2013), 524–535.
- [20] Z.-W. Sun, Congruences involving generalized trinomial coefficients, *Sci. China* 57 (2014), 1375–1400.
- [21] Z.-W. Sun and R. Tauraso, New congruences for central binomial coefficients, *Adv. Appl. Math.* 45 (2010), 125–148.
- [22] Z.-W. Sun and R. Tauraso, On some new congruences for binomial coefficients, *Int. J. Number Theory* 7(2011), 645–662.