Lectures on Structural Stability in Dynamics

Pierre Berger*

Abstract

These lectures present results and problems on the characterization of structurally stable dynamics. We will shed light those which do not seem to depend on the regularity class (holomorphic or differentiable). Furthermore, we will present some links between the problems of structural stability in dynamical systems and in singularity theory.

Contents

1	Uniformly hyperbolic dynamical systems	2
2	Properties of structurally stable dynamics	7
3	Hyperbolicity implies structural stability	11
4	Links between structural stability in dynamical systems and singularity theory	20
5	Structural stability of endomorphisms with singularities	24

Introduction

Structural stability is one of the most basic topics in dynamical systems and contains some of the hardest conjectures. Given a class of regularity \mathcal{C} , which can be C^r for $1 \leq r \leq \infty$ or holomorphic, and formed by diffeomorphisms or endomorphisms of a manifold M, the problem is to describe the structurally stable dynamics for the class \mathcal{C} . We recall that a dynamics f is \mathcal{C} -structurally stable if for any perturbation \hat{f} of f in the class \mathcal{C} , there exists a homeomorphism h of M so that $h \circ f = f \circ h$. Uniform hyperbolicity seems to provide a satisfactory way to describe the structurally stable dynamics. This observation goes back to the Fatou conjecture for quadratic maps of the Riemannian sphere in 1920 and the Smale conjecture for smooth diffeomorphisms in 1970. These conjectures have been deeply studied by many mathematicians and so they are difficult to tackle directly.

However at the interface of one-dimensional complex dynamics and differentiable dynamics, the field of two-dimensional complex dynamics grew up recently. It enables to study the structural

^{*}CNRS-LAGA, Université Paris 13.

stability problem thanks to ingredients of both fields. Also the mathematics developed in the 1970's for the structural stability in dynamics is very similar to the one developed for the structural stability in singularity theory. This led us to combine both in the study of the structurally stable endomorphisms, We will review some classical works in these beautiful fields, some works more recent, and we will present new open problems at these interfaces.

In section 1, we will recall some elementary definitions of uniform hyperbolic theory, and we will detail a few examples of such dynamics.

In section 2 we will state several theorems and conjectures suggesting the hyperbolicity of structurally stable dynamics. In particular we will recall the seminal work of Mañé [Mañ88] showing this direction in the C^1 -category. For holomorphic dynamical systems, we will present the work of Dujardin-Lyubich [DL] and our work with Dujardin [BD17] generalizing some aspects of Mañé-Sad-Sullivan and Lyubich theorems [MSS83, Lyu84] for polynomial automorphisms of \mathbb{C}^2 .

In section 3, we will present several results in the directions "hyperbolicty \Rightarrow stability". In §3.1, we will recall classical results, including the structural stability theorems of Anosov [Ano67], Moser[Mos69] and Shub [Shu69], and the proof of this direction of the Ω -stability conjecture by Smale[Sma70] and Przytycki [Prz77]. Then in §3.2, we will sketch the proof of this direction of the structural stability theorem by Robbin [Rob71] and Robinson [Rob76]; and we will relate a few works leading to a generalization of the Przytycki conjecture [Prz77], a description of the structurally stable local diffeomorphisms. Finally in §3.3, we will recall our conjecture with Rovella [BR13] stating a description of the endomorphisms (with possibly a non-empty critical set) whose inverse limit is structurally stable, and we will state our theorem with Kocsard [BK13] showing one direction of this conjecture.

In section 4, we will recall several results from singularity theory and we will emphasize on their similarities with those of structural stability.

In section 5, we will present the work [Ber12] which states sufficient conditions for a smooth map with non-empty critical set to be structurally stable. The statement involves developments of Mather's theorem on Singularity Theory of composed mappings. It suggests the problem of the description of the structurally stable, surface endomorphisms among those which display singularity but satisfy the axiom A.

These notes were written while I was giving lectures at Montevideo in 2009 and at the Banach Center in 2016. I am very grateful for their hospitality.

1 Uniformly hyperbolic dynamical systems

The theory of uniformly hyperbolic dynamical systems was constructed in the 1960's under the dual leadership of Smale in the USA, and Anosov and Sinai in the Soviet Union.¹

It encompasses various examples that we shall recall: expanding maps, horseshoes, solenoid maps, Plykin attractors, Anosov maps, DA, blenders all of which are *basic pieces*.

¹A few sentences of this section are taken from [BY14].

1.1 Uniformly hyperbolic diffeomorphisms

Let f be a C^1 -diffeomorphism f of a finite dimensional manifold M. A compact f-invariant subset $\Lambda \subset M$ is uniformly hyperbolic if the restriction to Λ of the tangent bundle TM splits into two continuous invariant subbundles

$$TM|\Lambda = E^s \oplus E^u$$
,

 E^s being uniformly contracted and E^u being uniformly expanded: $\exists \lambda < 1, \exists C > 0,$

$$||T_x f_{|E^s}^n|| < C \cdot \lambda^n$$
 and $||T_x f_{|E^u}^{-n}|| < C \cdot \lambda^n$, $\forall x \in \Lambda, \forall n \ge 0$.

Example 1.1 (Hyperbolic periodic point). A periodic point at which the differential has no eigenvalue of modulus 1 is called hyperbolic. It is a sink if all the eigenvalues are of modulus less than 1, a source if all of them are of modulus greater than 1, and a saddle otherwise.

Definition 1.2. A hyperbolic attractor is a hyperbolic, transitive compact subset Λ such that there exists a neighborhood N satisfying $\Lambda = \bigcap_{n \geq 0} f^n(N)$.

Example 1.3 (Anosov). If the compact hyperbolic set is equal to the whole compact manifold, then the map is called Anosov. For instance if a map $A \in SL_2(\mathbb{Z})$ has both eigenvalues of modulus not equal to 1, then it acts on the torus $\mathbb{R}^2/\mathbb{Z}^2$ as an Anosov diffeomorphism. The following linear map satisfies such a property:

$$A := \left[\begin{array}{cc} 2 & 1 \\ 1 & 1 \end{array} \right] .$$

Example 1.4 (Smale solenoid). We consider a perturbation of the map of the filled torus $\mathbb{T} := \{(\theta, z)\mathbb{R}/\mathbb{Z} \times \mathbb{C} : |z| < 1\}$:

$$(\theta, z) \in \mathbb{T} \mapsto (2\theta, 0) \in \mathbb{T}$$
,

which is a diffeomorphism onto its image. This is the case of the following:

$$(\theta, z) \in \mathbb{T} \mapsto (2\theta, \epsilon \cdot z + 2\epsilon \cdot \exp(2\pi i\theta)) \in \mathbb{T}$$
.

This defines a hyperbolic attractor called the *Smale solenoid*.

Example 1.5 (Derivated from Anosov (DA) and Plykin attractor). We start with a linear Anosov of the 2-torus $\mathbb{R}^2/\mathbb{Z}^2$. It fixes the point 0. In local coordinates ϕ of a neighborhood V of 0, it has the form for $0 < \lambda < 1$:

$$(x,y) \mapsto (\lambda x, y/\lambda).$$

For every $\epsilon > 0$, let ρ_{ϵ} be a smooth function so that:

- it is equal to $x \mapsto \lambda x$ outside of the interval $(-2\epsilon, 2\epsilon)$,
- ρ_{ϵ} displays exactly three fixed point: $-\epsilon$ and ϵ which are contracting and 0 which is expanding.

Let DA be the map of the two torus equal to A outside of V, and in the coordinate ϕ it has the form:

$$(x,y) \mapsto (\rho_{\epsilon}(x), y/\lambda).$$

We notice that 0 is an expanding the fixed point of DA. The complement of its repulsion basin is a hyperbolic attractor.




Figure 1: Derivated of Anosov (Credit Y. Coudene [Cou06])

The DA attractor project to a basic set of a surface attractor, the *Plykin attractor*.




Figure 2: Plykin attractor (Credit S. Crovisier)

Given a hyperbolic compact set Λ , for every $z \in \Lambda$, the sets

$$W^s(z) = \{z' \in M: \lim_{n \to +\infty} d(f^n(z), f^n(z')) = 0\},\$$

$$W^{u}(z) = \{z' \in M : \lim_{n \to -\infty} d(f^{n}(z), f^{n}(z')) = 0\}$$

are called the stable and unstable manifolds of z. They are immersed manifolds tangent at z to respectively $E^s(z)$ and $E^u(z)$.

The ϵ -local stable manifold $W^s_{\epsilon}(z)$ of z is the connected component of z in the intersection of $W^s(z)$ with a ϵ -neighborhood of z. The ϵ -local unstable manifold $W^u_{\epsilon}(z)$ is defined likewise.

Proposition 1.6. For $\epsilon > 0$ small enough, the subsets $W^s_{\epsilon}(z)$ and $W^u_{\epsilon}(z)$ are C^r -embedded manifolds which depend continuously on z and tangent at z to respectively $E^s(z)$ and $E^u(z)$.

A nice proof of this proposition can be found in [Yoc95].

Definition 1.7. A basic set is a compact, f-invariant, transitive, uniformly hyperbolic set Λ which is locally maximal: there exists a neighborhood N of Λ such that $\Lambda = \bigcap_{n \in \mathbb{Z}} f^n(N)$.

Example 1.8 (Horseshoe). A horseshoe is a basic set which is a Cantor set. For instance take two disjoint sub-intervals $I_+ \sqcup I_- \subset [0,1]$, and let $g: I_+ \sqcup I_- \to [0,1]$ be a locally affine map which sends each of the intervals I_{\pm} onto [0,1]. Let g_+ be its inverse branch with value in I_+ and let g_- be the other inverse branch. Let f be a diffeomorphism of the plane whose restriction to $I_{\pm} \times [0,1]$ is:

$$(x,y) \in (I_+ \sqcup I_-) \times [0,1] \to \begin{cases} (g(x), g_+(y)) & \text{if } x \in I_+ \\ (g(x), g_-(y)) & \text{if } x \in I_- \end{cases}$$




Figure 3: Smale's Horseshoe

Remark 1.9. Usually, one defines a basic piece as a hyperbolic set included in the closure of the set of its periodic points. Actually the three following assertion are equivalent for every uniformly hyperbolic, transitive, compact set K:

- K is locally maximal.
- K has a structure of local product : for $\epsilon > 0$ small enough, and any $x, y \in K$ close enough, the intersection point $W^u_{\epsilon}(x) \cap W^s_{\epsilon}(y)$ belongs to K.
- K is included in the closure of the set of periodic points in K: K = cl(Per(f|K)).

The equivalence of these conditions is proved in [Shu78].

Definition 1.10 (Axiom A). A diffeomorphism whose non-wandering set is a finite union of disjoint basic sets is called axiom A.

Example 1.11 (Morse-Smale). A Morse-Smale diffeomorphism is a diffeomorphism of a surface so that its non-wandering set consists of finitely many periodic hyperbolic points, and their stable and unstable manifolds are transverse.




Figure 4: Morse-Smale

1.2 Uniformly hyperbolic endomorphisms

A C^r -endomorphism of a manifold M is a differentiable map of class C^r of M, which is not necessarily injective, nor surjective, and that may possess points at which the differential is not onto (called critical points). The *critical set* is the subset of M of formed by the critical points.

A local C^r -diffeomorphism is a C^r -endomorphism without critical point.

A compact subset $\Lambda \subset M$ is *invariant* for an endomorphism f of M if $f^{-1}(\Lambda) = \Lambda$. A compact subset $\Lambda \subset M$ is *stable* for an endomorphism f of M if $f(\Lambda) = \Lambda$.

An invariant compact set is *hyperbolic* if there exists a subbundle $E^s \subset TM\Lambda$ which is left invariant and uniformly contracted by Df and so that the action of Df on TM/E^s is uniformly expanding.

Example 1.12 (Expanding map). Let $f \in End^1(M)$ and an invariant stable, compact subset K is expanded if there exists $n \geq 1$ s.t., for every $x \in K$, $D_x f^n$ is invertible and with contracting inverse. When K = M, f is said expanding.

Example 1.13 (Anosov endomorphism). If a hyperbolic set is equal to the whole manifold, then the endomorphism is called Anosov. For instance this is the case of the dynamics on the torus $\mathbb{R}^2/\mathbb{Z}^2$ induced by a linear maps in $M_2(\mathbb{Z})$ with eigenvalues of modulus not equal to 1. For instance, it the case of the following for every $n \geq 2$:

$$\left[\begin{array}{cc} n & 1 \\ 1 & 1 \end{array}\right]$$

The stable manifold of z in a hyperbolic set Λ of an endomorphism is defined likewise:

$$W^{s}(z) = \{ z' \in M : \lim_{n \to +\infty} d(f^{n}(z), f^{n}(z')) = 0 \}.$$

The unstable manifold depends on the preimages. For every orbit $\underline{z} = (z_n)_{n \in \mathbb{Z}} \in \Lambda^{\mathbb{Z}}$, has an unstable manifold:

$$W^{u}(\underline{z}) = \{z' \in M : \exists (z'_n)_n \text{ orbit s.t. } \lim_{n \to -\infty} d(z_n, z'_n) = 0\}.$$

If f is a local diffeomorphism then W^s and W^u are immersed, but in general only W^s is injectively immersed.

Definition 1.14. A hyperbolic set Λ is a basic piece if it is locally maximal.

Example 1.15 (Blender). A blender of surface endomorphism is a basic set so that C^1 -robustly its local unstable manifold cover an open subset of the surface.

For instance let I_{-} and I_{+} be two disjoint segments of [-1,1], and let Q be a map which sends affinely each of these segments onto [-1,1]. This is the case for instance of the following map:

$$(x,y) \in [-1,1]^2 \mapsto \begin{cases} (Q(x),(2y+1)/3) & x \in I_+ \\ (Q(x),(2y-1)/3) & x \in I_- \end{cases}$$




Figure 5: Blender of a surface local diffeomorphism

Definition 1.16. An endomorphism satisfies axiom A if its non-wandering set is a finite union of basic pieces.

2 Properties of structurally stable dynamics

Let us sate some definitions and conjectures on the structural stability in the C^r -category, $1 \le r \le \infty$ and in the holomorphic category denoted by \mathcal{H} . Let \mathcal{C} be a category in $\{C^r : 1 \le r \le \infty\} \cup \{\mathcal{H}\}$.

Definition 2.1 (Structural stability). A C-map f is structurally stable if every C-perturbation f' of the dynamics is conjugated: there exists a homeomorphism h of the manifold so that $h \circ f = f' \circ h$.

A weaker notion of structural stability focuses on the non-wandering set Ω_f of the dynamics f.

Definition 2.2 (Ω -stability). A C-map f is Ω -stable if for every C-perturbation f' of f, the dynamics of the restriction of f to Ω_f is conjugated (via a homeomorphism) to the restriction of f' to its non-wandering set $\Omega_{f'}$.

We recall that an axiom A diffeomorphism f satisfies the *strong transversality condition* if its stable and unstable manifolds intersect transversally. Here is an outstanding conjecture:

Conjecture 2.3 (Palis-Smale structural stability conjecture, 1970 [PS70]). A C-diffeomorphism is structurally stable if and only if it satisfies axiom A and the strong transversality condition.

For complex rational maps of the sphere, this conjecture takes the form:

Conjecture 2.4 (Fatou Conjecture, 1920). Structurally stable quadratic map are those which satisfy axiom A and whose critical points are not periodic.

Actually the initial Fatou conjecture stated the density of axiom A quadratic map. However, in section 2, we will recall the works of Mañé-Sad-Sullivan [MSS83] and Lyubich [Lyu84] showing the existence of an open and dense set of structurally stable rational maps. This implies the equivalence between the original Fatou Conjecture and the above conjecture. Among real quadratic maps, this conjecture² has been proved by Graczyk-Swiantek [GŚ97] and Lyubich [Lyu97].

The description of Ω -stable maps involves the no-cycle condition. We recall that any axiom A diffeomorphisms displays a non-wandering set Ω equal to a finite union of basic pieces $\Omega = \sqcup_i \Omega_i$. The family $(\Omega_i)_i$ is called the *spectral decomposition*.

Definition 2.5 (No-cycle condition). An axiom A diffeomorphism satisfies the no-cycle condition if given $\Omega_1, \Omega_2, \ldots, \Omega_n$ in the spectral decomposition, if $W^u(\Omega_i)$ intersects $W^s(\Omega_{i+1})$ for every i < n and if $W^u(\Omega_n)$ intersects $W^s(\Omega_1)$, then $\Omega_1 = \Omega_2 = \cdots = \Omega_n$.

Conjecture 2.6 (Smale Ω -Stability Conjecture, [Sma70]). A C-diffeomorphisms is structurally stable if and only if it satisfies axiom A and the no-cycle condition.

If the above conjectures turn out to be true then they would display a satisfactory description of structurally stable dynamics (for the axiom A diffeomorphisms are very well understood).

Let us define the probabilistic structural stability, which is implied by the Ω -stability. The definition involves the regular subset \mathcal{R}_f of Ω_f . This subset is formed by the points $p \in \Omega_f$ so that for every $a \in \{s, u\}$, there exist $\epsilon > 0$ and a sequence of periodic points $(p_n)_n$ satisfying:

- $(p_n)_n$ converges to p,
- $(W_{\epsilon}^{a}(p_{n}))_{n}$ is relatively compact in the \mathcal{C} -topology.

We showed in [BD17] thanks to Katok's closing Lemma, that the set \mathcal{R}_f has full measure for every ergodic, hyperbolic probability measure.

² The Fatou conjecture is implied by the Mandelbrot Locally connected (MLC) conjecture that we will not have the time to recall in this manuscript.

Definition 2.7. A C-map f is probabilistically structurally stable if for every C-perturbation f' of f, the restriction of f to \mathcal{R}_f is conjugated to the restriction of f' to its regular set $\mathcal{R}_{f'}$.

It is rather easy to see that probabilistic structural stability implies weak stability:

Definition 2.8. A map f is C-weakly stable if every C-perturbation f' of f displays only hyperbolic periodic points.

To sum it up, the above definitions are related as follows:

 Ω -Stability \Rightarrow Probabilistic Stability \Rightarrow Weak Stability

The Lambda Lemma Conjecture. This conjecture states that weak stability implies Ω -stability. For the category of rational functions of the Riemannian sphere, this Lemma has been shown independently by Mañé-Sad-Sullivan [MSS83] and Lyubich [Lyu84].

As the space of rational functions is finite dimensional, a neighborhood of a rational function f can be written as an analytic family $(f_{\lambda})_{\lambda \in \mathbb{D}^n}$, with \mathbb{D} the complex disk and $f_0 = f$. If $(f_{\lambda})_{\lambda}$ consists of weakly stable maps, then every periodic point p_0 of f_0 persists to as unique periodic point p_{λ} for f_{λ} . Moreover the map $\lambda \mapsto p_{\lambda}$ is holomorphic. The Lambda lemma asks the following question. Given p_0 in closure J_0^* of the set of periodic points of f_0 , for every sequence $(p_0^n)_n$ of periodic points converging to p_0 , does the family $(\lambda \mapsto p_{\lambda}^n)_n$ converges? If yes, the holomorphic motion is said well defined at p_0 .

Lemma 2.9 (Lambda-Lemma, Mañé-Sad-Sullivan [MSS83] and Lyubich [Lyu84]). If $(f_{\lambda})_{\lambda}$ is weakly stable, then the holomorphic motion is well defined at every point $p_0 \in J_0^*$.

We recall that every rational function J^* is equal to the non-wandering set and that any attracting periodic point displays a critical point in its basin. Furthermore if a rational function is not weakly stable, it displays a new attracting periodic point after a perturbation of the rational function. Hence the new critical point belongs to the basin of this attracting periodic point. As the number of critical points is finite, after a finite number of perturbations the rational function turns out to be weakly stable. This shows that weak stability is open and dense among the rational functions. By the Lambda Lemma 2.9, this implies:

Theorem 2.10 (Mañé-Sad-Sulivan [MSS83], Lyubich [Lyu84]). There is an open and dense subset of rational functions of degree $d \ge 2$ which are Ω -stable.

This result enables them to deduce a stronger result: the density of the set of structurally stable rational functions.

We recall that a polynomial automorphism of \mathbb{C}^2 is a polynomial mapping of \mathbb{C}^2 which is invertible and whose inverse is polynomial. Among polynomial automorphisms of \mathbb{C}^2 , Dujardin and Lyubich [DL] showed that the holomorphic motion is well defined on any uniformly hyperbolic compact set. We improved this result:

Lemma 2.11 (Berger-Dujardin [BD17]). If $(f_{\lambda})_{\lambda}$ is a weakly stable family of polynomial automorphisms of \mathbb{C}^2 , the holomorphic motion is uniquely defined on the regular set \mathcal{R}_0 of f_0 .

An immediate consequence of this result is that weak stability implies probabilistic stability for the category of polynomial automorphisms of \mathbb{C}^2 .

Unfortunately, there is no hope to get the density of Ω -stable polynomial automorphisms of \mathbb{C}^2 because in a non-empty open set [Buz97] of the parameter space is formed by automorphisms displaying a wild horseshoe. However, we will see below that if none pertubations of the dynamics display a homoclinic tangency, then the dynamics is weakly stable (under a mild hypothesis of dissipativeness).

As a corollary of the techniques, we showed that one connected component of the set of weakly stable polynomial automorphisms is formed by those which satisfy axiom A.

The Mañé Conjecture In 1982, Mañe conjectured in [Mañ82] that every C^r -weakly stable diffeomorphism satisfies axiom A for every $1 \le r \le \infty$. He proved this conjecture for r = 1 and surface diffeomorphisms. Mañé developed this technology to prove that C^1 -structurally stable diffeomorphisms satisfy axiom A and the strong transversality condition in [Mañ88]. This work enabled also Palis to prove the same direction for the C^1 - Ω -stability conjecture [Pal88]. By developing Mañé's works, Aoki and Hayashi proved the Mañé conjecture for r = 1 in any dimension [Aok92, Hay92].

Weak Stability
$$\stackrel{\text{Manñ\'e Conj.}}{\Longrightarrow}$$
 axiom A.

After the next section, it will be clear for the reader that the Mañé Conjecture implies the Lambda Lemma Conjecture in any category C.

A Palis Conjecture We recall that a hyperbolic periodic point displays a homocline tangency if its stable manifold $W^s(p)$ is tangent to its unstable manifold. Two saddle periodic points p, q display a heterocline tangency if $W^s(p)$ intersects transversally $W^u(q)$ whereas $W^s(q)$ is tangent to $W^u(p)$ (or vice versa). It is not hard to show that if a C^r -map is weakly stable then it cannot display a homoclinic nor a heteroclinic tangency, for every $1 \le r \le \infty$. The same is true for one dimensional complex maps. For polynomial automorphisms of \mathbb{C}^2 , it is a theorem [Buz97].

Let us recall also a famous Conjecture of Palis [Pal00] which states that if a dynamics which cannot be perturbed to one which displays a homoclinic nor a heteroclinic tangency, then it satisfies axiom A:

Weak Stability
$$\Rightarrow$$
 Far from tangencies $\stackrel{\text{Palis Conj.}}{\Longrightarrow}$ axiom A.

In the category of C^1 -surface diffeomorphisms, this conjecture has been proved by Pujals-Sambarino [PS00]. In the category of C^1 -diffeomorphisms of higher dimensional manifolds, a weaker version has been proved by Crovisier-Pujals [CP15].

We notice that the Palis conjecture implies the Mañé conjecture and so the Lambda lemma conjecture.

A description of structurally stable dynamics as those far from tangencies? This question is widely open in the C^r -category for r > 1 (for C^1 -surface diffeomorphisms it is a consequence of Mañé's theorem). It is also correct for the category of rational functions. This might be correct for polynomial automorphisms of \mathbb{C}^2 . Indeed, most of the work of Dujardin-Lyubich was dedicated to prove the following result:

Theorem 2.12 (Dujardin-Lyubich [DL]). Given a polynomial automorphism f of (dynamical) degree $d \geq 2$ and so that $|\det Df_0| \cdot d^2 < 1$, either f is weakly stable, either a perturbation of f' admits a homoclinic tangency.

From Lambda Lemma 2.11 we deduced:

Corollary 2.13 (Berger-Dujardin [DL]). Given a polynomial automorphism f of (dynamical) degree $d \ge 2$ and so that $|\det Df_0| \cdot d^2 < 1$, either f is probabilistically stable, either a perturbation of f' displays a homoclinic tangency.

Let us stress that this direction might be interesting since numerically we can see some local stable and unstable manifolds and observe if they display tangencies.




Figure 6: Summary of some Theorems and Conjectures on Structural Stability

3 Hyperbolicity implies structural stability

In the following subsection, we recall the proof ideas of several basic theorems showing the structural stability of subsets from hyperbolic hypotheses.

3.1 Ω -stability of maps satisfying axiom A and the no-cycle condition

First let us recall a generalization of the notion of structural stability for invariant subsets.

Definition 3.1 (Structurally stable subset). A compact set Λ left invariant by a differentiable map f of a manifold M is structurally stable if for every C^r -perturbation f' of f, there exists a continuous injection $i: \Lambda \to M$ so that $f' \circ i = i \circ f$.

We notice that M is structurally stable if and only if f is structurally stable.

Theorem 3.2 (Anosov [Ano67], proof by Moser [Mos69]). A uniformly hyperbolic compact set Λ for a C^1 -diffeomorphisms is structurally stable.

Proof. We want to solve the following equation:

$$(\star) f' \circ h \circ f^{-1} = h .$$

for f' C^1 -close to f and h C^0 -close to the canonical inclusion $i: \Lambda \hookrightarrow M$. We shall use the implicit function theorem with the map:

$$\Phi \colon (h, f') \in C^0(\Lambda, M) \times C^1(M, M) \to f' \circ h \circ f^{-1} \in C^0(\Lambda, M)$$
.

We notice that Φ is a C^1 -differentiable map of Banachic manifolds. Moreover it satisfies $\Phi(i, f) = i$. Hence to apply the implicit function theorem it suffices to prove that $id - \partial_h \Phi(i, f)$ is an isomorphism.

Note that the tangent space of the Banachic manifold $C^0(\Lambda, M)$ at the canonical inclusion i is the following Banach space:

$$\Gamma := \{ \gamma \in C^0(\Lambda, TM) : \forall x \in \Lambda \quad \gamma(x) \in T_x M \}.$$

The partial derivative of $\partial_h \Phi$ at (i, f) is:

$$\Psi := \partial_h \Phi(i, f) \colon \sigma \in \Gamma \mapsto Df \circ \sigma \circ f^{-1} \in \Gamma .$$

To compute the inverse of $id - \Psi$, we split Γ into two Ψ -invariant subspaces $\Gamma = \Gamma^u \oplus \Gamma^s$, with:

$$\Gamma^u := \{ \gamma \in C^0(\Lambda, TM) : \forall x \in \Lambda \quad \gamma(x) \in E^u_x \} \quad \text{and} \quad \Gamma^s := \{ \gamma \in C^0(\Lambda, TM) : \forall x \in \Lambda \quad \gamma(x) \in E^s_x \}.$$

As the norm of $\Psi | \Gamma^s$ is less than 1, the map $(id - \Psi) | \Gamma^s$ is invertible with inverse equal to

$$\sum_{n>0} (\Psi|\Gamma^s)^n .$$

As $\Psi | \Gamma^u$ is invertible with contracting inverse, the map $(id - \Psi) | \Gamma^u$ is invertible with inverse:

$$-(\Psi|\Gamma^{u}) \circ (id - (\Psi|\Gamma^{u})^{-1}) = -(\sum_{n \ge 1} (\Psi|\Gamma^{u})^{-n}) .$$

Hence by the implicit function theorem, for every f' C^1 -close to f, there exists a continuous map h C^0 -close to i which semi-conjugates the dynamics:

$$f' \circ h = h \circ f$$
.

As i is injective and close to h, if h(x) = h(y) then x and y are close. Also by semi-conjugacy, $h \circ f^n(x) = h \circ f^n(y)$ for every $n \in \mathbb{Z}$. Hence $f^n(x)$ is close to $f^n(y)$ for every n. By expansiveness (see below), we conclude that x = y and so that h is injective.

Lemma 3.3 (Expansiveness). Every hyperbolic compact set Λ for a diffeomorphism is expansive: there exists $\epsilon > 0$ so that if two orbits $(x_n)_{n \in \mathbb{Z}}$ and $(y_n)_{n \in \mathbb{Z}}$ are uniformly ϵ -close, then $x_0 = y_0$.

Proof. First we notice that for ϵ small enough, given two such orbits, $W_{2\epsilon}^s(y_n)$ intersects $W_{2\epsilon}^u(y_n)$ at a unique point z_n . We observe that $(z_n)_n$ is an orbit. As f is expanding along $W_{2\epsilon}^u(y_n)$ for every n and since $z_n \in W_{2\epsilon}^u(y_n)$, it comes that $z_n = y_n$ for every $n \geq 0$. Using the same argument for f^{-1} , it comes that $z_n = x_n$ for every $n \leq 0$ and so $x_0 = y_0$.

The image $\Lambda(f') := h(\Lambda)$ is called the *hyperbolic continuation* of Λ . Since the density of periodic points is preserved by conjugacy, it comes:

Corollary 3.4. If Λ is a basic piece, then its hyperbolic continuation is also a basic piece.

A similar result has been proved by Shub during his thesis:

Theorem 3.5 (Shub [Shu69]). An expanding compact set Λ for an endomorphisms f is C^1 -structurally stable.

Proof. First let us notice that f is a local diffeomorphism at a neighborhood of the compact set Λ . Hence there exists $\epsilon > 0$ so that for every f' C^1 -close to f, for every $x \in \Lambda$, the restriction $f'|B(x,\epsilon)$ is invertible. This enables us to look for a semi-conjugacy thanks to the map:

$$\Phi \colon (h, f') \in C^0(\Lambda, M) \times C^1(M, M) \to (f'|B(x, \epsilon))^{-1} \circ h \circ f \in C^0(\Lambda, M)$$

The latter is well defined and of class C^1 on the ϵ -neighborhood of the pair of the canonical inclusion $i \colon \Lambda \hookrightarrow M$ with f. Furthermore, it holds $\Phi(i, f) = i$ and the following partial derivative is contracting, with Γ the tangent space of $C^0(\Lambda, M)$ at i.

$$\partial_h \Phi(i,f) \colon \sigma \in \Gamma \to Df^{-1} \circ \sigma \circ f \in \Gamma$$
.

Thus, by the implicit function Theorem, for f' C^1 -close to f, there exists a unique solution with $h \in C^0(\Lambda, M)$ close to i for the semi-conjugacy equation:

$$\Phi(h, f') = h \Leftrightarrow h \circ f = f' \circ h$$
.

As h is close to the canonical inclusion, if h(x) = h(x') then x and x' must be close. Also by semi-conjugacy, it holds $h(f^n(x)) = h(f^n(x'))$ for every $n \ge 0$. Thus the orbits $(f^n(x))_{n\ge 0}$ and $(f^n(x'))_{n\ge 0}$ are uniformly close. By forward expansiveness (see below), it comes that x = x'.

One easily shows by a similar argument to Lemma 3.3:

Lemma 3.6 (Forward expansiveness). Every expanding compact set Λ is forward expansive: there exists $\epsilon > 0$ so that if two orbits $(x_n)_{n \geq 0}$ and $(y_n)_{n \geq 0}$ are uniformly ϵ -close, then $x_0 = y_0$.

The two latter theorems enable us to explain the proofs of Smale and Przytycki on Ω -stability. We recall that the local stable and unstable manifolds of the points of a hyperbolic set Λ for an endomorphism f (which might display a non-empty critical set) are uniquely defined, provided that:

- Either $f|\Lambda$ is bijective,
- Either Λ is injective.

On the other hand, the local stable manifold are always uniquely defined. Hence under these assumption, by looking at their images or preimages, the following is uniquely defined for $\epsilon > 0$ small enough:

$$W^s_{\epsilon}(\Lambda) = \bigcup_{x \in \Lambda} W^s_{\epsilon}(x) \quad W^u_{\epsilon}(\Lambda) = \bigcup_{x \in \Lambda} W^u_{\epsilon}(x) \quad W^s(\Lambda) = \bigcup_{n > 0} f^{-n}(W^s_{\epsilon}(\Lambda)) .$$

The following generalizes Smale's definion of axiom A diffeomorphisms:

Definition 3.7 (Axiom A in the sens of Przytycki). A C^1 -endomorphism satisfies axiom A-Prz, if its non-wandering set Ω is equal to the closure of the set of periodic points (or equivalently locally maximal), and if it is the disjoint union of an expanding compact set with a bijective, hyperbolic compact set.

For such maps we can generalize the notion of spectral decomposition. Indeed by local maximality and compactness, the non-wandering set Ω of such maps is the finite union of (maximal) transitive subsets Ω_i called basic pieces:

$$\Omega = \sqcup_i \Omega_i$$
.

The family $(\Omega_i)_i$ is called the *spectral decomposition* of the axiom A-Prz endomorphism. Let us generalize the no-cycle condition for such endomorphisms.

Definition 3.8 (No-cycle condition). An axiom A-Prz, C^1 -endomorphism satisfies the no-cycle condition if given $\Omega_1, \Omega_2, \ldots, \Omega_n$ in the spectral decomposition, if $W^u_{\epsilon}(\Omega_i)$ intersects $W^s(\Omega_{i+1})$ for every i < n and if $W^u_{\epsilon}(\Omega_n)$ intersects $W^s(\Omega_1)$, then $\Omega_1 = \Omega_2 = \cdots = \Omega_n$.

F. Przytycki generalized Smale's Theorem on the Ω -stability of axiom A diffeomorphisms which satisfy the no-cycle condition as follows:

Theorem 3.9 ([Sma70], [Prz77]). If a C^1 -endomorphism satisfies axiom A-Prz and the no-cycle condition, then it is $C^1 - \Omega$ -stable.

Sketch of proof of the Smale's Ω -stability Theorem. First let us recall that by Anosov Theorem, the non-wandering set Ω is structurally stable, and its hyperbolic continuation is still locally maximal (for a neighborhood uniformly large among an open set of perturbations of the dynamics).

Then the no-cycle condition is useful to construct a filtration $(M_i)_i$:

Proposition 3.10. If an axiom A, C^1 -diffeomorphism f satisfies the no-cycle condition, then there exists a chain of open subsets:

$$\emptyset = M_0 \subset M_1 \subset \cdots \subset M_N = M$$

so that $f(M_i) \in M_i$ and $\Omega_i \in M_i \setminus M_{i-1}$ for every $i \geq 1$.

The proof of this proposition involve Conway Theory and can be find in [Shu78, Thm 2.3 p. 9]. By using this filtration and the (uniform) local maximality of the hyperbolic continuation of the non-wandering set, one easily deduces the Ω -stability.

3.2 Structural stability of dynamics satisfying axiom A and the strong transversality condition

Structural stability of diffeomorphisms We recall that an axiom A diffeomorphism satisfies the strong transversality condition if for any non-wandering points x and y, the stable manifold of x is transverse to the unstable manifold of y.

Remark 3.11. By using the inclination lemma, one easily shows that the strong transversality condition implies the no-cycle condition.

The following theorem generalizes Anosov Theorems 3.2:

Theorem 3.12 (Robbin [Rob71], Robinson [Rob76]). For every $r \ge 1$, the diffeomorphisms which satisfy axiom A and the strong transversality condition are C^r -structurally stable.

Let us recall that the Mañe theorem [Mañ88] implies that a C^1 -structurally stable diffeomorphism satisfies also axiom A and the strong transversality condition, and so both solve the conjecture of C^1 -structural stability.

We will state Conjecture 3.15 generalizing this theorem for local diffeomorphisms. Hopefully the following will help the reader to tackle it.

Sketch of proof of Theorem 3.12. Again we want to solve the following semi-conjugacy equation:

$$(\star) \qquad \qquad f' \circ h \circ f^{-1} = h$$

for f' C^1 -close to f and h C^0 -close to the identity of M.

For f'=f and h=id, Equality (\star) is valid. The set of perturbations of the identity is isomorphic to $\Gamma=\{\gamma\in C^0(M,TM): \forall x\in M\quad \gamma(x)\in T_xM\}$ by using the exponential map (associated to a Riemannian metric on M). Let $\tilde{f}:=u\in T_xM\mapsto \exp_{f(x)}^{-1}\circ f\circ \exp_x(u)$.

Then Equation (\star) is equivalent to:

$$(\star\star) \qquad \qquad \tilde{f}'\circ\sigma\circ f^{-1}=\sigma, \quad \text{for } \sigma\in\Gamma \quad C^0\text{-small}.$$

As the map $\Phi \colon (\sigma, f') \in \Gamma \times C^1(M, M) \to \Phi_{f'}(\sigma) = \sigma - \tilde{f'} \circ \sigma \circ f^{-1} \in \Gamma$ is of class C^1 , and vanishes at (0, f), we shall show that $\partial_h \Phi$ is left-invertible.

Let

$$\Psi := \partial_h \Phi(0, f) \colon \sigma \in \Gamma \mapsto \sigma - Df \circ \sigma \circ f^{-1} \in \Gamma .$$

The following is shown in [Rob71]:

Proposition 3.13. For every i, there exists a neighborhood N_i of Ω_i and continuous extension E_i^s and E_i^u of respectively $E^s|\Omega_i$ and $E^u|\Omega_i$ to N_i , so that:

- There exists a filtration $(M_i)_i$ adapted to $(\Omega_i)_i$ so that $N_i = M_i \setminus M_{i-1}$. The subsets $(N_i)_i$ form an open covering of M,
- if $x \in N_i \cap f^{-1}(N_j)$, with $j \leq i$, then $Df(E_i^s(x)) \subset E_i^s(f(x))$, and $Df(E_i^u(x)) \supset E_i^u(f(x))$.

Let $(\gamma_i)_i$ be a partition of the unity adapted to $(N_i)_i$.

For every i let p_i^s and p_i^u be the projections onto respectively E_i^s and E_i^u parallely to E_i^u and E_i^s . Given $x \in M$ and $v \in T_xM$, we put $v_i^s := \gamma_i \cdot p_i^s(v)$ and $v_i^u := \gamma_i \cdot p_i^u(v)$. We observe that $v = \sum_i v_i^s + v_i^u$. Thus $Df(v) = \sum_i Df(v_i^s) + Df(v_i^u)$. As $(Df^n(v_i^s))_{n\geq 0}$ and $(Df^{-n}(v_i^u))_{n\geq 1}$ converge exponentially fast to 0, we consider:

$$J \colon \sigma \in \Gamma \mapsto \sum_{i} \sum_{n>0} Df^{n}(\sigma_{i}^{s} \circ f^{-n}(x)) - \sum_{n>1} Df^{-n}(\sigma_{i}^{u} \circ f^{n}(x)) .$$

We notice that J is a left inverse of Ψ :

$$J \circ \Psi = id$$

The following equations are equivalent:

$$\Phi_{f'}(\sigma) = 0 \Leftrightarrow (\Phi_{f'} - \Psi)(\sigma) + \Psi(\sigma) = 0,$$

$$\Leftrightarrow J \circ (\Phi_{f'} - \Psi)(\sigma) + J \circ \Psi(\sigma) = J(0).$$

Now observe that J(0) = 0 and $J \circ \Psi(\sigma) = \sigma$. Hence $(\star\star)$ is equivalent to

$$J \circ (\Psi - \Phi_{f'})(\sigma) = \sigma.$$

It is easy to see that whenever f' is C^1 -close to f, the map $\Phi_{f'}$ is C^1 -close to Ψ at a neighborhood of the 0-section. Hence the map $J \circ (\Psi - \Phi_{f'})$ is contracting and sends a closed ball about the zero section into itself. The contracting mapping theorem implies the existence of a fixed point σ . Hence (\star) displays a solution $h = \exp \circ \sigma$ close to the identity in the space of continuous maps.

It remains to show that the semi-conjugacy h is bijective. Contrarily to Anosov maps, in general axiom A diffeomorphisms are not expansive and the semi-conjugacy is not uniquely defined. Hence Robbin brought a new technique to construct a map h which is bijective. He defined the following metric:

$$d_f(x,y) = \sup_{n \in \mathbb{Z}} d(f^n(x), f^n(y)) ,$$

where d is the Riemannian metric of the manifold M.

Let us just notice that if the semi-conjugacy $h = \exp \circ \sigma$ satisfies that σ is C^0 -small and d_f -Lipschitz with a small constant η , then h is injective.

Indeed if h(x) = h(y), then by (\star) , $h(f^n(x)) = h(f^n(y))$ for every n. Since h is close to the identity, the orbits $(f^n(x))_n$ and $(f^n(y))_n$ are uniformly close, and so that $d_f(x,y)$ is small. As σ is η -Lipschitz, it comes:

$$0 = d(h(x), h(y)) \ge d(x, y) - \eta d_f(x, y)$$

The same holds at any n^{th} -iterate:

$$0 = d(h(f^n(x)), h(f^n(y))) \ge d(f^n(x), f^n(y)) - \eta d_f(f^n(x), f^n(y)) = d(f^n(x), f^n(y)) - \eta d_f(x, y).$$

Let n be such that $d(f^n(x), f^n(y)) \ge d_f(x, y)/2$. Then

$$0 = d(h(f^n(x)), h(f^n(y))) \ge (1 - 2\eta)d(f^n(x), f^n(y)) .$$

Thus $f^n(x) = f^n(y)$ and so x = y.

To obtain the section σ d_f -Lipschitz, Robbin assumed the diffeomorphism f of class C^2 . Then in Proposition 3.13, he constructs the section $(E_i^s)_i$ and $(E_i^u)_i$ d_f -Lipschitz, so that the map J preserves the d_f -Lipschitz sections. On the other hand the map $\Psi - \Psi_{f'}$ diminishes the d_f -Lipschitz constant for f' C^1 -close to f. Therefore the map $J \circ (\Psi - \Psi_{f'})$ preserves the space of continuous sections with small d_f -Lipschitz constant, and so its fixed point enjoys a small d_f -Lipschitz constant.

The C^1 -case was handled by Robinson. His trick was to smooth the map Df to a C^1 -map $\tilde{D}f$, and to replace Df by $\tilde{D}f$ in the definition of Ψ to define $\tilde{\Psi}$. Then he defined likewise $\tilde{D}f$ -pseudo invariant sections $(\tilde{E}_i^s)_i$ which are d_f -Lipschitz. By replacing $(E_i^s)_i$ by $(\tilde{E}_i^s)_i$ in the definition of J, he defined a left inverse \tilde{J} of $\tilde{\Psi}$. Then he showed likewise that the map $\tilde{J} \circ (\tilde{\Psi} - \Psi_{f'})$ admits a C^0 -small, d_f -Lipschitz fixed point, which is a solution of $(\star\star)$.

Structural stability of covering. We recall that every local diffeomorphism of a compact (connected) manifold is a covering.

F. Przytycki [Prz77] introduced an example of surface covering suggesting the following *strong* transversality condition.

Definition 3.14. A covering map f satisfies axiom A and the strong transversality condition if:

- (i) The non-wandering set is locally maximal.
- (ii) The non-wandering set Ω is the union of a hyperbolic set on which f acts bijectively with a repulsive set.
- (iii) $\forall x \in \Omega, \ y^1, \dots, y^k \in \overleftarrow{\Omega}$, the following multi-transversality condition holds:

$$W^s(x) \pitchfork W^u(y^1) \pitchfork \cdots \pitchfork W^u(y^k)$$
.

We recall that a finite family of submanifolds $(N_i)_i$ is multi-transverse if N_1 and N_2 are transverse, N_3 is transverse to $N_1 \cap N_2$, ..., and for every $i \geq 3$, N_i is transverse to $N_1 \cap N_2 \cap \cdots \cap N_{i-1}$. We notice that (iii) implies (ii).

Here is a generalization of a conjecture of Przytycki [Prz77]:

Conjecture 3.15. The C^1 -struturally stable coverings are those which satisfy axiom A and the strong transversality condition.

The fact that structurally stable coverings are axiom A has been proved by Aoki-Moriyasu-Sumi [AMS01], and the strong transversality condition has been proved by Iglesias-Portela-Rovella. The other direction is still open in the general case.

This conjecture has been proved in two special cases. The first one solves the initial Przytycki conjecture for surface coverings:

Theorem 3.16 (Iglesias-Portela-Rovella [IPR12]). If a covering map of a surface satisfies axiom A and the strong transversality condition then it is C^1 -structurally stable.

The other case is for attractor-repellor covering.

Theorem 3.17 (Iglesias-Portela-Rovella [IPR10]). Let M be a compact manifold. If f is a C^1 -covering map satisfying axiom A, and so that its basic pieces are either bijective attractors or expanding sets, then f is C^1 -structurally stable.

The strong transversality condition for these maps is certainly satisfied since, the unstable manifolds are either included in the attractor or form open subset of the manifold. They gave the following example:

$$f: (z, z') \in \mathbb{S}^1 \times \hat{\mathbb{C}} \mapsto (z^2, z/2 + z'/3),$$

where the non-wandering set consists of an expanding circle and of the Smale solenoid.

In [BK13], we constructed d_f -Lipschitz plane fields for endomorphisms which satisfies axiom A and the strong transversality condition. This might be useful to prove that under the hypothesis of Conjecture 3.15, the following map has a left inverse:

$$\sigma \in \Gamma^0(TM) \mapsto \sigma - Df^{-1} \circ \sigma \circ f \in \Gamma^0(TM)$$
.

3.3 Structural stability of the inverse limit

Given an endomorphism f of a compact manifold M, the inverse limit \overleftarrow{M}_f of f is the space of orbits :

$$\overleftarrow{M}_f := \left\{ \underline{x} = (x_n)_{n \in \mathbb{Z}} \colon x_{n+1} = f(x_n) \right\}.$$

It is a closed subset of $M^{\mathbb{Z}}$, which is compact endowed with the product metric:

$$d(\underline{x},\underline{x}') = \sum_{n \in \mathbb{Z}} 2^{-|n|} d(x_n, x_n') .$$

We notice that the inverse limit is homemorphic to M when f is a homeomorphism of M.

We notice also that the shift dynamics \overleftarrow{f} acts canonically on \overleftarrow{M}_f :

$$\overleftarrow{f}: (x_n)_n \mapsto (x_{n+1})_n.$$

With $\pi_0: (x_n)_n \mapsto x_0$ the zero coordinate projection, it holds:

$$\pi_0 \circ \overleftarrow{f} = f \circ \pi_0.$$

From this one easily deduces that the non-wandering sets $\overleftarrow{\Omega}_f$ and Ω_f of respectively \overleftarrow{f} and f satisfies the following relation:

$$\overleftarrow{\Omega}_f = \Omega_f^{\mathbb{Z}} \cap \overleftarrow{M}_f .$$

Definition 3.18. The endomorphism f is C^r -inverse limit stable if for every C^r -perturbation f' of f, there exists a homeomorphism h from \overline{M}_f onto $\overline{M}_{f'}$ so that:

$$h \circ \overleftarrow{f} = \overleftarrow{f}' \circ h.$$

We can define the unstable manifold of every point $\underline{x} = (x_i)_i \in \overleftarrow{\Omega}_f$:

$$W^{u}(\underline{x}; \overleftarrow{f}) := \{ y = (y_i)_i \in \overleftarrow{M}_f : d(x_i, y_i) \to 0, i \to -\infty \}$$

When f satisfies axiom A, it is an actual manifold embedded in M_f . Moreover, the 0-coordinate projection π_0 displays a differentiable restriction $\pi_0|W^u(\underline{x}; \overleftarrow{f})$.

On the other hand, there exists $\epsilon > 0$ so that the following local stable manifold is an embedded submanifold of M, for every $x \in \Omega_f$:

$$W^s_{\epsilon}(x;f) := \{ y \in M : d(f^n(x), f^n(y)) \to 0, \ n \to +\infty \}.$$

In [BR13], we notice that surprisingly, for certain axiom A endomorphisms, the presence of critical set (made by points with non surjective differential) does not interfere with the C^1 -inverse structural stability. This leads us to define:

Definition 3.19. An axiom A endomorphism f satisfies the weak transversality condition if for every $\underline{x} \in \overleftarrow{\Omega}_f$ and every $y \in \Omega_f$, the map $\pi_0 | W^u(\underline{x}; \overleftarrow{f})$ is transverse to $W^s_{\epsilon}(y)$.

There are many examples of endomorphisms which satisfy axiom A and the weak transversality condition. For instance:

- any axiom A map of the one point compactification $\hat{\mathbb{R}}$ of \mathbb{R} , in particular those of the form $x \mapsto x^2 + c$ and even the constant map $x \mapsto 0$.
- if f_1 and f_2 satisfy axiom A and the weak transversality condition, then the product dynamics (f_1, f_2) do so.
- By the two latter points, note that the map $(x, y, z) \mapsto (x^2, y^2, 0)$ of \mathbb{R}^3 satisfies axiom A and the weak transversality condition.

The latter map is not at all structurally stable, for its critical set is not and intersects moreover the non-wandering set. For this reason the following conjecture might sound irrealistic:

Conjecture 3.20 (Berger-Rovella [BR13]). The C^1 -inverse limit stable endomorphisms are those which satisfy axiom A and the weak transversality condition.

However in [BR13], we gave many evidences of veracity of this conjecture. Then in [BK13] we showed one direction of this conjecture; the other direction is still open.

Theorem 3.21 (Berger-Kocksard [BK13]). If a C^1 -endomorphisms of a compact manifold satisfies axiom A and the weak transversality condition, then it is inverse limit stable.

The proof of this theorem follows the strategy of the Robbin structural stability theorem. The main difficulty is the construction of pseudo-invariant plan fields $(E_i^s)_i$ and $(E_i^u)_i$, for the endomorphisms display in general a non-empty critical set.




Figure 7: Construction of $(E_i^s)_i$ for the map $(x,y,z) \mapsto (x^2,y^2,0)$




Figure 8: Construction of $(E_i^u)_i$ for the map $(x,y,z) \mapsto (x^2,y^2,0)$

4 Links between structural stability in dynamical systems and singularity theory

In the last section we saw how the inverse stability does not seem to involve any singularity theory. However let us notice that if a C^{∞} -endomorphism of a manifold M is structurally stable (that is conjugated to its perturbation via a homeomorphism of M), then its singularities are C^0 -equivalently, structurally stable:

Definition 4.1. Let f be a C^{∞} -map from a manifold M into a possibly different manifold N and $r \in \{0, \infty\}$. The map f is C^r -equivalently, structurally stable if for every f' C^{∞} -close to f, there are $h \in Diff^r(M)$ and $h' \in Diff^r(N)$ which are C^r -close to the identity and such that the following diagram commutes:

$$\begin{array}{cccc} & f' \\ M & \rightarrow & N \\ h & \uparrow & & \uparrow & h' \\ M & \rightarrow & N \\ & f \end{array}$$

The equivalently, structural stability has been deeply studied, in particular by Whitney, Thom and Mather. We shall recall some of the main results, by emphasizing their similarities with those

of structural stability in dynamical systems.

4.1 Infinitesimal stability

Let M, N be compact manifolds. For $r \in \{0, \infty\}$, let $\chi^r(M)$ and $\chi^r(N)$ be the space of C^r -sections of respectively TM and TN.

Definition 4.2. A Diffeomorphism $f \in Diff^1(M)$ is C^0 -infinitesimally stable if the following map is surjective:

$$\sigma \in \chi^0(M) \mapsto Tf \circ \sigma - \sigma \circ f \in \chi^0(f),$$

with $\chi^0(f)$ the space of continuous sections of the pull back bundle f^*TM .

In the Robbin-Robinson proofs of structural stability (Theorem 3.12), we saw the importance of the left-invertibility of $\sigma \mapsto Tf \circ \sigma - \sigma \circ f$. The latter implies the C^0 -infinitesimal stability which is equivalent to the C^1 -structural stability:

Theorem 4.3 (Robin-Robinson-Mañe [Rob71],[Rob76], [Mañ88]). The C^0 -infinitesimally stable diffeomorphisms are the C^1 -equivalently stable maps.

A similar definition exists in Singularity Theory:

Definition 4.4. Let $f \in C^{\infty}(M, N)$ is C^{∞} - equivalently infinitesimally stable if the following map is surjective:

$$(\sigma,\xi)\in\chi^\infty(M)\times\chi^\infty(N)\mapsto Tf\circ\sigma-\xi\circ f\in\chi^\infty(f)$$

with $\chi^{\infty}(f)$ the space of C^{∞} -sections of the pull back bundle f^*TM .

It turns out to be equivalent to the C^{∞} -equivalent stability.

Theorem 4.5 (Mather [Mat68a, Mat69a, Mat68b, Mat69b, Mat70]). The C^{∞} - infinitesimally equivalently stable maps are the C^{∞} -equivalently stable maps.

The latter might sound complicated to verify, but on concrete examples it is rather easy to check. That is why following Mather, it is a satisfactory description of C^{∞} -equivalently structurally stable maps.

4.2 Density structurally stable maps

Let us point out two similar results on structural stability:

Theorem 4.6 (Thom, Mather [Mat73, Mat76, GWdPL76]). For every manifolds M, N, the C^0 -equivalently structural stable maps form an open and dense set in $C^{\infty}(M, N)$.

Let us recall:

Theorem 4.7 (Mañe-Sad-Sullivan [MSS83], Lyubich [Lyu84]). For every $d \ge 2$, the set of structurally stable rational functions is open and dense.

In both cases, we do not know how to describe these structurally stable maps.

Still the axiom A condition is a candidate to describe the structurally stable rational functions, since the famous Fatou conjecture (1920). On the other hand, there is not even a conjecture for the description of the C^0 -equivalently structural stable maps.

Following Mather, a nice way to describe the equivalently structural stable maps would be (a similar way to) the C^{∞} -equivalently infinitesimal stability.

Nevertheless, Mather proved that C^{∞} -equivalently infinitesimal stable maps are dense if and only if the dimensions of M and N are not "nice" [Mat71]. We define the nice dimensions below. Thus one has to imagine a new criteria (at least of in "not nice" dimensions) to describe the C^0 -equivalently structural stable maps.

Definition 4.8 (Nice dimensions). If $m = \dim M$ and $n = \dim N$, the pair of dimensions (m; n) is nice if and only if one of the following conditions holds:

$$n-m \ge 4$$
 and $m < \frac{6}{7}n + \frac{8}{7}$, $3 \ge n - m \ge 0$ and $m < \frac{6}{7}n + \frac{9}{7}$, $n-m = -1$ and $n < 8$, $n-m = -2$ and $n < 6$, $n-m = -3$ and $n < 7$.

We notice that if $n := \dim M = \dim N$, then the pair of dimensions (m; n) is nice if and only if n < 8.

Let us finally recall an open question:

Problem 4.9. In nice dimensions, does a C^0 -equivalently structurally stable map is always C^{∞} -equivalently structurally stable map?

4.3 Geometries of the structural stability

The proof of the Thom-Mather Theorem 4.6 on the density of C^0 -equivalently structurally stable involves the concept of stratification (by analytic or smooth submanifolds).

Similarly, the set of stable and unstable manifolds of an axiom A diffeomorphisms form a stratification of laminations, as defined in [Ber13].

Let us recall these definitions.

4.3.1 Stratifications

A stratification is the pair of a locally compact subset A and a locally finite partition Σ by locally compact subsets $X \subset A$, called strata, and satisfying:

$$\forall (X,Y) \in \Sigma^2, \ cl(X) \cap Y \neq \emptyset \Rightarrow cl(X) \supset Y \ .$$
 We write then $X \geq Y$.

In practical, the set A will be embedded into a manifold M, and the strata X will be endowed with a structure of analytic manifold, differentiable manifold or even lamination, depending on the context.




Figure 9: Algebraic variety $x^2 + y^2 + z^2 + 2xyz - 1 = 0$

4.4 Whitney Stratification

The first use of stratification goes back to the work of Whitney to describe the algebraic varieties. Then it has been generalized by Thom and Lojasiewicz for the study of analytic variety and even semi-analytic variety.

Definition 4.10. An analytic variety of \mathbb{R}^n is the zero set of an analytic function on an open subset of \mathbb{R}^n . An analytic submanifold is a submanifold which is also an analytic variety. A semi-analytic variety is a subset A of \mathbb{R}^n which is covered by open subset U satisfying:

$$A \cap U = \bigcap_{i=1}^{N} \cup_{i=1}^{N} F_{ij}$$

with F_{ij} of the form $\{q_{ij} > 0\}$ or $\{q_{ij} = 0\}$ and q_{ij} a real analytic function on U.

Theorem 4.11 (Whitney-Lojasiewicz [Łoj71]). Any semi-analytic variety $S \subset \mathbb{R}^n$ splits into a stratification Σ by analytic manifolds.

One important property of the semi-analytic category is its stability by projection from the Seidenberg Theorem: given any projection p of $\mathbb{R}^n \to \mathbb{R}^p$, the image by p of any semi-analytic variety is a semi-analytic variety.

4.5 Thom-Mather Stratification

The following is a key step in the proof of the Thom-Mather Theorem 4.6.

Theorem 4.12 (Thom-Mather). For every C^{∞} -generic map from a compact manifold M into N, there exists a stratification on Σ_M of M and a stratification Σ_N on N such that:

- (i) The strata of Σ_M and Σ_N are smooth submanifolds,
- (ii) the restriction of f to each stratum of Σ_M is a submersion onto a stratum of N,

(iii) this stratification is structurally stable: for every perturbation f' of f there are stratifications Σ'_M and Σ'_N homeomorphic to respectively Σ_M and Σ_N , so that (ii) holds for f'.

The proof of this theorem is extremely interesting, it involves in particular the jet space, Thom's transversality theorem and Whitney stratifications in semi-analytics geometry.

4.6 Laminar stratification

Analogously to singularity theory, a structurally stable C^1 -diffeomorphism displays a stratification.

Definition 4.13. A lamination of M is a locally compact subset \mathcal{L} of M, which is locally homeomorphic to the product of a \mathbb{R}^d with a locally compact set T, so that $(\mathbb{R}^d \times \{t\})_{t \in T}$ corresponds to a continuous family of submanifolds.

Definition 4.14. A stratification of laminations is a stratification whose strata are endowed with a structure of lamination.

Proposition 4.15 ([Ber13]). Let f be a diffeomorphism M which satisfies Axiom A and the strong transversality condition. Then the stable set of every basic piece Λ_i of f has a structure of lamination X_i whose leaves are stable manifolds. Moreover the family $\Sigma_s := (X_i)_i$ forms a stratification of laminations such that $X_i \leq X_j$ iff $\Lambda_i \succeq \Lambda_j$ i.e. $W^u(\Lambda_i) \cap W^s(\Lambda_j) \neq \emptyset$.

5 Structural stability of endomorphisms with singularities

We are now ready to study the structural stability of endomorphisms which display a non empty critical set.

In dimension 2, Iglesias-Portela-Rovella [IPR08] showed the structural stability of C^3 -perturbations of the hyperbolic rational functions f which are equivalently stable and whose critical sets do not self-intersect along their orbits, nor intersect the non-wandering set.

In all these examples, the critical set does not self intersect along its orbit. J. Mather suggested me to generalize a study he did about structural stability of graph of maps.

Let G := (V, A) be a finite oriented graph with a manifold M_i associated to each vertex $i \in V$, and with a smooth map $f_{ij} \in C^{\infty}(M_i, M_j)$ associated to each arrow $[i, j] \in A$ from i to j.

For $k \in \{0, \infty\}$, such a graph is C^k -structurally stable if for every C^∞ -perturbation $(f'_{ij})_{[i,j]\in A}$ of $(f_{ij})_{[i,j]\in A}$, there exists a family of C^k diffeomorphisms $(h_i)_i \in \prod_{i\in V} Diff^k(M_i, M_i)$ such that the following diagram commutes:

$$\forall [i,j] \in A \quad \begin{array}{ccc} & f'_{ij} \\ M_i & \rightarrow & M_j \\ h_i & \uparrow & f_{ij} & \uparrow & h_j \\ M_i & \rightarrow & M_j \end{array}.$$

The graph (V, A) is convergent if for every $[i, j], [i', j'] \in A$ if i = i' then j = j'. The graph is without cycle if for every $n \ge 1$ and every $([i_k, i_{k+1}])_{0 \le k < n} \in V^n$ it holds $i_n \ne i_0$.

Theorem 5.1 (Mather). Let G be a graph of smooth proper maps, convergent and without cycle. The graph is C^{∞} - structurally stable if the following map is surjective:

$$(\sigma_i)_i \in \prod_{i \in V} \chi^{\infty}(M_i) \mapsto (Tf_{ij} \circ \sigma_i - \sigma_j \circ f_{ij})_{[ij]} \in \prod_{[i,j] \in A} \chi^{\infty}(f_{ij}).$$

Mather gave me an unpublished manuscript of Baas relating his proof, that I developed to study the structural stability of attractor-repellor endomorphisms with possibly a non-empty critical set.

Definition 5.2. Let f be a smooth endomorphism of a compact, non necessarily connected manifold. The endomorphism f is attractor-repellor if it satisfies axiom A, and its basic pieces are either expanding pieces or attractors which f acts bijectively.

The following theorem generalizes all the results I know (including [IPR08] and [IPR10]) about structurally stable maps with non-empty critical set.

Theorem 5.3 (Berger [Ber12]). Let f be an attractor-repellor, smooth endomorphism of a compact, non necessarily connected manifold M. If the following conditions are satisfied, then f is C^{∞} -structurally stable:

- (i) the singularities S of f have their orbits that do not intersect the non-wandering set Ω ,
- (ii) the restriction of f to $M \setminus \hat{\Omega}$ is C^{∞} -infinitesimally stable, with $\hat{\Omega} := cl(\cup_{n \geq 0} f^{-n}(\Omega))$. In other words, the following map is surjective:

$$\sigma \in \Gamma^{\infty}(M) \mapsto Df \circ \sigma - \sigma \circ f \in \Gamma^{\infty}(f)$$

(iii) f is transverse to the stable manifold of A's points: for any $y \in A$, for any point z in a local stable manifold W_y^s of y, for any $n \ge 0$, and for any $x \in f^{-n}(\{z\})$, we have:

$$Tf^n(T_xM) + T_zW_y^s = T_zM.$$

Hypothesis (ii) might seem difficult to verify, but it is not. In [Ber12] we apply it to many example, even for map for which the critical set does self intersect along its orbit.

It would be intersecting to investigate how the attractor-repeller could be relaxed to enjoy a greater generality. However the C^0 -equivalently stable singularities are not well classified and so an optimal theorem is today difficult to obtain. Nevertheless, it is not the case in dimension 2. Indeed it is well known that the structurally stable singularities are locally equivalent to one of the following polynomial (called resp. fold and cusp):

$$(x,y)\mapsto (x^2,y)$$
 and $(x,y)\mapsto (x^3+xy,y)$.

Hence here is a natural question:

Problem 5.4. Under which hypothesis an axiom A surface endomorphism with singularity is structurally stable?

References

- [AMS01] N. Aoki, K. Moriyasu, and N. Sumi. C^1 -maps having hyperbolic periodic points. Fund. Math., 169(1):1–49, 2001.
- [Ano67] D. V. Anosov. Geodesic flows on closed Riemannian manifolds of negative curvature. Trudy Mat. Inst. Steklov., 90:209, 1967.
- [Aok92] N. Aoki. The set of Axiom A diffeomorphisms with no cycles. *Bol. Soc. Brasil. Mat.* (N.S.), 23(1-2):21–65, 1992.
- [BD17] P. Berger and R. Dujardin. On stability and hyperbolicity for polynomial automorphisms of \mathbb{C}^2 . Ann. Sci. École Norm. Sup. (4), 2017.
- [Ber12] P. Berger. Structural stability of attractor-repellor endomorphisms with singularities. Ergodic Theory Dynam. Systems, 32(1):1–33, 2012.
- [Ber13] P. Berger. Persistence of stratifications of normally expanded laminations. *Mém. Soc. Math. Fr. (N.S.)*, (134):113, 2013.
- [BK13] P. Berger and A. Kocsard. Structural stability of the inverse limit of endomorphisms. $ArXiv\ e\text{-}prints,\ 2013.$
- [BR13] P. Berger and A. Rovella. On the inverse limit stability of endomorphisms. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 30(3):463–475, 2013.
- [Buz97] G. T. Buzzard. Infinitely many periodic attractors for holomorphic maps of 2 variables. Ann. of Math. (2), 145(2):389–417, 1997.
- [BY14] P. Berger and J.-C. Yoccoz. Strong regularity. *Manuscript*, 2014.
- [Cou06] Yves Coudene. Pictures of hyperbolic dynamical systems. *Notices Amer. Math. Soc.*, 53(1):8–13, 2006.
- [CP15] S. Crovisier and E. R. Pujals. Essential hyperbolicity and homoclinic bifurcations: a dichotomy phenomenon/mechanism for diffeomorphisms. *Invent. Math.*, 201(2):385–517, 2015.
- [DL] R. Dujardin and M. Lyubich. Stability and bifurcations for dissipative polynomial automorphisms of \mathbb{C}^2 . to appear in Invent. Math.
- [GŚ97] J. Graczyk and G. Światek. Generic hyperbolicity in the logistic family. *Ann. of Math.* (2), 146(1):1–52, 1997.
- [GWdPL76] C. G. Gibson, K. Wirthmüller, A. A. du Plessis, and E. J. N. Looijenga. Topological stability of smooth mappings. Lecture Notes in Mathematics, Vol. 552. Springer-Verlag, Berlin-New York, 1976.

- [Hay92] S. Hayashi. Diffeomorphisms in $\mathcal{F}^1(M)$ satisfy Axiom A. Ergodic Theory Dynam. Systems, 12(2):233–253, 1992.
- [IPR08] J. Iglesias, A. Portela, and A. Rovella. Structurally stable perturbations of polynomials in the Riemann sphere. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 25(6):1209–1220, 2008.
- [IPR10] J. Iglesias, A. Portela, and A. Rovella. C^1 stable maps: examples without saddles. Fund. Math., 208(1):23–33, 2010.
- [IPR12] J. Iglesias, A. Portela, and A. Rovella. C^1 stability of endomorphisms on two-dimensional manifolds. Fund. Math., 219(1):37–58, 2012.
- [Łoj71] S. Łojasiewicz. Sur les ensembles semi-analytiques. In Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2, pages 237–241. Gauthier-Villars, Paris, 1971.
- [Lyu84] M. Lyubich. Investigation of the stability of the dynamics of rational functions. *Teor. Funktsing Funktsional. Anal. i Prilozhen.*, (42):72–91, 1984. Translated in Selecta Math. Soviet. **9** (1990), no. 1, 69–90.
- [Lyu97] M. Lyubich. Dynamics of quadratic polynomials. I, II. Acta Math., 178(2):185–247, 247–297, 1997.
- [Mañ82] Ricardo Mañé. An ergodic closing lemma. Ann. of Math. (2), 116(3):503-540, 1982.
- [Mañ88] R. Mañé. A proof of the C^1 stability conjecture. Inst. Hautes Études Sci. Publ. Math., (66):161–210, 1988.
- [Mat68a] J. N. Mather. Stability of C^{∞} mappings. I. The division theorem. Ann. of Math. (2), 87:89–104, 1968.
- [Mat68b] J. N. Mather. Stability of C^{∞} mappings. III. Finitely determined mapgerms. Inst. Hautes Études Sci. Publ. Math., (35):279–308, 1968.
- [Mat69a] J. N. Mather. Stability of C^{∞} mappings. II. Infinitesimal stability implies stability. Ann. of Math. (2), 89:254–291, 1969.
- [Mat69b] J. N. Mather. Stability of C^{∞} mappings. IV. Classification of stable germs by R-algebras. Inst. Hautes Études Sci. Publ. Math., (37):223–248, 1969.
- [Mat70] J. N. Mather. Stability of C^{∞} mappings. V. Transversality. Advances in Math., 4:301–336 (1970), 1970.
- [Mat71] J. N. Mather. Stability of C^{∞} mappings. VI: The nice dimensions. In *Proceedings* of Liverpool Singularities-Symposium, I (1969/70), pages 207–253. Lecture Notes in Math., Vol. 192. Springer, Berlin, 1971.

- [Mat73] J. N. Mather. Stratifications and mappings. In *Dynamical systems (Proc. Sympos., Univ. Bahia, Salvador, 1971)*, pages 195–232. Academic Press, New York, 1973.
- [Mat76] J. N. Mather. How to stratify mappings and jet spaces. In Singularités d'applications différentiables (Sém., Plans-sur-Bex, 1975), pages 128–176. Lecture Notes in Math., Vol. 535. Springer, Berlin, 1976.
- [Mos69] J. Moser. On a theorem of Anosov. J. Differential Equations, 5:411–440, 1969.
- [MSS83] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. Ann. Sci. École Norm. Sup. (4), 16(2):193–217, 1983.
- [Pal88] J. Palis. On the C^1 Ω -stability conjecture. Inst. Hautes Études Sci. Publ. Math., (66):211–215, 1988.
- [Pal00] J. Palis. A global view of dynamics and a conjecture on the denseness of finitude of attractors. *Astérisque*, (261):xiii–xiv, 335–347, 2000.
- [Prz77] F. Przytycki. On *U*-stability and structural stability of endomorphisms satisfying Axiom A. *Studia Math.*, 60(1):61–77, 1977.
- [PS70] J. Palis and S. Smale. Structural stability theorems. In Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968), pages 223–231. Amer. Math. Soc., Providence, R.I., 1970.
- [PS00] E. R. Pujals and M. Sambarino. Homoclinic tangencies and hyperbolicity for surface diffeomorphisms. *Ann. of Math.* (2), 151(3):961–1023, 2000.
- [Rob71] J. W. Robbin. A structural stability theorem. Ann. of Math. (2), 94:447–493, 1971.
- [Rob76] C. Robinson. Structural stability of C^1 diffeomorphisms. J. Differential Equations, 22(1):28-73, 1976.
- [Shu69] M. Shub. Endomorphisms of compact differentiable manifolds. *Amer. J. Math.*, 91:175–199, 1969.
- [Shu78] M. Shub. Stabilité globale des systèmes dynamiques, volume 56 of Astérisque. Société Mathématique de France, Paris, 1978.
- [Sma70] S. Smale. The Ω-stability theorem. In Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968), pages 289–297. Amer. Math. Soc., Providence, r.I., 1970.
- [Yoc95] J.-C. Yoccoz. Introduction to hyperbolic dynamics. In *Real and complex dynamical systems (Hillerød, 1993)*, volume 464 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 265–291. 1995.