

Relating quantum coherence and correlations with entropy-based measures

Xiao-Li Wang · Qiu-Ling Yue ·
Chao-Hua Yu · Fei Gao · Su-Juan Qin

Received: date / Accepted: date

Abstract Quantum coherence as an important quantum resource plays a key role in quantum theory. In this paper, using entropy-based measures, we investigate the relations between quantum correlated coherence, which is the coherence between subsystems [K. C. Tan, H. Kwon, C. Y. Park, and H. Jeong, Phys. Rev. A **94**, 022329 (2016)], and two main kinds of quantum correlations as defined by quantum discord as well as quantum entanglement. In particular, we show that quantum discord and quantum entanglement can be well characterized by quantum correlated coherence. Moreover, we prove that the entanglement measure formulated by quantum correlated coherence is lower and upper bounded by the relative entropy of entanglement and the entanglement of formation, respectively, and equal to the relative entropy of entanglement for maximally correlated states.

Keywords quantum coherence · quantum correlated coherence · quantum discord · quantum entanglement

1 Introduction

Quantum coherence arising from quantum superposition [1], represents a fundamental feature that marks the departure of quantum mechanics from classical physics. Recently, many efforts have been devoted to develop the resource

X.-L. Wang · Q.-L. Yue · C.-H. Yu · F. Gao · S.-J. Qin
State Key Laboratory of Networking and Switching Technology, Beijing University of Posts and Telecommunications, Beijing 100876, China
E-mail: qsujuan@bupt.edu.cn

X.-L. Wang
School of Mathematics and Information Science, Henan Polytechnic University, Jiaozuo, 454000, China
State Key Laboratory of Information Security (Institute of Information Engineering, Chinese Academy of Sciences, Beijing 100093)

theory of quantum coherence [2-7]. Meanwhile, various properties of quantum coherence have been investigated such as the connections between quantum coherence and quantum correlations in multipartite systems [8-13], the distillation of coherence [5, 14, 15], the dynamics under noisy evolution of quantum coherence [16, 17], among others. The role of coherence in thermodynamics [18, 19] has also been discussed.

Quantum coherence in multipartite systems involves the essence of quantum correlations. One of the potential quantum correlations is quantum entanglement [20-24] which has been widely concerned. Another kind of quantum correlations is quantum discord [25-29], which may even exist in a separable state with vanished entanglement. Both of them are crucial resources for the development of quantum technologies, such as quantum communication [30, 31], quantum computation [32, 33], quantum metrology [34], and many more. In these cases, quantum correlations indicate an advantage of quantum methods over classical ones.

Note that previous results in Ref. [13] have established a unified view of quantum discord and quantum entanglement with the framework of quantum coherence based on the l_1 -norm of coherence. By contrast, we will adopt the relative entropy of coherence to explore the concise relations between quantum discord and quantum correlated coherence [13], which is the coherence between subsystems. In fact, quantum correlated coherence is a ‘correlation function’, which captures the correlation between subsystems. Besides, quantum correlated coherence can be used to construct an entanglement measure, which is called the entanglement of coherence (EOC) [13]. However, Many important properties, such as additivity, and relations to other entanglement measures, have not been investigated. In this paper, using entropy-based measures, we will show that the EOC is lower and upper bounded by the relative entropy of entanglement and the entanglement of formation, and equal to the relative entropy of entanglement for maximally correlated states. We also compare the EOC with the entanglement measure which is the minimal discord over state extensions [35]. Our work provides clear relations between quantum coherence and correlations with entropy-based measures.

The rest of this paper is organized as follows. Sec. 2 introduces some definitions of quantum coherence, quantum entanglement and quantum discord. In Sec. 3, we give the relations between quantum correlated coherence and quantum discord. In Sec.4, we prove the bounds of the entanglement of coherence (EOC), which is formulated by quantum correlated coherence with respect to the relative entropy of coherence. This paper is ended with the conclusion in Sec. 5.

2 Preliminaries

In the framework of coherence theory introduced in Ref. [4], let $\{|i\rangle\}$ be a fixed basis in the finite dimensional Hilbert space, and the incoherent states are those whose density matrices are diagonal in this fixed basis, being of the

form $\sum_i p_i |i\rangle\langle i|$ where p_i are probabilities. The set of all incoherent states is denoted by \mathcal{I} . It is known that quantum operations are characterized by a set of Kraus operators $\{K_l\}$ satisfying $\sum_l K_l^\dagger K_l = I$. In particular, an incoherent quantum operation is that for which there exists a Kraus representation $\{K_l\}$ such that $\frac{K_l \sigma K_l^\dagger}{\text{Tr}(K_l \sigma K_l^\dagger)} \in \mathcal{I}$ for all l and all $\sigma \in \mathcal{I}$. The von Neumann measurement with respect to the fixed basis $\{|i\rangle\}$ (otherwise called the dephasing operation) is a special incoherent quantum operation denoted by $\Pi = \{|i\rangle\langle i|\}$. For any state ρ , we have $\Pi(\rho) = \rho^{diag} = \sum_i |i\rangle\langle i| \rho |i\rangle\langle i|$. Remarkably, any state ρ will generate an incoherent state ρ^{diag} by removing all off-diagonal terms from its density matrix in the fixed basis through the von Neumann measurement Π . In this paper, we will employ the relative entropy of coherence as the coherence measure which is defined as $\mathcal{C}_{re}(\rho) = \min_{\sigma \in \mathcal{I}} S(\rho||\sigma)$, where $S(\rho||\sigma) = \text{Tr}(\rho \log_2 \rho) - \text{Tr}(\rho \log_2 \sigma)$ is the quantum relative entropy [36] and the minimization is taken over the set of incoherent states \mathcal{I} . It has been shown that $\mathcal{C}_{re}(\cdot)$ satisfies all the conditions mentioned in Ref. [4]. Crucially, this quantity can be evaluated exactly: $\mathcal{C}_{re}(\rho) = S(\rho^{diag}) - S(\rho)$, where $S(\rho) = -\text{Tr}(\rho \log_2 \rho)$ is the von Neumann entropy [36].

In this paper, unless otherwise stated, we will often refer to a bipartite system denoted by AB , where A and B are local subsystems. For convenience, we say the subsystems A and B are held by Alice and Bob, respectively. For a given state ρ_{AB} in system AB , the local states of Alice and Bob are denoted by $\rho_A = \text{Tr}_B(\rho_{AB})$ and $\rho_B = \text{Tr}_A(\rho_{AB})$, respectively, which are obtained by performing a partial trace on ρ_{AB} .

Quantum entanglement [20-24] is a popular kind of quantum correlations which can not be prepared by local operations and classical communication (LOCC). The states prepared by LOCC are called separable states which can be represented as a convex combination of product states, i.e., $\sigma_{AB} = \sum_k p_k |\varphi_k\rangle_A \langle \varphi_k| \otimes |\phi_k\rangle_B \langle \phi_k|$, where $\{|\varphi_k\rangle_A\}$ and $\{|\phi_k\rangle_B\}$ are normalized but not necessarily orthogonal. Here, we will refer to the relative entropy of entanglement defined as $E_{re}(\rho_{AB}) = \min_{\sigma \in \mathcal{S}} S(\rho||\sigma)$ with the minimization over the set of separable states \mathcal{S} [21, 22]. Another closely related quantity is entanglement of formation defined as $E_f(\rho_{AB}) = \min_{\{p_k, |\psi_k\rangle\}} \sum_k p_k E_{re}(|\psi_k\rangle\langle \psi_k|)$, where the minimization is taken over all decompositions of the state $\rho_{AB} = \sum_k p_k |\psi_k\rangle\langle \psi_k|$ [20].

Quantum discord measures the disturbance induced by local operations to multipartite states [25-29]. Let $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ be orthonormal bases of subsystems A and B , respectively. If Bob performs the von Neumann measurement $\Pi_B = \{|j\rangle_B \langle j|\}$ on his subsystem, the post-measurement state is denoted as

$$\Pi_B(\rho_{AB}) = \sum_j (I_A \otimes |j\rangle_B \langle j|) \rho_{AB} (I_A \otimes |j\rangle_B \langle j|). \quad (1)$$

The asymmetric quantum discord with respect to Π_B can be written in terms of a difference of relative entropies [25-27],

$$D_{\Pi_B}^{A|B}(\rho_{AB}) = S(\rho_{AB} || \rho_A \otimes \rho_B) - S(\Pi_B(\rho_{AB}) || \rho_A \otimes \Pi_B(\rho_B)). \quad (2)$$

In the classical-quantum dichotomy, the asymmetric quantum discord is defined as $D^{A|B}(\rho_{AB}) = \min_{\Pi_B} S(\rho_{AB} \| \rho_A \otimes \rho_B)$ with the minimization over all local von Neumann measurements Π_B to remove the measurement-basis dependence. If Alice only performs the von Neumann measurement $\Pi_A = \{|i\rangle_A \langle i|\}$ on her subsystem, the similar results are available. If both Alice and Bob perform local von Neumann measurements Π_A and Π_B on their respective subsystems, the symmetric quantum discord (global quantum discord in bipartite system [27]) with respect to $\Pi_A \otimes \Pi_B$ is defined as:

$$D_{\Pi_A \otimes \Pi_B}(\rho_{AB}) = S(\rho_{AB} \| \rho_A \otimes \rho_B) - S(\Pi_A \otimes \Pi_B(\rho_{AB}) \| \Pi_A(\rho_A) \otimes \Pi_B(\rho_B)).$$

Then, the standard form of symmetric quantum discord is defined as $D(\rho_{AB}) = \min_{\Pi_A \otimes \Pi_B} D_{\Pi_A \otimes \Pi_B}(\rho_{AB})$, with the minimization over all the local von Neumann measurements of Alice and Bob.

Recently, Yadin et al. [37] have studied the asymmetric basis-dependent discord $D_{\Pi_B}^{A|B}(\cdot)$ which can be seen as the basis-dependent measure of quantumness of correlation, and the properties of $D_{\Pi_B}^{A|B}(\cdot)$ under the strictly incoherent operations are investigated. Here, we will relate the basis-dependent discord and quantum correlated coherence.

3 Quantum correlated coherence and quantum discord

Let $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ be the fixed bases of subsystems A and B respectively, and we usually use their tensor product $\{|ij\rangle_{AB}\}$ as the fixed basis of the composite system AB . For a state ρ_{AB} in system AB , its total coherence is $\mathcal{C}_{re}(\rho_{AB})$, while $\mathcal{C}_{re}(\rho_A)$ and $\mathcal{C}_{re}(\rho_B)$ are known as local coherences. Whenever ρ_{AB} is a product state, the sum of local coherences is equal to the total coherence. Generally, the relative entropy of coherence admits the super-additive property [11]

$$\mathcal{C}_{re}(\rho_{AB}) \geq \mathcal{C}_{re}(\rho_A) + \mathcal{C}_{re}(\rho_B). \quad (3)$$

Thus, the definition of quantum correlated coherence with respect to the relative entropy of coherence is given as the following.

Definition 1. (K. C. Tan et al. [13]) Let $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ be the fixed bases of subsystems A and B respectively. For a given state ρ_{AB} in system AB , its quantum correlated coherence is defined as

$$\mathcal{C}_{re}^{cc}(\rho_{AB}) \equiv \mathcal{C}_{re}(\rho_{AB}) - \mathcal{C}_{re}(\rho_A) - \mathcal{C}_{re}(\rho_B). \quad (4)$$

Obviously, quantum correlated coherence is the total coherence between subsystems and non-negative. In fact, quantum correlated coherence is a ‘correlation function’ which is similar as quantum mutual information [36]. For arbitrary fixed bases of subsystems A and B , the quantum correlated coherence of ρ_{AB} vanishes if and only if ρ_{AB} has no correlations, i.e., $\rho_{AB} = \rho_A \otimes \rho_B$. The ‘only if’ part is directly derived from the theorem 2 below. In this sense, quantum correlated coherence can be seen as the basis-dependent measure of

quantumness of correlation and accounts for quantum correlations, for example, quantum discord.

With respect to the fixed bases of $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ of subsystems A and B respectively, the local von Neumann measurements of Alice and Bob are denoted by $\Pi_A = \{|i\rangle_A\langle i|\}$ and $\Pi_B = \{|j\rangle_B\langle j|\}$, respectively. By direct calculation, we get that the consumption of quantum correlated coherence for any state ρ_{AB} under Π_B coincides with the asymmetric basis-dependent discord $D_{\Pi_B}^{A|B}(\rho_{AB})$, i.e., $\mathcal{C}_{re}^{cc}(\rho_{AB}) - \mathcal{C}_{re}^{cc}(\Pi_B(\rho_{AB})) = D_{\Pi_B}^{A|B}(\rho_{AB})$. According to the condition for the asymmetric basis-dependent discord $D_{\Pi_B}^{A|B}(\rho_{AB})$ to vanish [37], we have the following result.

Theorem 1. Let $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ be the fixed bases of subsystems A and B , respectively, and the local von Neumann measurement in the basis $\{|j\rangle_B\}$ is denoted as Π_B . For a given state ρ_{AB} in system AB , its quantum correlated coherence remains unchanged under Π_B , i.e., $\mathcal{C}_{re}^{cc}(\rho_{AB}) = \mathcal{C}_{re}^{cc}(\Pi_B(\rho_{AB}))$, if and only if there exists a decomposition $\rho_{AB} = \sum_{\alpha} p_{\alpha} \rho_A^{\alpha} \otimes \rho_B^{\alpha}$ such that p_{α} are probabilities and all the states ρ_B^{α} are perfectly distinguishable by the von Neumann measurement in the fixed basis $\{|j\rangle_B\}$.

In theorem 1, two different states, which are perfectly distinguishable by the von Neumann measurement in the fixed basis $\{|j\rangle_B\}$, must have disjoint coherence support. The coherence support of a state is the set of incoherent basis vectors which have nonzero overlap with the state [37].

Using the very similar arguments as $D_{\Pi_B}^{A|B}(\rho_{AB})$, we obtain that quantum correlated coherence is corresponding to the symmetric basis-dependent discord $\mathcal{C}_{re}^{cc}(\rho_{AB}) = D_{\Pi_A \otimes \Pi_B}(\rho_{AB})$. Moreover, we also find the condition for quantum correlated coherence (the symmetric basis-dependent discord) to vanish.

Theorem 2. Let $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ be the fixed bases of subsystems A and B , respectively. For a given state ρ_{AB} in system AB , its quantum correlated coherence is equal to zero, i.e., $\mathcal{C}_{re}^{cc}(\rho_{AB}) = 0$, if and only if there exists a decomposition,

$$\rho_{AB} = \sum_{k,l} p_{kl} \rho_A^k \otimes \rho_B^l, \quad (5)$$

such that p_{kl} are probabilities, and all the states ρ_A^k and ρ_B^l are perfectly distinguishable by the local von Neumann measurements with respect to the fixed bases $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$, respectively.

Proof. To prove the sufficiency, we will use the following property of von Neumann entropy [36],

$$S\left(\sum_i p_i \rho_i\right) = h(\{p_i\}) + \sum_i p_i S(\rho_i), \quad (6)$$

where $h(\{p_i\})$ is Shannon entropy and all ρ_i have support on orthogonal subspaces. Since all ρ_A^k and ρ_B^l are perfectly distinguishable by the local von Neumann measurements in the fixed bases $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$, respectively,

$\{\rho_A^{k(l)}\}$, $\{\rho_A^k \otimes \rho_B^l\}$, $\{\rho_A^{k(l)diag}\}$, and $\{\rho_A^{kdiag} \otimes \rho_B^{ldiag}\}$ are sets of states with support on orthogonal subspaces. Direct calculation shows that $\mathcal{C}_{re}^{cc}(\rho_{AB}) = 0$.

Note that

$$\mathcal{C}_{re}^{cc}(\rho_{AB}) = S(\rho_{AB} \| \rho_A \otimes \rho_B) - S(\Pi_A \otimes \Pi_B(\rho_{AB}) \| \Pi_A(\rho_A) \otimes \Pi_B(\rho_B)),$$

where Π_A and Π_B are the local von Neumann measurements in the fixed bases $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$, respectively. To prove the necessity, we will use the condition for the quantum relative entropy which is unchanged under a quantum operation [38, 39] and the explicit proof is presented in the Appendix.

Theorem 2 has several implications. First, it implies that a state with vanished quantum correlated coherence is a especially classical-classical state [40] but not necessary to be a bipartite incoherent state [8, 13]. Particularly, a qubit-qubit state with vanished quantum correlated coherence is a product states or a bipartite incoherent state. More complex cases only emerge in higher dimension. For example, the following qutrit-qutrit state with vanished quantum correlated coherence, has yet local coherences:

$$\rho_{AB} = \frac{1}{2} \left(\frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |_{+01}\rangle\langle +_{01}| \right) \otimes \left(\frac{2}{3} |0\rangle\langle 0| + \frac{1}{3} |_{+02}\rangle\langle +_{02}| \right) + \frac{1}{2} |2\rangle\langle 2| \otimes |1\rangle\langle 1|,$$

where $|_{+ij}\rangle = \frac{1}{\sqrt{2}}(|i\rangle + |j\rangle)$ and the fixed basis of each subsystem is computable basis. Second, theorem 2 gives the structure of bipartite states which satisfy the super-additive property of the relative entropy of coherence with equality. This settles an important question left open in previous literature [11, 41] that whether the equality of formula (3) holds if and only if ρ_{AB} is product or incoherent. Finally, if we choose the local eigenbases of ρ_A and ρ_B as the fixed bases of subsystems A and B respectively, these two theorems reduce to the corresponding results in Ref. [13]. In this sense, our results somewhat generalize the previous results.

The above results show that quantum correlated coherence and the basis-dependent discord are closely related. With equality $\mathcal{C}_{re}^{cc}(\rho_{AB}) = D_{\Pi_A \otimes \Pi_B}(\rho_{AB})$, the symmetric quantum discord is rewritten as $D(\rho_{AB}) = \min_{\{|i\rangle_A, |j\rangle_B\}} \mathcal{C}_{re}^{cc}(\rho_{AB})$. Recall that the symmetric quantum discord based on the pseudo distance of relative entropy can be represented with the relative entropy of coherence, i.e., $\mathcal{C}^{free}(\rho_{AB}) = \min_{\{|i\rangle_A, |j\rangle_B\}} \mathcal{C}_{re}(\rho_{AB})$ [12, 28]. Both of the minimization are over all generic bases $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ of subsystems A and B , respectively. However, one may also consider defining a discord measure $D_{POVM}(\cdot)$ via general local positive-operator-valued measurements (POVMs) on each subsystem [25, 29]. Comparing these three quantifiers of quantum discord, we easily have the inequality $\mathcal{C}^{free}(\rho_{AB}) \geq D(\rho_{AB}) \geq D_{POVM}(\rho_{AB})$. Whenever these three quantities are zero, the corresponding states are classical-classical states [26, 40]. Similarly, the asymmetric quantum discord $D^{A|B}(\rho_{AB})$ can also be represented by quantum correlated coherence.

In multipartite systems, the global quantum discord (GQD) [27] can even be rewritten with quantum correlated coherence. It is worth noting that the

fixed basis of a multipartite system is the tensor product of the fixed bases of all the subsystems. For a N -partite state $\rho_{C_1 C_2 \dots C_N}$, its GQD is rewritten as

$$D(\rho_{C_1 C_2 \dots C_N}) = \min_{\{|i_1\rangle_{C_1}, |i_2\rangle_{C_2}, \dots, |i_N\rangle_{C_N}\}} \mathcal{C}_{re}^{cc}(\rho_{C_1 C_2 \dots C_N}),$$

where the minimization is taken over all generic fixed bases of the N -partite system denoted as $\{|i_1\rangle_{C_1} |i_2\rangle_{C_2} \dots |i_N\rangle_{C_N}\}$, and with respect to this fixed basis it holds that $\mathcal{C}_{re}^{cc}(\rho_{C_1 C_2 \dots C_N}) = \mathcal{C}_{re}(\rho_{C_1 C_2 \dots C_N}) - \sum_i \mathcal{C}_{re}(\rho_{C_i})$. With the super-additive property of the relative entropy of coherence as given in formula (3), the GQD is non-negative and for a multipartite classical state it is equal to zero. This provides a simple proof of the non-negativity of GQD in Ref. [27]. These results mean that quantum discord in multipartite systems can be better understood with the framework of quantum coherence.

4 Quantum correlated coherence and quantum entanglement

According to the above discussion, we know that if for arbitrary fixed bases of subsystems A and B the quantum correlated coherence of ρ_{AB} does not vanish there must exist some quantum correlation between subsystems A and B , for example, quantum discord. Moreover, it is also possible to characterize entanglement with quantum correlated coherence via state extensions [13], and then entanglement can be seen as the irreducible residue of quantum correlated coherence. This highlights the non-locality of quantum entanglement.

For a given state ρ_{AB} in system AB , a bipartite state $\rho_{AA'BB'}$ is an extension of ρ_{AB} if $\rho_{AA'BB'}$ satisfies $\text{Tr}_{A'B'}(\rho_{AA'BB'}) = \rho_{AB}$, where subsystems AA' and BB' are held by Alice and Bob, respectively [13, 42]. Via state extensions, the entanglement of ρ_{AB} formulated by quantum correlated coherence is given by definition 2 below.

Definition 2. (K. C. Tan et al. [13]) For a given state ρ_{AB} , $\rho_{AA'BB'}$ is its unitarily symmetric extension and let the local eigenbases of $\rho_{AA'}$ and $\rho_{BB'}$ be the fixed bases of subsystems AA' and BB' , respectively. The entanglement of coherence (EOC) of ρ_{AB} is defined as

$$E_{re}^{cc}(\rho_{AB}) \equiv \min \mathcal{C}_{re}^{cc}(\rho_{AA'BB'}), \quad (7)$$

where the minimization is taken over all possible unitarily symmetric extensions $\rho_{AA'BB'}$.

In definition 2, the extension $\rho_{AA'BB'}$ is unitarily symmetric if it remains invariant up to local unitary operations on AA' and BB' under a system swap between Alice and Bob. It has been shown that the EOC has the properties [13]: non-negative and vanished for separated states, invariant under local unitary operations, non-increasing under LOCC operations, and convex. Furthermore, using entropy-based measures, we can even give the bounds of EOC as the following theorem.

Theorem 3. For a given state ρ_{AB} , it holds that

$$E_{re}(\rho_{AB}) \leq E_{re}^{cc}(\rho_{AB}) \leq E_f(\rho_{AB}). \quad (8)$$

If ρ_{AB} is a pure state, these three quantities in inequality (8) are equal.

Proof. Taking some unitarily symmetric extension $\rho_{AA'BB'}$ of ρ_{AB} , we have

$$\mathcal{C}_{re}^{cc}(\rho_{AA'BB'}) = \mathcal{C}_{re}(\rho_{AA'BB'}) \geq E_{re}(\rho_{AA'BB'}) \geq E_{re}(\rho_{AB}), \quad (9)$$

where the first inequality is due to that the relative entropy of coherence is no less than the relative entropy of entanglement for a state [12], and the last inequality is due to that entanglement is un-increased under LOCC operations [21, 22]. With the definition of EOC, formula (9) means that $E_{re}(\rho_{AB}) \leq E_{re}^{cc}(\rho_{AB})$.

To prove the inequality $E_{re}^{cc}(\rho_{AB}) \leq E_f(\rho_{AB})$, we consider the optimal decomposition of the state $\rho_{AB} = \sum_i p_i^* |\psi_i^*\rangle\langle\psi_i^*|$ such that [20] $E_f(\rho_{AB}) = \sum_i p_i^* E_{re}(|\psi_i^*\rangle\langle\psi_i^*|)$. Every state $|\psi_i^*\rangle$ is represented with the Schmidt decomposition $|\psi_i^*\rangle = \sum_{j_i} \lambda_{j_i} |j_i\rangle_A |j_i\rangle_B$. For every i , $\{|j_i\rangle_A\}$ and $\{|j_i\rangle_B\}$ are expanded to be the orthonormal bases of subsystems A and B , respectively, but both of them are still labeled with original symbols. Define the state

$$\rho_{AA'BB'}^\Delta \equiv \sum_i p_i^* \sum_{j_i, j_i'} \lambda_{j_i} \lambda_{j_i'} |j_i\rangle_A \langle j_i'| \otimes |i\rangle_{A'} \langle i| \otimes |j_i\rangle_B \langle j_i'| \otimes |i\rangle_{B'} \langle i|,$$

where $\{|i\rangle_{A'(B')}\}$ is the orthonormal basis of system $A'(B')$. Note that $\{|j_i\rangle_A |i\rangle_{A'}\}$ and $\{|j_i\rangle_B |i\rangle_{B'}\}$ are local eigenbases of $\rho_{AA'}$ and $\rho_{BB'}$, respectively. Let $U_{AA'} = \sum_{i, j_i} |j_i\rangle_{BA} \langle j_i| \otimes |i\rangle_{B'A'} \langle i|$ and $U_{BB'} = \sum_{i, j_i} |j_i\rangle_{AB} \langle j_i| \otimes |i\rangle_{A'B'} \langle i|$ and a little thought shows that these two unitary operators satisfy

$$U_{AA'} \otimes U_{BB'} (T_{swap} \rho_{AA'BB'}^\Delta T_{swap}^\dagger) U_{AA'}^\dagger \otimes U_{BB'}^\dagger = \rho_{AA'BB'}^\Delta,$$

where T_{swap} denotes the swap operator with respect to the local eigenbases of $\rho_{AA'}$ and $\rho_{BB'}$, i.e., $\{|j_i\rangle_A |i\rangle_{A'}\}$ and $\{|j_i\rangle_B |i\rangle_{B'}\}$. Therefore, $\rho_{AA'BB'}^\Delta$ is unitarily symmetric. Consequently, we calculate the quantum correlated coherence of $\rho_{AA'BB'}^\Delta$,

$$\begin{aligned} \mathcal{C}_{re}^{cc}(\rho_{AA'BB'}^\Delta) &= \mathcal{C}_{re}(\rho_{AA'BB'}^\Delta) = S(\rho_{AA'BB'}^{\Delta diag}) - S(\rho_{AA'BB'}^\Delta) \\ &= \sum_i p_i^* S\left(\sum_{j_i} \lambda_{j_i}^2 |j_i\rangle_A \langle j_i| \otimes |j_i\rangle_B \langle j_i|\right) \\ &= \sum_i p_i^* S\left(\sum_{j_i} \lambda_{j_i}^2 |j_i\rangle_A \langle j_i|\right) = E_f(\rho_{AB}), \end{aligned}$$

where the third equality is due to the property of von Neumann entropy as given by formula (6). The above equality together with the definition of EOC implies that

$$E_{re}^{cc}(\rho_{AB}) \leq \mathcal{C}_{re}^{cc}(\rho_{AA'BB'}^\Delta) = E_f(\rho_{AB}).$$

If ρ_{AB} is a pure state, its relative entropy of entanglement is equal to its entanglement of formation, and then equal to its EOC. Hence, the desired results of theorem 3 are obtained.

From theorem 3, we conclude that the EOC is not strictly less than the relative entropy of entanglement for a bipartite state, since for pure states they

are equal. Moreover, for a maximally correlated states [15,43], which has the form:

$$\rho_{AB} = \sum_{i,j} \rho_{ij} |i\rangle_A \langle j| \otimes |i\rangle_B \langle j|, \quad (10)$$

where $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ are orthonormal bases of subsystems A and B respectively and ρ_{ij} are the matrix elements, its EOC is also equal to its relative entropy of entanglement. We show this result in the following theorem.

Theorem 4. For a maximally correlated state ρ_{AB} as given by formula (10), its EOC is equal to its relative entropy of entanglement, i.e., $E_{re}^{cc}(\rho_{AB}) = E_{re}(\rho_{AB})$.

Proof: For the maximally correlated state ρ_{AB} , it has the form as given by formula (10). Let the local eigenbases of ρ_A and ρ_B , i.e., $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$, be the fixed bases of subsystems A and B respectively. According to the Ref. [8], we have $\mathcal{C}_{re}(\rho_A^*) = E_{re}(\rho_{AB})$, where $\rho_A^* = \sum_{i,j} \rho_{ij} |i\rangle_A \langle j|$ in subsystem A . Direct calculation yields $\mathcal{C}_{re}^{cc}(\rho_{AB}) = S(\rho_{AB}^{diag}) - S(\rho_{AB}) = S(\rho_A^{*diag}) - S(\rho_A^*) = \mathcal{C}_{re}(\rho_A^*)$. Obviously, $\rho_{AB} \otimes |00\rangle_{A'B'} \langle 00|$ is a unitarily symmetric extension of ρ_{AB} . With respect to the local eigenbases of $\rho_{AA'}$ and $\rho_{BB'}$ as the fixed bases of subsystems AA' and BB' , respectively, we have the equality $\mathcal{C}_{re}^{cc}(\rho_{AB} \otimes |00\rangle_{A'B'} \langle 00|) = \mathcal{C}_{re}^{cc}(\rho_{AB})$. With the definition of EOC, it holds that $E_{re}^{cc}(\rho_{AB}) \leq \mathcal{C}_{re}^{cc}(\rho_{AB} \otimes |00\rangle_{A'B'} \langle 00|) = \mathcal{C}_{re}^{cc}(\rho_{AB})$. Combining the aforementioned results and theorem 3, we arrive at the result $E_{re}^{cc}(\rho_{AB}) = E_{re}(\rho_{AB})$.

Using the proof of theorem 4, we confirm that for a maximally correlated state ρ_{AB} , its EOC is even equal to its quantum correlated coherence with respect to the local eigenbases of ρ_A and ρ_B , respectively. Moreover, with theorem 4, it is easy to find a state for which the EOC is strictly less than the entanglement of formation, for example, the maximally correlated Bell diagonal state in the two-qubit system, $\rho_{AB}^{mc} = \frac{3}{4}|\Phi^+\rangle\langle\Phi^+| + \frac{1}{4}|\Phi^-\rangle\langle\Phi^-|$, where $|\Phi^\pm\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)$ [20, 21]. However, we do not know whether the EOC is equal to the relative entropy of entanglement for any mixed state. In addition, for any bipartite state ρ_{AB} and τ_{CD} , EOC satisfies the following sub-additivity,

$$\max\{E_{re}^{cc}(\rho_{AB}), E_{re}^{cc}(\tau_{CD})\} \leq E_{re}^{cc}(\rho_{AB} \otimes \tau_{CD}) \leq E_{re}^{cc}(\rho_{AB}) + E_{re}^{cc}(\tau_{CD}).$$

An alternative measure of entanglement formulated by quantum correlated coherence (quantum discord) is defined as $\bar{E}_{re}^{cc}(\rho_{AB}) \equiv \min D(\rho_{AA'BB'})$, where the minimization is taken over all possible unitarily symmetric extensions $\rho_{AA'BB'}$ of ρ_{AB} , and $D(\rho_{AA'BB'}) = \min_{\{|i\rangle_{AA'}, |j\rangle_{BB'}\}} \mathcal{C}_{re}^{cc}(\rho_{AA'BB'})$, with the minimization over all generic bases $\{|i\rangle_{AA'}\}$ and $\{|j\rangle_{BB'}\}$ of subsystems AA' and BB' , respectively. Removing the property of unitary symmetry of extension $\rho_{AA'BB'}$ in the definition $\bar{E}_{re}^{cc}(\rho_{AB})$, we denote this new measure of entanglement as $\tilde{E}_{re}^{cc}(\rho_{AB})$. Remarkably, $\tilde{E}_{re}^{cc}(\rho_{AB})$ is equivalent to the entanglement measure which is the minimal discord over state extensions [35]. Moreover, $\bar{E}_{re}^{cc}(\rho_{AB})$ and $\tilde{E}_{re}^{cc}(\rho_{AB})$ have the properties: non-negative and vanished for separated states, invariant under local unitary operations, non-increasing under local operations, convex and upper bounded by entanglement of formation $E_f(\rho_{AB})$. However, the properties of $\bar{E}_{re}^{cc}(\rho_{AB})$ and $\tilde{E}_{re}^{cc}(\rho_{AB})$, which

are the invariance (non-increasing property) under classical communication and the relation to the relative entropy of entanglement, are not clear. In this sense, the EOC is more advantageous than $\bar{E}_{re}^{cc}(\rho_{AB})$ and $\hat{E}_{re}^{cc}(\rho_{AB})$.

In multipartite systems, there exists an entanglement measure like the definition of EOC. For a N -partite state $\rho_{C_1 C_2 \dots C_N}$, $\rho_{C_1 C'_1 C_2 C'_2 \dots C_N C'_N}$ is its unitarily symmetric extension and the local fixed bases of subsystems are the eigenbases of $\rho_{C_1 C'_1}$, $\rho_{C_2 C'_2}, \dots$, and $\rho_{C_N C'_N}$, respectively. Then the entanglement of $\rho_{C_1 C_2 \dots C_N}$ is defined as $E_{re}^{cc}(\rho_{C_1 C_2 \dots C_N}) \equiv \min \mathcal{C}_{re}^{cc}(\rho_{C_1 C'_1 C_2 C'_2 \dots C_N C'_N})$, where the minimization is over all possible unitarily symmetric extensions $\rho_{C_1 C'_1 C_2 C'_2 \dots C_N C'_N}$ of $\rho_{C_1 C_2 \dots C_N}$, $Tr_{C'_1 C'_2 \dots C'_N}(\rho_{C_1 C'_1 C_2 C'_2 \dots C_N C'_N}) = \rho_{C_1 C_2 \dots C_N}$. Note that the extension $\rho_{C_1 C'_1 C_2 C'_2 \dots C_N C'_N}$ is unitarily symmetric if it remains invariant up to local unitary operations on $C_i C'_i$ and $C_j C'_j$ under a system swap between $C_i C'_i$ and $C_j C'_j$ for any $i, j = 1, 2, \dots, N$. Referring to the proofs of EOC as an entanglement measure [13], we can show that $E_{re}^{cc}(\rho_{C_1 C_2 \dots C_N})$ has the properties: non-negative and vanished for separated states, invariant under local unitary operations, un-increased under LOCC operations, and convex. These results show that the entanglement in multipartite systems can also be formulated by quantum correlated coherence via state extensions.

5 Conclusions

In this paper, using entropy-based measures, we have discussed the concise relationships between quantum coherence and quantum correlations as defined by quantum discord as well as quantum entanglement. The results mean that quantum discord and entanglement can be formulated by quantum correlated coherence. In particular, we gave the condition for quantum correlated coherence (symmetric basis-dependent discord) to vanish, and this condition provides the explicit structure of states which satisfy the super-additive property of the relative entropy of coherence with equality. We further proved the lower and upper bounds of EOC and showed that the EOC is equal to the relative entropy of entanglement in a large number of scenarios including all pure states and maximally correlated states. For pure states, the LOCC monotonicity (monotonicity on average under LOCC operations [24, 44]) of EOC is easily obtained with theorem 3. However, we do not know whether the EOC of a general mixed state is LOCC monotone [24, 44], and we leave it open for future research. Finally, we also generalized our results to multipartite settings.

These results suggest that the quantum properties of correlations originate from the quantum properties of coherence and quantum correlations are better understood with the framework of coherence. We hope that this work is helpful for further understanding quantum correlations and developing quantum technologies.

Acknowledgements We thank Jia-Jun Ma for useful comments. This work is supported by the National Natural Science Foundation of China (Grants No. 61572081, No. 61672110, No. 61671082, No. 61601171).

Appendix : The explicit structure of states with vanished quantum correlated coherence

Here we prove that a state ρ_{AB} with vanished quantum correlated coherence has a decomposition as given in formula (5) in the main text.

For a given state ρ_{AB} with vanished quantum correlated coherence, its symmetric quantum discord is equal to zero, i.e., $D(\rho_{AB}) = 0$. Then, ρ_{AB} is a classical-classical state [40] with the form

$$\rho_{AB} = \sum_{\alpha,\beta} \lambda_{\alpha\beta} |\psi_\alpha\rangle\langle\psi_\alpha| \otimes |\phi_\beta\rangle\langle\phi_\beta|, \quad (B1)$$

where $\{|\psi_\alpha\rangle\}$ and $\{|\phi_\beta\rangle\}$ are orthonormal bases of subsystems A and B , respectively. From the main text, we see that

$$\mathcal{C}_{re}^{cc}(\rho_{AB}) = 0 \Leftrightarrow S(\rho_{AB} \| \rho_A \otimes \rho_B) = S(\Pi_A \otimes \Pi_B(\rho_{AB}) \| \Pi_A(\rho_A) \otimes \Pi_B(\rho_B)). \quad (B2)$$

where the von Neumann measurements Π_A and Π_B are with respect to the fixed bases $\{|i\rangle_A\}$ and $\{|j\rangle_B\}$ of subsystems A and B , respectively.

Recall that the quantum relative entropy is unchanged under a quantum operation \mathcal{E} , meaning that $S(\rho \| \sigma) = S(\mathcal{E}(\rho) \| \mathcal{E}(\sigma))$, if and only if there is a recovery operation \mathcal{R} satisfying $\mathcal{R} \circ \mathcal{E}(\rho) = \rho$, $\mathcal{R} \circ \mathcal{E}(\sigma) = \sigma$ [38, 39]. Moreover, there is an explicit formula for the recovery operation: $\mathcal{R}(X) = \sigma^{\frac{1}{2}} \mathcal{E}^\dagger(\mathcal{E}(\sigma)^{-\frac{1}{2}} X \mathcal{E}(\sigma)^{-\frac{1}{2}}) \sigma^{\frac{1}{2}}$. Here, \mathcal{E} is the local von Neumann measurements $\Pi_A \otimes \Pi_B = (\Pi_A \otimes \Pi_B)^\dagger$. Applied to formula (B2), the recovery condition says that

$$\mathcal{R}(\Pi_A \otimes \Pi_B(\rho_{AB})) = \sum_{i,j} \frac{\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle i | \psi_\alpha \rangle|^2 |\langle j | \phi_\beta \rangle|^2}{\left(\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle i | \psi_\alpha \rangle|^2 \right) \left(\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle j | \phi_\beta \rangle|^2 \right)} \left(\rho_A^{\frac{1}{2}} |i\rangle\langle i| \rho_A^{\frac{1}{2}} \right) \otimes \left(\rho_B^{\frac{1}{2}} |j\rangle\langle j| \rho_B^{\frac{1}{2}} \right). \quad (B3)$$

By letting formula (B3) be equal to formula (B1) and pre- and post-multiplying by $\rho_A^{-\frac{1}{2}} \otimes \rho_B^{-\frac{1}{2}}$, we obtain

$$\begin{aligned} & \sum_{i,j} \frac{\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle i | \psi_\alpha \rangle|^2 |\langle j | \phi_\beta \rangle|^2}{\left(\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle i | \psi_\alpha \rangle|^2 \right) \left(\sum_{\alpha,\beta} \lambda_{\alpha\beta} |\langle j | \phi_\beta \rangle|^2 \right)} |i\rangle\langle i| \otimes |j\rangle\langle j| \\ &= \sum_{\alpha,\beta} \frac{\lambda_{\alpha\beta}}{\left(\sum_{\xi} \lambda_{\alpha\xi} \right) \left(\sum_{\gamma} \lambda_{\gamma\beta} \right)} |\psi_\alpha\rangle\langle\psi_\alpha| \otimes |\phi_\beta\rangle\langle\phi_\beta|. \end{aligned} \quad (B4)$$

Remarkably, $\rho_A^{-\frac{1}{2}}$ and $\rho_B^{-\frac{1}{2}}$ are defined as

$$\rho_A^{-\frac{1}{2}} \equiv \sum_{\alpha: \sum_{\beta} \lambda_{\alpha\beta} \neq 0} \frac{1}{\sqrt{\sum_{\beta} \lambda_{\alpha\beta}}} |\psi_\alpha\rangle\langle\psi_\alpha|, \quad \rho_B^{-\frac{1}{2}} \equiv \sum_{\beta: \sum_{\alpha} \lambda_{\alpha\beta} \neq 0} \frac{1}{\sqrt{\sum_{\alpha} \lambda_{\alpha\beta}}} |\phi_\beta\rangle\langle\phi_\beta|,$$

where all $\sum_{\beta} \lambda_{\alpha\beta}$ and $|\psi_{\alpha}\rangle$ are the eigenvalues and eigenvectors of ρ_A , and all $\sum_{\alpha} \lambda_{\alpha\beta}$ and $|\phi_{\beta}\rangle$ are the same requirements of ρ_B . Thus in formula (B4) we exclude terms on either side which are not in the support of ρ_A and ρ_B .

Let i', j', α' and β' be the values of i, j, α and β in formulas (B3) and (B1). After taking the inner product $\langle i' | \langle j' | (\) | \psi_{\alpha'} \rangle | \phi_{\beta'} \rangle$ on either side of formula (B4), we have

$$\frac{\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i' | \psi_{\alpha} \rangle|^2 |\langle j' | \phi_{\beta} \rangle|^2}{\left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i' | \psi_{\alpha} \rangle|^2 \right) \left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle j' | \phi_{\beta} \rangle|^2 \right)} \langle i' | \psi_{\alpha'} \rangle \langle j' | \phi_{\beta'} \rangle = \frac{\lambda_{\alpha' \beta'}}{\left(\sum_{\beta} \lambda_{\alpha' \beta} \right) \left(\sum_{\alpha} \lambda_{\alpha \beta'} \right)} \langle i' | \psi_{\alpha'} \rangle \langle j' | \phi_{\beta'} \rangle. \quad (B5)$$

If $\langle i' | \psi_{\alpha'} \rangle \langle j' | \phi_{\beta'} \rangle \neq 0$, Eq. (B5) means that

$$\frac{\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i' | \psi_{\alpha} \rangle|^2 |\langle j' | \phi_{\beta} \rangle|^2}{\left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i' | \psi_{\alpha} \rangle|^2 \right) \left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle j' | \phi_{\beta} \rangle|^2 \right)} = \frac{\lambda_{\alpha' \beta'}}{\left(\sum_{\beta} \lambda_{\alpha' \beta} \right) \left(\sum_{\alpha} \lambda_{\alpha \beta'} \right)}. \quad (B6)$$

Due to Eq. (B6), we confirm that the left sides of (B6) are the same for all i' and j' satisfying $\langle i' | \psi_{\alpha'} \rangle \langle j' | \phi_{\beta'} \rangle \neq 0$ when we fix α' and β' . As the same reason, the right sides of (B6) are the same for all α' and β' satisfying $\langle i' | \psi_{\alpha'} \rangle \langle j' | \phi_{\beta'} \rangle \neq 0$ when we fix i' and j' .

Expanding formula (B3) continuously, we obtain that

$$\begin{aligned} \mathcal{R}(\Pi_A \otimes \Pi_B(\rho_{AB})) &= \sum_{i, j} \frac{\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i | \psi_{\alpha} \rangle|^2 |\langle j | \phi_{\beta} \rangle|^2}{\left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle i | \psi_{\alpha} \rangle|^2 \right) \left(\sum_{\alpha, \beta} \lambda_{\alpha\beta} |\langle j | \phi_{\beta} \rangle|^2 \right)} \\ &\quad \left(\sum_{\alpha} \sqrt{\sum_{\beta} \lambda_{\alpha\beta}} \langle \psi_{\alpha} | i \rangle | \psi_{\alpha} \rangle \right) \left(\sum_{\alpha} \sqrt{\sum_{\beta} \lambda_{\alpha\beta}} \langle i | \psi_{\alpha} \rangle \langle \psi_{\alpha} | \right) \\ &\quad \otimes \left(\sum_{\beta} \sqrt{\sum_{\alpha} \lambda_{\alpha\beta}} \langle \phi_{\beta} | j \rangle | \phi_{\beta} \rangle \right) \left(\sum_{\beta} \sqrt{\sum_{\alpha} \lambda_{\alpha\beta}} \langle j | \phi_{\beta} \rangle \langle \phi_{\beta} | \right). \end{aligned} \quad (B7)$$

By letting formula (B7) be equal to formula (B1), we firstly consider the case that $|\psi_{\alpha_1}\rangle$ ($|\phi_{\beta_1}\rangle$) and all the other $|\psi_{\alpha}\rangle$ ($|\phi_{\beta}\rangle$) have disjoint coherence support. Without loss of generality, let $\{|i_1\rangle, |i_2\rangle\}$ and $\{|j_1\rangle, |j_2\rangle\}$ be the coherence support of $|\psi_{\alpha_1}\rangle$ and $|\phi_{\beta_1}\rangle$ respectively. A little thought shows that the sum of only the four terms (i_1, j_1) , (i_1, j_2) , (i_2, j_1) and (i_2, j_2) in formula (B7) coincides with the term $\lambda_{\alpha_1 \beta_1} |\psi_{\alpha_1}\rangle \langle \psi_{\alpha_1} | \otimes |\phi_{\beta_1}\rangle \langle \phi_{\beta_1} |$ in formula (B1).

Secondly, we consider the case that the coherence support of $|\psi_{\alpha_1}\rangle$ has some intersection with that of other $|\psi_{\alpha}\rangle$, or the coherence support of $|\phi_{\beta_1}\rangle$ has some intersection with that of other $|\phi_{\beta}\rangle$. Without loss of generality, let $\{|i_1\rangle, |i_2\rangle\}$ be the coherence support of $|\psi_{\alpha_1}\rangle$ and $|\psi_{\alpha_2}\rangle$, and the set of $\{|i_1\rangle, |i_2\rangle\}$ has no intersection with the coherence support of other $|\psi_{\alpha}\rangle$ except $|\psi_{\alpha_1}\rangle$ and $|\psi_{\alpha_2}\rangle$.

Similarly, let $\{|j_1\rangle, |j_2\rangle\}$ be the coherence support of $|\phi_{\beta_1}\rangle$ and $|\phi_{\beta_2}\rangle$, and the set of $\{|j_1\rangle, |j_2\rangle\}$ has no intersection with the coherence support of other $|\phi_\beta\rangle$ except $|\phi_{\beta_1}\rangle$ and $|\phi_{\beta_2}\rangle$. The sum of only the four terms (i_1, j_1) , (i_1, j_2) , (i_2, j_1) and (i_2, j_2) in formula (B7) will be written as

$$\begin{aligned}
& \sum_{\substack{i=i_1, i_2; \\ j=j_1, j_2}} \frac{\left(\sum_{\substack{\alpha=\alpha_1, \alpha_2; \\ \beta=\beta_1, \beta_2}} \lambda_{\alpha\beta} |\langle i|\psi_\alpha\rangle|^2 |\langle j|\phi_\beta\rangle|^2 \right) \left(\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta} \right) \left(\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2} \right)}{\left(\sum_{\alpha=\alpha_1, \alpha_2; \beta} \lambda_{\alpha\beta} |\langle i|\psi_\alpha\rangle|^2 \right) \left(\sum_{\alpha; \beta=\beta_1, \beta_2} \lambda_{\alpha\beta} |\langle j|\phi_\beta\rangle|^2 \right)} \\
& \left(\frac{\sum_{\beta} \lambda_{\alpha_1\beta}}{\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta}} |\langle i|\psi_{\alpha_1}\rangle|^2 |\psi_{\alpha_1}\rangle\langle\psi_{\alpha_1}| + \frac{\sum_{\beta} \lambda_{\alpha_2\beta}}{\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta}} |\langle i|\psi_{\alpha_2}\rangle|^2 |\psi_{\alpha_2}\rangle\langle\psi_{\alpha_2}| \right. \\
& + \frac{\sqrt{\left(\sum_{\beta} \lambda_{\alpha_1\beta}\right)\left(\sum_{\beta} \lambda_{\alpha_2\beta}\right)}}{\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta}} \langle\psi_{\alpha_1}|i\rangle\langle i|\psi_{\alpha_2}\rangle |\psi_{\alpha_1}\rangle\langle\psi_{\alpha_2}| \\
& \left. + \frac{\sqrt{\left(\sum_{\beta} \lambda_{\alpha_1\beta}\right)\left(\sum_{\beta} \lambda_{\alpha_2\beta}\right)}}{\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta}} \langle\psi_{\alpha_2}|i\rangle\langle i|\psi_{\alpha_1}\rangle |\psi_{\alpha_2}\rangle\langle\psi_{\alpha_1}| \right) \\
& \otimes \left(\frac{\sum_{\alpha} \lambda_{\alpha\beta_1}}{\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2}} |\langle j|\phi_{\beta_1}\rangle|^2 |\phi_{\beta_1}\rangle\langle\phi_{\beta_1}| + \frac{\sum_{\alpha} \lambda_{\alpha\beta_2}}{\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2}} |\langle j|\phi_{\beta_2}\rangle|^2 |\phi_{\beta_2}\rangle\langle\phi_{\beta_2}| \right. \\
& + \frac{\sqrt{\left(\sum_{\alpha} \lambda_{\alpha\beta_1}\right)\left(\sum_{\alpha} \lambda_{\alpha\beta_2}\right)}}{\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2}} \langle\phi_{\beta_1}|j\rangle\langle j|\phi_{\beta_2}\rangle |\phi_{\beta_1}\rangle\langle\phi_{\beta_2}| \\
& \left. + \frac{\sqrt{\left(\sum_{\alpha} \lambda_{\alpha\beta_1}\right)\left(\sum_{\alpha} \lambda_{\alpha\beta_2}\right)}}{\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2}} \langle\phi_{\beta_2}|j\rangle\langle j|\phi_{\beta_1}\rangle |\phi_{\beta_2}\rangle\langle\phi_{\beta_1}| \right)
\end{aligned} \tag{B8}$$

Using formula (B6), we know that the formulas

$$\frac{\sum_{\substack{\alpha=\alpha_1, \alpha_2; \\ \beta=\beta_1, \beta_2}} \lambda_{\alpha\beta} |\langle i|\psi_\alpha\rangle|^2 |\langle j|\phi_\beta\rangle|^2}{\left(\sum_{\alpha=\alpha_1, \alpha_2; \beta} \lambda_{\alpha\beta} |\langle i|\psi_\alpha\rangle|^2 \right) \left(\sum_{\alpha; \beta=\beta_1, \beta_2} \lambda_{\alpha\beta} |\langle j|\phi_\beta\rangle|^2 \right)},$$

are the same for any $i = i_1, i_2$ and $j = j_1, j_2$. Then using the orthonormality of states in sets $\{|\psi_{\alpha_1}\rangle, |\psi_{\alpha_2}\rangle\}$ and $\{|\phi_{\beta_1}\rangle, |\phi_{\beta_2}\rangle\}$, and removing the cross terms that contain $|\psi_{\alpha_1}\rangle\langle\psi_{\alpha_2}|, |\psi_{\alpha_2}\rangle\langle\psi_{\alpha_1}|, |\phi_{\beta_1}\rangle\langle\phi_{\beta_2}|$ or $|\phi_{\beta_2}\rangle\langle\phi_{\beta_1}|$ in formula (B8), we obtain the simplified form of formula (B8):

$$p_{\alpha_1\alpha_2\beta_1\beta_2} (\mu_1 |\psi_{\alpha_1}\rangle\langle\psi_{\alpha_1}| + \mu_2 |\psi_{\alpha_2}\rangle\langle\psi_{\alpha_2}|) \otimes (\eta_1 |\phi_{\beta_1}\rangle\langle\phi_{\beta_1}| + \eta_2 |\phi_{\beta_2}\rangle\langle\phi_{\beta_2}|), \tag{B9}$$

where $\mu_{1(2)}$, $\eta_{1(2)}$ and $p_{\alpha_1\alpha_2\beta_1\beta_2}$ are non-negative, and satisfy

$$\mu_{1(2)} = \frac{\sum_{\beta} \lambda_{\alpha_{1(2)}\beta}}{\sum_{\beta} \lambda_{\alpha_1\beta} + \sum_{\beta} \lambda_{\alpha_2\beta}}, \quad \eta_{1(2)} = \frac{\sum_{\alpha} \lambda_{\alpha\beta_{1(2)}}}{\sum_{\alpha} \lambda_{\alpha\beta_1} + \sum_{\alpha} \lambda_{\alpha\beta_2}},$$

$$\mu_1 + \mu_2 = 1, \quad \eta_1 + \eta_2 = 1, \quad p_{\alpha_1\alpha_2\beta_1\beta_2} = \sum_{\substack{\alpha=\alpha_1,\alpha_2; \\ \beta=\beta_1,\beta_2}} \lambda_{\alpha\beta}.$$

What's more, formula (B9) coincides with the sum of partial terms in formula (B1): $\sum_{\substack{\alpha=\alpha_1,\alpha_2; \\ \beta=\beta_1,\beta_2}} \lambda_{\alpha\beta} |\psi_{\alpha}\rangle\langle\psi_{\alpha}| \otimes |\phi_{\beta}\rangle\langle\phi_{\beta}|$. Finally, other cases that there exist some intersection of coherence support of $|\psi_{\alpha}\rangle$ or $|\phi_{\beta}\rangle$ can be discussed similarly, and the results like formula (B9) will be obtained. Hence, the equality of formula (B1) and (B7) means that ρ_{AB} has a decomposition as given in formula (5) in the main text.

References

1. S. D. Bartlett, T. Rudolph, and R.W. Spekkens, Reference frames, superselection rules, and quantum information, *Rev. Mod. Phys.* **79**, 555 (2007).
2. J. Aberg, Quantifying Superposition, arXiv:0612146V1.
3. F. Levi and F. Mintert, A quantitative theory of coherent delocalization, *New J. Phys.* **16**, 033007 (2014).
4. T. Baumgratz, M. Cramer, and M. B. Plenio, Quantifying Coherence, *Phys. Rev. Lett.* **113**, 140401 (2014).
5. A. Winter and D. Yang, Operational resource theory of coherence, *Phys. Rev. Lett.* **116**, 120404 (2016).
6. E. Chitambar and G. Gour, Critical Examination of Incoherent Operations and a Physically Consistent Resource Theory of Quantum Coherence, *Phys. Rev. Lett.* **117**, 030401 (2016).
7. A. Streltsov, G. Adesso, and M. B. Plenio, Colloquium: quantum coherence as a resource, arXiv:1609.02439.
8. A. Streltsov, U. Singh, H. S. Dhar, M. N. Bera, and G. Adesso, Measuring quantum coherence with entanglement, *Phys. Rev. Lett.* **115**, 020403 (2015).
9. J. Ma, B. Yadin, D. Girolami, V. Vedral, and M. Gu, Converting coherence to quantum correlations, *Phys. Rev. Lett.* **116**, 160407 (2016).
10. E. Chitambar and M.-H. Hsieh, Relating the resource theories of entanglement and quantum coherence, *Phys. Rev. Lett.* **117**, 020402 (2016).
11. Z. Xi, Y. Li, and H. Fan, Quantum coherence and correlations in quantum system, *Sci. Rep.* **5**, 10922 (2015).
12. Y. Yao, X. Xiao, L. Ge, and C. P. Sun, Quantum coherence in multipartite systems, *Phys. Rev. A* **92**, 022112 (2015).
13. K. C. Tan, H. Kwon, C.-Y. Park, and H. Jeong, Unified view of quantum correlations and quantum coherence, *Phys. Rev. A* **94**, 022329 (2016).
14. E. Chitambar, A. Streltsov, S. Rana, M.N. Bera, G. Adesso, and M. Lewenstein, Assisted distillation of quantum coherence, *Phys. Rev. Lett.* **116**, 070402 (2016).
15. A. Streltsov, S. Rana, M. Nath Bera, and M. Lewenstein, Towards resource theory of coherence in distributed scenarios, arXiv:1509.07456v3.
16. T. R. Bromley, M. Cianciaruso, and G. Adesso, Frozen quantum coherence, *Phys. Rev. Lett.* **114**, 210401 (2015).
17. X. D. Yu, D. J. Zhang, C. Liu, and D. Tong, Measure-independent freezing of quantum coherence, *Phys. Rev. A* **93**, 060303(R) (2016).

18. M. Lostaglio, D. Jennings, and T. Rudolph, Description of quantum coherence in thermodynamic processes requires constraints beyond free energy, *Nat. Commun.* **6**, 6383 (2015).
19. M. Lostaglio, K. Korzekwa, D. Jennings, and T. Rudolph, Quantum Coherence, Time-Translation Symmetry, and Thermodynamics, *Phys. Rev. X* **5**, 021001 (2015).
20. C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Mixed-state entanglement and quantum error correction, *Phys. Rev. A* **54**, 3824 (1996).
21. V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Quantifying entanglement, *Phys. Rev. Lett.* **78**, 2275 (1997).
22. V. Vedral and M. B. Plenio, Entanglement measures and purification procedures, *Phys. Rev. A* **57**, 1619 (1998).
23. G. Vidal, Entanglement monotones, *J. Mod. Opt.* **47**, 355 (2000).
24. M. B. Plenio and S. Virmani, An introduction to entanglement measures, *Quantum Inf. Comput.* **7**,1 (2007).
25. H. Ollivier and W. H. Zurek, Quantum discord: a measure of the quantumness of correlations, *Phys. Rev. Lett.* **88**, 017901 (2001).
26. S. L. Luo, Using measurement-induced disturbance to characterize correlations as classical or quantum, *Phys. Rev. A* **77**, 022301 (2008).
27. C. C. Rulli and M. S. Sarandy, Global quantum discord in multipartite systems, *Phys. Rev. A* **84**, 042109 (2011).
28. K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson, Unified view of quantum and classical correlations, *Phys. Rev. Lett.* **104**, 080501 (2010).
29. K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, The classical-quantum boundary for correlations: discord and related measures, *Rev. Mod. Phys.* **84**, 1655 (2012).
30. A. K. Ekert, Quantum cryptography based on Bells theorem, *Phys. Rev. Lett.* **67**, 661 (1991).
31. C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels, *Phys. Rev. Lett.* **70**, 1895 (1993).
32. R. Jozsa and N. Linden, On the role of entanglement in quantum computational speed-up, *Proc. R. Soc. Lon A.* **459**, 2011 (2003).
33. B. P. Lanyon, M. Barbieri, M. P. Almeida, and A. G. White, Experimental quantum computing without entanglement, *Phys. Rev. Lett.* **101**(20), 200501 (2008).
34. V. Giovannetti, S. Lloyd, and L. Maccone, Advances in quantum metrology, *Nature Photon.* **5**, 222 (2011).
35. S. L. Luo, Entanglement as minimal discord over state extensions, *Phys. Rev. A* **94**, 032129 (2016).
36. M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum information*, Cambridge University Press, Cambridge (2000).
37. B. Yadin, J. Ma, D. Girolami, M. Gu, and V. Vedral, Quantum Processes Which Do Not Use Coherence, *Phys. Rev. X* **6**, 041028 (2016).
38. D. Petz, Monotonicity of quantum relative entropy revisited, *Rev. Math. Phys.* **15**(01): 79-91 (2003).
39. P. Hayden, R. Jozsa, D. Petz, and A. Winter, Structure of states which satisfy strong subadditivity of quantum entropy with equality, *Commun. Math. Phys.* **246**, 359 (2004).
40. M. Piani, P. Horodecki, and R. Horodecki, No-local-broadcasting theorem for multipartite quantum correlations, *Phys. Rev. Lett.* **100**, 090502 (2008).
41. Z. F. Bai and S. P. Du, Maximally coherent states, *Quant. Inf. Coumput.* **15**: 1355-1364 (2015).
42. N. Li and S. Luo, Classical states versus separable states, *Phys. Rev. A* **78**, 024303 (2008).
43. E. Rains, A Semidefinite Program for Distillable Entanglement, *IEEE Trans. Inf. Theory* **47**, 2921 (2001).
44. M. Horodecki, Simplifying monotonicity conditions for entanglement measures, *Open Sys. Inf. Dyn.* **12**(3): 231-237 (2005).