# \*K-means and Cluster Models for Cancer Signatures

Zura Kakushadze $^{\$^{\dagger}1}$  and Willie  $\mathrm{Yu}^{\sharp2}$ 

§ Quantigic<sup>®</sup> Solutions LLC

1127 High Ridge Road #135, Stamford, CT 06905 <sup>3</sup>
<sup>†</sup> Free University of Tbilisi, Business School & School of Physics 240, David Agmashenebeli Alley, Tbilisi, 0159, Georgia
<sup>‡</sup> Centre for Computational Biology, Duke-NUS Medical School 8 College Road, Singapore 169857

(January 30, 2017)

#### Abstract

We present \*K-means clustering algorithm and source code by expanding statistical clustering methods applied in https://ssrn.com/abstract=2802753 to quantitative finance. \*K-means is statistically deterministic without specifying initial centers, etc. We apply \*K-means to extracting cancer signatures from genome data without using nonnegative matrix factorization (NMF). \*K-means' computational cost is a fraction of NMF's. Using 1,389 published samples for 14 cancer types, we find that 3 cancers (liver cancer, lung cancer and renal cell carcinoma) stand out and do not have cluster-like structures. Two clusters have especially high within-cluster correlations with 11 other cancers indicating common underlying structures. Our approach opens a novel avenue for studying such structures. \*K-means is universal and can be applied in other fields. We discuss some potential applications in quantitative finance.

<sup>&</sup>lt;sup>1</sup> Zura Kakushadze, Ph.D., is the President of Quantigic<sup>®</sup> Solutions LLC, and a Full Professor at Free University of Tbilisi. Email: zura@quantigic.com

<sup>&</sup>lt;sup>2</sup> Willie Yu, Ph.D., is a Research Fellow at Duke-NUS Medical School. Email: willie.yu@duke-nus.edu.sg

<sup>&</sup>lt;sup>3</sup> DISCLAIMER: This address is used by the corresponding author for no purpose other than to indicate his professional affiliation as is customary in publications. In particular, the contents of this paper are not intended as an investment, legal, tax or any other such advice, and in no way represent views of Quantigic<sup>®</sup> Solutions LLC, the website <u>www.quantigic.com</u> or any of their other affiliates.

# **1** Introduction and Summary

Every time we can learn something new about cancer, the motivation goes without saying. Cancer is different. Unlike other diseases, it is not caused by "mechanical" breakdowns, biochemical imbalances, etc. Instead, cancer occurs at the DNA level via somatic alterations in the genome structure. A common type of somatic mutations found in cancer is due to single nucleotide variations (SNVs) or alterations to single bases in the genome, which accumulate through the lifespan of the cancer via imperfect DNA replication during cell division or spontaneous cytosine deamination [Goodman and Fygenson, 1998], [Lindahl, 1993], or due to exposures to chemical insults or ultraviolet radiation [Loeb and Harris, 2008], [Ananthaswamy and Pierceall, 1990], etc. These mutational processes leave a footprint in the cancer genome characterized by distinctive alteration patterns or mutational signatures.

If we can identify all underlying signatures, this could greatly facilitate progress in understanding the origins of cancer and its development. Therapeutically, if there are common underlying structures across different cancer types, then a therapeutic for one cancer type might be applicable to other cancers, which would be a great news.<sup>4</sup> However, it all boils down to the question of usefulness, i.e., is there a small enough number of cancer signatures underlying all (100+) known cancer types, or is this number too large to be meaningful or useful? Indeed, there are only 96 SNVs,<sup>5</sup> so we cannot have more than 96 signatures.<sup>6</sup> Even if the number of true underlying signatures is, say, of order 50, it is unclear whether they would be useful, especially within practical applications. On the other hand, if there are only a dozen or so underlying signatures, then we could hope for an order of magnitude simplification.

To identify mutational signatures, one analyzes SNV patterns in a cohort of DNA sequenced whole cancer genomes. The data is organized into a matrix  $G_{is}$ , where the rows correspond to the N = 96 mutation categories, the columns correspond to d samples, and each element is a nonnegative occurrence count of a given mutation category in a given sample. Currently, the commonly accepted method for extracting cancer signatures from  $G_{is}$  [Alexandrov *et al*, 2013a] is via nonnegative matrix factorization (NMF) [Paatero and Tapper, 1994], [Lee and Seung, 1999]. Under NMF the matrix G is approximated via  $G \approx W H$ , where  $W_{iA}$  is an  $N \times K$  matrix,  $H_{As}$  is a  $K \times d$  matrix, and both W and H are nonnegative. The appeal of NMF is its biologic interpretation whereby the K columns of the matrix W are

<sup>&</sup>lt;sup>4</sup> Another practical application is prevention by pairing the signatures extracted from cancer samples with those caused by known carcinogens (e.g., tobacco, aflatoxin, UV radiation, etc).

<sup>&</sup>lt;sup>5</sup> In brief, DNA is a double helix of two strands, and each strand is a string of letters A, C, G, T corresponding to adenine, cytosine, guanine and thymine, respectively. In the double helix, A in one strand always binds with T in the other, and G always binds with C. This is known as base complementarity. Thus, there are six possible base mutations C > A, C > G, C > T, T > A, T > C, T > G, whereas the other six base mutations are equivalent to these by base complementarity. Each of these 6 possible base mutations is flanked by 4 possible bases on each side thereby producing  $4 \times 6 \times 4 = 96$  distinct mutation categories.

<sup>&</sup>lt;sup>6</sup> Nonlinearities could undermine this argument. However, again, it all boils down to usefulness.

interpreted as the weights with which the K cancer signatures contribute into the N = 96 mutation categories, and the columns of the matrix H are interpreted as the exposures to the K signatures in each sample. The price to pay for this is that NMF, which is an iterative procedure, is computationally costly and depending on the number of samples d it can take days or even weeks to run it. Furthermore, it does not automatically fix the number of signatures K, which must be either guessed or obtained via trial and error, thereby further adding to the computational cost.<sup>7</sup>

Some of the aforesaid issues were recently addressed in [Kakushadze and Yu, 2016b], to wit: i) by aggregating samples by cancer types, we can greatly improve stability and reduce the number of signatures;<sup>8</sup> ii) by identifying and factoring out the somatic mutational noise, or the "overall" mode (this is the "de-noising" procedure of [Kakushadze and Yu, 2016b]), we can further greatly improve stability and, as a bonus, reduce computational cost; and iii) the number of signatures can be fixed borrowing the methods from statistical risk models [Kakushadze and Yu, 2017b] in quantitative finance, by computing the effective rank (or eRank) [Roy and Vetterli, 2007] for the correlation matrix  $\Psi_{ij}$  calculated across cancer types or samples (see below). All this yields substantial improvements [Kakushadze and Yu, 2016b].

In this paper we push this program to yet another level. The basic idea here is quite simple (but, as it turns out, nontrivial to implement – see below). We wish to apply clustering techniques to the problem of extracting cancer signatures. In fact, we argue in Section 2 that NMF is, to a degree, "clustering in disguise". This is for two main reasons. The prosaic reason is that NMF, being a nondeterministic algorithm, requires averaging over many local optima it produces. However, each run generally produces a weights matrix  $W_{iA}$  with columns (i.e., signatures) not aligned with those in other runs. Aligning or matching the signatures across different runs (before averaging over them) is typically achieved via nondeterministic clustering such as k-means. So, not only is clustering utilized at some layer, the result, even after averaging, generally is both noisy<sup>9</sup> and nondeterministic! I.e., if this computationally costly procedure (which includes averaging) is run again and again on the same data, generally it will yield different looking cancer signatures every time!

The second, not-so-prosaic reason is that, while NMF generically does not produce exactly null weights, it does produce low weights, such that they are within error bars. For all practical purposes we might as well set such weights to zero. NMF requires nonnegative weights. However, we could as reasonably require that the weights should be, say, outside error bars (e.g., above one standard deviation –

<sup>&</sup>lt;sup>7</sup> Other issues include: i) out-of-sample instability, i.e., the signatures obtained from nonoverlapping sets of samples can be dramatically different; ii) in-sample instability, i.e., the signatures can have a strong dependence on the initial iteration choice; and iii) samples with low counts or sparsely populated samples (i.e., those with many zeros – such samples are ubiquitous, e.g., in exome data) are usually deemed not too useful as they contribute to the in-sample instability.

<sup>&</sup>lt;sup>8</sup> As a result, now we have the so-aggregated matrix  $G_{is}$ , where  $s = 1, \ldots, d$ , and d = n is the number of cancer types, not of samples. This matrix is much less noisy than the sample data.

 $<sup>^9</sup>$  By "noise" we mean the statistical errors in the weighs obtained by averaging. Typically, such error bars are not reported in the literature on cancer signatures. Usually they are large.

this would render the algorithm highly recursive and potentially unstable or computationally too costly) or above some minimum threshold (which would still further complicated as-is complicated NMF), or else the non-compliant weights are set to zero. As we increase this minimum threshold, the matrix  $W_{iA}$  will start to have more and more zeros. It may not exactly have a binary cluster-like structure, but it may at least have some substructures that are cluster-like. It then begs the question: are there cluster-like (sub)structures present in  $W_{iA}$  or, generally, in cancer signatures?

To answer this question, we can apply clustering methods directly to the matrix  $G_{is}$ , or, more, precisely, to its de-noised version  $G'_{is}$  (see below) [Kakushadze and Yu, 2016b]. The naïve, brute-force approach where one would simply cluster  $G_{is}$  or  $G'_{is}$  does not work for a variety of reasons, some being more nontrivial or subtle than others. Thus, e.g., as discussed in [Kakushadze and Yu, 2016b], the counts  $G_{is}$  have skewed, long-tailed distributions and one should work with log-counts, or, more precisely, their de-noised versions. This applies to clustering as well. Further, following a discussion in [Kakushadze and Yu, 2016c] in the context of quantitative trading, it would be suboptimal to cluster de-noised log-counts. Instead, it pays to cluster their normalized variants (see Section 2 hereof). However, taking care of such subtleties does not alleviate one big problem: nondeterminism!<sup>10</sup> If we run a vanilla nondeterministic algorithm such as k-means on the data however massaged with whatever bells and whistles, we will get random-looking disparate results every time we run k-means with no stability in sight. We need to address nondeterminism!

Our solution to the problem is what we term \*K-means. The idea behind \*Kmeans, which essentially achieves determinism *statistically*, is simple. Suppose we have an  $N \times d$  matrix  $X_{is}$ , i.e., we have N d-vectors  $\mathbf{X}_i$ . If we run k-means with the input number of clusters K but initially unspecified centers, every run will generally produce a new local optimum. \*K-means reduces and in fact essentially eliminates this indeterminism via two levels. At level 1 it takes clusterings obtained via Mindependent runs or samplings. Each sampling produces a binary  $N \times K$  matrix  $\Omega_{iA}$ , whose element equals 1 if  $\mathbf{X}_i$  belongs to the cluster labeled by A, and 0 otherwise. The aggregation algorithm and the source code therefor are given in Kakushadze and Yu, 2016c. This aggregation – for the same reasons as in NMF (see above) - involves aligning clusters across the M runs, which is achieved via k-means, and so the result is nondeterministic. However, by aggregating a large number M of samplings, the degree of nondeterminism is greatly reduced. The "catch" is that sometimes this aggregation yields a clustering with K' < K clusters, but this does not pose an issue. Thus, at level 2, we take a large number P of such aggregations (each based on M samplings). The occurrence counts of aggregated clusterings are not uniform but typically have a (sharply) peaked distribution around a few (or manageable) number of aggregated clusterings. So this way we can pinpoint the "ultimate" clustering, which is simply the aggregated clustering with the highest occurrence count. This is the gist of \*K-means and it works well for genome data.

<sup>&</sup>lt;sup>10</sup> Deterministic (e.g., agglomerative hierarchical) algorithms have their own issues (see below).

So, we apply \*K-mean to the same genome data as in [Kakushadze and Yu, 2016b] consisting of 1,389 (published) samples across 14 cancer types (see below). Our target number of clusters is 7, which was obtained in [Kakushadze and Yu, 2016b] using the eRank based algorithm (see above). We aggregated 1,000 samplings into clusterings, and we constructed 150,000 such aggregated clusterings (i.e., we ran 150 million k-means instances). We indeed found the "ultimate" clustering with 7 clusters. Once the clustering is fixed, it turns out that within-cluster weights can be computed via linear regressions (with some bells and whistles) and the weights are automatically positive. That is, we do not need NMF at all! Once we have clusters and weights, we can study reconstruction accuracy and within-cluster correlations between the underlying data and the fitted data that the cluster model produces.

We find that clustering works well for 10 out the 14 cancer types we study. The cancer types for which clustering does not appear to work all that well are Liver Cancer, Lung Cancer, and Renal Cell Carcinoma. Also, above 80% within-cluster correlations arise for 5 out of 7 clusters. Furthermore, remarkably, one cluster has high within-cluster correlations for 9 cancer types, and another cluster for 6 cancer types. These appear to be the leading clusters. Together they have high within-cluster correlations in 11 out of 14 cancer types. So what does all this mean?

Additional insight is provided by looking at the within-cluster correlations between signatures Sig1 through Sig7 extracted in [Kakushadze and Yu, 2016b] and our clusters. High within-cluster correlations arise for Sig1, Sig2, Sig4 and Sig7, which are precisely the signatures with "peaks" (or "spikes" – "tall mountain landscapes"), whereas Sig3, Sig5 and Sig6 do not have such "peaks" ("flat" or "rolling hills landscapes"); see Figures 14 through 20 of [Kakushadze and Yu, 2016b]. The latter 3 signatures simply do not have cluster-like structures. Looking at Figure 21 in [Kakushadze and Yu, 2016b], it becomes evident why clustering does not work well for Liver Cancer – it has a whopping 96% contribution from Sig5! Similarly, Renal Cell Carcinoma has a 70% contribution from Sig6. Lung Cancer is dominated by Sig3, hence no cluster-like structure. So, Liver Cancer, Lung Cancer and Renal Cell Carcinoma have little in common with other cancers (and each other)! However, 11 other cancers, to wit, B Cell Lymphoma, Bone Cancer, Brain Lower Grade Glioma, Breast Cancer, Chronic Lymphocytic Leukemia, Esophageal Cancer, Gastric Cancer, Medulloblastoma, Ovarian Cancer, Pancreatic Cancer and Prostate Cancer, have 5 (with 2 leading) cluster structures substantially embedded in them.

In Section 2 we i) discuss why applying clustering algorithms to extracting cancer signatures makes sense, ii) argue that NMF, to a degree, is "clustering in disguise", and iii) give the machinery for building cluster models via \*K-means, including various details such as what to cluster, how to fix the number of clusters, etc. In Section 3 we discuss i) cancer genome data we use, ii) our application of \*K-means to it, and iii) the interpretation of our empirical results. Section 4 contains some concluding remarks, including a discussion of potential applications of \*K-means in quantitative finance, where we outline some concrete problems where \*K-means can be useful. Appendix A contains R source code for \*K-means and cluster models.

# 2 Cluster Models

The chief objective of this paper is to introduce a novel approach to identifying cancer signatures using clustering methods. In fact, as we discuss below in detail, our approach is more than just clustering. Indeed, it is evident from the get-go that blindly using nondeterministic clustering algorithms,<sup>11</sup> which typically produce (unmanageably) large numbers of local optima, would introduce great variability into the resultant cancer signatures.<sup>12</sup> On the other hand, deterministic algorithms such as agglomerative hierarchical clustering<sup>13</sup> typically are (substantially) slower and require essentially "guessing" the initial clustering,<sup>14</sup> which in practical applications<sup>15</sup> can often turn out to be suboptimal. So, both to motivate and explain our new approach employing clustering methods, we first – so to speak – "break down" the NMF approach and argue that it is in fact a clustering method in disguise!

### 2.1 "Breaking Down" NMF

The current "lore" – the commonly accepted method for extracting K cancer signatures from the occurrence counts matrix  $G_{is}$  (see above) [Alexandrov *et al*, 2013a] - is via nonnegative matrix factorization (NMF) [Paatero and Tapper, 1994], [Lee and Seung, 1999]. Under NMF the matrix G is approximated via  $G \approx W H$ , where  $W_{iA}$  is an  $N \times K$  matrix of weights,  $H_{As}$  is a  $K \times d$  matrix of exposures, and both W and H are nonnegative. However, not only is the number of signatures K not fixed via NMF (and must be either guessed or obtained via trial and error), NMF too is a nondeterministic algorithm and typically produces a large number of local optima. So, in practice one has no choice but to execute a large number  $N_S$  of NMF runs – which we refer to as samplings – and then somehow extract cancer signatures from these samplings. Absent a guess for what K should be, one executes  $N_S$  samplings for a range of values of K (say,  $K_{\min} \leq K \leq K_{\max}$ , where  $K_{\min}$  and  $K_{\max}$  are basically guessed based on some reasonable intuitive considerations), for each K extracts cancer signatures (see below), and then picks K and the corresponding signatures with the best overall fit into the underlying matrix G. For a given K, different samplings generally produce different weights matrices W. So, to extract a single matrix W for each value of K one *averages* over the samplings. However, before averaging, one must match the K cancer signatures across different samplings - indeed, in a given sampling X the columns in the matrix  $W_{iA}$  are not necessarily aligned with

<sup>&</sup>lt;sup>11</sup> Such as k-means [Steinhaus, 1957], [Lloyd, 1957], [Forgy, 1965], [MacQueen, 1967], [Hartigan, 1975], [Hartigan and Wong, 1979], [Lloyd, 1982].

 $<sup>^{12}\,</sup>$  As we discuss below, in this regard NMF is not dissimilar.

<sup>&</sup>lt;sup>13</sup> E.g., SLINK [Sibson, 1973], etc. (see, e.g., [Murtagh and Contreras, 2011], [Kakushadze and Yu, 2016c], and references therein).

<sup>&</sup>lt;sup>14</sup> E.g., splitting the data into 2 initial clusters.

<sup>&</sup>lt;sup>15</sup> Such as quantitative trading, where out-of-sample performance can be objectively measured. There empirical evidence suggests that such deterministic algorithms underperform so long as nondeterministic ones are used thoughtfully [Kakushadze and Yu, 2016c].

the columns in the matrix  $W_{iA}$  in a different sampling Y. To align the columns in the matrices W across the  $N_S$  samplings, once often uses a clustering algorithm such as k-means. However, since k-means is nondeterministic, such alignment of the Wcolumns is not guaranteed to – and in fact does not – produce a unique answer. Here one can try to run multiple samplings of k-means for this alignment and aggregate them, albeit such aggregation itself would require another level of alignment (with its own nondeterministic clustering such as k-means).<sup>16</sup> And one can do this adinfinitum. In practice, one must break the chain at some level of alignment, either ad hoc (essentially by heuristically observing sufficient stability and "convergence") or via using a deterministic algorithm (see fn. 16). Either way, invariably all this introduces (overtly or covertly) systematic and statistical errors into the resultant cancer signatures and often it is unclear if they are meaningful without invoking some kind empirical biologic "experience" or "intuition" (often based on already well-known effects of, e.g., exposure to various well-understood carcinogens such as tobacco, ultraviolet radiation, aflatoxin, etc.). At the end of the day it all boils down to how useful – or *predictive* – the resultant method of extracting cancer signatures is, including signature stability. With NMF, the answer is not at all evident...

### 2.2 Clustering in Disguise?

So, in practice, under the hood, NMF already uses clustering methods. However, it goes deeper than that. While NMF generically does not produce vanishing weights for a given signature, some weights are (much) smaller than others. E.g., often one has several "peaks" with high concentration of weights, with the rest of the mutation categories having relatively low weights. In fact, many weights can even be within the (statistical plus systematic) error bars.<sup>17</sup> Such weights can for all practical purposes be set to zero. In fact, we can take this further and ask whether proliferation of low weights adds any explanatory power. One way to address this is to run NMF with an additional constraint that the weights (obtained via averaging - see above) should be higher than either i) some multiple of the corresponding error bars<sup>18</sup> or ii) some preset fixed minimum weight. This certainly sounds reasonable, so why is this not done in practice? A prosaic answer appears to be that this would complicate the already nontrivial NMF algorithm even further, require additional coding and computation resources, etc. However, arguendo, let us assume that we require, say, that the weights be higher than a preset fixed minimum weight  $w_{min}$  or else the weights are set to zero. As we increase  $w_{min}$ , the so-modified NMF would produce more and more zeros. This does not mean that the resulting matrix  $W_{iA}$ 

<sup>&</sup>lt;sup>16</sup> We should point out that at some level of alignment one may employ a deterministic (e.g., agglomerative hierarchical – see above) clustering algorithm to terminate the malicious circle, which can be a reasonable approach assuming there is enough stability in the data. However, this too adds a(n often hard to quantify and therefore hidden) systematic error to the resultant signatures.

<sup>&</sup>lt;sup>17</sup> And such error bars are rarely displayed in the prevalent literature...

<sup>&</sup>lt;sup>18</sup> This would require a highly recursive algorithm.

would have a binary cluster structure, i.e., that  $W_{iA} = w_i \, \delta_{G(i),A}$ , where  $\delta_{AB}$  is a Kronecker delta and  $G: \{1, \ldots, N\} \mapsto \{1, \ldots, K\}$  is a map from N = 96 mutation categories to K clusters. Put another way, this does not mean that in the resulting matrix  $W_{iA}$  for a given i (i.e., mutation category) we would have a nonzero element for one and only one value of A (i.e., signature). However, as we gradually increase  $w_{min}$ , generally the matrix  $W_{iA}$  is expected to look more and more like having a binary cluster structure, albeit with some "overlapping" signatures (i.e., such that in a given pair of signatures there are nonzero weights for one or more mutations). We can achieve a binary structure via a number of ways. Thus, a rudimentary algorithm would be to take the matrix  $W_{iA}$  (equally successfully before or after achieving some zeros in it via nonzero  $w_{min}$ ) and for a given value of i set all weights  $W_{iA}$  to zero except in the signature A for which  $W_{iA} = \max(W_{iA}|A=1,\ldots,K)$ . Note that this might result in some empty signatures (clusters), i.e., signatures with  $W_{iA} = 0$  for all values of i. This can be dealt with by i) ether simply dropping such signatures altogether and having fewer K' < K signatures (binary clusters) at the end, or ii) augmenting the algorithm to avoid empty clusters, which can be done in a number of ways we will not delve into here. The bottom line is that NMF essentially can be made into a clustering algorithm by reasonably modifying it, including via getting rid of ubiquitous and not-too-informative low weights. However, the downside would be an even more contrived algorithm, so this is not what we are suggesting here. Instead, we are observing that clustering is already intertwined in NMF and the question is whether we can simplify things by employing clustering methods directly.

## 2.3 Making Clustering Work

Happily, the answer is yes. Not only can we have much simpler and apparently more stable clustering algorithms, but they are also computationally much less costly than NMF. As mentioned above, the biggest issue with using popular nondeterministic clustering algorithms such as k-means<sup>19</sup> is that they produce a large number of local optima. For definiteness in the remainder of this paper we will focus on k-means, albeit the methods described herein are general and can be applied to other such algorithms. Fortunately, this very issue has already been addressed in [Kakushadze and Yu, 2016c] in the context of constructing statistical industry classifications (i.e., clustering models for stocks) for quantitative trading, so here we simply borrow therefrom and further expand and adapt that approach to cancer signatures.

### 2.3.1 K-means

A popular clustering algorithm is k-means [Steinhaus, 1957], [Lloyd, 1957], [Forgy, 1965], [MacQueen, 1967], [Hartigan, 1975], [Hartigan and Wong, 1979], [Lloyd, 1982]. The basic idea behind k-means is to partition N observations into K clusters such that each observation belongs to the cluster with the nearest mean. Each of the N

<sup>&</sup>lt;sup>19</sup> Which are preferred over deterministic ones for the reasons discussed above.

observations is actually a *d*-vector, so we have an  $N \times d$  matrix  $X_{is}$ , i = 1, ..., N, s = 1, ..., d. Let  $C_a$  be the K clusters,  $C_a = \{i | i \in C_a\}, a = 1, ..., K$ . Then k-means attempts to minimize<sup>20</sup>

$$g = \sum_{a=1}^{K} \sum_{i \in C_a} \sum_{s=1}^{d} \left( X_{is} - Y_{as} \right)^2 \tag{1}$$

where

$$Y_{as} = \frac{1}{n_a} \sum_{i \in C_a} X_{is} \tag{2}$$

are the cluster centers (i.e., cross-sectional means),<sup>21</sup> and  $n_a = |C_a|$  is the number of elements in the cluster  $C_a$ . In (1) the measure of "closeness" is chosen to be the Euclidean distance between points in  $\mathbf{R}^d$ , albeit other measures are possible.

One "drawback" of k-means is that it is not a deterministic algorithm. Generically, there are copious local minima of g in (1) and the algorithm only guarantees that it will converge to a local minimum, not the global one. Being an iterative algorithm, unless the initial centers are preset, k-means starts with a random set of the centers  $Y_{as}$  at the initial iteration and converges to a different local minimum in each run. There is no magic bullet here: in practical applications, typically, trying to "guess" the initial centers is not any easier than "guessing" where, e.g., the global minimum is. So, what is one to do? One possibility is to simply live with the fact that every run produces a different answer. In fact, this is acceptable in many applications. However, in the context of extracting cancer signatures this would result in an exercise in futility. We need a way to eliminate or greatly reduce indeterminism.

### 2.3.2 Aggregating Clusterings

The idea is simple. What if we *aggregate* different clusterings from multiple runs – which we refer to as samplings – into one? The question is how. Suppose we have M runs  $(M \gg 1)$ . Each run produces a clustering with K clusters. Let  $\Omega_{ia}^r = \delta_{G^r(i),a}$ ,  $i = 1, \ldots, N, a = 1, \ldots, K$  (here  $G^r : \{1, \ldots, N\} \mapsto \{1, \ldots, K\}$  is the map between – in our case – the mutation categories and the clusters),<sup>22</sup> be the binary matrix from each run labeled by  $r = 1, \ldots, M$ , which is a convenient way (for our purposes here) of encoding the information about the corresponding clustering; thus, each row of  $\Omega_{ia}^r$  contains only one element equal 1 (others are zero), and  $N_a^r = \sum_{i=1}^N \Omega_{ia}^r$  (i.e., column sums) is nothing but the number of mutations belonging to the cluster labeled by a (note that  $\sum_{a=1}^K N_a^r = N$ ). Here we are assuming that somehow we know how to properly order (i.e., align) the K clusters from each run. This is a nontrivial assumption, which we will come back to momentarily. However, assuming, for a second, that we know how to do this, we can aggregate the binary matrices

<sup>&</sup>lt;sup>20</sup> Below we will discuss what  $X_{is}$  should be for cancer signatures.

 $<sup>^{21}</sup>$  Throughout this paper "cross-sectional" refers to "over the index i ".

<sup>&</sup>lt;sup>22</sup> Note that here the superscript r in  $\Omega_{ia}^r$ ,  $G^r(i)$  and  $N_a^r$  (see below) is an index, not a power.

 $\Omega_{ia}^{r}$  into a single matrix  $\widetilde{\Omega}_{ia} = \sum_{r=1}^{M} \Omega_{ia}^{r}$ . Now, this matrix does not look like a binary clustering matrix. Instead, it is a matrix of occurrence counts, i.e., it counts how many times a given mutation was assigned to a given cluster in the process of M samplings. What we need to construct is a map G such that one and only one mutation belongs to each of the K clusters. The simplest criterion is to map a given mutation to the cluster in which  $\widetilde{\Omega}_{ia}$  is maximal, i.e., where said mutation occurs most frequently. A caveat is that there may be more than one such clusters. A simple criterion to resolve such an ambiguity is to assign said mutation to the cluster with most cumulative occurrences (i.e., we assign said mutation to the cluster with the largest  $\widetilde{N}_{a} = \sum_{i=1}^{N} \widetilde{\Omega}_{ia}$ ). Further, in the unlikely event that there is still an ambiguity, we can try to do more complicated things, or we can simply assign such a mutation to the cluster with the lowest value of the index a – typically, there is so much noise in the system that dwelling on such minutiae simply does not pay off.

However, we still need to tie up a loose end, to wit, our assumption that the clusters from different runs were somehow all aligned. In practice each run produces K clusters, but i) they are not the same clusters and there is no foolproof way of mapping them, especially when we have a large number of runs; and ii) even if the clusters were the same or similar, they would not be ordered, i.e., the clusters from one run generally would be in a different order than the clusters from another run.

So, we need a way to "match" clusters from different samplings. Again, there is no magic bullet here either. We can do a lot of complicated and contrived things with not much to show for it at the end. A simple pragmatic solution is to use k-means to align the clusters from different runs. Each run labeled by  $r = 1, \ldots, M$ , among other things, produces a set of cluster centers  $Y_{as}^r$ . We can "bootstrap" them by row into a  $(KM) \times d$  matrix  $\widetilde{Y}_{\tilde{a}s} = Y_{as}^r$ , where  $\tilde{a} = a + (r-1)K$  takes values  $\tilde{a} = 1, \ldots, (KM)$ . We can now cluster  $\widetilde{Y}_{\tilde{a}s}$  into K clusters via k-means. This will map each value of  $\widetilde{a}$  to  $\{1, \ldots, K\}$  thereby mapping the K clusters from each of the M runs to  $\{1,\ldots,K\}$ . So, this way we can align all clusters. The "catch" is that there is no guarantee that each of the K clusters from each of the M runs will be uniquely mapped to one value in  $\{1, \ldots, K\}$ , i.e., we may have some empty clusters at the end of the day. However, this is fine, we can simply drop such empty clusters and aggregate (via the above procedure) the smaller number of K' < K clusters. I.e., at the end we will end up with a clustering with K' clusters, which might be fewer than the target number of clusters K. This is not necessarily a bad thing. The dropped clusters might have been redundant in the first place. Another evident "catch" is that even the number of resulting clusters K' is not deterministic. If we run this algorithm multiple times, we will get varying values of K'. Malicious circle?

#### 2.3.3 Fixing the "Ultimate" Clustering

Not really! There is one other trick up our sleeves we can use to fix the "ultimate" clustering thereby rendering our approach essentially deterministic. The idea above is to aggregate a large enough number M of samplings. Each aggregation produces

a clustering with some  $K' \leq K$  clusters, and this K' varies from aggregation to aggregation. However, what if we take a large number P of aggregations (each based on M samplings)? Typically there will be a relatively large number of different clusterings we get this way. However, assuming some degree of stability in the data, this number is much smaller than the number of a priori different local minima we would obtain by running the vanilla k-means algorithm. What is even better, the occurrence counts of aggregated clusterings are not uniform but typically have a (sharply) peaked distribution around a few (or manageable) number of aggregated clusterings. In fact, as we will see below, in our empirical genome data we are able to pinpoint the "ultimate" clustering! So, to recap, what we have done here is this. There are myriad clusterings we can get via vanilla k-means with little to no guidance as to which one to pick.<sup>23</sup> We have reduced this proliferation by aggregating a large number of such clusterings into our aggregated clusterings. We then further zoom onto a few or even a unique clustering we consider to be the likely "ultimate" clustering by examining the occurrence counts of such aggregated clusterings, which turns out to have a (sharply) peaked distribution. Since vanilla k-means is a relatively fast-converging algorithm, each aggregation is not computationally taxing and running a large number of aggregations is nowhere as time consuming as running a similar number (or even a fraction thereof) of NMF computations (see below).

### 2.4 What to Cluster?

So, now that we know how to make clustering work, we need to decide what to cluster, i.e., what to take as our matrix  $X_{is}$  in (1). The naïve choice  $X_{is} = G_{is}$  is suboptimal for multiple reasons (as discussed in [Kakushadze and Yu, 2016b]).

First, the elements of the matrix  $G_{is}$  are populated by nonnegative occurrence counts. Nonnegative quantities with large numbers of samples tend to have skewed distributions with long tails at higher values. I.e., such distributions are not normal but (in many cases) roughly log-normal. One simple way to deal with this is to identify  $X_{is}$  with a (natural) logarithm of  $G_{is}$  (instead of  $G_{is}$  itself). A minor hiccup here is that some elements of  $G_{is}$  can be 0. We can do a lot of complicated and even convoluted things to deal with this issue. Here, as in [Kakushadze and Yu, 2016b], we will follow a pragmatic approach and do something simple instead – there is so much noise in the data that doing convoluted things simply does not pay off. So, as the first cut, we can take

$$X_{is} = \ln\left(1 + G_{is}\right) \tag{3}$$

This takes care of the  $G_{is} = 0$  cases; for  $G_{is} \gg 1$  we have  $R_{is} \approx \ln(G_{is})$ , as desired.

Second, the detailed empirical analysis of [Kakushadze and Yu, 2016b] uncovered what is termed therein the "overall" mode<sup>24</sup> unequivocally present in the occurrence count data. This "overall" mode is interpreted as somatic mutational *noise* unrelated

 $<sup>^{23}</sup>$  This is because things are pretty much random and the only "distribution" at hand is flat.

<sup>&</sup>lt;sup>24</sup> In finance the analog of this is the so-called "market" mode (see, e.g., [Bouchaud and Potters, 2011] and references therein) corresponding to the overall movement of the broad market, which

to (and in fact obscuring) the true underlying cancer signatures and must therefore be factored out somehow. Here is a simple way to understand the "overall" mode. Let the correlation matrix  $\Psi_{ij} = \text{Cor}(X_{is}, X_{js})$ , where  $\text{Cor}(\cdot, \cdot)$  is serial correlation.<sup>25</sup> I.e.,  $\Psi_{ij} = C_{ij}/\sigma_i \sigma_j$ , where  $\sigma_i^2 = C_{ii}$  are variances, and the serial covariance matrix<sup>26</sup>

$$C_{ij} = \text{Cov}(X_{is}, X_{js}) = \frac{1}{d-1} \sum_{s=1}^{d} Z_{is} \ Z_{js}$$
(4)

where  $Z_{is} = X_{is} - \overline{X}_i$  are serially demeaned, while the means  $\overline{X}_i = \frac{1}{d} \sum_{s=1}^d X_{is}$ . The average pair-wise correlation  $\rho = \frac{1}{N(N-1)} \sum_{i,j=1; i \neq j}^N \Psi_{ij}$  between different mutation categories is nonzero and is in fact high for most cancer types we study. This is the aforementioned somatic mutational noise that must be factored out. If we aggregate samples by cancer types (see below) and compute the correlation matrix  $\Psi_{ij}$  for the so-aggregated data (across the n = 14 cancer types we study – see below),<sup>27</sup> the average correlation  $\rho$  is over whopping 96%. Another way of thinking about this is that the occurrence counts in different samples (or cancer types, if we aggregate samples by cancer types) are not normalized uniformly across all samples (cancer types). Therefore, running NMF, a clustering or any other signature-extraction algorithm on the vanilla matrix  $G_{is}$  (or its "log"  $X_{is}$  defined in (3)) would amount to mixing apples with oranges thereby obscuring the true underlying cancer signatures.

Following [Kakushadze and Yu, 2016b], factoring out the "overall" mode (or "de-noising" the matrix  $G_{is}$ ) therefore most simply amount to cross-sectional (i.e., across the 96 mutation categories) demeaning of the matrix  $X_{is}$ . I.e., instead of  $X_{is}$  we use  $X'_{is}$ , which is obtained from  $X_{is}$  by demeaning its columns:<sup>28</sup>

$$X'_{is} = X_{is} - \overline{X}_s = X_{is} - \frac{1}{N} \sum_{j=1}^{N} X_{js}$$
(5)

We should note that using  $X'_{is}$  instead of  $X_{is}$  in (1) does not affect clustering. Indeed, g in (1) is invariant under the transformations of the form  $X_{is} \to X_{is} + \Delta_s$ , where  $\Delta_s$  is an arbitrary *d*-vector, as thereunder we also have  $Y_{as} \to Y_{as} + \Delta_s$ , so  $X_{is} - Y_{as}$  is unchanged. In fact, this is good: this means that de-noising does not introduce any additional errors into clustering itself. However, the actual *weights* in the matrix  $W_{iA}$  are affected by de-noising. We discuss the algorithm for fixing  $W_{iA}$  below. However, we need one more ingredient before we get to determining the weights, and with this additional ingredient de-noising does affect clustering.

affects all stocks (to varying degrees) – cash inflow (outflow) into (from) the market tends to push stock prices higher (lower). This is the market risk factor, and to mitigate it one can, e.g., hold a dollar-neutral portfolio of stocks (i.e., the same dollar holdings for long and short positions).

 $<sup>^{25}</sup>$  Throughout this paper "serial" refers to "over the index s".

<sup>&</sup>lt;sup>26</sup> The overall normalization of  $C_{ij}$ , i.e., d-1 (unbiased estimate) vs. d (maximum likelihood estimate) in the denominator in the definition of  $C_{ij}$  in (4), is immaterial for our purposes here.

<sup>&</sup>lt;sup>27</sup> So, in this case d = n = 14 in (4).

<sup>&</sup>lt;sup>28</sup> For the reasons discussed above, we should demean  $X_{is}$ , not  $G_{is}$ .

#### 2.4.1 Normalizing Log-counts

As was discussed in [Kakushadze and Yu, 2016c], clustering  $X_{is}$  (or equivalently  $X'_{is}$ ) would be suboptimal.<sup>29</sup> The issue is this. Let  $\sigma'_i$  be serial standard deviations, i.e.,  $(\sigma'_i)^2 = \text{Cov}(X'_{is}, X'_{is})$ , where, as above,  $\text{Cov}(\cdot, \cdot)$  is serial covariance. Here we assume that samples are aggregated by cancer types, so  $s = 1, \ldots, d$  with d = n = 14. Now,  $\sigma'_i$  are not cross-sectionally uniform and vary substantially across mutation categories. The density of  $\sigma'_i$  is depicted in Figure 1 and is skewed (tailed). The summary of  $\sigma'_i$  reads:<sup>30</sup> Min = 0.2196, 1st Qu. = 0.3409, Median = 0.4596, Mean = 0.4984, 3rd Qu. = 0.6060, Max = 1.0010, SD = 0.1917, MAD = 0.1859, Skewness = 0.8498. If we simply cluster  $X'_{is}$ , this variability in  $\sigma'_i$  will not be accounted for.

A simple solution is to cluster normalized demeaned log-counts  $\tilde{X}'_{is} = X'_{is}/\sigma'_i$ instead of  $X'_{is}$ . This way we factor out the nonuniform (and skewed) standard deviation out of the log-counts. Note that now de-noising does make a difference in clustering. Indeed, if we use  $\tilde{X}_{is} = X_{is}/\sigma_i$  (recall that  $\sigma_i^2 = \text{Cov}(X_{is}, X_{is})$ ) instead of  $\tilde{X}'_{is} = X'_{is}/\sigma'_i$  in (1) and (2), the quantity g (and also clusterings) will be different.

### 2.5 Fixing Cluster Number

Now that we know what to cluster (to wit,  $\widetilde{X}'_{is}$ ) and how to get to the "unique" clustering, we need to figure out how to fix the (target) number of clusters K, which is one of the inputs in our algorithm above.<sup>31</sup> In [Kakushadze and Yu, 2016b] it was argued that in the context of cancer signatures their number can be fixed by building a statistical factor model [Kakushadze and Yu, 2017b], i.e., the number of signatures is simply the number of statistical factors.<sup>32</sup> So, by the same token, here we identify the (target) number of clusters in our clustering algorithm with the number of statistical factors fixed via the method of [Kakushadze and Yu, 2017b].

#### 2.5.1 Effective Rank

So, following [Kakushadze and Yu, 2017b] and [Kakushadze and Yu, 2016b], we set<sup>33</sup>

$$K = \text{Round}(\text{eRank}(\Psi)) \tag{6}$$

<sup>&</sup>lt;sup>29</sup> More precisely, the discussion of [Kakushadze and Yu, 2016c] is in the financial context, to wit, quantitative trading, which has its own nuances (see below). However, some of that discussion is quite general and can be adapted to a wide variety of applications.

 $<sup>^{30}\,</sup>$  Qu. = Quartile, SD = Standard Deviation, MAD = Mean Absolute Deviation.

<sup>&</sup>lt;sup>31</sup> A variety of methods for fixing the number of clusters have been discussed in other contexts, e.g., [Rousseeuw, 1987], [Pelleg and Moore, 2000], [Steinbach *et al*, 2000], [Goutte *et al*, 2001], [Sugar and James, 2003], [Hamerly and Elkan, 2004], [Lleití *et al*, 2004], [De Amorim and Hennig, 2015].

<sup>&</sup>lt;sup>32</sup> In the financial context, these are known as statistical risk models [Kakushadze and Yu, 2017b]. For a discussion and literature on multifactor risk models, see, e.g., [Grinold and Kahn, 2000], [Kakushadze and Yu, 2016a] and references therein. For prior works on fixing the number of statistical risk factors, see, e.g., [Connor and Korajczyk, 1993] and [Bai and Ng, 2002].

<sup>&</sup>lt;sup>33</sup> Here Round(·) can be replaced by floor(·) =  $\lfloor \cdot \rfloor$ .

Here eRank(Z) is the effective rank [Roy and Vetterli, 2007] of a symmetric semipositive-definite (which suffices for our purposes here) matrix Z. It is defined as

$$eRank(Z) = exp(H) \tag{7}$$

$$H = -\sum_{a=1}^{L} p_a \ln(p_a) \tag{8}$$

$$p_a = \frac{\lambda^{(a)}}{\sum_{b=1}^L \lambda^{(b)}} \tag{9}$$

where  $\lambda^{(a)}$  are the *L* positive eigenvalues of *Z*, and *H* has the meaning of the (Shannon a.k.a. spectral) entropy [Campbell, 1960], [Yang et al, 2005]. Let us emphasize that in (6) the matrix  $\Psi_{ij}$  is computed based on the demeaned log-counts<sup>34</sup>  $X'_{is}$ .

The meaning of  $\operatorname{Rank}(\Psi_{ij})$  is that it is a measure of the effective dimensionality of the matrix  $\Psi_{ij}$ , which is not necessarily the same as the number L of its positive eigenvalues, but often is lower. This is due to the fact that many d-vectors  $X'_{is}$  can be serially highly correlated (which manifests itself by a large gap in the eigenvalues) thereby further reducing the effective dimensionality of the correlation matrix.

#### How to Compute Weights? $\mathbf{2.6}$

The one remaining thing to accomplish is to figure out how to compute the weights  $W_{iA}$ . Happily, in the context of clustering we have significant simplifications compared with NMF and computing the weights becomes remarkably simple once we fix the clustering, i.e., the matrix  $\Omega_{iA} = \delta_{G(i),A}$  (or, equivalently, the map  $G: \{i\} \mapsto$  $\{A\}, i = 1, \dots, N, A = 1, \dots, K$ , where for the notational convenience we use K to denote the number of clusters in the "ultimate" clustering – see above). Just as in NMF, we wish to approximate the matrix  $G_{is}$  via a product of the weights matrix  $W_{iA}$  and the exposure matrix  $H_{As}$ , both of which must be nonnegative. More precisely, since we must remove the "overall" mode, i.e., de-noise the matrix  $G_{is}$ , following [Kakushadze and Yu, 2016b], instead of  $G_{is}$  we will approximate the re-exponentiated demeaned log-counts matrix  $X'_{is}$ :

$$G'_{is} = \exp(X'_{is}) \tag{10}$$

We can include an overall normalization by taking  $G'_{is} = \exp(\operatorname{Mean}(X_{is}) + X'_{is})$ , or  $G'_{is} = \exp(\operatorname{Median}(X_{is}) + X'_{is}), \text{ or } G'_{is} = \exp(\operatorname{Median}(\overline{X}_s) + X'_{is}) \text{ (recall that } \overline{X}_s)$ is the vector of column means of  $X_{is}$  – see Eq. (5)), etc., to make it look more like the original matrix  $G_{is}$ ; however, this does not affect the extracted signatures.<sup>35</sup> Also, technically speaking, after re-exponentiating we should "subtract" the extra 1 we added in the definition (3) (assuming we include one of the aforesaid overall normalizations). However, the inherent noise in the data makes this a moot point.

<sup>&</sup>lt;sup>34</sup> Note that using normalized demeaned log-counts  $\widetilde{X}'_{is}$  gives the same  $\Psi_{ij}$ . <sup>35</sup> This is because each column of W, being weights, is normalized to add up to 1.

So, we wish to approximate  $G'_{is}$  via a product W H. However, with clustering we have  $W_{iA} = w_i \, \delta_{G(i),A}$ , i.e., we have a block (cluster) structure where for a given value of A all  $W_{iA}$  are zero except for  $i \in J(A) = \{j | G(j) = A\}$ , i.e., for the mutation categories labeled by i that belong to the cluster labeled by A. Therefore, our matrix factorization of  $G_{is}$  into a product W H now simplifies into a set of Kindependent factorizations as follows:

$$G'_{is} \approx w_i H_{As}, \quad i \in J(A), \quad A = 1 \dots, K$$
 (11)

So, there is no need to run NMF anymore! Indeed, if we can somehow fix  $H_{As}$  for a given cluster, then within this cluster we can determine the corresponding weights  $w_i$   $(i \in J(A))$  via a *serial* linear regression:

$$G'_{is} = \varepsilon_{is} + w_i \ H_{As}, \quad i \in J(A), \quad A = 1 \dots, K$$
(12)

where  $\varepsilon_{is}$  are the regression residuals. I.e., for each  $A \in \{1, \ldots, K\}$ , we regress the  $d \times n_A$  matrix<sup>36</sup>  $[(G')^T]_{si}$   $(i \in J(A), n_A = |J(A)|)$  over the *d*-vector  $H_{As}$   $(s = 1, \ldots, d)$ , and the regression coefficients are nothing but the  $n_A$ -vector  $w_i$   $(i \in J(A))$ , while the residuals are the  $d \times n_A$  matrix  $[(\varepsilon)^T]_{si}$ . Note that this regression is run without the intercept. Now, this all makes sense as (for each  $i \in J(A)$ ) the regression minimizes the quadratic error term  $\sum_{s=1}^d \varepsilon_{is}^2$ . Furthermore, if  $H_{As}$  are nonnegative, then the weights  $w_i$  are automatically nonnegative as they are given by:

$$w_i = \frac{\sum_{s=1}^d G'_{is} H_{G(i),s}}{\sum_{s=1}^d H^2_{G(i),s}}$$
(13)

Now, we wish these weights to be normalized:

$$\sum_{i \in J(A)} w_i = 1 \tag{14}$$

This can always be achieved by rescaling  $H_{As}$ . Alternatively, we can pick  $H_{As}$  without worrying about the normalization, compute  $w_i$  via (13), rescale them so that they satisfy (14), and simultaneously accordingly rescale  $H_{As}$ . Mission accomplished!

#### 2.6.1 Fixing Exposures

Well, almost... We still need to figure out how to fix the exposures  $H_{As}$ . The simplest way to do this is to note that we can use the matrix  $\Omega_{iA} = \delta_{G(i),A}$  to swap the index *i* in  $G'_{is}$  by the index *A*, i.e., we can take

$$H_{As} = \eta_A \sum_{i=1}^{N} \Omega_{iA} \ G'_{is} = \tilde{\eta}_A \ \frac{1}{n_A} \sum_{i \in J(A)} G'_{is}$$
(15)

<sup>&</sup>lt;sup>36</sup> The superscript T denotes matrix transposition.

That is, up to the normalization constants  $\tilde{\eta}_A$  (which are fixed via (14)) we simply take cross-sectional means of  $G'_{is}$  in each cluster. (Recall that  $n_A = J(A)$ .) The sodefined  $H_{As}$  are automatically positive as all  $G'_{is}$  are positive. Therefore,  $w_i$  defined via (13) are also all positive. This is a good news – vanishing  $w_i$  would amount to an incomplete weights matrix  $W_{iA}$  (i.e., some mutations would belong to no cluster.)

So, why does (15) make sense? Looking at (12), we can observe that, if the residuals  $\varepsilon_{is}$  cross-sectionally, within each cluster labeled by A, are random, then we expect that  $\sum_{i \in J(A)} \varepsilon_{is} \approx 0$ . If we had an exact equality here, then we would have (15) with  $\eta_A = 1$  (i.e.,  $\tilde{\eta}_A = n_A$ ) assuming the normalization (14). In practice, the residuals  $\varepsilon_{is}$  are not exactly "random". First, the number  $n_A$  of mutation categories in each cluster is not large. Second, as mentioned above, there is variability in serial standard deviations across mutation types. This leads us to consider variations.

#### 2.6.2 A Variation

Above we argued that it makes sense to cluster normalized demeaned log-counts  $\widetilde{X}'_{is} = X'_{is}/\sigma'_i$  due to the cross-sectional variability (and skewness) in the serial standard deviations  $\sigma'_i$ . We may worry about similar effects in  $G'_{is}$  when computing  $H_{As}$  and  $w_i$  as we did above. This can be mitigated by using normalized quantities  $\widetilde{G}'_{is} = G'_{is}/\omega_i$ , where  $\omega_i^2 = \text{Cov}(G'_{is}, G'_{is})$  are serial variances. That is, we can define<sup>37</sup>

$$H_{As} = \tilde{\eta}_A \ \frac{1}{\nu_A} \sum_{i \in J(A)} \tilde{G}'_{is} = \tilde{\eta}_A \ \frac{1}{\nu_A} \sum_{i \in J(A)} \frac{1}{\omega_i} \ G'_{is}$$
(16)

$$w_{i} = \omega_{i} \; \frac{\sum_{s=1}^{d} \widetilde{G}'_{is} \; H_{G(i),s}}{\sum_{s=1}^{d} H^{2}_{G(i),s}} = \frac{\sum_{s=1}^{d} G'_{is} \; H_{G(i),s}}{\sum_{s=1}^{d} H^{2}_{G(i),s}}$$
(17)

where  $\nu_A = \sum_{i \in J(A)} 1/\omega_i$ . So,  $1/\omega_i$  are the weights in the averages over the clusters.

#### 2.6.3 Another Variation

Here one may wonder, considering the skewed roughly log-normal distribution of  $G_{is}$  and henceforth  $G'_{is}$ , would it make sense to relate the exposures to within-cluster cross-sectional averages of demeaned log-counts  $X'_{is}$  as opposed to those of  $G'_{is}$ ? This is easily achieved. Thus, we can define (this ensures positivity of  $H_{As}$ ):

$$\ln(H_{As}) = \ln(\tilde{\eta}_A) + \frac{1}{n_A} \sum_{i \in J(A)} X'_{is}$$
(18)

Exponentiating we get

$$H_{As} = \tilde{\eta}_A \left[ \prod_{i \in J(A)} G'_{is} \right]^{1/n_A}$$
(19)

<sup>&</sup>lt;sup>37</sup> I.e., here we assume that  $\varepsilon_{is}/\omega_i$  are approximately random in (12).

I.e., instead of an arithmetic average as in (15), here we have a geometric average.

As above, here too we can introduce nontrivial weights. Note that the form of (17) is the same as (13), it is only  $H_{As}$  that is affected by the weights. So, we can introduce the weights in the geometric means as follows:

$$\ln(H_{As}) = \ln(\widetilde{\eta}_A) + \frac{1}{\mu_A} \sum_{i \in J(A)} \widetilde{X}'_{is} = \ln(\widetilde{\eta}_A) + \frac{1}{\mu_A} \sum_{i \in J(A)} \frac{1}{\sigma'_i} X'_{is}$$
(20)

where  $\mu_A = \sum_{i \in J(A)} 1/\sigma'_i$ . Recall that  $(\sigma'_i)^2 = \text{Cov}(X'_{is}, X'_{is})$ . Thus, we have:

$$H_{As} = \widetilde{\eta}_A \prod_{i \in J(A)} (G'_{is})^{1/\mu_A \sigma'_i}$$
(21)

So, the weights are the exponents  $1/\mu_A \sigma'_i$ . Other variations are also possible.

### 2.7 Implementation

We are now ready to discuss an actual implementation of the above algorithm, much of the R code for which is already provided in [Kakushadze and Yu, 2016b] and [Kakushadze and Yu, 2016c]. The R source code is given in Appendix A hereof.

# 3 Empirical Results

### 3.1 Data Summary

In our empirical analysis below we use the same genome data (from published samples only) as in [Kakushadze and Yu, 2016b]. This data is summarized in Table 1 (borrowed from [Kakushadze and Yu, 2016b]), which gives total counts, number of samples and the data sources, which are as follows: A1 = [Alexandrov *et al*, 2013b], A2 = [Love *et al*, 2012], B1 = [Tirode *et al*, 2014], C1 = [Zhang *et al*, 2013], D1 = [Nik-Zainal *et al*, 2012], E1 = [Puente *et al*, 2011], E2 = [Puente *et al*, 2015], F1 = [Cheng *et al*, 2016], G1 = [Wang *et al*, 2014], H1 = [Sung *et al*, 2012], H2 = [Fujimoto *et al*, 2016], I1 = [Imielinksi *et al*, 2012], J1 = [Jones *et al*, 2012], K1 = [Patch *et al*, 2015], L1 = [Waddell *et al*, 2015], M1 = [Gundem *et al*, 2015], N1 = [Scelo *et al*, 2014]. Sample IDs with the corresponding publication sources are given in Appendix A of [Kakushadze and Yu, 2016b]. In our analysis below we aggregate samples by the 14 cancer types. The resulting data is in Tables 2 and 3.

#### 3.1.1 Structure of Data

The underlying data consists of a matrix – call it  $G_{is}$  – whose elements are occurrence counts of mutation types labeled by i = 1, ..., N = 96 in samples labeled by s = 1, ..., d. More precisely, we can work with one matrix  $G_{is}$  which combines data from different cancer types; or, alternatively, we may choose to work with individual matrices  $[G(\alpha)]_{is}$ , where:  $\alpha = 1, \ldots, n$  labels *n* different cancer types; as before,  $i = 1, \ldots, N = 96$ ; and  $s = 1, \ldots, d(\alpha)$ . Here  $d(\alpha)$  is the number of samples for the cancer type labeled by  $\alpha$ . The combined matrix  $G_{is}$  is obtained simply by appending (i.e., bootstrapping) the matrices  $[G(\alpha)]_{is}$  together column-wise. In the case of the data we use here (see above), this "big matrix" turns out to have 1389 columns.

Generally, individual matrices  $[G(\alpha)]_{is}$  and, thereby, the "big matrix", contain a lot of noise. For some cancer types we can have a relatively small number of samples. We can also have "sparsely populated" data, i.e., with many zeros for some mutation categories. As mentioned above, different samples are not necessarily uniformly normalized. Etc. The bottom line is that the data is noisy. Furthermore, intuitively it is clear that the larger the matrix we work with, statistically the more "signatures" (or clusters) we should expect to get with any reasonable algorithm. However, as mentioned above, a large number of signatures would be essentially useless and defy the whole purpose of extracting them in the first place – we have 96 mutation categories, so it is clear that the number of signatures cannot be more than 96! If we end up with, say, 50+ signatures, what new or useful does this tell us about the underlying cancers? The answer is likely nothing other than that most cancers have not much in common with each other, which would be a disappointing result from the perspective of the applications. To mitigate the aforementioned issues, at least to a certain extent, following [Kakushadze and Yu, 2016b], we can aggregate samples by cancer types. This way we get an  $N \times n$  matrix, which we also refer to as  $G_{is}$ , where the index  $s = 1, \ldots, d$  now takes d = n values corresponding to the cancer types. In the data we use n = 14, the aggregated matrix  $G_{is}$  is much less noisy than the "big matrix", and we are ready to apply the above machinery to it.

### 3.2 Genome Data Results

The 96 × 14 matrix  $G_{is}$  given in Tables 2 and 3 is what we pass into the function bio.cl.sigs() in Appendix A as the input matrix x. We use: iter.max = 100 (this is the maximum number of iterations used in the built-in R function kmeans() – we note that there was not a single instance in our 150 million runs of kmeans() where more iterations were required);<sup>38</sup> num.try = 1000 (this is the number of individual k-means samplings we aggregate every time); and num.runs = 150000 (which is the number of aggregated clusterings we use to determine the "ultimate" – that is, the most frequently occurring – clustering). So, we ran k-means 150 million times. More precisely, we ran 15 batches with num.runs = 10000 as a sanity check, to make sure that the final result based on 150000 aggregated clusterings was consistent with the results based on smaller batches, i.e., that it was in-sample stable.<sup>39</sup> Based on Table 4, we identify Clustering-A as the "ultimate" clustering (cf. Clustering-B/C/D).

<sup>&</sup>lt;sup>38</sup> The R function kmeans() produces a warning if it does not converge within iter.max.

<sup>&</sup>lt;sup>39</sup> We ran these 15 batches consecutively, and each batch produced the same top-10 (by occurrence counts) clusterings as in Table 4; however, the actual occurrence counts are different across the batches with slight variability in the corresponding rankings. The results are pleasantly stable.

We give the weights for Clustering-A, Clustering-B, Clustering-C and Clustering-D using unnormalized and normalized regressions with exposures computed based on arithmetic averages (see Subsection 2.6) in Tables 5, 6, 7, 8, 9, 10, 11, 12, and Figures 2 through 55. We give the weights for Clustering-A using unnormalized and normalized regressions with exposures computed based on geometric averages (see Subsection 2.6) in Tables 13, 14, and Figures 56 through 69. The actual mutation categories in each cluster for a given clustering can be read off the aforesaid Tables with the weights (the mutation categories with nonzero weights belong to a given cluster), or from the horizontal axis labels in the aforesaid Figures. It is evident that Clustering-A, Clustering-B, Clustering-C and Clustering-D are essentially variations of each other (Clustering-D has only 6 clusters, while the other 3 have 7 clusters).

### **3.3** Reconstruction and Correlations

So, based on genome data, we have constructed clusterings and weights. Do they work? I.e., do they reconstruct the input data well? It is evident from the get-go that the answer to this question may not be binary in the sense that for some cancer types we might have a nice clustering structure, while for others we may not. The aim of the following exercise is to sort this all out. Here come the correlations...

#### 3.3.1 Within-cluster Correlations

We have our de-noised<sup>40</sup> matrix  $G'_{is}$ . We are approximating this matrix via the following factorized matrix:

$$G_{is}^* = \sum_{A=1}^{K} W_{iA} \ H_{As} = w_i \ H_{G(i),s}$$
(22)

We can now compute an  $n \times K$  matrix  $\Theta_{sA}$  of *within-cluster* cross-sectional correlations between  $G'_{is}$  and  $G^*_{is}$  defined via  $(\mathbf{xCor}(\cdot, \cdot) \text{ stands for "cross-sectional correla$  $tion" to distinguish it from "serial correlation" <math>\mathbf{Cor}(\cdot, \cdot)$  we use above)<sup>41</sup>

$$\Theta_{sA} = \operatorname{xCor}(G'_{is}, G^*_{is})|_{i \in J(A)} = \operatorname{xCor}(G'_{is}, w_i)|_{i \in J(A)}$$
(23)

We give this matrix for Clustering-A with weights using normalized regressions with exposures computed based on arithmetic means (see Subsection 2.6) in Table 15. Let us mention that, with exposures based on arithmetic means, weights using normalized regressions work a bit better than using unnormalized regressions. Using exposures based on geometric means changes the weights a bit, which in turn slightly affects the within-cluster correlations, but does not alter the qualitative picture.

<sup>&</sup>lt;sup>40</sup> De-noising per se does not affect cross-sectional correlations. Adding extra 1 in (3) (recall that we obtain  $G'_{is}$  by cross-sectionally demeaning  $X_{is}$  and then re-exponentiating) has a negligible effect. So, in the correlations below we can use the original data matrix  $G_{is}$  instead of  $G'_{is}$ .

<sup>&</sup>lt;sup>41</sup> Due to the factorized structure (22), these correlations do not directly depend on  $H_{As}$ .

#### 3.3.2 Overall Correlations

Another useful metric, which we use as a sanity check, is this. For each value of s (i.e., for each cancer type), we can run a linear cross-sectional regression (without the intercept) of  $G'_{is}$  over the matrix  $W_{iA}$ . So, we have n = 14 of these regressions. Each regression produces multiple  $R^2$  and adjusted  $R^2$ , which we give in Table 15. Furthermore, we can compute the *fitted* values  $\hat{G}^*_{is}$  based on these regressions, which are given by

$$\widehat{G}_{is}^* = \sum_{A=1}^{K} W_{iA} \ F_{As} = w_i \ F_{G(i),s}$$
(24)

where (for each value of s)  $F_{As}$  are the regression coefficients. We can now compute the overall cross-sectional correlations (i.e., the index *i* runs over all N = 96 mutation categories)

$$\Xi_s = \operatorname{xCor}(G'_{is}, \widehat{G}^*_{is}) \tag{25}$$

These correlations are also given in Table 15 and measure the overall fit quality.

#### 3.3.3 Interpretation

Looking at Table 15 a few things become immediately evident. Clustering works well for 10 out the 14 cancer types we study here. The cancer types for which clustering does not appear to work all that well are Breast Cancer (labeled by X4 in Table 15), Liver Cancer (X8), Lung Cancer (X9), and Renal Cell Carcinoma (X14). More precisely, for Breast Cancer we do have a high within-cluster correlation for Cl-5 (and also Cl-4), but the overall fit is not spectacular due to low within-cluster correlations in other clusters. Also, above 80% within-cluster correlations<sup>42</sup> arise for 5 clusters, to wit, Cl-1, Cl-3, Cl-4, Cl-5 and Cl-6, but not for Cl-2 or Cl-7. Furthermore, remarkably, Cl-1 has high within-cluster correlations for 9 cancer types, and Cl-5 for 6 cancer types. These appear to be the leading clusters. Together they have high within-cluster correlations in 11 cancer types. So what does all this mean?

Additional insight is provided by looking at the within-cluster correlations between the 7 cancer signature extracted in [Kakushadze and Yu, 2016b] and the clusters we find here. Let  $W_{i\alpha}$  be the weights for the 7 cancer signatures from Tables 13 and 14 of [Kakushadze and Yu, 2016b]. We can compute the following within-cluster correlations ( $\alpha = 1, ..., 7$  labels the cancer signatures of [Kakushadze and Yu, 2016b], which we refer to as Sig1 through Sig7):

$$\Delta_{\alpha A} = \operatorname{xCor}(\mathcal{W}_{i\alpha}, W_{iA})|_{i \in J(A)}$$
(26)

These correlations are given in Table 16. High within-cluster correlations arise for Cl-1 (with Sig1 and Sig7), Cl-5 (with Sig2) and Cl-6 (with Sig4). And this makes perfect sense. Indeed, looking at Figures 14 through 20 of [Kakushadze and

 $<sup>^{42}\,</sup>$  The 80% cutoff is somewhat arbitrary, but reasonable.

Yu, 2016b], Sig1, Sig2, Sig4 and Sig7 are precisely the cancer signatures that have "peaks" (or "spikes" – "tall mountain landscapes"), whereas Sig3, Sig5 and Sig6 do not have such "peaks" ("flat" or "rolling hills landscapes"). No wonder such signatures do not have high within-cluster correlations – they simply do not have cluster-like structures. Looking at Figure 21 in [Kakushadze and Yu, 2016b], it becomes evident why clustering does not work well for Liver Cancer (X8) – it has a whopping 96% contribution from Sig5! Similarly, Renal Cell Carcinoma (X14) has a 70% contribution from Sig6. Lung Cancer (X9) is dominated by Sig3, hence no cluster-like structure. Finally, Breast Cancer (X4) is dominated by Sig2, which has a high within-cluster correlation with Cl-5, which is why Breast Cancer has a high within-cluster correlation with Cl-5 (but poor overall correlation in Table 15). So, it all makes sense. The question is, what does all this tell us about cancer signatures?

Quite a bit! It tells us that cancers such as Liver Cancer, Lung Cancer and Renal Cell Carcinoma have little in common with other cancers (and each other)! At least at the level of mutation categories that dominate the genome structure of such cancers. On the other hand, 9 cancers, to wit, Bone Cancer (X2), Brain Lower Grade Glioma (X3), Chronic Lymphocytic Leukemia (X5), Esophageal Cancer (X6), Gastric Cancer (X7), Medulloblastoma (X10), Ovarian Cancer (X11), Pancreatic Cancer (X12) and Prostate Cancer (X13) apparently all have the Cl-1 cluster structure embedded in them substantially. Similarly, 6 cancers, to wit, B Cell Lymphoma (X1), Breast Cancer (X4), Esophageal Cancer(X6), Ovarian Cancer (X11), Pancreatic Cancer (X12) and Prostate Cancer (X13) apparently all have the Cl-5 cluster structure embedded in them substantially. Furthermore, note the overlap between these two lists, to wit, Esophageal Cancer(X6), Ovarian Cancer (X11), Pancreatic Cancer (X12) and Prostate Cancer (X13). We obtained this result purely statistically, with no biologic input, using our clustering algorithm and other statistical methods such as linear regression to obtain the actual weights. It is too early to know whether this insight will aid any therapeutic applications, but that is the hope - similarities in the underlying genomic structures of different cancer types raise hope that therapeutics for one cancer type could perhaps be applicable to other cancer types. On the other hand, our findings above relating to Liver Cancer, Lung Cancer and Renal Cell Carcinoma (and possibly also Breast Cancer, albeit the latter does appear to have a not-so-insignificant overlap with Cl-5, which differentiates it from the aforesaid 3 cancer types) suggest that these cancer types apparently stand out.

# 4 Concluding Remarks

Clustering ideas and techniques have been applied in cancer research in various incarnations and contexts aplenty – for a partial list of works at least to some extent related to our discussion here, see, e.g, [Chen *et al*, 2008a], [Chen *et al*, 2008b], [Kashuba *et al*, 2009], [Nik-Zainal *et al*, 2012], [Roberts *et al*, 2012], [Alexandrov *et al*, 2013a], [Alexandrov *et al*, 2013b], [Burns *et al*, 2013a], [Burns *et al*, 2013b], [Lawrence

et al, 2013], [Long et al, 2013] [Roberts et al, 2013], [Taylor et al, 2013], [Xuan et al, 2013], [Alexandrov and Stratton, 2014], [Bacolla et al, 2014], [Bolli et al, 2014], [Caval et al, 2014], [Davis et al, 2014], [Helleday et al, 2014], [Nik-Zainal et al, 2014], [Poon et al, 2014], [Qian et al, 2014], [Roberts and Gordenin, 2014a], [Roberts and Gordenin, 2014b], [Roberts and Gordenin, 2014c], [Sima and Gilbert, 2014], [Chan and Gordenin, 2015], [Pettersen et al, 2015] and references therein. As mentioned above, even in NMF clustering is used at some (perhaps not-so-evident) layer. What is new in our approach – and hence new results – is that: i) following [Kakushadze and Yu, 2016b], we apply clustering to aggregated by cancer types and de-noised data; ii) we use a tried-and-tested in quantitative finance bag of tricks from [Kakushadze and Yu, 2016c], which improves clustering; and iii) last but not least, we apply our \*K-means algorithm to cancer genome data. As mentioned above, \*Kmeans, unlike vanilla k-means or its other commonly used variations, is essentially deterministic, and it achieves determinism *statistically*, not by "guessing" initial centers or as in agglomerative hierarchical clustering, which basically "guesses" the initial (e.g., 2-cluster) clustering. Instead, via aggregating a large number of k-means clusterings and statistical examination of the occurrence counts of such aggregations, \*K-means takes a mess of myriad vanilla k-means clusterings and systematically reduces randomness and indeterminism without *ad hoc* initial "guesswork".

As mentioned above, consistently with the results of [Kakushadze and Yu, 2016b] obtained via improved NMF techniques, Liver Cancer, Lung Cancer and Renal Cell Carcinoma do not appear to have clustering (sub)structures. This could be both good and bad news. It is a good news because we learned something interesting about these cancer types – and in two complementary ways. However, it could also be a bad news from the therapeutic standpoint. Since these cancer types appear to have little in common with others, it is likely that they would require specialized therapeutics. On the flipside, we should note that it would make sense to exclude these 3 cancer types when running clustering analysis. However, it would also make sense to include other cancer types by utilizing the International Cancer Genome Consortium data, which we leave for future studies. (For comparative reasons, here we used the same data as in [Kakushadze and Yu, 2016b], which was limited to data samples published as of the date thereof.) This paper is not intended to be an exhaustive empirical study but a proof of concept and an opening of a new avenue for extracting and studying cancer signatures beyond the tools that NMF provides.

And we do find that 11 out of the 14 cancer types we study here have clustering structures substantially embedded in them and clustering overall works well for at least 10 out of these 11 cancer types.<sup>43</sup> Now, looking at Figure 14 of [Kakushadze and Yu, 2016b], we see that its "peaks" are located at ACGT, CCGT, GCGT and TCGT. The same "peaks" are present in our cluster Cl-1 (see Figures 2 and 3). Hence the high within-cluster correlation between Cl-1 and Sig1. On the other

<sup>&</sup>lt;sup>43</sup> Breast Cancer possibly being an exception. As mentioned above, it would make sense to exclude Liver Cancer, Lung Cancer and Renal Cell Carcinoma from the analysis, which may affect how well clustering works for Breast Cancer and possibly also the other 10 cancer types.

hand, Sig1 of [Kakushadze and Yu, 2016b] is essentially the same as the mutational signature 1 of [Nik-Zainal et al, 2012], [Alexandrov et al, 2013b], which is due to spontaneous cytosine deamination. So, this is what our cluster Cl-1 describes. Next, looking at Figure 15 of [Kakushadze and Yu, 2016b], we see that its "peaks" are located at TCAG, TCTG, TCAT and TCTT. The first two of these "peaks" TCAG and TCTG are present in our Cl-5 (see Figures 10 and 11), the third "peak" TCAT is present in our Cl-1 (see Figures 2 and 3), while the fourth "peak" TCTT is present in our Cl-4 (see Figures 8 and 9), which is consistent with the high within-cluster correlations between Sig2 and Cl-4 and Cl-5, albeit its within-cluster correlation with Cl-1 is poor. Note that Sig2 of [Kakushadze and Yu, 2016b] is essentially the same as the mutational signatures 2+13 of [Nik-Zainal et al, 2012], [Alexandrov et al, 2013b], which are due to APOBEC mediated cytosine deamination. In fact, it was reported as a single signature in [Alexandrov et al, 2013b], however, subsequently, it was split into 2 distinct signatures, which usually appear in the same samples.<sup>44</sup> Our clustering results indicate that grouping TCAG and TCTG into one signature makes sense as they belong to the same cluster Cl-5. However, grouping TCAT and TCTT together does not appear to make much sense. Looking at the Figures for Clustering-A, Clustering-B, Clustering-C and Clustering-D, we see that the TCAT "peak" invariably appears together with the ACGT, CCGT, GCGT and TCGT "peaks" as in Cl-1 in Clustering-A, Cl-2 in Clustering-B, Cl-1 in Clustering-C, and Cl-1 in Clustering-D, but never with TCTT. So, our clustering approach tells us something new beyond the NMF "intuition". This may have an important implication for Breast Cancer, which, as mentioned above, is dominated by Sig2. Thus, based on our results in Table 15, we see that Breast Cancer has high within-cluster correlations with Cl-4 and Cl-5, but not with Cl-1. This may imply that clustering simply does not work well for Breast Cancer, which would appear to put it in the same "standalone" league as Liver Cancer, Lung Cancer and Renal Cell Carcinoma. In any event, clustering invariably suggests that the TCAT "peak" belongs in Cl-1 with the 4 "peaks" ACGT, CCGT, GCGT and TCGT related to spontaneous cytosine deamination, rather than those related to APOBEC mediated cytosine deamination.

Now, let us check the remaining two signatures of [Kakushadze and Yu, 2016b] with "tall mountain landscapes" (see above), to wit, Sig4 and Sig7. Looking at Figure 17 of [Kakushadze and Yu, 2016b], we see that its "peaks" are at CTTC, TTTC, CTTG and TTTG. The same peaks appear in our Cl-6 (see Figures 12 and 13). Hence the high within-cluster correlation between Cl-6 and Sig4. Note that Sig4 is essentially the same as the mutational signature 17 of [Nik-Zainal *et al*, 2012], [Alexandrov *et al*, 2013b], and its underlying mutational process is unknown. Next, looking at Figure 20 of [Kakushadze and Yu, 2016b], we see that its "peaks" for the C > G mutations are essentially the same as in Cl-1. Hence the high within-cluster correlation between Cl-7 and Sig1. So, there are no surprises with Sig1, Sig4 and Sig7. However, based on our clustering results, as we discuss above, with Sig2

<sup>&</sup>lt;sup>44</sup> For detailed comments, see http://cancer.sanger.ac.uk/cosmic/signatures.

we do find – what we feel is a pleasant – surprise, that splitting it into two signatures (see above) might be inadequate and the TCAT "peak" might really belong with the Sig1 "peaks" (spontaneous v. APOBEC mediated cytosine deamination). This is exciting as it might be an indication of the limitations of NMF (or clustering...).<sup>45</sup>

In Introduction we promised that we would discuss some potential applications of \*K-means in quantitative finance, and so here it is. Let us mention that \*K-means is universal, oblivious to the input data and applicable in a variety of fields. In quantitative finance \*K-means a priori can be applied everywhere clustering methods are used with the added bonus of (statistical) determinism.<sup>46</sup> One evident example is statistical industry classifications discussed in [Kakushadze and Yu, 2016c], where one uses clustering methods to classify stocks. In fact, \*K-means is an extension of the methods discussed in [Kakushadze and Yu, 2016c]. One thing to keep in mind is that in \*K-means one sifts through a large number P of aggregations, which can get computationally costly when clustering 2000+ stocks into 100+ clusters.<sup>47</sup> Another potential application is in the context of combining alphas (trading signals) - see, e.g., [Kakushadze and Yu, 2017a]. Yet another application is when we have a term structure, such as a portfolio of bonds (e.g., U.S. Treasuries or some other bonds) with varying maturities, or futures (e.g., Eurodollar futures) with varying deliveries. These cases resemble the genome data more in the sense that the number N of instruments is relatively small (typically even fewer than the number of mutation categories). Another example with a relatively small number of instruments would be a portfolio of various futures for different FX (foreign exchange) pairs (even with the uniform delivery), e.g., USD/EUR, USD/HKD, EUR/AUD, etc., i.e., FX statistical arbitrage. One approach to optimizing risk in such portfolios is by employing clustering methods and a stable, essentially deterministic algorithm such as \*K-means can be useful. Hopefully \*K-means will prove a valuable tool in cancer research, quantitative finance as well as various other fields (e.g., image recognition).

# A R Source Code

In this appendix we give the R (R Package for Statistical Computing, http://www.rproject.org) source code for computing the clusterings and weights using the algorithms of Section 2. The code is straightforward and self-explanatory.<sup>48</sup> The main function is bio.cl.sigs(x, iter.max = 100, num.try = 1000, num.runs = 10000). Here: x is the  $N \times d$  occurrence counts matrix  $G_{is}$  (where N = 96 is the number of

 $<sup>^{45}</sup>$  Or both... Alternatively – and that would be truly exciting – perhaps there is a biologic explanation. In any event, it is too early to tell – yet another possibility is that this is merely an artifact of the dataset we use. More research and analyses on larger datasets (see above) is needed.

 $<sup>^{46}</sup>$  Albeit with the understanding that it requires additional computational cost.

<sup>&</sup>lt;sup>47</sup> This can be mitigated by employing top-down clustering [Kakushadze and Yu, 2016c].

<sup>&</sup>lt;sup>48</sup> The source code in Appendix A hereof is not written to be "fancy" or optimized for speed or in any other way. Its sole purpose is to illustrate the algorithms described in the main text in a simple-to-understand fashion. See Appendix B for some important legalese.

mutation categories, and d is the number of samples; or d = n, where n is the number of cancer types, when the samples are aggregated by cancer types); iter.max is the maximum number of iterations that are passed into the R built-in function kmeans(); num.try is the number M of aggregated clusterings (see Subsection 2.3.2); num.runs is the number of runs P used to determine the most frequently occurring clustering (the "ultimate" clustering) obtained via aggregation (see Subsection 2.3.3). The function bio.erank.pc() is defined in Appendix B of [Kakushadze and Yu, 2016b]. The function grm.stat.ind.class() is defined in Appendix A of [Kakushadze and Yu, 2016c]. This function internally calls another function grm.calc.norm.ret(), which we redefine here via the function bio.calc.norm.ret().49 The output is a list, whose elements are as follows: res\$ind is an  $N \times K$  binary matrix  $\Omega_{iA} = \delta_{G(i),A}$  $(i = 1, ..., N, A = 1, ..., K, \text{ the map } G : \{1, ..., N\} \mapsto \{1, ..., K\} - \text{see Section } 2),$ which defines the K clusters in the "ultimate" clustering;<sup>50</sup> resw is an N-vector of weights obtained via unnormalized regressions using arithmetic means for computing exposures (i.e., via (13), (14) and (15)); resv is an N-vector of weights obtained via normalized regressions using arithmetic means for computing exposures (i.e., via (17), (14) and (16)); resw.g is an N-vector of weights obtained via unnormalized regressions using geometric means for computing exposures (i.e., via (13), (14) and (19)); res\$v.g is an N-vector of weights obtained via normalized regressions using geometric means for computing exposures (i.e., via (17), (14) and (21)).

```
bio.calc.norm.ret <- function (ret)
{
    s <- apply(ret, 1, sd)
    x <- ret / s
    return(x)
}

qrm.calc.norm.ret <- bio.calc.norm.ret
bio.cl.sigs <- function(x, iter.max = 100,
    num.try = 1000, num.runs = 10000)
{
    cl.ix <- function(x) match(1, x)
    y <- log(1 + x)
    y <- t(t(y) - colMeans(y))</pre>
```

<sup>&</sup>lt;sup>49</sup> The definition of qrm.calc.norm.ret() in [Kakushadze and Yu, 2016c] accounts for some peculiarities and nuances pertinent to quantitative trading, which are not applicable here.

<sup>&</sup>lt;sup>50</sup> The code returns the K clusters ordered such that the number of mutation  $n_A$  (i.e., the column sum of  $\Omega_{iA}$ ) in the cluster labeled by A is in the increasing order. It also orders clusters with identical  $n_A$ . We note, however, that (for presentational convenience reasons) the order of such clusters in the tables and figures below is not necessarily the same as what this code returns.

```
x.d <- exp(y)
   k <- ncol(bio.erank.pc(y)$pc)</pre>
   n <- nrow(x)
   u <- rnorm(n, 0, 1)
   q <- matrix(NA, n, num.runs)</pre>
   p <- rep(NA, num.runs)</pre>
   for(i in 1:num.runs)
   {
       z <- qrm.stat.ind.class(y, k, iter.max = iter.max,</pre>
          num.try = num.try, demean.ret = F)
       p[i] \leftarrow sum((residuals(lm(u \sim -1 + z)))^2)
       q[, i] <- apply(z, 1, cl.ix)</pre>
   }
   p1 <- unique(p)</pre>
   ct <- rep(NA, length(p1))</pre>
   for(i in 1:length(p1))
       ct[i] <- sum(p1[i] == p)
   p1 <- p1[ct == max(ct)]</pre>
   i <- match(p1, p)[1]
   ix <- q[, i]
   k < - max(ix)
   z <- matrix(NA, n, k)</pre>
   for(j in 1:k)
       z[, j] <- as.numeric(ix == j)</pre>
   res <- bio.cl.wts(x.d, z)</pre>
   return(res)
bio.cl.wts <- function (x, ind)</pre>
   first.ix <- function(x) match(1, x)[1]</pre>
   calc.wts <- function(x, use.wts = F, use.geom = F)</pre>
   {
       if(use.geom)
       {
          if(use.wts)
```

}

{

```
s <- apply(log(x), 1, sd)</pre>
       else
          s \leftarrow rep(1, nrow(x))
       s <- 1 / s / sum(1 / s)
       fac <- apply(x^s, 2, prod)</pre>
   }
   else
   {
       if(use.wts)
          s <- apply(x, 1, sd)</pre>
       else
          s \leftarrow rep(1, nrow(x))
       fac <- colMeans(x / s)</pre>
   }
   w <- coefficients(lm(t(x) \sim -1 + fac))
   w <- 100 * w / sum(w)
   return(w)
}
n <- nrow(x)
w <- w.g <- v <- v.g <- rep(NA, n)
z <- colSums(ind)</pre>
z <- as.numeric(paste(z, ".", apply(ind, 2, first.ix), sep = ""))</pre>
dimnames(ind)[[2]] <- names(z) <- 1:ncol(ind)
z \leftarrow sort(z)
z < - names(z)
ind <- ind[, z]
dimnames(ind)[[2]] <- NULL</pre>
for(i in 1:ncol(ind))
{
   take <- ind[, i] == 1</pre>
   if(sum(take) == 1)
   {
       w[take] <- w.g[take] <- 1
       v[take] <- v.g[take] <- 1</pre>
       next
   }
   w[take] <- calc.wts(x[take, ], F, F)</pre>
   w.g[take] <- calc.wts(x[take, ], F, T)</pre>
   v[take] <- calc.wts(x[take, ], T, F)</pre>
```

```
v.g[take] <- calc.wts(x[take, ], T, T)
}
res <- new.env()
res$ind <- ind
res$w <- w
res$w.g <- w.g
res$v.g <- v.g
res$v.g <- v.g
return(res)
}</pre>
```

# **B** DISCLAIMERS

Wherever the context so requires, the masculine gender includes the feminine and/or neuter, and the singular form includes the plural and *vice versa*. The author of this paper ("Author") and his affiliates including without limitation Quantigic<sup>®</sup> Solutions LLC ("Author's Affiliates" or "his Affiliates") make no implied or express warranties or any other representations whatsoever, including without limitation implied warranties of merchantability and fitness for a particular purpose, in connection with or with regard to the content of this paper including without limitation any code or algorithms contained herein ("Content").

The reader may use the Content solely at his/her/its own risk and the reader shall have no claims whatsoever against the Author or his Affiliates and the Author and his Affiliates shall have no liability whatsoever to the reader or any third party whatsoever for any loss, expense, opportunity cost, damages or any other adverse effects whatsoever relating to or arising from the use of the Content by the reader including without any limitation whatsoever: any direct, indirect, incidental, special, consequential or any other damages incurred by the reader, however caused and under any theory of liability; any loss of profit (whether incurred directly or indirectly), any loss of goodwill or reputation, any loss of data suffered, cost of procurement of substitute goods or services, or any other tangible or intangible loss; any reliance placed by the reader on the completeness, accuracy or existence of the Content or any other effect of using the Content; and any and all other adversities or negative effects the reader might encounter in using the Content irrespective of whether the Author or his Affiliates is or are or should have been aware of such adversities or negative effects.

The R code included in Appendix A hereof is part of the copyrighted R code of Quantigic<sup>®</sup> Solutions LLC and is provided herein with the express permission of Quantigic<sup>®</sup> Solutions LLC. The copyright owner retains all rights, title and interest in and to its copyrighted source code included in Appendix A hereof and any and all copyrights therefor.

# References

Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Campbell, P.J. and Stratton, M.R. (2013a) Deciphering Signatures of Mutational Processes Operative in Human Cancer. *Cell Reports* 3(1): 246-259.

Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Aparicio, S.A., Behjati, S., Biankin, A.V., Bignell, G.R., Bolli, N., Borg, A., Børresen-Dale, A.L., Boyault, S., Burkhardt, B., Butler, A.P., Caldas, C., Davies, H.R., Desmedt, C., Eils, R., Eyfjörd, J.E., Foekens, J.A., Greaves, M., Hosoda, F., Hutter, B., Ilicic, T., Imbeaud, S., Imielinski, M., Jäger, N., Jones, D.T., Jones, D., Knappskog, S., Kool, M., Lakhani, S.R., López-Otín, C., Martin, S., Munshi, N.C., Nakamura, H., Northcott, P.A., Pajic, M., Papaemmanuil, E., Paradiso, A., Pearson, J.V., Puente, X.S., Raine, K., Ramakrishna, M., Richardson, A.L., Richter, J., Rosenstiel, P., Schlesner, M., Schumacher, T.N., Span, P.N., Teague, J.W., Totoki, Y., Tutt, A.N., Valdés-Mas, R., van Buuren, M.M., van 't Veer, L., Vincent-Salomon, A., Waddell, N., Yates, L.R.; Australian Pancreatic Cancer Genome Initiative; ICGC Breast Cancer Consortium; ICGC MMML-Seq Consortium; ICGC PedBrain, Zucman-Rossi, J., Futreal, P.A., McDermott, U., Lichter, P., Meyerson, M., Grimmond, S.M., Siebert, R., Campo, E., Shibata, T., Pfister, S.M., Campbell, P.J., Stratton, M.R. (2013b) Signatures of mutational processes in human cancer. Nature 500(7463): 415-421.

Alexandrov, L.B. and Stratton, M.R. (2014) Mutational signatures: the patterns of somatic mutations hidden in cancer genomes. Current Opinion in Genetics & Development 24: 52-60.

Ananthaswamy, H.N. and Pierceall, W.E. (1990) Molecular mechanisms of ultraviolet radiation carcinogenesis. *Photochemistry and Photobiology* 52(6): 1119-1136.

Bacolla, A., Cooper, D.N. and Vasquez, K.M. (2014) Mechanisms of base substitution mutagenesis in cancer genomes. *Genes* 5(1): 108-146.

Bai, J. and Ng, S. (2002) Determining the number of factors in approximate factor models. *Econometrica* 70(1): 191-221.

Bolli, N., Avet-Loiseau, H., Wedge, D.C., Van Loo, P., Alexandrov, L.B., Martincorena, I., Dawson, K.J., Iorio, F., Nik-Zainal, S., Bignell, G.R., Hinton, J.W., Li, Y., Tubio, J.M., McLaren, S., O' Meara, S., Butler, A.P., Teague, J.W., Mudie, L., Anderson, E., Rashid, N., Tai, Y.T., Shammas, M.A., Sperling, A.S., Fulciniti, M., Richardson, P.G., Parmigiani, G., Magrangeas, F., Minvielle, S., Moreau, P., Attal, M., Facon, T., Futreal, P.A., Anderson, K.C., Campbell, P.J., Munshi, N.C. (2014) Heterogeneity of genomic evolution and mutational profiles in multiple myeloma. *Nature Communications* 5: 2997. Bouchaud, J.-P. and Potters, M. (2011) Financial applications of random matrix theory: a short review. In: Akemann, G., Baik, J. and Di Francesco, P. (eds.) *The Oxford Handbook of Random Matrix Theory.* Oxford, United Kingdom: Oxford University Press.

Burns, M.B., Lackey, L., Carpenter, M.A., Rathore, A., Land, A.M., Leonard, B., Refsland, E.W., Kotandeniya, D., Tretyakova, N., Nikas, J.B., Yee, D., Temiz, N.A., Donohue, D.E., McDougle, R.M., Brown, W.L., Law, E.K., Harris, R.S. (2013a) APOBEC3B is an enzymatic source of mutation in breast cancer. *Nature* 494(7437): 366-370.

Burns, M.B., Temiz, N.A. and Harris, R.S. (2013b) Evidence for APOBEC3B mutagenesis in multiple human cancers. *Nature Genetics* 45(9): 977-983.

Caval, V., Suspène, R., Shapira, M., Vartanian, J.P. and Wain-Hobson, S. (2014) A prevalent cancer susceptibility APOBEC3A hybrid allele bearing APOBEC3B 3'UTR enhances chromosomal DNA damage. *Nature Communications* 5: 5129.

Campbell, L.L. (1960) Minimum coefficient rate for stationary random processes. *Information and Control* 3(4): 360-371.

Chan, K. and Gordenin, D.A. (2015) Clusters of Multiple Mutations: Incidence and Molecular Mechanisms. *Annual Review of Genetics* 49: 243-627

Chen, Z., Feng, J., Buzin, C.H. and Sommer, S.S. (2008a) Epidemiology of doublet/multiplet mutations in lung cancers: evidence that a subset arises by chronocoordinate events. *PloS One* 3(11): e3714.

Chen, Z., Feng, J., Saldivar, J.S., Gu, D., Bockholt, A., and Sommer, S.S. (2008b) EGFR somatic doublets in lung cancer are frequent and generally arise from a pair of driver mutations uncommonly seen as singlet mutations: one-third of doublets occur at five pairs of amino acids. *Oncogene* 27(31): 4336-4343.

Cheng, C., Zhou, Y., Li, H., Xiong, T., Li, S., Bi, Y., Kong, P., Wang, F., Cui, H., Li, Y., Fang, X., Yan, T., Li, Y., Wang, J., Yang, B., Zhang, L., Jia, Z., Song, B., Hu, X., Yang, J., Qiu, H., Zhang, G., Liu, J., Xu, E., Shi, R., Zhang, Y., Liu, H., He, C., Zhao, Z., Qian, Y., Rong, R., Han, Z., Zhang, Y., Luo, W., Wang, J., Peng, S., Yang, X., Li, X., Li, L., Fang, H., Liu, X., Ma, L., Chen, Y., Guo, S., Chen, X., Xi, Y., Li, G., Liang, J., Yang, X., Guo, J., Jia, J., Li, Q., Cheng, X., Zhan, Q., Cui, Y. (2016) Whole-Genome Sequencing Reveals Diverse Models of Structural Variations in Esophageal Squamous Cell Carcinoma. *The American Journal of Human Genetics* 98(2): 256-274.

Connor, G. and Korajczyk, R.A. (1993) A Test for the Number of Factors in an Approximate Factor Model. *The Journal of Finance* 48(4): 1263-1291.

Davis, C.F., Ricketts, C.J., Wang, M., Yang, L., Cherniack, A.D., Shen, H., Buhay, C., Kang, H., Kim, S.C., Fahey, C.C., Hacker, K.E., Bhanot, G., Gordenin, D.A., Chu, A., Gunaratne, P.H., Biehl, M., Seth, S., Kaipparettu, B.A., Bristow, C.A., Donehower, L.A., Wallen, E.M., Smith, A.B., Tickoo, S.K., Tamboli, P., Reuter, V., Schmidt, L.S., Hsieh, J.J., Choueiri, T.K., Hakimi, A.A.; Cancer Genome Atlas Research Network, Chin, L., Meyerson, M., Kucherlapati, R., Park, W.Y., Robertson, A.G., Laird, P.W., Henske, E.P., Kwiatkowski, D.J., Park, P.J., Morgan, M., Shuch, B., Muzny, D., Wheeler, D.A., Linehan, W.M., Gibbs, R.A., Rathmell, W.K., Creighton, C.J. (2014) The somatic genomic landscape of chromophobe renal cell carcinoma. *Cancer Cell* 26(3): 319-330.

De Amorim, R.C. and Hennig, C. (2015) Recovering the number of clusters in data sets with noise features using feature rescaling factors. *Information Sciences* 324: 126-145.

Forgy, E.W. (1965) Cluster analysis of multivariate data: efficiency versus interpretability of classifications. *Biometrics* 21(3): 768-769.

Fujimoto, A., Furuta, M., Totoki, Y., Tsunoda, T., Kato, M., Shiraishi, Y., Tanaka, H., Taniguchi, H., Kawakami, Y., Ueno, M., Gotoh, K., Ariizumi, S., Wardell, C.P., Hayami, S., Nakamura, T., Aikata, H., Arihiro, K., Boroevich, K.A., Abe, T., Nakano, K., Maejima, K., Sasaki-Oku, A., Ohsawa, A., Shibuya, T., Nakamura, H., Hama, H., Hosoda, F., Arai, Y., Ohashi, S., Urushidate, T., Nagae, G., Yamamoto, S., Ueda, H., Tatsuno, K., Ojima, H., Hiraoka, N., Okusaka, T., Kubo, M., Marubashi, S., Yamada, T., Hirano, S., Yamamoto, M., Ohdan, H., Shimada, K., Ishikawa, O., Yamaue, H., Chayama, K., Miyano, S., Aburatani, H., Shibata, T., Nakagawa, H. (2016) Whole-genome mutational landscape and characterization of noncoding and structural mutations in liver cancer. *Nature Genetics* 48(5): 500-509.

Goodman, M.F. and Fygenson, K.D. (1998) DNA polymerase fidelity: from genetics toward a biochemical understanding. *Genetics* 148(4): 1475-1482.

Goutte, C., Hansen, L.K., Liptrot, M.G. and Rostrup, E. (2001) Feature-Space Clustering for fMRI Meta-Analysis. *Human Brain Mapping* 13(3): 165-183.

Grinold, R.C. and Kahn, R.N. (2000) *Active Portfolio Management*. New York, NY: McGraw-Hill.

Gundem, G., Van Loo, P., Kremeyer, B., Alexandrov, L.B., Tubio, J.M., Papaemmanuil, E., Brewer, D.S., Kallio, H.M., Högnäs, G., Annala, M., Kivinummi, K., Goody, V., Latimer, C., O'Meara, S., Dawson, K.J., Isaacs, W., Emmert-Buck, M.R., Nykter, M., Foster, C., Kote-Jarai, Z., Easton, D., Whitaker, H.C.; ICGC Prostate UK Group, Neal, D.E., Cooper, C.S., Eeles, R.A., Visakorpi, T., Campbell, P.J., McDermott, U., Wedge, D.C., Bova, G.S. (2015) The evolutionary history of lethal metastatic prostate cancer. *Nature* 520(7547): 353-357.

Hamerly, G. and Elkan, C. (2004) Learning the k in k-means. In: Thrun, S. (ed.) Advances of the Neural Information Processing Systems, Vol. 16. Campridge, MA: MIT Press, pp. 281-289.

Hartigan, J.A. (1975) Clustering algorithms. New York, NY: John Wiley & Sons, Inc.

Hartigan, J.A. and Wong, M.A. (1979) Algorithm AS 136: A K-Means Clustering Algorithm. *Journal of the Royal Statistical Society, Series C (Applied Statistics)* 28(1): 100-108.

Helleday, T., Eshtad, S. and Nik-Zainal, S. (2014) Mechanisms underlying mutational signatures in human cancers. *Nature Reviews Genetics* 15(9): 585-598.

Imielinski, M., Berger, A.H., Hammerman, P.S., Hernandez, B., Pugh, T.J., Hodis, E., Cho, J., Suh, J., Capelletti, M., Sivachenko, A., Sougnez, C., Auclair, D., Lawrence, M.S., Stojanov, P., Cibulskis, K., Choi, K., de Waal, L., Sharifnia, T., Brooks, A., Greulich, H., Banerji, S., Zander, T., Seidel, D., Leenders, F., Ansén, S., Ludwig, C., Engel-Riedel, W., Stoelben, E., Wolf, J., Goparju, C., Thompson, K., Winckler, W., Kwiatkowski, D., Johnson, B.E., Jänne, P.A., Miller, V.A., Pao, W., Travis, W.D., Pass, H.I., Gabriel, S.B., Lander, E.S., Thomas, R.K., Garraway, L.A., Getz, G., Meyerson, M. (2012) Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* 150(6): 1107-1120.

Jones, D.T., Jäger, N., Kool, M., Zichner, T., Hutter, B., Sultan, M., Cho, Y.J.,
Pugh, T.J., Hovestadt, V., Stütz, A.M., Rausch, T., Warnatz, H.J., Ryzhova,
M., Bender, S., Sturm, D., Pleier, S., Cin, H., Pfaff, E., Sieber, L., Wittmann,
A., Remke, M., Witt, H., Hutter, S., Tzaridis, T., Weischenfeldt, J., Raeder, B.,
Avci, M., Amstislavskiy, V., Zapatka, M., Weber, U.D., Wang, Q., Lasitschka,
B., Bartholomae, C.C., Schmidt, M., von Kalle, C., Ast, V., Lawerenz, C., Eils,
J., Kabbe, R., Benes, V., van Sluis, P., Koster, J., Volckmann, R., Shih, D.,
Betts, M.J., Russell, R.B., Coco, S., Tonini, G.P., Schüller, U., Hans, V., Graf,
N., Kim, Y.J., Monoranu, C., Roggendorf, W., Unterberg, A., Herold-Mende,
C., Milde, T., Kulozik, A.E., von Deimling, A., Witt, O., Maass, E., Rössler,
J., Ebinger, M., Schuhmann, M.U., Frühwald, M.C., Hasselblatt, M., Jabado,
N., Rutkowski, S., von Bueren, A.O., Williamson, D., Clifford, S.C., McCabe,
M.G., Collins, V.P., Wolf, S., Wiemann, S., Lehrach, H., Brors, B., Scheurlen,
W., Felsberg, J., Reifenberger, G., Northcott, P.A., Taylor, M.D., Meyerson,
M., Pomeroy, S.L., Yaspo, M.L., Korbel, J.O., Korshunov, A., Eils, R., Pfister,

S.M., Lichter, P. (2012) Dissecting the genomic complexity underlying medulloblastoma. *Nature* 488(7409): 100-105.

Kakushadze, Z. and Yu, W. (2016a) Multifactor Risk Models and Heterotic CAPM. *The Journal of Investment Strategies* 5(4): 1-49. Available online: http://ssrn.com/abstract=2722093.

Kakushadze, Z. and Yu, W. (2016b) Factor Models for Cancer Signatures. *Physica A* 462: 527-559. Available online: http://ssrn.com/abstract=2772458.

Kakushadze, Z. and Yu, W. (2016c) Statistical Industry Classification. *Journal of Risk & Control* 3(1): 17-65. Available online: http://ssrn.com/abstract=2802753.

Kakushadze, Z. and Yu, W. (2017a) How to Combine a Billion Alphas. *Journal* of Asset Management 18(1): 64-80. Available online: http://ssrn.com/abstract=2739219.

Kakushadze, Z. and Yu, W. (2017b) Statistical Risk Models. *The Journal of Investment Strategies* 6(2): 1-40. Available online: http://ssrn.com/abstract=2732453.

Kashuba, V.I., Pavlova, T.V., Grigorieva, E.V., Kutsenko, A., Yenamandra, S.P., Li, J., Wang, F., Protopopov, A.I., Zabarovska, V.I., Senchenko, V., Haraldson, K., Eshchenko, T., Kobliakova, J., Vorontsova, O., Kuzmin, I., Braga, E., Blinov, V.M., Kisselev, L.L., Zeng, Y.-X., Ernberg, I., Lerman, M.I., Klein, G. and Zabarovsky, E.R. (2009) High mutability of the tumor suppressor genes RASSF1 and RBSP3 (CTDSPL) in cancer. *PloS One* 4(5): e5231.

Lawrence, M.S., Stojanov, P., Polak, P., Kryukov, G.V., Cibulskis, K., Sivachenko, A., Carter, S.L., Stewart, C., Mermel, C.H., Roberts, S.A., Kiezun, A., Hammerman, P.S., McKenna, A., Drier, Y., Zou, L., Ramos, A.H., Pugh, T.J., Stransky, N., Helman, E., Kim, J., Sougnez, C., Ambrogio, L., Nickerson, E., Shefler, E., Cortés, M.L., Auclair, D., Saksena, G., Voet, D., Noble, M., DiCara, D., Lin, P., Lichtenstein, L., Heiman, D.I., Fennell, T., Imielinski, M., Hernandez, B., Hodis, E., Baca, S., Dulak, A.M., Lohr, J., Landau, D.A., Wu, C.J., Melendez-Zajgla, J., Hidalgo-Miranda, A., Koren, A., McCarroll, S.A., Mora, J., Lee, R.S., Crompton, B., Onofrio, R., Parkin, M., Winckler, W., Ardlie, K., Gabriel, S.B., Roberts C.W., Biegel, J.A., Stegmaier, K., Bass, A.J., Garraway, L.A., Meyerson, M., Golub, T.R., Gordenin, D.A., Sunyaev, S., Lander, E.S., Getz, G. (2013) Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 499(7457): 214-208.

Lee, D.D. and Seung, H.S. (1999) Learning the parts of objects by non-negative matrix factorization. *Nature* 401(6755): 788-791.

Lindahl, T. (1993) Instability and decay of the primary structure of DNA. *Nature* 362(6422): 709-715.

Lletí, R, Ortiz, M.C., Sarabia, L.A. and Sánchez, M.S. (2004) Selecting Variables for k-Means Cluster Analysis by Using a Genetic Algorithm that Optimises the Silhouettes. *Analytica Chimica Acta* 515(1): 87-100.

Lloyd, S.P. (1957) Least square quantization in PCM. Working Paper. Bell Telephone Laboratories, Murray Hill, NJ.

Lloyd, S.P. (1982) Least square quantization in PCM. *IEEE Transactions on Information Theory* 28(2): 129-137.

Loeb, L.A. and Harris, C.C. (2008) Advances in chemical carcinogenesis: a historical review and perspective. *Cancer Research* 68(17): 6863-6872.

Long, J., Delahanty, R.J., Li, G., Gao, Y.T., Lu, W., Cai, Q., Xiang, Y.B., Li, C., Ji, B.T., Zheng, Y., Ali, S., Shu, X.O., Zheng, W. (2013) A common deletion in the APOBEC3 genes and breast cancer risk. *Journal of National Cancer Institute* 105(8): 573-579.

Love, C., Sun, Z., Jima, D., Li, G., Zhang, J., Miles, R., Richards, K.L., Dunphy, C.H., Choi, W.W., Srivastava, G., Lugar, P.L., Rizzieri, D.A., Lagoo, A.S., Bernal-Mizrachi, L., Mann, K.P., Flowers, C.R., Naresh, K.N., Evens, A.M., Chadburn, A., Gordon, L.I., Czader, M.B., Gill, J.I., Hsi, E.D., Greenough, A., Moffitt, A.B., McKinney, M., Banerjee, A., Grubor, V., Levy, S., Dunson, D.B., Dave, S.S. (2012) The genetic landscape of mutations in Burkitt lymphoma. *Nature Genetics* 44(12): 1321-1325.

MacQueen, J.B. (1967) Some Methods for classification and Analysis of Multivariate Observations. In: LeCam, L. and Neyman, J. (eds.) *Proceedings of the* 5th Berkeley Symposium on Mathematical Statistics and Probability. Berkeley, CA: University of California Press, pp. 281-297.

Murtagh, F. and Contreras, P. (2011) Algorithms for hierarchical clustering: An overview. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery* 2(1): 86-97.

Nik-Zainal, S., Alexandrov, L.B., Wedge, D.C., Van Loo, P., Greenman, C.D., Raine, K., Jones, D., Hinton, J., Marshall, J., Stebbings, L.A., Menzies, A., Martin, S., Leung, K., Chen, L., Leroy, C., Ramakrishna, M., Rance, R., Lau, K.W., Mudie, L.J., Varela, I., McBride, D.J., Bignell, G.R., Cooke, S.L., Shlien, A., Gamble, J., Whitmore, I., Maddison, M., Tarpey, P.S., Davies, H.R., Papaemmanuil, E., Stephens, P.J., McLaren, S., Butler, A.P., Teague, J.W., Jönsson, G., Garber, J.E., Silver, D., Miron, P., Fatima, A., Boyault, S., Langerød, A., Tutt, A., Martens, J.W., Aparicio, S.A., Borg, Å., Salomon, A.V., Thomas, G., Børresen-Dale, A.L., Richardson, A.L., Neuberger, M.S., Futreal, P.A., Campbell, P.J., Stratton, M.R.; Breast Cancer Working Group of the International Cancer Genome Consortium. (2012) Mutational processes molding the genomes of 21 breast cancers. *Cell* 149(5): 979-993.

Nik-Zainal, S., Wedge, D.C., Alexandrov, L.B., Petljak, M., Butler, A.P., Bolli, N., Davies, H.R., Knappskog, S., Martin, S., Papaemmanuil, E., Ramakrishna, M., Shlien, A., Simonic, I., Xue, Y., Tyler-Smith, C., Campbell, P.J., Stratton, M.R. (2014) Association of a germline copy number polymorphism of APOBEC3A and APOBEC3B with burden of putative APOBEC-dependent mutations in breast cancer. *Nature Genetics* 46(5): 487-491.

Paatero, P. and Tapper, U. (1994) Positive matrix factorization: A non-negative factor model with optimal utilization of error. *Environmetrics* 5(1): 111-126.

Patch, A.M., Christie, E.L., Etemadmoghadam, D., Garsed, D.W., George, J., Fereday, S., Nones, K., Cowin, P., Alsop, K., Bailey, P.J., Kassahn, K.S., Newell, F., Quinn, M.C., Kazakoff, S., Quek, K., Wilhelm-Benartzi, C., Curry, E., Leong, H.S.; Australian Ovarian Cancer Study Group, Hamilton, A., Mileshkin, L., Au-Yeung, G., Kennedy, C., Hung, J., Chiew, Y.E., Harnett, P., Friedlander, M., Quinn, M., Pyman, J., Cordner, S., O'Brien, P., Leditschke, J., Young, G., Strachan, K., Waring, P., Azar, W., Mitchell, C., Traficante, N., Hendley, J., Thorne, H., Shackleton, M., Miller, D.K., Arnau, G.M., Tothill, R.W., Holloway, T.P., Semple, T., Harliwong, I., Nourse, C., Nourbakhsh, E., Manning, S., Idrisoglu, S., Bruxner, T.J., Christ, A.N., Poudel, B., Holmes, O., Anderson, M., Leonard, C., Lonie, A., Hall, N., Wood, S., Taylor, D.F., Xu, Q., Fink, J.L., Waddell, N., Drapkin, R., Stronach, E., Gabra, H., Brown, R., Jewell, A., Nagaraj, S.H., Markham, E., Wilson, P.J., Ellul, J., McNally, O., Doyle, M.A., Vedururu, R., Stewart, C., Lengyel, E., Pearson, J.V., Waddell, N., deFazio, A., Grimmond, S.M., Bowtell, D.D. (2015) Whole-genome characterization of chemoresistant ovarian cancer. Nature 521(7553): 489-494.

Pelleg, D. and Moore, A.W. (2000) X-means: Extending K-means with Efficient Estimation of the Number of Clusters. In: Langley, P. (ed.) *Proceedings of the 17th International Conference on Machine Learning*. San Francisco, CA: Morgan Kaufman: pp. 727-734.

Pettersen, H.S., Galashevskaya, A., Doseth, B., Sousa, M.M., Sarno, A., Visnes, T., Aas, P.A., Liabakk, N.B., Slupphaug, G., Sætrom, P., Kavli, B., Krokan, H.E. (2015) AID expression in B-cell lymphomas causes accumulation of genomic uracil and a distinct AID mutational signature. *DNA Repair* 25: 60-71.

Poon, S., McPherson, J., Tan, P., Teh, B. and Rozen, S. (2014) Mutation signatures of carcinogen exposure: genome-wide detection and new opportunities for cancer prevention. *Genome Medicine* 6(3): 24. Puente, X.S., Pinyol, M., Quesada, V., Conde, L., Ordóñez, G.R., Villamor, N., Escaramis, G., Jares, P., Beà, S., González-Díaz, M., Bassaganyas, L., Baumann, T., Juan, M., López-Guerra, M., Colomer, D., Tubío, J.M., López, C., Navarro, A., Tornador, C., Aymerich, M., Rozman, M., Hernández, J.M., Puente, D.A., Freije, J.M., Velasco, G., Gutiérrez-Fernández, A., Costa, D., Carrió, A., Guijarro, S., Enjuanes, A., Hernández, L., Yagüe, J., Nicolás, P., Romeo-Casabona, C.M., Himmelbauer, H., Castillo, E., Dohm, J.C., de Sanjosé, S., Piris, M.A., de Alava, E., San Miguel, J., Royo, R., Gelpí, J.L., Torrents, D., Orozco, M., Pisano, D.G., Valencia, A., Guigó, R., Bayés, M., Heath, S., Gut, M., Klatt, P., Marshall, J., Raine, K., Stebbings, L.A., Futreal, P.A., Stratton, M.R., Campbell, P.J., Gut, I., López-Guillermo, A., Estivill, X., Montserrat, E., López-Otín, C., Campo, E. (2011) Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature* 475(7354): 101-105.

Puente, X.S., Beà, S., Valdés-Mas, R., Villamor, N., Gutiérrez-Abril, J., Martín-Subero, J.I., Munar, M., Rubio-Pérez, C., Jares, P., Aymerich, M., Baumann, T., Beekman, R., Belver, L., Carrio, A., Castellano, G., Clot, G., Colado, E., Colomer, D., Costa, D., Delgado, J., Enjuanes, A., Estivill, X., Ferrando, A.A., Gelpí, J.L., González, B., González, S., González, M., Gut, M., Hernández-Rivas, J.M., López-Guerra, M., Martín-García, D., Navarro, A., Nicolás, P., Orozco, M., Payer, Á.R., Pinyol, M., Pisano, D.G., Puente, D.A., Queirós, A.C., Quesada, V., Romeo-Casabona, C.M., Royo, C., Royo, R., Rozman, M., Russiñol, N., Salaverría, I., Stamatopoulos, K., Stunnenberg, H.G., Tamborero, D., Terol, M.J., Valencia, A., López-Bigas, N., Torrents, D., Gut, I., López-Guillermo, A., López-Otín, C., Campo, E. (2015) Non-coding recurrent mutations in chronic lymphocytic leukaemia. *Nature* 526(7574): 519-524.

Qian, J., Wang, Q., Dose, M., Pruett, N., Kieffer-Kwon, K.R., Resch, W., Liang, G., Tang, Z., Mathé, E., Benner, C., Dubois, W., Nelson, S., Vian, L., Oliveira, T.Y., Jankovic, M., Hakim, O., Gazumyan, A., Pavri, R., Awasthi, P., Song, B., Liu, G., Chen, L., Zhu, S., Feigenbaum, L., Staudt, L., Murre, C., Ruan, Y., Robbiani, D.F., Pan-Hammarström, Q., Nussenzweig, M.C., Casellas, R. (2014) B Cell Super-Enhancers and Regulatory Clusters Recruit AID Tumorigenic Activity. *Cell* 159(7): 1524-1537.

Roberts, S.A., Sterling, J., Thompson, C., Harris, S., Mav, D., Shah, R., Klimczak, L.J., Kryukov, G.V., Malc, E., Mieczkowski, P.A., Resnick, M.A., Gordenin, D.A. (2012) Clustered mutations in yeast and in human cancers can arise from damaged long single-strand DNA regions. *Molecular Cell* 46(4): 424-435.

Roberts, S.A., Lawrence, M.S., Klimczak, L.J., Grimm, S.A., Fargo, D., Stojanov, P., Kiezun, A., Kryukov, G.V., Carter, S.L., Saksena, G., Harris, S.,
Shah, R.R., Resnick, M.A., Getz, G. and Gordenin, D.A. (2013) An APOBEC cytidine deaminase mutagenesis pattern is widespread in human cancers. *Nature Genetics* 45(9): 970-976.

Roberts, S.A. and Gordenin, D.A. (2014a) Clustered Mutations in Human Cancer. In: *eLS (Genetics & Disease)*. Chichester, UK: John Wiley & Sons, Ltd.

Roberts S.A and Gordenin D.A. (2014b) Clustered and genome-wide transient mutagenesis in human cancers: Hypermutation without permanent mutators or loss of fitness. *BioEsseys* 36(4): 382-393.

Roberts S.A and Gordenin D.A. (2014c) Hypermutation in human cancer genomes: footprints and mechanisms. *Nature Reviews Cancer* 14(12): 786-800.

Rousseeuw, P.J. (1987) Silhouettes: a Graphical Aid to the Interpretation and Validation of Cluster Analysis. *Journal of Computational and Applied Mathematics* 20(1): 53-65.

Roy, O. and Vetterli, M. (2007) The effective rank: A measure of effective dimensionality. In: *European Signal Processing Conference (EUSIPCO)*. Poznań, Poland (September 3-7), pp. 606-610.

Scelo, G., Riazalhosseini, Y., Greger, L., Letourneau, L., Gonzàlez-Porta, M., Wozniak, M.B., Bourgey, M., Harnden, P., Egevad, L., Jackson, S.M., Karimzadeh, M., Arseneault, M., Lepage, P., How-Kit, A., Daunay, A., Renault, V., Blanché, H., Tubacher, E., Sehmoun, J., Viksna, J., Celms, E., Opmanis, M., Zarins, A., Vasudev, N.S., Seywright, M., Abedi-Ardekani, B., Carreira, C., Selby, P.J., Cartledge, J.J., Byrnes, G., Zavadil, J., Su, J., Holcatova, I., Brisuda, A., Zaridze, D., Moukeria, A., Foretova, L., Navratilova, M., Mates, D., Jinga, V., Artemov, A., Nedoluzhko, A., Mazur, A., Rastorguev, S., Boulygina, E., Heath, S., Gut, M., Bihoreau, M.T., Lechner, D., Foglio, M., Gut, I.G., Skryabin, K., Prokhortchouk, E., Cambon-Thomsen, A., Rung, J., Bourque, G., Brennan, P., Tost, J., Banks, R.E., Brazma, A., Lathrop, G.M. (2014) Variation in genomic landscape of clear cell renal cell carcinoma across Europe. Nature Communications 5: 5135.

Sibson, R. (1973) SLINK: an optimally efficient algorithm for the single-link cluster method. *The Computer Journal (British Computer Society)* 16(1): 30-34.

Sima, J. and Gilbert, D.M. (2014) Complex correlations: replication timing and mutational landscapes during cancer and genome evolution. *Current Opinion in Genetics & Developmen* 25: 93-100.

Steinbach, M., Karypis, G. and Kumar, V. (2000) A comparison of document clustering techniques. *KDD Workshop on Text Mining* 400(1): 525-526.

Steinhaus, H. (1957) Sur la division des corps matériels en parties. *Bull. Acad. Polon. Sci.* 4(12): 801-804.

Sugar, C.A. and James, G.M. (2003) Finding the number of clusters in a data set: An information theoretic approach. *Journal of the American Statistical Association* 98(463): 750-763.

Sung, W.K., Zheng, H., Li, S., Chen, R., Liu, X., Li, Y., Lee, N.P., Lee, W.H.,
Ariyaratne, P.N., Tennakoon, C., Mulawadi, F.H., Wong, K.F., Liu, A.M.,
Poon, R.T., Fan, S.T., Chan, K.L., Gong, Z., Hu, Y., Lin, Z., Wang, G., Zhang,
Q., Barber, T.D., Chou, W.C., Aggarwal, A., Hao, K., Zhou, W., Zhang, C.,
Hardwick, J., Buser, C., Xu, J., Kan, Z., Dai, H., Mao, M., Reinhard, C.,
Wang, J., Luk, J.M. (2012) Genome-wide survey of recurrent HBV integration
in hepatocellular carcinoma. *Nature Genetics* 44(7): 765-769.

Taylor, B.J.M., Nik-Zainal, S., Wu, Y.L., Stebbings, L.A., Raine, K., Campbell, P.J., Rada, C., Stratton, M.R., Neuberger, M.S. (2013) DNA deaminases induce break-associated mutation showers with implication of APOBEC3B and 3A in breast cancer kataegis. *eLife* 2: e00534.

Tirode, F., Surdez, D., Ma, X., Parker, M., Le Deley, M.C., Bahrami, A., Zhang, Z., Lapouble, E., Grossetête-Lalami, S., Rusch, M., Reynaud, S., Rio-Frio, T., Hedlund, E., Wu, G., Chen, X., Pierron, G., Oberlin, O., Zaidi, S., Lemmon, G., Gupta, P., Vadodaria, B., Easton, J., Gut, M., Ding, L., Mardis, E.R., Wilson, R.K., Shurtleff, S., Laurence, V., Michon, J., Marec-Bérard, P., Gut, I., Downing, J., Dyer, M., Zhang, J., Delattre, O.; St. Jude Children's Research Hospital – Washington University Pediatric Cancer Genome Project and the International Cancer Genome Consortium. (2014) Genomic Landscape of Ewing Sarcoma defines an aggressive subtype with co-association of STAG2 and TP53 mutations. *Cancer Discovery* 4(11): 1342-1353.

Waddell, N., Pajic, M., Patch, A.M., Chang, D.K., Kassahn, K.S., Bailey, P., Johns, A.L., Miller, D., Nones, K., Quek, K., Quinn, M.C., Robertson, A.J., Fadlullah, M.Z., Bruxner, T.J., Christ, A.N., Harliwong, I., Idrisoglu, S., Manning, S., Nourse, C., Nourbakhsh, E., Wani, S., Wilson, P.J., Markham, E., Cloonan, N., Anderson, M.J., Fink, J.L., Holmes, O., Kazakoff, S.H., Leonard, C., Newell, F., Poudel, B., Song, S., Taylor, D., Waddell, N., Wood, S., Xu, Q., Wu, J., Pinese, M., Cowley, M.J., Lee, H.C., Jones, M.D., Nagrial, A.M., Humphris, J., Chantrill, L.A., Chin, V., Steinmann, A.M., Mawson, A., Humphrey, E.S., Colvin, E.K., Chou, A., Scarlett, C.J., Pinho, A.V., Giry-Laterriere, M., Rooman, I., Samra, J.S., Kench, J.G., Pettitt, J.A., Merrett, N.D., Toon, C., Epari, K., Nguyen, N.Q., Barbour, A., Zeps, N., Jamieson, N.B., Graham, J.S., Niclou, S.P., Bjerkvig, R., Grützmann, R., Aust, D., Hruban, R.H., Maitra, A., Iacobuzio-Donahue, C.A., Wolfgang, C.L., Morgan, R.A., Lawlor, R.T., Corbo, V., Bassi, C., Falconi, M., Zamboni, G., Tor-

tora, G., Tempero, M.A.; Australian Pancreatic Cancer Genome Initiative, Gill, A.J., Eshleman, J.R., Pilarsky, C., Scarpa, A., Musgrove, E.A., Pearson, J.V., Biankin, A.V., Grimmond, S.M. (2015) Whole genomes redefine the mutational landscape of pancreatic cancer. *Nature* 518(7540): 495-501.

Wang, K., Yuen, S.T., Xu, J., Lee, S.P., Yan, H.H., Shi, S.T., Siu, H.C., Deng, S., Chu, K.M., Law, S., Chan, K.H., Chan, A.S., Tsui, W.Y., Ho, S.L., Chan, A.K., Man, J.L., Foglizzo, V., Ng, M.K., Chan, A.S., Ching, Y.P., Cheng, G.H., Xie, T., Fernandez, J., Li, V.S., Clevers, H., Rejto, P.A., Mao, M., Leung, S.Y. (2014) Whole-genome sequencing and comprehensive molecular profiling identify new driver mutations in gastric cancer. *Nature Genetics* 46(6): 573-582.

Xuan, D., Li, G., Cai, Q., Deming-Halverson, S., Shrubsole, M.J., Shu, X.O., Kelley, M.C., Zheng, W., Long, J. (2013) APOBEC3 deletion polymorphism is associated with breast cancer risk among women of European ancestry. *Carcinogenesis* 34(10): 2240-2243.

Yang, W., Gibson, J.D. and He, T. (2005) Coefficient rate and lossy source coding. *IEEE Transactions on Information Theory* 51(1): 381-386.

Zhang, J., Wu, G., Miller, C.P., Tatevossian, R.G., Dalton, J.D., Tang, B., Orisme, W., Punchihewa, C., Parker, M., Qaddoumi, I., Boop, F.A., Lu, C., Kandoth, C., Ding, L., Lee, R., Huether, R., Chen, X., Hedlund, E., Nagahawatte, P., Rusch, M., Boggs, K., Cheng, J., Becksfort, J., Ma, J., Song, G., Li, Y., Wei, L., Wang, J., Shurtleff, S., Easton, J., Zhao, D., Fulton, R.S., Fulton, L.L., Dooling, D.J., Vadodaria, B., Mulder, H.L., Tang, C., Ochoa, K., Mullighan, C.G., Gajjar, A., Kriwacki, R., Sheer, D., Gilbertson, R.J., Mardis, E.R., Wilson, R.K., Downing, J.R., Baker, S.J., Ellison, D.W.; St. Jude Children's Research Hospital-Washington University Pediatric Cancer Genome Project. (2013) Whole-genome sequencing identifies genetic alterations in pediatric low-grade gliomas. *Nature Genetics* 45(6): 602-612.

Cancer Type	Total Counts	# of Samples	Source
B Cell Lymphoma	43626	24	A1-2
Bone Cancer	36374	98	B1
Brain Lower Grade Glioma	3572	101	A1, C1
Breast Cancer	254381	119	A1, D1
Chronic Lymphocytic Leukemia	19489	134	A1, E1-2
Esophageal Cancer	1064	17	F1
Gastric Cancer	1996615	100	G1
Liver Cancer	3017487	389	H1-2
Lung Cancer	449527	24	I1
Medulloblastoma	44689	100	J1
Ovarian Cancer	668918	84	K1
Pancreatic Cancer	5087	100	L1
Prostate Cancer	29142	5	M1
Renal Cell Carcinoma	483329	94	N1
All Cancer Types	7053300	1389	Above

Table 1: Genome data summary. All Brain Lower Glioma samples are Pilocytic Astrocytoma samples. See Subsection 3.1 for the data source definitions.

Table 2: Occurrence counts for the first 48 mutation categories for the genome data summarized in Table 1 aggregated by cancer types. X1 = B Cell Lymphoma, X2 = B one Cancer, X3 = B rain Lower Grade Glioma, X4 = B reast Cancer, X5 = Ch ronic Lymphocytic Leukemia, X6 = E sophageal Cancer, X7 = G astric Cancer, X8 = L iver Cancer, X9 = Lung Cancer, X10 = M edulloblastoma, X11 = O varian Cancer, X12 = P ancreatic Cancer, X13 = P rostate Cancer, X14 = R enal Cell Carcinoma. The mutations are encoded as follows: XYZW = Y > W: XYZ.

Mutation	X1	X2	X3	X4	X5	X6	X7	X8	X9	X10	X11	X12	X13	X14
			-		-	-	-	-	-	-			-	
ACAA	466	716	59	3024	286	11	24884	51929	15865	801	13831	36	577	10734
ACCA	355	528	41	2600	249	3	14361	15227	9217	681	11192	42	422	4952
ACGA	43	112	6	446	55	4	2253	51156	3567	115	1804	9	59	1231
ACTA	309	577	38	2201	195	12	15833	33385	10038	557	10508	20	380	5573
CCAA	420	538	29	2814	238	11	21980	9507	19569	579	10943	59	459	7899
CCCA	245	361	30	2149	147	6	19624	26869	16889	424	9720	45	336	6037
CCGA	77	75	22	470	35	7	4034	8695	4995	105	1644	29	67	1069
CCTA	418	502	26	2380	211	10	43418	2967	15930	544	10037	80	388	6825
GCAA	452	624	41	2144	244	16	26007	42508	11084	532	6835	33	419	4795
GCCA	234	346	18	1677	149	6	10594	32252	9087	337	6423	33	247	3563
GCGA	67	66	7	375	29	5	2649	17426	3607	94	1295	17	67	853
GCTA	351	463	13	1684	197	11	21205	33670	7502	345	6543	30	332	3637
TCAA	584	628	38	5873	233	11	24288	33919	13358	766	9816	54	582	8973
TCCA	313	472	36	3735	145	19	17372	139576	13140	567	9697	53	479	7388
TCGA	66	57	7	543	28	4	3168	11569	2385	128	1334	17	87	1097
TCTA	916	849	54	4938	376	25	43364	20943	14028	1111	12202	56	817	8611
ACAG	232	260	19	2124	162	1	10057	42216	4369	445	8653	18	197	4551
ACCG	143	148	11	1307	92	4	4818	6202	2088	256	4506	21	188	2687
ACGG	48	82	10	641	32	2	1606	27957	953	121	2172	5	46	837
ACTG	290	240	25	2278	113	8	8670	90212	3091	394	8460	23	237	4457
CCAG	153	160	13	1726	78	7	4942	8975	3233	175	5559	30	125	2474
CCCG	131	154	14	1391	75	2	4117	27928	2902	163	4857	18	142	2601
CCGG	31	65	14	597	20	0	1176	87975	1210	122	1769	11	55	532
CCTG	213	227	19	2425	92	11	6840	14272	3348	238	7943	28	204	3846
GCAG	106	120	8	1128	52	2	3844	53180	2546	196	3621	12	95	1847
GCCG	147	148	7	1142	80	3	4174	11783	2406	160	3363	16	112	1630
GCGG	25	42	2	525	33	1	776	52658	1113	52	918	5	27	423
GCTG	225	146	9	1650	81	5	5116	43761	2187	177	5418	14	135	2045
TCAG	391	266	22	16099	79	46	12809	12172	8882	265	13033	49	359	3968
TCCG	279	185	18	4896	95	14	7208	41020	4566	244	7977	30	264	4441
TCGG	30	52	11	794	13	2	1294	9554	992	64	1423	11	33	553
TCTG	660	444	44	21847	202	53	21720	3519	10841	449	20292	75	563	7668
ACAT	950	931	97	3557	440	14	46364	28569	6530	1137	13367	66	542	8872
ACCT	482	482	49	1875	265	14	16724	47067	3418	590	6543	100	285	4786
ACGT	1085	2373	289	4978	792	70	78681	17833	3830	3694	17475	585	1603	9034
ACTT	603	628	57	2570	263	11	21545	63195	4179	827	10578	54	380	6567
CCAT	729	542	74	3344	294	16	19611	23743	7430	801	8689	79	428	7106
CCCT	607	545	78	2443	337	10	18553	52051	6155	810	7554	70	325	6038
CCGT	845	1095	180	3489	483	70	53496	7282	4362	1983	11157	498	888	4824
CCTT	784	774	124	3468	380	13	22137	29223	7932	854	11714	69	430	8606
GCAT	615	531	65	2815	285	16	31548	50399	4166	711	8492	107	420	6069
GCCT	583	484	64	2256	295	14	34372	10858	3715	825	7784	120	402	5474
GCGT	930	1382	152	3870	546	60	75814	35864	3544	2428	13447	629	1294	5232
GCTT	660	585	78	2254	316	15	26554	49322	3836	741	8329	93	378	5765
TCAT	1531	761	86	25251	268	69	30161	11611	13414	780	13880	131	681	9723
TCCT	1172	685	73	7268	319	25	23799	31779	7136	859	10287	112	544	8833
TCGT	628	903	127	4510	281	58	33759	53275	2998	1553	8055	300	708	3772
TCTT	1466	813	57	13615	313	43	29570	21922	11432	859	12692	79	662	8933

Mutation	X1	X2	X3	X4	X5	X6	X7	X8	X9	X10	X11	X12	X13	X14
ATAA	385	382	22	1375	248	2	9613	35715	3957	315	5776	7	313	9726
ATCA	241	274	13	1388	191	6	9419	7570	1917	235	4439	18	214	2973
ATGA	232	344	21	1459	211	4	7666	48117	3962	228	5983	18	197	7703
ATTA	580	516	29	2262	301	5	30067	32078	2928	590	9131	16	523	5833
CTAA	185	217	9	1009	98	3	5551	9978	4331	151	4473	10	175	12963
CTCA	211	256	17	1535	118	6	14584	41504	3430	224	6343	21	184	6465
CTGA	219	227	26	1406	146	12	8654	8508	7905	210	6473	28	207	14356
CTTA	379	408	27	1716	162	6	24273	2400	4255	303	8578	24	221	8077
GTAA	180	162	10	797	82	1	4092	36890	3147	104	3254	5	138	5769
GTCA	174	147	8	791	80	1	5919	25736	1399	113	3026	10	79	1954
GTGA	156	145	9	948	102	10	6717	9623	4021	122	4233	8	135	3924
GTTA	181	272	14	1168	129	3	10065	43279	2112	200	5741	10	156	2579
TTAA	576	454	29	2192	256	2	20211	13356	4086	510	6936	16	499	11629
TTCA	158	202	17	1095	110	3	7296	75515	1766	222	4859	12	142	4966
TTGA	179	156	4	774	144	3	5147	5013	2912	140	3493	10	129	7897
TTTA	434	467	37	2298	289	3	27521	12174	3117	481	10786	8	417	10311
ATAC	910	792	84	3342	519	10	34145	35102	7884	865	14628	36	552	11060
ATCC	382	255	32	1439	192	7	14365	4444	1952	354	6071	22	206	2821
ATGC	572	526	51	2354	329	14	28630	16789	5325	534	10017	48	331	6561
ATTC	792	734	94	3340	457	11	22018	48320	4344	769	13211	38	503	6714
CTAC	456	283	30	1378	187	1	21942	9143	4198	301	5972	21	202	4855
CTCC	427	295	31	1670	197	5	31474	13443	2361	331	7202	48	163	3092
CTGC	531	328	29	1659	222	6	38742	59589	3845	360	6553	60	206	4024
CTTC	795	328	48	1828	316	12	73708	10381	3566	383	8088	48	239	4982
GTAC	452	378	30	1533	215	4	25980	46065	3605	462	6804	48	345	4411
GTCC	404	332	25	981	176	10	15268	17372	1619	299	4123	20	195	1823
GTGC	370	284	33	1233	189	8	24550	54718	2912	332	5851	48	211	2962
GTTC	511	447	46	1680	269	7	22803	46663	2547	532	7513	36	367	3575
TTAC	541	428	48	1819	297	2	22540	16834	3732	497	7741	25	372	6232
TTCC	606	383	44	1610	211	2	25505	42903	1979	473	7086	24	280	3847
TTGC	306	242	22	1079	185	ā	17328	8016	2277	323	4419	24	210	2892
TTTC	818	437	60	2187	415	4	48109	2717	3140	617	10866	28	427	13766
ATAG	462	133	à	862	229	2	5956	20568	1055	164	3114	5	158	2747
ATCG	134	69	š	451	51	õ	5214	50363	476	92	1335	10	93	1949
ATGG	196	136	11	941	76	3	5505	35331	1134	165	3935	5	116	2726
ATTC	580	155	17	1030	166	2	31252	58405	1010	177	3455	11	221	4941
CTAC	416	110	1	601	134	3	5035	20104	779	85	1032	3	87	1381
CTCC	210	01	16	750	71	4	14376	86003	605	96	2754	15	96	1628
CTCC	213	137	7	1165	140	4	19511	13036	1684	142	1188	10	142	2080
CTGG	1720	202	21	1560	244	ч Б	142206	27101	1901	1942	4400	20	142	2080
CTAC	224	202	21	617	70	0	142200	45724	746	07	4424	30	111	2944
GIAG	122	64 E2	3	667	10	0	2002	40704	740 549	01 E 2	1792	3 10	45	690
GIUG	102	100	4	2080	43	0	5004	210333	042	171	1290	10	41	1212
GIGG	100	112	14	2989	140	2	27454	46081	2074	171	3407	10	100	1313
TTAC	1090	110	14	1094	142	3	3/434	40901	100	107	3439	13	122	1423
TIAG	004	199	19	1021	260	1	(0/4	11090	1230	187	3327	14	108	2799
TTCG	107	90	13	120	19	2 7	0310	32124 69446	1602	161	2007	12	120	3312
TIGG	200	132	10	0144	121	1	11452	02440	1093	101	4480	8	142	2974
TTTG	1349	296	43	2144	403	4	77262	28801	2361	379	7679	39	353	11415

Table 3: Table 2 continued: occurrence counts (aggregated by cancer types) for the next 48 mutation categories.

Table 4: Top 10 clusterings by occurrence counts (second column) in 150,000 runs (performed as 15 consecutive batches of 10,000 runs in each batch). Each run is based on 1,000 samplings (i.e., num.try = 1000 in the R function qrm.stat.ind.class(); also, the target number of clusters is k = 7; see Appendix A for details). The columns "Cl-1" through "Cl-7" give the numbers of mutations in each cluster (the total number of mutations in each clustering is 96). The entries "—" correspond to clusterings with fewer than 7 clusters. Note that Clustering-C and Clustering-H have the same numbers of mutations in their 7 clusters; however, these two clusterings are different, i.e., equally-sized clusters contain different mutations. While there was slight variability in the placement (by occurrence counts) of the top 10 clustering-J invariably were the top 10 in each batch. The weights are given in Tables 5 through 12 (for Clustering-A through Clustering-D) and 13 and 14 (for Clustering-A).

Count	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
12085	8	8	10	15	16	18	21
10962	7	8	8	12	15	17	29
10788	8	8	11	15	16	17	21
10328	8	8	15	16	18	31	_
6499	7	8	8	12	15	18	28
5451	8	8	15	17	17	31	_
5421	8	8	10	15	17	17	21
5302	8	8	11	15	16	17	21
4602	8	8	10	15	21	34	_
3698	8	8	15	31	34	_	_
	Count 12085 10962 10788 10328 6499 5451 5421 5302 4602 3698	CountCl-11208581096271078881032886499754518530284602836988	CountCl-1Cl-21208588109627810788881032888649978545188542188530288460288369888	CountCl-1Cl-2Cl-31208588101096278810788881110328881564997885451881554218810530288114602881036988815	CountCl-1Cl-2Cl-3Cl-4120858810151096278812107888811151032888151664997881254518815175421881015530288111546028810153698881531	CountCl-1Cl-2Cl-3Cl-4Cl-512085881015161096278812151078888111516103288815161864997881215545188151717542188101516460288101521369888153134	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

Table 5: Weights (in the units of 1%, rounded to 2 digits) for the first 48 mutation categories (this Table 5 is continued in Table 6 with the next 48 mutation categories) for the 7 clusters in Clustering-A (see Table 4) based on unnormalized (columns 2-8) and normalized (columns 9-15) regressions (see Subsection 2.6 for details). Each cluster is defined as containing the mutations with nonzero weights. For instance, cluster Cl-2 contains 8 mutations GCGA, TCGA, ACGG, GCCG, GCGG, TCGG, GTCA, GTCG. In each cluster the weights are normalized to add up to 100% (up to 2 digits due to the aforesaid rounding). In Tables 5 through 12 "weights based on unnormalized regressions" are given by (13), (14) and (15), while "weights based on normalized regressions" are given by (17), (14) and (16), i.e., the exposures are calculated based on arithmetic averages (see Subsection 2.6 for details).

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
	-	-		-				-	-		-			
ACAA	0.00	0.00	0.00	6.55	0.00	0.00	0.00	0.00	0.00	0.00	6.55	0.00	0.00	0.00
ACCA	0.00	0.00	0.00	0.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00	6.08	0.00	0.00
ACGA	0.00	0.00	0.00	0.00	0.00	0.00	4.06	0.00	0.00	0.00	0.00	0.00	0.00	4.00
ACTA	0.00	0.00	0.00	0.00	6.16	0.00	0.00	0.00	0.00	0.00	0.00	6.38	0.00	0.00
CCAA	0.00	0.00	0.00	0.00	7.91	0.00	0.00	0.00	0.00	0.00	0.00	8.10	0.00	0.00
CCCA	0.00	0.00	0.00	0.00	6.46	0.00	0.00	0.00	0.00	0.00	0.00	6.68	0.00	0.00
CCGA	0.00	0.00	7.21	0.00	0.00	0.00	0.00	0.00	0.00	7.23	0.00	0.00	0.00	0.00
CCTA	0.00	0.00	0.00	0.00	0.00	6.75	0.00	0.00	0.00	0.00	0.00	0.00	6.79	0.00
GCAA	4.05	0.00	0.00	0.00	0.00	0.00	0.00	4.65	0.00	0.00	0.00	0.00	0.00	0.00
GCCA	0.00	0.00	0.00	0.00	4.56	0.00	0.00	0.00	0.00	0.00	0.00	4.73	0.00	0.00
GCGA	0.00	13.81	0.00	0.00	0.00	0.00	0.00	0.00	13.89	0.00	0.00	0.00	0.00	0.00
GCTA	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00	0.00	5.20	0.00	0.00
TCAA	0.00	0.00	0.00	6.26	0.00	0.00	0.00	0.00	0.00	0.00	6.21	0.00	0.00	0.00
TCCA	0.00	0.00	0.00	0.00	8.94	0.00	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00
TCGA	0.00	11.87	0.00	0.00	0.00	0.00	0.00	0.00	12.24	0.00	0.00	0.00	0.00	0.00
TCTA	0.00	0.00	0.00	8.05	0.00	0.00	0.00	0.00	0.00	0.00	8.00	0.00	0.00	0.00
ACAG	0.00	0.00	0.00	0.00	3.96	0.00	0.00	0.00	0.00	0.00	0.00	4.18	0.00	0.00
ACCG	0.00	0.00	8.07	0.00	0.00	0.00	0.00	0.00	0.00	8.17	0.00	0.00	0.00	0.00
ACGG	0.00	12.62	0.00	0.00	0.00	0.00	0.00	0.00	12.22	0.00	0.00	0.00	0.00	0.00
ACTG	0.00	0.00	0.00	0.00	4.77	0.00	0.00	0.00	0.00	0.00	0.00	5.03	0.00	0.00
CCAG	0.00	0.00	9.26	0.00	0.00	0.00	0.00	0.00	0.00	9.35	0.00	0.00	0.00	0.00
CCCG	0.00	0.00	0.00	0.00	0.00	0.00	3.91	0.00	0.00	0.00	0.00	0.00	0.00	4.02
CCGG	0.00	0.00	0.00	0.00	0.00	0.00	5.37	0.00	0.00	0.00	0.00	0.00	0.00	5.12
CCTG	0.00	0.00	12.46	0.00	0.00	0.00	0.00	0.00	0.00	12.58	0.00	0.00	0.00	0.00
GCAG	0.00	0.00	0.00	0.00	0.00	0.00	4.61	0.00	0.00	0.00	0.00	0.00	0.00	4.57
GCCG	0.00	14.79	0.00	0.00	0.00	0.00	0.00	0.00	15.62	0.00	0.00	0.00	0.00	0.00
GCGG	0.00	15.50	0.00	0.00	0.00	0.00	0.00	0.00	13.92	0.00	0.00	0.00	0.00	0.00
GCTG	0.00	0.00	0.00	0.00	0.00	0.00	4.86	0.00	0.00	0.00	0.00	0.00	0.00	4.92
TCAG	0.00	0.00	0.00	0.00	10.31	0.00	0.00	0.00	0.00	0.00	0.00	9.03	0.00	0.00
TCCG	0.00	0.00	0.00	0.00	5.10	0.00	0.00	0.00	0.00	0.00	0.00	4.95	0.00	0.00
TCGG	0.00	8.40	0.00	0.00	0.00	0.00	0.00	0.00	8.65	0.00	0.00	0.00	0.00	0.00
TCTG	0.00	0.00	0.00	0.00	14.10	0.00	0.00	0.00	0.00	0.00	0.00	12.53	0.00	0.00
ACAT	0.00	0.00	0.00	7.67	0.00	0.00	0.00	0.00	0.00	0.00	7.71	0.00	0.00	0.00
ACCT	4.78	0.00	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00	0.00
ACGT	23.47	0.00	0.00	0.00	0.00	0.00	0.00	23.18	0.00	0.00	0.00	0.00	0.00	0.00
ACTT	0.00	0.00	0.00	5.43	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00	0.00
CCAT	0.00	0.00	0.00	6.02	0.00	0.00	0.00	0.00	0.00	0.00	6.02	0.00	0.00	0.00
CCCT	0.00	0.00	0.00	5.59	0.00	0.00	0.00	0.00	0.00	0.00	5.63	0.00	0.00	0.00
CCGT	17.66	0.00	0.00	0.00	0.00	0.00	0.00	17.12	0.00	0.00	0.00	0.00	0.00	0.00
CCTT	0.00	0.00	0.00	7.01	0.00	0.00	0.00	0.00	0.00	0.00	7.04	0.00	0.00	0.00
GCAI	0.00	0.00	0.00	5.98	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00
GCCT	5.74	0.00	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00	0.00	0.00	0.00
GCGT	20.46	0.00	0.00	0.00	0.00	0.00	0.00	19.80	0.00	0.00	0.00	0.00	0.00	0.00
GUII	0.00	0.00	0.00	0.00	0.00	0.00	0.00	10.00	0.00	0.00	0.93	0.00	0.00	0.00
TCCT	11.42	0.00	0.00	7.91	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00	0.00
TCCT	19.49	0.00	0.00	0.00	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTT	12.42	0.00	0.00	0.00	0.00	0.00	0.00	12.30	0.00	0.00	0.00	0.00	0.00	0.00
1011	0.00	0.00	0.00	9.47	0.00	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00	0.00

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
ATAA	0.00	0.00	0.00	0.00	4.18	0.00	0.00	0.00	0.00	0.00	0.00	4.52	0.00	0.00
ATCA	0.00	0.00	10.00	0.00	0.00	0.00	0.00	0.00	0.00	10.15	0.00	0.00	0.00	0.00
ATGA	0.00	0.00	0.00	0.00	4.02	0.00	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00
ATTA	0.00	0.00	0.00	0.00	0.00	5.54	0.00	0.00	0.00	0.00	0.00	0.00	5.66	0.00
CTAA	0.00	0.00	11.74	0.00	0.00	0.00	0.00	0.00	0.00	11.16	0.00	0.00	0.00	0.00
CTCA	0.00	0.00	0.00	0.00	3.79	0.00	0.00	0.00	0.00	0.00	0.00	3.98	0.00	0.00
CTGA	0.00	0.00	0.00	0.00	4.88	0.00	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00
CTTA	0.00	0.00	0.00	0.00	0.00	4.28	0.00	0.00	0.00	0.00	0.00	0.00	4.33	0.00
GTAA	0.00	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00	0.00	0.00	0.00	0.00	4.35
GTCA	0.00	15.20	0.00	0.00	0.00	0.00	0.00	0.00	15.36	0.00	0.00	0.00	0.00	0.00
GTGA	0.00	0.00	9.28	0.00	0.00	0.00	0.00	0.00	0.00	9.21	0.00	0.00	0.00	0.00
GTTA	0.00	0.00	0.00	0.00	0.00	0.00	5.13	0.00	0.00	0.00	0.00	0.00	0.00	5.19
TTAA	0.00	0.00	0.00	0.00	0.00	5.13	0.00	0.00	0.00	0.00	0.00	0.00	5.26	0.00
TTCA	0.00	0.00	0.00	0.00	0.00	0.00	6.64	0.00	0.00	0.00	0.00	0.00	0.00	6.58
TTGA	0.00	0.00	8.84	0.00	0.00	0.00	0.00	0.00	0.00	8.55	0.00	0.00	0.00	0.00
TTTA	0.00	0.00	0.00	0.00	0.00	5.27	0.00	0.00	0.00	0.00	0.00	0.00	5.38	0.00
ATAC	0.00	0.00	0.00	7.03	0.00	0.00	0.00	0.00	0.00	0.00	7.06	0.00	0.00	0.00
ATCC	0.00	0.00	0.00	0.00	0.00	3.30	0.00	0.00	0.00	0.00	0.00	0.00	3.39	0.00
ATGC	0.00	0.00	0.00	4.97	0.00	0.00	0.00	0.00	0.00	0.00	4.98	0.00	0.00	0.00
ATTC	0.00	0.00	0.00	6.30	0.00	0.00	0.00	0.00	0.00	0.00	6.34	0.00	0.00	0.00
CTAC	0.00	0.00	0.00	0.00	0.00	3.78	0.00	0.00	0.00	0.00	0.00	0.00	3.81	0.00
CTCC	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00	0.00	0.00	0.00	0.00	4.31	0.00
CTGC	0.00	0.00	0.00	0.00	0.00	5.37	0.00	0.00	0.00	0.00	0.00	0.00	5.41	0.00
CTTC	0.00	0.00	0.00	0.00	0.00	7.14	0.00	0.00	0.00	0.00	0.00	0.00	6.92	0.00
GTAC	0.00	0.00	0.00	0.00	0.00	4.84	0.00	0.00	0.00	0.00	0.00	0.00	4.96	0.00
GTCC	0.00	0.00	11.51	0.00	0.00	0.00	0.00	0.00	0.00	11.78	0.00	0.00	0.00	0.00
GTGC	0.00	0.00	0.00	0.00	0.00	4.32	0.00	0.00	0.00	0.00	0.00	0.00	4.43	0.00
GTTC	0.00	0.00	0.00	0.00	0.00	5.05	0.00	0.00	0.00	0.00	0.00	0.00	5.23	0.00
TTAC	0.00	0.00	0.00	0.00	0.00	4.97	0.00	0.00	0.00	0.00	0.00	0.00	5.10	0.00
TTCC	0.00	0.00	0.00	0.00	0.00	4.69	0.00	0.00	0.00	0.00	0.00	0.00	4.79	0.00
TTGC	0.00	0.00	11.62	0.00	0.00	0.00	0.00	0.00	0.00	11.82	0.00	0.00	0.00	0.00
TTTC	0.00	0.00	0.00	0.00	0.00	7.29	0.00	0.00	0.00	0.00	0.00	0.00	7.28	0.00
ATAG	0.00	0.00	0.00	0.00	0.00	0.00	3.98	0.00	0.00	0.00	0.00	0.00	0.00	4.09
ATCG	0.00	0.00	0.00	0.00	0.00	0.00	3.81	0.00	0.00	0.00	0.00	0.00	0.00	3.70
ATGG	0.00	0.00	0.00	0.00	0.00	0.00	3.97	0.00	0.00	0.00	0.00	0.00	0.00	3.99
ATTG	0.00	0.00	0.00	0.00	0.00	0.00	7.13	0.00	0.00	0.00	0.00	0.00	0.00	7.08
CTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.55	0.00	0.00	0.00	0.00	0.00	0.00	3.56
CTCG	0.00	0.00	0.00	0.00	0.00	0.00	6.52	0.00	0.00	0.00	0.00	0.00	0.00	6.31
CTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.67	0.00	0.00	0.00	0.00	0.00	0.00	3.83
CTTG	0.00	0.00	0.00	0.00	0.00	9.67	0.00	0.00	0.00	0.00	0.00	0.00	8.89	0.00
GTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.58	0.00	0.00	0.00	0.00	0.00	0.00	3.49
GTCG	0.00	7.80	0.00	0.00	0.00	0.00	0.00	0.00	8.11	0.00	0.00	0.00	0.00	0.00
GTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.82	0.00	0.00	0.00	0.00	0.00	0.00	3.98
GTTG	0.00	0.00	0.00	0.00	0.00	0.00	7.02	0.00	0.00	0.00	0.00	0.00	0.00	6.97
TTAG	0.00	0.00	0.00	0.00	0.00	0.00	4.24	0.00	0.00	0.00	0.00	0.00	0.00	4.43
TTCG	0.00	0.00	0.00	0.00	0.00	0.00	3.73	0.00	0.00	0.00	0.00	0.00	0.00	3.75
TTGG	0.00	0.00	0.00	0.00	0.00	0.00	6.10	0.00	0.00	0.00	0.00	0.00	0.00	6.06
TTTG	0.00	0.00	0.00	0.00	0.00	8.31	0.00	0.00	0.00	0.00	0.00	0.00	8.05	0.00

Table 6: Table 5 continued: weights for the next 48 mutation categories.

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7
ACAA	0.00	0.00	0.00	0.00	6.55	0.00	0.00	0.00	0.00	0.00	0.00	6.55	0.00	0.00
ACCA	0.00	0.00	0.00	6.66	0.00	0.00	0.00	0.00	0.00	0.00	7.03	0.00	0.00	0.00
ACGA	0.00	0.00	0.00	0.00	0.00	0.00	3.07	0.00	0.00	0.00	0.00	0.00	0.00	2.91
ACTA	0.00	0.00	0.00	6.99	0.00	0.00	0.00	0.00	0.00	0.00	7.33	0.00	0.00	0.00
CCAA	0.00	0.00	0.00	9.30	0.00	0.00	0.00	0.00	0.00	0.00	9.73	0.00	0.00	0.00
CCCA	0.00	0.00	0.00	7.52	0.00	0.00	0.00	0.00	0.00	0.00	7.93	0.00	0.00	0.00
CCGA	0.00	0.00	0.00	0.00	0.00	0.00	2.41	0.00	0.00	0.00	0.00	0.00	0.00	2.51
CCTA	0.00	0.00	0.00	8.69	0.00	0.00	0.00	0.00	0.00	0.00	9.10	0.00	0.00	0.00
GCAA	0.00	4.05	0.00	0.00	0.00	0.00	0.00	0.00	4.65	0.00	0.00	0.00	0.00	0.00
GCCA	0.00	0.00	0.00	5.20	0.00	0.00	0.00	0.00	0.00	0.00	5.46	0.00	0.00	0.00
GCGA	0.00	0.00	13.81	0.00	0.00	0.00	0.00	0.00	0.00	13.89	0.00	0.00	0.00	0.00
GCTA	0.00	0.00	0.00	5.70	0.00	0.00	0.00	0.00	0.00	0.00	5.97	0.00	0.00	0.00
TCAA	0.00	0.00	0.00	0.00	6.26	0.00	0.00	0.00	0.00	0.00	0.00	6.21	0.00	0.00
TCCA	0.00	0.00	0.00	9.80	0.00	0.00	0.00	0.00	0.00	0.00	10.21	0.00	0.00	0.00
TCGA	0.00	0.00	11.87	0.00	0.00	0.00	0.00	0.00	0.00	12.24	0.00	0.00	0.00	0.00
TCTA	0.00	0.00	0.00	0.00	8.05	0.00	0.00	0.00	0.00	0.00	0.00	8.00	0.00	0.00
ACAG	13.14	0.00	0.00	0.00	0.00	0.00	0.00	13.38	0.00	0.00	0.00	0.00	0.00	0.00
ACCG	0.00	0.00	0.00	0.00	0.00	0.00	2.76	0.00	0.00	0.00	0.00	0.00	0.00	2.91
ACGG	0.00	0.00	12.62	0.00	0.00	0.00	0.00	0.00	0.00	12.22	0.00	0.00	0.00	0.00
ACTG	17.06	0.00	0.00	0.00	0.00	0.00	0.00	17.11	0.00	0.00	0.00	0.00	0.00	0.00
CCAG	0.00	0.00	0.00	0.00	0.00	0.00	3.16	0.00	0.00	0.00	0.00	0.00	0.00	3.32
CCCG	0.00	0.00	0.00	0.00	0.00	0.00	3.22	0.00	0.00	0.00	0.00	0.00	0.00	3.24
CCGG	0.00	0.00	0.00	0.00	0.00	0.00	3.64	0.00	0.00	0.00	0.00	0.00	0.00	3.27
CCTG	0.00	0.00	0.00	0.00	0.00	0.00	4.32	0.00	0.00	0.00	0.00	0.00	0.00	4.51
GCAG	0.00	0.00	0.00	0.00	0.00	0.00	3.48	0.00	0.00	0.00	0.00	0.00	0.00	3.34
GCCG	0.00	0.00	14.79	0.00	0.00	0.00	0.00	0.00	0.00	15.62	0.00	0.00	0.00	0.00
GCGG	0.00	0.00	15.50	0.00	0.00	0.00	0.00	0.00	0.00	13.92	0.00	0.00	0.00	0.00
GCTG	0.00	0.00	0.00	0.00	0.00	0.00	3.82	0.00	0.00	0.00	0.00	0.00	0.00	3.78
TCAG	0.00	0.00	0.00	12.20	0.00	0.00	0.00	0.00	0.00	0.00	10.90	0.00	0.00	0.00
TCCG	0.00	0.00	0.00	5.76	0.00	0.00	0.00	0.00	0.00	0.00	5.60	0.00	0.00	0.00
TCGG	0.00	0.00	8.40	0.00	0.00	0.00	0.00	0.00	0.00	8.65	0.00	0.00	0.00	0.00
TCTG	0.00	0.00	0.00	16.63	0.00	0.00	0.00	0.00	0.00	0.00	15.02	0.00	0.00	0.00
ACAT	0.00	0.00	0.00	0.00	7.67	0.00	0.00	0.00	0.00	0.00	0.00	7.71	0.00	0.00
ACCT	0.00	4.78	0.00	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00
ACGT	0.00	23.47	0.00	0.00	0.00	0.00	0.00	0.00	23.18	0.00	0.00	0.00	0.00	0.00
ACTT	0.00	0.00	0.00	0.00	5.43	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00
CCAT	0.00	0.00	0.00	0.00	6.02	0.00	0.00	0.00	0.00	0.00	0.00	6.02	0.00	0.00
CCCT	0.00	0.00	0.00	0.00	5.59	0.00	0.00	0.00	0.00	0.00	0.00	5.63	0.00	0.00
CCGT	0.00	17.66	0.00	0.00	0.00	0.00	0.00	0.00	17.12	0.00	0.00	0.00	0.00	0.00
CCTT	0.00	0.00	0.00	0.00	7.01	0.00	0.00	0.00	0.00	0.00	0.00	7.04	0.00	0.00
GCAT	0.00	0.00	0.00	0.00	5.98	0.00	0.00	0.00	0.00	0.00	0.00	6.01	0.00	0.00
GCCT	0.00	5.74	0.00	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00	0.00	0.00
GCGT	0.00	20.46	0.00	0.00	0.00	0.00	0.00	0.00	19.80	0.00	0.00	0.00	0.00	0.00
GCTT	0.00	0.00	0.00	0.00	5.88	0.00	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00
TCAT	0.00	11.42	0.00	0.00	0.00	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00
TCCT	0.00	0.00	0.00	0.00	7.81	0.00	0.00	0.00	0.00	0.00	0.00	7.76	0.00	0.00
TCGT	0.00	12.42	0.00	0.00	0.00	0.00	0.00	0.00	12.30	0.00	0.00	0.00	0.00	0.00
TCTT	0.00	0.00	0.00	0.00	9.47	0.00	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00

Table 7: Weights for the first 48 mutation categories for the 7 clusters in Clustering-B (see Table 4). The conventions are the same as in Table 5.

Mutation	Cl-1	C1-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7
ATAA	14.61	0.00	0.00	0.00	0.00	0.00	0.00	14.91	0.00	0.00	0.00	0.00	0.00	0.00
ATCA	0.00	0.00	0.00	0.00	0.00	0.00	3.46	0.00	0.00	0.00	0.00	0.00	0.00	3.65
ATGA	13.85	0.00	0.00	0.00	0.00	0.00	0.00	14.13	0.00	0.00	0.00	0.00	0.00	0.00
ATTA	0.00	0.00	0.00	0.00	0.00	5.95	0.00	0.00	0.00	0.00	0.00	0.00	6.08	0.00
CTAA	0.00	0.00	0.00	0.00	0.00	0.00	3.93	0.00	0.00	0.00	0.00	0.00	0.00	4.01
CTCA	12.75	0.00	0.00	0.00	0.00	0.00	0.00	12.93	0.00	0.00	0.00	0.00	0.00	0.00
CTGA	0.00	0.00	0.00	5.54	0.00	0.00	0.00	0.00	0.00	0.00	5.73	0.00	0.00	0.00
CTTA	0.00	0.00	0.00	0.00	0.00	4.55	0.00	0.00	0.00	0.00	0.00	0.00	4.62	0.00
GTAA	0.00	0.00	0.00	0.00	0.00	0.00	3.44	0.00	0.00	0.00	0.00	0.00	0.00	3.36
GTCA	0.00	0.00	15.20	0.00	0.00	0.00	0.00	0.00	0.00	15.36	0.00	0.00	0.00	0.00
GTGA	0.00	0.00	0.00	0.00	0.00	0.00	3.13	0.00	0.00	0.00	0.00	0.00	0.00	3.26
GTTA	0.00	0.00	0.00	0.00	0.00	0.00	4.04	0.00	0.00	0.00	0.00	0.00	0.00	4.00
TTAA	0.00	0.00	0.00	0.00	0.00	5.49	0.00	0.00	0.00	0.00	0.00	0.00	5.64	0.00
TTCA	0.00	0.00	0.00	0.00	0.00	0.00	4.99	0.00	0.00	0.00	0.00	0.00	0.00	4.77
TTGA	0.00	0.00	0.00	0.00	0.00	0.00	2.94	0.00	0.00	0.00	0.00	0.00	0.00	3.05
TTTA	0.00	0.00	0.00	0.00	0.00	5.65	0.00	0.00	0.00	0.00	0.00	0.00	5.77	0.00
ATAC	0.00	0.00	0.00	0.00	7.03	0.00	0.00	0.00	0.00	0.00	0.00	7.06	0.00	0.00
ATCC	0.00	0.00	0.00	0.00	0.00	3.53	0.00	0.00	0.00	0.00	0.00	0.00	3.64	0.00
ATGC	0.00	0.00	0.00	0.00	4.97	0.00	0.00	0.00	0.00	0.00	0.00	4.98	0.00	0.00
ATTC	0.00	0.00	0.00	0.00	6.30	0.00	0.00	0.00	0.00	0.00	0.00	6.34	0.00	0.00
CTAC	0.00	0.00	0.00	0.00	0.00	4.03	0.00	0.00	0.00	0.00	0.00	0.00	4.08	0.00
CTCC	0.00	0.00	0.00	0.00	0.00	4.59	0.00	0.00	0.00	0.00	0.00	0.00	4.61	0.00
CTGC	0.00	0.00	0.00	0.00	0.00	5.75	0.00	0.00	0.00	0.00	0.00	0.00	5.79	0.00
CTTC	0.00	0.00	0.00	0.00	0.00	7.65	0.00	0.00	0.00	0.00	0.00	0.00	7.41	0.00
GTAC	0.00	0.00	0.00	0.00	0.00	5.18	0.00	0.00	0.00	0.00	0.00	0.00	5.32	0.00
GTCC	0.00	0.00	0.00	0.00	0.00	0.00	4.18	0.00	0.00	0.00	0.00	0.00	0.00	4.36
GTGC	0.00	0.00	0.00	0.00	0.00	4.63	0.00	0.00	0.00	0.00	0.00	0.00	4.76	0.00
GIIC	0.00	0.00	0.00	0.00	0.00	5.42	0.00	0.00	0.00	0.00	0.00	0.00	5.62	0.00
TIAC	0.00	0.00	0.00	0.00	0.00	5.32	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00
TICC	0.00	0.00	0.00	0.00	0.00	5.05	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00
TIGC	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.20
ATAC	0.00	0.00	0.00	0.00	0.00	1.62	0.00	0.00	0.00	0.00	0.00	0.00	1.62	0.00
ATAG	0.00	0.00	0.00	0.00	0.00	0.00	3.10	0.00	0.00	0.00	0.00	0.00	0.00	0.24
ATCC	0.00	0.00	0.00	0.00	0.00	0.00	2.70	0.00	0.00	0.00	0.00	0.00	0.00	2.04
ATTC	14 44	0.00	0.00	0.00	0.00	0.00	0.00	14.00	0.00	0.00	0.00	0.00	0.00	0.00
CTAC	0.00	0.00	0.00	0.00	0.00	0.00	2.69	14.09	0.00	0.00	0.00	0.00	0.00	2.66
CTCC	0.00	0.00	0.00	0.00	0.00	0.00	4.50	0.00	0.00	0.00	0.00	0.00	0.00	4.30
CTCC	0.00	0.00	0.00	0.00	0.00	0.00	3.07	0.00	0.00	0.00	0.00	0.00	0.00	3 10
CTTC	0.00	0.00	0.00	0.00	0.00	10.44	0.00	0.00	0.00	0.00	0.00	0.00	0.55	0.00
GTAG	0.00	0.00	0.00	0.00	0.00	0.00	2.52	0.00	0.00	0.00	0.00	0.00	0.00	2.36
GTCC	0.00	0.00	7.80	0.00	0.00	0.00	0.00	0.00	0.00	8 11	0.00	0.00	0.00	0.00
GTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.18	0.00	0.00	0.00	0.00	0.00	0.00	3 25
GTTG	14 14	0.00	0.00	0.00	0.00	0.00	0.00	13 44	0.00	0.00	0.00	0.00	0.00	0.00
TTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.48	0.00	0.00	0.00	0.00	0.00	0.00	3.62
TTCG	0.00	0.00	0.00	0.00	0.00	0.00	2.91	0.00	0.00	0.00	0.00	0.00	0.00	2.87
TTGG	0.00	0.00	0.00	0.00	0.00	0.00	4.62	0.00	0.00	0.00	0.00	0.00	0.00	4.49
TTTG	0.00	0.00	0.00	0.00	0.00	8.95	0.00	0.00	0.00	0.00	0.00	0.00	8.65	0.00
	0.00	0.00	0.00	0.00	0.00	5.55	0.00	5.00	0.00	5.00	0.00	0.00	0.00	0.00

Table 8: Table 7 continued: weights for the next 48 mutation categories.

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	C1-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
ACAA	0.00	0.00	0.00	6.55	0.00	0.00	0.00	0.00	0.00	0.00	6.55	0.00	0.00	0.00
ACCA	0.00	0.00	0.00	0.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00	6.08	0.00	0.00
ACGA	0.00	0.00	0.00	0.00	0.00	0.00	4.06	0.00	0.00	0.00	0.00	0.00	0.00	4.00
ACTA	0.00	0.00	0.00	0.00	6.16	0.00	0.00	0.00	0.00	0.00	0.00	6.38	0.00	0.00
CCAA	0.00	0.00	0.00	0.00	7.91	0.00	0.00	0.00	0.00	0.00	0.00	8.10	0.00	0.00
CCCA	0.00	0.00	0.00	0.00	6.46	0.00	0.00	0.00	0.00	0.00	0.00	6.68	0.00	0.00
CCGA	0.00	0.00	6.40	0.00	0.00	0.00	0.00	0.00	0.00	6.40	0.00	0.00	0.00	0.00
CCTA	0.00	0.00	0.00	0.00	0.00	6.97	0.00	0.00	0.00	0.00	0.00	0.00	7.02	0.00
GCAA	4.05	0.00	0.00	0.00	0.00	0.00	0.00	4.65	0.00	0.00	0.00	0.00	0.00	0.00
GCCA	0.00	0.00	0.00	0.00	4.56	0.00	0.00	0.00	0.00	0.00	0.00	4.73	0.00	0.00
GCGA	0.00	13.81	0.00	0.00	0.00	0.00	0.00	0.00	13.89	0.00	0.00	0.00	0.00	0.00
GCTA	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00	0.00	5.20	0.00	0.00
TCAA	0.00	0.00	0.00	6.26	0.00	0.00	0.00	0.00	0.00	0.00	6.21	0.00	0.00	0.00
TCCA	0.00	0.00	0.00	0.00	8.94	0.00	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00
TCGA	0.00	11.87	0.00	0.00	0.00	0.00	0.00	0.00	12.24	0.00	0.00	0.00	0.00	0.00
TCTA	0.00	0.00	0.00	8.05	0.00	0.00	0.00	0.00	0.00	0.00	8.00	0.00	0.00	0.00
ACAG	0.00	0.00	0.00	0.00	3.96	0.00	0.00	0.00	0.00	0.00	0.00	4.18	0.00	0.00
ACCG	0.00	0.00	7.23	0.00	0.00	0.00	0.00	0.00	0.00	7.29	0.00	0.00	0.00	0.00
ACGG	0.00	12.62	0.00	0.00	0.00	0.00	0.00	0.00	12.22	0.00	0.00	0.00	0.00	0.00
ACTG	0.00	0.00	0.00	0.00	4.77	0.00	0.00	0.00	0.00	0.00	0.00	5.03	0.00	0.00
CCAG	0.00	0.00	8.26	0.00	0.00	0.00	0.00	0.00	0.00	8.31	0.00	0.00	0.00	0.00
CCCG	0.00	0.00	0.00	0.00	0.00	0.00	3.91	0.00	0.00	0.00	0.00	0.00	0.00	4.02
CCGG	0.00	0.00	0.00	0.00	0.00	0.00	5.37	0.00	0.00	0.00	0.00	0.00	0.00	5.12
COTG	0.00	0.00	11.12	0.00	0.00	0.00	0.00	0.00	0.00	11.19	0.00	0.00	0.00	0.00
GCAG	0.00	0.00	0.00	0.00	0.00	0.00	4.61	0.00	0.00	0.00	0.00	0.00	0.00	4.57
GCCG	0.00	14.79	0.00	0.00	0.00	0.00	0.00	0.00	15.62	0.00	0.00	0.00	0.00	0.00
GCGG	0.00	15.50	0.00	0.00	0.00	0.00	0.00	0.00	13.92	0.00	0.00	0.00	0.00	0.00
TCAC	0.00	0.00	0.00	0.00	10.00	0.00	4.80	0.00	0.00	0.00	0.00	0.00	0.00	4.92
TCAG	0.00	0.00	0.00	0.00	5 10	0.00	0.00	0.00	0.00	0.00	0.00	9.05	0.00	0.00
TCCG	0.00	8.40	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.95	0.00	0.00
TCGG	0.00	0.40	0.00	0.00	14 10	0.00	0.00	0.00	0.00	0.00	0.00	12.52	0.00	0.00
ACAT	0.00	0.00	0.00	7.67	0.00	0.00	0.00	0.00	0.00	0.00	7 71	12.55	0.00	0.00
ACCT	4 78	0.00	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00	0.00
ACGT	23 47	0.00	0.00	0.00	0.00	0.00	0.00	23.18	0.00	0.00	0.00	0.00	0.00	0.00
ACTT	0.00	0.00	0.00	5.43	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00	0.00
CCAT	0.00	0.00	0.00	6.02	0.00	0.00	0.00	0.00	0.00	0.00	6.02	0.00	0.00	0.00
CCCT	0.00	0.00	0.00	5 59	0.00	0.00	0.00	0.00	0.00	0.00	5.63	0.00	0.00	0.00
CCGT	17 66	0.00	0.00	0.00	0.00	0.00	0.00	17 12	0.00	0.00	0.00	0.00	0.00	0.00
CCTT	0.00	0.00	0.00	7 01	0.00	0.00	0.00	0.00	0.00	0.00	7.04	0.00	0.00	0.00
GCAT	0.00	0.00	0.00	5.98	0.00	0.00	0.00	0.00	0.00	0.00	6.01	0.00	0.00	0.00
GCCT	5.74	0.00	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00	0.00	0.00	0.00
GCGT	20.46	0.00	0.00	0.00	0.00	0.00	0.00	19.80	0.00	0.00	0.00	0.00	0.00	0.00
GCTT	0.00	0.00	0.00	5.88	0.00	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00
TCAT	11.42	0.00	0.00	0.00	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00	0.00
TCCT	0.00	0.00	0.00	7.81	0.00	0.00	0.00	0.00	0.00	0.00	7.76	0.00	0.00	0.00
TCGT	12.42	0.00	0.00	0.00	0.00	0.00	0.00	12.30	0.00	0.00	0.00	0.00	0.00	0.00
TCTT	0.00	0.00	0.00	9.47	0.00	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00	0.00

Table 9: Weights for the first 48 mutation categories for the 7 clusters in Clustering-C (see Table 4). The conventions are the same as in Table 5.

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
ATAA	0.00	0.00	0.00	0.00	4.18	0.00	0.00	0.00	0.00	0.00	0.00	4.52	0.00	0.00
ATCA	0.00	0.00	8.99	0.00	0.00	0.00	0.00	0.00	0.00	9.10	0.00	0.00	0.00	0.00
ATGA	0.00	0.00	0.00	0.00	4.02	0.00	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00
ATTA	0.00	0.00	0.00	0.00	0.00	5.72	0.00	0.00	0.00	0.00	0.00	0.00	5.85	0.00
CTAA	0.00	0.00	10.32	0.00	0.00	0.00	0.00	0.00	0.00	9.83	0.00	0.00	0.00	0.00
CTCA	0.00	0.00	0.00	0.00	3.79	0.00	0.00	0.00	0.00	0.00	0.00	3.98	0.00	0.00
CTGA	0.00	0.00	0.00	0.00	4.88	0.00	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00
CTTA	0.00	0.00	0.00	0.00	0.00	4.42	0.00	0.00	0.00	0.00	0.00	0.00	4.47	0.00
GTAA	0.00	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00	0.00	0.00	0.00	0.00	4.35
GTCA	0.00	15.20	0.00	0.00	0.00	0.00	0.00	0.00	15.36	0.00	0.00	0.00	0.00	0.00
GTGA	0.00	0.00	8.23	0.00	0.00	0.00	0.00	0.00	0.00	8.16	0.00	0.00	0.00	0.00
GTTA	0.00	0.00	0.00	0.00	0.00	0.00	5.13	0.00	0.00	0.00	0.00	0.00	0.00	5.19
TTAA	0.00	0.00	0.00	0.00	0.00	5.30	0.00	0.00	0.00	0.00	0.00	0.00	5.43	0.00
TTCA	0.00	0.00	0.00	0.00	0.00	0.00	6.64	0.00	0.00	0.00	0.00	0.00	0.00	6.58
TTGA	0.00	0.00	7.82	0.00	0.00	0.00	0.00	0.00	0.00	7.57	0.00	0.00	0.00	0.00
TTTA	0.00	0.00	0.00	0.00	0.00	5.44	0.00	0.00	0.00	0.00	0.00	0.00	5.55	0.00
ATAC	0.00	0.00	0.00	7.03	0.00	0.00	0.00	0.00	0.00	0.00	7.06	0.00	0.00	0.00
ATCC	0.00	0.00	10.75	0.00	0.00	0.00	0.00	0.00	0.00	10.92	0.00	0.00	0.00	0.00
ATGC	0.00	0.00	0.00	4.97	0.00	0.00	0.00	0.00	0.00	0.00	4.98	0.00	0.00	0.00
ATTC	0.00	0.00	0.00	6.30	0.00	0.00	0.00	0.00	0.00	0.00	6.34	0.00	0.00	0.00
CTAC	0.00	0.00	0.00	0.00	0.00	3.91	0.00	0.00	0.00	0.00	0.00	0.00	3.94	0.00
CTCC	0.00	0.00	0.00	0.00	0.00	4.44	0.00	0.00	0.00	0.00	0.00	0.00	4.45	0.00
CTGC	0.00	0.00	0.00	0.00	0.00	5.56	0.00	0.00	0.00	0.00	0.00	0.00	5.61	0.00
CTTC	0.00	0.00	0.00	0.00	0.00	7.39	0.00	0.00	0.00	0.00	0.00	0.00	7.16	0.00
GTAC	0.00	0.00	0.00	0.00	0.00	5.00	0.00	0.00	0.00	0.00	0.00	0.00	5.14	0.00
GTCC	0.00	0.00	10.41	0.00	0.00	0.00	0.00	0.00	0.00	10.61	0.00	0.00	0.00	0.00
GTGC	0.00	0.00	0.00	0.00	0.00	4.47	0.00	0.00	0.00	0.00	0.00	0.00	4.59	0.00
GTTC	0.00	0.00	0.00	0.00	0.00	5.21	0.00	0.00	0.00	0.00	0.00	0.00	5.40	0.00
TTAC	0.00	0.00	0.00	0.00	0.00	5.13	0.00	0.00	0.00	0.00	0.00	0.00	5.26	0.00
TTCC	0.00	0.00	0.00	0.00	0.00	4.84	0.00	0.00	0.00	0.00	0.00	0.00	4.96	0.00
TTGC	0.00	0.00	10.48	0.00	0.00	0.00	0.00	0.00	0.00	10.62	0.00	0.00	0.00	0.00
TTTC	0.00	0.00	0.00	0.00	0.00	7.53	0.00	0.00	0.00	0.00	0.00	0.00	7.52	0.00
ATAG	0.00	0.00	0.00	0.00	0.00	0.00	3.98	0.00	0.00	0.00	0.00	0.00	0.00	4.09
ATCG	0.00	0.00	0.00	0.00	0.00	0.00	3.81	0.00	0.00	0.00	0.00	0.00	0.00	3.70
ATGG	0.00	0.00	0.00	0.00	0.00	0.00	3.97	0.00	0.00	0.00	0.00	0.00	0.00	3.99
ATTG	0.00	0.00	0.00	0.00	0.00	0.00	7.13	0.00	0.00	0.00	0.00	0.00	0.00	7.08
CTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.55	0.00	0.00	0.00	0.00	0.00	0.00	3.56
CTCG	0.00	0.00	0.00	0.00	0.00	0.00	6.52	0.00	0.00	0.00	0.00	0.00	0.00	6.31
CTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.67	0.00	0.00	0.00	0.00	0.00	0.00	3.83
CTTG	0.00	0.00	0.00	0.00	0.00	10.06	0.00	0.00	0.00	0.00	0.00	0.00	9.27	0.00
GTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.58	0.00	0.00	0.00	0.00	0.00	0.00	3.49
GTCG	0.00	7.80	0.00	0.00	0.00	0.00	0.00	0.00	8.11	0.00	0.00	0.00	0.00	0.00
GTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.82	0.00	0.00	0.00	0.00	0.00	0.00	3.98
GTTG	0.00	0.00	0.00	0.00	0.00	0.00	7.02	0.00	0.00	0.00	0.00	0.00	0.00	6.97
TTAG	0.00	0.00	0.00	0.00	0.00	0.00	4.24	0.00	0.00	0.00	0.00	0.00	0.00	4.43
TTCG	0.00	0.00	0.00	0.00	0.00	0.00	3.73	0.00	0.00	0.00	0.00	0.00	0.00	3.75
TTGG	0.00	0.00	0.00	0.00	0.00	0.00	6.10	0.00	0.00	0.00	0.00	0.00	0.00	6.06
TTTG	0.00	0.00	0.00	0.00	0.00	8.61	0.00	0.00	0.00	0.00	0.00	0.00	8.36	0.00

Table 10: Table 9 continued: weights for the next 48 mutation categories.

Mutation	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6
ACAA	0.00	0.00	6.55	0.00	0.00	0.00	0.00	0.00	6.55	0.00	0.00	0.00
ACCA	0.00	0.00	0.00	5.83	0.00	0.00	0.00	0.00	0.00	6.08	0.00	0.00
ACGA	0.00	0.00	0.00	0.00	0.00	2.75	0.00	0.00	0.00	0.00	0.00	2.63
ACTA	0.00	0.00	0.00	6.16	0.00	0.00	0.00	0.00	0.00	6.38	0.00	0.00
CCAA	0.00	0.00	0.00	7.91	0.00	0.00	0.00	0.00	0.00	8.10	0.00	0.00
CCCA	0.00	0.00	0.00	6.46	0.00	0.00	0.00	0.00	0.00	6.68	0.00	0.00
CCGA	0.00	0.00	0.00	0.00	0.00	2.13	0.00	0.00	0.00	0.00	0.00	2.25
CCTA	0.00	0.00	0.00	0.00	6.75	0.00	0.00	0.00	0.00	0.00	6.79	0.00
GCAA	4.05	0.00	0.00	0.00	0.00	0.00	4.65	0.00	0.00	0.00	0.00	0.00
GCCA	0.00	0.00	0.00	4.56	0.00	0.00	0.00	0.00	0.00	4.73	0.00	0.00
GCGA	0.00	13.81	0.00	0.00	0.00	0.00	0.00	13.89	0.00	0.00	0.00	0.00
GCTA	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00	5.20	0.00	0.00
TCAA	0.00	0.00	6.26	0.00	0.00	0.00	0.00	0.00	6.21	0.00	0.00	0.00
TCCA	0.00	0.00	0.00	8.94	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00
TCGA	0.00	11.87	0.00	0.00	0.00	0.00	0.00	12.24	0.00	0.00	0.00	0.00
TCTA	0.00	0.00	8.05	0.00	0.00	0.00	0.00	0.00	8.00	0.00	0.00	0.00
ACAG	0.00	0.00	0.00	3.96	0.00	0.00	0.00	0.00	0.00	4.18	0.00	0.00
ACCG	0.00	0.00	0.00	0.00	0.00	2.46	0.00	0.00	0.00	0.00	0.00	2.61
ACGG	0.00	12.62	0.00	0.00	0.00	0.00	0.00	12.22	0.00	0.00	0.00	0.00
ACTG	0.00	0.00	0.00	4.77	0.00	0.00	0.00	0.00	0.00	5.03	0.00	0.00
CCAG	0.00	0.00	0.00	0.00	0.00	2.81	0.00	0.00	0.00	0.00	0.00	2.98
CCCG	0.00	0.00	0.00	0.00	0.00	2.88	0.00	0.00	0.00	0.00	0.00	2.92
CCGG	0.00	0.00	0.00	0.00	0.00	3.29	0.00	0.00	0.00	0.00	0.00	2.97
CCTG	0.00	0.00	0.00	0.00	0.00	3.84	0.00	0.00	0.00	0.00	0.00	4.05
GCAG	0.00	0.00	0.00	0.00	0.00	3.13	0.00	0.00	0.00	0.00	0.00	3.02
GCCG	0.00	14.79	0.00	0.00	0.00	0.00	0.00	15.62	0.00	0.00	0.00	0.00
GCGG	0.00	15.50	0.00	0.00	0.00	0.00	0.00	13.92	0.00	0.00	0.00	0.00
GCTG	0.00	0.00	0.00	0.00	0.00	3.44	0.00	0.00	0.00	0.00	0.00	3.42
TCAG	0.00	0.00	0.00	10.31	0.00	0.00	0.00	0.00	0.00	9.03	0.00	0.00
TCCG	0.00	0.00	0.00	5.10	0.00	0.00	0.00	0.00	0.00	4.95	0.00	0.00
TCGG	0.00	8.40	0.00	0.00	0.00	0.00	0.00	8.65	0.00	0.00	0.00	0.00
TCTG	0.00	0.00	0.00	14.10	0.00	0.00	0.00	0.00	0.00	12.53	0.00	0.00
ACAT	0.00	0.00	7.67	0.00	0.00	0.00	0.00	0.00	7.71	0.00	0.00	0.00
ACCT	4.78	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00	0.00	0.00	0.00
ACGT	23.47	0.00	0.00	0.00	0.00	0.00	23.18	0.00	0.00	0.00	0.00	0.00
ACTT	0.00	0.00	5.43	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00	0.00
CCAT	0.00	0.00	6.02	0.00	0.00	0.00	0.00	0.00	6.02	0.00	0.00	0.00
CCCT	0.00	0.00	5.59	0.00	0.00	0.00	0.00	0.00	5.63	0.00	0.00	0.00
CCGT	17.66	0.00	0.00	0.00	0.00	0.00	17.12	0.00	0.00	0.00	0.00	0.00
CCTT	0.00	0.00	7.01	0.00	0.00	0.00	0.00	0.00	7.04	0.00	0.00	0.00
GCAT	0.00	0.00	5.98	0.00	0.00	0.00	0.00	0.00	6.01	0.00	0.00	0.00
GCCT	5.74	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00	0.00	0.00
GCGT	20.46	0.00	0.00	0.00	0.00	0.00	19.80	0.00	0.00	0.00	0.00	0.00
GCTT	0.00	0.00	5.88	0.00	0.00	0.00	0.00	0.00	5.93	0.00	0.00	0.00
TCAT	11.42	0.00	0.00	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00
TCCT	0.00	0.00	7.81	0.00	0.00	0.00	0.00	0.00	7.76	0.00	0.00	0.00
TCGT	12.42	0.00	0.00	0.00	0.00	0.00	12.30	0.00	0.00	0.00	0.00	0.00
TCTT	0.00	0.00	9.47	0.00	0.00	0.00	0.00	0.00	9.29	0.00	0.00	0.00

Table 11: Weights for the first 48 mutation categories for the 6 clusters in Clustering-D (see Table 4). The conventions are the same as in Table 5.

Mutation	Cl-1	Cl-2	C1-3	Cl-4	Cl-5	Cl-6	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6
ATAA	0.00	0.00	0.00	4.18	0.00	0.00	0.00	0.00	0.00	4.52	0.00	0.00
ATCA	0.00	0.00	0.00	0.00	0.00	3.11	0.00	0.00	0.00	0.00	0.00	3.29
ATGA	0.00	0.00	0.00	4.02	0.00	0.00	0.00	0.00	0.00	4.30	0.00	0.00
ATTA	0.00	0.00	0.00	0.00	5.54	0.00	0.00	0.00	0.00	0.00	5.66	0.00
CTAA	0.00	0.00	0.00	0.00	0.00	3 49	0.00	0.00	0.00	0.00	0.00	3.60
CTCA	0.00	0.00	0.00	3 79	0.00	0.00	0.00	0.00	0.00	3.98	0.00	0.00
CTGA	0.00	0.00	0.00	4 88	0.00	0.00	0.00	0.00	0.00	5.02	0.00	0.00
CTTA	0.00	0.00	0.00	0.00	4 28	0.00	0.00	0.00	0.00	0.00	4 33	0.00
GTAA	0.00	0.00	0.00	0.00	0.00	3.09	0.00	0.00	0.00	0.00	0.00	3.04
GTCA	0.00	15 20	0.00	0.00	0.00	0.00	0.00	15 36	0.00	0.00	0.00	0.00
GTGA	0.00	0.00	0.00	0.00	0.00	2 79	0.00	0.00	0.00	0.00	0.00	2.93
GTTA	0.00	0.00	0.00	0.00	0.00	3.65	0.00	0.00	0.00	0.00	0.00	3.63
TTAA	0.00	0.00	0.00	0.00	5.13	0.00	0.00	0.00	0.00	0.00	5.26	0.00
TTCA	0.00	0.00	0.00	0.00	0.10	4 50	0.00	0.00	0.00	0.00	0.00	4.22
TTCA	0.00	0.00	0.00	0.00	0.00	9.62	0.00	0.00	0.00	0.00	0.00	9.55
TTTA	0.00	0.00	0.00	0.00	5.00	2.03	0.00	0.00	0.00	0.00	5.29	2.74
ATAC	0.00	0.00	7.02	0.00	0.00	0.00	0.00	0.00	7.06	0.00	0.00	0.00
ATCC	0.00	0.00	0.00	0.00	2 20	0.00	0.00	0.00	0.00	0.00	2 20	0.00
ATCC	0.00	0.00	4.07	0.00	0.00	0.00	0.00	0.00	4.08	0.00	0.00	0.00
ATGC	0.00	0.00	6.20	0.00	0.00	0.00	0.00	0.00	6.94	0.00	0.00	0.00
CTAC	0.00	0.00	0.30	0.00	2.79	0.00	0.00	0.00	0.34	0.00	0.00	0.00
CTAC	0.00	0.00	0.00	0.00	3.10	0.00	0.00	0.00	0.00	0.00	3.81	0.00
CTCC	0.00	0.00	0.00	0.00	4.30	0.00	0.00	0.00	0.00	0.00	4.31	0.00
CIGC	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.00	0.00	0.00	0.41	0.00
CIIC	0.00	0.00	0.00	0.00	(.14	0.00	0.00	0.00	0.00	0.00	6.92	0.00
GIAC	0.00	0.00	0.00	0.00	4.84	0.00	0.00	0.00	0.00	0.00	4.96	0.00
GIUU	0.00	0.00	0.00	0.00	0.00	3.81	0.00	0.00	0.00	0.00	0.00	3.95
GIGC	0.00	0.00	0.00	0.00	4.32	0.00	0.00	0.00	0.00	0.00	4.43	0.00
GIIC	0.00	0.00	0.00	0.00	5.05	0.00	0.00	0.00	0.00	0.00	5.23	0.00
TIAC	0.00	0.00	0.00	0.00	4.97	0.00	0.00	0.00	0.00	0.00	5.10	0.00
TTCC	0.00	0.00	0.00	0.00	4.69	0.00	0.00	0.00	0.00	0.00	4.79	0.00
TTGC	0.00	0.00	0.00	0.00	0.00	3.61	0.00	0.00	0.00	0.00	0.00	3.80
TTTC	0.00	0.00	0.00	0.00	7.29	0.00	0.00	0.00	0.00	0.00	7.28	0.00
ATAG	0.00	0.00	0.00	0.00	0.00	2.91	0.00	0.00	0.00	0.00	0.00	2.95
ATCG	0.00	0.00	0.00	0.00	0.00	2.46	0.00	0.00	0.00	0.00	0.00	2.30
ATGG	0.00	0.00	0.00	0.00	0.00	2.80	0.00	0.00	0.00	0.00	0.00	2.76
ATTG	0.00	0.00	0.00	0.00	0.00	4.93	0.00	0.00	0.00	0.00	0.00	4.79
CTAG	0.00	0.00	0.00	0.00	0.00	2.47	0.00	0.00	0.00	0.00	0.00	2.43
CTCG	0.00	0.00	0.00	0.00	0.00	4.21	0.00	0.00	0.00	0.00	0.00	3.93
CTGG	0.00	0.00	0.00	0.00	0.00	2.80	0.00	0.00	0.00	0.00	0.00	2.90
CTTG	0.00	0.00	0.00	0.00	9.67	0.00	0.00	0.00	0.00	0.00	8.89	0.00
GTAG	0.00	0.00	0.00	0.00	0.00	2.30	0.00	0.00	0.00	0.00	0.00	2.16
GTCG	0.00	7.80	0.00	0.00	0.00	0.00	0.00	8.11	0.00	0.00	0.00	0.00
GTGG	0.00	0.00	0.00	0.00	0.00	2.85	0.00	0.00	0.00	0.00	0.00	2.93
GTTG	0.00	0.00	0.00	0.00	0.00	4.83	0.00	0.00	0.00	0.00	0.00	4.73
TTAG	0.00	0.00	0.00	0.00	0.00	3.20	0.00	0.00	0.00	0.00	0.00	3.30
TTCG	0.00	0.00	0.00	0.00	0.00	2.64	0.00	0.00	0.00	0.00	0.00	2.60
TTGG	0.00	0.00	0.00	0.00	0.00	4.20	0.00	0.00	0.00	0.00	0.00	4.08
TTTG	0.00	0.00	0.00	0.00	8.31	0.00	0.00	0.00	0.00	0.00	8.05	0.00

Table 12: Table 11 continued: weights for the next 48 mutation categories.

\_

Table 13: Weights (in the units of 1%, rounded to 2 digits) for the first 48 mutation categories for the 7 clusters in Clustering-A (see Table 4) based on unnormalized (columns 2-8) and normalized (columns 9-15) regressions with the exposures computed via geometric means (see Subsection 2.6 for details). Here "weights based on unnormalized regressions" are given by (13), (14) and (19), while "weights based on normalized regressions" are given by (17), (14) and (21). Other conventions are the same as in Table 5.

	01.1	C1 0	<u>Cl</u> 0	<u><u> </u></u>	C1 5	C1 0	C1 5	01.1	CI o	<u>(1)</u>	<u><u> </u></u>	C11 F	<u> </u>	<u></u>
Mutation	CI-1	CI-2	CI-3	CI-4	CI-5	CI-6	CI-7	CI-1	CI-2	CI-3	CI-4	CI-5	CI-6	CI-7
ACAA	0.00	0.00	0.00	6.54	0.00	0.00	0.00	0.00	0.00	0.00	6.54	0.00	0.00	0.00
ACCA	0.00	0.00	0.00	0.00	6.16	0.00	0.00	0.00	0.00	0.00	0.00	6.20	0.00	0.00
ACGA	0.00	0.00	0.00	0.00	0.00	0.00	4.12	0.00	0.00	0.00	0.00	0.00	0.00	4.05
ACTA	0.00	0.00	0.00	0.00	6.38	0.00	0.00	0.00	0.00	0.00	0.00	6.44	0.00	0.00
CCAA	0.00	0.00	0.00	0.00	8.27	0.00	0.00	0.00	0.00	0.00	0.00	8.27	0.00	0.00
CCCA	0.00	0.00	0.00	0.00	6.73	0.00	0.00	0.00	0.00	0.00	0.00	6.77	0.00	0.00
CCGA	0.00	0.00	7.32	0.00	0.00	0.00	0.00	0.00	0.00	7.24	0.00	0.00	0.00	0.00
CCTA	0.00	0.00	0.00	0.00	0.00	6.77	0.00	0.00	0.00	0.00	0.00	0.00	6.76	0.00
GCAA	4.31	0.00	0.00	0.00	0.00	0.00	0.00	4.68	0.00	0.00	0.00	0.00	0.00	0.00
GCCA	0.00	0.00	0.00	0.00	4.70	0.00	0.00	0.00	0.00	0.00	0.00	4.75	0.00	0.00
GCGA	0.00	13.79	0.00	0.00	0.00	0.00	0.00	0.00	13.76	0.00	0.00	0.00	0.00	0.00
GCTA	0.00	0.00	0.00	0.00	5.16	0.00	0.00	0.00	0.00	0.00	0.00	5.22	0.00	0.00
TCAA	0.00	0.00	0.00	6.22	0.00	0.00	0.00	0.00	0.00	0.00	6.20	0.00	0.00	0.00
TCCA	0.00	0.00	0.00	0.00	8.86	0.00	0.00	0.00	0.00	0.00	0.00	9.08	0.00	0.00
TCGA	0.00	11.96	0.00	0.00	0.00	0.00	0.00	0.00	12.13	0.00	0.00	0.00	0.00	0.00
TCTA	0.00	0.00	0.00	8.04	0.00	0.00	0.00	0.00	0.00	0.00	8.01	0.00	0.00	0.00
ACAG	0.00	0.00	0.00	0.00	4.08	0.00	0.00	0.00	0.00	0.00	0.00	4.16	0.00	0.00
ACCG	0.00	0.00	8.12	0.00	0.00	0.00	0.00	0.00	0.00	8.17	0.00	0.00	0.00	0.00
ACGG	0.00	12.58	0.00	0.00	0.00	0.00	0.00	0.00	12.32	0.00	0.00	0.00	0.00	0.00
ACTG	0.00	0.00	0.00	0.00	4.73	0.00	0.00	0.00	0.00	0.00	0.00	4.88	0.00	0.00
CCAG	0.00	0.00	9.34	0.00	0.00	0.00	0.00	0.00	0.00	9.36	0.00	0.00	0.00	0.00
CCCG	0.00	0.00	0.00	0.00	0.00	0.00	3.97	0.00	0.00	0.00	0.00	0.00	0.00	4.04
CCGG	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00	0.00	0.00	0.00	0.00	5.24
CCTG	0.00	0.00	12.56	0.00	0.00	0.00	0.00	0.00	0.00	12.61	0.00	0.00	0.00	0.00
GCAG	0.00	0.00	0.00	0.00	0.00	0.00	4.68	0.00	0.00	0.00	0.00	0.00	0.00	4.63
GCCG	0.00	14.96	0.00	0.00	0.00	0.00	0.00	0.00	15.53	0.00	0.00	0.00	0.00	0.00
GCGG	0.00	15.17	0.00	0.00	0.00	0.00	0.00	0.00	14.18	0.00	0.00	0.00	0.00	0.00
GCTG	0.00	0.00	0.00	0.00	0.00	0.00	4.92	0.00	0.00	0.00	0.00	0.00	0.00	4.94
TCAG	0.00	0.00	0.00	0.00	9.40	0.00	0.00	0.00	0.00	0.00	0.00	8.99	0.00	0.00
TCCG	0.00	0.00	0.00	0.00	4.93	0.00	0.00	0.00	0.00	0.00	0.00	4.90	0.00	0.00
TCGG	0.00	8.53	0.00	0.00	0.00	0.00	0.00	0.00	8.60	0.00	0.00	0.00	0.00	0.00
TCTG	0.00	0.00	0.00	0.00	13.10	0.00	0.00	0.00	0.00	0.00	0.00	12.56	0.00	0.00
ACAT	0.00	0.00	0.00	7.72	0.00	0.00	0.00	0.00	0.00	0.00	7.73	0.00	0.00	0.00
ACCT	4.86	0.00	0.00	0.00	0.00	0.00	0.00	5.01	0.00	0.00	0.00	0.00	0.00	0.00
ACGT	23.50	0.00	0.00	0.00	0.00	0.00	0.00	23.33	0.00	0.00	0.00	0.00	0.00	0.00
ACTT	0.00	0.00	0.00	5.45	0.00	0.00	0.00	0.00	0.00	0.00	5.47	0.00	0.00	0.00
CCAT	0.00	0.00	0.00	6.02	0.00	0.00	0.00	0.00	0.00	0.00	6.02	0.00	0.00	0.00
CCCT	0.00	0.00	0.00	5.60	0.00	0.00	0.00	0.00	0.00	0.00	5.62	0.00	0.00	0.00
CCGT	17.45	0.00	0.00	0.00	0.00	0.00	0.00	17.08	0.00	0.00	0.00	0.00	0.00	0.00
CCTT	0.00	0.00	0.00	7.03	0.00	0.00	0.00	0.00	0.00	0.00	7.05	0.00	0.00	0.00
GCAT	0.00	0.00	0.00	5.98	0.00	0.00	0.00	0.00	0.00	0.00	6.00	0.00	0.00	0.00
GCCT	5.85	0.00	0.00	0.00	0.00	0.00	0.00	5.97	0.00	0.00	0.00	0.00	0.00	0.00
GCGT	20.08	0.00	0.00	0.00	0.00	0.00	0.00	19.63	0.00	0.00	0.00	0.00	0.00	0.00
GCTT	0.00	0.00	0.00	5.90	0.00	0.00	0.00	0.00	0.00	0.00	5.92	0.00	0.00	0.00
TCAT	11.55	0.00	0.00	0.00	0.00	0.00	0.00	12.00	0.00	0.00	0.00	0.00	0.00	0.00
TOUT	0.00	0.00	0.00	7.77	0.00	0.00	0.00	0.00	0.00	0.00	7.75	0.00	0.00	0.00
TCGT	12.39	0.00	0.00	0.00	0.00	0.00	0.00	12.30	0.00	0.00	0.00	0.00	0.00	0.00
TCTT	0.00	0.00	0.00	9.35	0.00	0.00	0.00	0.00	0.00	0.00	9.27	0.00	0.00	0.00

$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Mutation	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7	Cl-1	Cl-2	Cl-3	Cl-4	C1-5	Cl-6	Cl-7
ATAA         0.00 <th< td=""><td>-</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></th<>	-														
ATCA0.000.001.0.060.000.000.000.000.001.0.150.000.000.000.000.00ATCA0.000.000.000.001.1.30.000	ATAA	0.00	0.00	0.00	0.00	4.41	0.00	0.00	0.00	0.00	0.00	0.00	4.51	0.00	0.00
ATGA         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.25         0.00         0.00           CTAA         0.00         0.	ATCA	0.00	0.00	10.06	0.00	0.00	0.00	0.00	0.00	0.00	10.15	0.00	0.00	0.00	0.00
ATTA         0.00 <th< td=""><td>ATGA</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.15</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.25</td><td>0.00</td><td>0.00</td></th<>	ATGA	0.00	0.00	0.00	0.00	4.15	0.00	0.00	0.00	0.00	0.00	0.00	4.25	0.00	0.00
CTAA         0.00         0.00         1.34         0.00         0.00         0.00         1.10         0.00         0.00         0.00           CTCA         0.00         0.	ATTA	0.00	0.00	0.00	0.00	0.00	5.59	0.00	0.00	0.00	0.00	0.00	0.00	5.64	0.00
CTCA         0.00 <th< td=""><td>CTAA</td><td>0.00</td><td>0.00</td><td>11.34</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>11.10</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td></th<>	CTAA	0.00	0.00	11.34	0.00	0.00	0.00	0.00	0.00	0.00	11.10	0.00	0.00	0.00	0.00
CTCA         0.00         0.00         0.00         0.00         0.00         0.00         0.00         5.07         0.00         0.00           GTTAA         0.00         0.00         0.00         0.00         4.33         0.00         0	CTCA	0.00	0.00	0.00	0.00	3.87	0.00	0.00	0.00	0.00	0.00	0.00	3.94	0.00	0.00
CTTA         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.31         0.00           GTAA         0.00         1.517         0.00         0	CTGA	0.00	0.00	0.00	0.00	5.08	0.00	0.00	0.00	0.00	0.00	0.00	5.07	0.00	0.00
GTAA         0.00 <th< td=""><td>CTTA</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.33</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.31</td><td>0.00</td></th<>	CTTA	0.00	0.00	0.00	0.00	0.00	4.33	0.00	0.00	0.00	0.00	0.00	0.00	4.31	0.00
GTCA         0.00         15.17         0.00         0.00         0.00         0.00         15.40         0.00         0.00         0.00         0.00           GTCA         0.00	GTAA	0.00	0.00	0.00	0.00	0.00	0.00	4.33	0.00	0.00	0.00	0.00	0.00	0.00	4.36
GTGA         0.00 <th< td=""><td>GTCA</td><td>0.00</td><td>15.17</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>15.40</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td></th<>	GTCA	0.00	15.17	0.00	0.00	0.00	0.00	0.00	0.00	15.40	0.00	0.00	0.00	0.00	0.00
GTTA         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         5.22         0.00           TTCA         0.00         0.	GTGA	0.00	0.00	9.30	0.00	0.00	0.00	0.00	0.00	0.00	9.24	0.00	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	GTTA	0.00	0.00	0.00	0.00	0.00	0.00	5.18	0.00	0.00	0.00	0.00	0.00	0.00	5.22
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTAA	0.00	0.00	0.00	0.00	0.00	5.21	0.00	0.00	0.00	0.00	0.00	0.00	5.21	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTCA	0.00	0.00	0.00	0.00	0.00	0.00	6.73	0.00	0.00	0.00	0.00	0.00	0.00	6.66
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTGA	0.00	0.00	8.62	0.00	0.00	0.00	0.00	0.00	0.00	8.51	0.00	0.00	0.00	0.00
ATAC         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00           ATCC         0.00         0.00         0.00         0.00         3.38         0.00         0.	TTTA	0.00	0.00	0.00	0.00	0.00	5.36	0.00	0.00	0.00	0.00	0.00	0.00	5.35	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATAC	0.00	0.00	0.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00	7.08	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATCC	0.00	0.00	0.00	0.00	0.00	3.38	0.00	0.00	0.00	0.00	0.00	0.00	3.40	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATGC	0.00	0.00	0.00	4.99	0.00	0.00	0.00	0.00	0.00	0.00	4.99	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATTC	0.00	0.00	0.00	6.34	0.00	0.00	0.00	0.00	0.00	0.00	6.36	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTAC	0.00	0.00	0.00	0.00	0.00	3.82	0.00	0.00	0.00	0.00	0.00	0.00	3.81	0.00
CTGC         0.00         0.00         0.00         5.27         0.00         0.00         0.00         0.00         5.35         0.00           CTTC         0.00         0.	CTCC	0.00	0.00	0.00	0.00	0.00	4.31	0.00	0.00	0.00	0.00	0.00	0.00	4.32	0.00
CTTC         0.00         0.00         0.00         7.09         0.00         0.00         0.00         0.00         0.00         7.01         0.00           GTAC         0.00         0.	CTGC	0.00	0.00	0.00	0.00	0.00	5.27	0.00	0.00	0.00	0.00	0.00	0.00	5.35	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTTC	0.00	0.00	0.00	0.00	0.00	7.09	0.00	0.00	0.00	0.00	0.00	0.00	7.01	0.00
GTCC         0.00         0.00         1.65         0.00 <th< td=""><td>GTAC</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.82</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>4.90</td><td>0.00</td></th<>	GTAC	0.00	0.00	0.00	0.00	0.00	4.82	0.00	0.00	0.00	0.00	0.00	0.00	4.90	0.00
GTGC         0.00         0.00         0.00         0.00         4.26         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.36         0.00           GTTC         0.00         0.	GTCC	0.00	0.00	11.65	0.00	0.00	0.00	0.00	0.00	0.00	11.80	0.00	0.00	0.00	0.00
GTTC         0.00         0.00         0.00         0.00         5.08         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         5.18         0.00           TTAC         0.00         0.00         0.00         0.00         5.06         0.00         0.00         0.00         0.00         0.00         5.09         0.00           TTCC         0.00<	GTGC	0.00	0.00	0.00	0.00	0.00	4.26	0.00	0.00	0.00	0.00	0.00	0.00	4.36	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	GTTC	0.00	0.00	0.00	0.00	0.00	5.08	0.00	0.00	0.00	0.00	0.00	0.00	5.18	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTAC	0.00	0.00	0.00	0.00	0.00	5.06	0.00	0.00	0.00	0.00	0.00	0.00	5.09	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTCC	0.00	0.00	0.00	0.00	0.00	4.69	0.00	0.00	0.00	0.00	0.00	0.00	4.76	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTGC	0.00	0.00	11.69	0.00	0.00	0.00	0.00	0.00	0.00	11.81	0.00	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	TTTC	0.00	0.00	0.00	0.00	0.00	7.37	0.00	0.00	0.00	0.00	0.00	0.00	7.31	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATAG	0.00	0.00	0.00	0.00	0.00	0.00	3.94	0.00	0.00	0.00	0.00	0.00	0.00	4.03
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATCG	0.00	0.00	0.00	0.00	0.00	0.00	3.83	0.00	0.00	0.00	0.00	0.00	0.00	3.74
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATGG	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	0.00	0.00	0.00	0.00	0.00	4.01
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	ATTG	0.00	0.00	0.00	0.00	0.00	0.00	6.98	0.00	0.00	0.00	0.00	0.00	0.00	7.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.50	0.00	0.00	0.00	0.00	0.00	0.00	3.52
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTCG	0.00	0.00	0.00	0.00	0.00	0.00	6.53	0.00	0.00	0.00	0.00	0.00	0.00	6.37
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.63	0.00	0.00	0.00	0.00	0.00	0.00	3.76
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	CTTG	0.00	0.00	0.00	0.00	0.00	9.36	0.00	0.00	0.00	0.00	0.00	0.00	9.13	0.00
GTCG         0.00         7.84         0.00         0.00         0.00         0.00         0.00         8.08         0.00 <th< td=""><td>GTAG</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>3.59</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>0.00</td><td>3.51</td></th<>	GTAG	0.00	0.00	0.00	0.00	0.00	0.00	3.59	0.00	0.00	0.00	0.00	0.00	0.00	3.51
GTGG         0.00         0.00         0.00         0.00         0.00         3.87         0.00         0.00         0.00         0.00         3.97           GTTG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         3.97           GTTG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         3.97           TTAG         0.00         0.00         0.00         0.00         0.00         6.71         0.00         0.00         0.00         0.00         6.77           TTAG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.32           TTCG         0.00         0.00         0.00         0.00         3.74         0.00         0.00         0.00         0.00         3.74           TTCG         0.00         0.00         0.00         0.00         6.11         0.00         0.00         0.00         0.00         3.69	GTCG	0.00	7.84	0.00	0.00	0.00	0.00	0.00	0.00	8.08	0.00	0.00	0.00	0.00	0.00
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	GTGG	0.00	0.00	0.00	0.00	0.00	0.00	3.87	0.00	0.00	0.00	0.00	0.00	0.00	3.97
TTAG         0.00         0.00         0.00         0.00         0.00         4.17         0.00         0.00         0.00         0.00         4.32           TTCG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.32           TTCG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         4.32           TTGG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         3.76           TTGG         0.00         0.00         0.00         0.00         6.11         0.00         0.00         0.00         0.00         6.09	GTTG	0.00	0.00	0.00	0.00	0.00	0.00	6.71	0.00	0.00	0.00	0.00	0.00	0.00	6.77
TTCG         0.00         0.00         0.00         0.00         3.74         0.00         0.00         0.00         0.00         3.76           TTGG         0.00         0.00         0.00         0.00         0.00         0.00         0.00         0.00         3.76           TTGG         0.00<	TTAG	0.00	0.00	0.00	0.00	0.00	0.00	4.17	0.00	0.00	0.00	0.00	0.00	0.00	4.32
TTGG 0.00 0.00 0.00 0.00 0.00 6.11 0.00 0.00	TTCG	0.00	0.00	0.00	0.00	0.00	0.00	3.74	0.00	0.00	0.00	0.00	0.00	0.00	3.76
	TTGG	0.00	0.00	0.00	0.00	0.00	0.00	6.11	0.00	0.00	0.00	0.00	0.00	0.00	6.09
TTTG 0.00 0.00 0.00 0.00 0.00 8.22 0.00 0.00	TTTG	0.00	0.00	0.00	0.00	0.00	8.22	0.00	0.00	0.00	0.00	0.00	0.00	8.12	0.00

Table 14: Table 13 continued: weights for the next 48 mutation categories.

Table 15: The within-cluster cross-sectional correlations  $\Theta_{sA}$  (columns 2-8), the overall correlations  $\Xi_s$  (column 11) based on the overall cross-sectional regressions, and multiple  $R^2$  and adjusted  $R^2$  of these regressions (columns 9 and 10). See Subsection 3.3 for details. Cancer types are labeled by X1 through X14 as in Table 2. All quantities are in the units of 1% rounded to 2 digits. The values above 80% are given in bold font.

Cancer Type	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7	r.sq	adj.r.sq	Overall Cor
X1	57.66	31.8	75.04	88.43	81.27	84.82	41.7	89.05	88.19	83.84
X2	90.57	66.35	81.97	79.64	41.42	-2.87	25.43	94.77	94.35	93.82
X3	93.29	-12.6	39.19	12.59	68.65	17.06	68.74	93.86	93.38	94.19
X4	9.88	16.97	52.94	79.11	81.85	46.74	7.34	58.18	54.9	61.53
X5	89.52	63.31	50.79	28.58	5.12	80.88	13.66	93.26	92.73	88.62
X6	86.53	34.07	48.92	76.77	85.01	19.59	34.54	89.57	88.75	91.28
X7	92.78	34.69	64.65	48.79	63.79	86.55	72.56	86.72	85.67	86.04
X8	-31.6	39.99	65.56	-46.21	-6.95	-3.36	61.8	69.52	67.12	41.88
X9	-28.63	53.86	-34.26	46.93	59.88	13.59	-12.39	77.76	76.02	70.18
X10	93.97	61.59	63.06	67.15	41.13	4.11	43.87	95.17	94.79	95.47
X11	88.16	56.6	66.76	55.12	90.27	16.33	26.3	95.02	94.63	89.62
X12	94.75	17.48	5.1	16.5	90	27.74	21.63	94.04	93.57	96.11
X13	97.05	58.21	75.77	78.67	88.42	20.28	44.07	96.31	96.02	95.35
X14	38.93	65.92	17.23	58.54	4.73	35.72	31.27	82.52	81.14	65.4

Table 16: The within-cluster cross-sectional correlations  $\Delta_{\alpha A}$  between the weights for 7 cancer signatures Sig1 through Sig7 of [Kakushadze and Yu, 2016b] and the weights (using normalized regressions with exposures based on arithmetic averages) for 7 clusters in Clustering A (see Subsection 3.3 for details). All quantities are in the units of 1% rounded to 2 digits. The values above 80% are given in bold font.

Signature	Cl-1	Cl-2	Cl-3	Cl-4	Cl-5	Cl-6	Cl-7
Sig1	92.05	10.29	-6.42	-8.33	51.12	29.06	20.61
$\operatorname{Sig2}$	-0.37	1.75	42.13	75.58	80.12	-27.92	-3.34
Sig3	-51.53	54.4	-37.16	28.19	32.98	12.37	-17.7
Sig4	31.56	11.97	54.43	56.83	-1.17	84.25	60.41
$\operatorname{Sig5}$	-42.53	40.31	62.96	-47.62	-8.34	-8.39	61.61
$\operatorname{Sig6}$	47.79	40.62	17.8	27.45	-27.96	16.87	16.97
$\operatorname{Sig7}$	80.94	19.87	55.03	33.4	13.89	-29.59	13.93



Figure 1: Horizontal axis: serial standard deviation  $\sigma'_i$  for N = 96 mutation categories (i = 1, ..., N) of cross-sectionally demeaned log-counts  $X'_{is}$  across n = 14cancer types (for samples aggregated by cancer types, so s = 1, ..., d, d = n). Vertical axis: density using R function density(). See Subsection 2.4.1 for details.



Figure 2: Cluster Cl-1 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6. Here and in all Figures below, for comparison and visualization convenience, we show all 96 channels on the horizontal axis even though the weights are nonzero only for the mutation categories belonging to a given cluster. Thus, in this cluster, only 8 weights are nonzero, to wit, for GCAA, ACCT, ACGT, CCGT, GCCT, GCGT, TCAT, TCGT.



Figure 3: Cluster Cl-1 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 4: Cluster Cl-2 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 5: Cluster Cl-2 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 6: Cluster Cl-3 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 7: Cluster Cl-3 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 8: Cluster Cl-4 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 9: Cluster Cl-4 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 10: Cluster Cl-5 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 11: Cluster Cl-5 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 12: Cluster Cl-6 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 13: Cluster Cl-6 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 14: Cluster Cl-7 in Clustering-A with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 15: Cluster Cl-7 in Clustering-A with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 5, 6.



Figure 16: Cluster Cl-1 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 17: Cluster Cl-1 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.


Figure 18: Cluster Cl-2 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 19: Cluster Cl-2 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 20: Cluster Cl-3 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 21: Cluster Cl-3 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 22: Cluster Cl-4 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 23: Cluster Cl-4 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 24: Cluster Cl-5 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 25: Cluster Cl-5 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 26: Cluster Cl-6 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 27: Cluster Cl-6 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 28: Cluster Cl-7 in Clustering-B with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 29: Cluster Cl-7 in Clustering-B with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 7, 8.



Figure 30: Cluster Cl-1 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 31: Cluster Cl-1 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 32: Cluster Cl-2 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 33: Cluster Cl-2 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 34: Cluster Cl-3 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 35: Cluster Cl-3 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 36: Cluster Cl-4 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 37: Cluster Cl-4 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 38: Cluster Cl-5 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 39: Cluster Cl-5 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 40: Cluster Cl-6 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 41: Cluster Cl-6 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 42: Cluster Cl-7 in Clustering-C with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 43: Cluster Cl-7 in Clustering-C with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 9, 10.



Figure 44: Cluster Cl-1 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 45: Cluster Cl-1 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 46: Cluster Cl-2 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 47: Cluster Cl-2 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 48: Cluster Cl-3 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 49: Cluster Cl-3 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 50: Cluster Cl-4 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 51: Cluster Cl-4 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 52: Cluster Cl-5 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 53: Cluster Cl-5 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.


Figure 54: Cluster Cl-6 in Clustering-D with weights based on unnormalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 55: Cluster Cl-6 in Clustering-D with weights based on normalized regressions with arithmetic means (see Subsection 2.6). See Tables 4, 11, 12.



Figure 56: Cluster Cl-1 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 57: Cluster Cl-1 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 58: Cluster Cl-2 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 59: Cluster Cl-2 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 60: Cluster Cl-3 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 61: Cluster Cl-3 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 62: Cluster Cl-4 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 63: Cluster Cl-4 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 64: Cluster Cl-5 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 65: Cluster Cl-5 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 66: Cluster Cl-6 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 67: Cluster Cl-6 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 68: Cluster Cl-7 in Clustering-A with weights based on unnormalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.



Figure 69: Cluster Cl-7 in Clustering-A with weights based on normalized regressions with geometric means (see Subsection 2.6). See Tables 4, 13, 14.