

General Monogamy Relations for Multiqubit W-class States in terms of convex-roof extended negativity of assistance and squared Rényi- α entanglement

Yanying Liang¹, Xiufang Feng¹ and Wei Chen²

¹ School of Mathematics, South China University of Technology, Guangzhou 510641, China

² School of Computer Science and Network Security, Dongguan University of Technology, Dongguan 523808, China

For multipartite entangled states, entanglement monogamy is an important property. We present some new analytical monogamy inequalities satisfied by the x -th power of the dual of convex-roof extended negativity (CREN), namely CREN of Assistance (CRENoA), with $x \geq 2$ and $x \leq 0$ for multiqubit generalized W-class states. We also provide the upper bound of the squared Rényi- α entanglement (SR α E) with α in the region $[(\sqrt{7}-1)/2, (\sqrt{13}-1)/2]$ for multiqubit generalized W-class states.

PACS numbers: 03.67.Mn, 03.65.Ud

I. INTRODUCTION

While classical correlation can be freely shared among parties in multi-party systems, quantum entanglement is restricted in its shareability. If a pair of parties are maximally entangled in multipartite systems, they cannot have any entanglement [1, 2] nor classical correlations [3] with the rest of the system. This restriction of entanglement shareability among multi-party systems is known as the monogamy of entanglement (MoE) [4–9].

The monogamy of entanglement (MoE) is one of the fundamental differences between quantum entanglement and classical correlations that a quantum system entangled with one of the other systems limits its entanglement with the remaining others. For example, MoE is a key ingredient to make quantum cryptography secure because it quantifies how much information an eavesdropper could potentially obtain about the secret key to be extracted [10].

Coffman, Kundu, and Wootters established the first quantitative characterization of the MoE for the squared concurrence (SC) [11–15] in an arbitrary three-qubit quantum state. Another two well-known entanglement measures are convex-roof extended negativity (CREN) [16] and Rényi- α entanglement (R α E) [17]. CREN is a good alternative for MoE without any known example violating its property even in higher-dimensional systems and R α E is the generalization of entanglement of formation. Recently, the general monogamy relations for the x -th power of CREN has been shown for a mixed state $\rho_{A_1 A_2 \dots A_N}$ in a N -qubit system [18],

$$\tilde{N}_{A_1|A_2 \dots A_N}^x \geq \tilde{N}_{A_1 A_2}^x + \dots + \tilde{N}_{A_1 A_N}^x, \quad (1)$$

for $x \geq 2$ and

$$\tilde{N}_{A_1|A_2 \dots A_N}^x < \tilde{N}_{A_1 A_2}^x + \dots + \tilde{N}_{A_1 A_N}^x, \quad (2)$$

for $x \leq 0$. Two years ago, Wei Song and Yan-Kui Bai showed the properties of the squared Rényi- α entanglement (SR α E) and proved that the lower bound of SR α E in an arbitrary N -qubit mixed state [19],

$$E_\alpha^2(\rho_{A_1|A_2 \dots A_N}) \geq E_\alpha^2(\rho_{A_1 A_2}) + \dots + E_\alpha^2(\rho_{A_1 A_N}), \quad (3)$$

where $E_\alpha^2(\rho_{A_1|A_2 \dots A_N})$ quantifies the entanglement in the partition $A_1|A_2 \dots A_N$ and $E_\alpha^2(\rho_{A_1 A_i})$ quantifies the one in two-qubit subsystem $A_1 A_i$ with the order $\alpha \geq (\sqrt{7}-1)/2$.

In this paper, we show the general monogamy relations for the x -th power of CRENoA of generalized multiqubit W-class states. This part provides a more efficient way for MoE. We also prove that the SR α E with the order α ranges in the region $[(\sqrt{7}-1)/2, (\sqrt{13}-1)/2]$ also obeys a general monogamy relation for arbitrary generalized multiqubit W-class states.

II. MONOGAMY OF CONCURRENCE AND CONVEX-ROOF EXTENDED NEGATIVITY

Given a bipartite pure state $|\psi\rangle_{AB}$ in a $d \otimes d'$ ($d \leq d'$) quantum system, its concurrence, $C(|\psi\rangle_{AB})$ is defined as [20]

$$C(|\psi\rangle_{AB}) = \sqrt{2[1 - \text{Tr}(\rho_A^2)]}, \quad (4)$$

where ρ_A is reduced density matrix by tracing over the subsystem B , $\rho_A = \text{Tr}_B(|\psi\rangle_{AB}\langle\psi|)$ (and analogously for ρ_B). For any mixed state ρ_{AB} , the concurrence is given by the minimum average concurrence taken over all decompositions of ρ_{AB} , the so-called convex roof

$$C(\rho_{AB}) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C(|\psi_i\rangle), \quad (5)$$

where the convex roof is notoriously hard to evaluate and therefore it is difficult to determine whether or not an arbitrary state is entangled.

Similarly, the concurrence of assistance (CoA) of any mixed state ρ_{AB} is defined as [21]

$$C_a(\rho_{AB}) = \max_{\{p_i, |\psi_i\rangle\}} \sum_i p_i C(|\psi_i\rangle), \quad (6)$$

where the maximum is taken over all possible pure state decompositions $\{p_i, |\psi_i\rangle\}$ of ρ_{AB} .

Another well-known quantification of bipartite entanglement is convex-roof extended negativity (CREN). For a bipartite mixed state ρ_{AB} , CREN is defined as

$$\tilde{N}(\rho_{AB}) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i N(|\psi_i\rangle), \quad (7)$$

where the minimum is taken over all possible pure state decompositions $\{p_i, |\psi_i\rangle\}$ of ρ_{AB} .

Similar to the duality between concurrence and CoA, we can also define a dual to CREN, namely CRENoA, by taking the maximum value of average negativity over all possible pure state decomposition, i.e.

$$\tilde{\mathcal{N}}_a(\rho_{AB}) = \max_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \mathcal{N}(|\psi_i\rangle), \quad (8)$$

where the maximum is taken over all possible pure state decompositions $\{p_i, |\psi_i\rangle\}$ of ρ_{AB} .

In the following we study the monogamy property of the CRENoA for the n -qubit generalized W-class states $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$ defined by

$$|\psi\rangle = a|000\dots\rangle + b_1|01\dots0\rangle + \dots + b_n|00\dots1\rangle, \quad (9)$$

with $|a|^2 + \sum_{i=1}^n |b_i|^2 = 1$.

Lemma 1. *For n -qubit generalized W-class states (9), we have*

$$\tilde{\mathcal{N}}(\rho_{A_1 A_i}) = \tilde{\mathcal{N}}_a(\rho_{A_1 A_i}), \quad (10)$$

where $\rho_{A_1 A_i} = \text{Tr}_{A_2 \dots A_{i-1} A_{i+1} \dots A_n}(|\psi\rangle\langle\psi|)$.

Proof. We assume $\rho_{A_1 A_i} = |x\rangle_{A_1 A_i} \langle x| + |y\rangle_{A_1 A_i} \langle y|$ [22], where

$$\begin{aligned} |x\rangle_{A_1 A_i} &= a|00\rangle_{A_1 A_i} + b_1|10\rangle_{A_1 A_i} + b_i|01\rangle_{A_1 A_i}, \\ |y\rangle_{A_1 A_i} &= \sqrt{\sum_{k \neq i} |b_k|^2} |00\rangle_{A_1 A_i}. \end{aligned}$$

From the HJW theorem in Ref. [22], for any pure-state decomposition of $\rho_{A_1 A_i} = \sum_{h=1}^r |\phi_h\rangle_{A_1 A_i} \langle\phi_h|$, one has $|\phi_h\rangle_{A_1 A_i} = u_{h1}|x\rangle_{A_1 A_i} + u_{h2}|y\rangle_{A_1 A_i}$ for some $r \times r$ unitary matrices u_{h1} and u_{h2} for each h . Consider the normalized bipartite pure state $|\tilde{\phi}_h\rangle_{A_1 A_i} = |\phi_h\rangle_{A_1 A_i} / \sqrt{p_h}$ with

$p_h = |\langle\phi_h|\phi_h\rangle|$. In Ref. [24], for any bipartite pure state $|\psi\rangle$, one has

$$C(|\psi\rangle) = \mathcal{N}(|\psi\rangle).$$

and combining with the Lemma 1 in Ref. [23], for $|\tilde{\phi}_h\rangle_{A_1 A_i}$, we have

$$\mathcal{N}(|\tilde{\phi}_h\rangle_{A_1 A_i}) = \frac{2}{p_h} |u_{hi}|^2 |b_1| |b_i|.$$

Then combining (7) and (8), we can obtain

$$\begin{aligned} \tilde{\mathcal{N}}(\rho_{A_1 A_i}) &= \min_{\{p_h, |\tilde{\phi}_h\rangle_{A_1 A_i}\}} \sum_h p_h \mathcal{N}(|\tilde{\phi}_h\rangle_{A_1 A_i}) \\ &= \max_{\{p_h, |\tilde{\phi}_h\rangle_{A_1 A_i}\}} \sum_h p_h \mathcal{N}(|\tilde{\phi}_h\rangle_{A_1 A_i}) \\ &= \tilde{\mathcal{N}}_a(\rho_{A_1 A_i}). \end{aligned}$$

□

Theorem 1. *For the n -qubit generalized W-class states $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$, the CRENoA satisfies*

$$\tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \geq \frac{x}{2^x - 1} \sum_{i=1}^{m-1} \tilde{\mathcal{N}}_a^x(\rho_{A_1 A_{j_i}}), \quad (11)$$

where $x \geq 2$ and $\rho_{A_1 A_{j_1} \dots A_{j_{m-1}}}$ is the m -qubit, $2 \leq m \leq n$, reduced density matrix of $|\psi\rangle$.

Proof. For the n -qubit generalized W-class state $|\psi\rangle$, according to the definitions of $\tilde{\mathcal{N}}(\rho)$ and $\tilde{\mathcal{N}}_a(\rho)$, one has $\tilde{\mathcal{N}}_a(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \geq \tilde{\mathcal{N}}(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})$. When $x \geq 2$, we have

$$\begin{aligned} \tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) &\geq \left(\frac{\tilde{\mathcal{N}}_a(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) + \tilde{\mathcal{N}}(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})}{2} \right)^x \\ &= \frac{1}{2^x} \tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \left(1 + \frac{\tilde{\mathcal{N}}(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})}{\tilde{\mathcal{N}}_a(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})} \right)^x \\ &\geq \frac{1}{2^x} \tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \left(1 + x \frac{\tilde{\mathcal{N}}(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})}{\tilde{\mathcal{N}}_a(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})} \right)^x \\ &= \frac{1}{2^x} \tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) + \frac{x}{2^x} \tilde{\mathcal{N}}^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \end{aligned}$$

Here we have used in the first inequality the inequality $a^x \geq (\frac{a+b}{2})^x$ for $a \geq b > 0$ and $x \geq 0$. The second inequality is due to $(1+t)^x \geq 1 + xt^x$ for $x \geq 1$ and $1 \geq t \geq 0$.

Then we have

$$\tilde{\mathcal{N}}_a^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}}) \geq \frac{x}{2^x - 1} \tilde{\mathcal{N}}^x(\rho_{A_1 | A_{j_1} \dots A_{j_{m-1}}})$$

Combining with Lemma 1, we have

$$\begin{aligned}\tilde{\mathcal{N}}_a^x(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) &\geq \frac{x}{2^x-1}\tilde{\mathcal{N}}^x(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) \\ &\geq \frac{x}{2^x-1}\sum_{i=1}^{m-1}\tilde{\mathcal{N}}^x(\rho_{A_1A_{j_i}}) \\ &= \frac{x}{2^x-1}\sum_{i=1}^{m-1}\tilde{\mathcal{N}}_a^x(\rho_{A_1A_{j_i}}).\end{aligned}$$

The second inequality is due to the monogamy relation for the x -th power of CREN (1). \square

Theorem 2. For the n -qubit generalized W -class state $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$ with $\tilde{\mathcal{N}}(\rho_{A_1A_{j_i}}) \neq 0$ for $1 \leq i \leq m-1$, we have

$$\tilde{\mathcal{N}}_a^y(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) < \frac{y}{2^y-1}\sum_{i=1}^{m-1}\tilde{\mathcal{N}}_a^y(\rho_{A_1A_{j_i}}), \quad (12)$$

where $y \leq 0$ and $\rho_{A_1A_{j_1}\dots A_{j_{m-1}}}$ is the m -qubit reduced density matrix as in Theorem 1.

Proof. Similar to the proof of Theorem 1, for $y \leq 0$, we get

$$\tilde{\mathcal{N}}_a^y(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) \leq \frac{y}{2^y-1}\tilde{\mathcal{N}}^y(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}})$$

Combining with Lemma 1, we have

$$\begin{aligned}\tilde{\mathcal{N}}_a^y(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) &\leq \frac{y}{2^y-1}\tilde{\mathcal{N}}^y(\rho_{A_1|A_{j_1}\dots A_{j_{m-1}}}) \\ &< \frac{y}{2^y-1}\sum_{i=1}^{m-1}\tilde{\mathcal{N}}^y(\rho_{A_1A_{j_i}}) \\ &= \frac{y}{2^y-1}\sum_{i=1}^{m-1}\tilde{\mathcal{N}}_a^y(\rho_{A_1A_{j_i}}).\end{aligned}$$

The second inequality is due to the monogamy relation for the x -th power of CREN (2). \square

As an example, consider the 5-qubit generalized W -class states (9) with $a = b_2 = \frac{1}{\sqrt{10}}$, $b_1 = \frac{1}{\sqrt{15}}$, $b_3 = \sqrt{\frac{2}{15}}$, $b_4 = \sqrt{\frac{3}{5}}$. We have

$$\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3}) \geq \sqrt{\frac{x}{2^x-1}} \frac{2}{\sqrt{15}} \sqrt{\left(\frac{1}{\sqrt{10}}\right)^x + \left(\sqrt{\frac{2}{15}}\right)^x}$$

and

$$\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3A_4}) \geq \sqrt{\frac{x}{2^x-1}} \frac{2}{\sqrt{15}} \sqrt{\left(\frac{1}{\sqrt{10}}\right)^x + \left(\sqrt{\frac{2}{15}}\right)^x + \left(\sqrt{\frac{3}{5}}\right)^x}$$

with $x \geq 2$. The optimal lower bounds can be obtained by varying the parameter x , see Fig. 1.

From Fig. 1, one gets that the optimal lower bounds of $\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3})$ and $\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3A_4})$ are 0.203 and 0.385, respectively, attained at $x = 2$ while the lower bounds of each in terms of CoA are given by 0.249 and 0.471 [23]. One can see that choosing CRENoA as a mathematical characterization of MoE is better than choosing CoA for $x \geq 2$.




Fig. 1: solid red line is the lower bound of $\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3})$ and solid blue line is the lower bound of $\tilde{\mathcal{N}}_a(\rho_{A_1|A_2A_3A_4})$ as functions of $x \geq 2$ from our result, red dashed line is the lower bound of $C_a(\rho_{A_1|A_2A_3})$ and blue dashed line is the lower bound of $C_a(\rho_{A_1|A_2A_3A_4})$ as functions of $x \geq 2$ from [23].

III. MONOGAMY OF RÉNYI- α ENTANGLEMENT

Rényi- α entanglement (R α E) is well-defined entanglement measure which is the generalization of entanglement of formation. For a bipartite pure state $|\psi\rangle_{AB}$, the R α E is defined as

$$E_\alpha(|\psi\rangle_{AB}) := S_\alpha(\rho_A) = \frac{1}{1-\alpha} \log_2(\text{tr} \rho_A^\alpha) \quad (13)$$

where the Rényi- α entropy is $S_\alpha(\rho_A) = [\log_2(\sum_i \lambda_i^\alpha)]/(1-\alpha)$ with α being a nonnegative real number and λ_i being the eigenvalue of reduced density matrix ρ_A . For a bipartite mixed state ρ_{AB} , the R α E is defined via the convex-roof extension

$$E_\alpha(\rho_{AB}) = \min \sum_i p_i E_\alpha(|\psi_i\rangle_{AB}) \quad (14)$$

where the minimum is taken over all possible pure state decompositions of $\rho_{AB} = \sum_i p_i |\psi_i\rangle_{AB} \langle \psi_i|$. In particular, for a two-qubit mixed state, the R α E with $\alpha \geq 1$ has an analytical formula which is expressed as a function of the SC [24]

$$E_\alpha(\rho_{AB}) = f_\alpha[C^2(\rho_{AB})] \quad (15)$$

where the function $f_\alpha(x)$ has the form

$$f_\alpha(x) = \frac{1}{1-\alpha} \log_2 \left[\left(\frac{1-\sqrt{1-x}}{2} \right)^\alpha + \left(\frac{1+\sqrt{1-x}}{2} \right)^\alpha \right]. \quad (16)$$

Recently, Wang *et al* further proved that the formula in (14) holds for the order $\alpha \geq (\sqrt{7}-1)/2 \simeq 0.823$ [25].

From Theorem 2 in Ref. [19], one has that for a bipartite $2 \otimes d$ mixed state ρ_{AC} , the Rényi- α entanglement has an analytical expression

$$E_\alpha(\rho_{AC}) = f_\alpha[C^2(\rho_{AC})] \quad (17)$$

where the order α ranges in the region $[(\sqrt{7} - 1)/2, (\sqrt{13} - 1)/2]$.

Theorem 3. For the n -qubit generalized W-class states $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$, we have

$$E_\alpha(|\psi\rangle_{A_1|A_2\dots A_n}) \leq \sum_{i=2}^n E_\alpha(\rho_{A_1 A_i}), \quad (18)$$

where $\rho_{A_1 A_i}$, $2 \leq i \leq n$, is the 2-qubit reduced density matrix of $|\psi\rangle$ and the order α ranges in the region $[(\sqrt{7} - 1)/2, (\sqrt{13} - 1)/2]$.

Proof. For the n -qubit generalized W-class states $|\psi\rangle$, we have

$$\begin{aligned} E_\alpha(|\psi\rangle_{A_1|A_2\dots A_n}) &= f_\alpha(C^2(|\psi\rangle_{A_1|A_2\dots A_n})) \\ &= f_\alpha\left(\sum_{i=2}^n C^2(\rho_{A_1 A_i})\right) \\ &\leq \sum_{i=2}^n f_\alpha(C^2(\rho_{A_1 A_i})) \\ &= \sum_{i=2}^n E_\alpha(\rho_{A_1 A_i}), \end{aligned}$$

where $f_\alpha(x) = \frac{1}{1-\alpha} \log_2 \left[\left(\frac{1-\sqrt{1-x}}{2} \right)^\alpha + \left(\frac{1+\sqrt{1-x}}{2} \right)^\alpha \right]$. We have used in the first and last equalities that the entanglement of formation obeys the relation (19). The second equality is due to the fact that $C^2(|\psi\rangle_{A_1\dots A_n}) = \sum_{i=2}^n C^2(\rho_{A_1 A_i})$. The inequality is due to the fact that the Rényi- α entanglement $E_\alpha(C^2)$ with $\alpha \in [(\sqrt{7} - 1)/2, (\sqrt{13} - 1)/2]$ is monotonic increasing and concave as a function of the squared concurrence C^2 [19]. \square

Next we will present an upper bound of SR α E. Before giving the result, we consider the following lemma.

Lemma 2. [19] Let $\psi_{A_1\dots A_n}$ be a generalized W-class state in (11). For any m -qubit subsystems $A_1 A_{j_1} \dots A_{j_{m-1}}$ of $A_1 \dots A_n$ with $2 \leq m \leq n - 1$, the reduced density matrix $\rho_{A_1 A_{j_1} \dots A_{j_{m-1}}}$ of $\psi_{A_1\dots A_n}$ is a mixture of a m -qubit generalized W-class state and vacuum.

In the following, we assume the m -qubit subsystems $A_1 A_{j_1} \dots A_{j_{m-1}}$ of $A_1 \dots A_n$ with $2 \leq m \leq n - 1$ is exactly $A_1 \dots A_n$. Then we can have the result below.

Theorem 4. For the n -qubit generalized W-class states $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$, we have

$$E_\alpha^2(\rho_{A_1|A_2\dots A_n}) \leq (n-1) \sum_{i=2}^n E_\alpha^2(\rho_{A_1 A_i}), \quad (19)$$

where $E_\alpha^2(\rho_{A_1|A_2\dots A_n})$ quantifies the entanglement in the partition $A_1|A_2\dots A_n$ and $E_\alpha^2(\rho_{A_1 A_i})$ quantifies the one in two-qubit subsystem $A_1 A_i$ with the order $\alpha \in [(\sqrt{7} - 1)/2, (\sqrt{13} - 1)/2]$.

Proof. We first consider the monogamy relation in an n -qubit pure state $\psi_{A_1 A_2 \dots A_n}$. Thus we can obtain

$$\begin{aligned} E_\alpha^2(|\psi\rangle_{A_1|A_2\dots A_n}) &\leq \left(\sum_{i=2}^n E_\alpha(\rho_{A_1 A_i}) \right)^2 \\ &\leq \left(\sum_{i=2}^n 1^2 \right) \left(\sum_{i=2}^n E_\alpha^2(\rho_{A_1 A_i}) \right) \\ &= (n-1) \left(\sum_{i=2}^n E_\alpha^2(\rho_{A_1 A_i}) \right), \end{aligned}$$

where in the first inequality we have used Theorem 3 and $a^2 \leq b^2$ for $0 \leq a \leq b$, and in the second inequality we have used the Cauchy-Schwarz inequality.

Next from Lemma 2, we consider $\rho_{A_1 \dots A_n}$ is a mixture of a n -qubit generalized W-class state and vacuum. Then since we have the pure decomposition of $\rho_{A_1 \dots A_n}$,

$$\rho_{A_1 A_2 \dots A_n} = \sum_j p_j |\psi_j\rangle_{A_1 A_2 \dots A_n} \langle \psi_j|,$$

Thus, we can obtain

$$\begin{aligned} E_\alpha^2(\rho_{A_1|A_2\dots A_n}) &= \left[\sum_j p_j E_\alpha(|\psi_j\rangle_{A_1|A_2\dots A_n}) \right]^2 \\ &\leq \left[\sum_j p_j \left(\sum_{i=2}^n E_\alpha(\rho_{A_1 A_i}) \right) \right]^2 \\ &= \left[\sum_{i=2}^n \left(\sum_j p_j E_\alpha(\rho_{A_1 A_i}) \right) \right]^2 \\ &\leq \left(\sum_{i=2}^n 1^2 \right) \left[\sum_{i=2}^n \left(\sum_j p_j E_\alpha(\rho_{A_1 A_i}) \right) \right]^2 \\ &= (n-1) \left(\sum_{i=2}^n E_\alpha^2(\rho_{A_1 A_i}) \right), \end{aligned}$$

where in the first inequality we have used Theorem 3 and $a^2 \leq b^2$ for $0 \leq a \leq b$, and in the second inequality we have used the Cauchy-Schwarz inequality. The last equality is due to $\sum_j p_j = 1$. \square

As an example, we still consider the 5-qubit generalized W-class states (9) with $a = b_2 = \frac{1}{\sqrt{10}}$, $b_1 = \frac{1}{\sqrt{15}}$, $b_3 = \sqrt{\frac{2}{15}}$, $b_4 = \sqrt{\frac{3}{5}}$. We have

$$E_\alpha^2(\rho_{A_1|A_2 A_3}) \leq 2 (E_\alpha^2(\rho_{A_1|A_2}) + E_\alpha^2(\rho_{A_1|A_3}))$$

and

$$E_\alpha^2(\rho_{A_1|A_2 A_3 A_4}) \leq 3 (E_\alpha^2(\rho_{A_1|A_2}) + E_\alpha^2(\rho_{A_1|A_3}) + E_\alpha^2(\rho_{A_1|A_4}))$$

where

$$E_\alpha(\rho_{A_1|A_2}) = \frac{1}{1-\alpha} \log_2 \left[\left(\frac{1-\sqrt{\frac{73}{75}}}{2} \right)^\alpha + \left(\frac{1+\sqrt{\frac{73}{75}}}{2} \right)^\alpha \right]$$

$$E_{\alpha}(\rho_{A_1|A_3}) = \frac{1}{1-\alpha} \log_2 \left[\left(\frac{1-\sqrt{\frac{217}{225}}}{2} \right)^{\alpha} + \left(\frac{1+\sqrt{\frac{217}{225}}}{2} \right)^{\alpha} \right]$$

and

$$E_{\alpha}(\rho_{A_1|A_4}) = \frac{1}{1-\alpha} \log_2 \left[\left(\frac{1-\sqrt{\frac{63}{75}}}{2} \right)^{\alpha} + \left(\frac{1+\sqrt{\frac{63}{75}}}{2} \right)^{\alpha} \right]$$

with the order $\alpha \in [(\sqrt{7}-1)/2, (\sqrt{13}-1)/2]$. See Fig. 2.




Fig. 2: solid red line is the upper bound of $E_{\alpha}^2(\rho_{A_1|A_2A_3})$ and solid blue line is the upper bound of $E_{\alpha}^2(\rho_{A_1|A_2A_3A_4})$ as functions of α when α ranges in the region $[(\sqrt{7}-1)/2, 0.99]$. When α ranges in the region $[1.001, (\sqrt{13}-1)/2]$, as the red dashed line and the blue dashed line show, we do not have an upper bound of $E_{\alpha}^2(\rho_{A_1|A_2A_3})$ and $E_{\alpha}^2(\rho_{A_1|A_2A_3A_4})$ in this example.

From Fig.2, one gets that the optimal upper bounds of $E_{\alpha}^2(\rho_{A_1|A_2A_3})$ and $E_{\alpha}^2(\rho_{A_1|A_2A_3A_4})$ are 0.02334 and 0.24211 attained at $\alpha = 0.971$ when $\alpha \in [(\sqrt{7}-1)/2, 0.99]$. This upper bounds can be easily generalized to arbitrary n -qubit generalized W-class states $|\psi\rangle \in H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_n}$.

IV. CONCLUSIONS AND REMARKS

We have investigated the monogamy relations of multi-qubit generalized W-class states in terms of CRENoA and SR α E. We have proved that the monogamy inequality of x -th power for CRENoA when $x \geq 2$ and $x \leq 0$. Our result shows that choosing CRENoA as a mathematical characterization of the monogamy of entanglement is better than choosing CoA for $x \geq 2$. We also show the monogamy inequality for SR α E when α ranges in the region $[(\sqrt{7}-1)/2, (\sqrt{13}-1)/2]$. We can find the optimal upper bound for $E_{\alpha}^2(\rho_{A_1|A_2\dots A_n})$ when the order $\alpha \in [(\sqrt{7}-1)/2, (\sqrt{13}-1)/2]$ by using our approach in Theorem 4. It is still an open problem to be answered that whether there exists the monogamy inequality for SR α E when $\alpha \geq (\sqrt{13}-1)/2$ in generalized W-class states.

Acknowledgments

This work is supported by the NSFC 11571119 and NSFC 11475178.

-
- [1] V. Coffman, J. Kundu and W. K. Wootters, Phys. Rev. A **61**, 052306 (2000).
 - [2] T. Osborne and F. Verstraete, Phys. Rev. Lett. **96**, 220503 (2006).
 - [3] M. Koashi and A. Winter, Phys. Rev. A **69**, 022309 (2004).
 - [4] F. Mintert, M. Kuš, and A. Buchleitner, Phys. Rev. Lett. **92**, 167902 (2004).
 - [5] K. Chen, S. Albeverio, and S. M. Fei, Phys. Rev. Lett. **95**, 040504 (2005).
 - [6] H. P. Breuer, J. Phys. A: Math. Gen. **39**, 11847 (2006).
 - [7] H. P. Breuer, Phys. Rev. Lett. **97**, 080501 (2006).
 - [8] J. I. de Vicente, Phys. Rev. A **75**, 052320 (2007).
 - [9] C. J. Zhang, Y. S. Zhang, S. Zhang, and G. C. Guo, Phys. Rev. A **76**, 012334 (2007).
 - [10] J. M. Renes and M. Grassl, Phys. Rev. A **74**, 022317 (2006).
 - [11] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [12] T. J. Osborne, and F. Verstraete, Phys. Rev. Lett. **96**, 220503 (2006).
 - [13] Y. K. Bai, M. Y. Ye, and Z. D. Wang, Phys. Rev. A **80**, 044301(2009).
 - [14] X. N. Zhu, and S. M. Fei, Physical Review A **90**, 024304 (2014)
 - [15] J. S. Kim, Phys. Rev. A **90**, 062306 (2014).
 - [16] S. Lee, D. P. Chi, S. D. Oh, and J. Kim, Phys. Rev. A **68**, 062304 (2003).
 - [17] R. Horodecki, P. Horodecki, and M. Horodecki, Phys. Rev. A **210**, 377 (1996).
 - [18] Y. Luo, Y. M. Li, Phys. Rev. A **362**, 511-520 (2015).
 - [19] W. Song, Y. K. Bai, M. Yang, M. Yang and Z. L. Cao, Phys. Rev. A **93**, 022306 (2015).
 - [20] P. Rungta, V. Bužek, C. M. Caves, M. Hillery, and G. J. Milburn, Phys. Rev. A **64**, 042315 (2001).
 - [21] C. S. Yu, and H. S. Song, Phys. Rev. A **77**, 032329 (2008).
 - [22] J. S. Kim, A. Das and B. C. Sanders, Phys. Rev. A **79**, 012329 (2008).
 - [23] X. N. Zhu, S. M. Fei, Quant. Inf. Process **16**, 53 (2017).
 - [24] J. S. Kim, and B. C. Sanders, Phys. Rev. A **43**, 445305 (2010).
 - [25] Y. X. Wang, L. Z. Mu, and V. Vedral, and H. Fen, Phys. Rev. A **93**, 022324 (2016).