# Thorup-Zwick Emulators are Universally Optimal Hopsets\*

Shang-En Huang University of Michigan

Seth Pettie University of Michigan

May 2, 2017

#### Abstract

A  $(\beta,\epsilon)$ -hopset is, informally, a weighted edge set that, when added to a graph, allows one to get from point a to point b using a path with at most  $\beta$  edges ("hops") and length  $(1+\epsilon)$  dist(a,b). In this paper we observe that Thorup and Zwick's sublinear additive emulators are also actually  $(O(k/\epsilon)^k,\epsilon)$ -hopsets for every  $\epsilon>0$ , and that with a small change to the Thorup-Zwick construction, the size of the hopset can be made  $O(n^{1+\frac{1}{2^{k+1}-1}})$ . As corollaries, we also shave "k" factors off the size of Thorup and Zwick's [20] sublinear additive emulators and the sparsest known  $(1+\epsilon,O(k/\epsilon)^{k-1})$ -spanners, due to Abboud, Bodwin, and Pettie [1].

## 1 Introduction

Let G = (V, E, w) be a weighted undirected graph. Define  $\operatorname{dist}_{G}^{(\beta)}(u, v)$  to be the length of the shortest path from u to v in G that uses at most  $\beta$  edges, or "hops." Whereas  $\operatorname{dist}_{G}^{(\infty)}$  is a metric,  $\operatorname{dist}_{G}^{(\beta)}$  is not in general. A set  $H \subset \binom{V}{2}$  of weighted edges is called a  $(\beta, \epsilon)$ -hopset if for every  $u, v \in V$ ,

$$\operatorname{dist}_{G}(u, v) \leq \operatorname{dist}_{G \cup H}^{(\beta)}(u, v) \leq (1 + \epsilon) \operatorname{dist}_{G}(u, v).$$

**Background.** Cohen [7] formally defined the notion of a hopset, but the idea was latent in earlier work [21, 14, 6, 18]. Cohen's  $(\beta, \epsilon)$ -hopset had size  $O(n^{1+1/\kappa} \log n)$  and  $\beta = (\epsilon^{-1} \log n)^{O(\log \kappa)}$ . Elkin and Neiman [9] showed that a constant hopbound  $\beta$  suffices (when  $\kappa, \epsilon$  are constants). In particular, their hopset has size  $O(n^{1+1/\kappa} \log n \log \kappa)$  and  $\beta = O(\epsilon^{-1} \log \kappa)^{\log \kappa}$ . Abboud, Bodwin, and Pettie [1] recently proved that the tradeoffs of [9] are essentially optimal: for any integer k, any hopset of size  $n^{1+\frac{1}{2k+1-1}-\delta}$  must have  $\beta = \Omega(c_k/\epsilon^{k+1})$ , where  $c_k$  is a constant depending only on k.<sup>1</sup> There are other constructions of hopsets [5, 11, 12, 16] that are designed for parallel or dynamic environments; their tradeoffs (between hopset size and hopbound) are worse than [7, 9] and the ones presented here. See Table 1.

Hopsets, Emulators, and Spanners. Recall that G is an undirected graph, possibly weighted. A spanner is a subgraph of G such that  $\operatorname{dist}_H(u,v) \leq f(\operatorname{dist}_G(u,v))$  for some nondecreasing stretch function f. An emulator of an unweighted graph G is a weighted edge set H such that  $\operatorname{dist}_H(u,v) \in [\operatorname{dist}_G(u,v), f(\operatorname{dist}_G(u,v))]$ . Syntactically, the definition of hopsets is closely related to emulators. The difference is that hopsets have a hopbound constraint but are allowed to use original edges in G whereas emulators must use only H. The purpose of emulators is to compress the graph metric  $\operatorname{dist}_G$ : ideally  $|H| \ll |E(G)|$ . Historically, the literature on hopset constructions [7, 9] has been noticeably more complex than those of spanners and emulators, many

<sup>\*</sup>Supported by NSF Grants CCF-1514383 and CCF-1637546.

Note that setting  $\kappa = 2^{k+1} - 1$  in the Elkin-Neiman construction gives  $\beta = O(k/\epsilon)^k$ , where  $\log \kappa = \lfloor \log \kappa \rfloor = k$ . Thus, saving any  $\delta$  in the exponent of the hopset increases  $\beta$  significantly. In general, the statement of [9] obscures the nature of the tradeoff: there are *not* distinct tradeoffs for each  $\kappa \in \{1, 2, 3, \ldots\}$ , but only for  $\kappa \in \{1, 3, 7, \ldots, 2^{k+1} - 1, \ldots\}$ .

Authors	Size	Hopbound	Stretch
Klein and Subramanian [14]	O(n)	$O(\sqrt{n}\log n)$	1
Thorup and Zwick [19]	$O(\kappa n^{1+1/\kappa})$	2	$2\kappa - 1$
Cohen [7]	$O(n^{1+\frac{1}{\kappa}} \cdot \log n)$	$((\log n)/\epsilon)^{O(\log \kappa)}$	$1+\epsilon$
Elkin and Neiman [9]	$O(n^{1+\frac{1}{\kappa}}\log n\log \kappa)$	$O((\log \kappa)/\epsilon)^{\log \kappa}$	$1 + \epsilon$
Abboud, Bodwin, and Pettie [1]	$n^{1+\frac{1}{2^{k+1}-1}-\delta} \longrightarrow$	$\Omega(c_k/\epsilon^{k+1})$	$1+\epsilon$
New	$O\left(n^{1+\frac{1}{2^{k+1}-1}}\right)$	$O(k/\epsilon)^k$	$1+\epsilon$

Table 1: Tradeoffs between size and hopbound of previous hopsets. Fix the parameter  $\kappa = 2^{k+1} - 1$  to compare [7, 9] against the lower bound [1] and the new result.

of which [3, 2, 8, 20, 4, 15, 1] are quite elegant. Our goal in this work is to demonstrate that there is nothing *intrinsically* complex about hopsets, and that a very simple construction improves on all prior constructions and matches the Abboud-Bodwin-Pettie lower bound.

New Results. Thorup and Zwick [20] designed their emulator for unweighted graphs, and proved that it has size  $O(kn^{1+\frac{1}{2^k+1}-1})$  and a sublinear additive stretch function  $f(d)=d+O(kd^{1-1/k})$ . In this paper we show that the Thorup-Zwick emulator, when applied to a weighted graph, produces a  $(\beta, \epsilon)$ -hopset that achieves every point on the Abboud-Bodwin-Pettie [1] lower bound tradeoff curve. Moreover, with two subtle modifications to the construction, we can reduce the size to  $O(n^{1+\frac{1}{2^k+1}-1})$ , shaving off a factor k. Our technique also applies to other constructions, and as corollaries we improve the size of Thorup and Zwick's emulator [20] and Abboud, Bodwin, and Pettie's  $(1+\epsilon,\beta)$ -spanners.<sup>2</sup>

**Theorem 1.** Fix any weighted graph G and integer  $k \geq 1$ . There is a  $(\beta, \epsilon)$ -hopset for G with size  $O(n^{1+\frac{1}{2^{k+1}-1}})$  and  $\beta = 2\left(\frac{(4+o(1))k}{\epsilon}\right)^k$ .

**Theorem 2.** (cf. [20]) Fix any unweighted graph G and integer  $k \ge 1$ . There is a sublinear additive emulator H for G with size  $O(n^{1+\frac{1}{2^{k+1}-1}})$  and stretch function  $f(d) = d + (4 + o(1))kd^{1-1/k}$ .

**Theorem 3.** (cf. [1]) Fix any unweighted graph G, integer  $k \ge 1$ , and real  $\epsilon > 0$ . There is a  $(1 + \epsilon, ((4 + o(1))k/\epsilon)^{k-1})$ -spanner H for G with size  $O((k/\epsilon)^{h}n^{1 + \frac{1}{2^{k+1}-1}})$ , where  $h = \frac{3 \cdot 2^{k-1} - (k+2)}{2^{k+1}-1} < 3/4$ .

Remark 1. In recent and independent technical report, Elkin and Neiman [10] also observed that Thorup and Zwick's emulator yields an essentially optimal hopset. They proposed a modification to Thorup and Zwick's construction that reduces the size to  $O(n^{1+\frac{1}{2^{k+1}-1}})$  (eliminating a factor k), but increases the hopbound  $\beta$  from  $O(k/\epsilon)^k$  to  $O((k+1)/\epsilon)^{k+1}$ . For example, their technique does not imply any of the improvements found in Theorems 1, 2, or 3.

## 2 The Hopset Construction

In this section, we present the construction of the hopset based on Thorup and Zwick's emulator [20], then analyze its size, stretch, and hopbound.

The construction is parameterized by an integer  $k \geq 1$  and a set  $\{q_i\}$  of sampling probabilities. Let  $V = V_0 \supseteq V_1 \supseteq V_2 \supseteq \cdots \supseteq V_k \supseteq V_{k+1} = \emptyset$  be the vertex sets in each layer. For each  $i \in [0, k)$ , each vertex in  $V_i$  is independently promoted to  $V_{i+1}$  with probability  $q_{i+1}/q_i$ . Thus  $E[|V_i|] = nq_i$ . For each vertex  $v \in V$ 

<sup>&</sup>lt;sup>2</sup>A  $(1 + \epsilon, \beta)$ -spanner of an unweighted graph is one with stretch function  $f(d) = (1 + \epsilon)d + \beta$ .

and  $i \in [1, k]$ , define  $p_i(v)$  to be any vertex in  $V_i$  such that  $\operatorname{dist}_G(v, p_i(v)) = \operatorname{dist}_G(v, V_i)$ . For any vertex  $v \in V_i \setminus V_{i+1}$ , define  $\mathcal{B}(v)$  to be:

$$\mathcal{B}(v) = \{ u \in V_i \mid \operatorname{dist}_G(v, u) < \operatorname{dist}_G(v, p_{i+1}(v)) \}$$

Note that  $p_{k+1}(v)$  does not exist; by convention  $\operatorname{dist}_G(v, p_{k+1}(v)) = \infty$ . The hopset is defined to be  $H = E_0 \cup E_1 \cup \cdots \cup E_k$ , where

$$E_i = \bigcup_{v \in V_i \setminus V_{i+1}} \{ (v, u) \mid u \in \mathcal{B}(v) \cup \{ p_{i+1}(v) \} \}.$$

The length of an edge in H is always the distance between its endpoints. This concludes the description of the construction.

### 2.1 Size Analysis

The expected size of  $E_i$  is at most  $\mathrm{E}[|V_i|](q_i/q_{i+1}) = nq_i^2/q_{i+1}$ , for each  $i \in [0,k)$ , and is  $(nq_k)^2$  if i=k. Following Pettie [17], we choose  $\{q_i\}$  such that the layers of the hopset have geometrically decaying sizes. Setting  $q_i = n^{-\frac{2^i-1}{2^k+1-1}} \cdot 2^{-2^i-i+1}$ , the expected size of  $E_i$ , for  $i \in [0,k)$ , is

$$\begin{split} nq_i^2/q_{i+1} &= n \cdot \left(n^{-\frac{2^{i}-1}{2^{k+1}-1}} \cdot 2^{-2^{i}-i+1}\right)^2 \bigg/ \left(n^{-\frac{2^{i+1}-1}{2^{k+1}-1}} \cdot 2^{-2^{i+1}-i}\right) \\ &= n^{1-\frac{2^{i+1}-2}{2^{k+1}-1} + \frac{2^{i+1}-1}{2^{k+1}-1}} \cdot 2^{-2^{i+1}-2i+2-(-2^{i+1}-i)} \\ &= n^{1+\frac{1}{2^{k+1}-1}} \cdot 2^{-i+2}. \end{split}$$

The expected size of  $E_k$  is

$$(nq_k)^2 = n^2 \cdot \left(n^{-\frac{2^k - 1}{2^{k+1} - 1}} \cdot 2^{-2^k - k + 1}\right)^2$$
$$= n^{1 + \frac{1}{2^{k+1} - 1}} 2^{-2^{k+1} - 2k + 2} \ll n^{1 + \frac{1}{2^{k+1} - 1}} \cdot 2^{-k + 2},$$

so the expected size of H is at most

$$\sum_{i=0}^{k} E[|E_i|] \le n^{1 + \frac{1}{2^{k+1} - 1}} \left( \sum_{i=0}^{k} 2^{-i+2} \right) = O(n^{1 + \frac{1}{2^{k+1} - 1}}).$$

#### 2.2 Stretch and Hopbound Analysis

Let us first give an informal sketch of the analysis. Let a,b be vertices. Choose an integer  $r \geq 2$ , and imagine dividing up the shortest a-b path into  $r^k$  intervals of length  $\mu = \operatorname{dist}_G(a,b)/r^k$ , where  $\mu$  defines one "unit" of length. Once r and  $\mu$  are fixed we prove that given any two vertices u,v at distance at most  $r^i\mu$ , there is either an  $h_i$ -hop path from u to v with additive stretch  $O(ir^{i-1}) \cdot \mu$ , or there is an  $h_i$ -hop path from u to a  $V_{i+1}$ -vertex with length  $(r^i + O(ir^{i-1})) \cdot \mu$ . Of course, when i = k the set  $V_{k+1} = \emptyset$  is empty, so we cannot be in the second case. Since, by definition of  $\mu$ ,  $\operatorname{dist}_G(a,b) \leq r^k\mu$ , there must be an  $h_k$ -hop path with additive stretch  $O(kr^{k-1}) \cdot \mu$ . In order for this stretch to be  $\epsilon \operatorname{dist}_G(a,b)$  we must set  $r = \Theta(k/\epsilon)$ .

So, to recap, the integer parameter  $r = \Theta(k/\epsilon)$  depends on the desired stretch  $\epsilon$ , and r determines the hopcount sequence  $(h_i)$ , which is defined inductively as follows.

$$h_0 = 1,$$
  
 $h_i = (r+1)h_{i-1} + r$  for  $i \in [1, k].$ 

The parameter  $\beta$  of the hopset is exactly  $h_k$ . It is straightforward to show that  $h_k < 2(r+1)^k$ . Once r and  $(h_i)$  are fixed, Theorem 4 is proved by induction.




Figure 1: The two cases depending on whether  $(u', v') \in H$  or not. The first case leads to (i) and the second case leads to (ii) in the statement of Theorem 4.

**Theorem 4.** For any fixed real  $\mu$  (the "unit"), for all  $i \in [0, k]$  and any pair  $u, v \in V$  such that  $\operatorname{dist}_G(u, v) \leq r^i \mu$ , at least one of the following statements holds.

(i) 
$$\operatorname{dist}_{G \cup H}^{(h_i)}(u, v) \le \operatorname{dist}_G(u, v) + ((r+4)^i - r^i)\mu$$
,

(ii) There exists 
$$u_{i+1} \in V_{i+1}$$
 such that  $\operatorname{dist}_{G \cup H}^{(h_i)}(u, u_{i+1}) \leq (r+4)^i \mu$ .

Proof. The proof is by induction on i. In the base case i=0 and  $h_0=1$ . Let  $u,v\in V$  with  $\mathrm{dist}_G(u,v)\leq r^0\mu=\mu$ . If  $(u,v)\in H$  then  $\mathrm{dist}_{G\cup H}^{(1)}(u,v)=\mathrm{dist}_G(u,v)$  so (i) holds. Otherwise,  $(u,v)\notin H$ , meaning  $v\notin \mathcal{B}(u)$ . If  $u\in V_0\setminus V_1$  then  $\mathrm{dist}_{G\cup H}^{(1)}(u,p_1(u))\leq \mathrm{dist}_G(u,v)\leq \mu$ , and if  $u\in V_1$  then  $p_1(u)=u$ , so  $\mathrm{dist}_{G\cup H}^{(1)}(u,p_1(u))=0$ . In either case, (ii) holds.

Now assume i > 0. Consider vertices  $u, v \in V$  with  $\operatorname{dist}_G(u, v) \leq r^i \mu$  and let P be a shortest u-v path in G. Then, as shown in Figure 1, we partition P into at most 2r-1 segments  $\langle u_0 = u, u_1 \rangle$ ,  $\langle u_1, u_2 \rangle$ , ...,  $\langle u_{\ell-1}, u_{\ell} = v \rangle$  as follows. Starting at  $u_0 = u$ , we pick  $u_1$  to be the farthest vertex on P such that  $\operatorname{dist}_G(u_0, u_1) \leq r^{i-1}\mu$ , and let  $(u_1, u_2)$  be the next edge on the path.<sup>3</sup> Repeat the process until we reach  $u_{\ell} = v$ , oscillating between selecting segments that have length at most  $r^{i-1}\mu$  and single edges.

- Multi-hop segment: the shortest path from  $u_s$  to  $u_{s+1}$  satisfies  $\operatorname{dist}_G(u_s, u_{s+1}) \leq r^{i-1}\mu$ .
- Single-hop segment: the segment is actually an edge  $(u_s, u_{s+1}) \in E$ .

By the induction hypothesis, each multi-hop segment satisfies (i) or (ii) within  $h_{i-1}$  hops. Moreover, in each greedy iteration the sum of the lengths from picked multi-hop segment and immediately followed single-hop segment is strictly greater than  $r^i\mu$  except the last one. Therefore, by the pigeonhole principle, there are at most r multi-hop segments on P and at most r-1 single-hop segments on P.

If condition (i) holds for all multi-hop segments, then in at most  $rh_{i-1} + r - 1 \le h_i$  hops,

$$\begin{aligned} \operatorname{dist}_{G \cup H}^{(h_i)}(u, v) &\leq \operatorname{dist}_{G}(u, v) + r((r+4)^{i-1} - r^{i-1})\mu \\ &\leq \operatorname{dist}_{G}(u, v) + ((r+4)^{i} - r^{i})\mu, \end{aligned}$$

<sup>&</sup>lt;sup>3</sup>Note that if the first edge has length more than  $r^{i-1}\mu$ , then  $u_1=u_0$ .

and condition (i) holds for P.

Otherwise, condition (i) does not hold for at least one multi-hop segment. Consider the first multi-hop segment  $\langle u_{j_1}, u_{j_1+1} \rangle$  and the last multi-hop segment  $\langle u_{j_2-1}, u_{j_2} \rangle$  that do not satisfy condition (i). By condition (ii), there exist u' and  $v' \in V_i$  satisfying

$$\operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') \leq (r+4)^{i-1}\mu$$
  
$$\operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_2}, v') \leq (r+4)^{i-1}\mu.$$

Now we have two cases depending on whether  $(u', v') \in H$  or not. If  $(u', v') \in H$ , then by the triangle inequality, we can get from  $u_{j_1}$  to  $u_{j_2}$  with  $2h_{i-1} + 1$  hops and additive stretch

$$\operatorname{dist}_{G \cup H}^{(2h_{i-1}+1)}(u_{j_1}, u_{j_2}) - \operatorname{dist}_{G}(u_{j_1}, u_{j_2}) \leq \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') + \operatorname{dist}_{H}^{(1)}(u', v') + \operatorname{dist}_{G \cup H}^{(h_{i-1})}(v', u_{j_2}) - \operatorname{dist}_{G}(u_{j_1}, u_{j_2})$$

$$\leq 2 \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') + 2 \operatorname{dist}_{G \cup H}^{(h_{i-1})}(v', u_{j_2})$$

$$\leq 4(r+4)^{i-1}\mu.$$

We know there are a total of at most r-1 multi-hop segments satisfying condition (i). Hence, within at most  $(r-1)h_{i-1}+r-1+2h_{i-1}+1 \le h_i$  hops, we can get from u to v with additive stretch

$$\begin{aligned} \operatorname{dist}_{G \cup H}^{(h_i)}(u,v) - \operatorname{dist}_G(u,v) &\leq (r-1)((r+4)^{i-1} - r^{i-1})\mu + \operatorname{dist}_{G \cup H}^{(2h_{i-1}+1)}(u_{j_1},u_{j_2}) - \operatorname{dist}_G(u_{j_1},u_{j_2}) \\ &\leq \left[ (r-1)((r+4)^{i-1} - r^{i-1}) + 4(r+4)^{i-1} \right]\mu \\ &= \left[ (r+3)(r+4)^{i-1} - r^i + r^{i-1} \right]\mu \\ &\leq ((r+4)^i - r^i)\mu \end{aligned} \qquad (r^{i-1} \leq (r+4)^{i-1})$$

and condition (i) holds for P in this case.

On the other hand, suppose that  $(u',v') \notin H$ . Since both  $u',v' \in V_i$  but  $(u',v') \notin H$ , we know that  $u'' = p_{i+1}(u') \in V_{i+1}$  must exist with  $\operatorname{dist}_H^{(1)}(u',u'') \leq \operatorname{dist}_G(u',v')$ . Hence, we can get from  $u_{j_1}$  to u'' via an  $(h_{i-1}+1)$ -hop path with length

$$\begin{aligned}
\operatorname{dist}_{G \cup H}^{(h_{i-1}+1)}(u_{j_1}, u'') &\leq \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') + \operatorname{dist}_{H}^{(1)}(u', u'') \\
&\leq \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') + \operatorname{dist}_{G}(u', v') \\
&\leq 2 \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_1}, u') + \operatorname{dist}_{G}(u_{j_1}, u_{j_2}) + \operatorname{dist}_{G \cup H}^{(h_{i-1})}(u_{j_2}, v') \\
&\leq 3(r+4)^{i-1}\mu + \operatorname{dist}_{G}(u_{j_1}, u_{j_2}).
\end{aligned}$$

Similar to the previous case, there are at most r-1 multi-hop segments appeared before  $u_{j_1}$ , and all of them are satisfying condition (i). Hence, the surplus

$$\operatorname{dist}_{G \cup H}^{((r-1)h_{i-1}+r-1)}(u, u_{j_1}) \leq \operatorname{dist}_G(u, u_{j_1}) + (r-1)((r+4)^{i-1} - r^{i-1})\mu.$$

Therefore, in at most  $(r-1)h_{i-1} + r - 1 + h_{i-1} + 1 \le h_i$  hops,

$$\operatorname{dist}_{G \cup H}^{(h_i)}(u, u'') \leq \operatorname{dist}_{G \cup H}^{((r-1)h_{i-1}+r-1)}(u, u_{j_1}) + \operatorname{dist}_{G \cup H}^{(h_{i-1}+1)}(u_{j_1}, u'')$$

$$\leq \left[ (r-1)((r+4)^{i-1} - r^{i-1}) + 3(r+4)^{i-1} \right] \mu + \operatorname{dist}_{G}(u, u_{j_2})$$

$$\leq \left[ (r+2)(r+4)^{i-1} - r^{i} + r^{i-1} \right] \mu + \operatorname{dist}_{G}(u, u_{j_2})$$

$$\leq \left[ (r+4)^{i} - r^{i} \right] \mu + \operatorname{dist}_{G}(u, u_{j_2}) \qquad (r^{i-1} \leq (r+4)^{i-1})$$

$$\leq (r+4)^{i} \mu \qquad (\operatorname{dist}_{G}(u, u_{j_2}) \leq \operatorname{dist}_{G}(u, v) \leq r^{i} \mu)$$

Proof of Theorem 1. Fix  $u, v \in V$  and  $d = \operatorname{dist}_G(u, v)$ . Define  $\epsilon' = \ln(1 + \epsilon)$ . Notice that  $1/\epsilon' = (1 + o(1))(1/\epsilon)$ . Set  $r = \lceil 4k/\epsilon' \rceil = \Theta(k/\epsilon)$  and  $\mu = d/r^k$ . By Theorem 4, since  $V_{k+1} = \emptyset$ , condition (i) must hold: within  $h_k < 2(r+1)^k$  hops we have

$$\begin{split} d_{G \cup H}^{(h_k)}(u,v) & \leq \mathrm{dist}_G(u,v) + ((r+4)^k - r^k)\mu \\ & = d + \left(\frac{4k}{r} + \frac{4^2 \binom{k}{2}}{r^2} + \frac{4^3 \binom{k}{3}}{r^3} + \cdots\right) d \\ & \leq \left(1 + \epsilon' + \frac{\epsilon'^2}{2!} + \frac{\epsilon'^3}{3!} + \cdots\right) d \qquad \qquad (\text{since } 4k/r \leq \epsilon') \\ & \leq e^{\epsilon'} d = (1 + \epsilon) d. \end{split}$$

Observe that if we set  $k = \log \log n - O(1)$  the size becomes linear.

Corollary 1. Every n-vertex graph has an O(n)-size  $(\beta, \epsilon)$ -hopset with  $\beta = 2(\frac{(4+o(1))k}{\epsilon})^k$  and  $k = \log \log n - O(1)$ .

## 3 Conclusion

In this paper our goal was to demonstrate that hopset constructions need not be complex, and that optimal hopsets can be constructed with a *simple* and elegant algorithm, namely a small modification to Thorup and Zwick's emulator construction [20]. From a purely quantitative perspective our hopsets also improve on the sparseness and/or hopbound of other constructions [7, 9, 10]. As a happy byproduct of our construction, we also shave small factors off the best sublinear additive emulators [20] and  $(1 + \epsilon, \beta)$ -spanners [1].

We now have a good understanding of the tradeoffs available between  $\beta$  and the hopset size when the stretch is fixed at  $1 + \epsilon$ ,  $\epsilon > 0$  being a small real. However, when  $\epsilon = 0$  or  $\epsilon$  is large, there are still gaps between the best upper and lower bounds. For example, when  $\epsilon = 0$  a trivial hopset<sup>4</sup> has size O(n) with  $\beta = O(\sqrt{n}\log n)$ . A construction of Hesse [13] (see also [1, §6]) implies that  $\beta$  must be at least  $n^{\delta}$  for some  $\delta$ , but it is open whether O(n)-size hopsets exist with  $\beta \ll \sqrt{n}$ . At the other extreme, Thorup and Zwick's distance oracles imply that  $O(\kappa n^{1+1/\kappa})$ -size hopsets exist with  $\beta = 2$  and stretch  $2\kappa - 1$ . Is this tradeoff optimal? Are there other tradeoffs available when  $\beta$  is a fixed constant (say 3 or 4), independent of  $\kappa$ ?

Acknowledgement. Thanks to Richard Peng for help with the references for zero-stretch hopsets.

### References

- [1] A. Abboud, G. Bodwin, and S. Pettie. A hierarchy of lower bounds for sublinear additive spanners. In *Proceedings 28th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 568–576, 2017.
- [2] D. Aingworth, C. Chekuri, P. Indyk, and R. Motwani. Fast estimation of diameter and shortest paths (without matrix multiplication). SIAM Journal on Computing, 28(4):1167–1181, 1999.
- [3] I. Althöfer, G. Das, D. Dobkin, D. Joseph, and J. Soares. On sparse spanners of weighted graphs. Discrete and Computational Geometry, 9:81–100, 1993.
- [4] S. Baswana, T. Kavitha, K. Mehlhorn, and S. Pettie. Additive spanners and  $(\alpha, \beta)$ -spanners. ACM Trans. Algorithms, 7(1), 2010.

<sup>&</sup>lt;sup>4</sup>Let H be a clique on a set of  $\sqrt{n}$  vertices chosen uniformly at random.

- [5] A. Bernstein. Fully dynamic  $(2 + \epsilon)$ -approximate all-pairs shortest paths with fast query and close to linear update time. In *Proceedings 50th Annual IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 693–702, 2009.
- [6] E. Cohen. Using selective path-doubling for parallel shortest-path computations. *Journal of Algorithms*, 22(1):30–56, 1997.
- [7] E. Cohen. Polylog-time and near-linear work approximation scheme for undirected shortest paths. *J. ACM*, 47(1):132–166, 2000.
- [8] D. Dor, S. Halperin, and U. Zwick. All-pairs almost shortest paths. SIAM Journal on Computing, 29(5):1740–1759, 2000.
- [9] M. Elkin and O. Neiman. Hopsets with constant hopbound, and applications to approximate shortest paths. In *Proceedings 57th Annual IEEE Symposium on Foundations of Computer Science*, (FOCS), pages 128–137, 2016.
- [10] M. Elkin and O. Neiman. Linear-size hopsets with small hopbound, and distributed routing with low memory. CoRR, abs/1704.08468, 2017.
- [11] M. Henzinger, S. Krinninger, and D. Nanongkai. Decremental single-source shortest paths on undirected graphs in near-linear total update time. In *Proceedings 55th Annual IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 146–155, 2014.
- [12] M. Henzinger, S. Krinninger, and D. Nanongkai. A deterministic almost-tight distributed algorithm for approximating single-source shortest paths. In *Proceedings 48th Annual ACM Symposium on Theory of Computing (STOC)*, pages 489–498, 2016.
- [13] W. Hesse. Directed graphs requiring large numbers of shortcuts. In *Proceedings 14th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 665–669, 2003.
- [14] P. N. Klein and S. Subramanian. A randomized parallel algorithm for single-source shortest paths. Journal of Algorithms, 25(2):205–220, 1997.
- [15] M. B. T. Knudsen. Additive spanners: A simple construction. In *Proceedings 14th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT)*, pages 277–281, 2014.
- [16] G. L. Miller, R. Peng, A. Vladu, and S. C. Xu. Improved parallel algorithms for spanners and hopsets. In Proceedings 27th ACM Symposium on Parallelism in Algorithms and Architectures (SPAA), pages 192–201, 2015.
- [17] S. Pettie. Low distortion spanners. ACM Trans. Algorithms, 6(1):7:1–7:22, 2009.
- [18] H. Shi and T. H. Spencer. Time-work tradeoffs of the single-source shortest paths problem. *Journal of Algorithms*, 30(1):19–32, 1999.
- [19] M. Thorup and U. Zwick. Approximate distance oracles. J. ACM, 52(1):1-24, 2005.
- [20] M. Thorup and U. Zwick. Spanners and emulators with sublinear distance errors. In Proceedings 17th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), pages 802–809, 2006.
- [21] J. D. Ullman and M. Yannakakis. High-probability parallel transitive-closure algorithms. SIAM J. Comput., 20(1):100–125, 1991.