

POSITIVE LYAPUNOV EXPONENTS FOR COCYCLES OVER TIME ONE ANOSOV MAPS

MAURICIO POLETTI

ABSTRACT. We prove that in an open and dense set, Symplectic linear cocycles over time one maps of Anosov flows, have positive Lyapunov exponents for SRB measures.

1. INTRODUCTION

Positiveness of Lyapunov exponents was widely studied in the past years. Avila [4] proved for several topologies, including C^r topology for $r \geq 0$, that there exists a dense, but not open, set of cocycles taking values in $SL(2, \mathbb{R})$ with non zero Lyapunov exponents. This was generalized to $Sp(2d, \mathbb{R})$ cocycles by Xu [14].

A result stated by Mañé and proved by Bochi [7] expose that the generic behaviour in the C^0 topology is not positiveness of Lyapunov exponents: C^0 generically $SL(2, \mathbb{R})$ cocycles are uniformly hyperbolic or have zero Lyapunov exponents.

In the case of more regular cocycles with some hyperbolic behaviour on the base map and bunching condition on the fibers, the generic behavior changes radically. Viana [11] proved that in this class the Lyapunov exponents are generically non-zero: When the base maps are non-uniformly hyperbolic and the cocycle take values in the group $SL(d, \mathbb{R})$. This was generalized by Bessa-Bochi-Cambrainha-Matheus-Varandas-Xu [6] for any non-compact semi-simple Lie group (for example $Sp(2d, \mathbb{R})$).

These results were extended in the partially hyperbolic setting in two cases: when the center leaves are compact, the map has zero exponent on the center direction and the cocycle takes values in $Sp(2d, \mathbb{R})$, by the author [9], and when the map is volume preserving dynamically coherent, accessible and the cocycle takes values in $SL(d, \mathbb{R})$ by Avila-Viana-Santamaria [2].

Here we deal with an especial case of partially hyperbolic maps, the time one map of a Anosov flow that preserves an SRB measure. As we do not need to assume that our map is accessible, we can deal with cases that can not be covered by [2], for example Anosov flows given by suspensions of hyperbolic diffeomorphism.

2010 *Mathematics Subject Classification.* 37H15,37D30,37D25.

Key words and phrases. Lyapunov Exponents, Partially Hyperbolic Diffeomorphism, Anosov flows, Linear Cocycles.

For an volume preserving Anosov flow, the volume is an SRB invariant measure, but we can also deal with dissipative flows (this means that do not preserve volume). For example any non-trivial attractor of an Anosov flows admits some SRB measure and our result cover this cases.

Acknowledgements. Thanks to C. Matheus and M. Shannon for the discussions and useful ideas, and F. Lenarduzzi for the commentaries.

2. DEFINITIONS AND RESULTS

A diffeomorphism $f : M \rightarrow M$ of a compact manifold M is *partially hyperbolic* if there exists a non-trivial splitting of the tangent bundle

$$TM = E^s \oplus E^c \oplus E^u$$

invariant under the derivative Df , a Riemannian metric $\|\cdot\|$ on M , and positive constants λ, γ with $\lambda < 1$, and $\lambda < \gamma$ such that, for any unit vector $v \in T_p M$,

$$(1) \quad \|Df(p)v\| < \lambda \quad \text{if } v \in E^s(p),$$

$$(2) \quad \gamma < \|Df(p)v\| < \gamma^{-1} \quad \text{if } v \in E^c(p),$$

$$(3) \quad \lambda^{-1} < \|Df(p)v\| \quad \text{if } v \in E^u(p).$$

All three sub bundles E^s, E^c, E^u are assumed to have positive dimension.

The stable and unstable bundles E^s and E^u are uniquely integrable and their integral manifolds form two transverse continuous foliations W^s and W^u , whose leaves are immersed sub manifolds of the same class of differentiability as f . These foliations are referred to as the *strong-stable* and *strong-unstable* foliations. They are invariant under f , in the sense that

$$f(W^s(x)) = W^s(f(x)) \quad \text{and} \quad f(W^u(x)) = W^u(f(x)),$$

where $W^s(x)$ and $W^u(x)$ denote the leaves of W^s and W^u , respectively, passing through any $x \in M$.

We take M to be endowed with the distance dist associated to such a Riemannian structure. The *Lebesgue class* is the measure class of the volume induced by this (or any other) Riemannian metric on M . These notions extend to any sub manifold of M , just considering the restriction of the Riemannian metric to the sub manifold. We say that f is *volume preserving* if it preserves some probability measure in the Lebesgue class of M .

A special class of partially hyperbolic maps are the *time one maps of Anosov flows*. We say that a flow $\phi_t : M \rightarrow M$ is an Anosov flow if there exists a splitting

$$TM = E^s \oplus X \oplus E^u,$$

where $X(p) = \frac{d}{dt}\phi_t(p) |_{t=0}$, invariant under the derivative $D\phi_t$, where E^s is contracted and E^u is expanded by the derivative of ϕ_t .

Let $f : M \rightarrow M$ be the time one map of a C^2 Anosov flow (i.e: $f = \phi_1$). This map is partially hyperbolic with center bundle $E^c = X$.

In this particular case the center bundle $E^c = X$ is integrable and C^1 (observe that $W^c(p) = \phi_{\mathbb{R}}(p)$). The bundles $E^c + E^u$ and $E^s + E^c$ are integrable and absolutely continuous (See [1]). We call them center stable W^{cs} and center unstable W^{cu} foliations.

We say that a measure μ is an *SRB measure* for the flow if the disintegration of the measure along the center unstable leaves is in the Lebesgue class along this sub-manifolds (for a more details see [1]). Observe that we are asking for the measure to be invariant for the flow not just for the time one map.

Recall that by the spectral decomposition theorem the non-wandering set of the flow can be decomposed into basic sets $\Omega(\phi_t) = \Delta_1 \cup \dots \cup \Delta_n$, where the Δ_i are invariant and the restriction ϕ_t is transitive in this sets, in particular the support of every ergodic measure is supported in one of this sets.

Also this basic sets coincide with the closure of the homoclinic class of its periodic points, as the support of an SRB measures is saturated by the center unstable foliation this implies that $\text{supp}(\mu) = \Delta_i$ for some $i \in \{1, \dots, n\}$, moreover this set is an attractor. Conversely, every non trivial attractor of an hyperbolic flow admits some SRB measure.

The symplectic group $Sp(2d, \mathbb{R})$ is the subgroup of matrices $B \in GL(2d, \mathbb{R})$ such that

$$B^T J B = J, \quad \text{and} \quad J = \begin{pmatrix} 0 & \text{id}_d \\ -\text{id}_d & 0 \end{pmatrix}.$$

Let $H^\alpha(M)$ be the Banach space of α -Hölder maps $A : M \rightarrow Sp(2d, \mathbb{R})$ with norm

$$\|A\|_\alpha = \sup_{x \in M} \|A(x)\| + \sup_{x \neq y} \frac{\|A(x) - A(y)\|}{\|x - y\|^\alpha}.$$

The *linear cocycle* defined by $A : M \rightarrow Sp(2d, \mathbb{R})$ over $f : M \rightarrow M$ is the (invertible) map $F_A : M \times \mathbb{R}^{2d} \rightarrow M \times \mathbb{R}^{2d}$ given by

$$F_A(x, v) = (f(x), A(x)v).$$

Its iterates are given by $F_A^n(x, v) = (f^n(x), A^n(x)v)$ where

$$A^n(x) = \begin{cases} A(f^{n-1}(x)) \dots A(f(x))A(x) & \text{if } n > 0 \\ Id & \text{if } n = 0 \\ A(f^n(x))^{-1} \dots A(f^{-1}(x))^{-1} & \text{if } n < 0 \end{cases}$$

Sometimes we denote this cocycle by (f, A) .

Let μ be an f -invariant probability measure on M and suppose that $\log \|A\|$ and $\log \|A^{-1}\|$ are integrable. By Kingman's sub-additive ergodic theorem, see [12], the limits

$$\lambda^+(A, \mu, x) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A^n(x)\| \quad \text{and} \\ \lambda^-(A, \mu, x) = \lim_{n \rightarrow \infty} -\frac{1}{n} \log \|(A^n(x))^{-1}\|,$$

exist for μ -almost every $x \in M$. When there is no risk of ambiguity we write just $\lambda^+(x) = \lambda^+(A, \mu, x)$.

By Oseledets [8], at μ -almost every point $x \in M$ there exist real numbers $\lambda^+(x) = \lambda_1(x) > \dots > \lambda_k(x) = \lambda^-(x)$ and a decomposition $\mathbb{R}^d = E_x^1 \oplus \dots \oplus E_x^k$ into vector subspaces such that

$$A(x)E_x^i = E_{f(x)}^i \text{ and } \lambda_i(x) = \lim_{|n| \rightarrow \infty} \frac{1}{n} \log \|A^n(x)v\|$$

for every non-zero $v \in E_x^i$ and $1 \leq i \leq k$.

Define $L(A, \mu) = \int \lambda^+ d\mu$, we say that A has *positive exponent* if $L(A, \mu) > 0$. When μ is ergodic, as we are going to assume later, we have $L(A, \mu) = \lambda^+(x)$ for μ -almost every $x \in M$.

We say that the cocycle is *fiber bunched* if there exist $C > 0$ and $\theta < 1$ such that

$$(4) \quad \|A^n(p)\| \|A^n(p)^{-1}\| \lambda^{n\alpha} \leq C\theta^n \quad \text{for every } p \in M \text{ and } n \geq 0,$$

where λ is a hyperbolicity constant for f as in conditions (1), (2), (3).

So now we can state the main theorem of this work.

Theorem A. Let $f : M \rightarrow M$ be a C^2 time one map of an Anosov flow and μ an SRB invariant measure for the flow, then $L(A, \mu) > 0$ in an open and dense subset of the fiber bunched cocycles of $H^\alpha(M)$.

3. HOLONOMIES

Suppose that $f : M \rightarrow M$ is partially hyperbolic. The stable and unstable foliations are, usually, *not* transversely smooth: the holonomy maps between any pair of cross-sections are not even Lipschitz continuous, in general, although they are always γ -Hölder continuous for some $\gamma > 0$. Moreover, if f is C^2 then these foliations are absolutely continuous, in the following sense.

Let $d = \dim M$ and \mathcal{F} be a continuous foliation of M with k -dimensional smooth leaves, $0 < k < d$. Let $\mathcal{F}(p)$ be the leaf through a point $p \in M$ and $\mathcal{F}(p, R) \subset \mathcal{F}(p)$ be the neighbourhood of radius $R > 0$ around p , relative to the distance defined by the Riemannian metric restricted to $\mathcal{F}(p)$. A *foliation box* for \mathcal{F} at p is the image of an embedding

$$\Phi : \mathcal{F}(p, R) \times \mathbb{R}^{d-k} \rightarrow M$$

such that $\Phi(\cdot, 0) = \text{id}$, every $\Phi(\cdot, y)$ is a diffeomorphism from $\mathcal{F}(p, R)$ to some subset of a leaf of \mathcal{F} (we call the image a *horizontal slice*), and these diffeomorphisms vary continuously with $y \in \mathbb{R}^{d-k}$. Foliation boxes exist at every $p \in M$, by definition of continuous foliation with smooth leaves. A *cross-section* to \mathcal{F} is a smooth co-dimension- k disk inside a foliation box that intersects each horizontal slice exactly once, transversely and with angle uniformly bounded from zero.

Then, for any pair of cross-sections Σ and Σ' , there is a well defined *holonomy map* $\Sigma \rightarrow \Sigma'$, assigning to each $x \in \Sigma$ the unique point of intersection

of Σ' with the horizontal slice through x . The foliation is *absolutely continuous* if all these homeomorphisms map zero Lebesgue measure sets to zero Lebesgue measure sets. That holds, in particular, for the strong-stable and strong-unstable foliations of partially hyperbolic C^2 diffeomorphisms and, in fact, the Jacobians of all holonomy maps are bounded by a uniform constant.

Let $f : M \rightarrow M$ be the time one map of an C^2 Anosov flow, his center stable and center unstable foliations are also absolutely continuous. If we take two center manifolds $W^c(p)$ and $W^c(z)$ in the same strong stable manifold, from every point $t \in W^c(p)$ we can define the stable holonomy locally $h_{p,z}^s : I \subset W^c(p) \rightarrow W^c(z)$ and the Jacobian of $h_{p,z}^s$ vary continuously with the points p and z . Actually the Jacobian is given by

$$\lim_{n \rightarrow \infty} \frac{Jf_c^n(t)}{Jf_c^n(h_{p,z}^s(t))},$$

where $Jf_c^n(t) = \det Df^n|_{E_t^c}(t)$. Analogously for the unstable holonomies.

3.1. Measure disintegration. Given a measurable partition \mathcal{P} of M , by Rokhlin disintegration theorem (see [12]) there exists a family of measures $\{\mu_P\}_{P \in \mathcal{P}}$ such that for every measurable set $B \in M$

- $P \rightarrow \mu_P(B)$ is measurable,
- $\mu_P(P) = 1$ and
- $\mu(B) = \int_M \mu_P(B) d\tilde{\mu}(P)$.

Moreover, such disintegration is essentially unique [10].

In general the partition by the invariant foliations is not measurable, to overcome this problem we disintegrate our measure locally as we explain next.

We call $C_x \in M$ a foliated box centred in $x \in M$ if there exists a continuous function $\Phi : [-1, 1]^d \rightarrow C$ such that

- $\Phi(0) = x$,
- for every $y \in [-1, 1]^{d-1}$,

$$W_{C_x}^c(\Phi(y, 0)) := \Phi(y, [-1, 1]) \subset W^c(\Phi(y, 0)),$$

- for every $y \in [-1, 1]^u$,

$$W_{C_x}^{cs}(\Phi(0_s, y, 0)) := \Phi([-1, 1]^s \times \{y\} \times [-1, 1]) \subset W^{cs}(\Phi(0_s, y, 0)),$$

- for every $x \in [-1, 1]^s$,

$$W_{C_x}^{cu}(\Phi(x, 0_u, 0)) := \Phi(\{x\} \times [-1, 1]^u \times [-1, 1]) \subset W^{cu}(\Phi(x, 0_u, 0)),$$

where s is the dimension of E^s and u the dimension of E^u .

So μ is an SRB measure if and only if for every x the disintegration of $\mu|_{C_x}$ given by the partition $\{W_{C_x}^{cu}(y)\}_{y \in C_x}$ gives measures absolutely continuous with respect to the Lebesgue measure on $W^{cu}(y)$.

Given a foliated box C_x lets call $\Sigma_x = \Phi([-1, 1]^{d-1} \times \{0\})$ and $\pi : C_x \rightarrow \Sigma_x$ the natural projection given by $\pi(\Phi(x_s, y_u, t)) = \Phi(x_s, y_u, 0)$, observe that Σ_x has a product structure $\Sigma^s \times \Sigma^u$ given by Φ . We say that μ has *projective*

product structure if $\pi_*\mu \mid C_x$ is absolutely continuous with a product measure $\mu^s \times \mu^u$, where μ^* is a measure on Σ^* for $* = s, u$.

We say that a measure μ has a *good product structure* if it has projective product structure and also for every foliated box the disintegration in center manifolds $\{W_{C_x}^c(y)\}_{y \in C_x}$ is absolutely continuous with respect to the Lebesgue measure.

Every SRB measure for the flow has a good projective product structure, the absolute continuity in the center manifold is just because the measure is invariant by the flow and the projective product structure is because of the absolute continuity of the center stable foliation (see [13, Proposition 3.4] for a proof of this fact).

The good product structure of the SRB measure will be the key property to prove theorem 4.1 in section 4, that is the main tool for proving our result.

3.2. Linear holonomies.

Definition 3.1. We say that A admits *strong stable holonomies* if there exist, for every p and $q \in M$ with $p \in W_{loc}^s(q)$, linear transformations $H_{p,q}^{s,A} : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ with the properties:

- (1) $H_{f^j(p), f^j(q)}^{s,A} = A^j(q) \circ H_{p,q}^{s,A} \circ A^j(p)^{-1}$ for every $j \geq 1$,
- (2) $H_{p,p}^{s,A} = id$ and, for $w \in W^s(p)$, $H_{p,q}^{s,A} = H_{w,q}^{s,A} \circ H_{p,w}^{s,A}$,
- (3) there exists some $L > 0$ (that does not depend on p, q) such that $\|H_{p,q}^{s,A} - id\| \leq L \text{dist}(p, q)^\alpha$.

Moreover, these holonomies vary continuously with the base points p and q .

These linear transformations are called *strong stable holonomies*.

Observe that if p and q are in the same strong stable manifold (not necessarily local) there exists some $n \geq 0$ such that $f^n(p)$ and $f^n(q)$ are in the same local strong stable manifold. So we can extend the holonomies to these points defining $H_{p,q}^{s,A} = A^n(q)^{-1} \circ H_{f^n(p), f^n(q)}^{s,A} \circ A^n(p)$, by property (1) this does not depend on the choice of n .

Analogously if $p \in W_{loc}^u(q)$ we have the *strong unstable holonomies*. When there is no ambiguity we write $H_{p,q}^s$ instead of $H_{p,q}^{s,A}$.

Remark 3.2. The fiber bunched condition (4) implies the existence of the strong stable and strong unstable holonomies (see [2]).

Let $\mathbb{R}P^{2d-1}$ be the real projective space, i.e: the quotient space of $\mathbb{R}^{2d} \setminus \{0\}$ by the equivalence relation $v \sim u$ if there exist $c \in \mathbb{R} \setminus \{0\}$, such that, $v = cu$. Every invertible linear transformation B induces a projective transformation, that we also denote by B ,

$$B : \mathbb{R}P^{2d-1} \rightarrow \mathbb{R}P^{2d-1} \quad \text{and} \quad B[v] = [B(v)].$$

We also denote by $F : M \times \mathbb{R}P^{2d-1} \rightarrow M \times \mathbb{R}P^{2d-1}$ the induced projective cocycle.

3.3. Invariance principle. Let $P : M \times \mathbb{R}P^{2d-1} \rightarrow M$ be the projection to the first coordinate, $P(x, v) = x$, and let m be a F -invariant measure on $M \times \mathbb{R}P^{2d-1}$ such that $P_*m = \mu$. By Rokhlin [10] we can disintegrate the measure m into $\{m_x, x \in M\}$ with respect to the partitions

$$\mathcal{P} = \{\{p\} \times \mathbb{R}P^{2d-1}, p \in M\},$$

A measure m that projects on μ is called:

- (a) *u-invariant* if there exists a full μ -measure subset $M^s \subset M$ such that $m_z = (H_{y,z}^s)_*m_y$ for every $y, z \in M^s$ in the same strong-stable leaf;
- (b) *s-invariant* if there exists a full μ -measure subset $M^u \subset M$ such that $m_z = (H_{y,z}^u)_*m_y$ for every $y, z \in M^u$ in the same strong-unstable leaf.

If both *a* and *b* are satisfied we say that m is *su-invariant*.

We recall the following result whose proof can be found in [2].

Theorem 3.3. *Let $f : M \rightarrow M$ be a C^1 partially hyperbolic diffeomorphism and $A : M \rightarrow GL(2d, \mathbb{R})$ a linear cocycle defined over f . If μ is an f -invariant measure, such that $\lambda^+(A, \mu, x) = \lambda^-(A, \mu, x) = 0$ almost everywhere, then every F_A -invariant measure m such that $P_*m = \mu$ is *su-invariant*.*

4. PROOF OF THE MAIN RESULT

We say that a disintegration of m is *u/c-invariant* if, for every x and y in the support of μ and in the same unstable manifold, for Lebesgue almost every $t \in W^c(x)$, $H_{t, h(t)}^u m_t = m_{h(t)}$ where $h(t) = W^c(y) \cap W^u(t)$. Analogously, we call a disintegration *s/c-invariant* changing unstable by stable. We say that the disintegration is *su/c-invariant* if the disintegration is both *u/c* and *s/c*-invariant.

The primary tool to prove our main theorem is given by the next theorem:

Theorem 4.1. *Let $f : M \rightarrow M$ be a C^2 partially hyperbolic, dynamically coherent diffeomorphism and μ an invariant measure with a good product structure, then if m is an *su-invariant* measure then there exists a disintegration that is *su/c-invariant*.*

Proof. For any topological space N , we denote by $\mathcal{M}(N)$ the space of measures on N with the weak* topology.

There exist M^s and M^u of total measure with *s*-invariance and *u*-invariance respectively. Take $M' = M^s \cap M^u$, $x \in \text{supp}(\mu)$ and a foliated box $C_x \subset M$; via a local chart we can write $C = \Sigma \times D_R^c$ where Σ is a transversal section to the center foliation and D_R^c is a disc of radius R in a center manifold, let $\pi : C_x \rightarrow \Sigma$ be the natural projection given by the center discs, observe that the center stable and center unstable manifolds gives a product structure of $\Sigma = \Sigma^s \times \Sigma^u$ and by hypothesis $\mu_\Sigma \sim \mu^s \times \mu^u$.

By the absolute continuity of the center foliation and the continuity of the Jacobians of the stable and unstable holonomies of f we have that the disintegration $\Sigma \rightarrow \mathcal{M}(D_R^c)$, $x \mapsto \mu_x^c$ is of the form $\mu_x^c = \rho \mu^c$ where μ^c is the Lebesgue measure in D_R^c and ρ is continuous.

We write $x \sim^s y$ for $x, y \in \Sigma$ in the same center stable manifold, and $x \sim^u y$ if they are in the same center unstable manifold. Take $r > 0$ smaller than R such that for every $x \sim^s y$, $h_{x,y}^s : D_r^c \rightarrow D_R^c$ is well defined.

Fix some $x^s \in \Sigma^s$ such that $\mu^u(\{x^s\} \times \Sigma^u \cap \pi(M')) = 1$ and fix some $x^u \in \Sigma^u$ such that $\mu_{(x^s, x^u)}^c((x^s, x^u) \times D_R^c \cap M') = 1$, by the absolute continuity in the center direction this implies that Lebesgue almost every $t \in (x^s, x^u) \times D_R^c$ is in M' ; for simplicity denote $x_0 = (x^s, x^u)$.

Take $\epsilon > 0$ such that

$$D_\epsilon^c \subset \bigcap_{z \sim^u y \sim^s x_0} h_{y,z}^u h_{x_0,y}^s(D_r^c).$$

Lets call $m_x^c = \int_{D_r^c} m_t d\mu_x^c(t)$. Taking $x \sim^s y$ for $t \in D_r^c$ lets call $h(t) = h_{x,y}^s(t)$ and define $(\mathcal{H}_{x,y}^s m_x^c)_{h(t)} = H_{(x,t),(y,h(t))}^s m_t$. If m_t is well defined for almost every $t \in D_r^c$ then by the absolute continuity of the holonomies $(\mathcal{H}_{x,y}^s m_x^c)_{t'}$ is defined for almost every $t' \in h_{x,y}^s(D_r^c)$. Analogously we can define $(\mathcal{H}_{x,y}^u m_x^c)_{t'}$

Now define a new disintegration of m in the box $\Sigma \times D_\epsilon^c$ in the following way: Fix the restriction of the original disintegration defined for almost every $(x_0, t) \in \{x_0\} \times D_r^c$, (in particular $m_{x_0}^c$ is well defined) extend the disintegration to every $y \sim^u x_0$ and $t' \in h_{x_0,y}^u(D_r^c)$ by $(\mathcal{H}_{x_0,y}^u m_{x_0}^c)_{t'}$. Lets call this disintegration m_y^c , and then extend it for every $z \sim^s y \sim^u x_0$ by $(\mathcal{H}_{y,z}^s m_y^c)_{t''}$ for almost every $t'' \in h_{y,z}^s h_{x_0,y}^u(D_r^c)$. Now by definition of ϵ this disintegration is well defined in $\Sigma \times D_\epsilon^c$, and it coincides with the original disintegration almost everywhere, lets denote this new disintegration by $t \mapsto m_t^s$.

By construction this disintegration is s/c -invariant. We claim that this new disintegration, with respect to $\{x\} \times D_\epsilon^c \times \mathbb{R}P^{2d-1}$, defined by

$$\Sigma \rightarrow \mathcal{M}(D_r^c \times \mathbb{R}P^{2d-1}), \quad m_x^{cs} = \int_{D_r^c} m_t^s d\mu_x^c(t),$$

is continuous. Assuming this claim, we can define analogously, reducing ϵ if necessary, a disintegration $x \mapsto m_x^u$ that is u/c -invariant on $\Sigma \times \epsilon$. We have that $m_x^{cs} = m_x^{cu}$ for Lebesgue almost every $x \in \Sigma$, then as both are continuous we have the equality for every $x \in \Sigma$. Then by the uniqueness of the Rokhlin disintegration, for every $x \in \Sigma$, $m_t^s = m_t^u$ for μ^c almost every $t \in D_\epsilon^c$. As this boxes cover $\text{supp}(\mu)$ we conclude the theorem.

We are left to prove the claim.

We will prove that $x \mapsto m_x^{cs}$ is uniformly continuous varying $y \sim^s x$ and $z \sim^u y$. Fix some continuous $\varphi : D_\epsilon^c \times \mathbb{R}P^{2d-1} \rightarrow \mathbb{R}$ and let $y_n \rightarrow x$, $y_n \sim^s x$.

$$\begin{aligned} \int \varphi(t, v) dm_{y_n}^{cs} &= \int_{D_\epsilon^c} \int_{\mathbb{R}P^{2d-1}} \varphi(t, v) dm_{y_n, t}^s(v) d\mu_{y_n}^c(t) \\ \int \varphi(t, v) dm_{y_n}^{cs} &= \int_{D_\epsilon^c} \int_{\mathbb{R}P^{2d-1}} \varphi(t, v) dH_{n, t} m_{x, h_n(t)}^s(v) d\mu_{y_n}^c(t), \end{aligned}$$

where $h_n(t) = h_{y_n, x}^s(t)$, and $H_{n, t} = H_{(x, h_n(t)), (y_n, t)}^s$ so we can write the integral as

$$\int_{D_\epsilon^c} \int_{\mathbb{R}P^{2d-1}} \varphi(t, H_{n, t} v) dm_{x, h_n(t)}^s(v) \rho(y_n, t) d\mu^c(t),$$

changing variables and we have

$$\int_{h_n(D_\epsilon^c)} \int_{\mathbb{R}P^{2d-1}} \varphi(h_n^{-1}(t), H_{n, h_n^{-1}(t)} v) dm_{x, t}^s(v) J_{\mu^c} h_n^{-1}(t) \rho(y_n, h_n^{-1}(t)) d\mu^c(t),$$

where $J_{\mu^c} h_n^{-1}$ is the Jacobian of h_n^{-1} with respect to the Lebesgue measure μ^c . Hence using that $h_n^{-1} \rightarrow \text{id}_{W^c(x)}$, $J_{\mu^c} h_n^{-1} \rightarrow 1$, $H_{n, t} \rightarrow \text{id}_{\mathbb{R}P^{2d-1}}$ uniformly and the continuity of ρ and φ , we conclude that $\int \varphi(t, v) dm_{y_n}^{cs} \rightarrow \int \varphi(t, v) dm_x^{cs}$.

For $z_n \sim^u x$, such that $z_n \rightarrow x$, we take $z_n \sim^s y_n \sim^u x_0$ and $x \sim^s x' \sim^u x_0$, by construction $y_n \rightarrow x'$ so

$$\int \varphi(t, v) dm_{y_n}^{cs} \rightarrow \int \varphi(t, v) dm_{x'}^{cs},$$

observe that,

$$\int \varphi(t, v) dm_{z_n}^{cs} = \int_{D_\epsilon^c} \int_{\mathbb{R}P^{2d-1}} \varphi(t, v) dH_{n, t}^u m_{z_n, h_n^u(t)}^s(v) d\mu_{z_n}^c(t),$$

where $H_{n, t}^u$ and h_n^u are defined as before just changing stable by unstable in the corresponding point. Therefore, using the same calculations as before, it follows that $\int \varphi(t, v) dm_{z_n}^{cs} \rightarrow \int \varphi(t, v) dm_x^{cs}$.

This proves the claim and concludes the proof. \square

As we mention before, the support of the SRB measure μ coincide with a non-trivial attractor of the spectral decomposition of $\Omega(\phi_t)$. This implies that we have density of periodic orbits and the homoclinic intersection of this periodic orbits is also dense and contained in the support of μ .

Now we prove that after a perturbation we have stably positive Lyapunov exponents for the cocycle. For this lets first find some center manifold with positive Lyapunov exponents.

Observe that, as the center manifolds of f are the lines flows of ϕ_t , the density of periodic orbits of the flow implies density of compact invariant center leaves. Moreover, the restriction of f to one of this compact leaves,

$W^c(p)$, is a rotation in $\mathbb{R}/T\mathbb{Z}$ where T is the period p by the flow. So after a C^1 change of coordinates we can suppose that $f_p = f|_{W^c(p)} = r_{\frac{1}{T}}$ where

$$r_{\frac{1}{T}} : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z}, \quad t \mapsto t + \frac{1}{T}.$$

The restriction of the cocycle to $W^c(p)$ is a cocycle over a rotation with Lebesgue μ_p^c as invariant measure.

Now we are going to define $h : W^c(p) \rightarrow W^c(p)$ that is a composition of stable and unstable holonomies (see figure 1).

Fix a flow periodic point $p \in \text{supp}(\mu)$, lets take $z \in W^{cs}(p) \cap W^{cu}(p)$, by invariance of the center stable and center unstable manifolds $\phi_t(z) \in W^{cs}(p) \cap W^{cu}(p)$ for every $t \in \mathbb{R}$. Observe also that the orbit of z is non recurrent.




FIGURE 1. definition $h : W^c(p) \rightarrow W^c(p)$.

Suppose that actually $z \in W^u(p)$ and define $h(p) = W^c(p) \cap W^s(z)$, now for $t \in [1, T]$ define $h(\phi_t(p)) := \phi_t(h(p))$ and observe that by the invariance of the stable and unstable manifolds $\phi_t(h(p)) \in W^c(\phi_t(p)) \cap W^s(\phi_t(z))$ and $\phi_t(z) \in W^u(\phi_t(p))$.

So h is a composition of stable and unstable holonomies. Also in the circle coordinates, identifying p with 0,

$$h : \mathbb{R}\mathbb{Z} \rightarrow \mathbb{R}\mathbb{Z}, \quad t \mapsto t + \omega,$$

where ω is such that $\phi_\omega(p) = h(p)$.

Call $h_{p,z}^u : W^c(p) - \{p\} \rightarrow W^c(z)$ the map given by $t \mapsto \phi_t(z)$, by the same reasoning as before this map is given by an unstable holonomy, if there is no risk of ambiguity we just write $h^u = h_{p,z}^u$. This map is not well defined in 0 because $\phi_T(z) \neq z$.

Observe that as the center stable and center unstable are dense in $\text{supp}(\mu)$ we can find points z_1, \dots, z_d as before such that any pair z_i, z_j , with $i \neq j$,

are in different orbits of the flow. Hence, we can define maps h_1, \dots, h_d with the properties above.

Lemma 4.2. *There exists A' arbitrarily close to A such that for some center leave $W^c(p)$, $L(A_p, f_p, \mu_p^c) > 0$.*

Proof. Suppose that there exists $p \in \text{supp}(\mu)$ a periodic point for the flow such that $\frac{1}{T_p}$ is irrational then the Lebesgue measure is non periodic, so by [14], we can perturb A such that $L(A_p, f_p, \mu_p^c) > 0$.

If for every periodic point $p \in \text{supp}(\mu)$, $\frac{1}{T_p}$ is rational we have that as T_p can be taken arbitrarily large, we have rational rotations with arbitrarily large period. Now, taking

$$A_\theta = \begin{pmatrix} \cos(\theta) \text{id}_d & \sin(\theta) \text{id}_d \\ -\sin(\theta) \text{id}_d & \cos(\theta) \text{id}_d \end{pmatrix} A,$$

for a generic A we have that for an n -periodic point there exists an $O(\frac{1}{n})$ dense set with $L(A_\theta(t)) > 0$ (see [14, Lemma 3.4]), so taking p such that T_p is very large we can take θ small such that A_θ is close to A . \square

We recall two results whose proofs can be found in [9].

Lemma 4.3. *Suppose there exists some sequence A_k converging to A , such that $L(A_k, \mu) = 0$ then A admits some some su -invariant measure m .*

Lemma 4.4. *Let (E, ω) be a 2d-dimensional symplectic vector space and let $\{(V_j, W_j) : 1 \leq j \leq m\}$ be a finite collection of pairs subspaces of E with complementary dimensions (i.e., $\dim(V_j) + \dim(W_j) = 2d$ for all $1 \leq j \leq m$). Then, there exists a symplectic automorphism σ arbitrarily close to the identity such that*

$$\sigma(V_j) \cap W_j = \{0\} \quad \forall j = 1, \dots, m.$$

As a consequence if the subspaces V and W have not complementary dimensions we can always make $\dim(\sigma(V) + W) = \dim V + \dim W$.

Lemma 4.5. *There exists A' arbitrarily close to A such that A' does not admit any su -invariant measure.*

Proof. Given a neighbourhood of A by Lemma 4.2 we can find A' in this neighbourhood such that there exists some compact center leave $W^c(p)$ with $L(A'_p, f_p, \mu_p^c) > 0$. Take z in $W^u(p) \cap W^{cs}(p)$ and $h : W^c(p) \rightarrow W^c(p)$ defined as before, observe that by the absolute continuity of the holonomies this map is absolutely continuous.

Lets call the Lyapunov exponents of (f_p, A'_p) by $\lambda_c^1, \dots, \lambda_c^k$. By the positivity of the integrated Lyapunov exponents there exists i such that $\lambda_c^i(t) > \lambda_c^{i+1}(t)$ for some $1 \leq i \leq 2d$, as we have finite possibilities of i we can take some positive measure set $I \subset W^c(p)$ such that, for every $t \in I$, there exists an invariant decomposition $\mathbb{R}^{2d} = V_t^u + V_t^s$ with constant dimensions such that the smaller Lyapunov exponent in V_t^u is larger than the largest Lyapunov exponent in V_t^s .

We can also assume, reducing I if necessary, that this decomposition varies continuously in I . As h preserves the Lebesgue measure μ_p^c (h is a rotation in the circle coordinates) we can find an iterate j such that $h^j(I) \cap I$ has positive measure, take j to be the smaller integer with this property.

Suppose that A' admits some su -invariant measure, by Theorem 4.1 we can take the disintegration to be su/c -invariant. By [5, Proposition 3.1] we have that, for every $t \in I$, $m_t = m_t^u + m_t^s$ where $\text{supp}(m_t^u) \subset V_t^u$ and $\text{supp}(m_t^s) \subset V_t^s$. Taking $t' \in I$ density point of $h^j(I) \cap I$ we can find a neighbourhood $N \subset W^c(p)$ of t' such that the sets $h_{p,z}^u(h^i(N)) \subset W^c(z)$, $i = 0, \dots, j-1$ are disjoint.

Let $H_t^{A'}$ be the composition of the stable linear holonomy from t to the point $h^u(t)$ and the unstable linear holonomy from $h^u(t)$ to $h(t)$.

The su/c -invariance implies that $m_{h^j(t)} = (H_t^{A'})_* m_t$ for μ_p^c -almost every point. This implies that

$$\text{supp}(m_{h^j(t)}) \subset \{V_{h^j(t)}^u, V_{h^j(t)}^s\} \cap \{H_t^j V_t^u, H_t^j V_t^s\}.$$

Fix a neighbourhood \mathcal{N} of $h^u(t')$ that does not contain $f^n(h^u(t'))$ for $n \neq 0$, making a small perturbation we find $\hat{A} : M \rightarrow Sp(2d, \mathbb{R})$ such that $\hat{A}(h^u(t')) = \sigma A'(h^u(t'))$ and $\hat{A}(h^u(r)) = A'(h^u(r))$ for every $r \in W^c(z) - N$ and also $\hat{A}(x) = A'(x)$ outside \mathcal{N} . We have that

$$H_{t'}^{\hat{A}^j} = H_{h(t')}^{j-1, A'} \circ H_{h^u(t'), h(t')}^{s, A'} \circ \sigma \circ H_{t', h^u(t')}^{u, A'}.$$

Then using 4.4 we can find σ arbitrarily close to the identity such that $V_{h^j(t')}^s \cap H_{t'}^j V_{t'}^u = \emptyset$ and $V_{h^j(t')}^u \cap H_{t'}^j V_{t'}^s = \emptyset$. So

$$\text{supp}(m_{h^j(t')}) \subset V'_{h^j(t')} = V_{h^j(t')}^s \cap H_{t'}^j V_{t'}^s$$

using 4.4 again we can make $\dim V' < \dim V^s$. As t' and $h^j(t')$ are density points of a set where the V^u and V^s varies continuously then there exists a positive measure subset of $t \in I$ such that

$$\text{supp}(m_{h^j(t)}) \subset V'_{h^j(t)} \quad \text{and} \quad \dim V' < \dim V^s.$$

If \hat{A} still admits some m su -invariant, we can repeat the argument taking a different homoclinic point z_1, \dots, z_d (i.e the orbit of z_i is not in the orbit of z_j for $i \neq j$) with V' instead of V^s such that the new perturbation does not affect the cocycle in the orbit of z . Inductively, if the perturbed cocycles always admit some su -invariant measure, we can do this with less than d -point z_i to get that $\text{supp}(m_t) = \emptyset$, for t in a positive measure set, a contradiction.

Then we can find A'' arbitrarily close to A does not admit any su -invariant measure. \square

proof of Theorem A. By Lemma 4.5 the set of cocycles that do not admit any su -invariant measure is dense, by the invariance principle [3] and by




FIGURE 2. perturbation of H

Lemma 4.3 every cocycle in this set has a neighbourhood of cocycles with positive Lyapunov exponents. \square

REFERENCES

- [1] Vitor Araújo and Maria José Pacifico. *Three Dimensional Flows*, volume 114. Apr 2012.
- [2] A. Avila, J. Santamaria, and M. Viana. Holonomy invariance: rough regularity and applications to Lyapunov exponents. *Astérisque*, 358:13–74, 2013.
- [3] A. Avila and M. Viana. Extremal Lyapunov exponents: an invariance principle and applications. *Invent. Math.*, 181(1):115–189, 2010.
- [4] Artur Avila. Density of positive Lyapunov exponents for $sl(2,r)$ -cocycles. *J. Amer. Math. Soc.*, 24:9991014, 2011.
- [5] L. Backes and M. Poletti. Continuity of Lyapunov exponents is equivalent to continuity of Oseledets subspaces. Preprint <http://arxiv.org/pdf/1511.08423v1.pdf>.
- [6] M. Bessa, J. Bochi, M. Cambrainha, C. Matheus, P. Varandas, and D. Xu. Positivity of the top lyapunov exponent for cocycles on semisimple lie groups over hyperbolic bases. Preprint <https://arxiv.org/pdf/1611.10158.pdf>.
- [7] J. Bochi. Genericity of zero Lyapunov exponents. *Ergod. Th. & Dynam. Sys.*, 22:1667–1696, 2002.
- [8] V. I. Oseledets. A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems. *Trans. Moscow Math. Soc.*, 19:197–231, 1968.
- [9] M. Poletti. Stably positive lyapunov exponents for symplectic linear cocycles over partially hyperbolic diffeomorphisms. Preprint <https://arxiv.org/pdf/1605.00044.pdf>.
- [10] V. A. Rokhlin. On the fundamental ideas of measure theory. *A. M. S. Transl.*, 10:1–54, 1962. Transl. from *Mat. Sbornik* 25 (1949), 107–150. First published by the A. M. S. in 1952 as Translation Number 71.
- [11] M. Viana. Almost all cocycles over any hyperbolic system have nonvanishing Lyapunov exponents. *Ann. of Math.*, 167:643–680, 2008.
- [12] M. Viana and K. Oliveira. *Foundations of Ergodic Theory*. Cambridge University Press, 2015.
- [13] M. Viana and J. Yang. Physical measures and absolute continuity for one-dimensional center direction. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 30:845–877, 2013.

- [14] Disheng Xu. Density of positive lyapunov exponents for symplectic cocycles. Preprint <https://arxiv.org/pdf/1506.05403.pdf>.

LAGA – UNIVERSITÉ PARIS 13, 99 Av. JEAN-BAPTISTE CLÉMENT, 93430 VILLETANEUS, FRANCE.

E-mail address: `mpoletti@impa.br`