

Bayesian Portfolio Selection

Sourish Das
Chennai Mathematical Institute

Rituparna Sen,
Indian Statistical Institute

April 15, 2017

Abstract

We present a Bayesian portfolio selection strategy, which uses the Capital Asset Pricing Model (CAPM). We propose a strategy which will mimic the market if the market is information efficient. However, the strategy will outperform the market if the market is not efficient. The strategy depends on the selection of a portfolio via Bayesian methodology for the parameters of the CAPM, which turns out to be multiple testing problems. We present the “discrete-mixture prior” model and “hierarchical Bayes model” for the intercept and slope parameters of CAPM. In hierarchical Bayes model, we use the half-Cauchy prior on the global shrinkage parameter of the model. We establish the Bayesian optimality properties of multiple testing rules from the Bayesian decision-theoretic point of view. The risk for the Bayesian decision rule up to $O(1)$ attains the risk of Bayes oracle. We present detailed empirical study, where 500 stocks from the New York Stock Exchange (NYSE) are considered and S&P 500 index is taken as the proxy for the market. The study of portfolio selection via four different strategies are examined over the period from the year 2006 to 2014. The out of the sample performance of the portfolio selected by the various methods is presented. Empirical results indicate that market is not efficient and it is possible to propose a strategy which can outperform the market.

Key words: CAPM, Efficient Market, Discrete Mixture Prior, Hierarchical Bayes, Oracle

1 Introduction

Markowitz portfolio theory [19] is a portfolio optimization problem in finance; where an investor allocates the wealth among securities in such a way that the portfolio guarantees a certain level of expected returns and minimizes the ‘risk’ associated with it. The term ‘expected returns’ implies expectation in a statistical sense. This means actual realized return of the portfolio could be less or more than the expected return.

Although Markowitz portfolio selection analytically formalizes the risk-return tradeoff in selecting optimal portfolios, it is very sensitive [12] to errors in the estimates of the expected return vector and the covariance matrix. The problem is severe when the portfolio size is large. Several techniques have been suggested to reduce the sensitivity of the Markowitz optimal portfolios. One approach is to use a James-Stein estimator for means (i.e., expected return) [7] and shrink the sample covariance matrix [17, 8]. Still curse of dimensionality kicks-in for a typically large portfolio and the procedure underestimates the risk profile of the portfolio [11]. This results in a need for dimension reduction.

In this paper, to address the dimension reduction problem, we take the alternative route for portfolio selection, which goes through the capital asset pricing model (CAPM). The milestone paper [19], leads to the CAPM, [22, 18, 1]. Although CAPM fails to explain several features, including rationality of the investors; however, the CAPM has become a standard tool in corporate finance [15]. The CAPM splits a portfolio return into *systematic* return and *idiosyncratic* return and models it as the linear regression of portfolio's 'excess returns' on market's 'excess returns'. In section 2, we discuss that when the maximum weight of the large portfolio is small, the idiosyncratic risk of a portfolio is washed out. Then the portfolio return can be mostly explained by market movements and corresponding regression coefficients, popularly known as portfolio's beta (β). The intercept of the model indicates if the asset is fairly valued or not. If the value of the intercept is zero, then the asset is fairly valued. The efficient market hypothesis indicates that the intercept of the CAPM must be zero.

Bayesian methods are proposed [21, 16] to test the restriction imposed in CAPM that the intercepts in the regression of 'excess returns' on the 'market excess return' are equal to zero. The non-zero intercept indicating the asset is either underpriced or overpriced, implies that the market is not efficient. Shanken's methodology [21] relies on the prior induced on functions of intercept and sampling distribution of F -statistics. Harvey and Zhou [16] proposed full Bayesian specification of the CAPM test with diffuse prior and conjugate prior structure. Black and Litterman [2] presented an informal Bayesian approach to economic views and equilibrium relations.

In this paper, we present the problem as multiple testing problems, under sparsity when the number of assets is high. Intercept for most of the assets under consideration are zero, and only very few of them are non-zero. That is, the null hypothesis; *the market is efficient*, means all the intercepts of all the assets are zero. An alternative hypothesis is at least one intercept is non-zero. The existing methodology concentrates on the test for intercept (or α) only. In our proposed methodology we present a joint test for both intercept and slope, (i.e., α and β); where α for most assets are zero and for β is one. The motivation for the joint test is provided in the next section.

Rest of the paper is organized as follows. In Section 2, we propose a portfolio selection strategy, which will mimic the market if the market is efficient and/or outperform the market if the market is not efficient. In Section 3 we presented the discrete-mixture prior model and hierarchical Bayes model with half-Cauchy distribution on the scale parameters. In Section 4 we presented the results of the asymptotic Bayes optimality for the multiple testing methodology proposed in section 3. In Section 5, we present a detailed simulation study. In Section 6, we present the empirical study based on data from New York Stock Exchange (NYSE). Then we conclude the paper with discussion in section 7.

2 Proposed Strategy

In matrix notation, the CAPM is presented as follows:

$$\mathbf{r} = \mathbf{X}\mathbf{B} + \boldsymbol{\epsilon}, \tag{2.1}$$

where $\mathbf{r} = ((r_{i,j}))_{n \times P}$ is the matrix of excess return over risk free rate for P many assets that are available in the market over n days; $\mathbf{X}\mathbf{B}$ is the systematic return due to market index, where

$$\mathbf{X} = ((\mathbf{1} \quad \mathbf{r}_m))_{n \times 2}$$

is the design matrix with first column is the unit vector or the place holder for intercept and the second column is the vector of n -days excess return of the market index over risk free rate;

$$\mathbf{B} = \begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_P \\ \beta_1 & \beta_2 & \dots & \beta_P \end{pmatrix}_{2 \times P};$$

if market is efficient then according to [22, 18, 1], the intercepts $\alpha_i = 0 \quad \forall i = 1(1)P$ and β_i is the measure of systematic risk due to market movement; $\boldsymbol{\epsilon} = ((\epsilon_{i,j}))_{n \times P}$ is the idiosyncratic return of the asset. The covariance of \mathbf{r} is $\boldsymbol{\Sigma}$, which can be decomposed into

$$\boldsymbol{\Sigma} = \mathbf{B}\boldsymbol{\Sigma}_m\mathbf{B}^T + \boldsymbol{\Sigma}_\epsilon,$$

$\boldsymbol{\Sigma}_\epsilon = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_P^2)$. The portfolio covariance can be decomposed into two parts as,

$$\begin{aligned} \mathbf{w}'\boldsymbol{\Sigma}\mathbf{w} &= \mathbf{w}'[\mathbf{B}\boldsymbol{\Sigma}_m\mathbf{B}^T + \boldsymbol{\Sigma}_\epsilon]\mathbf{w} \\ &= \mathbf{w}'\mathbf{B}\boldsymbol{\Sigma}_m\mathbf{B}^T\mathbf{w} + \mathbf{w}'\boldsymbol{\Sigma}_\epsilon\mathbf{w}, \end{aligned}$$

where first part explains the portfolio volatility due to market volatility and the second part explains portfolio volatility due to idiosyncratic behaviour of the stock. We assume σ_i 's are bounded $\forall i$. Then

$$\begin{aligned} \mathbf{w}'\boldsymbol{\Sigma}_\epsilon\mathbf{w} &= \sum_{i=1}^P \omega_i^2 \sigma_i^2 \\ &\leq \sigma_{max}^2 \sum_{i=1}^P \omega_i^2, \quad \sigma_{max}^2 = \max\{\boldsymbol{\Sigma}_\epsilon\} < \infty, \\ &\leq \sigma_{max}^2 M_{\omega:P} \sum_{i=1}^P \omega_i, \quad M_{\omega:P} = \max\{\|\mathbf{w}\|\}, \\ &= \sigma_{max}^2 M_{\omega:P}. \end{aligned}$$

Clearly, if $P \rightarrow \infty$ and $M_{\omega:P} \rightarrow 0 \implies \mathbf{w}'\boldsymbol{\Sigma}_\epsilon\mathbf{w} \rightarrow 0$. Hence we have the following result.

Result 2.1. *If $P \rightarrow \infty$ and $M_{\omega:P} \rightarrow 0$ and $\sigma_{max}^2 = \max\{\boldsymbol{\Sigma}_\epsilon\} < \infty$, then*

$$\lim_{P \rightarrow \infty} \mathbf{w}'\boldsymbol{\Sigma}_\epsilon\mathbf{w} = 0.$$

Remark 2.1. *Thus we can select $\tilde{P} (\ll P)$ many assets for the portfolio (out of P many assets available in the market), such that the idiosyncratic risk is washed out, i.e., for $\tilde{P} > \tilde{P}_0, \exists \epsilon > 0$, such that*

$$\|\mathbf{w}'\boldsymbol{\Sigma}_\epsilon\mathbf{w}\| < \epsilon;$$

and portfolio return is mostly explained by α and β only. Note that here \tilde{P} is the effective size of the portfolio. Therefore, Markowitz optimization (6.1) can be expressed as

$$\text{Min } \mathbf{w}'\mathbf{B}\boldsymbol{\Sigma}_m\mathbf{B}^T\mathbf{w} \quad \text{subject to } \mathbf{w}'\mathbf{1}_{\tilde{P}} = 1 \quad (2.2)$$

$$\text{and } \mathbf{w}'\mathbf{B}'\mathbf{X}_\mu = \mu_k,$$

where $\mathbf{X}_\mu = (1 \quad \mu_m)_{2 \times 1}^T$; μ_m is the expected excess market return, and the expected portfolio return $\mathbf{w}'\mathbf{B}'\mathbf{X}_\mu = \mu_k$ can be expressed as

$$\mu_k = \alpha_{\tilde{P}} + \beta_{\tilde{P}}\mu_m; \quad \alpha_{\tilde{P}} = \sum_{i=1}^{\tilde{P}} \omega_i \alpha_i \quad \text{and} \quad \beta_{\tilde{P}} = \sum_{i=1}^{\tilde{P}} \omega_i \beta_i \quad (2.3)$$

Remark 2.2. In (2.3), if $\alpha_{\tilde{P}} = 0$ and $\beta_{\tilde{P}} = 1$, then portfolio return will mimic the market return.

Remark 2.3. However, if $\alpha_{\tilde{P}} = 0$ and $\beta_{\tilde{P}} > 1$, then the portfolio return will mimic the behaviour of market return by $\beta_{\tilde{P}}$ factor. That is, if the $\beta_{\tilde{P}} = 1.25$ and market return increases by 3%, the the portfolio return increases by 3.75%. Similarly, if the market drops by 2%, then the portfolio return drops by 2.5%. Here portfolio will perform worse than the market specially during the recession period.

Remark 2.4. However, if $\alpha_{\tilde{P}} > 0$ and $\beta_{\tilde{P}} = 1$ then the portfolio will make superior return than market. That is if $\alpha_{\tilde{P}} = 0.01$, $\beta_{\tilde{P}} = 1$ and the market return increases by 3%, then from (2.3) the portfolio return is 3.01%, which is above the market return. Similarly, if the market drops by 2%, then the portfolio return drops by 1.99%. During the recession period, the portfolio return will be superior to the market return. It will also indicate the inefficiency of the market.

Remark 2.5. Oracle Set: If market is not efficient and there are q many assets whose $\alpha > 0$, where $q < \tilde{P} \ll P$, i.e.,

$$A_q = \{\alpha_j : \alpha_j > 0, j = 1, 2, \dots, q\},$$

then we can construct a portfolio $B_{\tilde{P}}$ with \tilde{P} many assets, such that

$$\mathbb{P}(A_q \subset B_{\tilde{P}}) \geq 1 - \eta, \quad (2.4)$$

where $0 < \eta < 1$ and $\mathbf{w}'_{\tilde{P}} \Sigma_{\tilde{P}, \epsilon} \mathbf{w}_{\tilde{P}} < \epsilon$. Note that $\mathbf{w}'_{\tilde{P}} = \{\omega_i : \max_{i=1(1)\tilde{P}} \|\omega_i\| < \epsilon\}$ and $\Sigma_{\tilde{P}, \epsilon} = \text{diag}(\sigma_1^2, \dots, \sigma_{\tilde{P}}^2)$ is the covariance matrix of idiosyncratic return. If the market is efficient the then A_q will be a null set.

The problem reduces to identifying the oracle set A_q . Essentially, it is a multiple testing problem, where we select those stocks in the portfolio $B_{\tilde{P}}$ for which the following hypothesis is rejected:

$$H_{0i} : \begin{pmatrix} \alpha_i \\ \beta_i \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \text{vs.} \quad H_{Ai} : \begin{pmatrix} \alpha_i \\ \beta_i \end{pmatrix} \neq \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad i = 1(1)P.$$

We would like to define an optimal test rule, such that (2.4) satisfies. Then run Markowitz's optimization technique to allocate weights (ω_i) on selected portfolio, such that $\max \omega_i < \epsilon \quad \forall i$.

Here, the structure of the multiple testing problem is very different, compare to typical multiple testing problem in the literature [5, 6, 9], which are mainly motivated from genome wide association study. In the next section, we present the Bayesian methodology to identify $B_{\tilde{P}}$.

3 Methodology

In this section, we propose Bayesian methodologies for testing $(\alpha_i = 0, \beta_i = 1) \quad \forall i = 1(1)P$. First we propose the discrete mixture prior and then we propose the hierarchical Bayes model. We know that the least squares estimator $\hat{\theta}_i = (\hat{\alpha}_i, \hat{\beta}_i)^T$, $i = 1(1)P$, is the MLE of $\theta_i = (\alpha_i, \beta_i)^T$, which is also sufficient statistics and the sampling distribution is $\hat{\theta}_i \sim N_2\left(\theta_i, \sigma_i^2 \Sigma_X^{-1}\right)$ where $\hat{\beta}_i = \hat{\rho}_i * \frac{s_{r_i}}{s_{r_M}}$, $\hat{\alpha}_i = \bar{r}_i - \hat{\beta}_i \bar{r}_M$, $\hat{\rho}_i$ is the correlation between r_i and r_M and

$$\Sigma_X = X^T X = \begin{bmatrix} n & \sum_{t=1}^n r_{tM} \\ \sum_{t=1}^n r_{tM} & \sum_{t=1}^n r_{tM}^2 \end{bmatrix}.$$

3.1 Discrete-mixture prior

We propose to use a discrete mixture prior, commonly known as spike and slab prior introduced by [20]. The prior puts probability $1 - p$ on $\theta_i = \mu_0$ and p on an absolutely continuous alternative as $[\theta_i | \Lambda_0] \sim N_2(\mu_0, \Lambda_0^{-1})$. Unconditionally, θ_i 's are independently distributed as

$$\theta_i \sim (1 - p)\delta_{\mu_0} + pN_2(\mu_0, \Lambda_0^{-1}),$$

where δ_{μ_0} is the degenerate distribution at zero. As $\theta_i = (\alpha_i, \beta_i)$, the corresponding $\mu_0 = (0, 1)$ implies prior mean of $\mathbb{E}(\alpha_i) = 0$ and $\mathbb{E}(\beta_i) = 1$. The parameter p is often known as the sparsity parameter and as $p \rightarrow 0$ the model becomes a sparse model and as $p \rightarrow 1$ the model is known as dense model. This implies that the marginal distribution of $\hat{\theta}_i$ is the scale mixture of normals, that is

$$\hat{\theta}_i \sim (1 - p)N_2(\mu_0, \sigma_i^2 \Sigma_X^{-1}) + pN_2(\mu_0, \sigma_i^2 \Sigma_X^{-1} + \Lambda_0^{-1}). \quad (3.1)$$

The conditional posterior distribution under the alternative is

$$[\theta_i | \sigma_i, \Lambda_0, \hat{\theta}_i] \sim N_2(\mu_{ni}, \Lambda_{ni}^{-1})$$

$$\text{where } \Lambda_{ni} = \Lambda_0 + \frac{\Sigma_X}{\sigma_i^2} \text{ and } \mu_{ni} = \Lambda_{ni}^{-1} \left(\Lambda_0 \mu_0 + \frac{\Sigma_X}{\sigma_i^2} \hat{\theta}_i \right).$$

Under a sparse mixture model, the Bayes oracle has the rejection region \mathcal{C} on which the Bayes factor exceeds $\frac{(1-p)\delta_0}{p\delta_A}$, see [3]. In this case, the Bayes factor can be computed as

$$\begin{aligned} & (\det(I - Q_i))^{T/2} \exp\left(\frac{S_i}{2}\right) \\ \text{where } & S_i = (\mu_{ni} - \mu_0)^T \Lambda_{ni} (\mu_{ni} - \mu_0) \\ & \text{and } Q_i = \frac{X}{\sigma_i} \Lambda_{ni}^{-1} \frac{X^T}{\sigma_i} \end{aligned}$$

The optimal rule is to reject H_{0i} if

$$S_i \geq c^2 = T \log(\det(I - Q_i)) + 2 \log(f\delta) \quad (3.2)$$

where $f = \frac{(1-p)}{p}$ and $\delta = \frac{\delta_0}{\delta_A}$. We call this rule *Bayes Oracle* since it makes use of unknown parameters p, Λ_0 and cannot be attained in finite samples. The posterior inclusion probability is

$$\tilde{\pi}_i = Pr(M_a | D) = \left(1 + \frac{1-p}{p} \exp\left(-\frac{S_i}{2}\right) \right)^{-1}, \quad \text{by Bayes theorem.}$$

Under symmetric loss, the rejection region coincides with $\tilde{\pi}_i > 1/2$. The test statistic involves

$$S_i = \frac{(y - X\mu_0)^T}{\sigma_i} Q_i \frac{(y - X\mu_0)}{\sigma_i}.$$

Under the null hypothesis, $(y - X\mu_0)/\sigma_i \sim \mathcal{N}(0, I)$. Thus S_i is a quadratic form in multivariate normal with Q_i symmetric non-idempotent of rank 2. The distribution is weighted sum of central χ^2 random variables of 1 df with weights being the eigenvalues of matrix Q , see [24]. In summary,

$$S_i \sim \lambda_1 \chi_1^2 + \lambda_2 \chi_2^2$$

where λ_1 and λ_2 are the two non-zero eigenvalues of Q and χ_1^2 and χ_2^2 are two independent central chi-square random variables with 1df. The distribution denoted by $M_2(\cdot; \lambda)$ is well studied for eg [23]. The probability of type I error is

$$t_{i1} = P(\lambda_1 \chi_1^2 + \lambda_2 \chi_2^2 \geq c^2).$$

The probability of type II error is

$$t_{i2} = P\left(\frac{\lambda_1}{1 - \lambda_1} \chi_1^2 + \frac{\lambda_2}{1 - \lambda_2} \chi_2^2 \leq c^2\right).$$

The derivation is as follows: Under the alternative hypothesis the marginal of the data is, $(y - X\mu_0) \sim \mathcal{N}(0, \sigma^2 A_i)$ where $A_i = (I - Q_i)^{-1}$. The distribution of S_i is weighted sum of central χ^2 random variables of 1 df with weights being the eigenvalues of matrix $A_i^{1/2} Q_i A_i^{1/2} = (I - Q_i)^{-1} - I$. Eigenvalues of this matrix are $\frac{\lambda_j}{1 - \lambda_j}$, where $\lambda_j, j = 1, 2$ are eigenvalues of Q as before. Under additive loss function, the Bayes risk of the Bayes Oracle is

$$R_{\text{opt}} = (1 - p)\delta_0 \sum_{i=1}^P t_{i1} + p\delta_A \sum_{i=1}^P t_{i2}.$$

3.2 Hierarchical Bayesian Model

We present the hierarchical Bayesian model in the spirit of [13], which is as follows:

$$\begin{aligned} r_{ti} &\sim N(\alpha_i + \beta_i r_{tM}, \sigma_i^2), \\ \theta_i | \Lambda &\sim N_2(\theta_0, \tau^2 \Lambda^{-1}), \end{aligned}$$

where $\theta_i = (\alpha_i, \beta_i)$, and $\theta_0 = (\alpha_0, \beta_0)$

$$\begin{aligned} \sigma_i^2 &\propto \text{InvGamma}\left(\frac{\nu_0}{2}, \frac{\nu_0}{2}\right), \\ \Lambda &\sim \text{Wishart}\left((\rho R)^{-1}, \rho\right), \\ \theta_0 &\sim N_2(\mu_0, C), \\ \tau^2 &\sim C^+(0, 1) \end{aligned}$$

where R is the prior scale matrix, ρ is prior degrees of freedom of the Wishart distribution, $\mu_0 = (0, 1)^T$ and $C^+(0, 1)$ denotes half-Cauchy distribution with location parameter 0 and scale parameter 1 with corresponding pdf as

$$f(\tau) = \frac{2\mathbb{I}(\tau > 0)}{\pi(1 + \tau^2)}.$$

The parameter τ plays a crucial role in controlling the shrinkage behavior of the estimator. It is known as ‘‘global shrinkage parameter’’ [5, 6, 9], as it adjust to to the overall sparsity in the data. The posterior probability of τ is concentrated near zero when the data is very sparse ($p \rightarrow 0$). We define $\bar{\theta} = P^{-1} \sum_{i=1}^P \theta_i$, $n = \sum_{i=1}^P n_i$,

$$\begin{aligned} D &= [(X^T X) + \Lambda]^{-1}, \\ V &= (P\Lambda + C^{-1})^{-1}. \end{aligned}$$

The Gibbs sampler for θ_i , σ_i^2 , Λ and θ_0 is straight forward as

$$[\theta_i|X, r_i, \mu_c, \Lambda, \sigma_i^2] \sim N_2(m_i, \Sigma_i),$$

where $m_i = D(X^T r_i + \Sigma_c \mu_c)$ and $\Sigma_i = \sigma_i^2 D$,

$$[\sigma_i^2|n_i, r_i, \theta_i] \sim \text{InvGamma}\left(\frac{\nu_0 + n_i}{2}, \frac{\nu_0 + (r_i - X\theta_i)^T(r_i - X\theta_i)}{2}\right),$$

$$[\Lambda|\theta_i, P, R, \rho, \mu_c] \sim \text{Wishart}\left\{\left(\sum_{i=1}^P (\theta_i - \mu_c)(\theta_i - \mu_c)^T + \rho R\right)^{-1}, P + \rho\right\},$$

$$[\theta_0|V, P, \Lambda, C, \mu_0] \sim N_2\left(V(P\Lambda\bar{\theta} + C^{-1}\mu_0), V\right).$$

We implemented a Metropolis-Hastings update for τ . Here τ acts as global shrinkage parameter and Λ behaves as local shrinkage parameters. Note that we consider the half-Cauchy prior [14] over the scale parameter τ . $\tau \sim C^+(0, 1)$. Later [5, 6] showed that such prior specification is suited for high-dimension sparse solution problem and named it as ‘horseshoe prior’.

4 Asymptotic Optimality

[3] introduced the notion of ‘Asymptotic Bayes Optimality under Sparsity’ (ABOS). To our knowledge, this is the only notion of optimality for multiple testing. This has been extended to show optimality of one-group models in [9]. In particular, it is shown that if the global shrinkage parameter τ of the horse-shoe prior is chosen to be the same order as p , then the natural decision rule induced by the Horseshoe prior attains the risk of the Bayes oracle upto $O(1)$ with a constant close to the constant in the oracle. There have been several studies following this in the one-parameter setting. Our aim is to extend these results to the multi-parameter setting.

The asymptotic framework that we work under is motivated by [3]. For the two-dimensional setting, Assumption A is modified to the set of $\gamma_t = \{p_t, \Lambda_{0t}, \sigma_{it}, \delta_t\}$ satisfying

$$p_t \rightarrow 0, \sigma^2 \Lambda_0 \rightarrow 0, v := \det(I - Q)^{T/2} f \delta \rightarrow \infty, \\ \sqrt{(1 - \lambda_1)(1 - \lambda_2)} \log(v) \rightarrow C \in (0, \infty).$$

Theorem 1. *Under assumption A, $t_{i1} \rightarrow 0$ and $t_{i2} \rightarrow 1 - e^{-C/2}$.*

Proof. It has been seen in section 3.1 that, under the alternative hypothesis, S_i is a linear combination of 2 independent χ_1^2 random variables with weights λ_1 and λ_2 the non-zero eigenvalues of Q_i . Under assumption A, λ_1 and λ_2 converge to 1 as $t \rightarrow \infty$. Hence $S_i \Rightarrow \chi_2^2$. Since $c_i^2 \rightarrow \infty, t_{i1} \rightarrow 0$.

Under the alternative hypothesis S_i is a linear combination of 2 independent χ_1^2 random variables with weights $\lambda_1/(1 - \lambda_1)$ and $\lambda_2/(1 - \lambda_2)$. Under assumption A, λ_1 and λ_2 converge to 1. Hence $\sqrt{(1 - \lambda_1)(1 - \lambda_2)} S_i \Rightarrow \chi_2^2$. Also, $\sqrt{(1 - \lambda_1)(1 - \lambda_2)} \log(v) \rightarrow C$. Hence $t_{i2} \rightarrow P(X \leq C)$ where X is a χ_2^2 random variable, hence the result. \square

Theorem 2. *Under the Bayes oracle, the risk takes the form $R_{\text{opt}} = mp\delta_A(1 - e^{-C/2})$*

Definition 1. *Consider a sequence of parameters γ_t satisfying Assumption A. We call a multiple testing rule ABOS for γ_t if its risk R satisfies*

$$\frac{R}{R_{\text{opt}}} \rightarrow 1 \quad \text{as } t \rightarrow \infty$$

We propose an alternative test with rejection region $\tilde{S}_i > c^2$, where c^2 is as defined in equation (3.2) and

$$\tilde{S}_i = \frac{(y - X\mu_0)^T}{\sigma_i} X(X^T X)^{-1} X^T \frac{(y - X\mu_0)}{\sigma_i}.$$

The advantage of \tilde{S}_i is that it does not depend on Λ_0 . We show that this new test is ABOS.

Theorem 3. *The test that rejects H_0 when $\tilde{S}_i > c_{it}^2$ is ABOS if and only if $c_{it} \rightarrow \infty$ and $c_{it}^2 \sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})} \rightarrow C$.*

Proof. Under H_0 , $\tilde{S}_i = \frac{(y - X\mu_0)^T}{\sigma_i} \tilde{Q}_i \frac{(y - X\mu_0)}{\sigma_i}$ is a quadratic form in standard multivariate normal with $\tilde{Q}_i = X(X^T X)^{-1} X^T$ symmetric idempotent of rank 2. Hence $\tilde{S}_i \sim \chi_2^2$. $t_{i1} = P(\tilde{S}_i > c_{it}^2) \rightarrow 0$ if and only if $c_{it} \rightarrow \infty$.

Under the alternative, $Z = \frac{(y - X\mu_0)^T}{\sigma_i} A^{-1/2}$ is standard normal and $\tilde{S}_i = Z^T A^{1/2} \tilde{Q}_i A^{1/2} Z$. So $(\det(A))^{-1/2} \tilde{S}_i \Rightarrow \chi_2^2$. Also, $\det(A) = (1 - \lambda_1)(1 - \lambda_2)$. The same argument as in Thm 1 shows that $t_{i2} \rightarrow 1 - e^{-C/2}$ if and only if $c_{it}^2 \sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})} \rightarrow C$.

The ABOS property is now established using the Theorem 2. □

The Bayesian False Discovery Rate (BFDR) was introduced by [10] as

$$\text{BFDR} = P(H_{0i} \text{ is true} \mid H_{0i} \text{ is rejected}) = \frac{(1 - p)t_1}{(1 - p)t_1 + pt_2}.$$

It has been seen that multiple testing procedures controlling BFDR at a small level α behave very well in terms of minimizing the misclassification error, see eg [4].

Consider a fixed threshold rule based on \tilde{S}_i with BFDR equal to α . Under the mixture model (3.1), a corresponding threshold value c^2 can be obtained by solving the equation

$$\frac{(1 - p)e^{-c^2/2}}{(1 - p)e^{-c^2/2} + pe^{-c^2/2}\sqrt{(1 - \lambda_1)(1 - \lambda_2)}} = \alpha \quad (4.1)$$

Theorem 4. *Consider a fixed threshold rule with $\text{BFDR} = \alpha = \alpha_t$. The rule is ABOS if and only if it satisfies the following two conditions*

$$r_\alpha/f \rightarrow 0, \text{ where } r_\alpha = \alpha/(1 - \alpha) \quad (4.2)$$

and

$$\frac{2 \log(r_\alpha/f)}{1 - \frac{1}{\sqrt{(1 - \lambda_1)(1 - \lambda_2)}}} \rightarrow C. \quad (4.3)$$

The threshold for this rule is of the form

$$c_t^2 = C - 2 \log(r_\alpha/f) + o_t. \quad (4.4)$$

Proof. Suppose the test is ABOS. Equation 4.1 is equivalent to

$$\frac{p}{1 - p} \frac{\alpha}{1 - \alpha} = e^{-c^2/2(1 - \sqrt{(1 - \lambda_1)(1 - \lambda_2)})}$$

By theorem 3, the right hand side goes to zero. This implies left hand side = r_α/f goes to zero, establishing the first condition.

Simplifying Equation 4.1 and using $c_{it}^2 \sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})} \rightarrow C$ we have,

$$2 \log(r_\alpha/f) = c_t^2 + C + o(t) \tag{4.5}$$

that is, the threshold is of the form given by equation 4.4.

Furthermore, $c_{it}^2 = C/\sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})} + o(t)$. Combining this with equation 4.5, we get the second condition.

Now we prove the converse.

Suppose a test with $\text{BFDR}=\alpha$ satisfies the two conditions. Let us define z_t as $z_t = c_{it}^2 \sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})}$. Such a test satisfies 4.1. Hence,

$$2 \log(r_\alpha/f) = z_t \left(1 - \frac{1}{\sqrt{(1 - \lambda_{1t})(1 - \lambda_{2t})}}\right).$$

Combining this with 4.3, we have $z_t \rightarrow C$.

Also, from 4.4, $2 \log(r_\alpha/f) = z_t - c_t^2$. Combining this with 4.2 and $z_t \rightarrow C$, we have $c_t^2 \rightarrow \infty$.

Now by using theorem 3, the test is ABOS. □

5 Simulation Study

In this section, we present two different simulation studies. In the study 1, we compare the performance of S_i and \tilde{S}_i . In the study 2, we compare the performance of Bayes Oracle estimator \tilde{S}_i with the other methods like diffuse prior and LARS-LASSO [12]. For both studies we simulate the data from a true model given by equation (2.1) with $\sigma_i = \sigma$ for all i . Without loss of generality we consider first $[pP]$ many stocks are not fairly priced. That is we simulate α_i and β_i for those stocks from $N(0, 0.1)$ and $N(1, 0.1)$ respectively. Rest of the stock's α and β are being set as $(0,1)$.

Study 1: In this study, we consider two different choices of P , i.e., $P = 100$ and 500 and the sample size is varied from $n = 20$ to $n = 50$ by an interval of 5. *Note that due to space constraint we present the result for $n = 20$ and $n = 50$ in figure 1.* We allow the sparsity parameter p to vary from 0.01 to 0.9 by an interval of 0.01. We choose two different values of $\sigma = (0.1, 0.05)$, and $\Lambda_0 = \begin{pmatrix} 0.5 & 0.3 \\ 0.3 & 0.7 \end{pmatrix}$. For all these different choices of $n, P, \alpha, \beta, \sigma$; we simulate 1000 datasets. For each dataset, we compute S_i and \tilde{S}_i and make a decision. Based on the decision over 1000 datasets we compute type-I error, type-II error, BFDR and the probability of misclassification (PMC) and present the results in Figure 1 and 2. We report the following observations.

Observations

1. As sparsity tends to 0 the type-I error goes to 0 in all four panels of the Figure 1. In all panels of the Figure 1, sparsity near to 0, the type-II error does not shoot to 1. The bounded type-II error guarantees particular statistical power of the ABOS. This observation verifies the Theorem 1
2. From all four panels of the figure 1, we observe for different choices of n, P, σ and sparsity p , all the four metrics of S_i and \tilde{S}_i overlap. This verifies the Theorem 3.
3. With increasing n , the probability of type-II error and the probability of misclassification drop. This indicates increasing statistical power even when sparsity parameter p is near zero.

4. As $p \rightarrow 1$, i.e., the model becomes dense, one should not use this test, as type-I error increases. However, up to 0.5 of the sparsity, the type-I error stays below 5% level.
5. In all four panels of the figure 1 the BFDR is about 0.05 irrespective the value of n , P , σ and p .
6. Figure 2 indicates that with increasing number of stocks from $P = 10$ to $P = 500$, all the metrics of the test becomes smoother.

Study 2: In this study, we consider $P = 500$, the sample size $n = 20$, $\sigma = 0.1$ and $\Lambda_0 = \begin{pmatrix} 0.5 & 0.3 \\ 0.3 & 0.7 \end{pmatrix}$.

For all these choices of parameters, we simulate 1000 datasets. For each dataset, we compute \tilde{S}_i and make a decision. We also make the decision using diffuse-prior and LARS-LASSO technique and compare against original decision. Based on the decision on 1000 datasets we compute type-I error, type-II error, Bayesian False Discover Rate (BFDR) and the probability of misclassification and present the results in Figure 3.

Observations

1. As the sparsity tends to 0, the type-I error of ABOS goes to 0. In likelihood testing, the type-I error is fixed at 5% level throughout the different values of sparsity. The LARS-LASSO method also demonstrates a flat behavior. However, it is more than the likelihood method.
2. If we compare the type-II error, BFDR and the probability of misclassification for all three methods, ABOS is uniformly better than other two methods.

Note: We tried to compare ABOS against Hierarchical Bayes (HB) method. However, given the computational power it took about 3 days to implement the HB method for 1000 simulated datasets, for one fixed sparsity parameter. For each dataset we simulated 25000 MCMC simulations after 5000 burn-in. For the study 2, we consider the sparsity ranges from 0.01 to 0.9 by an interval of 0.01. For one sparsity value it was taking approximately 3 days and for all 90 possible values it will take approximately 270 days, assuming no possible disruption in the systems. Hence we could not implemented the comparison for lack of computational resources. Hence we leave this task as future research project.

6 Empirical Study

In this empirical study we considered nine years of daily return from Jan 2006 to Dec 2014. The purpose of choosing this is to study the behavior of the methods specially during the stress period of 2008 and 2011. On the t^{th} month, we run the modeling procedure described in section 3 over the daily return. As there are P many assets with excess returns over risk-free rate r_1, \dots, r_P in the market and $P \gg n$. Here n is typically 22 or 23 days of return, as there are only that many business days in a month. We select \tilde{P} many assets using the methodology described above and use the daily return of t^{th} month. Note that still \tilde{P} could be larger than n . We select the stocks which are under-priced and our revised portfolio for $(t + 1)^{st}$ month would be with these under-priced stocks. Once we select \tilde{P} many assets for the portfolio then the problem reduces to portfolio allocation and we solve it as risk minimization problem (i.e., quadratic programming problem) described in equation (6.1). We allocate weights based on Markowitz's solution applied on the selected portfolio.

6.1 Portfolio Optimization

The theory of Markowitz portfolio selection analytically formalizes the risk return tradeoff in selecting optimal portfolios. However, the Markowitz portfolio is very sensitive [12] to errors in the estimates of the expected return and the covariance matrix. The problem is severe when the portfolio size is large. Several techniques have been suggested to reduce the sensitivity of the Markowitz optimal portfolios to input uncertainty. One approach is to use a James-Stein estimator for means (i.e., expected return) [7] and shrink the sample covariance matrix [17, 8].

In standard framework this risk is measured by the variance of the portfolio whereas the expectation by the mean of the portfolio. There are P many assets with excess returns over risk-free rate r_1, \dots, r_P to be managed. Let \mathbf{r} be the return vector, Σ is the corresponding covariance matrix and \mathbf{w} be its portfolio allocation vector. In ideal situation the means, variances and covariances are known and the problem is the following quadratic programming problem:

$$\text{Min } \mathbf{w}'\Sigma\mathbf{w} \quad \text{subject to } \mathbf{w}'\mathbf{1}_P = 1 \quad \text{and} \quad \mathbf{w}'\boldsymbol{\mu} = \mu_k \quad (6.1)$$

Here $\mathbf{1}_P$ is a P -dimensional vector with one in every entry and μ_k is the desired level of return.

In practice, Σ and $\boldsymbol{\mu}$ are unknown. The most common procedure known as ‘‘plug-in’’ implementation replaces them with their method of moments estimators, $\hat{\boldsymbol{\mu}}$ and $\hat{\Sigma}$ to obtain the optimal weights \mathbf{w}_{opt} . The curve $\mathbf{w}'_{opt}\hat{\Sigma}\mathbf{w}_{opt}$ seen as a function of μ_k is called the *efficient frontier*.

In the Markowitz setting, let us assume that n and P both go to infinity and each $\mathbf{X}_i \sim \mathcal{N}_P(\boldsymbol{\mu}, \Sigma)$ independently and identically. The parameters of the distribution are estimated using sample estimators. We have from Corollary 3.3 of [11],

$$\begin{aligned} \mathbf{w}'_{emp}\hat{\Sigma}\mathbf{w}_{emp} &= \left(1 - \frac{P-2}{n-1}\right) \left(\mathbf{w}'_{theo}\Sigma\mathbf{w}_{theo} - \frac{P}{n} \frac{(U'M^{-1}\mathbf{e}_2)^2}{(1 + \frac{P}{n}\mathbf{e}'_2M^{-1}\mathbf{e}_2)}\right) \\ &\quad + o_P(\mathbf{w}'_{theo}\hat{\Sigma}\mathbf{w}_{theo} \vee n^{-1/2}) \end{aligned}$$

where $M = V'\Sigma^{-1}V$ and \mathbf{w}_{emp} represents the weights obtained from the empirical data at hand while \mathbf{w}_{theo} is it’s population counterpart. \mathbf{e}_k denote the canonical basis vectors in \mathbb{R}^2 . The corollary shows that the effects of covariance and mean estimation are to underestimate the risk.

6.2 Analysis

We invest in the selected stocks for month $t+1$ and calculate the out-sample return of the portfolio and S&P 500 Index at the same time. We considered the out-sample period to be from Jan 01, 2006 to Dec 31, 2014. For example: we run the statistical processes on the stock price of Dec 2007 and identify the 25 stocks and construct the portfolio using the Markowitz weights for Jan 2008 and invest only in these stocks. We repeat the process for each month.

Figure 4, shows the performance of four different strategies in out-sample. Here S&P 500 index is being considered as benchmark portfolio, as many investors invest in S&P500 index fund as passive investors. If we look at the table 1, except three years (2008,2013,2014) the hierarchical-Bayes selection strategy outperforms the benchmark index in terms of annual return. Out of 9 years, there are four years (2007,2011,2012,2014) where LARS-LASSO selection process underperforms compared to benchmark in terms of annual return. Similarly, out of the same 9 years, only during 2012 the Bayesian method with diffuse prior selection process underperforms compared to benchmark in terms of annual return. In terms of annual return all three portfolio selection process are outperforming the benchmark in most of the years; but we cannot clearly say one

outperform the other for all years. There is no clear winner in terms of annual return. However, all four strategies indicate possible existence for inefficiency in the market.

In table 2 we present the out-sample annualized volatility of all the selection strategies. The annualized volatility of hierarchical Bayes is either lower or marginally higher than the benchmark. The out-sample annualized volatility of LARS-LASSO strategy is marginally higher than the benchmark for all years. However, the annualized volatility of Bayes with diffuse prior selection strategy is significantly higher than other three alternatives. Figure 5 presents the daily annualized volatility of all four selection strategies in out-of the sample return. This volatility is being estimated by GARCH(1,1) model and clearly we can see the daily annualized volatility for diffuse prior is significantly higher than other three strategies. The daily annualized volatility of benchmark S&P 500 index fund and the hierarchical selection strategies have similar level. The same for LARS-LASSO strategy is slightly higher but consistently lower than diffuse prior strategy. Therefore we can say that in terms of the volatility risk in out-sample diffuse prior strategy is most risky compared to other three alternatives.

Table 3 presents the ‘Value at Risk’ (VaR) of different strategies in out-sample. Out of 9 years, there are five years (2006,2008-11) where hierarchical Bayes strategy has lowest VaR among all four strategies. The benchmark S&P 500 index fund has lowest VaR in another three years (2012-2014). The LARS-LASSO strategy experiences lowest VaR only during 2007. The VaR of Bayes method with diffuse prior strategy is always highest among the four strategies we consider which implies that in terms of extreme risk measures like VaR, the diffuse prior strategy is the most risky compared to others.


Table 4 presents the Sharpe ratio or ‘risk adjusted return’ of different strategies in out-sample. Clearly there is no single winner. There are five years where hierarchical Bayes strategy has highest Sharpe ratio, another five years where diffuse prior strategy has the highest Sharpe ratio. There are three years where the Sharpe ratio is highest for the LARS-LASSO strategy.

7 Discussion

We presented Bayesian portfolio selection strategy, via the Capital Asset Pricing Model (CAPM). If the market is information efficient, the proposed strategy will mimic the market; otherwise, the strategy will outperform the market. The strategy depends on the selection of a portfolio via Bayesian multiple testing methodologies for the parameter of the CAPM. We present the “discrete-mixture prior” model and “hierarchical Bayes model” for the intercept and slope parameters of CAPM. We prove that under the asymptotic framework of [9] the Bayes rule attains the risk of Bayes Oracle up to $O(1)$ with a constant close to the constant in the Oracle.

We present detail empirical study, where 500 stocks from the New York Stock Exchange (NYSE) are considered, and S&P 500 index is considered as the benchmark. We present a detailed study of portfolio selection via different methodologies over the period from the year 2006 to 2014. The out of the sample risk and return performance of the portfolio selected by the various methods are presented. Empirical results indicate the existence of inefficiency of the market, and it is possible to propose a strategy which can outperform the market.

Figure 1: Performance of S_i and \tilde{S}_i with 100 stocks, i.e., $P = 100$ with 20 and 50 days of data, i.e., sample size $n = 20$ and $n = 50$



References

- [1] F. Black. Capital market equilibrium with restricted borrowing. *Journal of Business*, 45: 444–454, 1972.
- [2] F. Black and R. Litterman. Global portfolio optimization. *Financial Analysts Journal*, 48: 28–43, 1992.
- [3] M. Bogdan, A. Chakraborty, F. Frommlet, and J. Ghosh. Asymptotic bayes optimality under sparsity of some multiple testing procedures. *The Annals of Statistics*, 39(3):1551–1579, 2011.

Figure 2: Performance of S_i and \tilde{S}_i with 10 and 500 stocks, i.e., $P = 10$ and $P = 500$ with 20 days of data, i.e., sample size $n = 20$




Table 1: Out-sample Annual Return

	Benchmark	Hierarchical Bayes	Diffuse Prior	LARS-LASSO	Bayes Oracle
2006	11.24	23.02	12.66	39.43	17.28
2007	2.34	4.50	27.51	-5.77	7.97
2008	-42.67	-46.81	-56.22	-41.09	-43.19
2009	15.37	27.83	34.61	23.98	23.18
2010	9.22	10.35	12.52	17.91	23.54
2011	-3.78	1.22	-2.35	-3.41	2.30
2012	10.79	22.34	7.65	7.63	14.52
2013	25.64	19.49	41.12	31.75	33.53
2014	11.66	11.48	20.80	9.82	5.35

- [4] Małgorzata Bogdan, Jayanta K Ghosh, Aleksandra Ochman, and Surya T Tokdar. On the empirical bayes approach to the problem of multiple testing. *Quality and Reliability Engineering International*, 23(6):727–739, 2007.
- [5] C. Carvalho, N. Polson, and J. Scott. Handling sparsity via the horseshoe. *Journal of Machine Learning Research*, 5:73–80, 2009.
- [6] C. Carvalho, N. Polson, and J. Scott. The horseshoe estimator for sparse signals. *Biometrika*, 97(2):465–480, 2010.
- [7] V. K. Chopra and W. T. Ziemba. The effect of errors in means, variance and covariances on optimal portfolio choice. *The Journal of Portfolio Management*, 19:6–11, 1993.
- [8] S. Das, A. Halder, and D. K. Dey. Regularizing portfolio risk analysis: A bayesian approach. *Methodology and Computing in Applied Probability*, 2016. doi: 10.1007/s11009-016-9524-5.

Table 2: Annualized Volatility of out-sample return

	Benchmark	Hierarchical Bayes	Diffuse Prior	LARS-LASSO	Bayes Oracle
2006	9.90	10.10	16.24	11.10	12.46
2007	16.09	15.36	21.48	16.26	18.53
2008	41.26	34.79	53.24	43.22	47.99
2009	27.16	23.12	49.33	30.49	34.46
2010	18.05	16.58	24.86	19.59	20.17
2011	23.48	21.62	29.14	23.57	28.29
2012	12.69	12.59	18.01	14.92	14.77
2013	10.84	11.85	14.81	13.07	12.37
2014	11.38	11.55	14.06	12.84	13.03

Table 3: Value at Risk of out-sample return

	Benchmark	Hierarchical Bayes	Diffuse Prior	LARS-LASSO
2006	1.66	1.55	2.57	1.58
2007	2.88	2.79	3.30	2.49
2008	8.95	7.32	11.37	8.49
2009	4.83	3.78	8.47	5.36
2010	3.28	2.69	4.56	3.55
2011	4.65	3.88	5.61	4.16
2012	2.02	2.13	3.03	2.13
2013	1.74	1.88	2.44	2.09
2014	2.12	2.13	2.32	2.33

- [9] J. Datta and J. K. Ghosh. Asymptotic properties of bayes risk for the horseshoe prior. *Bayesian Analysis*, 8(1):111–132, 2013.
- [10] Bradley Efron and Robert Tibshirani. Empirical bayes methods and false discovery rates for microarrays. *Genetic epidemiology*, 23(1):70–86, 2002.
- [11] N. El Karoui. High-dimensionality effects in the markowitz problem and other quadratic programs with linear constraints: risk underestimation. *The Annals of Statistics*, 38(6):3487–3566, 2010.
- [12] J. Fan, J. Zhang, and K. Yu. Vast portfolio selection with gross-exposure constraints. *Journal of the American Statistical Association*, 107:592–606, 2012.
- [13] A. E. Gelfand, S. E. Hills, A. Racine-Poon, and A. F. M. Smith. Illustration of bayesian inference in normal data models using gibbs sampling. *Journal of American Statistical Association*, 85:972–985, 1990.
- [14] A. Gelman. Prior distributions for variance parameters in hierarchical models(comment on article by browne and draper). *Bayesian Analysis*, 1:515–534, 2006.
- [15] A. Goyal. Empirical cross-sectional asset pricing: a survey. *Financial Market Portfolio Management*, 26:3 – 38, 2012.
- [16] R. C Harvey and G. Zhou. Bayesian inference in asset pricing tests. *Journal of Financial Economics*, 26:221–254, 1990.

Table 4: Sharpe Ratio of out-sample return

	Benchmark	Hierarchical Bayes	Diffuse Prior	LARS-LASSO
2006	0.07	0.13	0.05	0.19
2007	0.01	0.02	0.07	-0.02
2008	-0.08	-0.11	-0.10	-0.08
2009	0.03	0.07	0.04	0.04
2010	0.03	0.04	0.03	0.05
2011	-0.01	0.00	-0.01	-0.01
2012	0.05	0.10	0.03	0.03
2013	0.13	0.10	0.15	0.13
2014	0.06	0.06	0.09	0.05

- [17] O. Ledoit and M. Wolf. Improved estimation of the covariance matrix of stock returns with an application to portfolio selection. *Journal of Empirical Finance*, 10:603–621, 2003.
- [18] J. Lintner. The valuation of risky assets and the selection of risky investments in stock portfolios and capital budgets. *Review of Economics and Statistics*, 47:13–37, 1965.
- [19] H. Markowitz. Portfolio selection. *The Journal of Finance*, 7(1):77–91, 1952.
- [20] T J Mitchell and J J Beauchamp. Bayesian variable selection in linear regression. *Journal of the American Statistical Association*, 83(404):1023–1032, 1988.
- [21] J. Shanken. A bayesian approach to testing portfolio efficiency. *Journal of Financial Economics*, 19:195–216, 1987.
- [22] W. Sharpe. Capital asset prices:a theory of market equilibrium under conditions of risk. *Journal of Finance*, 19:425–442, 1964.
- [23] H. Solomon and M. A. Stephens. Distribution of a sum of weighted chi-square variables. *Journal of the American Statistical Association*, 72(360):881–885, 1977.
- [24] Q. H. Vuong. Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica*, 57(2):307–333, 1989.

Figure 3: Performance Comparison of ABOS, Diffuse prior and LARS-LASSO, with 100 stock and 20 days of data and simulation size 100




Figure 4: Performance of a different portfolios in outsample




Figure 5: Volatility of a different portfolios in outsample modeled through GARCH(1,1)

